Monocyte Metabolic Plasticity and Cytokine Production Differentiate Latent TB Infection from Active Disease

Running title: Monocyte Metabolism and Function in Latent vs Active TB

Gráinne Jameson[1], Isabella Batten[1], Adam H. Dyer[1], Caoimhe Geoghegan[3], Moninne Murray[3], Niamh McDonnell[3], Dearbhla M. Murphy[1], Sarah A. Connolly[1], Anne Marie McLaughlin[2,3], Cilian Ó Maoldomhnaigh[1], Laura E. Gleeson[1,3], Joseph Keane[1,2,3], Sharee A. Basdeo[1]

[1]School of Medicine, Trinity Translational Medicine Institute, St James's Hospital, Trinity College Dublin, Dublin, Ireland.
[2]National Tuberculosis Centre, Department of Respiratory Medicine, St James's Hospital, Dublin, Ireland.
[3]Respiratory Department, St James's Hospital, Dublin 8, Dublin, Ireland.

Corresponding author:
Gráinne Jameson
Trinity Translational Medicine Institute,
Trinity Centre for Health Sciences,
St. James's Hospital,
Dublin 8
GJAMESON@TCD.IE, +353 1 896 3263

Authors Contributions:
G.J. and S.A.B. conceived the study and hypotheses. G.J., I.B., A.H.D., C.G., M.M., N.M., A.M., L.E.G. and J.K. participated in data collection. J.K, L.E.G., C.O.M. and A.M. provided domain expert advice and evaluation of the project. G.J., I.B., A.H.D., D.M., S.A.C., C.O.M. and S.A.B. performed data and statistical analyses. All authors interpreted the data. G.J. and S.A.B. wrote the initial draft of the manuscript. All authors provided substantive critical reviews and approved of the submitted manuscript.

Sources of Support: This work was funded by the Health Research Board Ireland (HRB-ILP-POR-2022-033 and HRB-EIA-2019-010; awarded to S.A.B.) and The Royal City of Dublin Hospital Trust (RCDH app 185; Awarded to J.K.). The following reagent was obtained through BEI Resources, NIAID, NIH: Mycobacterium tuberculosis, Strain H37Rv, Gamma-Irradiated Whole Cells, NR-49098).

Total word count: 38033

Abstract:

Rationale:

Monocytes are central to host defence against *Mycobacterium tuberculosis* (Mtb), yet their

functional and metabolic profiles during latent TB infection (TBI) and active TB disease (TBD)

remain poorly defined. Immunometabolic dysfunction may underlie ineffective responses in TB,

but cell-specific mechanisms are unclear.

Objectives:

To compare the phenotypic, functional, and metabolic profiles of circulating monocytes from

individuals with TBI, TBD, and healthy controls (HC), and assess the impact of treatment.

Measurements:

Peripheral blood monocytes were profiled using high-dimensional flow cytometry, Luminex

cytokine/chemokine assays, and SCENITH™, a flow-based metabolic assay. Unstimulated and

Mtb-stimulated monocytes from treatment-naïve and treated individuals were analysed.

Main Results:

Monocytes from TBI and TBD showed distinct phenotypes from HC, marked by elevated CD14

and CD45RA. HLA-DR was reduced in TBI versus HC and further decreased in TBD. TNF

receptors were downregulated in TBI but unchanged in TBD. Baseline cytokine and chemokine

profiles in TBI and TBD were similar (yet distinct from HC), but Mtb stimulation elicited a stronger

cytokine response in TBI. Metabolically, TBI and TBD monocytes exhibited increased glycolysis

and reduced mitochondrial dependence versus HC. Treatment partially restored mitochondrial

function. Upon Mtb challenge, TBI monocytes had higher glycolytic capacity than TBD.

Conclusions:

Monocyte metabolic plasticity and cytokine production distinguish latent from active TB and are

partially reversible with treatment. Circulating monocyte metabolism reflects TB immune status

and may serve as a biomarker or therapeutic target. Reprogrammed glycolytic profiles in TBI

contrast with impaired adaptability in TBD, suggesting dysfunctional myeloid activation during

disease.

Introduction

Tuberculosis (TB) is the leading cause of death by an infectious agent, with an estimated 1.25 million deaths and 10.8 million cases in 2023 (1). While most individuals infected with *Mycobacterium tuberculosis* (Mtb) do not progress to active disease, those with latent TB infection (TBI) carry approximately twice the risk of developing tuberculosis disease (TBD compared with uninfected individuals), yet also demonstrate a 79% lower risk of progressive disease following reinfection (2). Understanding the immunological mechanisms that confer protection in TBI or promote progression to TBD is a central question in TB research.

Monocytes are short-lived, circulating myeloid cells that play a critical role in the peripheral immune response to infection. They contribute to antimicrobial defence through phagocytosis, cytokine production, and antigen presentation (3–7). In TB, monocytes are recruited from the circulation to the lungs, the primary site of infection, where they can differentiate into alveolar macrophages (AMs) (8–12). The AM niche almost exclusively harbours early Mtb infection and acts as the first line of defence (13,14).

Cellular metabolism has emerged as a key regulator of myeloid function. Glycolysis, in particular, drives effective antimicrobial activity in both human AM and monocyte-derived macrophages responding to Mtb (15–18). Similar metabolic reprogramming occurs in healthy monocytes upon infection (19). In contrast, monocytes from individuals with TBD exhibit impaired ex vivo metabolic function, characterised by reduced glycolysis and oxidative phosphorylation compared with healthy controls (20). These deficits are partially reversible with TB treatment, suggesting that Mtb infection directly alters monocyte bioenergetics and function.

Together, these findings highlight the importance of immunometabolism in shaping host responses

to TB and support ongoing efforts to develop host-directed therapies. However, how monocyte

metabolism and function differ between stages of TB, particularly between TBI and TBD, remains

incompletely understood. In this study, we aimed to characterise the phenotypic, functional, and

metabolic profiles of circulating monocytes from individuals across the TB spectrum. We used

SCENITH™, a single-cell metabolic profiling assay, to uncover immune and bioenergetic features

that may distinguish infection from disease(21) (Figure 1).

Methods

Human Participants

All peripheral blood patient samples were collected in the out-patient department of St James's

Hospital Dublin when patients were attending routine appointments (Table 1). Written informed

consent was obtained prior to collection. Exclusion criteria included age under 18 years, inability

to provide written informed consent, multi-drug-resistant tuberculosis, or a known (or ensuing)

diagnosis of malignancy, HIV or any other infection. Individuals with TBI were selected based on

a positive QuantiFERON-TB Gold Plus IGRA test with absence of any clinical symptoms of TB

disease and were subdivided into those pre-treatment (TBI/-; n=6) and those on-treatment (TBI/+;

n=4). Individuals clinically diagnosed of TBD were grouped into intensive (TBD/I, <2 months

treatment of 3-4 drugs; n=4) and continuation (TBD/C, >2 months reduced to 1-2 drugs; n=4)

treatment phases. Peripheral blood samples were also obtained from healthy controls who have

all received BCG vaccination previously.

 PBMC and Monocyte Isolations

PBMC were isolated from peripheral blood samples by density centrifugation over Lymphoprep

(StemCell Technologies). Monocytes were separated by hyperosmotic density gradient

centrifugation by layering PBMC on top of a fresh hyperosmotic Percoll solution (48.5% Percoll,

41.5% sterile H2O, 0.16 M NaCl) and centrifuged for 15 minutes at 580*g*. The interphase layer

was isolated, and cells were washed with cold PBS and resuspended at a concentration of 1 ×

10^6 cells/mL. Cells were plastic adhered for 1hr in serum-free medium and washed twice with PBS

to remove any non-adherent cells.

Irradiated H37Rv (iH37Rv; gifted by BEI Resources) stocks were thawed, sonicated for 15 mins

and syringed three times through a 25G needle break up clumps. iH37Rv was centrifuged at 30

xg for 5 mins to pellet any further clumps. Supernatant was aspirated and the concentration of

iH37Rv was calculated using Pierce™ BCA Protein Assay with Albumin protein as a control

(Thermo Scientific™). Cells were stimulated with 10 µg/mL of iH37Rv overnight for 18hours. Cells

were left unstimulated as a control.

Cytokine and Chemokine Analyses

A Human Luminex Discovery Kit (Cat LXSAHM-25) was designed to assess the following analytes

in monocyte supernatants: CCL1/I-309/TCA-3 (BR15), CCL2/JE/MCP-1 (BR25), CCL3/MIP-1

alpha (BR35), CCL4/MIP-1 beta (BR37), CXCL1/GRO alpha/KC/CINC-1 (BR77), CXCL2/GRO

beta/MIP-2/CINC-3 (BR27), CXCL9/MIG (BR52), CXCL10/IP-10/CRG-2 (BR67), G-CSF (BR54),

GM-CSF (BR46), Granzyme B (BR57), IFN-alpha (BR63), IFN-beta (BR21), IFN-gamma (BR29),

IL-1 alpha/IL-1F1 (BR38), IL-1 beta/IL-1F2 (BR28), IL-1ra/IL-1F3 (BR30), IL-6 (BR13), IL-

8/CXCL8 (BR18), IL-10 (BR22), IL-12 p70 (BR56), IL-17/IL-17A (BR42), IL-18/IL-1F4 (BR78), IL-

23 (BR76), TNF-alpha (BR12). The assay was carried out as per manufacturers protocol and

analysed on Luminex MAGPIX System. Unstimulated and Mtb-stimulated cells from each donor

were analyzed concurrently on the same plate, thereby eliminating potential inter-assay variability

and ensuring the accurate calculation of fold-change values. Analyte concentrations were

determined based on standards supplied in the kit, with a 5 parameter logistic regression (5-PL)

curve used on xPONENT(TM) software.

Flow cytometry

For phenotypical analyses PBMC were resuspended in PBS (Gibco) and stained immediately.

Anti-CD86 BV421 (1:100, IT2.2), anti-CD45RA BV711 (1:100, HI100), anti-CD40 APCCy7 (1:100,

5C3), anti-HLADR APC (1:100, L243), anti-CD14 AlexaFlour 488 (1:100, HCD14), anti-CD16

PerCpCy5.5 (1:100, 3G8), anti-CD80 PeCy7 (1:100, 2D10), anti-TNFRI PE (1:50, W15099A), and

anti-TNFRII PeDazzle594 (1:100, 3G7A02) were purchased from BioLegend and added to the

cells for 10 mins at RT. Cells were washed, fixed and acquired or stored at 4°C until acquisition.

Cultured PBMC were resuspended in 100 µl of FcR block (1µl/test; Miltenyi Biotec) and

ZombieNIR stain (BioLegend) in 1X PBS (Gibco). Cells were then surface stained with anti-CD14

and permeabilized with Foxp3 staining kit (eBioscience) according to the manufacturer's

instructions. Cells were incubated for 30 mins at 4°C with anti-puromycin (1:100, 2A4; BioLegend).

Cells were acquired on a FACS Canto (BD Biosciences), BD Fortessa, (BD Biosciences) or an

Amnis CellStream (Luminex Corporation, Austin, TX, USA). All data analysis was performed using

FlowJo software (BD Biosciences).

SCENITH (Single Cell mEtabolism by profiling Translation inHibition)

This protocol uses puromycin as a surrogate for protein synthesis, as it becomes incorporated into

newly formed proteins. Since protein synthesis represents around 50% of total cellular metabolic

activity, measuring puromycin incorporation provides a robust indicator of overall metabolic

activity. By measuring puromycin incorporation post-oligomycin treatment we can estimate the

cells dependency on mitochondrial respiration and their capacity for glycolysis. By measuring

puromycin incorporation post 2-Deoxy-D-Glucose (2-DG) treatment we can estimate the cells

dependency on glucose as a fuel and therefore their capacity for other fuels such as fatty acid and

amino acid oxidation. A detailed explanation of these calculations was described in our previous

work and in the original protocol publication (21,22). Post 18 hours of culture, SCENITH metabolic

function profiling was carried out on PBMCs. Cells were treated for 40 mins at 37°C, 5% CO2 with

DMSO Control (Co, 0.02% DMSO), 2-DG (100 mM; Sigma-Aldrich), Oligomycin (O; 1µM; Sigma-

Aldrich), a combination of both drugs (DGO), or with Harringtonine, which inhibits protein synthesis

(Sigma-Aldrich; 2 µg/ml). Puromycin (Puro; 10 µg/mL; Sigma-Aldrich) was added for the final 35

mins. Cells were washed in PBS and flow cytometry staining was carried out. The following

calculations were used:

Co = GeoMFI of anti-Puro-Fluorochrome upon DMSO Control treatment.

DG = GeoMFI of anti-Puro-Fluorochrome upon 2-Deoxy-D-Glucose treatment.

O = GeoMFI of anti-Puro-Fluorochrome upon Oligomycin A treatment.

DGO = GeoMFI of anti-Puro-Fluorochrome upon DG+O treatment.

Mitochondrial Dependence (MD)=100(Co-O)/(Co-DGO)

Glycolytic Capacity (GC)= 100-100(Co-O)/(Co-DGO)

Fatty Acid and Amino Acid Oxidation Capacity (FAOC)= 100-(100(Co-DG)/(Co-DGO)

Glucose Dependence (GD)= 100(Co-DG)/(Co-DGO)

Therefore, GC is the inverse data of MD (i.e. GC= 100- MD) and FAOC is the inverse data of GD

(i.e. FAOC= 100-GD). Patient samples with sufficient cell numbers underwent metabolic profiling.

No data was omitted.

Statistics

Statistical analysis was performed using GraphPad Prism version 10. The statistical test used is

indicated in each figure legend. A P-value of <0.05 was considered significant.

Study Approval

Our study protocol was approved by the Joint Research Ethics Committee of St. James's and

Tallaght Hospitals and by Trinity College Dublin. Written informed consent was received from

participants prior to inclusion in the study.

Data availability

Data is available directly from the corresponding author.

Results

Altered Expression of HLA-DR, CD14, and TNFRs in Monocytes Across the TB Spectrum

We next investigated whether TB infection influences monocyte frequency and phenotype across

the disease spectrum. No significant differences were observed in the frequencies of classical,

intermediate, or non-classical monocyte subsets between groups (Figure 2A–B). However,

phenotypic analysis revealed elevated expression intensity of CD14 and CD45RA on classical

monocytes from individuals with both TBI (CD14: P = 0.0039; CD45RA: P = 0.0070) and TBD

(CD14: P = 0.0010; CD45RA: P = 0.0007) compared with healthy controls (HC) (Figure 2C–D).

In contrast, expression of HLA-DR was significantly reduced on classical monocytes from TBI (P

= 0.0374) and even further reduced in TBD (P < 0.0001) compared with HC, with significantly

lower levels in TBD than TBI (P = 0.0068; Figure 2F). Expression of TNF receptors was decreased

in TBI but not TBD; both TNFRI (P = 0.0185) and TNFRII (P = 0.0414) were downregulated on

classical monocytes from TBI compared with HC (Figure 2G–H). Similar phenotypic changes were

observed in intermediate and non-classical monocyte subsets across the TB spectrum

(Supplemental Figure 2).

These data demonstrate that while monocyte subset frequencies remain stable, TB infection—

both latent and active—induces distinct phenotypic reprogramming of monocytes. Notably, the

progressive reduction in HLA-DR expression from TBI to TBD may reflect functional impairment,

with implications for antigen presentation and immune regulation during infection (23–27).

Monocytes from TBI and TBD individuals exhibit distinct functional responses to Mtb

stimulation.

To assess functional differences in monocytes across the TB spectrum, CD14$^+$ monocytes were

isolated from PBMC and cultured for 18 hours with or without irradiated-H37Rv (iMtb). Culture

supernatants were collected, and cytokine and chemokine concentrations were measured using

a custom Luminex multiplex bead-based immunoassay (Figure 3).

Unstimulated monocytes from individuals with latent TB infection (TBI, light grey) and active TB

disease (TBD, dark grey) displayed cytokine/chemokine profiles that were distinct from healthy

controls (HC, blue) (Figure 3A, C and Supplemental Figure 1A). Compared with TBD monocytes,

unstimulated monocytes from HC produced significantly higher levels of IL-1β (P=0.0061), GM-

CSF (P=0.0265), TNF-α (P=0.0237), and CXCL2 (P=0.0191), and lower levels of IL-17A

(P=0.0098) and CXCL9 (P=0.0353) (Figure 3A, C). Similarly, HC monocytes produced higher

levels of IL-1β (P=0.0053) and CXCL2 (P=0.0191), and lower levels of IL-17A (P=0.0018) than

monocytes from TBI individuals.

Following iMtb stimulation, monocytes from both TBI and TBD groups showed significantly

increased production of several cytokines and chemokines compared with HC, including IL-1α

(TBI: P=0.0153; TBD: P=0.0228), IL-8 (TBI: P=0.0129; TBD: P=0.0278), IL-23 (TBI: P=0.0158;

TBD: P=0.0400), and GM-CSF (TBI: P=0.0346; TBD: P=0.0203) (Figure 3B, D; Supplemental

Figure 1). Notably, despite comparable baseline cytokine production, only monocytes from TBI

individuals showed significantly increased levels of IL-1RA (P=0.0084), IL-17A (P=0.0171), IL-18

(P=0.0438), IFN-γ (P=0.0382), CXCL1 (P=0.0194), CXCL9 (P=0.0129), and CCL2 (P=0.0420)

compared with HC (Figure 3B, E). These findings demonstrate that while both TBI and TBD

monocytes respond robustly to iMtb stimulation, monocytes from individuals with TBI exhibit a

broader and more distinct cytokine and chemokine response compared with healthy controls. This

suggests that TBI monocytes may be functionally distinct from TBD monocytes, enabling a

heightened response to iMtb. In contrast, the blunted baseline and narrower inducible responses

observed in TBD may reflect monocyte dysfunction or exhaustion associated with active disease.

Reduced mitochondrial metabolism in monocytes from TBI and TBD individuals

We next compared the metabolic function of monocytes from HC, TBI, and TBD individuals using

the SCENITH assay, which measures puromycin incorporation as a proxy for protein synthesis, a

process that accounts for approximately 50% of a cell's total metabolic activity (Figure 4A). The

addition of metabolic inhibitors 2-DG and oligomycin enabled us to evaluate cellular dependence

on glycolysis and mitochondrial respiration, respectively.

In unstimulated conditions, monocyte metabolic activity (measured by puromycin mean

fluorescence intensity [MFI]) was reduced following treatment with either 2-DG or oligomycin

across all groups (Figure 4B), confirming sensitivity to both glycolytic and oxidative metabolic

pathways.

Glucose dependency, as calculated by SCENITH, was comparably high in monocytes from HC,

TBI, and TBD individuals (Figure 4C), indicating a shared reliance on glucose to fuel glycolysis

under basal conditions. In contrast, mitochondrial dependency was significantly reduced in

monocytes from both TBI ($P = 0.01$) and TBD ($P = 0.0078$) individuals compared with HC (Figure

4D), suggesting a metabolic shift away from oxidative phosphorylation. This reduction in

mitochondrial dependence is consistent with a compensatory increase in glycolytic capacity

(Figure 4E).

TBI monocytes exhibit enhanced glycolytic response to Mtb

Given the distinct clinical and immunological profiles of TBI and TBD, we hypothesized that

monocyte metabolic responses to Mtb would differ between TBI and TBD individuals. Following

iMtb stimulation, treatment with 2-DG significantly reduced monocyte metabolic activity in all

groups, confirming a strong dependence on glucose to fuel the activation-induced response

(Figure 4F–G).

In contrast, oligomycin treatment had no significant impact on metabolic activity post-stimulation

(Figure 4F), and calculated mitochondrial dependency remained low across all groups (Figure

4H). Notably, monocytes from TBI individuals exhibited significantly lower mitochondrial

dependence and higher glycolytic capacity compared with those from TBD individuals (P = 0.031),

indicating a more flexible metabolic profile.

Collectively, these data reveal a clear divergence in metabolic responsiveness to Mtb between

TBI and TBD monocytes. Monocytes from TBI individuals display enhanced glycolytic capacity

and reduced reliance on mitochondrial metabolism, both at baseline and in response to

stimulation; hallmarks of a metabolically reprogrammed, activated phenotype. In contrast, the

limited metabolic adaptability observed in monocytes from TBD individuals may reflect impaired

functional reprogramming during active disease.

Treatment restores monocyte mitochondrial function toward a healthy baseline

To determine whether treatment influenced monocyte phenotype or metabolic function, we further

stratified participants based on treatment status (Figure 5). Individuals with TBI were grouped as

pre-treatment (TBI/-) or on-treatment (TBI/+), and those with TBD were classified into intensive-

phase treatment (TBD/I, <2 months, multi-drug) or continuation-phase treatment (TBD/C, >2

months, reduced-drug) cohorts. CD14 expression is significantly elevated in TBI/+ (P=0.0228) and

TBD/I (P=0.0069) individuals (Figure 5A). CD45RA expression levels were significantly increased

in TBI/+ (P=0.0208), TBD/I (P=0.0310), and TBD/C (P=0.0078; Figure 5B). Of note, HLA-DR

expression is only significantly decreased in TBD/I (P=0.0069) and TBD/+ (P=0.0023) individuals

(Figure 5D), and TNFRI expression is only reduced in TBI/+ individuals (P=0.0162; Figure 5E).

Assessment of their metabolic activity revealed no differences in baseline protein synthesis

between these subgroups (Figure 5G). However, monocytes from TBI/- individuals showed

significantly reduced mitochondrial dependence compared with HC, while those from TBI/+ did

not (P = 0.0258; Figure 5H). Similarly, TBD/I monocytes displayed reduced mitochondrial

dependence compared with HC (P = 0.0198), whereas mitochondrial dependence in TBD/C

monocytes was not significantly different from HC (Figure 5H).

Upon iMtb stimulation, only monocytes from TBI individuals significantly increased their overall

metabolic activity (TBI/−: P = 0.0137; TBI/+: P = 0.0024; Figure 5I). Again, a corresponding

decrease in mitochondrial dependence was only observed in TBI individuals on treatment

(P=0.0185) and HC (P=0.0044; Figure 5J). In contrast, monocytes from TBD individuals showed

no significant change in either metric following iMtb stimulation, regardless of treatment phase

(Figure 5I-J).

These data suggest that both TBI and TBD are associated with reduced mitochondrial metabolism

in circulating monocytes, particularly in untreated individuals, and that both TB preventative

treatment (TPT) and TBD treatment may partially restore metabolic function toward a healthy

baseline.

Discussion

Our study reveals that Mtb infection induces distinct alterations in circulating monocyte phenotype,

function, and metabolism, that differ between individuals with TBI and TBD. Despite similar

monocyte subset frequencies across groups, phenotypic profiling showed that monocytes in TBI

individuals exhibit an upregulation of activation markers (e.g., CD14, CD45RA) alongside

downregulation of HLA-DR and TNF receptors, key mediators of antigen presentation and

inflammation. Notably, HLA-DR expression declined progressively from TBI to TBD, suggesting a

trajectory of monocyte dysfunction with disease progression which has been highlighted as a

biomarker of infection in TB, sepsis and paediatric infection (24–28). Functionally, monocytes from

both groups remained responsive to iMtb stimulation, but TBI monocytes exhibited a more robust

cytokine and metabolic response compared with TBD monocytes.

Among the analytes elevated in TBI compared with TBD and healthy controls, IL-1RA, IL-17A,

and CXCL9 showed the most pronounced increase, unaffected by outlier values. IL-1RA, an

antagonist of IL-1α/β signalling produced predominantly by monocytes and neutrophils, has been

proposed as a biomarker for TB spectrum stratification (29). In our study, IL-1RA was significantly

upregulated in monocytes from individuals with TBI, but not from those with TBD, supporting a

potential role for monocyte-derived IL-1RA in maintaining infection containment. However,

previous findings on IL-1RA expression in TB have been inconsistent, with some studies reporting

higher IL-1RA levels in response to TB antigens in TBD relative to TBI, and others showing the

opposite (30–33). Notably, these prior studies were based on whole blood or PBMC cultures,

making it difficult to resolve cell-specific contributions. By focusing on purified monocytes, our data

provide more direct evidence that monocyte-specific IL-1RA production may differentiate TBI from

TBD.

Similarly, we observed significant upregulation of IL-17A and CXCL9 in monocytes from TBI but

not TBD individuals (Figure 1E). CXCL9 acts as a critical chemoattractant for CXCR3[+] T cells,

facilitating the recruitment and activation of IFN-γ–producing effector T cells (34). IL-17A is a

potent inducer of chemokines that recruit immune cells such as T cells and neutrophils, and

stimulates antimicrobial peptide production from epithelial cells (35). Collectively, this monocyte

cytokine and chemokine profile characterized by elevated IL-1RA, IL-17A, and CXCL9 in TBI may

reflect an immunological environment conducive to effective containment of Mtb infection.

We hypothesize that the distinct effector cytokine and chemokine profile observed in monocytes from individuals with TBI is underpinned by metabolic reprogramming that favours glycolysis over mitochondrial respiration. TBI monocytes displayed enhanced glycolytic capacity and reduced mitochondrial dependence both at rest and upon stimulation, consistent with an activated immunometabolic phenotype (Figure 4D-E). In contrast, TBD monocytes failed to undergo this shift, suggesting impaired metabolic plasticity which supports previous reports of their overall diminished bioenergetics (20). Given the known link between glycolysis and effector function, this metabolic rewiring is likely associated with the enhanced production of cytokines and chemokine observed specifically in TBI monocytes, which may facilitate the effective immune containment of Mtb. Of note, to assess differential sustained metabolic responses of monocytes to Mtb between TBI and TBD individuals, cells were cultured for 18 hours with iMtb followed by the SCENITH protocol. We acknowledge that future studies incorporating Seahorse Analyzer assays on freshly isolated monocytes may provide further insight into the dynamic and immediate metabolic changes that occur upon iMtb stimulation.

These data resonate with and extend the work of other research groups investigating host metabolic responses in TB. Notably, bioenergetic dysfunction of monocytes in South African TBD cohorts have been reported (20). In addition, myeloid cell metabolic rewiring governs innate immune cell function in the context of BCG vaccination (36) and heterologous innate immunity to Mtb elicited by adenoviral vectored COVID-19 vaccination (37). Our findings complement and build upon these studies by focusing specifically on purified circulating monocytes metabolic capacity and effector cytokine output across the TB spectrum.

Importantly, our findings also indicate that treatment, including TPT in TBI and multidrug regimens

in TBD, partially restores mitochondrial function in circulating monocytes, shifting their metabolic

baseline profile closer to that of healthy individuals. This suggests that monocyte metabolism is

dynamic and responsive to therapeutic intervention, indicating that mitochondrial dependency

metrics hold potential as a novel biomarker for treatment efficacy and disease resolution.

Moreover, these insights underscore the therapeutic potential of targeting monocyte metabolic

pathways to enhance host immune function in TB.

These results contribute to a growing body of evidence that immunometabolism shapes infection

outcomes in TB. Since circulating monocytes give rise to macrophages, including lung-resident

alveolar macrophages, targeting metabolic pathways in monocytes may enhance both peripheral

and pulmonary immune responses (38–40). Although our study focused on peripheral blood

monocytes, it is reasonable to hypothesize that some aspects of monocyte immunometabolic

reprogramming are reflected in lung-resident or recruited mononuclear phagocytes, given the

dynamic trafficking of monocytes from blood to tissue. However, the lung microenvironment may

also induce additional, tissue-specific metabolic adaptations that could modify these peripheral

phenotypes. Recent evidence indicates that human airway macrophages are metabolically and

functionally distinct from monocyte-derived macrophages in response to Mtb (41,42) and murine

studies highlight the metabolic heterogeneity across alveolar and interstitial lung macrophages in

response to Mtb (18). If future studies reveal differences between systemic and local immune

programming, this will underscore compartmentalized immune responses, helping to delineate

distinct mechanisms of immune protection or pathogenesis in TB. Importantly, such insights

reinforce the potential of monocyte metabolism as a promising target for host-directed therapies,

particularly those aimed at preventing progression from TBI to TBD by enhancing both systemic

and lung-localized immunity.

While our study provides novel insights into monocyte immunometabolism across the TB

spectrum, several limitations should be acknowledged. First, the relatively small sample size may

limit the generalizability of our findings and the statistical power to detect more subtle differences

or correlations with clinical outcomes. Second, we did not establish a direct link between the

observed monocyte metabolic and phenotypic changes and disease progression or treatment

success, which will require longitudinal studies with larger cohorts. Third, our functional assays

utilized irradiated H37Rv. This well-characterized lab strain facilitates standardization however,

future studies incorporating live, virulent strains are essential to fully elucidate how pathogen

variability shapes host monocyte responses. Despite these limitations, our data provide a valuable

foundation for future investigations aimed at elucidating the functional benefits of this metabolic

plasticity in TBI, and its impact on bacterial killing. Such studies will be essential for identifying

robust biomarkers and therapeutic targets through detailed mechanistic analyses of monocyte

immunometabolism in tuberculosis.

In summary, our data reveal that Mtb infection drives distinct immunometabolic changes in

circulating monocytes with TBI characterised by functional responsiveness and glycolytic, pro-

inflammatory profile, while TBD is marked by impaired metabolic plasticity. Furthermore, we show

that TB treatment of both TBI and TBD individuals may restore monocyte mitochondrial function.

These findings offer important insight into the cellular mechanisms of immune containment in TB

and support the rationale for treating TBI. Moreover, they underscore the potential of monocyte

metabolism as a biomarker of disease state and a tractable target for host-directed therapies.

Continued research building on these insights will be crucial for advancing TB diagnosis,

monitoring treatment efficacy, and developing novel interventions to improve patient outcomes.

Acknowledgements

We sincerely thank all the individuals who participated in this study. We are especially grateful to

the patients for their time, generosity, and willingness to contribute to research aimed at improving

the understanding and treatment of tuberculosis. Their participation made this work possible. The

following reagent was obtained through BEI Resources, NIAID, NIH: Mycobacterium tuberculosis,

Strain H37Rv, Gamma-Irradiated Whole Cells, NR-49098. We gratefully acknowledge the support

of the Core Flow Cytometry Facility and Prof. Nollaig Bourke's research team at the Trinity

Translational Medicine Institute.

References

1.   Geneva WHO. *Global Tuberculosis Report 2024.*; 2024.

2.   Andrews JR, Noubary F, Walensky RP, Cerda R, Losina E, Horsburgh CR. Risk of

Progression to Active Tuberculosis Following Reinfection With Mycobacterium

tuberculosis. *Clin Infect Dis An Off Publ Infect Dis Soc Am*. 2012;54(6):784.

doi:10.1093/CID/CIR951

3.   Sampath P, Moideen K, Ranganathan UD, Bethunaickan R. Monocyte Subsets:

Phenotypes and Function in Tuberculosis Infection. *Front Immunol*. 2018;9:1726.

doi:10.3389/FIMMU.2018.01726

4.   Wolf Y, Shemer A, Polonsky M, et al. Autonomous TNF is critical for in vivo monocyte

survival in steady state and inflammation. *J Exp Med*. 2017;214(4):905.

doi:10.1084/JEM.20160499

5.   Netea MG, Nold-Petry CA, Nold MF, et al. Differential requirement for the activation of the

inflammasome for processing and release of IL-1beta in monocytes and macrophages.

Blood. 2009;113(10):2324-2335. doi:10.1182/BLOOD-2008-03-146720

6.   Affandi AJ, Olesek K, Grabowska J, et al. CD169 Defines Activated CD14+ Monocytes

With Enhanced CD8+ T Cell Activation Capacity. *Front Immunol*. 2021;12:697840.

doi:10.3389/FIMMU.2021.697840/FULL

7.   Guilliams M, Mildner A, Yona S. Developmental and Functional Heterogeneity of

Monocytes. *Immunity*. 2018;49(4):595-613. doi:10.1016/J.IMMUNI.2018.10.005

8.   Serbina N V., Pamer EG. Monocyte emigration from bone marrow during bacterial

infection requires signals mediated by chemokine receptor CCR2. *Nat Immunol*.

2006;7(3):311-317. doi:10.1038/NI1309

9.   McQuattie-Pimentel AC, Budinger GRS, Ballinger MN. Monocyte-derived Alveolar

Macrophages: The Dark Side of Lung Repair? *Am J Respir Cell Mol Biol*. 2018;58(1):5-6.

doi:10.1165/RCMB.2017-0328ED

10.   Misharin A V., Morales-Nebreda L, Reyfman PA, et al. Monocyte-derived alveolar

macrophages drive lung fibrosis and persist in the lung over the life span. *J Exp Med*.

2017;214(8):2387-2404. doi:10.1084/JEM.20162152

11.   Machiels B, Dourcy M, Xiao X, et al. A gammaherpesvirus provides protection against

allergic asthma by inducing the replacement of resident alveolar macrophages with

regulatory monocytes. *Nat Immunol*. 2017;18(12):1310-1320. doi:10.1038/NI.3857

12.   Aegerter H, Kulikauskaite J, Crotta S, et al. Influenza-induced monocyte-derived alveolar

macrophages confer prolonged antibacterial protection. *Nat Immunol 2020 212*.

2020;21(2):145-157. doi:10.1038/s41590-019-0568-x

13.   Srivastava S, Ernst JD, Desvignes L. Beyond macrophages: the diversity of mononuclear

cells in tuberculosis. *Immunol Rev*. 2014;262(1):179. doi:10.1111/IMR.12217

14.   Cohen SB, Gern BH, Delahaye JL, et al. Alveolar Macrophages Provide an Early

Mycobacterium tuberculosis Niche and Initiate Dissemination. *Cell Host Microbe*.

2018;24(3):439-446.e4. doi:10.1016/J.CHOM.2018.08.001

15.   Gleeson LE, Sheedy FJ, Palsson-McDermott EM, et al. Cutting Edge: Mycobacterium

tuberculosis Induces Aerobic Glycolysis in Human Alveolar Macrophages That Is

Required for Control of Intracellular Bacillary Replication . *J Immunol*. 2016;196(6):2444-

2449. doi:10.4049/jimmunol.1501612

16.   Mendonca LE, Pernet E, Khan N, et al. Human alveolar macrophage metabolism is

compromised during Mycobacterium tuberculosis infection. *Front Immunol*. 2023;13.

doi:10.3389/FIMMU.2022.1044592/PDF

17.   Cumming BM, Addicott KW, Adamson JH, Steyn AJC. Mycobacterium tuberculosis

induces decelerated bioenergetic metabolism in human macrophages. *Elife*. 2018;7.

doi:10.7554/eLife.39169

18.   Huang L, Nazarova E V., Tan S, Liu Y, Russell DG. Growth of Mycobacterium

464   tuberculosis in vivo segregates with host macrophage metabolism and ontogeny. *J Exp*

465   *Med*. 2018;215(4):1135-1152. doi:10.1084/JEM.20172020,

466   19.   Lachmandas E, Beigier-Bompadre M, Cheng SC, et al. Rewiring cellular metabolism via

467   the AKT/mTOR pathway contributes to host defence against Mycobacterium tuberculosis

468   in human and murine cells. *Eur J Immunol*. 2016;46(11):2574-2586.

469   doi:10.1002/EJI.201546259;PAGE:STRING:ARTICLE/CHAPTER

470   20.   Cumming BM, Addicott KW, Maruri F, et al. Longitudinal mitochondrial bioenergetic

471   signatures of blood monocytes and lymphocytes improve during treatment of drug-

472   susceptible pulmonary tuberculosis patients Monocyte/lymphocyte bioenergetic

473   signatures post-TB treatment. *Front Immunol*. 2024;15:1465448.

474   doi:10.3389/fimmu.2024.1465448

475   21.   Argüello RJ, Combes AJ, Char R, et al. SCENITH: A Flow Cytometry-Based Method to

476   Functionally Profile Energy Metabolism with Single-Cell Resolution. *Cell Metab*.

477   2020;32(6):1063-1075.e7. doi:10.1016/J.CMET.2020.11.007

478   22.   Jameson G, Walsh A, Woods R, et al. Human tissue-resident NK cells in the lung have a

479   higher glycolytic capacity than non-tissue-resident NK cells in the lung and blood. *Proc*

480   *Natl Acad Sci*. 2024;121(42):2412489121. doi:10.1073/PNAS.2412489121

481   23.   Ansari MA, Das S, Rai G, et al. Low monocytic HLA-DR expression in critically ill patients

482   of sepsis: An indicator for antimicrobial and/or immunomodulatory intervention. *Transpl*

483   *Immunol*. 2023;81. doi:10.1016/j.trim.2023.101942

484   24.   Quadrini KJ, Patti-Diaz L, Maghsoudlou J, Cuomo J, Hedrick MN, McCloskey TW. A flow

485   cytometric assay for HLA-DR expression on monocytes validated as a biomarker for

486   enrollment in sepsis clinical trials. *Cytom Part B - Clin Cytom*. 2021;100(1):103-114.

487   doi:10.1002/CYTO.B.21987;CTYPE:STRING:JOURNAL

25.   Fumeaux T, Pugin J. Role of interleukin-10 in the intracellular sequestration of human

leukocyte antigen-DR in monocytes during septic shock. *Am J Respir Crit Care Med*.

2002;166(11):1475-1482. doi:10.1164/RCCM.200203-

217OC;JOURNAL:JOURNAL:AJRCCM;WGROUP:STRING:DEFAULT

26.   Brandes-Leibovitz R, Riza A, Yankovitz G, et al. Sepsis pathogenesis and outcome are

shaped by the balance between the transcriptional states of systemic inflammation and

antimicrobial response. *Cell Reports Med*. 2024;5(11). doi:10.1016/j.xcrm.2024.101829

27.   Leventogiannis K, Kyriazopoulou E, Antonakos N, et al. Toward personalized

immunotherapy in sepsis: The PROVIDE randomized clinical trial. *Cell Reports Med*.

2022;3(11). doi:10.1016/j.xcrm.2022.100817

28.   Juskewitch JE, Abraham RS, League SC, et al. Monocyte HLA-DR expression and

neutrophil CD64 expression as biomarkers of infection in critically ill neonates and infants.

Pediatr Res. 2015;78(6):683-690.

doi:10.1038/PR.2015.164;SUBJMETA=2421,250,255,53,631,692;KWRD=DIAGNOSTIC+

MARKERS,INFECTIOUS+DISEASES

29.   Chegou NN, Heyckendorf J, Walzl G, Lange C, Ruhwald M. Beyond the IFN-γ horizon:

Biomarkers for immunodiagnosis of infection with Mycobacterium tuberculosis. *Eur Respir*

J. 2014;43(5):1472-1486. doi:10.1183/09031936.00151413,

30.   Wergeland I, Assmus J, Dyrhol-Riise AM. Cytokine Patterns in Tuberculosis Infection; IL-

1ra, IL-2 and IP-10 Differentiate Borderline QuantiFERON-TB Samples from Uninfected

Controls. *PLoS One*. 2016;11(9):e0163848. doi:10.1371/JOURNAL.PONE.0163848

31.   Sanchez C, Jaramillo-Valverde L, Capristano S, et al. Antigen-Induced IL-1RA Production

Discriminates Active and Latent Tuberculosis Infection. *Microorganisms*. 2023;11(6).

doi:10.3390/MICROORGANISMS11061385,

32.   Akashi S, Suzukawa M, Takeda K, et al. IL-1RA in the supernatant of QuantiFERON-TB

513  Gold In-Tube and QuantiFERON-TB Gold Plus is useful for discriminating active

514  tuberculosis from latent infection. *J Infect Chemother*. 2021;27(4):617-624.

515  doi:10.1016/J.JIAC.2020.11.023

516 33. Llibre A, Smith N, Rouilly V, et al. Tuberculosis alters immune-metabolic pathways

517  resulting in perturbed IL-1 responses. *Front Immunol*. 2022;13:897193.

518  doi:10.3389/FIMMU.2022.897193/BIBTEX

519 34. Lugassy J, Abdala-Saleh N, Jarrous G, et al. Development of DPP-4-resistant CXCL9-Fc

520  and CXCL10-Fc chemokines for effective cancer immunotherapy. *Proc Natl Acad Sci U S*

521  *A*. 2025;122(16):e2501791122.

522  doi:10.1073/PNAS.2501791122/SUPPL_FILE/PNAS.2501791122.SAPP.PDF

523 35. Lindén A, Laan M, Anderson GP. Neutrophils, interleukin-17A and lung disease. *Eur*

524  *Respir J*. 2005;25(1):159-172. doi:10.1183/09031936.04.00032904

525 36. Arts RJW, Carvalho A, La Rocca C, et al. Immunometabolic Pathways in BCG-Induced

526  Trained Immunity. *Cell Rep*. 2016;17(10):2562-2571. doi:10.1016/j.celrep.2016.11.011

527 37. Murphy DM, Cox DJ, Connolly SA, et al. Trained immunity is induced in humans after

528  immunization with an adenoviral vector COVID-19 vaccine. *J Clin Invest*. 2023;133(2).

529  doi:10.1172/JCI162581

530 38. Suzuki H, Hisamatsu T, Chiba S, et al. Glycolytic pathway affects differentiation of human

531  monocytes to regulatory macrophages. *Immunol Lett*. 2016;176:18-27.

532  doi:10.1016/J.IMLET.2016.05.009

533 39. Russell DG, Simwela N V., Mattila JT, Flynn J, Mwandumba HC, Pisu D. How

534  macrophage heterogeneity affects tuberculosis disease and therapy. *Nat Rev Immunol*.

535  Published online January 7, 2025. doi:10.1038/S41577-024-01124-3

536 40. Niec RE, Rudensky AY, Fuchs E. Inflammatory adaptation in barrier tissues. *Cell*.

537  2021;184(13):3361-3375. doi:10.1016/J.CELL.2021.05.036

538   41.   Mendonca LE, Pernet E, Khan N, et al. Human alveolar macrophage metabolism is

539   compromised during Mycobacterium tuberculosis infection. *Front Immunol*. 2023;13.

540   doi:10.3389/fimmu.2022.1044592

541   42.   Cox DJ, Connolly SA, Maoldomhnaigh CÓ, et al. Human airway macrophages are

542   metabolically reprogrammed by IFN-γ resulting in glycolysis dependent functional

543   plasticity. *Elife*. 2024;13. doi:10.7554/ELIFE.98449.3

547   Figure 1. Schematic of study design for analysis of monocyte phenotype, function and

548   metabolism. Peripheral blood samples were collected from healthy controls (HC), people with

TBI and people with TBD and PBMC were isolated by density centrifugation. Isolated PBMCs

were stained for flow cytometry analysis of monocyte expression levels of CD14, CD45RA,

CD40, HLA-DR, TNFRI and TNFRII. Monocytes were isolated from PBMCs using density

centrifugation with a Percoll gradient followed by 1hr plastic adherence. Purified monocytes were

then cultured, with or without irradiated-H37Rv (iMtb), for 18 hours. The supernatants were

harvested and analysed for cytokine and chemokine concentrations. PBMC were plated for 24

hours with or without iMtb. Puromycin and metabolic inhibitors (2-DG and oligomycin) were

added for 40 mins and then the cells were stained for flow cytometry analysis. Cells were

washed and stained with fluorochrome-conjugated antibodies against CD14 and puromycin and

analyzed by flow cytometry. Puromycin MFI levels measure overall protein synthesis and act as

a surrogate for metabolic activity. Created in https://BioRender.com.

561   Figure 2. Distinct phenotype in monocytes from individuals with TBI. PBMC were isolated

from peripheral blood and stained for flow cytometry analysis. **A-B)** Representative flow plot of

gating strategy (A) and bar chart of monocyte subset frequencies as percentage of total

monocytes (B). **C-H)** Expression levels of CD14 (C), CD45RA (D) and CD40 (E) and HLA-DR

(F), TNFRI (G) and TNFRII (H) on classical monocytes across HC, TBI and TBD. P values

generated using Two-Way ANOVA with Tukey's multiple comparison test (B) and Kruskal Wallis

test with Dunn's multiple comparison (C-H), *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$ or

Mann Whitney test, ##, $P<0.01$ (F).

Figure 3. Monocytes from TBI individuals' distinct cytokine and chemokine production in response to Mtb. Monocytes were isolated from PBMC and cultured for 18 hours with and without irradiated H37Rv (iMtb). [Healthy controls (HC, n=5)), latent (TBI, n=8) and Active (TBD, n=5) TB patients]. **A-B)** Mean levels of cytokine and chemokine concentrations (pg/mL) in supernatants collected from unstimulated (A) and iH37Rv (iMtb) stimulated (B) monocyte cultures [HC (blue), TBI (light grey) and TBD (dark grey]. **C)** Differential concentration of cytokines and chemokines in unstimulated monocytes. **D-E)** Analytes that are significantly increased in both TBI and TBD individual's monocytes (D) and in only TBI individual's monocytes (E). P values calculated using Kruskal Wallis test and Dunn's multiple comparison test.

Figure 4. Monocyte metabolic response to Mtb is distinct in TBI. PBMC were isolated from peripheral blood and cultured with or without irradiated H37Rv (iMtb) 10ug/mL for 18 hr. Post-18hr PBMC cultures were treated with DMSO control or metabolic inhibitors 2-deoxyglucose (2DG), oligomycin (Oligo), or combinations of both, or harringtonine, which inhibits protein synthesis and acts as a negative control for this assay. Puromycin was then added, and cells were cultured for 40 min. Cells were washed and stained with fluorochrome-conjugated antibodies against CD14 and puromycin and analyzed by flow cytometry. Puromycin MFI levels measure overall protein synthesis and act as a surrogate for metabolic activity. **A)** Representative flow cytometry gating strategy showing the analyses of monocytes and their puromycin staining across metabolic treatments. **B)** Puromycin MFI values measuring overall metabolic activity in unstimulated monocytes across untreated (-, DMSO control), 2DG, and

oligomycin treatments, from HC (n=10) and people with TBI (n=10) and TBD (n=8). **C-E)**

Unstimulated monocytes percentage glucose (C) and mitochondrial (D) dependence and

glycolytic capacity (E) across all cohorts. **F)** Puromycin MFI values measuring overall metabolic

activity in iMtb-stimulated monocytes across untreated, 2DG, and oligomycin treatments from

HC (n=6) and people with TBI (n=8) and TBD (n=6). **G-I)** iMtb-stimulated monocytes percentage

glucose (G) and mitochondrial (H) dependence and glycolytic capacity (I) across all cohorts. P

values generated using Two-Way ANOVA with Dunnett's (B, F), Kruskal Wallis test with Dunn's

multiple comparison (D, E) and Mann Whitney test (H).

Figure 5. Treatment impact on monocyte phenotype and function.

PBMC were isolated from peripheral blood and analysed by flow cytometry or cultured with or

without irradiated H37Rv (iMtb) 10ug/mL for 18 hr. **A-F)** Expression levels of CD14 (A), CD45RA

(B) and CD40 (C) and HLA-DR (D), TNFRI (E) and TNFRII (F) on classical monocytes ex vivo.

Post-18hr PBMC cultures were treated with DMSO control or metabolic inhibitors 2-deoxyglucose

(2DG), oligomycin (Oligo), or combinations of both, or harringtonine, which inhibits protein

synthesis and acts as a negative control for this assay. Puromycin was then added, and cells were

cultured for 40 min. Cells were washed and stained with fluorochrome-conjugated antibodies

against CD14 and puromycin and analyzed by flow cytometry. Puromycin MFI levels measure

overall protein synthesis and act as a surrogate for metabolic activity. **G)** Puromycin MFI values

measuring overall metabolic activity in unstimulated, DMSO control-treated, monocytes in HC

(n=10), TBI pre-treatment (TBI/-, n=6) and TBI on-treatment (TBI/+, n=4) and TBD in the intensive

(TBD/I, n=4)) and continuation (TBD/C, n=4) treatment phase. **H)** Percent mitochondrial

dependence in unstimulated monocytes. **I)** Puromycin MFI values measuring overall change in

metabolic activity from unstimulated (white) to iH37Rv-stimulated (blue) monocytes. **J)** Change in

percentage mitochondrial dependence in iH37Rv (iMtb)-stimulated monocytes. P values

generated using Two-Way ANOVA with Sidak's multiple comparison test.

Supplemental Figure 1. Raw concentration values for monocyte cytokine and chemokine

production.

Monocytes were isolated from PBMC and cultured for 18 hrs with and without irradiated H37Rv

(iMtb). **A-B)** Concentration of cytokines and chemokines (pg/mL) in supernatants collected from

unstimulated (A) and iMtb-stimulated (B) monocyte cultures. P values generated using 2 Way

ANOVA with Tukey's multiple comparison test.

Supplemental Figure 2. TB infection alters intermediate and non-classical monocyte

phenotype.

PBMC were isolated from peripheral blood and stained for flow cytometry analysis. **A-B)** MFI

values for CD14, HLA-DR, CD45RA, CD40, TNFRI, and TNFRII expression on intermediate (A)

and non-classical (B) monocytes across HC, TBI and TBD. P values generated using Two-Way

ANOVA with Tukey's multiple comparison test.

	HC	TBI	TBD
Age	38+/-15	44.90+/-12.81	41.25+/-18.43
Gender	F (9) M (1)	M (11) F (10)	M (10) F (6)
Ethnicity (self-reported)	European (9) African (1)	Asian (10), European (4), African (2), Indian (2), Not reported (3)	Asian (4), European (5), African (7),
Site of Mtb infection	N/A	N/A	Pulmonary (6), Extra-pulmonary (7), Both (2)

641   Table 1: Patient information. HC=Healthy Controls, TBI= Tuberculosis Infection, TBD=

642   tuberculosis disease, F= female, M=male.

Figure 1

Figure 2

652 **Figure 3**

654 **Figure 4**

Figure 5

660   **Supplemental Figure 1**

A

Intermediate Monocytes

B

Non classical Monocytes

662   **Supplemental Figure 2**