Undecaprenyl Pyrophosphate Phosphatase (UppP) is a pivotal element in *Salmonella* Intramacrophage Survival

Rhea Vij[1], Debapriya Mukherjee[1#], Kirti Parmar[1#], Krishna Chaitanya Nallamotu[2,3], Manjula Reddy[2,3], Dipshikha Chakravortty[1,4*]

[1]Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore 560012, India.

[2]CSIR-Centre for Cellular and Molecular Biology, Hyderabad, Telangana 500007, India

[3]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, Uttar Pradesh, India

[4]Adjunct Faculty, Indian Institute of Science Research and Education, Thiruvananthapuram, Kerala, India

[#]Equal contribution

[*]Corresponding author: dipa@iisc.ac.in

Tel: 0091 80 2293 2842

Fax: 0091 80 2360 2697

Abstract

To establish infection, *Salmonella* confronts a dynamic barrage of host-induced stresses. The peptidoglycan layer, essential for maintaining cell integrity and counteracting environmental stress, relies on the lipid carrier undecaprenyl phosphate, generated by Undecaprenyl pyrophosphate phosphatase (UppP). While UppP is linked to virulence in other pathogens, its role in *Salmonella* remains unclear. We show that an *uppP* mutant in *S*. Typhimurium exhibits

altered cell morphology, reduced stiffness, and impaired survival in RAW 264.7

macrophages. The mutant is also attenuated in systemic infection in C57BL/6 mice. These

defects are associated with increased sensitivity to nitrosative stress. Notably, iNOS

inhibition or deficiency restores intracellular survival of the *uppP* mutant in both RAW 264.7

macrophages and the mouse model, implicating UppP in resistance to nitrosative stress. Our

findings reveal a critical role for UppP in promoting *Salmonella* survival within macrophages

and contributing to systemic pathogenesis.

Introduction

Salmonella is a Gram-negative foodborne pathogen belonging to the *Enterobacteriaceae*

family [1]. The genus, comprising more than 2600 serovars, can infect various hosts,

including plants, birds, and mammals[2], [3]. Clinical manifestations of *Salmonella* infection

range from self-limiting gastroenteritis associated with non-typhoidal serovars to systemic

infection caused by typhoidal serovars and invasive non-typhoidal serovars [4]. The

significant burden of *Salmonella*-associated enterocolitis, affecting 95.1 million people in

2017, coupled with the rising threat of drug-resistant strains as recognised by its inclusion on

the WHO priority pathogen list, underscores *Salmonella*'s growing danger to public health

[5], [6], [7].

Salmonella enters the host through contaminated food and water. *Salmonella* encounters

several stresses within the host, including oxidative and nitrosative stress, osmotic stress, acidic

pH, nutrient starvation, metal ions, and host antimicrobials [8]. Out of all these stresses, the

bacterial peptidoglycan layer helps counter osmotic stress by acting as a barrier to the external

environment and resisting internal turgor pressure, thereby preventing the cell from bursting.

It also helps in maintaining the bacterial shape and virulence. The synthesis of the

peptidoglycan layer is a concerted effort of many proteins and takes place in the cytosolic and

periplasmic compartments of the bacterial cell [9]. The biosynthesis of the peptidoglycan

building block, N-acetylglucosamine-N-acetyl muramyl-pentapeptide, takes place in the

cytoplasm on the lipid carrier undecaprenyl phosphate (C_{55}-P), forming Lipid II. Lipid II is

then flipped by MurJ to the periplasmic phase [9], [10]. In the periplasmic phase, the

disaccharide pentapeptide part of Lipid-II undergoes polymerisation by SEDS (Shape,

Elongation, Division, and Sporulation) proteins (such as FtsW and RodA) and Penicillin-

binding proteins (PBPs) for peptidoglycan synthesis, releasing undecaprenyl pyrophosphate

[11], [12], [13], [14]. The undecaprenyl pyrophosphate (C_{55}-PP) is dephosphorylated by

undecaprenyl pyrophosphate phosphatase (UppP) to undecaprenyl phosphate in the

periplasmic phase, which is recycled for another round of peptidoglycan synthesis [15]. While

UppP accounts for approximately 75% of this phosphatase activity in *E. coli*, the remaining

activity is provided by the functionally redundant PAP2 family proteins, PgpB, YbjG, and LpxT

[16], [17].

Certain studies have highlighted the crucial role of undecaprenyl pyrophosphate phosphatases

in the virulence and pathogenesis of different bacterial species. In *E. coli,* the undecaprenyl

pyrophosphate phosphatase encoded by the gene *uppP*, also known as *bacA*, leads to bacitracin

resistance when it is overexpressed. Bacitracin strongly binds to undecaprenyl pyrophosphate

and prevents its dephosphorylation [18]. Mutant strains lacking the *bacA* homologue in

Staphylococcus aureus and *Streptococcus pneumoniae* exhibited reduced virulence and severe

attenuation, respectively, in the mouse model of infection [19]. HupA, an undecaprenyl

pyrophosphate phosphatase of the PAP2 family, belonging to *Helicobacter pylori*, is essential

for colonisation of the stomach [20]. Other reports have shown that deletion of *uppP* in

Mycobacterium smegmatis impairs its ability to form biofilms [21]. However, the role of

undecaprenyl phosphate in *Salmonella* Typhimurium (STM) pathogenesis has remained

unexplored.

Recent studies suggest that intracellular *Salmonella* actively remodels its peptidoglycan layer

to survive in the intracellular niche, differing significantly from those grown in LB medium

[22], [23]. Intracellular *Salmonella* Typhimurium actively edits its peptidoglycan to incorporate

elevated levels of L-alanine-D-glutamic acid (L, D-) cross-links, replacing the terminal D-

alanine with the amino alcohol alaninol and cleaving γ-D-glutamyl-meso-diaminopimelic acid

(D, D-) crosslinks, which reduce the release of immunogenic muropeptides, thereby limiting

recognition by innate immune receptors, such as NOD1/2, and attenuating the inflammatory

response [24], [25]. Additionally, in *Salmonella* Typhi, the elevated levels of L, D-crosslinks

play a dual role by facilitating the secretion of the typhoid toxin [22]. UppP's crucial function

in peptidoglycan synthesis suggests that it may play a role in cell wall adaptation during stress,

thereby enhancing *Salmonella*'s survival. In this study, we generated an *uppP* knockout strain

of *S.* Typhimurium and evaluated its impact on bacterial survival and virulence. Our findings

demonstrate that UppP is essential for *S.* Typhimurium proliferation in RAW 264.7

macrophages and systemic dissemination in C57BL/6 mice by promoting resistance to host-

induced nitrosative stress.

Results

UppP affects bacterial cell dimensions and strength

To investigate the role of undecaprenyl pyrophosphate phosphatase in *Salmonella*

pathogenesis, we generated an *uppP*-deleted strain of *Salmonella* Typhimurium by a one-step

chromosomal gene inactivation method [26]. We observed that the *uppP* deleted strain

demonstrated no significant difference in growth in nutrient-rich LB broth (Fig. 1A), M9

minimal media (Fig. 1B) and acidic F-media (pH-5.4, similar to the pH of the *Salmonella*-

containing vacuole [SCV]) [27], [28] (Fig. 1C).

A continuous supply of undecaprenyl phosphate is crucial for normal cell shape maintenance

in *Escherichia coli*, and its depletion induces morphological defects [29], [30], [31]. To

dissect the contribution of UppP in bacterial cell morphology of *Salmonella*, we performed

Scanning Electron Microscopy (SEM) to examine the logarithmic (1:100 dilution) and

stationary-phase cultures of STM WT and STM Δ*uppP*. The quantitative analysis of cell

length revealed that STM Δ*uppP* (1.319 ± 0.3145 μm) exhibits reduced cell length in

comparison to STM WT (1.509 ± 0.5406 μm) during the log phase, which was restored upon

complementation of *uppP* (1.768 ± 0.4086 μm) (Fig. 1D-E). Whereas there is an increase in

cell length of STM Δ*uppP* (1.236 ± 0.2295 μm) in comparison to STM WT (1.067 ± 0.2360

μm) during the stationary phase (S1A-B). Interestingly, when STM Δ*uppP* cells were

subcultured at a higher dilution of 1:1000, allowing them to grow for a longer period in the

logarithmic phase, we observed that the mutant (1.963 ± 0.5102 μm) became notably more

elongated compared to the wild type (1.642 ± 0.4547 μm). Additionally, cell size returned to

normal upon complementation (1.704 ± 0.4999 μm) (Fig. S1C-D).

Furthermore, we performed atomic force microscopy (AFM) to evaluate the stiffness of the

bacterial cell envelope. Young's modulus, quantified from AFM micrographs, was reduced in

the STM Δ*uppP* (39.55 ± 23.26 MPa) strain compared to the wild type (137.5 ± 115.5 MPa),

and this reduction was also restored upon complementation (210.3 ± 126.2 MPa) (Fig. 1F-G,

S1E). Additionally, we evaluated the porosity of the outer membrane through a bisbenzimide

uptake assay, as it is a major determinant of the stiffness of the bacterial cell envelope [32]. We

observed no significant difference in fluorescence uptake between the wild-type and the

mutant, either during the stationary or logarithmic phase (Fig. S1F). Our results indicate that

uppP contributes to the maintenance of bacterial cell size and strength.

To investigate whether the observed defects in bacterial cell length and stiffness of STM Δ*uppP*

can be attributed to biochemical changes in the cell envelope, we evaluated the composition of

two major components: the peptidoglycan layer and lipopolysaccharide (LPS), both of which

are dependent on undecaprenyl phosphate for their biosynthesis. The muropeptide composition

of STM WT and STM Δ*uppP* analysed using HPLC revealed no significant differences in the

muropeptide profile or peptidoglycan cross-links (Figure S2A-C). Similarly, silver staining of

LPS showed no differences in the banding pattern or intensity between the wild-type and the

mutant (Fig. S2D). Collectively, our results suggest that deletion of *uppP* has an impact on

bacterial cell morphology and stiffness, without influencing the biochemical composition of

either the peptidoglycan layer or lipopolysaccharide (LPS).

***uppP* plays a role in the intracellular survival of *Salmonella* in RAW 264.7 macrophages**

As macrophages play a crucial role in the dissemination of *Salmonella* within the host, we

performed an intracellular survival assay using the murine macrophage-like cell line RAW

264.7. We assessed the percentage of phagocytosis/invasion, which represents the percentage

of bacteria from the total pre-inoculum that invade the cell, and fold proliferation, which

accounts for the extent of bacterial replication occurring inside the cell with respect to an early

time point, respectively. We found that STM Δ*uppP* exhibits impaired invasion and fold

proliferation compared to STM WT in RAW264.7 cells. The decreased percentage invasion

and fold proliferation of the mutant is rescued upon complementation of the *uppP* gene (Fig.

2A, S3A). Since the intestinal epithelial barrier is *Salmonella*'s initial point of contact, we also

evaluated the intracellular survival of STM Δ*uppP* in Caco-2 cells. However, we observed no

significant difference in the invasion or fold proliferation of STM Δ*uppP* compared to wild-

type in Caco-2 cells (Fig. S3B-C).

As SPI-1 can also contribute to macrophage invasion [33], we evaluated the transcript levels

of SPI-1 genes. We observed increased *sipA* (SPI-1 effector), *invF*, *hilA* (transcriptional

regulators), and *prgH* (structural translocon component) transcripts in the mutant during both

the logarithmic (6h) and stationary phase (12h) (Fig. S3D-E). Increased expression of SPI-1

genes is also associated with macrophage apoptosis during *Salmonella* infection [34], [35],

[36]. We hypothesised that the reduced invasion and fold proliferation of STM *ΔuppP* may be

attributed to the infected cells undergoing apoptosis. To evaluate the same, 16 hours post-

infection, RAW 264.7 cells were treated with tetracycline, followed by an MTT (3-(4,5-

dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay to determine cell viability. No

significant difference was observed in the cell viability between the wild-type, mutant and

uninfected control (Fig. S3F). This suggests that apoptosis, or other forms of cell death, may

not account for the decreased invasion and proliferation of STM *ΔuppP.*

Salmonella Pathogenicity Island (SPI)-2 is a key player in governing *Salmonella* survival in

macrophages [37]. Since STM *ΔuppP* exhibited reduced fold proliferation in RAW 264.7

macrophages, we evaluated the transcript levels of SPI-2 effectors *sifA* and *spiC*. Interestingly,

STM *ΔuppP* exhibited elevated transcripts of *sifA* and *spiC* during the stationary phase,

compared to the wild-type (Fig. S3G).

Multiple host defences within the macrophage, such as acidic pH, reactive oxygen species,

reactive nitrogen species, antimicrobial peptides, nutrient deprivation, etc, influence

Salmonella survival [8]. We assessed the susceptibility of STM *ΔuppP* to exogenous stresses

of hydrogen peroxide, nitrite and peroxynitrite by a resazurin dye-assisted cell viability assay.

Despite exhibiting elevated reactive oxygen species (ROS) levels, as measured by H_2DCFDA

staining, STM *ΔuppP* displayed only a marginal reduction in viability when subjected to

hydrogen peroxide stress (Fig. 2B-C). In contrast, treatment with increasing concentrations of

sodium nitrite and peroxynitrite resulted in significantly decreased viability of STM *ΔuppP*

compared to the wild-type. (Fig. 2D-G). Additionally, complementing *uppP* in the mutant strain

restored bacterial viability when exposed to nitrite stress (Fig. S3H). We evaluated intracellular

nitric oxide production in STM Δ*uppP* under sodium nitrite stress using DAF_2DA staining.

Flow cytometry-mediated visualisation showed that STM Δ*uppP* demonstrates an increased

percentage of DAF_2DA-positive cells upon nitrite stress (Fig. 2H-I). Moreover, on exposure to

nitrite stress, STM Δ*uppP* exhibited an elevated median fluorescence intensity (MFI) compared

to the wild type, indicating elevated nitric oxide levels (Fig. 2J). Our findings suggest that UppP

contributes to *Salmonella* survival in RAW264.7 cells by aiding the bacterial defence against

nitrosative stress.

UppP deletion leads to increased nitric oxide (NO) production and nitrotyrosine

localisation with intracellular bacteria within macrophages

To evaluate if the reduced fold proliferation of STM Δ*uppP* is attributed to nitrosative stress,

we measured nitric oxide production by infected macrophages. We performed the Griess assay

to evaluate the levels of extracellular NO in the culture supernatants of RAW 264.7 cells

infected with STM WT, STM Δ*uppP* and STM Δ*uppP:uppP* for 16 hours. There was an 8 to

10-fold increase in the extracellular nitrite concentration in the culture supernatants of RAW

264.7 cells infected with STM Δ*uppP* (38.19 ± 5.025 μM) in comparison to the wild-type

control (3.949 ± 1.418 μM). Moreover, upon complementation of *uppP*, the extracellular nitrite

levels were restored (3.172 ± 1.006 μM), similar to that of the wild-type (Fig. 3A). We infected

RAW 264.7 cells with STM WT and STM Δ*uppP,* transformed with the pFPV-mCherry

plasmid, responsible for the constitutive expression of mCherry [38]. This was followed by

DAF_2DA staining 16h post-infection. The STM Δ*uppP*-mCherry infected cells showed

increased MFI compared to STM WT-mCherry infected cells, indicating elevated nitric oxide

levels (Fig. 3B-C, S4A-C). To verify that the elevated NO production was specifically STM

Δ*uppP:uppP* for 16h, followed by DAF_2DA staining. We observed increased intracellular NO

levels in STM Δ*uppP*-infected cells, which were restored to wild-type levels upon

complementation (Fig. S4D-E). NO exerts its bactericidal activity by oxidising to form NO

adducts, such as nitrites and peroxynitrites, with peroxynitrite nitrosylating tyrosine and

impairing protein function [39]. To assess this, we used confocal laser scanning microscopy to

estimate nitrotyrosine levels in RAW 264.7 cells 16h post-infection with eGFP-tagged STM

WT and STM Δ*uppP*. Our results show that in comparison to STM WT, there is an increase in

the colocalisation of nitrotyrosine residues with STM Δ*uppP*. Moreover, cells infected with

STM Δ*uppP* demonstrated increased levels of nitrotyrosine residues, consistent with our flow

cytometry findings (Fig. 3D-F). To further solidify our findings, we performed an ELISA to

quantify nitrotyrosine residues in the lysates of RAW 264.7 cells infected with STM WT, STM

Δ*uppP* and STM Δ*uppP:uppP* for 16h. The cell lysates of STM Δ*uppP*-infected cells exhibited

increased levels of nitrotyrosine residues, which reverted to wild-type levels upon

complementation of *uppP* (Fig. S4F). Taken together, the elevated nitric oxide and nitrotyrosine

levels in RAW 264.7 cells infected with STM Δ*uppP* suggest that *uppP* may have a protective

role against nitrosative stress.

Elevated NO production is also a prominent indicator of M1 polarised proinflammatory

macrophages [40]. To determine whether the increased NO production resulted from a

heightened immune response against the mutant, hence aiding in bacterial clearance, we

quantified the levels of *Nos2* transcript and two other proinflammatory markers, CD86 and

TNF-α, using flow cytometry and ELISA, respectively. The *Nos2* gene encodes for the enzyme

inducible nitric oxide synthase (iNOS), which generates NO. We observed that the *Nos2* mRNA

transcript was elevated in RAW 264.7 cells upon infection with STM Δ*uppP* in comparison to

STM WT (Fig. 3G). Flow cytometry analysis showed that there was an increase in the MFI of

STM Δ*uppP*-infected RAW 264.7 cells in comparison to the STM WT and STM Δ*uppP:uppP*-

infected cells, indicating a higher cell surface expression of CD86 upon infection with STM

Δ*uppP* (Fig. S5A-B). Additionally, we observed that the spent media of RAW 264.7 cells

infected with STM Δ*uppP* showed a minimal increase in TNF-α levels compared to STM WT-

infected cells (Fig. S5C). Altogether, these data indicate that STM Δ*uppP* may exhibit increased

immunogenicity, inducing a stronger proinflammatory immune response within host

macrophages in comparison to STM WT. This NO-driven immune response, along with the

mutant's innate sensitivity to nitrosative stress, may facilitate its clearance in macrophages.

UppP mutant shows increased survival on alleviating nitrosative stress

To corroborate our findings of *uppP*'s role in withstanding nitrosative stress, we inhibited

inducible nitric oxide synthase (iNOS) by an irreversible inhibitor, 1400W [27], [41], and

observed a rescue in the fold proliferation of STMΔ*uppP* in RAW 264.7 cells (Fig. 4A).

To investigate the role of *uppP* in *Salmonella* pathogenesis *in vivo*, C57BL/6 mice were orally

challenged with a dose of 10^7 CFU of STM WT, and STM Δ*uppP* and bacterial burden was

assessed 5 days post-infection (Fig. S6A). STM Δ*uppP* displayed reduced dissemination by

blood and decreased bacterial loads in the intestine, MLN (mesenteric lymph node), liver and

spleen (Fig. S6B-F). Additionally, STM Δ*uppP*-infected mice demonstrated a modest increase

in survival compared to STM WT (Fig. S6G). Furthermore, to evaluate whether the mutant's

attenuation in the mouse model was due to its increased susceptibility to nitrosative stress,

iNOS$^{-/-}$ C57BL/6 mice were orally challenged with 10^5 CFU of STM WT and STM Δ*uppP*

(Fig. S6H). Notably, STM Δ*uppP* still exhibited attenuation in dissemination and reduced organ

burden compared to the wild type (Fig. S6I-M). These findings suggest that UppP may have

additional roles in bacterial invasion of the intestine, beyond its role in conferring resistance to

nitrosative stress. To specifically evaluate UppP role in countering nitrosative stress during

systemic infection, *iNOS*$^{+/+}$ and *iNOS*$^{-/-}$ C57BL/6 mice were intraperitoneally injected with 10^4

CFU of STM WT and STM Δ*uppP* (Fig. 4B). We observed a significantly increased burden of

STM Δ*uppP* in the blood, liver and spleen of *iNOS*$^{-/-}$ mice compared to *iNOS*$^{+/+}$ mice,

substantiating the role of UppP in shielding *Salmonella* from nitrosative stress (Fig. 4C-E).

Induction of iNOS and NO production is triggered by downstream signalling from multiple

pattern recognition receptors (PRRs). These include but are not limited to TLR4, TLR2, NOD1,

and NOD2, out of which the latter of the three receptors detects various components associated

with the bacterial peptidoglycan layer and TLR4 recognises lipopolysaccharide (LPS) [42],

[43], [44], [45]. Also, both NOD1 and NOD2 have been reported to localise on *Salmonella*-

containing vacuole [46], [47]. As UppP is involved in the biosynthesis of both peptidoglycan

and LPS, we aimed to determine whether signalling through their cognate PRRs contributes to

the enhanced susceptibility of STM Δ*uppP* in macrophages. We inhibited TLR2, TLR4 and

RIPK2 (the immediate downstream kinase of NOD receptors) using small molecule inhibitors

C29, Restoravid and WEHI-345, respectively [48], [49], [50]. We observed a rescue in the fold

proliferation of STM Δ*uppP* in RAW 264.7 cells upon treatment with RIPK2 inhibitor WEHI-

345, but not with either C29 or Restoravid, suggesting that NOD signalling may play a role in

the increased susceptibility of STM Δ*uppP* in macrophages (Fig. 4F, S7A). Furthermore, to

identify the NOD receptor which may be involved, we estimated the transcript levels of *Nod1*

and *Nod2* in RAW264.7 cells infected with STM WT and STM Δ*uppP*. Although we did not

observe any significant difference in the transcript levels of *Nod1*, we detected elevated *Nod2*

transcript levels in RAW 264.7 cells, 2 hours post-infection with STM Δ*uppP* (Fig. S7B-C).

We performed immunofluorescence microscopy in STM WT and STM Δ*uppP*-infected

RAW264.7 cells to extend our findings to the translational level. We found that STM Δ*uppP*-

infected RAW264.7 cells at 6 and 16 hours post-infection exhibited a higher corrected total cell

fluorescence (CTCF) of NOD1 and NOD2, respectively, compared to the wild-type and

uninfected control (Fig. 4G-H, S7D-E). To further validate our findings, we treated RAW 264.7

cells with Nodinitib-1 and GSK717, which inhibit NOD1 and NOD2, respectively [51]. We

observed a rescue in the fold proliferation of STM Δ*uppP* in RAW264.7 cells, treated with

either Nodinitib-1 or GSK717, suggesting that both NOD1 and NOD2-mediated signalling may

contribute to the attenuated proliferation of STM Δ*uppP* within macrophages (Fig. 4I).

Discussion

Cell wall integrity is vital for bacterial structure, protection, and survival. Reports in

Escherichia coli, *Salmonella* Typhimurium and *Bacillus subtilis* have shown that cell size and

shape are affected by changes in growth rate and nutritional availability [52], [53], [54], [55].

Although the STM Δ*uppP* mutant exhibited similar growth rate kinetics to the wild type in

LB medium, similar to the observations of *uppP* mutant in *E. coli* [15]. However, we found

structural defects in STM Δ*uppP* through SEM and AFM-assisted visualisation. Quantitative

analysis of SEM images confirmed a biphasic effect: STM Δ*uppP* cells were shorter in

logarithmic phase (1:100 dilution) but longer in stationary phase than wild-type cells.

Interestingly, STM Δ*uppP* cells were notably more elongated than wild-type cells when

allowed to grow in a longer logarithmic phase (1:1000 dilution). This may stem from altered

surface area to volume (SA/V) ratios; increased SA/V in the low dilution (1:100) log phase

could permit division at smaller sizes, while the reduced surface area synthesis rate in the

stationary phase and the high dilution (1:1000) logarithmic phase might facilitate cell

lengthening, resulting in a decrease in the SA/V ratio [56], [57]. Additionally, the AFM

results revealed a reduced Young's modulus of STM Δ*uppP* cells. Undecaprenyl phosphate,

produced by the undecaprenyl pyrophosphate phosphatase activity of *uppP*, is a shared lipid

carrier in the biosynthesis pathways of peptidoglycan, O antigen and enterobacterial common

antigen (ECA), all of which contribute to the stiffness of the cell envelope to varying degrees

[30], [31]. Recent work has demonstrated that the outer membrane comprising O antigen and

ECA is a major determinant of mechanical stiffness in Gram-negative bacteria [32]. In

conjunction with our SEM and AFM findings, this led us to hypothesise that STM Δ*uppP*

may have an altered cell envelope. Interestingly, the biochemical analysis of the composition

of both the peptidoglycan and LPS revealed no significant differences between the mutant

and the wild-type strains. This suggests that the reduced Young's modulus of the mutant and

its biphasic morphological defects may be attributed to the availability dynamics of free

undecaprenyl phosphate, which may in turn affect the biosynthesis rate of these

macromolecules, rather than their composition or structure. Moreover, our bisbenzimide

uptake assay demonstrated that there was no increased outer membrane porosity in the

mutant, suggesting that the loss of *uppP* impairs the mechanical properties of the cell

envelope without affecting the outer membrane's barrier function.

The changes observed in the bacterial cell dimensions of STM Δ*uppP* may also contribute to

its attenuated virulence. Maintaining the correct cellular shape is imperative for the

pathogenicity of various Gram-negative bacteria. For example, peptidoglycan hydrolase

deletion mutants of *Helicobacter pylori* and *Campylobacter jejuni* deviate from their standard

helical shape to a rod or curved rod morphology, which leads to impaired colonisation in the

mouse and chick model, respectively [58], [59], [60]. Furthermore, Uropathogenic *E. coli*

(UPEC) subverts neutrophil-mediated phagocytosis by undergoing filamentation [61]. Thus,

the morphological alterations in STM Δ*uppP* are likely to influence its survival in *in vitro* and

in vivo settings.

To further explore the functional consequence of UppP during *Salmonella* pathogenesis, we

infected RAW 264.7 murine macrophages and Caco-2 colon carcinoma cells with the STM

Δ*uppP*. Though STM Δ*uppP* demonstrated no growth defects in Caco-2, its proliferation was

attenuated in RAW 264.7 cells. Resazurin-assisted live/ dead assay revealed that STM Δ*uppP*

had increased susceptibility to nitrosative stress. Moreover, inhibition of inducible nitric

oxide synthetase restored the proliferation of STM Δ*uppP* in macrophages. Even though

Salmonella avoids trafficking of iNOS-containing vesicles to the *Salmonella*-containing

vacuole (SCV) in an SPI-2-dependent manner, the freely diffusible nature of nitric oxide

(NO) can facilitate bacterium killing [62], [63]. The altered cell structure of the null mutant

may further facilitate the entry of NO within the bacterium. Also, STM Δ*uppP*-infected

macrophages exhibited elevated levels of NO and CD86, along with increased secretion of

TNF-α, suggesting that the mutant elicits a robust proinflammatory immune response within

macrophages. This enhanced immune response, driven by increased NO production and the

mutant's inherent nitrite sensitivity, may promote its attenuated survival within macrophages.

The nucleotide oligomerisation domain (NOD) - like receptors 1 and 2 (NOD1/2) are

cytoplasmic pattern recognition receptors (PRRs) recognising γ-d-glutamyl-meso-

diaminopimelic acid (iE-DAP) and muramyl dipeptide (MDP), which are constituents of the

peptidoglycan layer [64]. NOD1 and NOD2, and RIPK2 (downstream effector protein

kinase) are reported to colocalise with the SCV, along with endosomal transporters SLC15A3

and SLC15A4, which translocate MDP to NOD receptors [46], [47]. In the *Salmonella* colitis

model, mice deficient for NOD1 and NOD2 have higher bacterial loads in the mucosal tissue

[65]. *Mycobacterium tuberculosis* growth in human macrophages is controlled by NOD2-

dependent NO production [66]. Similarly, NOD2 induces NO, which mediates bacterial

clearance in the mouse model of *Chlamydophila pneumoniae*-induced pneumonia [67].

NOD1-induced NO production provides resistance against *Trypanosoma cruzi* in the murine

model [68]. TLR2 and TLR4, which recognise peptidoglycan and LPS, respectively, also

signal the production of NO during bacterial infections [42], [45]. Upon inhibiting RIPK2,

NOD1 or NOD2, but not TLR2 and TLR4, we observed a rescue in the proliferation of STM

Δ*uppP* in macrophages. Undecaprenyl phosphate is a common substrate for synthesising

peptidoglycan, O antigen, and ECA, all of which function as essential virulence determinants

[17]. Initially, we hypothesised that an insufficient influx of undecaprenyl phosphate may

lead to the accumulation of precursors and intermediates, which may act as ligands to directly

or indirectly activate PRRs. In our study, upon deletion of *uppP,* we did not observe any difference in the composition of either LPS or peptidoglycan. We propose an alternative hypothesis that the deletion of *uppP* may slow the synthesis rate of both peptidoglycan and lipopolysaccharide; however, the synthesis of the latter will be more significantly impacted than that of the former, as peptidoglycan is more essential to the cell envelope. The slowed synthesis of peptidoglycan and LPS may explain the mutant's reduced stiffness and altered morphology, leading to its lysis more quickly within macrophages. Furthermore, upon bacterial lysis of the *uppP* mutant, the peptidoglycan fragments may be relatively more abundant, leading to the preferential activation of NOD receptors and driving macrophage activation and elevated NO levels.

In C57BL/6 mice with functional iNOS (*iNOS*$^{+/+}$), intraperitoneal infection with STM Δ*uppP* showed reduced dissemination by blood and decreased colonisation in the spleen and liver. This phenotype was reversed in iNOS-deficient mice, confirming the role of UppP in promoting *Salmonella* dissemination and colonisation by counteracting nitrosative stress.

To the best of our knowledge, our study is one of the first to report a unique function of undecaprenyl pyrophosphate phosphatase in mitigating nitrosative stress in *Salmonella* (Fig. 5), in addition to its canonical roles in cell envelope biogenesis and maintenance of morphology. Our findings potentially highlight undecaprenyl phosphate metabolism as a promising target for developing new antimicrobial strategies to enhance the efficacy of macrophage-mediated clearance of *Salmonella* infection.

Material and methods

Bacterial strains and growth condition: *Salmonella enterica* serovar Typhimurium 14028S (STM WT) was used in all experiments. Bacterial strains were cultured in Luria Bertani (LB) broth (HiMedia) with constant shaking (160 rpm) at 37°C (wild type, knockout strains) or 30°C

364 (strains having temperature-sensitive pKD46 plasmid). Ampicillin (50μg/ml), Kanamycin

365 (50μg/ml), Chloramphenicol (25μg/ml) and IPTG (500μM) were used wherever required. A

366 complete list of bacterial strains and plasmids used in this study is provided in Table 1.

367 Table 1. Strains and plasmids used in this study

Strains/ Plasmids	Characteristics	Source/ Information
Salmonella enterica subspecies *enterica* serovar Typhimurium strain 14028S	Wild type (WT)	Gift from Prof. M. Hensel (Division of Microbiology, University of Osnabrück, Germany)
STMΔ*uppP*	Chl^R	This study
STM Δ*uppP:uppP* (STMΔ*uppP*:pQE60:*uppP*)	Chl^R, Amp^R	This study
STM:pFPV-mCherry	Constitutive expression of mCherry in STM WT, Amp^R	Laboratory stock
STMΔ*uppP*:pFPV-mCherry	Constitutive expression of mCherry in STM Δ*uppP*, Amp^R, Chl^R	This study
STM:pFPV25.1	Constitutive expression of GFP in STM WT, Amp^R	Laboratory stock
STMΔ*uppP*:pFPV25.1	Constitutive expression of GFP in STM Δ*uppP*, Amp^R, Chl^R	This study
pKD3	Plasmid with FRT flanked by Chloramphenicol-resistance gene, Chl^R	Laboratory stock
pKD46	Plasmid expressing λ red recombinase system, Amp^R	Laboratory stock
pQE60	Low copy number plasmid, Amp^R	Laboratory stock
pFPV-mCherry	Constitutive expression of mCherry, Amp^R	Laboratory stock
pFPV25.1	Constitutive expression of GFP, Amp^R	Laboratory stock

Construction of *uppP* knockout strain of *Salmonella*: The knockout strain of *Salmonella* enterica serovar Typhimurium (strain 14028S) was generated using the one-step chromosomal gene inactivation method developed by Datsenko and Wanner [26]. In a concise overview, STM WT was transformed with pKD46 plasmid, housing a 'lambda red recombinase system' controlled by an arabinose-inducible promoter. The transformed cells were cultivated in LB broth with ampicillin (50 μg/mL) and 50 mM arabinose at 30°C until an optical density of 0.35 to 0.4 at 600nm. Electrocompetent STM pKD46 cells were prepared by washing the bacterial cell pellet thrice with double autoclaved chilled Milli Q water and 10% (v/v) glycerol. Finally, the electrocompetent STM pKD46 cells were resuspended in 50 μL of 10% glycerol. For the knockout procedure, the chloramphenicol-resistant gene cassette (Chl^R, 1.1 kb, used for knocking out *uppP*) was amplified from the pKD3 plasmid, utilising knockout primers. The amplified Chl^R gene cassettes underwent chloroform-isopropanol extraction and were electroporated into STM WT pKD46. The transformed cells were then plated on LB agar with chloramphenicol (25 μg/mL) to select the STM*ΔuppP* strain. Subsequently, the plates were incubated overnight at 37°C. The confirmation of knockout colonies was achieved through expression primers. The primers used for generating the knockout strain are listed in Table 2.

Construction of *uppP* complemented strain of *Salmonella*: The *uppP* gene was amplified using cloning primers through colony PCR. The resulting PCR product, purified via chloroform extraction, was then subjected to EcoRI (NEB) and HindIII (NEB) restriction digestion in CutSmart buffer (NEB) at 37°C overnight, alongside the empty pQE60 vector backbone. The double-digested insert and vector were ligated using T4 DNA ligase in 10X ligation buffer (NEB) overnight at 16°C. The ligated product was subsequently transformed into the knockout strains to generate the complemented strain. Verification of complementation was accomplished through the restriction digestion of the recombinant plasmid and Sanger sequencing. The primers used for generating the complement strain are listed in Table 2.

393   Table 2. List of primer sequences (5' to 3')

Primer Name	Sequence (5' to 3')
uppP KO FP	CCAAACGGTTATATATGGGGTCTTAACGCGAGATCCATTTCATATGAATATCCTCCTTAG
uppP KO RP	GCCATCCGGCACGGTCAATCATCAGAAGAAGACAACGTAAGTGTAGGCTGGAGCTGCTTC
uppP confirmatory FP	ATTGAGCGTGGCAATAATCTG
uppP confirmatory RP	CTGCGCTTATCAGACCGACAA
uppP cloning FP	GCGAATTCAGGAGAATGAGCGATATGCACTCGCTGCTG
uppP cloning RP	GCAAGCTTTCAGAAGAAGACAACGTAAACGG
16S rRNA RT FP	CGCTTCTCTTTGTATGCGCC
16S rRNA RT RP	TCGTCAGCTCGTGTTGTGAA
spiC RT FP	AGGGCCGAAGGTAATAGCCGA
spiC RT RP	CCCATCCGCTGTGAGCTGTA
sifA RT FP	TCGCATCCACAAATGACGGC
sifA RT RP	TTCGCTGCCAGGTAGGTTGC
hilA RT FP	GCCGGTGACCATTACGAAGA
hilA RT RP	AAGAGAGAAGCGGGTTGGTG
sipA RT FP	TTCGGATGAAGCGTTGGTCA
sipA RT RP	GTCATTCGCGTGTGGATTCG
invF RT FP	CGTCGTTTGTGCAGCAGAGC
invF RT RP	TAATTTCCGCGGCGAAACGC
prgH RT FP	CGTTTTCAGGTGTTGCCAGG
prgH RT RP	TACGCGGCTCATCGAAATGA
β-actin RT FP	GGTGGGAATGGGTCAGAAGG

β-actin RT RP	AGGTCTCAAACATGATCTGGGT
Nod1 RT FP	CCTCAAGCACCTTAAACTAGGGA
Nod1 RT RP	TCCTTCAATGCCTCTGCGAA
Nod2 RT FP	ACTGTCCAACAATGGCATCAC
Nod2 RT RP	TGTGTTCCCTCGAAGCCAAA
Nos2 RT FP	AGCCACAGTCCTCTTTGCT
Nos2 RT RP	ACAACCTGTGTGACAACCGT

Growth Curve Analysis: For *in vitro* growth curve experiments, a single colony of each strain

was inoculated in Luria Bertani (LB) broth (HiMedia) with constant shaking (160 rpm) at 37°C,

with the corresponding antibiotic. Overnight-grown stationary phase bacteria were sub-

cultured at a ratio of 1: 100 in freshly prepared LB broth, M9 minimal media and F-media

(50mM KCl, 75mM $(NH_4)SO_4$, 5mM K_2SO_4 10mM K_2HPO_4, 1Mm Bis-Tris, 8μM $MgCl_2$,

38mM glycerol and 0.1% cas amino acids, pH- 5.4) kept at 37ºC, with constant shaking (160

rpm). At different time intervals, aliquots were taken to measure [OD]595nm by TECAN 96

well microplate reader or for CFU analysis.

Scanning electron microscopy (SEM): Overnight cultures were subcultured in LB broth at a

1:100 or 1:1000 ratio for 5-6 hours to obtain logarithmic-phase cultures. Logarithmic phase

bacterial cultures were drop-casted and immobilised over coverslips by air-drying, followed by

overnight fixation with 3.5% glutaraldehyde at 4°C. The following day, the samples were

washed with MilliQ water and treated with an increasing ethanol gradient (10%, 30%, 50%,

70%, 90%, and 100%) to facilitate dehydration. The samples were desiccated for 30 minutes,

mounted on copper stubs using carbon tape and coated with gold particles under high vacuum.

Images were acquired with an ESEM Quanta microscope at a magnification of 5000X. ImageJ was used to analyse bacterial size.

AFM (Atomic force microscopy): AFM-mediated assessment of bacterial morphology was performed by adopting the protocol from [69]. Overnight cultures were subcultured in LB broth at a 1:100 ratio for 5-6 hours to obtain logarithmic-phase cultures. Logarithmic phase bacterial cultures were drop-casted and immobilised over coverslips by air-drying, followed by fixation with 3.5% paraformaldehyde. The samples were washed thrice with MilliQ water, followed by desiccation for 30 minutes. Images were acquired from an NX10 AFM. XEI software was utilised to calculate Young's modulus.

Bisbenzimide assay to determine the outer membrane porosity of bacteria: The outer membrane porosity of the bacterial strains (STM WT, STM $\Delta uppP$, and STM $\Delta uppP{:}uppP$) in both the stationary phase and logarithmic phase was measured using bisbenzimide (Sigma-Aldrich), as modified from a previously specified protocol [27], [70]. At the end of the incubation period, the OD_{600nm} of the cultures was adjusted to 0.1 with sterile 1x PBS. Subsequently, 180 μL of 0.1 OD culture was added to a 96-well microplate containing 20 μl of bisbenzimide (10 μg/mL) solution and further incubated for 10 minutes at 37°C in a shaker incubator. Due to enhanced outer membrane porosity, when bisbenzimide is taken up by bacterial cells, it binds to the bacterial DNA and begins to fluoresce. The fluorescence intensity of DNA-bound bisbenzimide was measured in a TECAN 96-well microplate reader using 346 nm excitation and 460 nm emission filter.

Peptidoglycan isolation and analysis: The isolation and analysis of peptidoglycan were performed as described in [71], [72], [73]. Overnight bacterial cultures were subcultured at a 1:100 dilution in 500 mL of LB broth. Bacterial strains were grown till the logarithmic phase ($OD_{600nm} \approx 1$) and subsequently harvested by centrifugation at 8000 RPM for 20 minutes at

4°C. The cell pellet was resuspended in chilled MQ water, and the suspension was added

dropwise to an equal volume of a constantly boiling solution of 8% SDS. The suspension was

boiled for an additional 40 minutes, with continuous stirring and allowed to cool overnight at

RT. The sample was mixed with MQ water and pelleted by ultracentrifugation at 3,00,000g,

for 90 minutes, at RT. After multiple washes to completely remove SDS, the pellet was

resuspended in 10 mM Tris Cl pH 7, 10mM NaCl buffer and treated with 0.1 mg/mL of α-

amylase for 2 hours at 37°C, followed by treatment with 0.2 mg/mL of pronase for 2 hours at

60°C. The enzymes were inactivated by boiling with an equal volume of 8% SDS for 20

minutes, followed by removal of SDS, with several washes of MQ water by centrifugation at

25,000g at RT. The peptidoglycan pellets were finally resuspended in 25 mM Tris-HCl, pH 8,

and stored at -30°C.

For analysing the peptidoglycan composition, muropeptide fragments were obtained by

digesting the peptidoglycan with 10 U of mutanolysin in a 25 mM Tris-HCl buffer, pH 8, at

37°C for 16 hours. The samples were centrifuged at 25,000g for 10 minutes at RT, and the

supernatant comprising the soluble muropeptides was reduced with 1 mg of sodium

borohydride in 50 mM sodium-borate buffer, pH 9, for 30 min. 88% ortho-phosphoric acid was

added to decrease the pH to 2, thereby eliminating the excess borohydride. The sample was

centrifuged to remove any additional particulate matter, and the supernatant was injected into

a preheated (55°C) Zorbax 300SB RP-C18 column, connected to an Agilent Technologies

RRLC 1200 HPLC system. Muropeptides were eluted using a 0 to 20% gradient of acetonitrile

containing 0.1% trifluoroacetic acid with a flow rate of 0.5 mL/min for 80 minutes.

Muropeptides were detected by measuring absorbance at 205 nm.

LPS isolation and quantification: The isolation and quantification of LPS were performed

according to the protocol described by Mahalakshmi et al [74]. Cell lysates were prepared by

sonication of logarithmic-phase strains (STM WT, STM Δ*uppP*, and STM Δ*uppP:uppP*)

resuspended in 10mM HEPES buffer (pH 7.4). The protein concentration of the cell lysates

was estimated using the Bradford Reagent (Bio-Rad) with standard solutions comprising

bovine serum albumin (BSA) concentrations ranging from 50 μg/μL to 0.31 μg/μL. Equal

volumes of cell lysates, normalised to protein content, were mixed with an equal volume of

tricine sample buffer (comprising 100 mM Tris-HCl, pH 6.8, 24% w/v glycerol, 8% w/v SDS,

5% v/v 2-β-mercaptoethanol, and 0.02% w/v bromophenol blue) and boiled for 10 minutes. 10

μL of proteinase K solution (2.5 mg/mL in sample buffer) was added to 50 μL of the boiled

sample, followed by incubation at 60°C for 1 hour. Subsequently, the sample was centrifugation

at 16,000 g for 30 minutes. The supernatants were loaded onto 16.5% tricine-SDS

polyacrylamide gels, and LPS was visualised through silver staining.

Cell culture protocol: DMEM- Dulbecco's Modified Eagle Medium (Lonza) supplemented

with 10% FBS (Gibco) was used to culture RAW 264.7 murine macrophages and the human

colorectal adenocarcinoma cell line, Caco-2. The cells were maintained at 37°C in a humified

incubator with 5% CO_2. Additionally, Caco-2 cells were supplemented with 1% sodium

pyruvate (Sigma) and 1% non-essential amino acids (Sigma). Cells were seeded at a density of

1.5×10^5 cells per well in a 24-well plate for intracellular survival assay preceding infection.

Gentamicin protection assay: Macrophages or epithelial cells were infected with stationary-

phase or logarithmic phase bacterial culture with an MOI of 10. For synchronisation of the

infection and enhancing bacterial adhesion, tissue culture plates were centrifugated at 600 rpm

for 5 minutes, followed by incubation for 25 minutes at 37°C in a humified incubator with 5%

CO_2. Post-incubation cells were washed with 1X PBS, and fresh media (DMEM + 10% FBS)

containing 100μg/ml gentamicin was added, followed by 1h of incubation at 37°C and 5% CO_2.

Subsequently, the gentamicin concentration was reduced to 25μg/ml and maintained until the

cells were harvested.

Percentage phagocytosis and Intracellular proliferation: Following infection of cells with STM at an MOI of 10, cells were treated with DMEM (Sigma) supplemented with 10% FBS (Gibco) containing 100μg/ml gentamicin for 1h. Subsequently, the gentamicin concentration was reduced to 25μg/ml and maintained until the specified time point.

Additionally, in some experiments, 1400W dihydrochloride (Sigma), an inhibitor for inducible nitric oxide synthetase (iNOS), or WEHI-345 (MedChemExpress, HY-18937), a RIPK2 inhibitor which delays NOD signalling [50], Nodinitib-1 (MedChemExpress, HY-18639), an inhibitor of NOD1, and GSK717 (MedChemExpress, HY-136555), an inhibitor of NOD2, were administered to cells at concentrations of 10μM, 1μM, 5μM, and 5μM respectively, alongside 25μg/ml gentamicin treatment. TLR2 and TLR4 were inhibited using C29 (MedChemExpress, HY-100461) and Resatorvid (MedChemExpress, HY-11109), respectively, at concentrations of 25 μM and 100 nm. The cells were treated with either C29 or Resatorvid for 4 hours prior to infection, followed by the addition of the inhibitor alongside the 25μg/ml gentamicin during the entire infection period. 2h and 16h post-infection; cells were lysed in 0.1% triton-X-100. Lysed cells and the pre-inoculum were serially diluted and plated onto LB agar or *Salmonella-Shigella* agar to obtain colony-forming units (CFU). Calculations for percentage phagocytosis/ invasion and fold proliferation are as follows:

$$Percentage\ Phagocytosis\ or\ Invasion\ = \frac{No.\ of\ CFU\ at\ 2h}{No.\ of\ CFU\ in\ Pre\ inoculum} \times 100$$

$$Fold\ Proliferation\ = \frac{No.\ of\ CFU\ at\ 16h}{No.\ of\ CFU\ at\ 2h}$$

RNA isolation and quantitative RT PCR: Bacterial strains were inoculated in LB broth, followed by subculture in LB broth at a ratio of 1:100. Bacterial cells were pelleted and lysed in TRIzol (RNAiso plus Takara). Total RNA was isolated from the TRIzol lysed sample as per the manufacturer's protocol. Quantification of RNA was performed in NanoDrop (Thermo-Fischer Scientific). The quality of isolated RNA was detected by performing gel electrophoresis

with 2% agarose gel. 1μg of RNA was subjected to Turbo DNase (Thermo-Fischer Scientific)

treatment at 37°C for 30 minutes, followed by the addition of EDTA and heat inactivation at

65°C for 10 mins. The mRNA was reverse transcribed to cDNA using oligo dT primer, random

hexamers, buffer, dNTPs, and reverse transcriptase (Takara) as per the manufacturer's protocol.

The expression profile of target genes was evaluated using specific primers by using TB green

RT-qPCR master mix (Takara) in the Applied Biosystem QuantStudio-5 system. 16S rRNA and

β-actin were used as internal controls for bacterial and host genes, respectively. All the

reactions were set up in 384-well plates with three replicates for each sample. The primers used

for RT-qPCR are listed in Table 2 [75], [76], [77].

MTT Assay: Post 16h of infection, cells were treated with complete media containing 10μg/ml

tetracycline for one hour to kill the intracellular bacteria, to negate the influence of MTT (3-

(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) metabolisation by the bacteria

[78]. On completion of the tetracycline treatment, fresh incomplete media (i.e. DMEM only)

containing 0.5mg/ml MTT (Sigma) was added to cells, followed by 3h of incubation at 37°C

and 5% CO2 in the dark. After incubation, media is removed, and DMSO is added, followed

by proper aspiration to dissolve the formazan dye. The optical density was measured at 590nm.

Intracellular Reactive oxygen species (ROS) determination using H₂DCFDA staining:

Overnight cultures were inoculated into fresh LB media. Upon reaching an optical density of

0.1 (logarithmic phase), H₂DCFDA was added at a final concentration of 10μM to 10^8 CFU/ml

of each strain in 1xPBS (pH 7.2) and incubated at 37°C for 30 minutes. Subsequently, 10^7

H₂DCFDA-stained bacterial cells were suspended in 1xPBS (pH 7.2) containing varying

concentrations of hydrogen peroxide (1-5mM) and incubated at 37°C with orbital shaking for

2 hours. The samples were then transferred to a 96-well ELISA plate, and fluorescence was

quantified using a Tecan-ELISA plate reader. (Ex-490nm/ Em-520nm).

Bacterial nitric oxide (NO) determination using DAF-2DA staining: The production of nitric oxide (NO) indicates nitrosative stress. We evaluated the NO generation of the strains to *in vitro* nitrite stress by staining with DAF2-DA (4, 5- diamino fluorescein diacetate). DAF2-DA is a cell membrane-permeable dye that forms a membrane-impermeable adduct emitting fluorescence in NO presence. To mimic nitrite stress *in vitro*, 10^8 CFU/ml of bacteria were treated with a range (500μM-5mM) of acidified nitrite (NaNO2 in PBS of pH 5.4) and incubated for 12 hours at 37°C. Post incubation, the cells are treated with 5μM DAF-DA and incubated for 30 minutes at 37°C. BD FACSverse flow cytometer was used to acquire the samples (Ex-491nm/ Em-513nm).

Resazurin assay: Resazurin dye was used to evaluate the viability of bacteria when subjected to *in vitro* peroxide stress, acidified nitrite and peroxynitrite stress. 10^8 CFU/ml of bacteria were treated with varying concentrations of H_2O_2 (1-5mM) and acidified nitrite (500μM-5mM), simulating oxidative and nitrosative stress, respectively. For peroxynitrite stress, a combination of H_2O_2 and acidified nitrite was used in a combined concentration range of 62.5μM -1mM. The treated bacteria were incubated for 12 hours at 37°C. On completion of treatment, resazurin was added to the bacteria at a final concentration of 1μg/ml, followed by an incubation at 37°C, for 2 hours. Post incubation, fluorescence intensity was measured in TECAN plate reader (Ex-540nm/ Em-590nm). Percentage viability was calculated by the formula:

$$Percentage\ Viability = \frac{Treated\ Flourescence\ Intensity}{Untreated\ Flourescence\ Intensity} \times 100$$

Griess assay for estimating extracellular NO production by macrophages on infection: Griess assay detects the presence of nitrite, a stable product of NO breakdown. The protocol for the assay was adapted from [27], [79]. Briefly, 50μl of culture supernatants of infected macrophages or sodium nitrite standards (0.6, 1.2, 2.4, 4.8, 9.7, 19.5, 39, 78.1 and 156.2 μM)

were treated with 50μl of 1% sulphanilamide (prepared in 5% phosphoric acid) and 50μl of

0.1% NED (N-1-naphthyl ethylene diamine dihydrochloride), followed by an incubation for

10 minutes at room temperature in the dark. On the appearance of a pink-purple colour product

absorbance was measured at 545 nm.

Measurement of NO production in infected macrophages: Intracellular NO production by

RAW264.7 cells upon infection was measured by staining with DAF-2DA. 16h post-infection

cells were treated with 5μM DAF-2DA and incubated for 30 minutes at 37°C. The cells were

washed with 1xPBS and acquired in BD FACSverse flow cytometer (Ex-491nm/ Em-513nm).

Immunofluorescence microscopy for nitrotyrosine residues, NOD1 and NOD2:

RAW264.7 cells were seeded on coverslips and infected with eGFP-tagged strains [80]. The

cells were fixed with 3.5% paraformaldehyde (PFA) 16h post-infection. The sample was

stained with the rabbit-raised anti-nitrotyrosine or goat-raised anti-NOD1 or NOD2 antibody

at a dilution of 1:100 (in 2.5% BSA, 0.01% saponin) overnight at 4°C, followed by staining

with the corresponding secondary antibody tagged with a fluorophore at a dilution of 1:200 in

2.5% BSA, 0.01% saponin for one hour at room temperature. The antibodies used in this study

are listed in Table 3. The coverslips were mounted on glass slides and sealed with transparent

nail polish. The coverslips were imaged by using a Zeiss 880 microscope. Zeiss ZEN Black

2012 software was used to analyse the images. The formula used for calculating corrected total

cell fluorescence (CTCF) is as follows:

$CTCF = Integrated\ density$

$- (area\ of\ cell \times mean\ flourescent\ intensity\ of\ background)$

Table 3. List of antibodies

Sr. No.	Antibody	Catalogue No

1	Anti-Nitrotyrosine Antibody	AB5411 (Sigma-Aldrich)
2	NOD2 (P-18) Antibody	sc-23587 (SCBT)
3	NOD1 (E-14) Antibody	sc-22045 (SCBT)
4	Goat raised anti-rabbit tagged with Cyanine5	Jackson Laboratory
5	Donkey raised anti-goat tagged with TRITC	Jackson Laboratory
6	CD16/CD32 Monoclonal Antibody (93)	14-0161-82 (eBioscience)
7	CD86 (B7-2) Monoclonal Antibody (GL1), PE	12-0862-82 (eBioscience)

ELISA for estimating Nitrotyrosine and TNF-α: For the estimation of Nitrotyrosine, 16h

post-infection, RAW264.7 cells were washed with 1x PBS and pelleted at 500g for 5 minutes.

The cells were lysed with native lysis buffer (150 mM NaCl, 2 mM EDTA, 20 mM Tris, pH 8,

1% NP-40) at 4°C, with intermittent vortexing for 30 minutes, followed by centrifugation at

13,000 RPM for 15 minutes at 4°C. The protein concentration of the supernatant was estimated

using the Bradford Reagent (Bio-Rad). Equal volumes of the supernatant, normalised to protein

content, were applied to the pre-coated ELISA plates of the Nitrotyrosine ELISA kit (Abcam,

ab210603), and the ELISA was performed according to the manufacturer's protocol. To

evaluate levels of TNF-α, the spent medium was collected from infected RAW264.7 cells 16

hours post-infection and subjected to centrifugation at 13,000 RPM, 4°C, for 10 minutes. The

supernatant was added to the pre-coated TNF-α ELISA plates of the GENLISA Mouse TNF-α

ELISA kit (Krishgen BioSystems, KB2145), and the ELISA was performed according to the

manufacturer's protocol. The final absorbance was measured at 450 nm wavelength using a

Tecan Plate reader, and the concentrations of Nitrotyrosine and TNF-α were interpolated from

a standard curve.

Flow cytometry for assessing macrophage cell surface markers: The protocol was adapted

from [81] and is as follows: Infected RAW 264.7 cells were washed and harvested in 1x PBS,

16 hours post-infection. Cells were centrifuged at 500g at 4°C, and the pellet was resuspended

in Fc blocking solution, comprising anti-mouse CD16/CD32 antibody at a dilution of 1:200 in

FACS buffer (1% BSA, 2 mM EDTA in 1x PBS), followed by a 30-minute incubation on ice.

Post-incubation, the cells were washed and stained with a PE-conjugated CD86 antibody at a

dilution of 1:200 for 1 hour on ice. Subsequently, the cells were washed and fixed with 2% PFA

in PBS for 10 minutes. The cells were washed with 1x PBS, resuspended in FACS buffer, and

acquired using CytoFLEX flow cytometer by Beckman Coulter Life Sciences. The data were

then analysed in CytExpert software.

Bacterial burden in murine model: Specific pathogen-free (SPF) 6-8 weeks old C57BL/6

(*iNOS*$^{+/+}$) and *iNos*$^{-/-}$ (*iNOS* knockout in C57BL/6 background) mice were segregated in

cohorts of n=5 or 6. For oral infection, C57BL/6 mice were orally gavaged with 10^7 CFU of

STM WT or STM*ΔuppP*. For *iNOS*$^{-/-}$ mice, a dose of 10^5 CFU was administered orally. The

infected mice were ethically euthanised 5 days post-infection. For intraperitoneal injection,

C57BL/6 (*iNOS*$^{+/+}$) and *iNOS*$^{-/-}$ mice were injected with a dose of 10^4 CFU, followed by ethical

euthanasia 3 days post-infection. Bacterial burden was enumerated in the homogenised lysate

of organs by serial dilution plating on *Salmonella-Shigella* agar.

Mouse survival assay: Specific pathogen-free (SPF) 6-8 weeks C57BL/6 mice were orally

gavaged with 10^7 CFU of STM WT or STM*ΔuppP* (n=6) and monitored daily for mortality.

The data is represented as survival percentage.

Statistical analysis: Data was analysed and graphed using the GraphPad Prism 8 software

(San Diego, CA). Statistical significance was determined by Student's t-test, One-way

ANOVA or Two-way ANOVA to obtain p values. Adjusted p-values below 0.05 are

considered statistically significant. The results are expressed as mean ± SD unless mentioned

otherwise.

Ethics statement

All the animal experiments have been approved by the Institutional Animal Ethics Clearance Committee (IAEC), Indian Institute of Science, Bangalore. The registration number is 48/1999/CPCSEA. The guidelines formulated by the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) were strictly followed. The ethical clearance number for this study is CAF/Ethics/853/2021.

Funding Information

We sincerely appreciate the financial support provided by the Department of Biotechnology (DBT) and the Department of Science and Technology (DST) from the Ministry of Science and Technology. DC acknowledges the Department of Atomic Energy-Scientific Research Council (DAE-SRC) for the Outstanding Investigator Award and funds, ASTRA Chair Professorship, and TATA Innovation fellowship funds. The authors collectively acknowledge the DBT-IISc Partnership Program. We gratefully acknowledge the infrastructure support from the Indian Council of Medical Research (Centre for Advanced Study in Molecular Medicine), DST-FIST (Fund for Improvement of Science and Technology Infrastructure), and the University Grants Commission (special assistance). RV acknowledges the Council of Scientific and Industrial Research (CSIR) for the fellowship. DM and KP acknowledge the Ministry of Human Resource Development (MHRD) for the fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We acknowledge the Real-Time Facility, Departmental Confocal Microscopy Facility and Central Instrumentation Facility, at the Department of Microbiology and Cell Biology (MCB), the Division of Biological Sciences Flow Cytometry Facility, the Atomic Force

Microscopy (AFM) Facility at the Department of Bioengineering (BE), Advanced Facility for

Microscopy and Microanalysis (AFMM) and Central Animal Facility at the Indian Institute of

Science (IISc). Mrs. Deepika and Mrs. Monisha are acknowledged for their help in acquiring

Scanning Electron Microscopy (SEM) images and AFM images, respectively. We are grateful

to Mr. Munish and Mrs. Navya for helping with the data acquisition for flow cytometry. We

thank Mr. Arjun Suresh for his assistance with confocal microscopy image acquisition. Dr

Abhilash Vijay Nair is acknowledged for providing us with the gene of interest for our study.

Author Contributions

Conceptualisation: RV, DM, KP, and DC; Methodology: RV, DM, KP, KCN (PG isolation and

analysis); Investigation: RV, DM, KP, KCN (PG isolation and analysis); Data curation: RV,

Formal analysis: RV, DM, KP, Fund acquisition: DC, MR (PG isolation and analysis) Project

administration: RV and DC, Writing-original draft: RV, Editing and proofreading: RV, DM,

KP, KCN, MR and DC, Supervision: DC.

References

[1]   A. Fàbrega and J. Vila, "Salmonella enterica serovar Typhimurium skills to succeed in
      the host: Virulence and regulation," *Clin Microbiol Rev*, vol. 26, no. 2, pp. 308–341,
      2013, doi: 10.1128/CMR.00066-12.

[2]   R. G. Ferrari, D. K. A. Rosario, A. Cunha-Neto, S. B. Mano, E. E. S. Figueiredo, and
      C. A. Conte-Juniora, "Worldwide epidemiology of Salmonella serovars in animal-
      based foods: A meta-analysis," *Appl Environ Microbiol*, vol. 85, no. 14, Jul. 2019, doi:
      10.1128/AEM.00591-19/SUPPL_FILE/AEM.00591-19-S0001.PDF.

[3]   S. Issenhuth-Jeanjean *et al.*, "Supplement 2008–2010 (no. 48) to the White–
      Kauffmann–Le Minor scheme," *Res Microbiol*, vol. 165, no. 7, pp. 526–530, Sep.
      2014, doi: 10.1016/J.RESMIC.2014.07.004.

[4]   O. Gal-Mor, "Persistent infection and long-term carriage of typhoidal and
      nontyphoidal salmonellae," *Clin Microbiol Rev*, vol. 32, no. 1, Jan. 2019, doi:
      10.1128/CMR.00088-18.

[5]   J. D. Stanaway *et al.*, "The global burden of non-typhoidal salmonella invasive
      disease: a systematic analysis for the Global Burden of Disease Study 2017," *Lancet*

Infect Dis, vol. 19, no. 12, pp. 1312–1324, Dec. 2019, doi: 10.1016/S1473-3099(19)30418-9.

[6] "WHO bacterial priority pathogens list, 2024: Bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance." Accessed: Apr. 14, 2025. [Online]. Available: https://www.who.int/publications/i/item/9789240093461

[7] WHO, "WHO bacterial priority pathogens list, 2024: Bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance," *Bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance*, p. 72, 2024, Accessed: Apr. 14, 2025. [Online]. Available: https://www.who.int/publications/i/item/9789240093461

[8] P. Garai, D. P. Gnanadhas, and D. Chakravortty, "Salmonella enterica serovars typhimurium and typhi as model organisms: Revealing paradigm of host-pathogen interactions," *Virulence*, vol. 3, no. 4, pp. 377–388, 2012, doi: 10.4161/viru.21087.

[9] S. Garde, P. K. Chodisetti, and M. Reddy, "Peptidoglycan: Structure, Synthesis, and Regulation," *EcoSal Plus*, vol. 9, no. 2, Dec. 2021, doi: 10.1128/ecosalplus.esp-0010-2020.

[10] G. Manat *et al.*, "Deciphering the metabolism of undecaprenyl-phosphate: The bacterial cell-wall unit carrier at the membrane frontier," in *Microbial Drug Resistance*, Mary Ann Liebert Inc., Jun. 2014, pp. 199–214. doi: 10.1089/mdr.2014.0035.

[11] S. Kumar, A. Mollo, D. Kahne, and N. Ruiz, "The Bacterial Cell Wall: From Lipid II Flipping to Polymerization," *Chem Rev*, vol. 122, no. 9, pp. 8884–8910, May 2022, doi: 10.1021/ACS.CHEMREV.1C00773/ASSET/IMAGES/LARGE/CR1C00773_0011.JPEG.

[12] S. Booth and R. J. Lewis, "Structural basis for the coordination of cell division with the synthesis of the bacterial cell envelope," *Protein Science*, vol. 28, no. 12, pp. 2042–2054, Dec. 2019, doi: 10.1002/PRO.3722;.

[13] K. Emami *et al.*, "RodA as the missing glycosyltransferase in Bacillus subtilis and antibiotic discovery for the peptidoglycan polymerase pathway," *Nature Microbiology 2017 2:3*, vol. 2, no. 3, pp. 16253-, Jan. 2017, doi: 10.1038/nmicrobiol.2016.253.

[14] A. Taguchi *et al.*, "FtsW is a peptidoglycan polymerase that is functional only in complex with its cognate penicillin-binding protein," *Nature Microbiology 2019 4:4*, vol. 4, no. 4, pp. 587–594, Jan. 2019, doi: 10.1038/s41564-018-0345-x.

[15] M. El Ghachi, A. Bouhss, D. Blanot, and D. Mengin-Lecreulx, "The bacA gene of Escherichia coli encodes an undecaprenyl pyrophosphate phosphatase activity,"

Journal of Biological Chemistry, vol. 279, no. 29, pp. 30106–30113, Jul. 2004, doi: 10.1074/jbc.M401701200.

[16] M. El Ghachi, A. Derbise, A. Bouhss, and D. Mengin-Lecreulx, "Identification of multiple genes encoding membrane proteins with undecaprenyl pyrophosphate phosphatase (UppP) activity in Escherichia coli," *Journal of Biological Chemistry*, vol. 280, no. 19, pp. 18689–18695, May 2005, doi: 10.1074/jbc.M412277200.

[17] L. D. Tatar, C. L. Marolda, A. N. Polischuk, D. van Leeuwen, and M. A. Valvano, "An Escherichia coli undecaprenyl-pyrophosphate phosphatase implicated in undecaprenyl phosphate recycling," *Microbiology (N Y)*, vol. 153, no. 8, pp. 2518–2529, Aug. 2007, doi: 10.1099/mic.0.2007/006312-0.

[18] B. D. Cain, P. J. Norton, W. Eubanks, H. S. Nick, and C. M. Allen, "Amplification of the bacA Gene Confers Bacitracin Resistance to Escherichia coli," 1993.

[19] A. F. Chalker *et al.*, "The bacA gene, which determines bacitracin susceptibility in Streptococcus pneumoniae and Staphylococcus aureus, is also required for virulence," 2000. [Online]. Available: www.tigr.org

[20] E. Gasiorowski *et al.*, "HupA, the main undecaprenyl pyrophosphate and phosphatidylglycerol phosphate phosphatase in Helicobacter pylori is essential for colonization of the stomach," *PLoS Pathog*, vol. 15, no. 9, 2019, doi: 10.1371/journal.ppat.1007972.

[21] L. Röse, S. H. E. Kaufmann, and S. Daugelat, "Involvement of Mycobacterium smegmatis undecaprenyl phosphokinase in biofilm and smegma formation," *Microbes Infect*, vol. 6, no. 11, pp. 965–971, Sep. 2004, doi: 10.1016/J.MICINF.2004.05.011.

[22] T. Geiger, M. Pazos, M. Lara-Tejero, W. Vollmer, and J. E. Galán, "Peptidoglycan editing by a specific ld-transpeptidase controls the muramidase-dependent secretion of typhoid toxin," *Nature Microbiology 2018 3:11*, vol. 3, no. 11, pp. 1243–1254, Sep. 2018, doi: 10.1038/s41564-018-0248-x.

[23] J. C. Quintela, M. A. De Pedro, P. Zöllner, G. Allmaier, and F. G. Del Portillo, "Peptidoglycan structure of Salmonella typhimurium growing within cultured mammalian cells," *Mol Microbiol*, vol. 23, no. 4, pp. 693–704, Feb. 1997, doi: 10.1046/J.1365-2958.1997.2561621.X;

[24] G. Rico-Pérez, A. Pezza, M. G. Pucciarelli, M. A. de Pedro, F. C. Soncini, and F. García-del Portillo, "A novel peptidoglycan D,L-endopeptidase induced by Salmonella inside eukaryotic cells contributes to virulence," *Mol Microbiol*, vol. 99, no. 3, pp. 546–556, Feb. 2016, doi: 10.1111/MMI.13248;

[25] S. B. Hernández *et al.*, "Peptidoglycan editing in non-proliferating intracellular Salmonella as source of interference with immune signaling," *PLoS Pathog*, vol. 18, no. 1, p. e1010241, Jan. 2022, doi: 10.1371/JOURNAL.PPAT.1010241.

[26] K. A. Datsenko and B. L. Wanner, "One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products," *Proc Natl Acad Sci U S A*, vol. 97, no. 12,
pp. 6640–6645, Jun. 2000, doi: 10.1073/PNAS.120163297/ASSET/103C1B8D-D302-
4337-8E67-9F9084156407/ASSETS/GRAPHIC/PQ1201632006.JPEG.

[27] A. R. Chowdhury, S. Sah, U. Varshney, and D. Chakravortty, "Salmonella
Typhimurium outer membrane protein A (OmpA) renders protection from nitrosative
stress of macrophages by maintaining the stability of bacterial outer membrane," *PLoS
Pathog*, vol. 18, no. 8, p. e1010708, Aug. 2022, doi:
10.1371/JOURNAL.PPAT.1010708.

[28] J. Delwick, T. Nikolaus, S. Erdogan, and M. Hensel, "Environmental regulation of
Salmonella pathogenicity island 2 gene expression," *Mol Microbiol*, vol. 31, no. 6, pp.
1759–1773, Apr. 1999, doi: 10.1046/J.1365-2958.1999.01312.X.

[29] M. A. Jorgenson and J. C. Bryant, "A genetic screen to identify factors affected by
undecaprenyl phosphate recycling uncovers novel connections to morphogenesis in
Escherichia coli," *Mol Microbiol*, vol. 115, no. 2, pp. 191–207, Feb. 2021, doi:
10.1111/mmi.14609.

[30] M. A. Jorgenson, S. Kannan, M. E. Laubacher, and K. D. Young, "Dead-end
intermediates in the enterobacterial common antigen pathway induce morphological
defects in Escherichia coli by competing for undecaprenyl phosphate," *Mol Microbiol*,
vol. 100, no. 1, pp. 1–14, Apr. 2016, doi: 10.1111/mmi.13284.

[31] M. A. Jorgenson and K. D. Young, "Interrupting biosynthesis of O antigen or the
lipopolysaccharide core produces morphological defects in Escherichia coli by
sequestering undecaprenyl phosphate," *J Bacteriol*, vol. 198, no. 22, pp. 3070–3079,
2016, doi: 10.1128/JB.00550-16.

[32] E. R. Rojas *et al.*, "The outer membrane is an essential load-bearing element in Gram-
negative bacteria," *Nature 2018 559:7715*, vol. 559, no. 7715, pp. 617–621, Jul. 2018,
doi: 10.1038/s41586-018-0344-3.

[33] G. Wood *et al.*, "Salmonella injectisome penetration of macrophages triggers rapid
translation of transcription factors and protection against cell death," *bioRxiv*, p.
2023.07.21.550113, Jul. 2024, doi: 10.1101/2023.07.21.550113.

[34] D. Hersh, D. M. Monack, M. R. Smith, N. Ghori, S. Falkow, and A. Zychlinsky, "The
Salmonella invasin SipB induces macrophage apoptosis by binding to caspase-1," *Proc
Natl Acad Sci U S A*, vol. 96, no. 5, pp. 2396–2401, Mar. 1999, doi:
10.1073/PNAS.96.5.2396/ASSET/F6D52141-9395-4E16-A419-
3F95A9B47CA4/ASSETS/GRAPHIC/PQ0494845006.JPEG.

[35] Y. Zhang *et al.*, "Programmed cell death and Salmonella pathogenesis: an interactive
overview," *Front Microbiol*, vol. 14, p. 1333500, Jan. 2023, doi:
10.3389/FMICB.2023.1333500/BIBTEX.

[36]   A. McIntosh *et al.*, "SipA activation of caspase-3 is a decisive mediator of host cell
survival at early stages of Salmonella enterica serovar Typhimurium infection," *Infect*
Immun, vol. 85, no. 9, Sep. 2017, doi: 10.1128/IAI.00393-
17/SUPPL_FILE/ZII999092114SM5.AVI.

[37]   E. Jennings, T. L. M. Thurston, and D. W. Holden, "Salmonella SPI-2 Type III
Secretion System Effectors: Molecular Mechanisms And Physiological
Consequences," *Cell Host Microbe*, vol. 22, no. 2, pp. 217–231, 2017, doi:
10.1016/j.chom.2017.07.009.

[38]   D. Drecktrah *et al.*, "Dynamic Behavior of Salmonella-Induced Membrane Tubules in
Epithelial Cells," *Traffic*, vol. 9, no. 12, pp. 2117–2129, Dec. 2008, doi:
10.1111/J.1600-0854.2008.00830.X.

[39]   C. A. Henard and A. Vázquez-Torres, "Nitric oxide and salmonella pathogenesis,"
Front Microbiol, vol. 2, no. APR, p. 9906, Apr. 2011, doi:
10.3389/FMICB.2011.00084/BIBTEX.

[40]   S. Chen *et al.*, "Macrophages in immunoregulation and therapeutics," Dec. 01, 2023,
Springer Nature. doi: 10.1038/s41392-023-01452-1.

[41]   E. P. Garvey *et al.*, "1400W Is a Slow, Tight Binding, and Highly Selective Inhibitor of
Inducible Nitric-oxide Synthase in Vitro and in Vivo," *Journal of Biological*
Chemistry, vol. 272, no. 8, pp. 4959–4963, Feb. 1997, doi: 10.1074/JBC.272.8.4959.

[42]   S. Mukherjee, S. Karmakar, and S. P. S. Babu, "TLR2 and TLR4 mediated host
immune responses in major infectious diseases: a review," *The Brazilian Journal of*
Infectious Diseases, vol. 20, no. 2, pp. 193–204, Mar. 2016, doi:
10.1016/J.BJID.2015.10.011.

[43]   L. O. Moreira and D. S. Zamboni, "NOD1 and NOD2 signaling in infection and
inflammation," *Front Immunol*, vol. 3, no. NOV, p. 35118, Nov. 2012, doi:
10.3389/FIMMU.2012.00328/XML/NLM.

[44]   S. Tötemeyer *et al.*, "IFN-γ Enhances Production of Nitric Oxide from Macrophages
via a Mechanism That Depends on Nucleotide Oligomerization Domain-2," *The*
Journal of Immunology, vol. 176, no. 8, pp. 4804–4810, Apr. 2006, doi:
10.4049/JIMMUNOL.176.8.4804.

[45]   S. Thoma-Uszynski *et al.*, "Induction of direct antimicrobial activity through
mammalian toll-like receptors," *Science (1979)*, vol. 291, no. 5508, pp. 1544–1547,
Feb. 2001, doi: 10.1126/SCIENCE.291.5508.1544/ASSET/AF8C5909-6776-4874-
9ADC-0D81695FA094/ASSETS/GRAPHIC/SE0619197003.JPEG.

[46]   Y. Lu *et al.*, "Palmitoylation of NOD1 and NOD2 is required for bacterial sensing,"
Science (1979), vol. 366, no. 6464, pp. 460–467, Oct. 2019, doi:
10.1126/SCIENCE.AAU6391/SUPPL_FILE/AAU6391-LU-SM.PDF.

[47] N. Nakamura *et al.*, "Endosomes are specialized platforms for bacterial sensing and NOD2 signalling," *Nature 2014 509:7499*, vol. 509, no. 7499, pp. 240–244, Mar. 2014, doi: 10.1038/nature13133.

[48] P. Mistry *et al.*, "Inhibition of TLR2 signaling by small molecule inhibitors targeting a pocket within the TLR2 TIR domain," *Proc Natl Acad Sci U S A*, vol. 112, no. 17, pp. 5455–5460, Apr. 2015, doi: 10.1073/pnas.1422576112.

[49] N. Matsunaga, N. Tsuchimori, T. Matsumoto, and M. Ii, "TAK-242 (resatorvid), a small-molecule inhibitor of Toll-like receptor (TLR) 4 signaling, binds selectively to TLR4 and interferes with interactions between TLR4 and its adaptor molecules," *Mol Pharmacol*, vol. 79, no. 1, pp. 34–41, Jan. 2011, doi: 10.1124/mol.110.068064.

[50] U. Nachbur *et al.*, "A RIPK2 inhibitor delays NOD signalling events yet prevents inflammatory cytokine production," *Nature Communications 2015 6:1*, vol. 6, no. 1, pp. 1–13, Mar. 2015, doi: 10.1038/ncomms7442.

[51] M. Jin *et al.*, "EGFR-MEK1/2 cascade negatively regulates bactericidal function of bone marrow macrophages in mice with Staphylococcus aureus osteomyelitis," *PLoS Pathog*, vol. 20, no. 8 August, Aug. 2024, doi: 10.1371/journal.ppat.1012437.

[52] M. Schaechter, O. MaalØe, and N. O. Kjeldgaard, "Dependency on Medium and Temperature of Cell Size and Chemical Composition during Balanced Growth of Salmonella typhimurium," *Microbiology (N Y)*, vol. 19, no. 3, pp. 592–606, Dec. 1958, doi: 10.1099/00221287-19-3-592.

[53] H. Bremer and P. P. Dennis, "Modulation of Chemical Composition and Other Parameters of the Cell at Different Exponential Growth Rates," *EcoSal Plus*, vol. 3, no. 1, Jan. 2008, doi: 10.1128/ECOSAL.5.2.3/ASSET/EC65E8C7-8572-4870-BAE2-75BD45F44790/ASSETS/GRAPHIC/FIG3.GIF.

[54] M. G. Sargent, "Control of cell length in Bacillus subtilis," *J Bacteriol*, vol. 123, no. 1, pp. 7–19, Jul. 1975, doi: 10.1128/JB.123.1.7-19.1975.

[55] S. Vadia and P. A. Levin, "Growth rate and cell size: a re-examination of the growth law," *Curr Opin Microbiol*, vol. 24, pp. 96–103, Apr. 2015, doi: 10.1016/J.MIB.2015.01.011.

[56] L. K. Harris and J. A. Theriot, "Surface Area to Volume Ratio: A Natural Variable for Bacterial Morphogenesis," *Trends Microbiol*, vol. 26, no. 10, pp. 815–832, Oct. 2018, doi: 10.1016/J.TIM.2018.04.008.

[57] L. K. Harris and J. A. Theriot, "Relative Rates of Surface and Volume Synthesis Set Bacterial Cell Size," *Cell*, vol. 165, no. 6, pp. 1479–1492, Jun. 2016, doi: 10.1016/J.CELL.2016.05.045.

[58] L. K. Sycuro *et al.*, "Peptidoglycan Crosslinking Relaxation Promotes Helicobacter pylori's Helical Shape and Stomach Colonization," *Cell*, vol. 141, no. 5, pp. 822–833, May 2010, doi: 10.1016/J.CELL.2010.03.046.

[59] E. Frirdich et al., "Peptidoglycan-Modifying Enzyme Pgp1 Is Required for Helical Cell Shape and Pathogenicity Traits in Campylobacter jejuni," *PLoS Pathog*, vol. 8, no. 3, p. e1002602, Mar. 2012, doi: 10.1371/JOURNAL.PPAT.1002602.

[60] E. Frirdich and E. C. Gaynor, "Peptidoglycan hydrolases, bacterial shape, and pathogenesis," *Curr Opin Microbiol*, vol. 16, no. 6, pp. 767–778, Dec. 2013, doi: 10.1016/J.MIB.2013.09.005.

[61] S. S. Justice, D. A. Hunstad, P. C. Seed, and S. J. Hultgren, "Filamentation by Escherichia coli subverts innate defenses during urinary tract infection," *Proc Natl Acad Sci U S A*, vol. 103, no. 52, pp. 19884–19889, Dec. 2006, doi: 10.1073/PNAS.0606329104;.

[62] D. Chakravortty, I. Hansen-Wester, and M. Hensel, "Salmonella pathogenicity island 2 mediates protection of intracellular Salmonella from reactive nitrogen intermediates," *Journal of Experimental Medicine*, vol. 195, no. 9, pp. 1155–1166, 2002, doi: 10.1084/jem.20011547.

[63] D. Chakravortty and M. Hensel, "Inducible nitric oxide synthase and control of intracellular bacterial pathogens," *Microbes Infect*, vol. 5, no. 7, pp. 621–627, Jun. 2003, doi: 10.1016/S1286-4579(03)00096-0.

[64] D. J. Philpott, M. T. Sorbara, S. J. Robertson, K. Croitoru, and S. E. Girardin, "NOD proteins: regulators of inflammation in health and disease," *Nature Reviews Immunology 2013 14:1*, vol. 14, no. 1, pp. 9–23, Dec. 2013, doi: 10.1038/nri3565.

[65] K. Geddes et al., "Nod1 and Nod2 regulation of inflammation in the Salmonella colitis model," *Infect Immun*, vol. 78, no. 12, pp. 5107–5115, Dec. 2010, doi: 10.1128/IAI.00759-10.

[66] M. B. Landes, † Murugesan, V. S. Rajaram, H. Nguyen, and L. S. Schlesinger, "Role for NOD2 in Mycobacterium tuberculosis-induced iNOS expression and NO production in human macrophages," *J Leukoc Biol*, vol. 97, no. 6, pp. 1111–1119, Jun. 2015, doi: 10.1189/JLB.3A1114-557R.

[67] K. Shimada et al., "The NOD/RIP2 Pathway Is Essential for Host Defenses Against Chlamydophila pneumoniae Lung Infection," *PLoS Pathog*, vol. 5, no. 4, p. e1000379, Apr. 2009, doi: 10.1371/JOURNAL.PPAT.1000379.

[68] G. K. Silva et al., "Cutting Edge: Nucleotide-Binding Oligomerization Domain 1-Dependent Responses Account for Murine Resistance against Trypanosoma cruzi Infection," *The Journal of Immunology*, vol. 184, no. 3, pp. 1148–1152, Feb. 2010, doi: 10.4049/JIMMUNOL.0902254.

[69] A. R. Chowdhury, D. Mukherjee, A. K. Singh, and D. Chakravortty, "Loss of outer membrane protein A (OmpA) impairs the survival of Salmonella Typhimurium by inducing membrane damage in the presence of ceftazidime and meropenem," *Journal*

897   *of Antimicrobial Chemotherapy*, vol. 77, no. 12, pp. 3376–3389, Nov. 2022, doi:
898   10.1093/JAC/DKAC327.

899   [70]   N. G. Coldham, M. Webber, M. J. Woodward, and L. J. V. Piddock, "A 96-well plate
900   fluorescence assay for assessment of cellular permeability and active efflux in
901   Salmonella enterica serovar Typhimurium and Escherichia coli," *Journal of*
902   *Antimicrobial Chemotherapy*, vol. 65, no. 8, pp. 1655–1663, Aug. 2010, doi:
903   10.1093/JAC/DKQ169.

904   [71]   S. K. Singh, L. Saisree, R. N. Amrutha, and M. Reddy, "Three redundant murein
905   endopeptidases catalyse an essential cleavage step in peptidoglycan synthesis of
906   Escherichia coli K12," *Mol Microbiol*, vol. 86, no. 5, pp. 1036–1051, Dec. 2012, doi:
907   10.1111/mmi.12058.

908   [72]   K. C. Nallamotu, R. Bahadur, M. Kaul, and M. Reddy, "Peptidoglycan Remodeling by
909   an L,D-Transpeptidase, LdtD during Cold Shock in Escherichia coli," *J Bacteriol*, vol.
910   205, no. 1, Jan. 2023, doi: 10.1128/jb.00382-22.

911   [73]   B. Glauner, J. V. Holtje, and U. Schwarz, "The composition of the murein of
912   Escherichia coli.," *Journal of Biological Chemistry*, vol. 263, no. 21, pp. 10088–
913   10095, Jul. 1988, doi: 10.1016/S0021-9258(19)81481-3.

914   [74]   S. Mahalakshmi, M. R. Sunayana, L. Saisree, and M. Reddy, "YciM is an essential
915   gene required for regulation of lipopolysaccharide synthesis in Escherichia coli," *Mol*
916   *Microbiol*, vol. 91, no. 1, pp. 145–157, Jan. 2014, doi: 10.1111/mmi.12452.

917   [75]   A. Datey *et al.*, "Rewiring of one carbon metabolism in Salmonella serves as an
918   excellent live vaccine against systemic salmonellosis," *Vaccine*, vol. 36, no. 50, pp.
919   7715–7727, Nov. 2018, doi: 10.1016/J.VACCINE.2018.10.079.

920   [76]   A. V. Nair, A. Singh, R. S. Rajmani, and D. Chakravortty, "Salmonella Typhimurium
921   employs spermidine to exert protection against ROS-mediated cytotoxicity and rewires
922   host polyamine metabolism to ameliorate its survival in macrophages," *Redox Biol*,
923   vol. 72, p. 103151, Jun. 2024, doi: 10.1016/J.REDOX.2024.103151.

924   [77]   D. Mukherjee and D. Chakravortty, "A Tale of Twins: The Distinct Roles of
925   Intracellular and Extracellular Formate in Regulating Flagellar and Pathogenicity
926   Island-1 Genes in Salmonella Typhimurium," *bioRxiv*, p. 2024.01.30.577997, Nov.
927   2024, doi: 10.1101/2024.01.30.577997.

928   [78]   S. Grille *et al.*, "Salmonella enterica serovar Typhimurium immunotherapy for B-cell
929   lymphoma induces broad anti-tumour immunity with therapeutic effect," *Immunology*,
930   vol. 143, no. 3, pp. 428–437, Nov. 2014, doi: 10.1111/IMM.12320.

931   [79]   P. Das, A. Lahiri, A. Lahiri, and D. Chakravorthy, "Novel role of the nitrite transporter
932   NirC in Salmonella pathogenesis: SPI2-dependent suppression of inducible nitric
933   oxide synthase in activated macrophages," *Microbiology (N Y)*, vol. 155, no. 8, pp.
934   2476–2489, Aug. 2009, doi: 10.1099/MIC.0.029611-0/CITE/REFWORKS.

[80] R. H. Valdivia, A. E. Hromockyj, D. Monack, L. Ramakrishnan, and S. Falkow,
"Applications for green fluorescent protein (GFP) in the study of hostpathogen
interactions," *Gene*, vol. 173, no. 1, pp. 47–52, Jan. 1996, doi: 10.1016/0378-
1119(95)00706-7.

[81] D. Hajra, R. S. Rajmani, A. D. Chaudhary, S. K. Gupta, and D. Chakravortty,
"Salmonella-induced SIRT1 and SIRT3 are crucial for maintaining the metabolic
switch in bacteria and host for successful pathogenesis," *Elife*, vol. 13, Dec. 2024, doi:
10.7554/eLife.93125.

Figure Legends

Main Figures

Figure 1. Loss of *uppP* affects the bacterial cell dimensions and strength

A-C) Growth kinetics of STM WT and STM Δ*uppP* by measuring optical density at 595nm

(OD_{595nm}) in LB (Luria Bertani) broth (A), M9 minimal media (B), and acidic F-media (C).

Data are representative of N=3, n=3 ± SD (standard deviation).

D) Representative scanning electron micrographs of STM WT, STM Δ*uppP* and STM

Δ*uppP*:*uppP* logarithmic phase cells (1:100 dilution). Red and yellow arrows represent cells

with increased and decreased length, respectively. Scale bar = 5μm

E) Quantification of the length of the logarithmic phase (1:100 dilution) bacterial cells captured

in the SEM images (D). Data are representative of N=2, n≥50 ± SD (standard deviation). One-

way ANOVA was performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq$

0.01, *$P \leq 0.05$ ns- non-significant)

F) Representative atomic force images of STM WT, STM Δ*uppP* and STM Δ*uppP*:*uppP*.

G) Quantification of Young's modulus (MPa- Megapascal) of the bacterial cells captured in the

AFM images. Data are representative of n≥30 ± SD (standard deviation). One-way ANOVA

was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05

ns- non-significant)

Figure 2. *uppP* facilitates *Salmonella* to withstand nitrite and peroxynitrite stress

A) Fold proliferation of STM WT, STM Δ*uppP*, and STM Δ*uppP:uppP*, within RAW 264.7

macrophages (MOI=10). Results are mean of N=4, n=3 ± SEM (standard error of mean). One-

way ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤

0.01, *P ≤ 0.05 ns- non-significant)

B) Percentage viability of STM WT and STM *ΔuppP* in the presence of varying concentrations

of H_2O_2 measured by resazurin assay. Results are representative of N=2, n=4 ± SD. Two-way

ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P

≤ 0.05 ns- non-significant)

C) Estimation of reactive oxygen species (ROS) levels in STM WT, STM *ΔuppP* and STM

Δ*uppP:uppP* by H_2DCFDA staining in the presence of varying concentrations of H_2O_2. Results

are representative of N=5, n=4 ± SD. Two-way ANOVA was performed to obtain the p-values.

(****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

D-E) Representative image (D) and percentage viability (E) of resazurin assay of STM WT and

STM Δ*uppP* in the presence of varying concentrations of $NaNO_2$ (*in vitro* nitrite stress). Results

are representative of N=3, n=3 ± SD. Two-way ANOVA was performed to obtain the p-values.

(****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

F-G) Representative image (F) and percentage viability (G) of resazurin assay of STM WT and

STM Δ*uppP* in the presence of varying concentrations of $NaNO_2$ + H_2O_2 (*in vitro* peroxynitrite

stress). Results are representative of N=2, n=4 ± SD. Two-way ANOVA was performed to

obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

H-J) Estimation of nitric oxide in STM WT and STM $\Delta uppP$ by DAF$_2$DA staining in the

presence of varying concentrations of NaNO$_2$ by flow cytometry. Representative SSCA vs

DAF$_2$DA dot plots of STM WT and STM $\Delta uppP$ (H). Quantification of DAF$_2$DA positive

percentage population (I) and median fluorescence intensity (MFI) (J). Results are

representative of N=3, n=4 $\pm$ SD. Two-way ANOVA was performed to obtain the p-values.

(****P $\leq$ 0.0001, ***P $\leq$ 0.001, **P $\leq$ 0.01, *P $\leq$ 0.05 ns- non-significant)

Figure 3. *uppP* sustains *Salmonella* survival in macrophages by countering host-induced

nitrosative stress

A) Estimation of extracellular nitrite by Griess assay in culture supernatants of RAW264.7

cells, 16h post-infection with STM WT, STM $\Delta uppP$ and STM $\Delta uppP:uppP$, MOI=10, N=3,

n=3 $\pm$ SD. One-way ANOVA was performed to obtain the p-values. (****P $\leq$ 0.0001, ***P $\leq$

0.001, **P $\leq$ 0.01, *P $\leq$ 0.05 ns- non-significant)

B-C) Estimation of nitric oxide in macrophages infected with mCherry-tagged STM WT and

STM $\Delta uppP$ at MOI=20 by DAF$_2$DA staining followed by flow cytometry. Representative

mCherry vs SSC-A, DAF$_2$DA vs SSC-A, dot plots and histograms of STM WT and STM

$\Delta uppP$ (B). Quantification of DAF$_2$DA median fluorescence intensity (MFI) of the mCherry-

positive population (C). Results are representative of N=3, n=4 $\pm$ SD. An unpaired two-tailed

Student's t-test was performed to obtain the p-values. (****P $\leq$ 0.0001, ***P $\leq$ 0.001, **P $\leq$

0.01, *P $\leq$ 0.05 ns- non-significant)

D-F) Representative immunofluorescence images of RAW 264.7 cells infected with STM WT

and STM $\Delta uppP$ for 16h, MOI=10, immunostained for nitrotyrosine (D). Quantification of

corrected total cell fluorescence (CTCF) of nitrotyrosine in infected cells (E) and colocalisation

of nitrotyrosine with bacteria (F). Results are representative of N=3, n$\geq$50 $\pm$ SD. Unpaired two-

tailed Student's t-test was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001,

**P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

G) Quantification of *Nos2* mRNA expression in RAW264.7 cells infected with STM WT and

STM Δ*uppP* for 16h. Results are representative of N=2, n=3 ± SD. One-way ANOVA was

performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns-

non-significant)

Figure 4. *uppP* counters nitrosative stress during systemic infection

A) Fold proliferation of STM WT and STM Δ*uppP* within RAW 264.7 macrophages on

treatment with 1400W MOI=10. Results are representative of N=3, n=3 ± SD (standard

deviation). One-way ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤

0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

B) Schematic representation of the experimental protocol followed for intraperitoneal (I.P.) and

infection, enumerating bacterial burden in tissues

C-E) Enumeration of bacterial burden in the blood (C), liver (D), and spleen (E) on infection

via intraperitoneal injection in C57BL/6 (*iNOS*$^{+/+}$) and *iNOS*$^{-/-}$ mice. Results are representative

of N=2 n=7 ± SEM (standard deviation). Mann-Whitney test was performed to obtain the p

values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

F) Fold proliferation of STM WT and STM Δ*uppP* within RAW 264.7 macrophages on

treatment with WEHI-345 MOI=10. Results are representative of N=3, n=3 ± SD (standard

deviation). One-way ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤

0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

G-H) Representative immunofluorescence images of RAW 264.7 cells infected with STM WT

and STM Δ*uppP* for 16h, MOI=10, immunostained for NOD2 (G). Quantification of corrected

total cell fluorescence (CTCF) of NOD2 in infected cells (H). Results are representative of N=2, n≥50 ± SD. One-way ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

I) Fold proliferation of STM WT and STM Δ*uppP* within RAW 264.7 macrophages on treatment with Nodinitib-1 and GSK717, MOI=10. Results are mean of N=3, n=3 ± SEM (standard error of mean). One-way ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

Figure 5. Graphical Abstract

Increased susceptibility of STM Δ*uppP* to host nitrosative stress is indicated by elevated NO levels in RAW 264.7 cells and a greater accumulation of nitrotyrosine residues on the mutant. The elevated NO levels may be due to activation of NOD receptors, which is one of the factors contributing to STM Δ*uppP*'s reduced fold proliferation in RAW 264.7 cells and attenuated survival in the mouse model. NO: Nitric oxide, ONO_2^-: Peroxynitrite, NOD: Nucleotide-binding oligomerisation domain-containing protein, GlcNac: N-acetylglucosamine, MurNac: N-Acetylmuramic acid. Created with BioRender.com

Supplementary Figures

Figure S1. Loss of *uppP* affects the bacterial cell dimensions

A) Representative scanning electron micrographs of STM WT and STM Δ*uppP* stationary phase cells. Red and yellow arrows represent cells with increased and decreased length, respectively. Scale bar = 5μm

B) Quantification of the length of the stationary phase bacterial cells captured in the SEM images (A). Data are representative of N=2, n≥50 ± SD (standard deviation). An unpaired two-

tailed Student's t-test was performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$,

**$P \leq 0.01$, *$P \leq 0.05$ ns- non-significant)

C) Representative scanning electron micrographs of STM WT, STM Δ*uppP* and STM

Δ*uppP*:*uppP* logarithmic phase cells (1:1000 dilution). Red and yellow arrows represent cells

with increased and decreased length, respectively. Scale bar = 5μm

D) Quantification of the length of the logarithmic phase (1:1000 dilution) bacterial cells

captured in the SEM images (C). Data are representative of N=2, n≥50 ± SD (standard

deviation). One-way ANOVA was performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq$

0.001, **$P \leq 0.01$, *$P \leq 0.05$ ns- non-significant)

E) Representative atomic force micrographs of STM WT, STM Δ*uppP* and STM Δ*uppP*:*uppP*.

A 20 x 20μm^2 area from each coverslip was selected for imaging.

F) Measurement of outer membrane porosity of STM WT, STM Δ*uppP* and STM Δ*uppP*:*uppP*,

in the stationary and logarithmic phase by bisbenzimide uptake assay. Data are representative

of N=3, n=4 ± SD (standard deviation). Two-way ANOVA was performed to obtain the p-

values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$ ns- non-significant)

Figure S2. Peptidoglycan and LPS composition of STM Δ*uppP*

A-B) HPLC chromatograms showing petidoglycan composition of STM WT and STM

Δ*uppP* strains grown in Luria Bertani broth. Strains were grown till they reached the early

logarithmic phase (OD$_{600nm}$ ~1). Data are representative of N=2. The identities of the peaks,

according to their mass, are represented in the table (C).

C) Table summarising the identity of the muropeptide fragment peaks observed in HPLC

chromatograms (A) and (B).

D) Silver staining gel of LPS isolated from STM WT, STM Δ*uppP* and STM Δ*uppP:uppP*.

Data are representative of N=2.

Figure S3. *uppP* facilitates the intracellular survival of *Salmonella* in macrophages

A) Percentage phagocytosis of STM WT, STM Δ*uppP*, and STM Δ*uppP:uppP*, within RAW

264.7 macrophages (MOI=10). Results are mean of N=3, n=3 ± SEM (standard error of mean).

One-way ANOVA was performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **P

≤ 0.01, *$P \leq 0.05$ ns- non-significant)

B-C) Percentage invasion (B) and fold proliferation (C) of STM WT, STM Δ*uppP* and STM

Δ*uppP:uppP* within colorectal adenocarcinoma cell line Caco-2 (MOI=10). Results are mean

of N=3, n=3 ± SEM (standard error of mean). An unpaired two-tailed student's t-test was

performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$ ns-

non-significant)

D-E) Quantification of the expression of *hilA*, *prgH*, *invF* and *sipA* during logarithmic (E) and

stationary (F) phase in STM WT and STM Δ*uppP* cultured in LB broth. Results are

representative of N=2, n=3 ± SD. Two-way ANOVA was performed to obtain the p-values.

(****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$ ns- non-significant)

F) Cell viability assay of RAW 264.7 cells, 16h post-infection with STM WT, STM Δ*uppP*,

and STM Δ*uppP:uppP*. Results are representative of N=2, n=3 ± SD. Two-way ANOVA was

performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$ ns-

non-significant)

G) Quantification of the expression of *spiC* and *sifA* during the stationary phase in STM WT

and STM Δ*uppP* cultured in LB broth. Results are representative of N=2, n=3 ± SD. Two-way

ANOVA was performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *P

≤ 0.05 ns- non-significant)

H) Percentage viability of resazurin assay of STM WT, STM Δ*uppP* and STM Δ*uppP:uppP* in

the presence of varying concentrations of $NaNO_2$ (*in vitro* nitrite stress). Results are

representative of N=3, n=3 ± SD. Two-way ANOVA was performed to obtain the p-values.

(****$P \le 0.0001$, ***$P \le 0.001$, **$P \le 0.01$, *$P \le 0.05$ ns- non-significant)

Figure S4. STM Δ*uppP* upregulates nitric oxide (NO) levels in RAW 264.7 cells

A-C) Alternative gating strategy for Fig. 3A-B. Estimation of NO in macrophages infected

with mCherry-tagged STM WT and STM Δ*uppP* by DAF_2DA staining followed by flow

cytometry. Representative DAF_2DA vs mCherry scatter plots of STM WT and STM Δ*uppP*

(A). Quantification of mCherry- DAF_2DA double-positive percentage population (B) and

DAF_2DA median fluorescence intensity (MFI) (C). Results are representative of N=3, n=4 ±

SD. Unpaired two-tailed Student's t-test was performed to obtain the p-values. (****$P \le$

0.0001, ***$P \le 0.001$, **$P \le 0.01$, *$P \le 0.05$ ns- non-significant)

D-E) Estimation of NO, 16h post-infection with STM WT, STM Δ*uppP* and STM Δ*uppP:uppP*

in RAW264.7 cells by DAF_2DA staining followed by flow cytometry. Representative SSC-A

vs DAF_2DA dot plots and Count vs DAF_2DA histogram plots of STM WT, STM Δ*uppP*, and

STM Δ*uppP:uppP* infected RAW 264.7 cells (D). Quantification of median fluorescence

intensity (MFI) of DAF_2DA-positive population (E). Results are representative of N=2, n=4 ±

SD. One-way ANOVA was performed to obtain the p-values. (****$P \le 0.0001$, ***$P \le 0.001$,

**$P \le 0.01$, *$P \le 0.05$ ns- non-significant)

F) Estimation of nitrotyrosine through ELISA, in cell lysates of RAW 264.7 cells infected with

STM WT, STM Δ*uppP*, and STM Δ*uppP:uppP*. Results are representative of N=2, n=2 ± SD.

One-way ANOVA was performed to obtain the p-values. (****$P \le 0.0001$, ***$P \le 0.001$, **$P

\le 0.01$, *$P \le 0.05$ ns- non-significant)

Figure S5. STM Δ*uppP* triggers an enhanced immune response in RAW 264.7 cells

A-B) Flow cytometric analysis of cell surface levels of CD86 in macrophages, 16h post-infection with STM WT, STM Δ*uppP* and STM Δ*uppP:uppP*. Representative Count vs CD86 (PE) histogram plots and SSC-A vs CD86 (PE) dot plots of STM WT, STM Δ*uppP* and STM Δ*uppP:uppP* infected RAW 264.7 cells (A). Quantification of median fluorescence intensity (MFI) of CD86 (PE) positive population (B). Results are representative of N=2, n=4 ± SD. One-way ANOVA was performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$ ns- non-significant)

C) Estimation of pro-inflammatory cytokine TNF-α production through ELISA in cell supernatants of RAW 264.7 macrophages infected with STM WT, STM Δ*uppP*, and STM Δ*uppP:uppP* for 16 hours. Results are mean of N=2, n=2 ± SEM (standard error of mean). One-way ANOVA was performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$ ns- non-significant)

Figure S6. UppP facilitates *Salmonella* survival in the *in vivo* murine model

A) Schematic representation of the experimental protocol followed for enumerating bacterial burden in tissues upon oral gavage in C57BL/6 (*iNOS*$^{+/+}$) mice.

B-F) Enumeration of bacterial burden in the blood (B), intestine (C), MLN (D), spleen (E) and liver (F) on infection via oral route in C57BL/6 (*iNOS*$^{+/+}$) mice. Results are representative of N=2, n=6 ± SD (standard deviation). Mann-Whitney test was performed to obtain the p-values. (****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$ ns- non-significant)

G) Percentage survival of C57BL/6 (*iNOS*$^{+/+}$) mice infected with STM WT and STMΔ*uppP* via the oral route.

H) Schematic representation of the experimental protocol followed for enumerating bacterial burden in tissues upon oral gavage in $iNOS^{-/-}$ mice.

I-M) Enumeration of bacterial burden in the blood (I), intestine (J), MLN (K), liver (L) and spleen (M) on infection via oral route in $iNOS^{-/-}$ mice. Results are representative of n=5 ± SD (standard deviation). Mann-Whitney test was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

Figure S7. NOD receptors contribute to the attenuated survival of STM Δ*uppP* in RAW 264.7 cells

A) Fold proliferation of STM WT and STM Δ*uppP* within RAW 264.7 macrophages on treatment with C29 and Resatorvid, MOI=10. Results are mean of N=3, n=3 ± SEM (standard error of mean). One-way ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

B-C) Quantification of mRNA expression of *Nod1* (B) and *Nod2* (C) in RAW264.7 cells infected with STM WT and STM Δ*uppP* for 2h, 6h and 16h. Results are representative of N=3, n=3 ± SD. Two-way ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

D-E) Representative immunofluorescence images of RAW 264.7 cells infected with STM WT and STM Δ*uppP* for 6h, MOI=10, immunostained for NOD1 (D). Quantification of corrected total cell fluorescence (CTCF) of NOD1 in infected cells (E). Results are representative of N=2, n≥50 ± SD. One-way ANOVA was performed to obtain the p-values. (****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05 ns- non-significant)

Figure 1

Figure 2

Figure 3

Figure 4

Graphical Summary

Figure S1

Figure S2

A) STM WT

B) STM Δ*uppP*

C)

Peak no	Muropeptide Species	Cross-link type
1	Tri	Un-crosslinked
2	Di	Un-crosslinked
3	Tetra(Gly^4)	Un-crosslinked with Glycine in 4th position
4	Terta	Un-crosslinked
5	Tri-lys-arg	Tri peptide bound to Braun's lipoprotein (Lpp)
6	Tri-Tri	3-3 cross-links
7	Tetra-Tri	4-3 cross-links
8	Tri-Tetra	3-3 cross-links
9	Tetra-Tetra	4-3 cross-links
10	Tetra-Tri-lys-arg	4-3 cross linked muropeptide bound to Lpp
11	Tetra-Tetra-Tetra	4-3 cross-links
12	Tetra-Tetra-Anh	4-3 cross-links

D)

Figure S3

Figure S4

Figure S5

A)

Autofluorescence Uninfected STM WT STM ΔuppP STM ΔuppP:uppP

CD86 (PE)

B)

RAW 264.7

C)

RAW 264.7

Figure S6

Figure S7