**Separating anorexia -dependent and -independent effects in cancer cachexia**

Yanshan Liang [1*], Young-Yon Kwon [1*], Sheng Hui [1 2 #]

[1] Department of Molecular Metabolism, Harvard T. H. Chan School of Public Health, Boston, MA, USA

[2] Lead contact

[*] The authors contributed equally

[#] Correspondence: shui@hsph.harvard.edu

**Abstract**

Cancer cachexia is characterized by unintentional weight loss and wasting away of fat and

muscle tissues. Anorexia, or reduced food intake, is often implicated as a contributor to the

negative energy balance in this condition. However, to what extent anorexia alone accounts for

body weight loss and wasting of different tissues, and whether anorexia is responsible for other

cachectic phenotypes such as physical performance impairment remains insufficiently

characterized in preclinical models and patients. In this study, we demonstrate the critical need to

address these questions in cancer cachexia research. Using the colon carcinoma 26 (C26) model

of cancer cachexia as an example, we systematically determined how much each of the key

phenotypes of cancer cachexia is driven by anorexia. Anorexia was the predominant driver for

body weight loss, adipose tissue wasting, and muscle wasting, strikingly suggesting the lack of

other mechanisms for causing these phenotypes in this model. In contrast, anorexia had no

impact on physical performance, pointing to the existence of anorexia-independent mechanisms

in causing fatigue. Thus, for a given preclinical model or patient group, anorexia can be the main

cause for certain cachectic phenotypes and play no role in causing other cachectic phenotypes.

Discriminating between anorexia-mediated and independent effects is essential for guiding

research focus and ultimately unraveling the causal pathways of cancer cachexia.

**Introduction**

Cancer cachexia is an involuntary wasting condition prevalent in cancer patients, contributing to poor quality of life and reduced survival [1,2]. However, the etiology of the disease remains to be fully understood, and no effective treatment is currently available [3]. One challenge in establishing the causal pathways of cancer cachexia is to distinguish causality from mere association, as evidenced in our previous work showing that the widely assumed cachectic factor IL-6 likely only associates with cancer cachexia [4]. Another often-cited factor contributing to cancer cachexia is anorexia. However, to what extent anorexia contributes to the body weight loss and also to other phenotypes of cancer cachexia, such as loss of fat mass and muscle mass, impairment of physical performance, and alteration of systemic metabolism, has been rarely rigorously assessed in preclinical models or human patients. This is a critical question because the answer dictates the focus of downstream investigations. For example, food intake reduction is known to cause reduced muscle mass even in healthy animals [5,6]. If anorexia alone accounts for most muscle mass loss in cancer cachexia, research should be focused on unraveling mechanisms causing anorexia rather than some active mechanisms of muscle wasting. The same question can be posed for other phenotypes of cancer cachexia, such as adipose tissue wasting and metabolic alterations, both of which are heavily influenced by food intake [7,8]. Thus, in this study, we focus on the causality of food intake reduction in different cachectic phenotypes and demonstrate the importance of separating anorexia-dependent and - independent effects in cancer cachexia research.

Among previous studies of cancer cachexia, a marginal fraction of them controlled for food intake. While the lack of food intake control in patient research is likely due to practical infeasibility, we were only able to find on the order of 10 preclinical studies that used a pair-

feeding strategy (matching daily food intake between cachectic and non-cachectic groups) [9–14]

and a couple that employed a caloric restriction strategy (matching total food intake between

cachectic and non-cachectic groups) [15,16]. These studies focused on one or two specific

phenotypes such as body weight or adipose tissue wasting, and missed key phenotypes such as

physical performance. Thus, a study is needed to systematically delineate the causal roles of

anorexia in all key phenotypes of cancer cachexia.

Here, we systematically measured how reduced food intake contributes to cachexia

phenotypes and metabolic changes in the syngeneic murine colon 26 carcinoma (C26) cancer

cachexia model [17] using an isocaloric feeding strategy. Our findings revealed that anorexia is a

primary driver for body weight loss, fat mass loss, and muscle protein loss in this model,

suggesting the lack of additional mechanisms for these cachectic phenotypes. In contrast,

anorexia does not affect physical performance, highlighting anorexia-independent cachectic

pathways. Additionally, anorexia is responsible for key alterations of systemic metabolism,

including reduced circulating glucose and increased ketone body levels, suggesting no need to

invoke additional mechanisms to explain these metabolic changes. Therefore, distinguishing

anorexia-induced effects from other cachectic symptoms is crucial for guiding research efforts to

identify the causal pathways of cancer cachexia.

**Results**

**Anorexia accounts for almost all body weight loss and body composition changes in the**

**C26 model**

We first aimed to investigate the contribution of anorexia to body weight loss, a primary

phenotype of cancer cachexia, in the C26 model. The C26 model was set up by subcutaneously

injecting the C26 cells into the syngeneic CD2F1 mice[4,17]. Two weeks post-tumor inoculation,

these mice exhibited over 15% body weight loss manifested by fat and lean mass reductions. We

refer to this group as the cachectic C26 (cxC26) group. To control for effects due to tumor

growth, we included a mouse group using a non-cachectic C26 cell line (ncxC26), which

maintained body weight and composition despite similar or bigger tumor size (Supplementary

Fig. S1).

The cxC26 mice showed clear anorexia (Fig. 1a). Consistent with previous observations

[18–20], these mice produced a large amount of masticated food debris (Supplementary  Data

Fig. S2). For accurate food intake measurement, mice were single-housed in cages with paper

bedding, which allows reliable separation of food debris and bedding materials. To determine the

extent to which anorexia contributes to body weight loss, we initially employed a pair-feeding

approach by providing a group of ncxC26 mice with a single portion of food equal to the amount

consumed by cxC26 cachectic mice ad lib. The pair-fed ncxC26 mice showed similar body

weight loss as the cxC26 mice under comparable tumor burden (Supplementary Fig. S1a, c). Fat

and lean mass reductions were consistently comparable between the pair-fed ncxC26 and cxC26

mice, suggesting a predominant role for anorexia in body weight loss and composition

(Supplementary Fig. S1b). A potential caveat in the pair-feeding experiment is that pair feeding

alters the diurnal food intake patterns and subjects the animals to a fixed feeding schedule, which

can induce caloric deprivation and lead to different metabolic and physiological responses in

freely-fed animals [21,22]. Thus, to minimize potential differences in feeding behavior between

the two groups, we adopted an isocaloric feeding approach in which both ncxC26 and cxC26

mice were provided daily the same amount of food matching the food intake of a group of cxC26

mice ad lib as shown in Fig. 1a. Similar to the pair-feeding experiment, the body weight curves

were in close match between the isocaloric-fed ncxC26 and cxC26 groups (Fig. 1b-c).

Furthermore, the extent of both lean mass loss and fat mass loss was also the same between the

two isocaloric-fed groups (Fig. 1d). Importantly, there was no difference in tumor size between

the two groups (Fig. 1e). These results showed that anorexia accounts for almost all of the three

key cachectic phenotypes: body weight loss, fat mass loss, and lean mass loss.

**Neither energy expenditure nor fecal energy excretion contributes to body weight loss**

Our observation that anorexia is the sole driver for body weight loss implies a limited

role for energy expenditure (EE) in body weight loss, which is often cited to be concomitant with

cancer cachexia [23,24]. To test this, we measured the EE using a metabolic cage system for the

isocaloric-fed ncxC26 and cxC26 mice. With identical caloric intake, the time-course EE data in

cxC26 mice closely followed that of ncxC26 mice, which was lower than the ncxC26 mice ad lib

(Fig. 2a). The lack of difference in EE between the two isocaloric-fed groups was also evident in

the cumulative EE data (Fig. 2b). The metabolic cage data further showed no difference between

the two groups in the respiratory exchange ratio (RER), indicating their similar substrate

utilization at the whole-body level (Fig. 2c-d).

Another important variable in the energy balance equation is energy excretion, which can

be affected by possible malabsorption complications [25,26] in cancer cachexia. To determine its

role in energy balance and body weight loss, we collected all feces produced within the 24 hours

before the endpoint from the three groups. We then measured fecal calories using bomb

calorimetry. Our results indicated that the two isocaloric-fed groups had similar fecal production

though significantly less than the ncxC26 mice ad lib (Fig. 3a). As the fecal energy density was

not changed across the groups (Fig. 3b), there was reduced but identical total fecal energy

excretion in the two isocaloric-fed groups (Fig. 3c). Together, these results demonstrated that

neither energy expenditure nor fecal energy excretion contributes to body weight loss; instead,

anorexia accounts for almost all body weight loss as well as body composition in the C26 model.

**Anorexia is the main driver for muscle protein mass loss in cancer cachexia**

Skeletal muscle wasting is a key hallmark of cancer cachexia, and the tumor-secreted

factors have been reported to have direct effects on muscle [27,28]. Our body composition data,

however, suggested that the muscle wasting could be simply due to anorexia. To investigate

further the effects of anorexia on skeletal muscle mass, at the endpoint, we dissected and

weighed the lower limb muscles, including gastrocnemius, soleus, tibialis anterior (TA), and

extensor digitorum longus (EDL). Consistent with the lean mass result, reduced food intake for

the ncxC26 group resulted in generally reduced muscle mass (comparing the isocaloric-fed

ncxC26 group to the ncxC26 ad lib group), albeit to slightly less degree than the cxC26 group in

gastrocnemius and soleus (Fig. 4a). The muscle mass change was not due to glycogen mass

change because of low muscle glycogen content (<1%) (Fig. 4b). As proteins are the main

component of muscle tissue, we next assessed the protein content of the gastrocnemius muscle.

We observed no significant difference in the protein content of gastrocnemius muscle between

isocaloric-fed cxC26 and ncxC26 mice (Fig. 4c), indicating similar magnitude of muscle protein

mass loss in the two groups (Fig. 4d). This finding was corroborated by the gene expression

levels of muscle atrophy biomarkers *Murf1 and Atrogin1* (Fig. 4e), which were elevated in the

ncxC26 group to the same level as in the cxC26 group. Altogether, these results showed that it is

reduced food intake, not some specific cachectic factor, that is the main driver for muscle protein

wasting in the C26 model.

**Physical performance is independent of anorexia-induced muscle mass loss in cancer**

**cachexia**

To continue assessing the effects of anorexia on key phenotypes in cancer cachexia, we

next determined the impact of anorexia on physical performance. For this, we first measured grip

strength and found surprisingly that compared to the ncxC26 ad lib group, despite reduced

muscle mass, the isocaloric-fed ncxC26 group did not show a decrease in grip strength (Fig. 4e).

In contrast, the isocaloric-fed cxC26 group showed clear reduced grip strength (Fig. 4e). Grip

strength impairment has been reported to be closely associated with fatigue [29–31]. To subject

this finding to further testing, we next measured physical activity, which is well related to fatigue

[32] and used for fatigue evaluation [33]. Both daily pedestrian locomotion and daily distance

covered in the cage showed increased activity in the isocaloric-fed ncxC26 group, reflecting a

foraging behavior under starvation, but not in the isocaloric-fed cxC26 group (Fig. 4f). This

observation, in line with the grip strength results, suggests that anorexia has a limited role in

contributing to fatigue. Finally, we performed an additional assay to evaluate fatigue-like

behavior [34]. In this test, mice were forced to run on a treadmill until exhaustion to assess their

maximal endurance performance. Again, consistent with the grip strength and the physical

activity data, the isocaloric-fed cxC26 group showed a substantial reduction in performance

while the isocaloric-fed ncxC26 group did not (Fig. 4g). Together, these results pointed to

limited role of anorexia in contributing to fatigue.

The independence of fatigue on muscle mass is a striking result. A potential caveat for

this conclusion is that cachectic animals exhibited slightly but significantly smaller muscle mass

compared to their isocaloric-fed non-cachectic counterparts (Fig. 4a). Thus, to eliminate this

potential caveat and more rigorously test our finding, we subjected a group of ncxC26 mice to a

severe caloric restriction, reducing their food intake by 30% relative to that of the cxC26 mice ad

lib starting from day 7 post-tumor injection (Fig. 5a). With this approach, the body weight, lean

and fat mass, and muscle mass of the caloric-restricted ncxC26 mice were aligned with, and in

some cases slightly lower than, those of the cxC26 mice ad lib (Fig. 5b-d). We employed the grip

strength assay for fatigue evaluation and observed that the grip strength of ncxC26 caloric-

restricted mice was comparable to that of the freely fed ncxC26 mice despite substantial food

intake reduction (Fig. 5e). As the caloric-restricted ncxC26 group now unambiguously had

matched muscle mass with the cxC26 group, their largely intact physical performance clearly

demonstrated that fatigue is independent of anorexia-induced muscle mass loss.

**Anorexia can account for important metabolic alterations observed in cancer cachexia.**

To probe the extent to which anorexia contributes to cancer cachexia-associated

metabolic changes, we performed a metabolomics analysis on serum samples collected from the

ncxC26 ad lib, isocaloric-fed ncxC26, and isocaloric-fed cxC26 mice. Based on principal

component analysis (PCA), isocaloric restriction caused significant serum metabolome changes,

demonstrated by the distinct separation between the ncxC26 ad lib cluster and the isocaloric-fed

ncxC26 cluster (Fig. 6a). Note that the isocaloric-fed cxC26 cluster diverged from the isocaloric-

fed ncxC26 cluster, suggesting the presence of tumor-dependent metabolic effects. We further

identified significantly altered metabolites (adjusted p-values < 0.05) across comparisons with

isocaloric-fed cxC26 versus ncxC26 ad lib representing the conventional comparison, isocaloric-

fed ncxC26 versus ncxC26 ad lib showing the caloric restriction-induced metabolic changes, and

isocaloric-fed cxC26 versus isocaloric-fed ncxC26 showing anorexia-independent metabolic

changes. As shown in the Venn diagram, 31 out of 51 caloric restriction-induced metabolites

overlapped with those identified in the conventional comparison (Fig. 6b), suggesting that

metabolomics results on cancer cachexia can be misleading in terms of reflecting metabolic

abnormalities if food intake is not controlled for. Only 26 out of 72 significantly altered

metabolites from the conventional comparison dataset overlapped with net cancer cachexia-

associated metabolic changes (Supplementary Fig. S3). Changes in important metabolic

indicators such as glucose and ketone bodies (acetoacetate and 3-hydroxybutyrate) (Fig. 6c) and

liver glycogen content (Fig. 6d) are likely consequences of reduced food intake instead of

reflecting metabolic abnormalities in cancer cachexia.

**Discussion**

This study demonstrates the importance of separating anorexia-dependent and -

independent effects in cancer cachexia research. The findings that key cachectic phenotypes,

including body weight loss, mass loss of adipose tissue and muscle, and alterations of key

metabolites, are mostly contributable to anorexia in the C26 model inform that a search for new

mechanisms for these phenotypes may not be justified in this model. Instead, the finding that

fatigue occurs independently of anorexia highlights novel mechanisms that warrant further

investigation. The extent to which anorexia contributes to cachectic phenotypes is generally

different in different preclinical models and different subtypes of patients. Thus, we propose to

make it a standard procedure to get accurate food intake data and perform rigorous control for

food intake in studies of cancer cachexia.

Our results caution using the expression of the genes *Murf1* and *Atrogin1* as markers of

muscle atrophy in cancer cachexia. Consistently with the muscle mass data, food intake

reduction alone induces their expression to the same level as in the cachectic C26 mice. Thus,

while these gene markers may reflect muscle mass loss, their use as an indication of active

muscle wasting processes is not validated.

It was surprising to find that the loss of adipose tissue and muscle mass was simply due

to the food intake reduction in the C26 model. This is because the model has often been used for

studying active mechanisms of adipose tissue wasting [14,35] and muscle wasting [36–38].

These studies generally did not check food intake or conduct food intake control, and thus cannot

distinguish whether the wasting was primarily due to food intake reduction or some active

mechanisms. While we cannot rule out the complete lack of additional mechanisms of tissue

wasting, our results indicate a minor role for them in this model.

Our study showed no elevation of energy expenditure in the C26 model. Despite being

commonly cited as a contributor to body weight loss in cancer cachexia, energy expenditure

elevation has only been observed in limited number of cases, including some patients with lung

and pancreatic cancer [39], as well as  preclinical models such as Lewis lung cancer (LLC)

cachexia model [23] and a skin cancer cachexia model [40].

Our results revealed that fatigue does not necessarily depend on food intake reduction or

muscle mass loss in cancer cachexia. With comparable anorexia-induced muscle loss, only the

cachectic animals exhibited reductions in grip strength, physical activities, and maximum

exercise capacity, while non-cachectic animals remained as competent as the ad lib control. In

line with our results, a study using a lung cancer cachexia model compared caloric-restricted

mice with cachectic mice and observed similar reductions in fat mass, lean mass, and skeletal

muscle mass, but no change of muscle function in the caloric-restricted group [41]. Grip strength

has been reported to be preserved [42–44], and exercise endurance has been shown to increase

[45,46] in humans after caloric restriction, which aligns with what we found in the calorie-

restricted non-cachectic mice. Thus, different from muscle wasting or muscle atrophy, fatigue is

a distinct phenotype in cancer cachexia. Fatigue could be due to either muscle function

impairment [47,48] or neural dysfunction [49–51], which so far has been understudied in cancer

cachexia research and deserves more attention in the field.

By isolating out the effects of reduced food intake, the goal is to simplify the complex

problem of cancer cachexia and ultimately establish the causal pathways of cancer cachexia. We

do not downplay the role of anorexia in cancer cachexia. Instead, as shown in our results,

anorexia plays a dominant role in multiple phenotypes of cachexia and thus deserves intensive

study. We simply argue for separate treatments of anorexia-dependent and -independent

phenotypes while searching for the underlying mechanisms (Fig. 7). In this structure of causal

pathway, a successful strategy for therapeutical development might be to act separately on

appetite and physical performance. In line with our view, several clinical trials targeting the

ghrelin pathway for increased food take had no effects on hand grip strength and overall survival

despite reported improved lean body mass [52–54]. While inhibition of GDF15 signaling was

reported to have mitigated not only body weight loss but also physical performance impairment

through rescuing appetite in preclinical cancer cachexia models (HT-1080 [55] and TOV21G

(ovarian cancer) [56]), the mechanism by which GDF15 impacts physical performance is unclear

and the physical performance in a recent clinical trial of an anti-GDF15 antibody had

inconsistent dose-dependence [57]. Thus, a future effective treatment is likely a combination of

two drugs, targeting anorexia and fatigue.

**Methods**

**Animals**

CD2F1 male mice were purchased from Charles River Laboratories and were housed (4

per cage) under regular light-dark cycles of 12 h with ad lib access to water and food (PicoLab

5053, LabDiet). Prior to tumor implantation, mice were single-housed and acclimated to

experimental conditions for 1 week. Tumor implantation was conducted when animals were 12-

16 weeks old and reached a stable stage in growth and weight gain. One million of cxC26 or

ncxC26 cells per animal were implanted subcutaneously in a 200-µL injection of saline cell

suspension into the right flank of mice. The length of the studies was 2 weeks. Animal care and

experimental procedures were conducted with the approval of the Institutional Animal Care and

Use Committees (IACUC) of Harvard Medical School and Harvard T.H. Chan School of Public

Health.

**Cell culture**

The cachectic C26 (cxC26) and non-cachectic (ncxC26) cells were kind gifts from

Andrea Bonetto [17] and Nicole Beauchemin, respectively. Nicole Beauchemin obtained the

ncxC26 cells from Brattain MG, who established the C26 (also named CT26 or colon carcinoma

26) cell line [58]. All the cell lines were maintained using high glucose DMEM (Corning) with

10% fetal bovine serum (FBS, R&D Systems) and 1% penicillin/streptomycin (P/S, Hyclone).

Cells were maintained in incubators at 37 °C and 5% $CO_2$.

**Food intake and isocaloric feeding**

For food intake measurement and during the isocaloric feeding, animals were housed

individually on alpha pad bedding (6" × 10", scottPharm solution) with virgin kraft crinkled

paper strips as nesting material (Enviro-Dri™, scottPharm solution) (See Extended Data Fig. 2a).

Food pellets were weighed and placed each day on the lid. The mass of food pellets on the lid

from the previous day was checked and recorded the next day. Any visible food debris (Extended

Data Fig. 2a) on the bedding was collected, weighed, and discarded. The food surplus would be

the sum of pellets on the lid and the bedding. Daily food intake was calculated as the weight

difference in food pellets between 2 consecutive days.

For the isocaloric feeding, the mice in all three groups were body weight-matched. A

group of ncxC26 tumor-bearing mice were on ad lib as a control. From day 6 to day 12 after

tumor injection, the cxC26 mice and another group of ncxC26 mice were fed the same amount of

food daily following the food intake of the cxC26 mice we measured in Fig. 1a. Food debris on

the bedding will be checked and weighed daily and returned to the mice. On day 12, any mice

with more than 1g of food (about 5% of the total food intake of the isocaloric feeding) left would

be excluded from the experiment.

**Body weight and body composition assessment**

Body weight was measured every other day following implantation (approximately 11

am and 1 pm). Whole-body composition (fat mass and lean mass) analysis was performed using

an EchoMRI-100 (EchoMRI LLC) on each animal at day 0 and day 12 prior to their euthanasia.

Each mouse was individually placed in a specialized plastic tube for mild conscious restraint and

inserted into a magnetic resonance imager (MRI) to measure body composition. Each mouse

took approximately 5 min to complete the procedure. Mice were then returned to their home cage.

**Metabolic cage analyses**

Mice were individually housed at 22 °C and subjected to indirect calorimetry between

day 5 and day 12 after tumor inoculation under a 12 h light-dark cycle using the Promethion

Metabolic Cage system (Sable Systems, USA). During this period, spontaneous activity, and

volume of oxygen consumed and carbon dioxide produced were measured. This data allows us to

calculate energy expenditure (EE) and the respiratory exchange ratio (RER). Fold changes in

daily locomotion and cage distance were calculated by comparing values from day 12 to day 0.

**Fecal energy determination**

Feces produced 24 hours before the endpoint were collected and stored at -80 °C. The feces were dehydrated at 60 °C for 48 hours in an Eppendorf tube with lid open. Then, the calorie content of the dry feces was determined by using Bomb Calorimetry (Parr Oxygen Bomb). Wet and dry weights were measured for later calculations of total fecal production (g/day) and fecal caloric content (cal/g).

**Muscle mass and grip strength**

At the end of the study, animals were euthanized, and gastrocnemius, soleus, tibialis anterior (TA), and extensor digitorum longus (EDL) were isolated and weighed. To measure grip strength, mice were allowed to grab the metal grid and then were pulled backward by an experimenter until the grasp was released. The peak tension was measured using a Grip Strength meter (Bioseb).

**Maximum exercise capacity test**

Mice were acclimated to a motorized treadmill for three consecutive days from day 4-6 before the maximal exercise capacity test. Shocking grids with frequency set at 2 Hz and intensity at 1.22 mA were located at the end of the treadmill to force the mice to run at their maximum capacity. The treadmill was set at 10° incline. The running test was conducted at the end of day 12. Animals ran at 6 m/min for the first 2 min, at 10 m/min for the following 8 min, followed by 2 m/min increases every 5 min until exhaustion, defined by the inability to remain on the treadmill for >5 s.

**RNA extraction and RT-qPCR**

Fresh gastrocnemius samples were isolated from ncxC26 ad lib, isocaloric-fed ncxC26, and isocaloric-fed cxC26 tumor-bearing mice and immediately snap-frozen using liquid nitrogen.

Total RNA was extracted using the Trizol reagent (Invitrogen, #15596026) from the 20-30 mg of

gastrocnemius and TA muscle tissues, following the manufacturer's protocol. RNA

concentration was determined using NanoDrop One (Thermofisher Scientific). One microgram

of purified RNA was converted to cDNA using the iScript Reverse Transcription Supermix for

RT-qPCR (Bio-Rad, #1708840). The diluted cDNA was used for RT-qPCR using SsoAdvanced

Universal SYBR Green Supermix (Bio-Rad, #1725270) with the CFX384 Real-Time PCR

detection System (Bio-Rad). The following primer sequences were used to determine the

Atrogin-1 and MuRF1 mRNA expression. Cyclophilin was used as a housekeeping gene for

normalization.

Table 1. Primer sequences

| Primer | Sequence 5'-3' |
|---|---|
| Atrogin1 forward | TCA GAG AGG CAG ATT CGC AA |
| Atrogin1 reverse | GGG TGA CCC CAT ACT GCT CT |
| MuRF-1 forward | TCC TGA TGG AAA CGC TAT GGA G |
| MuRF-1 reverse | ATT CGC AGC CTG GAA GAT GT |
| Cyclophilin forward | GGA GAT GGC ACA GGA GGA A |
| Cyclophilin reverse | GCC CGT AGT GCT TCA GCT T |

**Measurement of glycogen in tissues**

Mice were fasted for 7-8 hours before the tissue collection. They were euthanized by

cervical dislocations, and tissues (liver, gastrocnemius, and soleus) were immediately dissected,

clamped with a pre-chilled Wollenberger clamp, and rapidly submerged in liquid nitrogen for

snap-1freezing. Frozen tissues were then ground under cryogenic condition using a CryoMill

(Retsch). For glycogen quantification, 10 – 20 µg of ground tissue were used in the Glycogen-

Glo assay kit (Promega, #J5051), following the manufacturer's instructions.

**Serum metabolomics**

Serum metabolites were extracted using extraction buffer (acetonitrile: methanol: water,

4: 4: 2, *v: v: v*) at a volume-to-volume ratio of 1:20 (µL to µL), followed immediately by

vortexing for 10 seconds. The mixture was then incubated at 4 °C for 10 min before

centrifugation at 16,000 g at 4 °C for 10 min. The supernatant was transferred to an LC-MS vial

and analyzed on the LC-MS. A pooled quality control (QC) sample was prepared by mixing 20

µL from each experimental sample.

Metabolite extracts were analyzed using a quadrupole-orbitrap mass spectrometer

coupled with hydrophilic interaction chromatography (HILIC). Chromatographic separation was

achieved on an XBridge BEH Amide XP Column (2.5 µm, 2.1 mm × 150 mm) with a guard

column (2.5 µm, 2.1 mm × 5 mm) (Waters, Milford, MA). For the gradient, mobile phase A was

water: acetonitrile 95:5, and mobile phase B was water: acetonitrile 20:80, both phases

containing 10 mM ammonium acetate and 10 mM ammonium hydroxide. The linear elution

gradient was: 0 ~ 3 min, 100% B; 3.2 ~ 6.2 min, 90% B; 6.5. ~ 10.5 min, 80% B; 10.7 ~ 13.5

min, 70% B; 13.7 ~ 16 min, 45% B; and 16.5 ~ 22 min, 100% B, with a flow rate of 0.3 mL/ min.

The autosampler was at 4˚C. The column temperature was at 30˚C. The injection volume was 5

µL. Needle wash was applied between samples using acetonitrile: methanol: water at 4: 4: 2 (*v: v:*

*v*). For mass spectrometry, a Q Exactive HF (Thermo Fisher Scientific, San Jose, CA) mass

spectrometer was used. For MS1 acquisition, the MS scanned from 70 to 1000 m/z with

switching polarity at a resolution of 120,000 for all experimental samples. The relevant

parameters were: sheath gas, 40; auxiliary gas, 10; sweep gas, 2; spray voltage, 3.5 kV; capillary

temperature, 300 °C; S-lens, 45. The resolution was set at 120,000 (at m/z 200). Maximum

injection time (max IT) was set at 500 ms, and automatic gain control (AGC) was set at $3 \times 10^6$.

MS1 raw data files were converted into mzxML using msconvert and imported to EI-

Maven (Elucidata, Cambridge, MA) for targeted metabolomics. Metabolites were identified

using an in-house library based on accurate mass and retention time.

**Statistical analysis**

All statistical analyses for murine data were performed in GraphPad Prism 10 or R studio.

Quantitative data are reported as mean with standard deviation (s.d.). Three-group comparisons

were analyzed using one-way ANOVA followed by Tukey post hoc. Metabolites with adjusted

p-values less than 0.05 are considered significant. For all analyses, a p-value of $<0.05$ was

considered significant (*$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$, ****$p < 0.0001$).

**Figure captions**

**Figure 1. Anorexia is the main driver of body weight loss and body composition changes.**

CD2F1 male mice were inoculated with $1{\times}10^{6}$ ncxC26 or cxC26 cells subcutaneously. **(a)** Food

intake of ncxC26 and cxC26 mice ad lib.  **(b)** Percentage change in body weight, **(c)** absolute

initial and endpoint body weight, **(d)** body composition measured using EchoMRI at the

endpoint, and **(e)** tumor size at the endpoint, for ncxC26 ad lib, isocaloric-fed ncxC26, and

isocaloric-fed cxC26 groups. **(a)** n=16 for ncxC26 ad lib and n=16 for cxC26 ad lib. **(b)** n=20 for

ncxC26 ad lib, n=20 for isocaloric-fed ncxC26, n=15 for isocaloric-fed cxC26. **(c-d)** n=18 for

ncxC26 ad lib, n=18 for isocaloric-fed ncxC26, n=14 for isocaloric-fed cxC26. **(e)** n=21 for

ncxC26 ad lib, n=22 for isocaloric-fed ncxC26, n=22 for isocaloric-fed cxC26. All data are

shown as the mean ± s.d. Significance of the differences: ns not significant, $*P < 0.05$, $**P <$

$0.01$, $***P < 0.001$, $****$ $P<0.0001$ between groups by one-way ANOVA followed by Tukey

post hoc comparison.

**Figure 2. Energy expenditure does not contribute to body weight loss.**

**(a)** Hourly energy expenditure from day 6 to day 12 post-inoculation. **(b)** Cumulative energy

expenditure during full day, daytime, and nighttime. **(c)** Hourly respiratory exchange ratio from

day 6 to day 12 post-inoculation. **(d)** Average respiratory exchange ratio. **(a-d)** n=4 per group. **(b**

**and d)** are shown as the mean ± s.d. Significance of the differences: ns not significant, $*P < 0.05$,

$**P < 0.01$, $***P < 0.001$, $****$ $P < 0.0001$ between groups by one-way ANOVA followed by

Tukey post hoc comparison.

**Figure 3. Energy excretion does not contribute to body weight loss.**

Feces produced by mice within 24 hours before euthanasia were collected. **(a)** Fecal production,

**(b)** Fecal energy density **(c)** Total fecal energy excretion. **(a-c)** n=8 for ncxC26 ad lib, n=8 for

isocaloric-fed ncxC26, and n=5 for isocaloric-fed cxC26. All data are shown as the mean ± s.d.

Significance of the differences: ns not significant, *P < 0.05, **P < 0.01, ***P < 0.001, **** P <

0.0001 between groups by one-way ANOVA followed by Tukey post hoc comparison.

**Figure 4. Food intake reduction induces muscle protein wasting but not fatigue in cancer**

**cachexia.**

**(a)** Muscle mass. **(b)** Muscle glycogen content. **(c)** Muscle protein content. **(d)** Total protein

mass in gastrocnemius calculated by multiplying averaged protein content with muscle mass. **(e)**

mRNA expression levels of atrophy markers in gastrocnemius. **(f)** Grip strength. **(g)** Changes in

daily pedestrian locomotion and distance in cages comparing D12 to D0. **(h)** Maximum exercise

capacity (running on a treadmill until exhaustion). **(a, d)** n=23 for ncxC26 ad lib, n=24 for

isocaloric-fed ncxC26, n=28 for isocaloric-fed cxC26. **(b)** n=6 for ncxC26 ad lib, n=7 for

isocaloric-fed ncxC26, n=6 for isocaloric-fed cxC26. **(c)** n=10 for ncxC26 ad lib, n=12 for

isocaloric-fed ncxC26, n=8 for isocaloric-fed cxC26. **(e)** n=10 for ncxC26 ad lib, n=12 for

isocaloric-fed ncxC26, n=9 for isocaloric-fed cxC26. **(f)** n=22 for ncxC26 ad lib, n=23 for

isocaloric-fed ncxC26, n=23 for isocaloric-fed cxC26. **(g)** n=4 for ncxC26 ad lib, n=4 for

isocaloric-fed ncxC26, n=6 for isocaloric-fed cxC26. **(h)** n=6 for ncxC26 ad lib, n=8 for

isocaloric-fed ncxC26, n=7 for isocaloric-fed cxC26. All data are shown as the mean ± s.d.

Significance of the differences: ns not significant, *P < 0.05, **P < 0.01, ***P < 0.001, **** P <

0.0001 between groups by one-way ANOVA followed by Tukey post hoc comparison.

**Figure 5. Fatigue is independent of anorexia-induced muscle mass loss in cancer cachexia.**

A group of ncxC26 mice was under severe caloric restriction (severeCR), receiving 30% less

food intake than a group of cxC26 mice ad lib starting from day 7 post-tumor inoculation. **(a)**

Food intake. **(b)** Percentage change in body weight. **(c)** Percentage change of body composition

**(d)** Muscle mass. **(e)** Grip strength at the endpoint. **(a-e)** n=6 for ncxC26 ad lib, n=6 for ncxC26

severeCR, and n=5 for cxC26 ad lib. All data are shown as the mean ± s.d. Significance of the

differences: ns not significant, *P < 0.05, **P < 0.01, ***P < 0.001, **** P < 0.0001 between

groups by one-way ANOVA followed by Tukey post hoc comparison.

**Figure 6. Anorexia accounts for important metabolic alterations observed in cancer**

**cachexia.**

**(a)** Principle component analysis of the serum metabolomics. **(b)** Venn diagram of the significant

metabolites found in different comparisons. **(c)** Serum glucose and ketone bodies, acetoacetate

and 3-hydroxybutyrate. **(d)** Liver glycogen content. **(e)** Diagram summarizing the anorexia-

dependent and -independent cachectic phenotypes in the C26 model. **(a, c)** n=11 for ncxC26 ad

lib, n=10 for isocaloric-fed ncxC26, n=12 for isocaloric-fed cxC26. **(d)** n=6 for ncxC26 ad lib,

n=7 for isocaloric-fed ncxC26, n=6 for isocaloric-fed cxC26.

**Figure 7. The causal pathway of cancer cachexia consists of mechanisms causing anorexia**

**and anorexia-independent mechanisms.**

**Supplementary Figure S1. Pair-feeding recapitulates the body weight loss and body**

**composition of the cachectic C26 mice ad lib.**

CD2F1 male mice were inoculated with $1\times10^6$ cxC26 or ncxC26 cells. A group of ncxC26 mice

were pair-fed to match the food intake of cxC26 mice ad lib. **(a)** Percentage change of body

weight. **(b)** Body composition. **(c)** Tumor mass. **(a-c)** n=8 for ncxC26 ad lib, n=8 for pair-fed

ncxC26, and n=8 for cxC26 ad lib.

**Supplementary Figure S2. Cage setup for food intake measurement and masticated food**

**debris in cages of cxC26 and ncxC26 mice.** **(a)** Cage setup. Mice were single-housed with

alpha paper bedding and paper strips as nesting material. **(b)** Mass of food debris in cages of

ncxC26 and cxC26 mice (n=8 per group). The cxC26 mice produced significantly more food

debris than the ncxC26 mice. The mass of food debris should be considered when calculating the

daily food intake.

**Supplementary Figure S3. Volcano plot for serum metabolites between isocaloric-fed**

**cxC26 and ncxC26 groups.** Metabolites with adjusted p-values $< 0.05$ were considered as

significant changes, with elevated metabolites shown in red and decreased ones in green. N=10

for isocaloric-fed ncxC26, n=12 for isocaloric-fed cxC26.

## Reference

[1]   Fearon, K., Strasser, F., Anker, S.D., Bosaeus, I., Bruera, E., Fainsinger, R.L., et al., 2011. Definition and classification of cancer cachexia: an international consensus. The Lancet Oncology 12(5): 489–95, Doi: 10.1016/S1470-2045(10)70218-7.

[2]   Peixoto Da Silva, S., Santos, J.M.O., Costa E Silva, M.P., Gil Da Costa, R.M., Medeiros, R., 2020. Cancer cachexia and its pathophysiology: links with sarcopenia, anorexia and asthenia. Journal of Cachexia, Sarcopenia and Muscle 11(3): 619–35, Doi: 10.1002/jcsm.12528.

[3]   Fearon, K., Arends, J., Baracos, V., 2013. Understanding the mechanisms and treatment options in cancer cachexia. Nature Reviews Clinical Oncology 10(2): 90–9, Doi: 10.1038/nrclinonc.2012.209.

[4]   Kwon, Y.-Y., Hui, S., 2024. IL-6 promotes tumor growth through immune evasion but is dispensable for cachexia. EMBO Reports 25(6): 2592–609, Doi: 10.1038/s44319-024-00144-3.

[5]   Weiss, E.P., Jordan, R.C., Frese, E.M., Albert, S.G., Villareal, D.T., 2017. Effects of Weight Loss on Lean Mass, Strength, Bone, and Aerobic Capacity. Medicine & Science in Sports & Exercise 49(1): 206–17, Doi: 10.1249/MSS.0000000000001074.

[6]   He, Y., Yang, W., Huang, L., Mever, M.A., Ramautar, R., Harms, A., et al., 2024. Metabolomic analysis of dietary□restriction□induced attenuation of sarcopenia in prematurely aging DNA repair□deficient mice. Journal of Cachexia, Sarcopenia and Muscle 15(3): 868–82, Doi: 10.1002/jcsm.13433.

[7]   Lane, M.A., Baer, D.J., Rumpler, W.V., Weindruch, R., Ingram, D.K., Tilmont, E.M., et al., 1996. Calorie restriction lowers body temperature in rhesus monkeys, consistent with a postulated anti-aging mechanism in rodents. Proceedings of the National Academy of Sciences 93(9): 4159–64, Doi: 10.1073/pnas.93.9.4159.

[8]   Most, J., Redman, L.M., 2020. Impact of calorie restriction on energy metabolism in humans. Experimental Gerontology 133: 110875, Doi: 10.1016/j.exger.2020.110875.

[9]   Wang, W., Lonnroth, C., Svanberg, E., Lundholm, K., 2001. Cytokine and Cyclooxygenase-2 Protein in Brain Areas of Tumor-bearing Mice with Prostanoid-related Anorexia. Cancer Research 61: 4707–15.

[10]  Belury, 2010. Cardiac alterations in cancer-induced cachexia in mice. International Journal of Oncology 37(2), Doi: 10.3892/ijo_00000683.

[11]  Tsoli, M., Moore, M., Burg, D., Painter, A., Taylor, R., Lockie, S.H., et al., 2012. Activation of Thermogenesis in Brown Adipose Tissue and Dysregulated Lipid Metabolism Associated with Cancer Cachexia in Mice. Cancer Research 72(17): 4372–82, Doi: 10.1158/0008-5472.CAN-11-3536.

[12]  Kliewer, K.L., Ke, J.-Y., Tian, M., Cole, R.M., Andridge, R.R., Belury, M.A., 2015. Adipose tissue lipolysis and energy metabolism in early cancer cachexia in mice. Cancer Biology & Therapy 16(6): 886–97, Doi: 10.4161/15384047.2014.987075.

[13]  Reddel, C.J., Allen, J.D., Ehteda, A., Taylor, R., Chen, V.M.Y., Curnow, J.L., et al., 2017. Increased thrombin generation in a mouse model of cancer cachexia is partially interleukin□6 dependent. Journal of Thrombosis and Haemostasis 15(3): 477–86, Doi: 10.1111/jth.13612.

[14] Chen, X., Wu, Q., Gong, W., Ju, S., Fan, J., Gao, X., et al., 2024. GRP75 triggers white adipose tissue browning to promote cancer-associated cachexia. Signal Transduction and Targeted Therapy 9(1): 253, Doi: 10.1038/s41392-024-01950-w.

[15] Flint, T.R., Janowitz, T., Connell, C.M., Roberts, E.W., Denton, A.E., Coll, A.P., et al., 2016. Tumor-Induced IL-6 Reprograms Host Metabolism to Suppress Anti-tumor Immunity. Cell Metabolism 24(5): 672–84, Doi: 10.1016/j.cmet.2016.10.010.

[16] Sudo, Y., Otsuka, H., Miyakawa, R., Goto, A., Kashiwase, Y., Terawaki, K., et al., 2018. Differential Metabolic Responses to Adipose Atrophy Associated with Cancer Cachexia and Caloric Restriction in Rats and the Effect of Rikkunshito in Cancer Cachexia. International Journal of Molecular Sciences 19(12): 3852, Doi: 10.3390/ijms19123852.

[17] Bonetto, A., Rupert, J.E., Barreto, R., Zimmers, T.A., 2016. The Colon-26 Carcinoma Tumor-bearing Mouse as a Model for the Study of Cancer Cachexia. Journal of Visualized Experiments (117): 54893, Doi: 10.3791/54893.

[18] Villars, F., Pietra, C., Giuliano, C., Lutz, T., Riediger, T., 2017. Oral Treatment with the Ghrelin Receptor Agonist HM01 Attenuates Cachexia in Mice Bearing Colon-26 (C26) Tumors. International Journal of Molecular Sciences 18(5): 986, Doi: 10.3390/ijms18050986.

[19] Zhang, W., Sun, W., Gu, X., Miao, C., Feng, L., Shen, Q., et al., 2022. GDF-15 in tumor-derived exosomes promotes muscle atrophy via Bcl-2/caspase-3 pathway. Cell Death Discovery 8(1): 162, Doi: 10.1038/s41420-022-00972-z.

[20] Flint, T.R., Janowitz, T., Connell, C.M., Roberts, E.W., Denton, A.E., Coll, A.P., et al., 2016. Tumor-Induced IL-6 Reprograms Host Metabolism to Suppress Anti-tumor Immunity. Cell Metabolism 24(5): 672–84, Doi: 10.1016/j.cmet.2016.10.010.

[21] Rupp, H., Maisch, B., Brilla, C.G., 1997. Schedule-induced psychological stress and molecular structures of cardiomyocytes. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology 272(3): R776–82, Doi: 10.1152/ajpregu.1997.272.3.R776.

[22] Russell, J.C., Proctor, S.D., Kelly, S.E., Brindley, D.N., 2008. Pair feeding-mediated changes in metabolism: stress response and pathophysiology in insulin-resistant, atherosclerosis-prone JCR:LA-cp rats. American Journal of Physiology-Endocrinology and Metabolism 294(6): E1078–87, Doi: 10.1152/ajpendo.90257.2008.

[23] Kir, S., White, J.P., Kleiner, S., Kazak, L., Cohen, P., Baracos, V.E., et al., 2014. Tumour-derived PTH-related protein triggers adipose tissue browning and cancer cachexia. Nature 513(7516): 100–4, Doi: 10.1038/nature13528.

[24] Palesty, J.A., Dudrick, S.J., 2003. What We Have Learned about Cachexia in Gastrointestinal Cancer. Digestive Diseases 21(3): 198–213, Doi: 10.1159/000073337.

[25] Riner, A.N., Herremans, K.M., Vudatha, V., Han, S., Qu, X., Liu, J., et al., 2024. Heterogeneity of weight loss and transcriptomic signatures in pancreatic ductal adenocarcinoma. Journal of Cachexia, Sarcopenia and Muscle 15(1): 149–58, Doi: 10.1002/jcsm.13390.

[26] Hendifar, A.E., Petzel, M.Q.B., Zimmers, T.A., Denlinger, C.S., Matrisian, L.M., Picozzi, V.J., et al., 2019. Pancreas Cancer-Associated Weight Loss. The Oncologist 24(5): 691–701, Doi: 10.1634/theoncologist.2018-0266.

[27] Bilgic, S.N., Domaniku, A., Toledo, B., Agca, S., Weber, B.Z.C., Arabaci, D.H., et al., 2023. EDA2R–NIK signalling promotes muscle atrophy linked to cancer cachexia. Nature 617(7962): 827–34, Doi: 10.1038/s41586-023-06047-y.

[28] Liu, X., Li, S., Cui, Q., Guo, B., Ding, W., Liu, J., et al., 2024. Activation of GPR81 by lactate drives tumour-induced cachexia. Nature Metabolism, Doi: 10.1038/s42255-024-01011-0.

[29] Kilgour, R.D., Vigano, A., Trutschnigg, B., Hornby, L., Lucar, E., Bacon, S.L., et al., 2010. Cancer□related fatigue: the impact of skeletal muscle mass and strength in patients with advanced cancer. Journal of Cachexia, Sarcopenia and Muscle 1(2): 177–85, Doi: 10.1007/s13539-010-0016-0.

[30] Cantarero-Villanueva, I., Fernández-Lao, C., Díaz-Rodríguez, L., Fernández-de-las-Peñas, C., Ruiz, J.R., Arroyo-Morales, M., 2012. The Handgrip Strength Test as a Measure of Function in Breast Cancer Survivors: Relationship to Cancer-Related Symptoms and Physical and Physiologic Parameters. American Journal of Physical Medicine & Rehabilitation 91(9): 774–82, Doi: 10.1097/PHM.0b013e31825f1538.

[31] Veni, T., Boyas, S., Beaune, B., Bourgeois, H., Rahmani, A., Landry, S., et al., 2019. Handgrip fatiguing exercise can provide objective assessment of cancer-related fatigue: a pilot study. Supportive Care in Cancer 27(1): 229–38, Doi: 10.1007/s00520-018-4320-0.

[32] Romero, S.A.D., Jones, L., Bauml, J.M., Li, Q.S., Cohen, R.B., Mao, J.J., 2018. The association between fatigue and pain symptoms and decreased physical activity after cancer. Supportive Care in Cancer 26(10): 3423–30, Doi: 10.1007/s00520-018-4203-4.

[33] Dougherty, J.P., Springer, D.A., Cullen, M.J., Gershengorn, M.C., 2019. Evaluation of the effects of chemotherapy-induced fatigue and pharmacological interventions in multiple mouse behavioral assays. Behavioural Brain Research 360: 255–61, Doi: 10.1016/j.bbr.2018.12.011.

[34] Dougherty, J.P., Springer, D.A., Gershengorn, M.C., 2016. The Treadmill Fatigue Test: A Simple, High-throughput Assay of Fatigue-like Behavior for the Mouse. Journal of Visualized Experiments (111): 54052, Doi: 10.3791/54052-v.

[35] Rohm, M., Schäfer, M., Laurent, V., Üstünel, B.E., Niopek, K., Algire, C., et al., 2016. An AMP-activated protein kinase–stabilizing peptide ameliorates adipose tissue wasting in cancer cachexia in mice. Nature Medicine 22(10): 1120–30, Doi: 10.1038/nm.4171.

[36] Pranzini, E., Muccillo, L., Nesi, I., Santi, A., Mancini, C., Lori, G., et al., 2024. Limiting serine availability during tumor progression promotes muscle wasting in cancer cachexia. Cell Death Discovery 10(1): 510, Doi: 10.1038/s41420-024-02271-1.

[37] Lin, K., Wei, L., Wang, R., Li, L., Song, S., Wang, F., et al., 2024. Disrupted methionine cycle triggers muscle atrophy in cancer cachexia through epigenetic regulation of REDD1. Cell Metabolism: S1550413124004133, Doi: 10.1016/j.cmet.2024.10.017.

[38] Drinkwater, K., 2023. Interleukin-11 (IL11) inhibits myogenic differentiation of C2C12 cells through activation of extracellular signal-regulated kinase (ERK). Cellular Signalling.

[39] Tisdale, M.J., 1999. Wasting in Cancer. The Journal of Nutrition 129(1): 243S-246S, Doi: 10.1093/jn/129.1.243S.

[40] Petruzzelli, M., Schweiger, M., Schreiber, R., Campos-Olivas, R., Tsoli, M., Allen, J., et al., 2014. A Switch from White to Brown Fat Increases Energy Expenditure in Cancer-Associated Cachexia. Cell Metabolism 20(3): 433–47, Doi: 10.1016/j.cmet.2014.06.011.

[41] Queiroz, A.L., Dantas, E., Ramsamooj, S., Murthy, A., Ahmed, M., Zunica, E.R.M., et al., 2022. Blocking ActRIIB and restoring appetite reverses cachexia and improves survival in mice with lung cancer. Nature Communications 13(1): 4633, Doi: 10.1038/s41467-022-32135-0.

[42] Jakobsdottir, S., Van Nieuwpoort, I.C., Van Bunderen, C.C., De Ruiter, M.B., Twisk, J.W.R., Deijen, J.B., et al., 2016. Acute and short-term effects of caloric restriction on metabolic profile and brain activation in obese, postmenopausal women. International Journal of Obesity 40(11): 1671–8, Doi: 10.1038/ijo.2016.103.

[43] Kolnes, K.J., Nilsen, E.T.F., Brufladt, S., Meadows, A.M., Jeppesen, P.B., Skattebo, Ø., et al., 2025. Effects of seven days' fasting on physical performance and metabolic adaptation during exercise in humans. Nature Communications 16(1): 122, Doi: 10.1038/s41467-024-55418-0.

[44] Consolazio, C.F., Nelson, R.A., Johnson, H.L., Matoush, L.O., Krzywicki, H.J., Isaac, G.J., 1967. Metabolic Aspects of Acute Starvation In Normal Humans: Performance and Cardiovascular Evaluation1. The American Journal of Clinical Nutrition 20(7): 684–93, Doi: 10.1093/ajcn/20.7.684.

[45] Krotkiewski, M., Grimby, G., Holm, G., Szczepanik, J., 1990. Increased muscle dynamic endurance associated with weight reduction on a very-low-calorie diet. The American Journal of Clinical Nutrition 51(3): 321–30, Doi: 10.1093/ajcn/51.3.321.

[46] Racette, S.B., Rochon, J., Uhrich, M.L., Villareal, D.T., Das, S.K., Fontana, L., et al., 2017. Effects of Two Years of Calorie Restriction on Aerobic Capacity and Muscle Strength. Medicine & Science in Sports & Exercise 49(11): 2240–9, Doi: 10.1249/MSS.0000000000001353.

[47] Brown, J.L., Rosa-Caldwell, M.E., Lee, D.E., Blackwell, T.A., Brown, L.A., Perry, R.A., et al., 2017. Mitochondrial degeneration precedes the development of muscle atrophy in progression of cancer cachexia in tumour-bearing mice: Mitochondrial degeneration precedes cancer cachexia. Journal of Cachexia, Sarcopenia and Muscle 8(6): 926–38, Doi: 10.1002/jcsm.12232.

[48] Delfinis, L.J., Bellissimo, C.A., Gandhi, S., DiBenedetto, S.N., Garibotti, M.C., Thuhan, A.K., et al., 2022. Muscle weakness precedes atrophy during cancer cachexia and is linked to muscle-specific mitochondrial stress. JCI Insight 7(24): e155147, Doi: 10.1172/jci.insight.155147.

[49] Campos, C.A., Bowen, A.J., Han, S., Wisse, B.E., Palmiter, R.D., Schwartz, M.W., 2017. Cancer-induced anorexia and malaise are mediated by CGRP neurons in the parabrachial nucleus. Nature Neuroscience 20(7): 934–42, Doi: 10.1038/nn.4574.

[50] Olson, B., Diba, P., Korzun, T., Marks, D.L., 2021. Neural Mechanisms of Cancer Cachexia. Cancers 13(16): 3990, Doi: 10.3390/cancers13163990.

[51] Zhu, X.A., Starosta, S., Ferrer, M., Hou, J., Chevy, Q., Lucantonio, F., et al., 2025. A neuroimmune circuit mediates cancer cachexia-associated apathy. Science 388(6743): eadm8857, Doi: 10.1126/science.adm8857.

[52] Katakami, N., Uchino, J., Yokoyama, T., Naito, T., Kondo, M., Yamada, K., et al., 2018. Anamorelin (ONO☐7643) for the treatment of patients with non–small cell lung cancer and cachexia: Results from a randomized, double☐blind, placebo☐controlled, multicenter study of Japanese patients (ONO☐7643☐04). Cancer 124(3): 606–16, Doi: 10.1002/cncr.31128.

[53] Takayama, K., Katakami, N., Yokoyama, T., Atagi, S., Yoshimori, K., Kagamu, H., et al., 2016. Anamorelin (ONO-7643) in Japanese patients with non-small cell lung cancer and cachexia: results of a randomized phase 2 trial. Supportive Care in Cancer 24(8): 3495–505, Doi: 10.1007/s00520-016-3144-z.

[54] Temel, J.S., Abernethy, A.P., Currow, D.C., Friend, J., Duus, E.M., Yan, Y., et al., 2016. Anamorelin in patients with non-small-cell lung cancer and cachexia (ROMANA 1 and

ROMANA 2): results from two randomised, double-blind, phase 3 trials. The Lancet Oncology 17(4): 519–31, Doi: 10.1016/S1470-2045(15)00558-6.

[55] Suriben, R., Chen, M., Higbee, J., Oeffinger, J., Ventura, R., Li, B., et al., 2020. Antibody-mediated inhibition of GDF15–GFRAL activity reverses cancer cachexia in mice. Nature Medicine 26(8): 1264–70, Doi: 10.1038/s41591-020-0945-x.

[56] Kim-Muller, J.Y., Song, L., LaCarubba Paulhus, B., Pashos, E., Li, X., Rinaldi, A., et al., 2023. GDF15 neutralization restores muscle function and physical performance in a mouse model of cancer cachexia. Cell Reports 42(1): 111947, Doi: 10.1016/j.celrep.2022.111947.

[57] Groarke, J.D., Crawford, J., Collins, S.M., Lubaczewski, S., Roeland, E.J., Naito, T., et al., 2024. Ponsegromab for the Treatment of Cancer Cachexia. New England Journal of Medicine: NEJMoa2409515, Doi: 10.1056/NEJMoa2409515.

[58] Brattain, M.G., Strobel-Stevens, J., Fine, D., Webb, M., Sarrif, A.M., 1980. Establishment of mouse colonic carcinoma cell lines with different metastatic properties. Cancer Research 40(7): 2142–6.

**Acknowledgements**

We would like to thank Dr. Andrea Bonetto (cxC26) and Dr. Nicole Beauchemin (ncxC26) for graciously providing the cell lines used for our studies. We are grateful to Dr. Tobias Janowitz, Dr. Eileen White, and Dr. Marcus Goncalves for their critical reading and valuable suggestions of the manuscript. We thank Karen E. Inouye for the technical support with the metabolic cages and EchoMRI, and Guangru Jiang for the help in fecal energy excretion measurement. This work was delivered as part of the CANCAN team supported by the Cancer Grand Challenges partnership funded by Cancer Research UK (CGCATF-2021/100034) and the National Cancer Institute (OT2CA278654).

**Author contributions**

Yanshan Liang: Conceptualization; Resources; Investigation; Data curation; Formal analysis; Visualization; Methodology; Writing—original draft; Project administration; Writing—review and editing. Young-Yon Kwon: Conceptualization; Resources; Investigation; Writing—original draft; Writing—review and editing. Sheng Hui: Conceptualization; Supervision; Funding acquisition; Investigation; Writing original draft; Project administration; Writing—review and editing.

**Disclosure and competing interests statement**

The authors declare no competing interests.

Figure 1

**Figure 2**

**a** Energy expenditure

kcal/hr

Day 6  Day 7  Day 8  Day 9  Day 10  Day 11  Day 12

**b** Cumulative energy expenditure

kcal

ncxC26 ad lib
ncxC26 isocaloric
cxC26 isocaloric

Full day    Day    Night

**c** Respiratory exchange ratio

Ratio

Day 6  Day 7  Day 8  Day 9  Day 10  Day 11  Day 12

**d** Average respiratory exchange ratio

Ratio

ncxC26 ad lib
ncxC26 isocaloric
cxC26 isocaloric

Full day    Day    Night

**Figure 3**

Figure 4

Figure 5

**Figure 7**