1 Complement receptor *C3ar1* deficiency does not alter brain
2 structure or functional connectivity across early life development

Hanna Lemmik[1], Eugene Kim[1], Eilidh MacNicol[1], Davide Maselli[2], Michel
Bernanos[1], Zhuoni Li[3], Dauda Abdullahi[1], Esther Walters[1], Maria Elisa Serrano
Navacerrada[1], Wuding Zhou[4], Aleksandar Ivetic[2], Diana Cash[1†*], Laura Westacott[5†*]

[1]Department of Neuroimaging, King's College London, UK [2]School of
Cardiovascular and Metabolic Medicine & Sciences, King's College London, UK
[3]Department of Forensic and Neurodevelopmental Sciences, King's College London,
UK [4]Peter Gorer Department of Immunobiology, King's College London, UK
[5]Neuroscience and Mental Health Innovation Institute, Cardiff University, UK

[†]These authors contributed equally

For correspondence: westacottlj@cardiff.ac.uk (LJW), diana.cash@kcl.ac.uk (DC)

13 Abstract

Genetic deletion of the complement C3a anaphylatoxin chemotactic receptor
(*C3ar1*), a key component of the innate immune response, is reported to induce
behavioural phenotypes consistent with psychiatric symptomatology in mice, but
when and where *C3ar1* is needed in the brain is unresolved. These questions are
significant because, as a G-protein-coupled receptor, human C3AR1 serves as a
potential therapeutic target for disorders associated with complement dysregulation,
such as schizophrenia. To provide a brain-wide (where) assessment of developmental
C3ar1 activity, we used longitudinal (when) tensor-based morphometry, diffusion-
weighted magnetic resonance imaging (MRI) and blood oxygen-level dependent
functional MRI in male and female *C3ar1*-deficient mice and wild-type littermates,
with behavioural assessment in adulthood. Unexpectedly, we did not find a robust
C3ar1-dependent phenotype in any of these measures. Therefore, our study does not
support neurodevelopmental hypotheses for *C3ar1*, which is encouraging for
therapeutic strategies targeting this receptor since interventions are unlikely to
disrupt brain development.

29 Introduction

The complement system is a conserved immune pathway that participates in host
defence through pathogen clearance and regulating inflammation (Chaplin 2020) as
well as tissue homeostasis (Kunz and Kemper 2021; West and Kemper 2023), with
emerging roles in neurodevelopment, psychiatric disorders and neurodegeneration
(Stevens et al. 2007; Hong et al. 2016; Sekar et al. 2016).

The most convincing evidence for the involvement of the complement system in
neurodevelopment so far is its genetic association with schizophrenia. Schizophrenia
is a complex and highly heritable neurodevelopmental disorder characterised by
hallucinations, delusions, and impaired cognition, with symptoms typically emerging
in late adolescence or early adulthood (McCutcheon, Reis Marques, and Howes
2020; Howes, Bukala, and Beck 2024). Neurobiological hallmarks of schizophrenia
include grey matter loss (Vita et al. 2012) and a reduction in synaptic density (Osimo
et al. 2019). Genome-wide association studies (GWAS) of schizophrenia have helped
to identify two complement-related risk loci; a structural variant in the complement
component 4 A (*C4A*) gene (Sekar et al. 2016), which encodes the C4A protein
responsible for propagation of complement activation, and a variant in the CUB and
Sushi Multiple Domains 1 (*CMSD1*) gene, which encodes a putative complement
inhibitor protein (Schizophrenia Working Group of the Psychiatric Genomics
Consortium 2014; Baum et al. 2024). Preclinical studies link these variants to
increased brain-specific complement activation and synapse loss (Sekar et al. 2016;
Yilmaz et al. 2021; Baum et al. 2024), potentially tying complement activation to
synaptic pathology in schizophrenia. Indeed, the *C4A* schizophrenia-risk genotype
associates with MRI markers of grey matter loss and reduced cognitive performance
in humans, even in the absence of neurological disorders (O'Connell et al. 2021).
Further, patients with elevated complement proteins have more severe negative
symptoms in psychosis (Byrne et al. 2024), which typically do not respond to anti-
psychotic medication. Complement modulation therefore has potential in
addressing this unmet therapeutic need.

C3a anaphylatoxin chemotactic receptor (C3aR1), a G-protein coupled receptor
(GPCR) bound by complement activation product C3a and the granin family
neuropeptide TLQP-21 (Rodriguez et al. 2023), acts downstream of complement
activation and stands out as a pharmacologically tractable target for modifying
complement activity in the brain (Hauser et al. 2017). In mice, *C3ar1* transcripts are
predominantly expressed by microglia, with minimal neuronal expression observed
in both healthy adult humans and mice according to single-cell transcriptomic
analyses (Hammond et al. 2019; Tasic et al. 2016). While its temporal expression
patterns are not yet well characterised, C3aR1 appears to be active during early
embryonic development, potentially influencing progenitor cell proliferation
(Coulthard et al. 2018; Bénard et al. 2008; Hammond et al. 2019). C3aR1 also
appears to facilitate developmental astrocyte phagocytosis by microglia in the retina
(Gnanaguru et al. 2023), as well as to regulate microglial reactivity and
neuroinflammation more broadly (Gedam et al. 2023; Ge, Guan, and Wang 2024;

Zheng et al. 2021; Lian et al. 2015; Vasek et al. 2016; Chew and Petretto 2019). Brain
morphological changes observed in *C3ar1*-deficient mice further support its
neurodevelopmental relevance (Westacott et al. 2021; Pozo-Rodrigálvarez et al.
2021), although there is no consensus on the precise neurodevelopmental actions of
C3aR1. Addressing this gap is important given this receptor's potential as a
pharmacological target.

C3aR1 signalling may impact brain functions relevant to psychiatric
symptomatology since a range of behavioural phenotypes have been reported in
C3ar1-deficienct mice. These include abnormal anxiety-like behaviours (Westacott
et al. 2022), hyperactivity (Pozo-Rodrigálvarez et al. 2021), cognitive impairment
(Coulthard et al. 2018) but also a resilience to depressive-like behaviours induced by
chronic stress or inflammation (Crider et al. 2018; Zhang et al. 2022; Sun et al. 2024).
Although the observed involvement of C3aR1 in behaviour suggests that it is needed
for normal brain function, previous studies have not addressed the question of
whether these phenotypes arise because of a C3aR1 deficit during development or
because it is continuously needed, which can be resolved through longitudinal
assessment. Another important yet overlooked aspect is the use of the appropriate
wild-type *littermate* control animals that none of the aforementioned studies
included. Instead, these studies used separately raised cohorts of control and mutant
mice which represents a known confound due to the rapidly diverging genetic
backgrounds in small inbred colonies (Fitch and Atchley 1985), but also because
litter environment affects behaviour and brain development (Crews et al. 2009;
Jiménez and Zylka 2021; Valiquette et al. 2023).

To investigate the potential consequences of *C3ar1* deficiency during development,
we adopted a global, unbiased approach, conducting a longitudinal study of male
and female *C3ar1*-deficient mice and their wild-type littermates during adolescence
(postnatal day, PND30) and adulthood (PND90). Using structural and diffusion
magnetic resonance imaging (MRI and dMRI), as well as resting state fMRI
(rsfMRI), we aimed to assess whether the absence of C3aR1 affects brain
development during adolescence—a critical period for psychiatric vulnerability
(Westacott and Wilkinson 2022; Paus, Keshavan, and Giedd 2008)—or whether this
requirement only becomes evident by adulthood.

Our imaging measures included regional brain volumes derived from tensor-based
morphometry (TBM), fractional anisotropy (FA) from dMRI to evaluate white
matter organisation, which is influenced by microglial activity during development
(Chan et al. 2024; Falangola et al. 2023), and graph theoretical analysis of functional

connectivity (FC) correlates of blood oxygenation level dependent (BOLD) signal,
such as global efficiency and clustering coefficient to characterise brain network
topology (Zhu et al. 2017; Forlim et al. 2024; Hadley et al. 2016). These techniques
were complemented by behavioural testing in adult mice that measured cognition
and emotional reactivity, as we sought to replicate previously reported behavioural
experiments in *C3ar1*-deficient mice (Pozo-Rodrigálvarez et al. 2021; Westacott et al.
2022; Coulthard et al. 2018). Unexpectedly, we found no robust brain or
behavioural phenotype in our datasets, challenging the previous assumptions of a
neurodevelopmental role for C3aR1 under physiological conditions.

Results

C3ar1^{tm1Cge} homozygous mice lack detectable *C3ar1* mRNA

We used an established *C3ar1* mutant line, *C3ar1*^{tm1Cge} (Humbles et al. 2000) and
performed our own validation of the mutation. For this purpose, we designed a 79-
base pair (bp) amplicon targeting the putatively deleted region (**Figure 1a**). PCR
analysis of cDNA derived from bone marrow-derived macrophages—a cell type
consistently reported to express high levels of *C3ar1* mRNA (Tao et al. 2021;
Mommert et al. 2018; Mamane et al. 2009)—showed no detectable transcript in this
region in the mutant animals (**Figure 1c**), confirming the absence of the canonical
transcript. The *C3ar1* gene contains an in-frame start codon after the deletion
(**Supplemental figure 1**). We also confirmed that no transcript is made that
includes this region (**Figure 1b**). These results were further corroborated by
quantitative PCR (qPCR, n = 8) in homozygous knockout animals in either M0-like
or interleukin 4 (IL4)-induced M2-like macrophages (**Supplemental figures 2a–e**).
Together, these results confirm that *C3ar1*^{tm1Cge} is a true loss-of-function or a
"knockout" allele resulting in no detectable *C3ar1* transcript.

Figure 1 | *C3ar1*^{tm1Cge} homozygous mice lack detectable *C3ar1* mRNA. (a)
Schematic shows *C3ar1* mRNA with its only protein coding exon and the design of
the 79 bp PCR amplicon which targets an exon-1/5' untranslated region (UTR)-
exon-2 junction of the canonical *C3ar1*$^{+/+}$ transcript. Also shown is another 372 bp
amplicon which targets a region downstream of the deletion and after an alternative
start codon. The start of exon 2 is deleted in *C3ar1*$^{-/-}$ mice, so PCR should result in
no amplification. Similarly, if no alternative transcript is present, there should be no
amplification**. (b–c)** Gel images show PCR products of cDNA from bone-marrow
derived interleukin 4 (IL4)-induced M2-like macrophages. Each sample well has a
297 bp Glyceraldehyde 3-phosphate dehydrogenase (*Gapdh*) control band. NEG: no
reverse transcriptase negative control carried over from cDNA synthesis. **(a)**
Canonical transcript. **(b)** Hypothesised alternative transcript. The bands appear
hollow due to over-saturation. -/- = *C3ar1*$^{-/-}$, +/+ = *C3ar1*$^{+/+}$.

***C3ar1*-deficiency does not influence total or regional brain volume**

We conducted a longitudinal MRI study (referred to as Cohort 1 hereafter or
implied when cohort is not specified) to investigate potential genotype-related
differences in brain structure and function using *C3ar1*^{tmCge} homozygous knockout
mice (*C3ar1*$^{-/-}$, *C3ar1*-deficient) and their littermate wild-type controls
(*C3ar1*$^{+/+}$) (**Figure 2a**) on C57BL6J (Charles River, UK) background. Both groups
underwent *in vivo* MRI in adolescence (PND27–31) and adulthood (PND81–92).
Structural MR images were additionally collected *ex vivo* from the same mice
sacrificed in adulthood immediately after the *in vivo* scan to achieve higher isotropic
resolution (0.1 mm *ex vivo* vs 0.15 mm *in vivo*) and increased signal-to-noise ratio
(SNR). To corroborate these results, we also conducted MRI in adulthood (PND74-
110) in an independent study cohort, referred to as Cohort 2.

We used tensor-based morphometry (TBM) analysis to estimate total brain volume
and to map regional brain volume differences. There were no genotype-dependent
differences in total brain volume *in vivo* in adolescence (**Figure 2b**) nor in adulthood
(**Figure 2c**). In all adolescent mice, irrespective of genotype, female mice exhibited
significantly smaller total brain volumes compared to males. These sex differences
disappeared by adulthood—a finding that aligns with previous observations in
wildtype mice (Guma et al. 2024).

Still using TBM, no significant genotype-dependent differences in regional brain
volumes were detected in adolescence (**Figure 2d-i**). In adulthood (**Figure 2d-ii**),
C3ar1$^{-/-}$ mice showed a significant ($p < 0.05$) volume increase in the right pretectal
area, and subthreshold ($0.05 < p < 0.5$) increases in the left pretectal area and the right
lateral thalamus *in vivo*, but these differences were no longer observed in the same

study cohort *ex vivo* despite improved spatial resolution (not shown since these
group-level data would be presented as empty study-template maps), nor did we
observe any genotype-dependent differences using TBM analysis in Cohort 2 (not
shown). We also did not observe any significant genotype-by-sex interaction effects
(not shown).

While no reproducibly significant genotype effects in regional volumes were
observed in Cohort 1, we nevertheless detected sexually dimorphic effects (**Figure
2d-iii** and **2d-iv**). Adolescent female mice had significantly larger volumes (relative
to total brain volume) bilaterally in the agranular insular cortex (AI), superior
colliculus (SC), medial septum (MS) and in the CA1 region of the hippocampus,
whereas male mice had increased relative volumes in white matter areas including the
olfactory tract, corpus callosum, hippocampal commissure, and notably also in the
median preoptic nucleus (MEPO) which is well known to be larger in male rodents
(Gorski et al. 1978) (**Figure 2d-iii, Supplemental figure 3** for absolute volume).
Echoing total brain volume measures, many of these sex differences were no longer
observed in adulthood (see also **Supplemental figure 4** for Cohort 2 data), but
females showed larger relative volumes in the dorsal anterior cingulate cortex
(ACAd), secondary motor cortex (MOs) and primary somatosensory cortex (SSp).
Adult males had larger relative volumes in areas including the medial amygdala
(MEA) and the bed nucleus of stria terminalis (BST) which, like the MEPO, are
previously documented sexual dimorphisms (Hines, Allen, and Gorski 1992) and
which were also observed *ex vivo* in this study cohort (**Supplemental figure 5**).

There were no genotype nor sex-by-genotype interactions on regional brain volumes
at either time-point (not shown). Overall, female somatosensory and motor cortices
increased more in volume between adolescence and adulthood than male
(**Supplemental figure 6**), in line with the observed smaller differences in these areas
in adulthood compared to adolescence.

If variability was high within our study sample, particularly in the *C3ar1*-deficient
group, it would have hampered our ability to detect significant differences. The
coefficient of variation for grey matter region-of-interest (ROI) volume showed no
differences between genotypes *in vivo* in adolescence (**Figure 2e**) or adulthood
(**Figure 2f**), nor *ex vivo* in adulthood (**Figure 2g**). The overall variability was low,
ranging from 4.2–5.5% *in vivo* and 4.2% *ex vivo*, while hippocampal variability which
has previously been repeatedly measured establishing a neuroimaging gold standard
at 5% variability (Lerch et al. 2012), was 3.7–3.8% *in vivo* and 4.2% *ex vivo* in our

study. These values suggest that our study was well-positioned to detect a genotype

effect if one had been present.

a)

| $C3ar1^{+/+}$
 n = 35: 17 M, 18 F | $C3ar1^{-/-}$
 n = 34: 18 M, 16 F |

in vivo MRI   EPM   OF   *in vivo* MRI   *ex vivo* MRI

PND   30   80   90

b) Total brain volume adolescence *in vivo*
mm³
genotype *p* val = ns
sex *p* val < 0.001, η² = 0.2 (large)
M   F

c) Total brain volume adulthood *in vivo*
mm³
genotype *p* val = ns
sex *p* val = ns
M   F

d)   TBM: difference in regional volume (%)

i) adolescence $C3ar1^{-/-}$ vs $C3ar1^{+/+}$

ii) adulthood $C3ar1^{-/-}$ vs $C3ar1^{+/+}$ PRT

iii) adolescence female vs male   SC, CA1, CC, CP, AI, MEPO, MS

iv) adulthood female vs male   PAG, DG, MPO, SSp, MEA, ACAd, MOs, BST, OB

$p < 0.05$ 20   relative volume (%)   −20   0 0.5 FWER

e) Grey matter volume variability adolescence *in vivo*
Coefficient of variation
ns
v. hipp, d. hipp
intergroup mean 5.5%
$C3ar1^{+/+}$ $C3ar1^{-/-}$

f) Grey matter volume variability adulthood *in vivo*
Coefficient of variation
ns
v. hipp, d. hipp
4.3%
$C3ar1^{+/+}$ $C3ar1^{-/-}$

g) Grey matter volume variability adulthood *ex vivo*
Coefficient of variation
ns
d. hipp, v. hipp
4.2%
$C3ar1^{+/+}$ $C3ar1^{-/-}$

Figure 2 | Sex but not *C3ar1* status influences regional brain volume. (a)
Schematic of the MRI study shown in **(d–g)**. 69 mice were scanned twice *in vivo*; in
adolescence at ~ postnatal day (PND) 30 (range 27-31) and at adulthood ~ PND90
(range 81-92), as well as once *ex vivo* after sacrifice after the adulthood scanning
session. Behavioural tests were carried out before the adulthood scanning session.
EPM = elevated plus maze, OF = open field. **(b)** Total brain volume in adolescence.
Two-way ANOVA, genotype, sex, genotype * sex, $F_{[1,65]}$ = 0.35, 16.72, 3.26, p = 0.56,
<0.001, 0.76. **(c)** Total brain volume in adulthood. Two-way ANOVA, genotype,
sex, genotype * sex, $F_{[1,65]}$ = 0.74, 0.26, 3.33, p = 0.39, 0.61, 0.07. **(b–c)** Data
presented as mean ± 95% CI. **(d)** Panels showing relative regional volume changes (%)
overlaid on study-specific coronal templates (grey). Red hues signify areas larger in
C3ar1$^{-/-}$ **(i–ii)** or females **(iii–iv)** and blue hues signify areas larger in *C3ar1$^{+/+}$* **(i–ii)**
or males **(iii–iv)**. Transparency of the colour overlay shows the statistical
significance, ranging from family wise error (FWE)-corrected p value 0.5 to 0
(transparent to opaque, respectively). Areas where FWE-corrected p value < 0.05 are
demarcated with a black line, and where the p value > 0.5 are grey (no overlay),
meaning that in adolescence genotype comparison **(i)**, no voxels had a p value < 0.5.
The locations of the coronal slices in relation to bregma in the left most column from
top: -7.6, -4.6, -1.6, 1.4 mm. *C3ar1$^{+/+}$* n = 35, 17 males and 18 females; *C3ar1$^{-/-}$* n =
34, 18 males and 16 females. ACAd = dorsal anterior cingulate cortex, AI = agranular
insular cortex, BST = bed nucleus of stria terminalis, CA = cornu ammonis, CP =
caudoputamen, DG = dentate gyrus, MEA = medial amygdala, MEPO = medial
preoptic nucleus, MOs = secondary motor cortex (MOs), MPO = medial preoptic
area, MS = medial septum, OB = olfactory bulb, PAG = periaqueductal grey, PRT =
pretectal area, SC = superior colliculus, SSp = primary somatosensory cortex. **(e)**
Coefficient of variation (CV) in adolescence *in vivo* for 72 regional volumes across all
animals in the experiment. **(f)** CV in adulthood *in vivo*. **(g)** CV in adulthood *ex vivo*.
(e–g) Data are shown with quartiles and whiskers show the extent of the
distribution. Dotted line shows the intergroup mean excluding cerebrospinal fluid
(CSF) areas. Kruskal Wallis p values, all ns. D. hipp = dorsal hippocampus, v. hipp =
ventral hippocampus.

C3ar1-deficiency does not influence fractional anisotropy

To assess the potential impact of *C3ar1* deficiency on white matter integrity, we
measured fractional anisotropy (FA) as an indirect marker of axonal microstructure
(Friedrich et al. 2020) (**Figure 3**). Using voxel-wise analysis, we observed sub-
threshold, non-significant ($0.05 < p < 0.5$) decreases in FA in *C3ar1*-deficient mice
compared to wild-types (**Figure 3a**). In adolescence, sub-threshold reductions were
noted in the corpus callosum (CC) and optic tract (OPT, **Figure 3a**, upper panel).
In adulthood, they were primarily localised to the CC (**Figure 3a**, bottom panel).
These sub-threshold differences were not observed *ex vivo*. Compared to males,
female mice had sub-threshold decreases in FA *in vivo* in adulthood in the internal

capsule and in the third ventricle (**Supplemental figure 7**), but this was again not
observed *ex vivo* (not shown). We did not observe any sex-by-genotype interactions
in our FA datasets (not shown).

In our ROI-based FA analysis, we focused on predefined white matter regions,
hypothesising that FA alterations would primarily occur in areas containing axonal
tracts due to reported increased microglial phagocytosis in *C3ar1*-deficient mice
during development (Gnanaguru et al. 2023; Falangola et al. 2023; Chan et al. 2024).
In this analysis also, no significant genotype effects were observed in adolescence
(**Figure 3b-i**) or adulthood (**Figure 3b-ii**). There were no genotype-dependent
differences in the change in FA over time, but FA increased between adolescence and
adulthood in all white matter areas except the olfactory tract (**Figure 3b-iii**), which
is in line with the reported early maturation of the olfaction system in mice
(Gretenkord et al. 2019).

Figure 3 | Fractional anisotropy does not depend on *C3ar1* status but increases with age. (a) Panels showing voxel-wise fractional anisotropy analysis in adolescence (top) and adulthood (bottom) corrected for FWE. There are no significant voxels (no black contour) where genotype effect is significant ($p < 0.05$). CC = corpus callosum, OPT = optic tract. **(b)** *In vivo* ROI-based median fractional anisotropy (FA) values in white matter regions for **(i)** adolescence, **(ii)** adulthood and **(iii)** change over time (adulthood – adolescence). Two-sided one-sample *t* test (difference from 0) *p* values that were adjusted with Benjamini-Hochberg (BH) procedure (### *p* value < 0.001). **(b-i–iii)** Mixed ANOVA with genotype and genotype-by-region interaction effects, all ns. Data are presented as mean ± 95% CI.

C3ar1-deficiency does not influence global functional brain connectivity

To evaluate global brain functional connectivity (FC), which has been reported to be altered in other mice who lack a specific microglial gene (Filipello et al. 2018; Deivasigamani et al. 2023; Zhan et al. 2014), we estimated FC through analysis of BOLD signal time-courses with the assumption that the magnitude of correlation between these time-courses corresponds to the strength of FC. We calculated pairwise correlation coefficients between 36 (18 per hemisphere) grey matter regions. Non-zero correlation values were averaged across proportional progressively decreasing sparsity thresholds to preserve biologically meaningful weak correlations while minimising noise (Bassett et al. 2009). No genotype-dependent differences in global FC were observed in adolescence (**Figure 4a**), adulthood (**Figure 4d**) or in the change over time (**Figure 4g**).

We applied graph theory to characterise global brain network connectivity across all regions, focusing on two key metrics: clustering coefficient (**Figure 4b**, **e** and **h**) and global efficiency (**Figure 4c**, **f** and **i**), and applying the same strategy for proportional thresholding as for FC. The clustering coefficient reflects the tendency of nodes to form connected local clusters, with higher values indicating the presence of more highly interconnected subnetworks within the brain (Bullmore and Sporns 2009). Global efficiency refers to the average of shortest paths linking nodes in a network and can be a proxy of information integration abilities since it decreases with cognitive deficit (Berlot et al. 2016; Hawkins et al. 2020) and increases with development (Jiang et al. 2023). Clustering coefficient and global efficiency appeared higher in *C3ar1*-deficient animals at both time-points, but this was not significant (**Supplemental table 1**). Further, we found no effects of genotype across global connectivity measures when we treated males and females as separate groups (**Table 1**, for *p* values, see **Supplemental table 2**).

FC, global efficiency and clustering coefficient appeared to increase with brain
maturation when genotype groups were combined, but this was only significant in
the case of global efficiency when graph sparsity was lower, that is, when 30–50% of
the strongest connections were retained, with small effect sizes observed (Cohen's *d*
= 0.30–0.32). These findings indicate that while *C3ar1* deficiency did not result in
alterations in global network properties under our experimental conditions,
developmental changes in global efficiency were detectable.

Figure 4 | Global functional connectivity is not changed in *C3ar1*-deficient
mice. FC, clustering coefficient and global efficiency at decreasing graph sparsity
levels in adolescence **(a–c)**, and adulthood **(d–f)**, and change over time **(g–i)**. **(a–i)**
Statistical significance was determined with sexes combined using AUC permutation
testing (10,000 iterations) and the resulting *p* values were corrected for multiplicity
with the BH method (n = 3 tests per outcome measure). Adolescence: *C3ar1*[+/+] n =
32 (15 males, 17 females); *C3ar1*[-/-] n = 32 (18 males, 14 females); Adulthood:
C3ar1[+/+] n = 33 (16 males, 17 females) and *C3ar1*[-/-] n = 32 (17 males, 15 females);

change: *C3ar1*[+/+] n = 30 (14 males and 16 females); *C3ar1*[-/-] n = 31 (17 males and 14 females). Data are shown as mean ± 95% CI. **(g–i)** Two-sided one-sample *t* tests (difference from 0) for global connectivity changes at each sparsity level with genotypes combined, corrected using the BH procedure (n = 10 sparsity levels), # = adjusted *p* value < 0.05.

Table 1 | Mean network connectivity metrics ± 95% CI in males and females of both genotypes

Age	Adolescence				Adulthood			
C3ar1 status	+/+	-/-	+/+	-/-	+/+	-/-	+/+	-/-
Sex	F	F	M	M	F	F	M	M
Connectivity metric (AUC ± 95% CI)								
FC	25.5 ± 2	27.5 ± 3.1	25.9 ± 5	29.5 ± 4.5	27.6 ± 3.3	28.9 ± 2.9	29.9 ± 5.3	31.9 ± 4.7
GE	11.5 ± 0.8	12.7 ± 1.2	11.4 ± 1.5	13.2 ± 1.7	12.17 ± 1.2	12.9 ± 1.3	13.3 ± 2	14.5 ± 2.1
CC	13.5 ± 1.1	14.9 ± 1.9	13.9 ± 2.8	15.9 ± 2.6	14.8 ± 2	15.8 ± 1.9	16.3 ± 3.2	17.4 ± 2.6

C3ar1-deficiency does not influence functional brain networks

Since *C3ar1*-deficient mice did not show statistically significant changes in global connectivity metrics, we next examined specific networks after conducting *t*-tests for each pairwise ROI FC between genotypes. For this we used two hypothesis-free approaches; false discovery rate (FDR) correction to identify strongly differing edges between genotypes, and network-based statistics (NBS) (Zalesky, Fornito, and Bullmore 2010) to detect network-level differences while controlling for family-wise error rate. Thresholding the resulting *t*-value matrices at |*t*| ≥ 2 for adulthood and adolescence, and their changes over time (adulthood – adolescence; **Figure 5a**), no edges remained significant after FDR correction. Then, using NBS, which tests the likelihood of detecting a connected component of a specific size, we determined that the network sizes observed after thresholding at |*t*| ≥ 2 (n = 68 at adolescence, n = 15 at adulthood, and n = 1 for change) could occur by chance in these datasets.

Given that neither one of the hypothesis-free approaches, FDR correction and NBS, detected any genotype-related differences, we next focused on anxiety-related

regions. This decision was motivated by prior evidence of anxiety-like behaviour in
C3ar1-deficient animals (Westacott et al. 2022) and the inclusion of anxiety-specific
tests in our behavioural battery. We calculated the mean FC, or nodal strength, of 20
a priori selected brain regions relevant to anxiety. We did not detect effects of
genotype (**Figure 5b**), sex (**Table 2**) or sex-by-genotype interaction (**Supplemental**
table 3).

Next, we examined FC and global efficiency within two resting state networks that
have been linked to anxiety and emotionality in humans (Coutinho et al. 2016; Geng
et al. 2015; Schimmelpfennig et al. 2023) and that have been observed in mice
(Grandjean et al. 2020; Sforazzini et al. 2014; Hikishima et al. 2023); the default
mode network (DMN) and the salience network (**Figure 5c**). Additionally, we
analysed network connectivity within a third, anxiety network (regions in **Figure**
5b). Consistent with our earlier findings, we did not observe genotype-dependent
differences in these networks at either time-point.

The only significant genotype-related differences were seen in voxel-wise seed-to-
brain connectivity analysis. Two anxiety-related regions, the left prefrontal cortex
and left ventral hippocampus (**Figure 5d**; **Supplemental figure 8** for the right
hemisphere), had a weak but significant and widespread higher connectivity in
C3ar1-deficient animals compared to controls at both time-points. Seed-based
analysis further revealed higher FC across the brain when other anxiety-related
regions were used as seeds, mostly in adolescence (**Supplemental figure 9**).
However, this effect was not confined to anxiety-related regions or specific networks
(**Supplemental figure 10**). Although the increase of FC in *C3ar1*-deficient mice
appears widespread, this is not robust, since our seed-based analysis controls for
voxel-wise comparisons within subjects but does not correct for testing multiple
seeds.

a)

C3ar1⁻/⁻ vs C3ar1⁺/⁺ functional connectivity *t* test
t-value matrices thresholded at |*t*| > 2

Cortex Limbic system Basal ganglia Midbrain Brain stem

adolescence

Remaining connections after FDR correction = 0

NBS *p* value = ns

adulthood

Remaining connections after FDR correction = 0

NBS *p* value = ns

adulthood-adolescence

Remaining connections after FDR correction = 0

NBS *p* value = ns

t values

b)

Nodal strength of anxiety-related regions

adolescence

ns { *C3ar1⁺/⁺*
    C3ar1⁻/⁻

adulthood

ns { *C3ar1⁺/⁺*
    C3ar1⁻/⁻

L cingulate cx
R cingulate cx
L prefrontal cx
R prefrontal cx
L amygdala
R amygdala
L pallidum / accumbens
R pallidum / accumbens
L striatum
R striatum
L hypothalamus
R hypothalamus
L dorsal hippocampus
R dorsal hippocampus
L ventral hippocampus
R ventral hippocampus
L pag
R pag
L brain stem
R brain stem

Mean FC with all other regions

c)

Network FC
Student's *t* tests with
BH correction
all ns

adolescence   adulthood

DMN   DMN

Salience   Salience

Anxiety   Anxiety

C3ar1: +/+   -/-   +/+   -/-

Network GE
Student's *t* tests with
BH correction
all ns

adolescence   adulthood

DMN   DMN

Salience   Salience

Anxiety   Anxiety

+/+   -/-   +/+   -/-

d)

C3ar1⁻/⁻ vs C3ar1⁺/⁺ seed-based FC analysis examples

adolescence

Left prefrontal cortex   Left ventral hippocampus

adulthood

Left prefrontal cortex   Left ventral hippocampus

Higher FC
in *C3ar1⁻/⁻*

contrast value

Higher FC
in *C3ar1⁺/⁺*

TFCE *p*-value
transformed (0.81 − *p*) and scaled

black line = *p*-value < 0.05

Figure 5 | *C3ar1*-deficiency does not influence functional brain networks. (a)
Thresholded adjacency matrices comparing *C3ar1*[+/+] vs *C3ar1*[-/-] groups where $|t| \geq 2$
in adolescence and adulthood, and the change between time points. No connections
with $|t| \geq 2$ remained significant after the false discovery rate (FDR) correction.
Additionally, the number of connections in the $|t| \geq 2$ adjacency matrix components
were not significant when assessed using network-based statistics (NBS) correction.
(b) Nodal strength (mean absolute connectivity) of *a priori* anxiety-related seeds (e.g.
L cingulate cortex correlation coefficients with all other regions). *P* values were
calculated using a mixed ANOVA with between-subjects factor of genotype and
within-subjects factor of region. Only the region effect was significant at both time-
points (both *p* values < 0.001, $\eta_p^2 = 0.69$ in adolescence and $\eta_p^2 = 0.73$ in adulthood).
(c) Mean functional connectivity (FC) and global efficiency (GE). Student's *t* tests
within a metric (n = 4) corrected for multiple comparisons using the BH method.
DMN = default mode network **(d)** Examples of seed-based FC maps showing voxel-
wise group differences between *C3ar1*[-/-] and *C3ar1*[+/+] mice (using *t* tests) with seeds
placed in the left ventral hippocampus and left prefrontal cortex in adolescence and
adulthood datasets. The dual scale bar displays contrast value on the x-axis and
TFCE *p*-values (transformed 0.81 – p) on the y-axis. The transformed *p*-values have
been re-scaled to range from 0 to 1 for visualisation, with darker colours representing
greater statistical significance. Black outlines demarcate regions where TFCE *p*-values
are below 0.05.

**Table 2 | Mean anxiety-associated node connectivity in males and females of
both genotypes.**

Age	Adolescence				Adulthood			
C3ar1 status	+/+	-/-	+/+	-/-	+/+	-/-	+/+	-/-
Sex	F	F	M	M	F	F	M	M
Brain area (mean node connectivity strength)								
L cingulate cx	0.39	0.41	0.39	0.44	0.45	0.49	0.48	0.54
R cingulate cx	0.38	0.41	0.38	0.43	0.45	0.49	0.48	0.53
L prefrontal cx	0.29	0.33	0.32	0.37	0.41	0.44	0.37	0.49
R prefrontal cx	0.28	0.28	0.33	0.37	0.41	0.45	0.39	0.47
L amygdala	0.14	0.17	0.15	0.2	0.2	0.2	0.24	0.28
R amygdala	0.11	0.16	0.15	0.18	0.19	0.2	0.25	0.25
L pallidum & accumbens	0.23	0.27	0.23	0.31	0.3	0.29	0.35	0.4
R pallidum & accumbens	0.24	0.26	0.24	0.3	0.31	0.31	0.35	0.39
L striatum	0.28	0.32	0.3	0.36	0.32	0.36	0.4	0.43
R striatum	0.26	0.31	0.32	0.35	0.33	0.36	0.42	0.41
L hypothalamus	0.28	0.29	0.27	0.34	0.32	0.32	0.34	0.38
R hypothalamus	0.25	0.27	0.26	0.32	0.31	0.32	0.33	0.37
L dorsal hippocampus	0.4	0.43	0.42	0.49	0.44	0.46	0.48	0.52
R dorsal hippocampus	0.41	0.43	0.42	0.49	0.44	0.47	0.5	0.52
L ventral hippocampus	0.18	0.18	0.17	0.26	0.21	0.23	0.2	0.27

R ventral hippocampus	0.18	0.23	0.19	0.25	0.22	0.25	0.25	0.26
L PAG	0.28	0.3	0.3	0.38	0.27	0.3	0.32	0.36
R PAG	0.28	0.31	0.3	0.39	0.27	0.29	0.32	0.37
L brain stem	0.11	0.11	0.12	0.18	0.1	0.14	0.16	0.16
R brain stem	0.11	0.09	0.12	0.18	0.11	0.13	0.17	0.18

C3ar1-deficient mice do not have discernible behavioural phenotypes

Behavioural testing is another way of assessing functional consequences of genetic manipulations with expected neurodevelopmental sequelae (Crawley 2007). We aimed to evaluate the impact of *C3ar1* deficiency on anxiety-like behaviour, locomotion and recognition memory, which were chosen based on prior reports of abnormalities in *C3ar1*-deficient mice. We used a battery of well-established behavioural tests, including the OF test and EPM for anxiety-like behaviour and locomotion, as well as NOR for recognition memory. We also tested PPI which is a sensorimotor reflex consistently found to be attenuated in schizophrenia (Ludewig, Geyer, and Vollenweider 2003; Mena et al. 2016) but which has hitherto not been tested in *C3ar1*-deficient mice before.

We performed behavioural testing in both cohorts described in the structural MRI results sections. In Cohort 1 (**Figure 2a**), behavioural testing (EPM, then OF) occurred shortly before the adulthood MRI scan (and two months after adolescence MRI). Behaviour of Cohort 2 was also tested in adulthood (OF, NOR, EPM, PPI, in that order) followed by MRI, except that these mice were not previously scanned under anaesthesia. Behavioural data from each cohort were analysed separately to account for differences in study design (see also Methods).

To assess multiple measures of anxiety-like behaviour and locomotion along one another (**Figure 6a–b, Supplemental table 4** for description of anxiety-like metric selection), we calculated z-scores for each behavioural outcome measure for *C3ar1*-deficient mice relative to wild-type controls (**Table 3** for untransformed means). In Cohort 1, no significant genotype effects were detected in anxiety-like behaviour or locomotion at either time-point (**Supplemental table 5**). To rule out any potential confounding effects from co-housing littermate mutants and wild-types (Kalbassi et al. 2017), we also examined whether the number of *C3ar1*-deficient cage-mates influenced wild-type behaviour but found no consistent pattern or evidence of systematic anxiety-like effects in wild-types (**Supplemental figure 11**).

In Cohort 2, most anxiety-related and locomotion measures showed no genotype differences (**Supplemental table 6**), except for reduced distance travelled in the

centre of the OF arena by *C3ar1*-deficient mice (uncorrected two-sample *t*-test $p <$
0.01, Cohen's $d = 0.92$, **Supplemental table 6**). However, no genotype effect on
distance was detected in the core of the OF arena or in the centre and core of the OF
arena in Cohort 1.

The sample size (n = 38) of Cohort 2 was too small to reliably test for interactions
between sex and genotype, meaning we only had enough statistical power to detect
very large effects (Cohen's $f = 0.5$ at alpha = 0.05 and 80% power), which were clearly
not observed across behavioural measures. No sex-by-genotype interactions were
observed in Cohort 1 (**Supplemental table 5**).

Additionally, PPI testing (performed exclusively on Cohort 2) showed no effect of
C3ar1 deletion, though PPI increased with prepulse intensity as expected (**Figure**
6c), confirming that the test was set up to reliably measure this phenomenon. NOR
testing (also performed exclusively on Cohort 2) revealed no genotype differences,
with all groups demonstrating successful learning based on recognition indices
significantly above chance levels (**Figure 6d**).

Overall, these findings suggest that *C3ar1* deficiency does not result in robust
anxiety-like or hyperactive phenotypes, nor deficits in recognition memory or PPI.

Figure 6 | *C3ar1* deficiency does not cause behavioural abnormalities. (a) Z-
scored (normalised to *C3ar1*$^{+/+}$ control mean = 0) locomotion metrics for *C3ar1*$^{-/-}$

animals from EPM and OF tests, showing results from Cohort 1 and 2. Two-way

ANOVA for Cohort 1 (4 groups, males and females separated) or Student's *t* tests

for Cohort 2 (2 groups, males and females combined due to smaller sample size)

where parametric assumptions were met, otherwise the Kruskal Wallis test (with

Dunn in Cohort 1); uncorrected for multiplicity, all ns. **(b)** Same as (a) but for

anxiety-like metrics from EPM and OF tests; *p* values all ns. **(c)** Prepulse inhibition in

Cohort 2. PPI increased with increasing prepulse intensity in both, *C3ar1*[+/+] vs

C3ar1[-/-] mice (paired *t* tests p value ### < 0.001). Dotted line at 0 indicates inhibition

threshold. A score above 0 indicates inhibition (shaded area). Slope plot shows means

and 95% confidence intervals. Individual mice are plotted three times at increasing

prepulse intensity. Prepulse inhibition did not differ by genotype at any prepulse

intensity (uncorrected Student's *t* tests at 3 dB, 6 dB and 12 dB, $t_{[37]}$ = 1.58, 1.99,

0.28, *p* = 0.12, 0.54, 0.17). **(d)** Novel object recognition (NOR) recall 1 hr after

acquisition in Cohort 2. Both groups showed novelty preference (one-sample *t* test,

value > 50/chance, # = *p* < 0.05, ## = *p* < 0.01 ### = *p* < 0.001). There were no

differences between groups (Student's *t* test $t_{[37]}$ = 0.42, *p* = 0.67). The dotted line

marks the chance threshold. **(a-d)** Cohort 1: *C3ar1*[-/-] n = 33 (17 male, 16 female),

C3ar1[+/+] n = 33 (16 male, 17 female), Cohort 2: *C3ar1*[-/-] n = 20, *C3ar1*[+/+] n = 19

(males and females combined). Data are expressed as mean ± 95% CI.

Table 3 | Mean behavioural outcome measures ± 95% CI in males and females

of both genotypes

Cohort	Cohort 1				Cohort 2			
Sex	F		M		F		M	
C3ar1 status	+/+	-/-	+/+	-/-	+/+	-/-	+/+	-/-
Locomotion-related								
OF locomotion speed (mm/s)	79.65± 3.47	79.48± 3.91	75.61± 4.31	74.72± 4.08	94.21± 6.64	84.33± 7.26	84.93± 7.55	84.93± 5.68
OF locomotion duration (s)	213.38 ±22.53	223.73 ±26.85	216.60 ±21.94	217.97 ±20.82	315.66 ±50.93	264.33 ±49.90	297.85 ±39.56	283.61 ±17.86
OF locomotion distance (cm)	1777.8 9±210. 25	1847.7 1±271. 41	1746.4 5±257. 47	1747.5 2±225. 53	2984.9 9±589. 73	2231.4 5±473. 78	2562.3 6±515. 37	2412.0 3±268. 13
EPM speed (mm/s)	67.07± 3.43	65.89± 2.40	67.71± 3.74	69.90± 2.60	75.27± 5.08	75.38± 6.40	73.61± 4.21	76.63± 3.76
EPM locomotion duration (s)	69.25± 15.36	61.22± 11.89	73.29± 13.14	67.01± 7.48	78.93± 13.23	82.53± 18.41	71.11± 17.69	62.51± 8.88
EPM locomotion distance (cm)	473.32 ±124.8 7	401.52 ±88.01	502.39 ±109.2 2	465.16 ±54.60	591.26 ±122.2 0	620.88 ±155.6 9	518.89 ±140.0 6	472.57 ±68.28
Anxiety-related								
OF duration core (s)	25.92± 12.12	18.74± 4.60	19.81± 5.35	22.99± 7.73	25.06± 10.65	29.96± 24.36	32.19± 18.93	16.71± 3.87
OF duration centre (s)	106.94 ±18.40	102.01 ±16.36	97.18± 13.68	109.40 ±17.99	105.42 ±40.78	120.94 ±39.65	133.80 ±31.20	95.62± 13.10

OF duration periphery (s)	493.13 ±18.41	498.07 ±16.36	502.90 ±13.68	490.66 ±17.98	493.16 ±42.58	478.89 ±39.60	466.11 ±31.25	504.15 ±13.10
EPM duration open (s)	29.52± 11.52	28.92± 12.57	28.06± 12.03	28.19± 14.72	18.87± 8.77	20.45± 16.56	15.13± 7.96	12.15± 7.29
OF latency core (s)	30.66± 17.23	44.47± 34.34	29.89± 16.18	40.95± 19.45	34.19± 36.91	24.15± 20.50	25.86± 25.45	29.44± 52.19
OF latency centre (s)	10.02± 3.52	13.26± 8.07	12.35± 7.40	11.97± 4.95	5.79±6 .00	7.45±7 .79	5.22±5 .01	5.28±4 .26
EPM duration middle (s)	109.53 ±17.89	102.17 ±19.15	112.33 ±24.24	122.79 ±17.27	85.38± 21.11	72.59± 29.02	71.13± 14.54	74.93± 17.85
EPM duration closed (s)	176.11 ±22.18	198.69 ±19.25	188.78 ±24.17	178.29 ±17.22	214.24 ±21.04	207.31 ±38.82	227.88 ±14.86	224.17 ±17.79
EPM bouts open (#)	4.65±1 .92	3.19±1 .66	3.44±1 .33	2.94±1 .18	2.62±1 .48	3.43±2 .77	2.30±1 .12	1.91±1 .49
EPM head dips (#)	21.53± 6.20	18.12± 5.17	19.62± 6.29	22.31± 5.36	16.75± 4.55	17.57± 13.16	11.90± 3.14	16.73± 8.52
EPM stretch-attend postures (#)	19.00± 4.15	21.81± 2.98	23.62± 3.14	23.38± 2.87	16.62± 6.16	9.43±6 .04	13.50± 2.68	14.73± 4.10
EPM latency open (s)	65.31± 27.63	106.76 ±53.38	92.95± 48.81	117.27 ±51.75	75.66± 81.38	83.26± 103.55	95.40± 71.29	137.52 ±70.66
Other								
PPI 3 dB (%)					21.11± 13.2	15.31± 6.40	22.87± 7.41	17.14± 7.70
PPI 6 dB (%)					40.63± 10.16	26.36± 8.74	39.49± 11.24	34.37± 10.65
PPI 12 dB (%)					56.81± 13.18	49.81± 13.28	57.07± 9.51	60.40± 8.38
Acoustic startle response at 120 dB (A.U.)					876.35 ±432.3 6	1068.3 2±284. 31	1472.9 9±360. 84	1908.7 9±655. 66
NOR recognition index (%)					62.49± 11.20	62.60± 9.77	65.02± 7.66	68.16± 8.87
NOR total exploration training (s)					39.45± 15.62	38.58± 10.66	38.50± 12.16	41.12± 11.07
NOR total exploration test (s)					31.68± 10.07	32.41± 6.83	30.81± 9.17	33.58± 5.39

Discussion

Here we used longitudinal neuroimaging and behavioural testing to investigate the

impacts of *C3ar1* deletion on brain structure and function across early life

development in mice. We found no robust evidence that *C3ar1* deficiency affects

total or regional brain volume, white matter FA, global FC, global efficiency,

clustering coefficient, or specific functional networks.

We also found no altered behavioural phenotype in either male or female *C3ar1*-
deficient mice in adulthood. These results raise questions regarding the source of
discrepancies between these data, our previous behavioural work (Westacott et al.
2022; Westacott et al. 2021), and that of others (Crider et al. 2018; Coulthard et al.
2018; Pozo-Rodrigálvarez et al. 2021; Sun et al. 2024), which may be attributed to
our use of littermate controls in this study, as will be elaborated upon in the
following sections. Overall, the lack of a behavioural phenotype combined with the
paucity of genotype effects on brain structure across time-points means that the role
of C3aR1 in neural development and behaviour is not as significant as originally
thought, at least in the absence of exacerbating environmental or immune triggers.

Absence of *C3ar1*-dependent effects on brain structure

Our morphometric analysis showed that total and regional brain volumes were not
changed between *C3ar1*-deficient and wild-type mice. We were, however, able to
detect sex-specific neurodevelopmental changes that have previously been reported
in rodents, such as brain growth occurring later in females, minimal total and
regional brain volume differences in adulthood and larger volume of MEPO and
BNST in males (Qiu et al. 2018; Guma et al. 2024; Gorski et al. 1978; Hines, Allen,
and Gorski 1992; de Courten-Myers 1999), supporting the sensitivity of our
methodology. Together with the low variability in our sample, our ability to detect
these known sex differences means that we were well-positioned to identify potential
genotype effects had they been present.

Absence of *C3ar1*-dependent effects on white matter fractional anisotropy

C3ar1 is predominantly expressed by microglia in the brain (Quell et al. 2017; Tasic
et al. 2018), where it influences their reactivity and phagocytosis (Vasek et al. 2016;
Gedam et al. 2023; Zheng et al. 2021; Lian et al. 2015; Gnanaguru et al. 2023).
Notably, changes in microglial characteristics have been linked to developmental
alterations in FA in both mice and humans (Falangola et al. 2023; Chan et al. 2024).
We found no *C3ar1*-dependent changes in the white matter despite showing the
overall effect of time on the brain maturation on FA, which is known to increase
during development between adolescence and adulthood in both humans and mice
(Brouwer et al. 2012; Hagmann et al. 2010; Reynolds et al. 2019; Piekarski et al.
2023; Chahboune et al. 2007). Lack of a phenotype after the deletion of a
predominantly microglial gene may not be surprising since mice that completely lack
microglia for their entire lifespan (achieved by deleting the super enhancer for
macrophage colony-stimulating factor receptor, *Csf1r*) do not have overt
neurodevelopmental phenotypes (Rojo et al. 2019), and only show vulnerability in a

pathological context involving neuroinflammation (Kiani Shabestari et al. 2022).
This raises the possibility that neurodevelopmental phenotypes associated with
C3ar1 deficiency may likewise only become apparent under neuroinflammatory
conditions.

Widespread, weak seed-to-brain functional connectivity increase in *C3ar1*-deficient mice in adolescence

While we did not detect genotype-dependent global FC changes nor changes to
clustering coefficient and global efficiency at either time-point, we did observe a
genotype-independent developmental increase in global efficiency with time, which
has been reported before (Jiang et al. 2023; Hagmann et al. 2010; Koenis et al. 2018),
particularly at lower sparsities (Cai, Dong, and Niu 2018). This additionally suggests
that our method was sensitive enough to pick up developmentally relevant effects.

Similarly to whole brain analyses, we did not observe any genotype-dependent
changes to specific brain networks, including anxiety networks (**Figure 5**), which we
hypothesised to be affected based on our previous behavioural results (Westacott et
al. 2022). However, our anxiety network connectivity null findings are internally
consistent with the absence of an anxiety-like phenotype in the cohorts tested in this
study. Combined with a recent null report for an anxiety-like effect in the EPM in
C3ar1-deficient animals by another group (Sun et al. 2024), our data indicate that
the effect of *C3ar1* on anxiety-networks and anxiety-like behaviour is context-
dependent if present.

Although we did not detect network-specific effects, *C3ar1*-deficient mice exhibited
subtle but widespread increases in resting-state FC in voxel-wise seed-based analyses
during adolescence across nearly all seeds examined, with a weaker effect persisting
into adulthood (**Figure 5** and **Supplemental figures 8-10**). These increases in
connectivity may reflect developmental alterations in circuit properties caused by the
absence of C3aR1. Similar findings have been reported for other microglial
receptors—such as TREM2, CX3CR1, and CR3—where genetic deficiency altered
adult brain FC (Filipello et al. 2018; Zhan et al. 2014; Deivasigamani et al. 2023). In
these reports, TREM2 or CX3CR1 knockouts led to impaired synapse elimination
alongside decreased FC, accompanied by social behaviour deficits and increased
repetitive behaviour (Filipello et al. 2018; Zhan et al. 2014). In contrast, CR3
knockout mice had no deficits in synapse or axon refinement but showed reduced
phagocytosis of perinatal cortical neurons and higher cortical FC (Deivasigamani et
al. 2023). It is indeed possible that similarly to CR3, C3aR1 is also involved in

perinatal neuron phagocytosis so that its deletion results in higher number of
neurons in early embryonic development and a weak brain-wide increase in FC,
particularly as *C3ar1*-deficiency leads to a deficit of developmental astrocyte
phagocytosis in the retina (Gnanaguru et al. 2023). For now, we urge caution with
this interpretation for two reasons that should be considered together. These
connectivity changes were isolated findings–that is, they were not accompanied by
any structural alterations, which would be unexpected if a larger number of neurons
were present, nor were they associated with any behavioural changes. Second, the
voxel-wise seed-based analysis is the least robust presented here, as the observed
changes were largely transient, weak, not confined to a specific subnetwork, and not
corrected for multiple comparisons across seeds. Future studies investigating FC in
C3ar1-deficient mice should incorporate repeated measures from the same
individual during adolescence to improve robustness as well as assessing synaptic
transmission more directly, for example by electrophysiological assessment in brain
slices, although it is still unclear which brain regions should be targeted currently
with the latter approach.

Behavioural outcomes were unaffected by *C3ar1* deficiency

Neither anxiety-like behaviours nor locomotor activity showed significant
differences between genotypes across multiple testing paradigms, apart from a small
decrease in ambulation of *C3ar1*-deficient mice in the aversive central zone of the OF
arena, which in the absence of a locomotor phenotype could index anxiety-like
behaviour (Prut and Belzung 2003). This latter change, however, was not
reproducible across study cohorts and was no longer observed in the same study in
the core of the OF arena. Tests for recognition memory and sensorimotor gating,
NOR and PPI tests, also yielded null results.

Differences between our current findings and previous reports of anxiety-like
behaviours (Westacott et al. 2022), hyperactivity (Pozo-Rodrigálvarez et al. 2021)
and cognitive deficits (Coulthard et al. 2018) may stem from variations in
experimental design, particularly environmental factors such as lighting and testing
time. Our behavioural experiments were conducted during the active phase (7 PM–
11 PM, lights off) to minimise stress from sleep deprivation, which is known to
upregulate complement in the brain (Tillmon et al. 2024; Crider et al. 2018; Li et al.
2023; Tripathi et al. 2021). It is possible that previous studies, which tested animals
during their inactive phase, introduced a stress-dependent "second hit" to *C3ar1*
deficiency. However, this seems unlikely since inhibiting complement, including
C3ar1, has been shown to improve resilience to chronic stress by preventing

anhedonia-like behaviour and neuroinflammation (Li et al. 2023; Crider et al. 2018; Tripathi et al. 2021; Madeshiya et al. 2022; Tillmon et al. 2024).

Unlike previous studies of *C3ar1*-deficient mice, our study employed littermate control animals, meaning that wild-type mice were co-housed with their *C3ar1*-deficient siblings. This approach minimises potential environmental differences between groups, which is especially important in neurobehavioral studies where subtle environmental factors can significantly affect outcomes. While the use of littermate controls is widely regarded as best practice in such contexts (Holmdahl and Malissen 2012; Valiquette et al. 2023; Bailey, Rustay, and Crawley 2006), it is not without its own potential confounds. For instance, abnormal behaviours exhibited by genetically altered animals can influence the behaviour of co-housed wild-type mice, particularly when the phenotype is pronounced, such as in cases of increased aggression (Kalbassi et al. 2017). However, as aggression has not been reported in *C3ar1*-deficient mice, and their previously described behavioural phenotype was not notably severe, we considered such an influence unlikely in our study, especially as the anxiety-like behaviour of wild-type mice did not seem to depend on the number of *C3ar1*-deficient cage-mates.

The complement system plays an essential role in normal pregnancy and parturition (Girardi et al. 2020). A notable aspect of using a littermate design is that litters of mixed genotype are born from exclusively heterozygous parents, whereas in most prior studies of *C3ar1*-deficiency, both the mother and offspring were homozygous knockouts. This raises the possibility that behavioural effects in *C3ar1*-deficient mice may have resulted from altered intrauterine environments or care by *C3ar1*-deficient mothers. In line with this, the inbred homozygous *C3ar1* knockout colony used in our previous studies showed increased pre-weaning deaths and maternal cannibalism (unpublished data). If this is indeed the reason for the previously observed adult behavioural phenotypes, two scenarios should be considered. In the first scenario, the *C3ar1*-deficient offspring are *uniquely sensitive* to C3aR1-dependent *in utero* conditions or maternal care deficits. In the second scenario, the adult phenotypes are not specific to *C3ar1*-deficiency meaning that wild-type mice would have been similarly affected. These scenarios can be tested by using pup transfer and *in-vitro* fertilisation experiments. For now, we were able to avoid measuring these non-specific effects on behaviour by using a littermate design.

The genetic background of mutants is another factor that varies between laboratories and is known to affect phenotypic expression. For example, heterozygous knockout of the autism-associated gene *CHD8* has varying effects in 33 sub-strains (Tabbaa,

Knoll, and Levitt 2023), mirroring heterogeneity observed in human *CHD8*-haploinsufficiency. While that study identified significant variability across sub-strains, these profiles consistently differed from that of wild-type littermate controls. In contrast, our study found no robust genotype-dependent differences in the brain globally and across measures or in behaviour, apart from a single internally non-reproducible behavioural outcome measure, meaning that the effect of C3aR1 on behaviour would have to be entirely dependent on modifying genes or environmental factors.

Finally, genetic drift could have precluded reproducibility of previously reported phenotypes. Mice accumulate spontaneous mutations that rapidly reach homozygosity within small colonies, speeding up drift, a phenomenon recognised as early as the 1980s (Fitch and Atchley 1985). So far, our current work is the first to study *C3ar1* deficiency in behaviour using a littermate design, which minimises the number of genetic loci that differ between the mutant and control mice, all of which could influence measured phenotypes.

Future directions

Aside from the role of C3aR1 in brain structure and function, many aspects of the basic biology of C3aR1 remain unclear, including its expression pattern in the healthy brain, with little information available on the cell types where it is expressed nor its expression contexts and time-points. It is also unclear which G-proteins C3aR1 signals through in different brain cell types. Future studies could integrate toolkits like TRUPATH, a suite of Gαβγ biosensors for analysing G-protein coupling preferences (Olsen et al. 2020), with transcriptomics to address these questions. This approach would be particularly valuable, as it could enable the use of chemogenetics to activate the same G-protein as C3aR1 in specific cell types and contexts to study its function.

Most importantly, we propose that *C3ar1*-deficiency in development should be studied after an immune challenge. Unlike humans, laboratory mice live in an immune-privileged environment, hence investigating the consequences of a combined immune insult with the genetic deficit would be relevant; or better yet, in the context of *C4A* overexpression, which is a known genetic risk factor for schizophrenia (Sekar et al. 2016) and is known to impact white matter integrity (Caseras et al. 2024).

Limitations

In this study, we provided high-level global data on *C3ar1*-deficient mice. While our study had notable strengths, including the use of a littermate design and the inclusion of female animals, it is not an exhaustive characterisation of the mutant.

While we measured brain region volumes, we cannot definitively address more reductionist questions, such as cellular composition which would have required single cell RNASeq or antibody staining and microscopy. Similarly, for axonal integrity analysis for which we used the proxy of dMRI-derived FA, the gold standard would have been electron microscopy. However, a more granular approach would have necessitated a trade-off with throughput—something that is hard to justify in the absence of strong hypotheses regarding the developmental expression pattern of *C3ar1*. Additionally, our fMRI analysis relied on a specific parcellation of the brain consisting of 36 regions, and different parcellations could yield varying results (Thirion et al. 2014; Zalesky et al. 2010). We encourage others exploring our datasets to therefore experiment with alternative parcellations.

Conclusion

Contrary to expectations, we found no evidence for *C3ar1*-dependent effects across imaging measures at either time-point, nor could we replicate previously seen behavioural phenotypes. The absence of detectable phenotypes suggests that C3aR1 plays a minimal role in brain structure and function, at least in the absence of an immune trigger. These findings challenge prior assumptions about its neurodevelopmental significance and necessitate further investigation into C3aR1 function on sensitized backgrounds, such as those involving immune activation.

Methods and materials

Animals

All animal procedures complied with the UK Animals and Scientific Procedures Act 1986 and were approved by the local ethical committee at King's College London (KCL). Homozygous *C3ar1*$^{-/-}$ mice were generated by homologous recombination in embryonic stem cells and kindly provided by Dr. Bao Lu and Prof. Craig Gerard (Harvard Medical School, Boston, MA) (Humbles et al. 2000). These mice were subsequently backcrossed onto the C57BL/6J strain for at least 12 generations and maintained on a C57BL/6J background in Professor Wuding Zhou's laboratory at KCL. For this study, cryopreserved stocks were rederived at KCL and crossed to

C57BL/6J mice purchased from Charles River to refresh the genetic background
following Jackson's Laboratories line refreshing protocol.

Experimental animals (*C3ar1*$^{-/-}$ and *C3ar1*$^{+/+}$ littermates) were generated through
heterozygote incrosses and resulting genotypes followed Mendelian ratios. The
heterozygote breeders were generated by outcrossing heterozygous mice to bought
wild-type Charles River C57BL/6Js. The breeders used to produce the experimental
animals were derived either from the first, second or third of these outcrosses. Sibling
crosses were not conducted, and parental age was between 2–4 months to minimise
genetic drift.

Experimental mice were housed in individually ventilated cages under controlled
temperature (20–25°C), humidity (50–60%), and a 12-hour light-dark cycle (lights
on at 7:00 AM, lights off at 7:00 PM). Environmental enrichment included nesting
materials, tunnels, and chew sticks. Mice had *ad libitum* access to irradiated rodent
chow and autoclaved water. Animals were group-housed (2–4 mice per cage), with
males and females housed separately after weaning (PND21±2). Genotyping was
conducted on ear biopsy DNA by Transnetyx using probes targeting the neomycin
cassette for the mutant allele and intron 1 for the wild-type allele. No mismatches
were identified through double-genotyping 20% of the study cohorts.

Validation of mutation
Bone marrow-derived macrophages were obtained from tibias and femurs of eight
three-month-old mice (n = 4 *C3ar1*$^{+/+}$, n = 4 *C3ar1*$^{-/-}$) following standard protocols.
Bone marrow suspension was filtered through a 40 μm mesh, centrifuged at 450 x g
for 5 minutes at 4°C, and treated with NH_4Cl haemolysis buffer (NH_4Cl 0.15M,
$KHCO_3$ 0.01M, EDTA 0.0001M). After a second centrifugation under the same
conditions, cells were washed with PBS and resuspended in Gibco RPMI 1640
Medium (Thermo Fisher, #21875-034) supplemented with 50 ng/ml recombinant
mouse macrophage colony-stimulating factor (M-CSF, R&D Systems, #416-ML-
010/CF), 1% penicillin, 1% streptomycin, and 10% heat-inactivated foetal bovine
serum (Sigma, #F9665-50ml).

Cells were seeded at 1x10^6 cells/ml in six-well plates (six wells per animal) and
incubated at 37°C with 5% CO_2 for 72 hours. On day three, the medium was
refreshed, and 50 ng/ml recombinant mouse IL4 (R&D Systems, #404-ML-
010/CF) was added to half of the wells to skew them towards M2 phenotype.
Incubation continued for an additional 48–72 hours, depending on cell confluence.

RNA was extracted using the ReliaPrep™ miRNA Cell and Tissue Miniprep
System (Promega, #Z6211) according to the manufacturer's instructions. RNA
concentration was determined with a NanoDrop spectrophotometer (Thermo
Scientific, NanoDrop 2000), yielding values between 25.5 ng/µl and 227.5 ng/µl.
Reverse transcription PCR (RT-PCR) was performed using the LunaScript® RT
SuperMix Kit (NEB, #E3010L), following the manufacturer's instructions.
Depending on RNA yield, either 100 or 500 ng of RNA was used per reaction.

For cDNA PCR and gel electrophoresis, we used GoTaq® G2 Master Mix (NEB,
#M7822). We amplified a 79 basepair (bp) fragment in the deleted region alongside a
372 bp region in the *C3ar1* cDNA that was located outside the deleted region and
downstream of an alternative start codon identified through Benchling. *Gapdh*
primers were included in each reaction to confirm amplification efficiency. PCR
products were analysed on a 2% agarose gel stained with GelRed.

For qPCR, we amplified the previously mentioned 79 bp fragment in the deleted
region. Samples were analysed in triplicate on 96-well plates (Applied Biosystems,
#4346906) with Luna® Universal qPCR Master Mix (NEB, #M3003L), and
readings were obtained using an Applied Biosystems StepOnePlus plate reader. The
amplification data were processed with the $^{\Delta\Delta}$Ct method, normalising against the
housekeeping gene Hypoxanthine phosphoribosyltransferase 1 (*Hprt*). Genotypes
were arranged alternately across the plate to minimise bias. To verify M2-like
polarisation, the expression of M2-specific markers Arginase 1 (*Arg1*) and the
mannose receptor Cluster of differentiation 206 (*Cd206*) were assessed.

Table 4 | Primer sequences

Primer name	Forward	Reverse	Application
C3ar1	AGGATTTGTTGGTG GCTCGCA	CTCCATGGCTCAGTC AAGCACA	PCR and qPCR deleted region
C3ar1	GCTTCCTGGTGCCG TTTTTC	AGTTGGTAGAGTGCG TGAGC	PCR of putative alternative transcript (3' end of exon 2 after alternative start codon)
Hprt	AGTCCCAGCGTCGT GATTAGCG	TTGAGCACACAGAGG GCCACAA	qPCR housekeeping gene
Arg1	GGCTTGCGAGACGT AGACCC	GTCCAGCCCGTCGAC ATCAAA	qPCR verification of M2 polarisation
Cd206	CCGGAGGGTGCAGA CAAAGG	TCGTCCACAGTCCAC CGAAAC	qPCR verification of M2 polarisation

Gapdh	CCTAGACAAAATGG TGAA	GACTCCACGACATAC TCAGC	PCR amplification efficiency control

Table 5 | PCR reaction

Amount (µl)	Reagent
12.5	GoTaq® G2 Master Mix (Green)
1	*C3ar1* forward primer, 10 mM
1	*C3ar1* reverse primer, 10 mM
1	*Gapdh* forward primer, 10 mM
1	*Gapdh* reverse primer, 10 mM
7.5	H₂O
1	Template DNA
Final volume: 25 µl	

Table 6 | Cycling conditions

Step	Time	Temperature
1. Initial denaturation	5 min	94°C
2. Denaturation	30 sec	94°C
3. Annealing	30 sec	60°C
4. Extension	1 min	72°C
5. Repeat steps 2-4	Repeat 35 x	
6. Final extension	5 min	72°C

Study design

This study used two separate cohorts of male and female *C3ar1*-deficient and

littermate wild-type mice. The main, longitudinal MRI cohort, termed Cohort 1,

had *in vivo* MRI performed in adolescence (range 27–31 days) and adulthood (range

81–92 days), and the adulthood MRI scan was preceded by OF and EPM tests. The

adolescence time-point was chosen because mice reach puberty approximately

between PND24–34 (Semaan and Kauffman 2015; Brust, Schindler, and

Lewejohann 2015; Pintér et al. 2007). *Ex vivo* imaging was conducted in Cohort 1's

perfusion-fixed brains after the final adulthood scan.

For Cohort 2, behavioural testing was conducted similarly to Cohort 1 in adulthood

only (range 74–110), and consisted of OF, NOR, EPM and PPI followed by *in vivo*

structural and diffusion MRI (note that MRI was not conducted in adolescence in

this cohort). For both cohorts, behavioural testing was conducted 2–7 days before

the adulthood scan.

The sample size of Cohort 1 was statistically powered to detect medium effect sizes in
regional volume using TBM across four groups (males and females analysed
separately), with a minimum sample size of n = 15 per group based on previously
observed variance with this method by our group (Serrano et al. 2023). Cohort 2
sample size was powered to detect medium effect sizes in regional TBM with sexes
combined, n = 16 per group.

In vivo **MRI**

Two to three days after behavioural testing, mice were imaged using a Bruker
BioSpec 9.4 T scanner with an 86-mm volume resonator for transmission and a 4-
channel surface array coil. Anaesthesia was induced with 4% isoflurane in medical air
(1 L/min) and oxygen (0.4 L/min), maintained at 2% but adjusted based on
respiration rates. For BOLD fMRI in Cohort 1, we used a medetomidine and
isoflurane anaesthesia optimised for mouse fMRI (Grandjean et al. 2014). This
consisted of a subcutaneous medetomidine bolus (0.05 mg/kg) followed ten minutes
later by its continuous infusion (0.1 mg/kg/h), with isoflurane levels gradually
reduced to 0.45-0.65% over 15 minutes from the start of the infusion. BOLD fMRI
was conducted after the structural scans, which took a further 45-60 minutes after
reducing isoflurane level. The respiration rate was monitored with a pressure sensor,
and temperature was monitored with a rectal thermometer and maintained at 36–
37°C using a water circulation system.

Ex vivo **MRI**

Following the adulthood *in vivo* scan in Cohort 1, mice were perfused transcardially
with 20 mL phosphate-buffered saline (PBS) followed by 4% paraformaldehyde
(PFA). Heads were stored in PFA for 48 hours, then transferred to PBS containing
0.05% sodium azide and 2 mM gadolinium-based contrast agent (Gd-DO3A-butrol).
Brains were scanned *in cranio* in groups of four using a custom-made holder
immersed in perfluoropolyether (Galden®, Solvay).

MRI acquisition parameters

For *in vivo* imaging, we first acquired an Actual Flip Angle Imaging (AFI) sequence
for B1 mapping. Then we acquired three types of 3D multi-gradient-echo images
that were used for creating study-specific templates: magnetization-transfer weighted
(MTw), proton-density weighted (PDw), and T1-weighted (T1w). Subsequently,
T2-weighted Rapid Acquisition with Relaxation Enhancement (RARE) images were
obtained. Diffusion-weighted images were then acquired using a single-shot spin-
echo echo planar imaging (EPI) sequence. Finally, for the longitudinal MRI study,

BOLD rsfMRI data were acquired using a single-shot gradient-echo EPI sequence
with 720 repetitions. Additionally, spin-echo EPI image pairs with opposing phase-
encoding polarity were recorded to enable correction of susceptibility-induced
distortions. The *in vivo* scanning session lasted 1-1.5 hours.

For *ex vivo* morphometric analysis, 3D T2-weighted images were acquired using
RARE sequences, with a total scan duration of 1 hour and 5 minutes. *Ex vivo*
diffusion-weighted images were obtained using Stejskal-Tanner pulsed gradient spin-
echo sequences with a 3D segmented EPI readout. Three b0 images were collected at
the beginning of three blocks of 30 diffusion-weighted images. The total scan time
for this acquisition was 14 hours and 15 minutes.

Table 7 | MRI acquisition parameters

Image type	TR (ms)	TE (ms)	Flip angle (°)	Averages	Bandwidth (kHz)	FOV	Matrix	Other
AFI	20/ 100	2.85	55	1	25	16.2 x 16.2 x 9	42 x 42 x 24	
MTw	24	2.5	6	2	100	16.2 x 16.2 x 9	108 × 108 × 60	6 echoes with 2.1 ms spacing; MT pulse: gaussian, 4 ms, amplitude 10 μT, offset -3 kHz, bandwidth 685 Hz
PDw	20	2.5	4	2	100	16.2 x 16.2 x 9	108 × 108 × 60	7 echoes with 2.1 ms spacing
T1w	20	2.5	20	2	100	16.2 x 16.2 x 9	108 × 108 × 60	7 echoes with 2.1 ms spacing
T2w in vivo	5000	42	90/ 180	4	50	16 x 12	128 × 96	32 slices, slice thickness 0.5, RARE factor 8
T2w ex vivo	300	30	90/ 180	1	50	25 x 24 x 18	250 x 240 x 180	RARE factor 4, scan time = 1 h 5 min

DWI in vivo	3000	21	90/180	2	357	19.2 x12	96 x 60	Single shot spin-echo planar imaging, 30 slices, slice thickness 0.5 mm, three diffusion shells: b = 350 s/mm² with 9 directions, b = 1000 s/mm² with 34 directions, b = 2000 s/mm² with 78 directions (δ = 3 ms, Δ = 11 ms), 3 b0 images per shell
DWI ex vivo	300	28.5	90/180		300	25 × 24 × 18	200 × 192 × 144	Stejskal-Tanner pulsed gradient spin echo sequences with a 3D segmented EPI readout. 12 segments and a total of 90 diffusion-weighted images acquired at a b-value of 4000 s/mm² (δ = 4 ms, Δ = 13 ms)
BOLD fMRI	1000	15	55	1	200	20 x 20	64 x 64	Single-shot gradient-echo EPI sequence, 16 slices, slice thickness 0.5, 720 repetitions

Structural MR image processing

For preprocessing MTw, T1w, and PDw images, de-ringing was conducted using the MRtrix3's (Tournier et al. 2019) `mrdegibbs` command. MTw, T1w, and PDw images were averaged across echo times, rigidly co-registered using Advanced Normalization Tools (Avants et al. 2011) `antsRegistration`, and used for template construction (see below).

For DWI, MRtrix3 `dwidenoise` was used for de-noising, MRtrix3 `mrdegibbs` for de-ringing, and FSL's (Jenkinson et al. 2012; Smith et al. 2004) `topup` and `eddy` for susceptibility and eddy-current distortion and motion correction. FSL's `dtifit` was used for diffusion tensor imaging (DTI) model fitting, enabling the calculation of

809 fractional anisotropy (FA), mean diffusivity (MD) and axial diffusivity (AD), the
810 latter two which are not reported in this manuscript for brevity. For a more detailed
811 diffusion processing protocol see (Kim et al. 2023).

Study templates

The `antsMultivariateTemplateConstruction2.sh` script from ANTs was used to
create study specific templates from processed images. For *in vivo* scans of Cohort 1,
MTw, T1w, R2* map (generated from the multi-gradient-echo PDw, T1w and
MTw images using the `qi mpm_r2s` command in the QUIT package), S0, FA and
MD images were used. For the *ex vivo* scans of Cohort 1, separate T2w and DTI
(comprising S0, FA, and MD images) templates were created. For Cohort 2, PDw,
T1w, MTw, S0 (estimated non-diffusion-weighted image from `dtifit`), FA, and
MD images were used.

Jacobian determinant maps

To estimate volume, Jacobian determinant maps were generated from the
deformation fields corresponding to the transformation of each subject to the study
template using the `CreateJacobianDeterminantImage` command from ANTs. The
Jacobian determinant values of all voxels within the template brain mask were
summed to obtain the total brain volume of each subject. Jacobian determinants
were calculated from the combined rigid, affine, and Symmetric Normalization
(SyN) transforms as well as from only the SyN transforms to obtain maps of absolute
and relative volume (accounting for differences in global brain volume), respectively.
For TBM, the Jacobian determinants were subsequently log-transformed.

Voxel-wise analysis

For voxel-wise statistics, FSL randomise with permutation testing was used (10,000
for Cohort 1, 5000 iterations for Cohort 2) followed by a threshold-free cluster
enhancement (TFCE) and family-wise error (FWE) correction as described in (Wood
et al. 2016; Kim et al. 2023). Given the absence of genotype differences in total brain
volume, TBM regional volumes are reported relative to total brain size for greater
accuracy (Lerch et al. 2012), while absolute volume maps are reported in the
Supplementary files where sex differences in total brain volume were present.

Common coordinate space

The study template was registered to the Allen Mouse Brain Common Coordinate Framework (Wang et al. 2020) using ANTs, and the Allen atlas was subsequently transformed to the study template space with the inverse transform.

ROI-based analysis of volume

The images were segmented using an in house modified version of the Allen atlas of 72 regions (Serrano et al. 2023; Wang et al. 2020). These segmentations were subsequently used to compute regional volumes by summing the Jacobian determinants within each parcellation.

Regional volume variability was estimated by calculating a coefficient of variation (CV) for each region with normalised root-mean-square method for each genotype in each experiment, using the calculation: $CV = \frac{\sigma}{\mu}$, where σ is the standard deviation and μ the population mean for each region in the atlas (n = 72 regions). Group differences in CV were calculated with a Kruskal-Wallis test (SciPy.stats, `kruskal`).

Fractional anisotropy

For mass-univariate voxel-wise analysis of fractional anisotropy, values from `dtifit` for Cohort 1 were again analysed with FSL randomise with permutation testing (10,000 permutations) followed by TFCE and FWE-correction. FA is not reported for Cohort 2 in this manuscript for brevity. For ROI-based analysis, voxel FA medians within parcellation were used for all white matter regions.

To calculate the change over time in fractional anisotropy in the longitudinal study, adolescence values for each voxel value or regional median were subtracted from adulthood values. Differences between genotypes were calculated with a mixed ANOVA with between-subjects factor of genotype and within-subjects factor of region. Change from 0 was calculated with a two-sided one-sample t-test (SciPy.stats, `ttest_1samp`), which was corrected with the Benjamini–Hochberg method.

BOLD fMRI pre-processing

Images were largely pre-processed using the Analysis of Functional NeuroImages (AFNI) toolkit. Slice timing correction was performed using the `3dTshift` package, despiking with `3dDespike`, and motion correction was applied with `3dvolreg`. The motion-corrected time average was then registered to each subject's own T2w image using ANTs, followed by registration to a template T2w image derived from a

separate mouse study conducted at the BRAIN Centre (KCL). The use of this

external template was justified by the low resolution of fMRI, which does not

benefit from creating a study-specific template.

The images were also distortion corrected using FSL's `topup`, with the distortion

estimated using an auxiliary pair of spin-echo images with opposite phase encoding

polarity and otherwise matching acquisition parameters to the gradient echo EPI

used for the BOLD signal. Prior to analysis, corrected images were band-pass filtered

at 0.01–0.2 Hz with AFNI's `3dTproject` to remove low-frequency scanner drift

noise and high-frequency physiological noise. Nuisance variables (movement and

CSF signal) were also simultaneously regressed out of the signal at this stage. Finally,

spatial smoothing was applied using AFNI's `3dBlurInMask` with a FWHM kernel.

Functional connectivity and graph theory analysis

For functional analysis, a high-level parcellation scheme was applied to segment 36

unilateral regions (18 per hemisphere) excluding white matter from the acquired 3D

volume. The BOLD signal time-courses were averaged within each ROI, and the

mean time-courses were extracted using FSL's `fslmeants` tool. Pearson correlation

coefficients were calculated for each time-course pair, producing a 36×36 correlation

matrix for each subject. These matrices were analysed as functional connectivity (FC)

graphs, with edge strength determined by the Fisher z transformed Pearson

correlation coefficient. To identify the strongest connections, graphs were

thresholded at 5% intervals from 5% to 50%. At each threshold level, referred to as the

graph sparsity interval, connections below threshold were set to zero, generating 10

sparsity graphs per subject. FC was calculated as the average non-zero connectivity at

each threshold level. Global graph metrics, global efficiency and clustering

coefficient, were computed at each sparsity level using Brain Connectivity Toolbox

algorithms (Rubinov and Sporns 2010) implemented with Network X (3.4.2)

`global_efficiency` and `clustering` passing binary thresholded Pearson matrices to

prioritise topology in the presence of noise. For global efficiency and clustering

coefficient, random curves were calculated by shuffling the thresholded binary

matrix positions. The area under the curve (AUC) was computed per subject using

trapezoidal numerical integration (`numpy.trapz`), and the likelihood of observed

values was estimated with permutation testing (10,000 permutations).

To estimate changes over time, global graph metric values at adolescence were

subtracted from adulthood values at each sparsity interval. Group differences were

again tested by calculating the AUC and applying permutation testing. Changes

from baseline (zero) were evaluated using two-sided one-sample t-tests (difference

from 0), corrected for multiple comparisons with the Benjamini-Hochberg (BH)

method.

FDR correction and network-based statistics

Matrices for *C3ar1*-deficient and wild-type mice were compared with Student's t-

tests for each pairwise connection, resulting in 36×36 t-statistic and p-value matrices.

To control for multiple comparisons, FDR correction was applied to the upper

triangle of the p-value matrix using the BH procedure

(`statsmodels.stats.multitest.fdrcorrection`, $\alpha = 0.05$).

For network-based statistics (NBS), the above-described t matrices were thresholded

at $|t| \geq 2$. To identify connected components within the FC graph, adjacency

matrices representing significant connections ($|t| \geq 2$) were converted into graph

objects using NetworkX. Regions of interest (ROIs) were treated as nodes, and

significant connections as edges. Connected components were identified using a

breadth-first search (BFS) algorithm, which explores all neighbouring nodes before

moving deeper into the graph. Only components containing more than one ROI

were retained for further analysis. To generate a null distribution of maximal

component sizes, group labels were randomly shuffled across subjects for each

permutation while preserving matrix structure (10,000 permutation). The p-value

for an observed component was calculated as the proportion of permutations where

the maximal component size exceeded that of the observed component.

***A priori* node strength analysis**

We selected 20 (10 left +10 right) anxiety and fear related regions (cingulate cortex,

prefrontal cortex, amygdala, pallidum and nucleus accumbens, striatum,

hypothalamus, dorsal hippocampus, ventral hippocampus, periaqueductal grey,

brain stem) and calculated their average absolute connectivity to all other regions

using Fisher z-transformed Pearson correlation coefficients. We then used a mixed

ANOVA with between-subjects factor of genotype and within-subjects factor of

region followed by pairwise testing with BH FDR p value correction.

Within-network analysis

Two mouse resting state networks were subset from correlation matrices: the default

mode network (DMN) and the salience network (Grandjean et al. 2020; Sforazzini et

al. 2014). The DMN included bilateral prefrontal cortex, cingulate cortex, and dorsal

hippocampus, while the salience network comprised bilateral cingulate cortex,

amygdala, and striatum. Additionally, a third anxiety-related network was defined,
consisting of the 20 regions described in the node-strength analysis above.

For each network, mean FC was calculated as the average of all pairwise connections
between nodes within the network, without applying a threshold. For individual
network global efficiency analysis, thresholding was not used due to the small
number of nodes. Instead, Pearson matrices normalised to range from 0-1 were
passed to bctpy (0.6.1) `efficiency_wei` which calculated (not binary) efficiency.

Student's t tests were used to test significance, and the comparisons were corrected
within outcome measure with the BH method.

Voxel-wise seed-to-brain analysis

We conducted voxel-wise seed-based FC analyses for anxiety-related regions, as well
as the colliculus and sensory cortex, which served as control regions not specific to
anxiety. For each seed region, the time-course of the BOLD signal was extracted and
regressed with the BOLD signal of every voxel in the brain, resulting in a 3D spatial
map of the connectivity with the seed. Group-level comparisons of these maps were
performed between genotypes using voxel-wise permutation tests with FSL's
randomise (5000 permutation), converted with TFCE and statistical significance
corrected for multiple comparisons using FWE (seed-to-brain), but they were not
corrected for the presence of multiple seeds.

General behavioural procedures

For Cohort 1, EPM was administered as the first test, followed by OF. For Cohort 2,
OF was the first test, followed by NOR test after two days of low light habituation (4
lux), EPM, and PPI.

Handling of the mice began 2–3 days prior to the behavioural testing battery. By the
start of the experiments, the mice sat comfortably on the experimenter's hand. Mice
were handled using cardboard tunnels to minimise stress, and tail handling was
avoided.

Behavioural tests were conducted during the dark phase (between 7:00 PM and
11:00 PM) of the light-dark cycle to align with the active period of mice. Mice were
randomised by genotype and counter-balanced by sex, with the experimenter
systematically blinded to genotype throughout testing and analysis though the
allocation of a study ID and test order ID, respectively.

For cleaning of the test apparatus, we used 70% EtOH for all arenas except for the
EPM arena for which we used Virusolve (Amity International) to avoid damage to
the material. Behaviour was recorded using a Google Pixel 5a camera at 1080p/60
fps. Videos were then cropped and down-sampled using FFmpeg.

Open field test

978 Mice were placed in a 40×40×40 cm white arena and allowed to explore freely. For
979 Cohort 1, dim red light (4 lux) was used, while for Cohort 2, the OF test was
980 conducted under bright overhead lighting (500 lux). 10-minute videos were analysed
981 using CleverSys (VA, USA). Outcome parameters included time spent in the centre,
982 total distance travelled, velocity, and thigmotaxis (edge exploration).

Novel object recognition

Cohort 2 mice were initially habituated to the arena under low light conditions (4
lux) over two days, five minutes per session. On the training day, two identical
objects were introduced, and mice were allowed to explore for five minutes. After a
one-hour delay, one object was replaced with a novel one. The videos were analysed
using CleverSys software stereotypic event "Sniffing" module. Objects were
manually outlined with the polygon tool. An interaction with an object was recorded
when the mouse's nose was within 5 mm of the object. The novelty preference was
determined by calculating the proportion of time spent exploring the novel object
relative to the total exploration time of both objects, with a recognition index chance
level of 50%.

Elevated plus maze

For Cohort 1, EPM was administered to behaviourally naïve animals, while for
Cohort 2, it was conducted after OF and NOR tests. In both cases, mice were placed
in the closed arm of the arena (65×65×55 cm, elevated 40 cm) under full overhead
lighting (500 lux) and allowed to explore for five minutes. The arena was divided into
closed, middle, and open areas for analysis with CleverSys software. The number of
head dips and stretch-attend postures was recorded using BORIS software, and
testing accuracy was compared with the results of an independent scorer.

Prepulse inhibition

Like NOR, PPI was only conducted in Cohort 2. We used an acoustic startle
chamber (SR-LAB, San Diego Instruments, San Diego, CA, USA) with a cylindrical
Plexiglas enclosure horizontally mounted on a mobile platform within a sound-

proofed isolation chamber. A high-frequency loudspeaker positioned above the
enclosure emitted continuous background noise at 65 dB, along with the
experimental acoustic stimuli. The startle response was recorded by converting
Plexiglas enclosure vibrations into millivolt signals using a piezoelectric unit.

Each session started with a five-minute acclimatisation period to the 65 dB
background noise, followed by five startle-alone trials at 120 dB to ensure
habituation. Mice then received 10 prepulse stimuli at 3, 6, and 12 dB above
background, each preceding a 120 dB pulse in a pseudo-randomised order,
interspersed with 10 no-stimulus trials and 10 startle-alone trials. The session
concluded with five final startle-alone pulses. The inter-trial interval was randomised
between 9-15 seconds to prevent expectation-based modulation of the startle
response. %PPI was calculated for each prepulse using the formula: $\%PPI =$
$\dfrac{pulse\ alone\ (mV_{Max}) - pulse\ with\ prepulse\ (mV_{Max})}{pulse\ alone\ (mV_{Max})}$.

Statistical procedure for behavioural outcome measures

All statistical analyses were conducted using Python 3.11.7, using relevant libraries
such as SciPy and statsmodels. For both studies, the Shapiro-Wilk test was used to
assess the normality of data distributions, while Levene's test was used to evaluate
homogeneity of variances. When either assumption of normality or equal variance
was violated, the Kruskal-Wallis test was used, followed by Dunn's test for post hoc
pairwise comparisons with Bonferroni correction for multiple testing where
applicable. For datasets meeting parametric assumptions, different approaches were
used based on the study design. For Cohort 1, a two-way ANOVA was performed to
analyse the effect of sex, genotype and sex-by-genotype interaction, followed by
Tukey's test for post hoc comparisons. Bonferroni correction was applied to adjust
for multiple comparisons. Due to the relatively small sample size (n = 38) in Cohort
2, sex-by-genotype interactions were not analysed. Instead, Student's *t*-test was used
for comparisons between groups.

To balance statistical rigor with preserving power in exploratory contexts,
behavioural measures were not universally corrected for multiplicity. Instead,
robustness was inferred through replication across independent cohorts and broad
phenotypic consistency. The threshold for statistical significance was set at $p < 0.05$.
Individual mice served as the experimental units in all analyses.

Code availability

Data and code to reproduce figures will be added to
https://github.com/hannalemmik upon publication. The MRI processing code is
available upon request from Eugene Kim.

Data availability

Behaviour videos will be added to figshare, and MRI data will be added to
openneuro. The intermediary analysis files will be added to github at
https://github.com/hannalemmik.

Acknowledgements

We thank Bao Lu and Craig Gerard for providing the *C3ar1* knockout mice, Marija
M. Petrinovic for providing prepulse inhibition testing equipment, KCL Biological
Service Unit staff for animal care and François Kroll for comments on this
manuscript. This work was funded by the MRC grant "Complement C3aR in
adolescent synaptic pruning and risk for anxiety" (MR/W004607/1). Hanna
Lemmik was funded by the Wellcome Trust as part of the "Neuro-Immune
Interactions in Health & Disease" Wellcome Trust PhD Programme
(218452/Z/19/Z).

Author contributions

HL - writing - original draft, writing - review & editing, conceptualization,
methodology, software, formal analysis, data curation, investigation, visualization,
funding acquisition; EK - methodology, software, formal analysis, investigation, data
curation, writing - review & editing, visualization, funding acquisition; EM -
methodology, software, investigation, writing - review & editing; DM - validation,
methodology, investigation; MB - software, formal analysis, data curation,
visualization; ZL - investigation, validation; DA - investigation, Esther - investigation;
MES - funding acquisition; WZ - resources, supervision; AI - resources, supervision;
DC - writing - original draft, writing - review & editing, supervision, project
administration, funding acquisition, conceptualization, methodology; LW - writing -
original draft, writing - review & editing, supervision, funding acquisition,
conceptualization, methodology.

References

Avants, Brian B., Nicholas J. Tustison, Gang Song, Philip A. Cook, Arno Klein, and James C. Gee. 2011. "A Reproducible Evaluation of ANTs Similarity Metric Performance in Brain Image Registration." *NeuroImage* 54 (3): 2033–44.

Bailey, Kathleen R., Nathan R. Rustay, and Jacqueline N. Crawley. 2006. "Behavioral Phenotyping of Transgenic and Knockout Mice: Practical Concerns and Potential Pitfalls." *ILAR Journal* 47 (2): 124–31.

Bassett, Danielle S., Edward T. Bullmore, Andreas Meyer-Lindenberg, José A. Apud, Daniel R. Weinberger, and Richard Coppola. 2009. "Cognitive Fitness of Cost-Efficient Brain Functional Networks." *Proceedings of the National Academy of Sciences of the United States of America* 106 (28): 11747–52.

Baum, Matthew L., Daniel K. Wilton, Rachel G. Fox, Alanna Carey, Yu-Han H. Hsu, Ruilong Hu, Henna J. Jäntti, et al. 2024. "CSMD1 Regulates Brain Complement Activity and Circuit Development." *Brain, Behavior, and Immunity*, March.

Bénard, Magalie, Emilie Raoult, David Vaudry, Jérome Leprince, Anthony Falluel-Morel, Bruno J. Gonzalez, Ludovic Galas, Hubert Vaudry, and Marc Fontaine. 2008. "Role of Complement Anaphylatoxin Receptors (C3aR, C5aR) in the Development of the Rat Cerebellum." *Molecular Immunology* 45 (14): 3767–74.

Berlot, Rok, Claudia Metzler-Baddeley, M. Arfan Ikram, Derek K. Jones, and Michael J. O'Sullivan. 2016. "Global Efficiency of Structural Networks Mediates Cognitive Control in Mild Cognitive Impairment." *Frontiers in Aging Neuroscience* 8 (December): 292.

Brouwer, Rachel M., René C. W. Mandl, Hugo G. Schnack, Inge L. C. van Soelen, G. Caroline van Baal, Jiska S. Peper, René S. Kahn, Dorret I. Boomsma, and H. E. Hulshoff Pol. 2012. "White Matter Development in Early Puberty: A Longitudinal Volumetric and Diffusion Tensor Imaging Twin Study." *PloS One* 7 (4): e32316.

Brust, Vera, Philipp M. Schindler, and Lars Lewejohann. 2015. "Lifetime Development of Behavioural Phenotype in the House Mouse (Mus Musculus)." *Frontiers in Zoology* 12 Suppl 1 (S1): S17.

Bullmore, Ed, and Olaf Sporns. 2009. "Complex Brain Networks: Graph Theoretical
Analysis of Structural and Functional Systems." *Nature Reviews. Neuroscience* 10 (3):
186–98.

Byrne, Jonah F., Colm Healy, Melanie Föcking, Meike Heurich, Subash Raj Susai,
David Mongan, Kieran Wynne, et al. 2024. "Plasma Complement and Coagulation
Proteins as Prognostic Factors of Negative Symptoms: An Analysis of the NAPLS 2
and 3 Studies." *Brain, Behavior, and Immunity* 119 (July): 188–96.

Cai, Lin, Qi Dong, and Haijing Niu. 2018. "The Development of Functional
Network Organization in Early Childhood and Early Adolescence: A Resting-State
FNIRS Study." *Developmental Cognitive Neuroscience* 30 (April): 223–35.

Caseras, Xavier, Emily Simmonds, Antonio F. Pardiñas, Richard Anney, Sophie E.
Legge, James T. R. Walters, Neil A. Harrison, Michael C. O'Donovan, and
Valentina Escott-Price. 2024. "Common Risk Alleles for Schizophrenia within the
Major Histocompatibility Complex Predict White Matter Microstructure."
Translational Psychiatry 14 (1): 194.

Chahboune, Halima, Laura R. Ment, William B. Stewart, Xiaoxian Ma, Douglas L.
Rothman, and Fahmeed Hyder. 2007. "Neurodevelopment of C57B/L6 Mouse
Brain Assessed by in Vivo Diffusion Tensor Imaging." *NMR in Biomedicine* 20 (3):
375–82.

Chan, Shi Yu, Eamon Fitzgerald, Zhen Ming Ngoh, Janice Lee, Jasmine Chuah,
Joanne S. M. Chia, Marielle V. Fortier, et al. 2024. "Examining the Associations
between Microglia Genetic Capacity, Early Life Exposures and White Matter
Development at the Level of the Individual." *Brain, Behavior, and Immunity* 119
(July): 781–91.

Chaplin, Hugh, Jr. 2020. "Review: The Burgeoning History of the Complement
System 1888-2005." *Immunohematology / American Red Cross* 21 (3): 85–93.

Chew, Gabriel, and Enrico Petretto. 2019. "Transcriptional Networks of Microglia
in Alzheimer's Disease and Insights into Pathogenesis." *Genes* 10 (10).

Coulthard, L. G., O. A. Hawksworth, J. Conroy, J. D. Lee, and T. M. Woodruff.
2018. "Complement C3a Receptor Modulates Embryonic Neural Progenitor Cell
Proliferation and Cognitive Performance." *Molecular Immunology* 101 (September):
176–81.

Courten-Myers, G. M. de. 1999. "The Human Cerebral Cortex: Gender Differences
in Structure and Function." *Journal of Neuropathology and Experimental Neurology*
58 (3): 217–26.

Coutinho, Joana Fernandes, Sara Veiga Fernandesl, José Miguel Soares, Liliana Maia,
Óscar Filipe Gonçalves, and Adriana Sampaio. 2016. "Default Mode Network
Dissociation in Depressive and Anxiety States." *Brain Imaging and Behavior* 10 (1):
147–57.

Crawley, J. N. 2007. "What's Wrong with My Mouse?: Behavioral Phenotyping of
Transgenic and Knockout Mice." John Wiley & Sons.

Crews, David, David Rushworth, Francisco Gonzalez-Lima, and Sonoko Ogawa.
2009. "Litter Environment Affects Behavior and Brain Metabolic Activity of Adult
Knockout Mice." *Frontiers in Behavioral Neuroscience* 3 (August): 12.

Crider, A., T. Feng, C. D. Pandya, T. Davis, A. Nair, A. O. Ahmed, B. Baban, D.
Turecki, and A. Pillai. 2018. "Complement Component 3a Receptor Deficiency
Attenuates Chronic Stress-Induced Monocyte Infiltration and Depressive-like
Behavior." *Brain, Behavior, and Immunity* 70 (May): 246–56.

Deivasigamani, Senthilkumar, Mariya T. Miteva, Silvia Natale, Daniel Gutierrez-
Barragan, Bernadette Basilico, Silvia Di Angelantonio, Laetitia Weinhard, et al. 2023.
"Microglia Complement Signaling Promotes Neuronal Elimination and Normal
Brain Functional Connectivity." *Cerebral Cortex* 33 (21): 10750–60.

Falangola, Maria Fatima, Siddhartha Dhiman, Joshua Voltin, and Jens H. Jensen.
2023. "Quantitative Microglia Morphological Features Correlate with Diffusion
MRI in 2-Month-Old 3xTg-AD Mice." *Magnetic Resonance Imaging* 103
(November): 8–17.

Filipello, Fabia, Raffaella Morini, Irene Corradini, Valerio Zerbi, Alice Canzi,
Bernadeta Michalski, Marco Erreni, et al. 2018. "The Microglial Innate Immune
Receptor TREM2 Is Required for Synapse Elimination and Normal Brain
Connectivity." *Immunity* 48 (5): 979-991.e8.

Fitch, W. M., and W. R. Atchley. 1985. "Evolution in Inbred Strains of Mice
Appears Rapid." *Science (New York, N.Y.)* 228 (4704): 1169–75.

Forlim, Caroline Garcia, Leonie Klock, Jürgen Gallinat, and Simone Kühn. 2024. "Altered Resting-State Functional Connectivity in a Thalamo-Cortico-Cerebellar Network in Patients with Schizophrenia." *Scientific Reports* 14 (1): 26284.

Friedrich, Patrick, Christoph Fraenz, Caroline Schlüter, Sebastian Ocklenburg, Burkhard Mädler, Onur Güntürkün, and Erhan Genç. 2020. "The Relationship between Axon Density, Myelination, and Fractional Anisotropy in the Human Corpus Callosum." *Cerebral Cortex (New York, N.Y.: 1991)* 30 (4): 2042–56.

Ge, Tong-Qi, Pei-Pei Guan, and Pu Wang. 2024. "Complement 3a Induces the Synapse Loss via C3aR in Mitochondria-Dependent NLRP3 Activating Mechanisms during the Development and Progression of Alzheimer's Disease." *Neuroscience and Biobehavioral Reviews* 165 (105868): 105868.

Gedam, Manasee, Michele M. Comerota, Nicholas E. Propson, Tao Chen, Feng Jin, Meng C. Wang, and Hui Zheng. 2023. "Complement C3aR Depletion Reverses HIF-1α-Induced Metabolic Impairment and Enhances Microglial Response to Aβ Pathology." *The Journal of Clinical Investigation* 133 (12).

Geng, Haiyang, Xuebing Li, Jie Chen, Xinying Li, and Ruolei Gu. 2015. "Decreased Intra- and Inter-Salience Network Functional Connectivity Is Related to Trait Anxiety in Adolescents." *Frontiers in Behavioral Neuroscience* 9: 350.

Girardi, Guillermina, Joshua J. Lingo, Sherry D. Fleming, and Jean F. Regal. 2020. "Essential Role of Complement in Pregnancy: From Implantation to Parturition and Beyond." *Frontiers in Immunology* 11 (July): 1681.

Gnanaguru, Gopalan, Steven J. Tabor, Gracia M. Bonilla, Ruslan Sadreyev, Kentaro Yuda, Jörg Köhl, and Kip M. Connor. 2023. "Microglia Refine Developing Retinal Astrocytic and Vascular Networks through the Complement C3/C3aR Axis." *Development* 150 (5).

Gorski, R. A., J. H. Gordon, J. E. Shryne, and A. M. Southam. 1978. "Evidence for a Morphological Sex Difference within the Medial Preoptic Area of the Rat Brain." *Brain Research* 148 (2): 333–46.

Grandjean, J., C. Canella, C. Anckaerts, C. Ayrancı, S. Bougacha, T. Bienert, D. Buehlmann, et al. 2020. "Common Functional Networks in the Mouse Brain Revealed by Multi-Centre Resting-State FMRI Analysis." *NeuroImage* 205 (January).

Grandjean, Joanes, Aileen Schroeter, Imene Batata, and Markus Rudin. 2014. "Optimization of Anesthesia Protocol for Resting-State FMRI in Mice Based on Differential Effects of Anesthetics on Functional Connectivity Patterns." *NeuroImage* 102 Pt 2 (November): 838–47.

Gretenkord, Sabine, Johanna K. Kostka, Henrike Hartung, Katja Watznauer, David Fleck, Angélica Minier-Toribio, Marc Spehr, and Ileana L. Hanganu-Opatz. 2019. "Coordinated Electrical Activity in the Olfactory Bulb Gates the Oscillatory Entrainment of Entorhinal Networks in Neonatal Mice." *PLoS Biology* 17 (1): e2006994.

Guma, Elisa, Antoine Beauchamp, Siyuan Liu, Elizabeth Levitis, Jacob Ellegood, Linh Pham, Rogier B. Mars, Armin Raznahan, and Jason P. Lerch. 2024. "Comparative Neuroimaging of Sex Differences in Human and Mouse Brain Anatomy." *ELife* 13 (March).

Hadley, Jennifer Ann, Nina Vanessa Kraguljac, David Matthew White, Lawrence Ver Hoef, Janell Tabora, and Adrienne Carol Lahti. 2016. "Change in Brain Network Topology as a Function of Treatment Response in Schizophrenia: A Longitudinal Resting-State FMRI Study Using Graph Theory." *Npj Schizophrenia* 2 (1): 16014.

Hagmann, P., O. Sporns, N. Madan, L. Cammoun, R. Pienaar, V. J. Wedeen, R. Meuli, J-P Thiran, and P. E. Grant. 2010. "White Matter Maturation Reshapes Structural Connectivity in the Late Developing Human Brain." *Proceedings of the National Academy of Sciences of the United States of America* 107 (44): 19067–72.

Hammond, Timothy R., Connor Dufort, Lasse Dissing-Olesen, Stefanie Giera, Adam Young, Alec Wysoker, Alec J. Walker, et al. 2019. "Single-Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes." *Immunity* 50 (1): 253-271.e6.

Hauser, Alexander S., Misty M. Attwood, Mathias Rask-Andersen, Helgi B. Schiöth, and David E. Gloriam. 2017. "Trends in GPCR Drug Discovery: New Agents, Targets and Indications." *Nature Reviews. Drug Discovery* 16 (12): 829–42.

Hawkins, R., A. S. Shatil, L. Lee, A. Sengupta, L. Zhang, S. Morrow, and R. I. Aviv. 2020. "Reduced Global Efficiency and Random Network Features in Patients with Relapsing-Remitting Multiple Sclerosis with Cognitive Impairment." *AJNR. American Journal of Neuroradiology* 41 (3): 449–55.

Hikishima, Keigo, Tomokazu Tsurugizawa, Kazumi Kasahara, Ryo Takagi, Kiyoshi
Yoshinaka, and Naotaka Nitta. 2023. "Brain-Wide Mapping of Resting-State
Networks in Mice Using High-Frame Rate Functional Ultrasound." *NeuroImage*
279 (120297): 120297.

Hines, M., L. S. Allen, and R. A. Gorski. 1992. "Sex Differences in Subregions of the
Medial Nucleus of the Amygdala and the Bed Nucleus of the Stria Terminalis of the
Rat." *Brain Research* 579 (2): 321–26.

Holmdahl, Rikard, and Bernard Malissen. 2012. "The Need for Littermate
Controls." *European Journal of Immunology* 42 (1): 45–47.

Hong, Soyon, Victoria F. Beja-Glasser, Bianca M. Nfonoyim, Arnaud Frouin,
Shaomin Li, Saranya Ramakrishnan, Katherine M. Merry, et al. 2016. "Complement
and Microglia Mediate Early Synapse Loss in Alzheimer Mouse Models." *Science* 352
(6286): 712–16.

Howes, Oliver D., Bernard R. Bukala, and Katherine Beck. 2024. "Schizophrenia:
From Neurochemistry to Circuits, Symptoms and Treatments." *Nature Reviews.*
Neurology 20 (1): 22–35.

Humbles, Alison A., Bao Lu, Christy A. Nilsson, Craig Lilly, Elliot Israel, Yuko
Fujiwara, Norma P. Gerard, and Craig Gerard. 2000. "A Role for the C3a
Anaphylatoxin Receptor in the Effector Phase of Asthma." *Nature* 406 (6799): 998–
1001.

Jenkinson, Mark, Christian F. Beckmann, Timothy E. J. Behrens, Mark W.
Woolrich, and Stephen M. Smith. 2012. "FSL." *NeuroImage* 62 (2): 782–90.

Jiang, Weixiong, Zhen Zhou, Guoshi Li, Weiyan Yin, Zhengwang Wu, Li Wang,
Maryam Ghanbari, et al. 2023. "Mapping the Evolution of Regional Brain Network
Efficiency and Its Association with Cognitive Abilities during the First Twenty-
Eight Months of Life." *Developmental Cognitive Neuroscience* 63 (101284): 101284.

Jiménez, Jessica A., and Mark J. Zylka. 2021. "Controlling Litter Effects to Enhance
Rigor and Reproducibility with Rodent Models of Neurodevelopmental Disorders."
Journal of Neurodevelopmental Disorders 13 (1): 2.

Kalbassi, Shireene, Sven O. Bachmann, Ellen Cross, Victoria H. Roberton, and
Stéphane J. Baudouin. 2017. "Male and Female Mice Lacking Neuroligin-3 Modify
the Behavior of Their Wild-Type Littermates." *ENeuro* 4 (4).

Kiani Shabestari, Sepideh, Samuel Morabito, Emma Pascal Danhash, Amanda McQuade, Jessica Ramirez Sanchez, Emily Miyoshi, Jean Paul Chadarevian, et al. 2022. "Absence of Microglia Promotes Diverse Pathologies and Early Lethality in Alzheimer's Disease Mice." *Cell Reports* 39 (11): 110961.

Kim, Eugene, Ines Carreira Figueiredo, Camilla Simmons, Karen Randall, Loreto Rojo Gonzalez, Tobias Wood, Brigida Ranieri, et al. 2023. "Mapping Acute Neuroinflammation in Vivo with Diffusion-MRI in Rats given a Systemic Lipopolysaccharide Challenge." *Brain, Behavior, and Immunity* 113 (July): 289–301.

Koenis, Marinka M. G., Rachel M. Brouwer, Suzanne C. Swagerman, Inge L. C. van Soelen, Dorret I. Boomsma, and Hilleke E. Hulshoff Pol. 2018. "Association between Structural Brain Network Efficiency and Intelligence Increases during Adolescence." *Human Brain Mapping* 39 (2): 822–36.

Kunz, Natalia, and Claudia Kemper. 2021. "Complement Has Brains-Do Intracellular Complement and Immunometabolism Cooperate in Tissue Homeostasis and Behavior?" *Frontiers in Immunology* 12 (February): 629986.

Lerch, Jason P., Lisa Gazdzinski, Jurgen Germann, John G. Sled, R. Mark Henkelman, and Brian J. Nieman. 2012. "Wanted Dead or Alive? The Tradeoff between in-Vivo versus Ex-Vivo MR Brain Imaging in the Mouse." *Frontiers in Neuroinformatics* 0 (MARCH): 6.

Li, Chenchen, Bo Liu, Jingyi Xu, Bin Jing, Lijie Guo, Liyong Wang, Milin Wang, et al. 2023. "Phloretin Decreases Microglia-Mediated Synaptic Engulfment to Prevent Chronic Mild Stress-Induced Depression-like Behaviors in the MPFC." *Theranostics* 13 (3): 955–72.

Lian, Hong, Li Yang, Allysa Cole, Lu Sun, Angie C-A Chiang, Stephanie W. Fowler, David J. Shim, et al. 2015. "NFκB-Activated Astroglial Release of Complement C3 Compromises Neuronal Morphology and Function Associated with Alzheimer's Disease." *Neuron* 85 (1): 101–15.

Ludewig, Katja, Mark A. Geyer, and Franz X. Vollenweider. 2003. "Deficits in Prepulse Inhibition and Habituation in Never-Medicated, First-Episode Schizophrenia." *Biological Psychiatry* 54 (2): 121–28.

Madeshiya, Amit Kumar, Carl Whitehead, Ashutosh Tripathi, and Anilkumar Pillai. 2022. "C1q Deletion Exacerbates Stress-Induced Learned Helplessness Behavior and Induces Neuroinflammation in Mice." *Translational Psychiatry* 12 (1): 50.

Mamane, Yaël, Chi Chung Chan, Genevieve Lavallee, Nicolas Morin, Li-Jing Xu, Jingqi Huang, Robert Gordon, et al. 2009. "The C3a Anaphylatoxin Receptor Is a Key Mediator of Insulin Resistance and Functions by Modulating Adipose Tissue Macrophage Infiltration and Activation." *Diabetes* 58 (9): 2006–17.

McCutcheon, Robert A., Tiago Reis Marques, and Oliver D. Howes. 2020. "Schizophrenia-an Overview." *JAMA Psychiatry (Chicago, Ill.)* 77 (2): 201–10.

Mena, Auxiliadora, Juan C. Ruiz-Salas, Andrea Puentes, Inmaculada Dorado, Miguel Ruiz-Veguilla, and Luis G. De la Casa. 2016. "Reduced Prepulse Inhibition as a Biomarker of Schizophrenia." *Frontiers in Behavioral Neuroscience* 10 (October): 202.

Mommert, Susanne, Derya Aslan, Lisanne Ratz, Holger Stark, Ralf Gutzmer, and Thomas Werfel. 2018. "The Anaphylatoxin C3a Receptor Expression on Human M2 Macrophages Is Down-Regulated by Stimulating the Histamine H4 Receptor and the IL-4 Receptor." *Journal of Innate Immunity* 10 (4): 349–62.

O'Connell, Kevin S., Ida E. Sønderby, Oleksandr Frei, Dennis van der Meer, Lavinia Athanasiu, Olav B. Smeland, Dag Alnæs, et al. 2021. "Association between Complement Component 4A Expression, Cognitive Performance and Brain Imaging Measures in UK Biobank." *Psychological Medicine* 52 (15): 1–11.

Olsen, Reid H. J., Jeffrey F. DiBerto, Justin G. English, Alexis M. Glaudin, Brian E. Krumm, Samuel T. Slocum, Tao Che, et al. 2020. "TRUPATH, an Open-Source Biosensor Platform for Interrogating the GPCR Transducerome." *Nature Chemical Biology* 16 (8): 841–49.

Osimo, Emanuele Felice, Katherine Beck, Tiago Reis Marques, and Oliver D. Howes. 2019. "Synaptic Loss in Schizophrenia: A Meta-Analysis and Systematic Review of Synaptic Protein and MRNA Measures." *Molecular Psychiatry* 24 (4): 549–61.

Paus, T., M. Keshavan, and J. N. Giedd. 2008. "Why Do Many Psychiatric Disorders Emerge during Adolescence?" *Nature Reviews. Neuroscience* 9 (12): 947–57.

Piekarski, David J., Natalie M. Zahr, Qingyu Zhao, Uran Ferizi, Kilian M. Pohl,
Edith V. Sullivan, and Adolf Pfefferbaum. 2023. "White Matter Microstructural
Integrity Continues to Develop from Adolescence to Young Adulthood in Mice and
Humans: Same Phenotype, Different Mechanism." *Neuroimage. Reports* 3 (3).

Pintér, O., Zoltan Beda, Z. Csaba, and I. Gerendai. 2007. "Differences in the Onset
of Puberty in Selected Inbred Mouse Strains." *Endocrine Abstracts* 14 (April).

Pozo-Rodrigálvarez, Andrea, Roosa Ollaranta, Jenny Skoog, Milos Pekny, and
Marcela Pekna. 2021. "Hyperactive Behavior and Altered Brain Morphology in
Adult Complement C3a Receptor Deficient Mice." *Frontiers in Immunology* 12
(February): 406.

Prut, Laetitia, and Catherine Belzung. 2003. "The Open Field as a Paradigm to
Measure the Effects of Drugs on Anxiety-like Behaviors: A Review." *European*
Journal of Pharmacology 463 (1–3): 3–33.

Qiu, Lily R., Darren J. Fernandes, Kamila U. Szulc-Lerch, Jun Dazai, Brian J.
Nieman, Daniel H. Turnbull, Jane A. Foster, Mark R. Palmert, and Jason P. Lerch.
2018. "Mouse MRI Shows Brain Areas Relatively Larger in Males Emerge before
Those Larger in Females." *Nature Communications* 9 (1): 1–15.

Quell, Katharina M., Christian M. Karsten, Anna Kordowski, Larissa Nogueira
Almeida, Daria Briukhovetska, Anna V. Wiese, Jing Sun, et al. 2017. "Monitoring
C3aR Expression Using a Floxed TdTomato-C3aR Reporter Knock-in Mouse." *The*
Journal of Immunology 199 (2): 688–706.

Reynolds, Jess E., Melody N. Grohs, Deborah Dewey, and Catherine Lebel. 2019.
"Global and Regional White Matter Development in Early Childhood."
NeuroImage 196 (August): 49–58.

Rodriguez, Pedro, Lauren J. Laskowski, Jean Pierre Pallais, Hailey A. Bock, Natalie
G. Cavalco, Emilie I. Anderson, Maggie M. Calkins, et al. 2023. "Functional
Profiling of the G Protein-Coupled Receptor C3aR1 Reveals Ligand-Mediated
Biased Agonism." *The Journal of Biological Chemistry* 300 (1): 105549.

Rojo, Rocío, Anna Raper, Derya D. Ozdemir, Lucas Lefevre, Kathleen Grabert, Evi
Wollscheid-Lengeling, Barry Bradford, et al. 2019. "Deletion of a Csf1r Enhancer
Selectively Impacts CSF1R Expression and Development of Tissue Macrophage
Populations." *Nature Communications* 10 (1): 3215.

Rubinov, Mikail, and Olaf Sporns. 2010. "Complex Network Measures of Brain Connectivity: Uses and Interpretations." *NeuroImage* 52 (3): 1059–69.

Schimmelpfennig, Jakub, Jan Topczewski, Wojciech Zajkowski, and Kamila Jankowiak-Siuda. 2023. "The Role of the Salience Network in Cognitive and Affective Deficits." *Frontiers in Human Neuroscience* 17 (March): 1133367.

Schizophrenia Working Group of the Psychiatric Genomics Consortium. 2014. "Biological Insights from 108 Schizophrenia-Associated Genetic Loci." *Nature* 511 (7510): 421–27.

Sekar, A., A. R. Bialas, H. de Rivera, A. Davis, T. R. Hammond, Kamitaki N, Tooley K, et al. 2016. "Schizophrenia Risk from Complex Variation of Complement Component 4." *Nature* 530 (7589): 177–83.

Semaan, Sheila J., and Alexander S. Kauffman. 2015. "Daily Successive Changes in Reproductive Gene Expression and Neuronal Activation in the Brains of Pubertal Female Mice." *Molecular and Cellular Endocrinology* 401 (February): 84–97.

Serrano, Maria Elisa, Eugene Kim, Bernard Siow, Da Ma, Loreto Rojo, Camilla Simmons, Darryl Hayward, et al. 2023. "Investigating Brain Alterations in the Dp1Tyb Mouse Model of Down Syndrome - ScienceDirect." *Neurobiology of Disease* 188 (November): 106336.

Sforazzini, Francesco, Adam J. Schwarz, Alberto Galbusera, Angelo Bifone, and Alessandro Gozzi. 2014. "Distributed BOLD and CBV-Weighted Resting-State Networks in the Mouse Brain." *NeuroImage* 87 (February): 403–15.

Smith, Stephen M., Mark Jenkinson, Mark W. Woolrich, Christian F. Beckmann, Timothy E. J. Behrens, Heidi Johansen-Berg, Peter R. Bannister, et al. 2004. "Advances in Functional and Structural MR Image Analysis and Implementation as FSL." *NeuroImage* 23 Suppl 1 (January): S208-19.

Stevens, Beth, Nicola J. Allen, Luis E. Vazquez, Gareth R. Howell, Karen S. Christopherson, Navid Nouri, Kristina D. Micheva, et al. 2007. "The Classical Complement Cascade Mediates CNS Synapse Elimination." *Cell* 131 (6): 1164–78.

Sun, Rui, Meng-Yu Tang, Dan Yang, Yan-Yi Zhang, Yi-Heng Xu, Yong Qiao, Bin Yu, et al. 2024. "C3aR in the Medial Prefrontal Cortex Modulates the Susceptibility to LPS-Induced Depressive-like Behaviors through Glutamatergic Neuronal Excitability." *Progress in Neurobiology*, April, 102614.

Tabbaa, Manal, Allison Knoll, and Pat Levitt. 2023. "Mouse Population Genetics
Phenocopies Heterogeneity of Human Chd8 Haploinsufficiency." *Neuron* 111 (4):
539-556.e5.

Tao, Juan, Jing Zhao, Xiang-Ming Qi, and Yong-Gui Wu. 2021. "Complement-
Mediated M2/M1 Macrophage Polarization May Be Involved in Crescent
Formation in Lupus Nephritis." *International Immunopharmacology* 101 (Pt A):
108278.

Tasic, Bosiljka, Vilas Menon, Thuc Nghi Nguyen, Tae Kyung Kim, Tim Jarsky,
Zizhen Yao, Boaz Levi, et al. 2016. "Adult Mouse Cortical Cell Taxonomy Revealed
by Single Cell Transcriptomics." *Nature Neuroscience* 19 (2): 335–46.

Tasic, Bosiljka, Zizhen Yao, Lucas T. Graybuck, Kimberly A. Smith, Thuc Nghi
Nguyen, Darren Bertagnolli, Jeff Goldy, et al. 2018. "Shared and Distinct
Transcriptomic Cell Types across Neocortical Areas." *Nature* 563 (7729): 72–78.

Thirion, Bertrand, Gaël Varoquaux, Elvis Dohmatob, and Jean-Baptiste Poline.
2014. "Which FMRI Clustering Gives Good Brain Parcellations?" *Frontiers in*
Neuroscience 8 (July): 167.

Tillmon, Haven, Breeanne M. Soteros, Liang Shen, Qifei Cong, Mackenna Wollet,
Julianne General, Hanna Chin, et al. 2024. "Complement and Microglia Activation
Mediate Stress-Induced Synapse Loss in Layer 2/3 of the Medial Prefrontal Cortex in
Male Mice." *Nature Communications* 15 (1): 9803.

Tournier, J-Donald, Robert Smith, David Raffelt, Rami Tabbara, Thijs Dhollander,
Maximilian Pietsch, Daan Christiaens, Ben Jeurissen, Chun-Hung Yeh, and Alan
Connelly. 2019. "MRtrix3: A Fast, Flexible and Open Software Framework for
Medical Image Processing and Visualisation." *NeuroImage* 202 (116137): 116137.

Tripathi, Ashutosh, Carl Whitehead, Katelyn Surrao, Ananya Pillai, Amit
Madeshiya, Yong Li, Hesam Khodadadi, et al. 2021. "Type 1 Interferon Mediates
Chronic Stress-Induced Neuroinflammation and Behavioral Deficits via
Complement Component 3-Dependent Pathway." *Molecular Psychiatry* 26 (7):
3043–59.

Valiquette, Vanessa, Elisa Guma, Lani Cupo, Daniel Gallino, Chloe Anastassiadis,
Emily Snook, Gabriel A. Devenyi, and M. Mallar Chakravarty. 2023. "Examining
Litter Specific Variability in Mice and Its Impact on Neurodevelopmental Studies."
NeuroImage 269 (April): 119888.

Vasek, Michael J., Charise Garber, Denise Dorsey, Douglas M. Durrant, Bryan Bollman, Allison Soung, Jinsheng Yu, et al. 2016. "A Complement-Microglial Axis Drives Synapse Loss during Virus-Induced Memory Impairment." *Nature* 534 (7608): 538–43.

Vita, A., L. De Peri, G. Deste, and E. Sacchetti. 2012. "Progressive Loss of Cortical Gray Matter in Schizophrenia: A Meta-Analysis and Meta-Regression of Longitudinal MRI Studies." *Translational Psychiatry* 2 (11): e190.

Wang, Quanxin, Song-Lin Ding, Yang Li, Josh Royall, David Feng, Phil Lesnar, Nile Graddis, et al. 2020. "The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas." *Cell* 181 (4): 936-953.e20.

West, Erin E., and Claudia Kemper. 2023. "Complosome - the Intracellular Complement System." *Nature Reviews. Nephrology* 19 (7): 426–39.

Westacott, Laura J., Niels Haan, Claudia Evison, Omar Marei, Jeremy Hall, Timothy R. Hughes, Malik Zaben, et al. 2021. "Dissociable Effects of Complement C3 and C3aR on Survival and Morphology of Adult Born Hippocampal Neurons, Pattern Separation, and Cognitive Flexibility in Male Mice." *Brain, Behavior, and Immunity* 98 (November): 136–50.

Westacott, Laura J., Trevor Humby, Niels Haan, Sophie A. Brain, Emma-Louise Bush, Margarita Toneva, Andreea-Ingrid Baloc, et al. 2022. "Complement C3 and C3aR Mediate Different Aspects of Emotional Behaviours; Relevance to Risk for Psychiatric Disorder." *Brain, Behavior, and Immunity* 99 (January): 70–82.

Westacott, Laura J., and Lawrence S. Wilkinson. 2022. "Complement Dependent Synaptic Reorganisation During Critical Periods of Brain Development and Risk for Psychiatric Disorder." *Frontiers in Neuroscience* 16 (May): 840266.

Wood, Tobias C., Camilla Simmons, Samuel A. Hurley, Anthony C. Vernon, Joel Torres, Flavio Dell'Acqua, Steve C. R. Williams, and Diana Cash. 2016. "Whole-Brain Ex-Vivo Quantitative MRI of the Cuprizone Mouse Model." *PeerJ* 4 (November): e2632.

Yilmaz, Melis, Esra Yalcin, Jessy Presumey, Ernest Aw, Minghe Ma, Christopher W. Whelan, Beth Stevens, Steven A. McCarroll, and Michael C. Carroll. 2021. "Overexpression of Schizophrenia Susceptibility Factor Human Complement C4A Promotes Excessive Synaptic Loss and Behavioral Changes in Mice." *Nature Neuroscience* 24 (2): 214–24.

Zalesky, Andrew, Alex Fornito, and Edward T. Bullmore. 2010. "Network-Based Statistic: Identifying Differences in Brain Networks." *NeuroImage* 53 (4): 1197–1207.

Zalesky, Andrew, Alex Fornito, Ian H. Harding, Luca Cocchi, Murat Yücel, Christos Pantelis, and Edward T. Bullmore. 2010. "Whole-Brain Anatomical Networks: Does the Choice of Nodes Matter?" *NeuroImage* 50 (3): 970–83.

Zhan, Y., R. C. Paolicelli, F. Sforazzini, L. Weinhard, G. Bolasco, F. Pagani, A. I. Vyssotski, et al. 2014. "Deficient Neuron-Microglia Signaling Results in Impaired Functional Brain Connectivity and Social Behavior." *Nature Neuroscience* 17 (3): 400–406.

Zhang, Man-Man, Min-Xia Guo, Qiu-Ping Zhang, Xue-Qin Chen, Na-Zhi Li, Qing Liu, Jie Cheng, et al. 2022. "IL-1R/C3aR Signaling Regulates Synaptic Pruning in the Prefrontal Cortex of Depression." *Cell & Bioscience* 12 (1): 90.

Zheng, Jingwei, Jia'nan Lu, Shuhao Mei, Haijian Wu, Zeyu Sun, Yuanjian Fang, Shenbin Xu, et al. 2021. "Ceria Nanoparticles Ameliorate White Matter Injury after Intracerebral Hemorrhage: Microglia-Astrocyte Involvement in Remyelination." *Journal of Neuroinflammation* 18 (1): 43.

Zhu, Hongru, Changjian Qiu, Yajing Meng, Minlan Yuan, Yan Zhang, Zhengjia Ren, Yuchen Li, et al. 2017. "Altered Topological Properties of Brain Networks in Social Anxiety Disorder: A Resting-State Functional MRI Study." *Scientific Reports* 7 (1): 43089.