Growth rates and Metabolic Traits Differentiate *Campylobacter jejuni* Strains by Diarrheal

Manifestation

Jennifer M. Bosquez[1], Craig T. Parker[2], Ben Pascoe[3,4], Kerry K. Cooper[1, 5]*

[1]School of Animal and Comparative Biomedical Sciences, University of Arizona, Tucson, AZ

85721, USA

[2]Produce Safety and Microbiology Research Unit, Agricultural Research Service, U.S. Department

of Agriculture, Albany, California, USA

[3]Ineos Oxford Institute for Antimicrobial Research, Department of Biology, University of Oxford,

Oxford, United Kingdom

[4]Faculty of veterinary Medicine, Chiang Mai University, Chiang Mai, Thailand

[5]BIO5 Institute, University of Arizona, Tucson, AZ 85719, USA

Corresponding author:

Kerry K. Cooper, School of Animal and Comparative Biomedical Sciences, 1117 E. Lowell St.,

University of Arizona, Tucson, AZ, kcooper@arizona.edu, Phone: (520) 621-3342

Key words: Campylobacter jejuni, L-fucose, L-glutamine, biofilm, motility, generation time,

growth

Short title: Metabolism and growth differences drive *Campylobacter jejuni* diarrheal

manifestations

Abstract

Campylobacter jejuni is the leading bacterial cause of gastroenteritis worldwide, and can result in two different diarrheal manifestations in humans: a bloody/inflammatory diarrhea or a watery diarrhea. Currently, little is known about the factors leading to these two distinct diarrheal manifestations; however, five strains of *C. jejuni* associated with bloody/inflammatory diarrhea and five strains associated with watery diarrhea have been characterized in the neonatal piglet diarrhea model system. Here, we demonstrate individual differences between *C. jejuni* strains and found that these two groups of *C. jejuni* strains exhibit differences in temperature dependent and independent growth rate and carbon source utilization. We show that that watery diarrhea associated *C. jejuni* strains grow significantly faster and have enhanced biofilm production compared to bloody/inflammatory diarrhea associated *C. jejuni* strains at human body temperature (37°C). Whereas bloody/inflammatory diarrhea associated strains grew faster at 42°C compared to 37°C. Using phenotypic microarrays to examine 192 carbon source utilization as a preliminary screening, we show that L-fucose is utilized differently by strains within the two diarrheal manifestation groups. L-fucose utilization is significantly higher among watery diarrhea associated strains at both 37°C and 42°C compared to bloody/inflammatory associated strains. While L-glutamine utilization is significantly higher among bloody/inflammatory diarrhea associated strains at 37°C compared to watery associated strains. Overall, these findings suggest that there are distinct metabolic adaptations of bloody/inflammatory and watery diarrhea-associated *C. jejuni* strains to human (37°C) and poultry (42°C) temperatures that may contribute to their differing diarrheal manifestations.

Introduction

Campylobacter jejuni is a major cause of bacterial gastroenteritis, and is responsible for an estimated 550 million cases per year worldwide[1]. Patients with campylobacteriosis caused by *C. jejuni* typically experience symptoms within 2-5 days after exposure, which can include abdominal cramping, nausea, and fever along with diarrhea that can be watery or bloody/inflammatory[2-4]. The diarrheal manifestations of *C. jejuni*, watery and/or bloody/inflammatory, have been noted for several decades[5; 6], yet little has been explored concerning the role of *C. jejuni* phenotypic characteristics in these outcomes. Watery diarrhea is generally characterized by loose non-bloody stools, while the more severe bloody/inflammatory diarrhea includes the presence of blood in the stool and/or inflammation of the ileum or colon[4; 7]. Campylobacteriosis is associated with several post-infectious sequelae including Guillain-Barré syndrome and post-infectious irritable bowel syndrome (PI-IBS)[8; 9]. Previous studies have suggested that severe, bloody diarrhea is associated with increased invasiveness[10-12] and cytotoxin production[13-15], while watery diarrhea has been linked to secretory mechanisms and enterotoxin activity[16-20]. However, fundamental bacterial traits such as growth rate, carbon source utilization, motility, and biofilm formation, key factors in colonization and persistence, have not been systematically examined in the context of diarrheal manifestation.

C. jejuni isolates are phenotypically diverse including in their LOS[21; 22], biofilm/aerotolerance[23], antimicrobial resistance, in their ability to produce PI-IBS[24], immune response[25], phase-variation[26] or within a single sequence type (ST)[27; 28], among agriculture strains[29] and even laboratory strains demonstrate considerable variability following multiple growth passages[30-34]. Unlike many enteric bacteria, *C. jejuni* generally lacks key pathways necessary for the metabolism of most carbohydrates in the intestinal tract such as glucose and

galactose[35]. *C. jejuni* has a relatively small genome and few identified metabolic pathways

compared to other enteric pathogens, such as *Salmonella,* but despite this, it has unique metabolic

flexibility and can utilize substrates such as amino acids or citric acid cycle intermediates[31; 36;

37]. Metabolic heterogeneity has been observed between *C. jejuni* strains with some strains able

to metabolize glutamine and asparagine[38]. In fact, the amino acid L-glutamine can be utilized as

a primary energy source in *C. jejuni* strains containing γ-glutamyl transpeptidase (GGT)[39].

Strains of *C. jejuni* that contain GGT, such as the sequence type 45 clonal complex (ST-45 CC)[40]

have been shown to have a colonization advantage in murine models. However, other *C. jejuni*

strains may be able to compensate for the lack of GGT by utilizing other metabolic pathways[38;

41]. Additionally, certain *C. jejuni* strains, particularly those of ST-22 CC[40], can contain a

genomic island for L-fucose metabolism, a sugar present in mucin glycoproteins in the intestinal

tract[42; 43]. L-fucose metabolism provides a competitive advantage during colonization of

certain hosts, however, its role in human pathogenesis remains unclear[42; 44]. Metabolic diversity

among *C. jejuni* strains may have a role in colonization and virulence differences within different

hosts, including strain variation of diarrheal manifestations in humans. Colonization factors like

growth, motility, biofilm production, and carbon utilization have rarely been studied for their

potential role in the diarrheal manifestation of *C. jejuni* yet understanding them is crucial to

furthering understanding the pathogenesis of *C. jejuni*.

Here, we investigated whether growth characteristics and metabolic capacity differ between strains

of *C. jejuni* associated with distinct diarrheal manifestations, and whether these differences are

consistent with host adaptation or zoonotic potential. Using a panel of ten well-characterized *C.*

jejuni strains shown to cause either watery or bloody/inflammatory diarrhea in the neonatal piglet

model - a system that reliably recapitulates human disease phenotypes[45; 46], we compared

generation times, carbon source utilization (L-fucose, L-glutamine, and mucin), motility, and

biofilm formation across temperatures representative of human (37°C) and avian (42°C) hosts.

Materials and Methods

Bacterial Culturing and Growth

Strains used in this study, their source and associated diarrheal manifestation in the neonatal piglet

model are listed in **Table 1**. The strains selected for this study were chosen based on their

documented diarrheal manifestations in neonatal piglets, offering a more controlled way to

investigate the mechanisms behind the observed clinical variability. All watery diarrhea associated

strains were sourced from poultry, while two strains associated with bloody/inflammatory diarrhea

originated from poultry and the other three were isolated from human clinical cases. In addition to

the *in vivo* work that has been performed with these strains, all have been characterized either

genotypically or phenotypically to varying degrees in previous studies, with strains NCTC11168,

81-176 and M129 the most well characterized. However, none of these strains have been

investigated as part of two diarrheal manifestation groups for their growth and metabolic

capabilities as has been done in this study. All *C. jejuni* strains were grown and maintained on

Mueller-Hinton agar plates supplemented with 5% defibrinated bovine blood (MHB; Becton,

Dickinson and Company Difco) for 48 hours at 37°C with 10% CO_2. Strains for assays that

required liquid culture were grown in Mueller-Hinton (MH; Becton, Dickinson and Company

Difco) broth with constant shaking (100 rpm) for 16-18 hours at 37°C with 10% CO_2.

Growth Curves

Growth curves in MH broth were conducted for each of the *C. jejuni* strains. Overnight cultures

were grown with shaking (100 rpm) at 37°C with 10% CO_2, OD_{600} measured, and then diluted to

a final OD_{600} of 0.001 (~1 x 10^6 CFU/ml) with MH broth. Next, 300 µl aliquots of the diluted

inoculum were added in triplicate to wells in a sterile 96-well polystyrene plate (Fisher Scientific),

and the plate incubated at 37°C with 10% CO_2 for 1 hour before being sealed tightly with an optical

plate film cover (Applied Biosystems) to prevent evaporation during incubation. The sealed plate

was then placed in a SpectraMax M2 plate reader at either 37°C or 42°C, to investigate body

temperature differences between humans (37°C) and poultry (42°C; the primary *C. jejuni*

reservoir), with continuous shaking (100 rpm) for 36 hours with OD_{600} readings recorded every 15

minutes. Generation times for each strain were calculated using readings from the logarithmic

phase of growth. Three biological replicates were performed for all 10 *C. jejuni* strains with two

technical replicates (n = 3).

Carbon Utilization Microarray

Phenotype carbon utilization microarrays were conducted as described previously by Line et al

[47]. Briefly, phenotype microarrays (PMs) were conducted using the PM1 and PM2A plates

(Biolog Inc., CA, USA) for variation in carbon utilization sources between strains. Each strain of

C. jejuni was grown on MHB agar for 12 – 16 hours at 37°C with 10% CO_2. The bacteria were

collected from the MHB agar plate using an inoculating loop, resuspended in 17 ml of IF-0a

medium (Biolog Inc., CA, USA), and then diluted to a final OD_{600} of 0.8 in IF-0a medium. Next,

the inoculum suspension was mixed with IF-0a, a phenotype microarray additive solution

containing tetrazolium violet (Redox dye D), according to manufacturer's instructions. Then 100

µl of the inoculum suspension was added to each well of either the PM1 or PM2A plates. The

plates were incubated at 37°C with 10% CO_2 for 48 hours before taking OD_{550} measurements in a

microplate reader (SpectraMax M2) and visually recording a color change from clear to violet in

each well on a scale of 0-3. A negative control plate for both types of plates was also incubated in

the absence of bacteria to determine false positives. Each strain was screened a single time with

both the PM1 and PM2A carbon utilization plates to serve as an initial screening method to indicate

potential carbon sources of interest to study further for the strains as described below (n = 1).

L- Fucose, L- Glutamine, and Mucin Utilization Assays

The phenotype microarray initial screening results indicated carbon utilization differences between

the diarrheal manifestation groups for both L-fucose and L-glutamine, thus we focused on these

carbon sources for additional studies. Studies on L-fucose and L-glutamine were done at both 37°C

and 42°C. Mucin utilization at 37°C was added as an additional assay as it has been shown to be

important to *Campylobacter* pathogenesis and L-fucose is a component of mucus. L-fucose, L-

glutamine, and mucin utilization assays were adapted from Stahl et al [42]. Briefly, overnight MH

broth cultures of the different *C. jejuni* strains were grown with shaking (100 rpm) then diluted to

a final OD_{600} of 0.01 with minimum essential medium (MEM; Gibco) without glutamine or phenol

red and supplemented with 20 μM $FeSO_4$. The media was also supplemented with either no

supplement, 25 mM L-fucose (Thermo Scientific), 0.01 mg/ml bovine submaxillary gland mucin

(Millipore) or 20 mM L-glutamine (Alfa Aesar) depending on the assay. Next, 300 μl aliquots of

the diluted inoculum with or without the appropriate supplement were inoculated in triplicate into

a sterile 96-well plate, incubated at 37°C with 10% CO_2 for 1 hour, then sealed tightly with an

optical plate film cover (Applied Biosystems) to prevent evaporation during incubation. The sealed

plate was then placed in a SpectraMax M2 plate reader at either 37°C or 42°C (for L-fucose and

L-glutamine) with continuous shaking (100 rpm) for 36 hours with OD_{600} reading recorded every

15 minutes. The plate included a blank control well containing 300 µl of uninoculated medium.

Results were recorded as the maximum OD_{600} obtained for the culture during 36 hours of growth

minus the blank media OD_{600}[39]. Each strain was tested in three biological replicates with three

technical replicates for each of the different carbon utilization assays with either L-fucose, mucin

or L-glutamine or without supplements (n = 3). There was no growth without any L-fucose, mucin

or L-glutamine present during any of the assays.

Motility Assay (Soft Agar Swarming)

Motility assays were conducted based on assays described previously by Reuter et al [48] with the

following modifications: (1) using an OD_{600} of 1.0; (2) checking at time points at 24 hours and 48

hours; (3) using a 10 µl inoculation into the agar; and (4) incubating the agar plates at either 37°C

or 42°C. Briefly, each *C. jejuni* strain was grown overnight in MH broth, the OD_{600} measured, and

the culture diluted in MH broth to a final OD_{600} of 1.0. Mueller Hinton (MH) soft agar plates

(Mueller Hinton broth supplemented with 0.4% agar) were inoculated with 10 µl of the diluted

inoculum directly into the center of the plate by stabbing into the agar. Inoculated plates were

incubated at either 37°C or 42°C with 10% CO_2 for 48 hours. The radius of growth was measured

after 24 hours and 48 hours of incubation using a digital calibrator starting from the edge of the

inoculum spot to the edge of growth. Each strain was inoculated onto two separate plates for two

technical replicates and each assay was conducted in three biologically independent assays (n =

3).

Biofilm Assay

Biofilm assays were performed as previously described by Reeser et al[49] except for the

following modification: (1) strains were inoculated at an OD_{600} of 0.25. Briefly, *C. jejuni* strains

were grown overnight in MH broth with shaking (100 rpm) at 37°C with 10% CO_2, the OD_{600} was

measured, and the culture diluted with MH broth to a final OD_{600} of 0.25. Then 100 µl of diluted

inoculum were then added to the corresponding well of a sterile 24-well polystyrene plate (Fisher

Scientific, Los Angeles, CA, USA) containing 1 ml of MH broth. The plate was then statically

incubated at 37°C with 10% CO_2 for 72 hours. After 72 hours incubation, all liquid media was

carefully removed from each well by pipetting to avoid disrupting the pellicle, and then the plate

was dried at 55°C for 30 minutes. The pellicle in each well was then stained by adding 1 ml of

0.1% crystal violet (100 mg of crystal violet powder (Acros Organics), 8 ml of sterile double

distilled water (ddH_2O), and 2 ml 100% methanol (Fisher Bioreagents)) and incubating at room

temperature for 5 minutes. Next, each well was washed twice with 1 ml of sterile ddH_2O to remove

any excess crystal violet, the plate was dried at 55°C for 15 minutes, and then decolorized by

adding 1 ml of decolorizer solution (80% ethanol (Fisher Bioreagents) and 20% acetone solution

(Fisher Bioreagents)). Finally, 100 µl of the decolorized solution for each strain was transferred to

a corresponding well of a 96-well polystyrene plate, and the absorbance at 570 nm was measured

using a SpectraMax M2 microplate reader. Each strain was tested in two technical replicates during

three biologically independent assays (n = 3).

Statistical Analysis

Statistical analyses were performed using GraphPad Prism (v10.4.1) for comparisons between

diarrheal manifestations or RStudio (v2022.07.1) for comparisons between strains. All data were

tested for normality (Shapiro–Wilk test) and equal variance (Levene's test). For comparisons

between strains associated with watery and bloody/inflammatory diarrhea (n = 15 per group), the

non-parametric Mann–Whitney U test was applied. Statistical significance was defined as $p \leq 0.05$.

For comparisons across multiple strains or conditions, non-parametric Kruskal–Wallis tests with

Dunn's post hoc test were used. Compact Letter Display (CLD) notation was used to visualize the

results of the pairwise comparisons. Strains that do not share a letter are significantly different

from each other ($p < 0.05$), while strains sharing the same letter are not significantly different.

Letters were assigned using the multcompView package in R, based on adjusted p-values.

Statistical significance was defined as $p \leq 0.05$. Each assay included a minimum of three

independent biological replicates per strain (n = 3), with technical replicates as noted. Data are

presented as mean ± SEM unless otherwise stated.

Results

Watery diarrhea-associated C. jejuni strains grow faster than bloody/inflammatory strains at

37°C

To evaluate whether differences in growth dynamics could explain divergent clinical

manifestations, we compared the growth kinetics of five watery diarrhea-associated and five

bloody/inflammatory-associated *C. jejuni* strains at two host-relevant temperatures: 37°C (human)

and 42°C (poultry). Growth was monitored over 36 hours using OD_{600} measurements taken every

15 minutes. Strain to strain differences were observed widely among both 37°C and 42°C, and

strains D42a, M129, D33a, A9a and NCTC11168 all grew significantly slower at 37°C than at

42°C ($p < 0.05$; **Figure 1A**).

At both temperatures, watery diarrhea associated strains entered logarithmic growth phase earlier

and reached higher OD_{600} values compared to bloody/inflammatory strains (**Supplementary**

Figure 1A–D). One watery strain (D42a) displayed a slower growth profile, similar to that of

bloody/inflammatory strains. Generation times calculated from logarithmic growth phase found

significantly faster generation times for watery diarrhea associated strains compared to

bloody/inflammatory associated strains at 37°C but not at 42°C (37°C watery diarrhea associated

115.8 minutes, 37°C bloody/inflammatory diarrhea associated 139.99 minutes p = 0.018; 42°C

watery diarrhea associated 95.23 minutes, bloody/inflammatory diarrhea associated 42°C 111.47

minutes p = 0.38; **Figure 1B**). There was no significant difference in generation times at 37°C

compared to 42°C for watery diarrhea associated strains (p = 0.12) but bloody/inflammatory

diarrhea associated strains grew significantly faster at 42°C than 37°C (p = 0.0057).

Strains differ in utilization of L-fucose and L-glutamine but not mucin

To determine whether nutrient utilization profiles differ between strains associated with distinct

diarrheal outcomes, we screened all ten *C. jejuni* strains for their ability to metabolize 192 carbon

sources using BIOLOG PM1 and PM2A phenotype microarray plates. There were a few carbon

sources used solely by one or two bloody/inflammatory diarrhea associated strains and not watery

associated strains and vice versa, including D-tagatose (D33a), succinnamic acid (D33a, 81-176),

citric acid (NCTC11168) and tricatballyic Acid (S3). Several carbon sources exhibited distinct

utilization patterns between the two diarrheal manifestation groups (**Supplementary Figure 2**).

L-serine was utilized by only three bloody/inflammatory diarrhea associated strains (M129, 81-

176 and NCTC11168) and no watery diarrhea associated strains, while L-glycyl-proline was

utilized by watery diarrhea associated strains A13a, A14a, S3 but only bloody/inflammatory

diarrhea associated strain A9a. Notably, L-glutamine and L-fucose were utilized by multiple

strains in only one group: L-glutamine was used by three bloody/inflammatory strains (81-176,

M129, D33a) and only one watery strain (S3), while L-fucose was used by four watery strains

(A13a, A14a, D42a, S3) and just one bloody/inflammatory strain (A9a). L-fucose and L-glutamine

were selected for further phenotypic analysis, along with mucin due to its relevance in host

colonization[50-52].

All five watery strains showed robust growth in the presence of L-fucose, with strain S3 reaching

the highest OD_{600} at 42°C (**Figure 2A**). At 37°C, the mean OD_{600} for watery strains was 0.077

compared to 0.035 for bloody/inflammatory strains ($p < 0.0001$); at 42°C, the difference was even

more pronounced (mean OD_{600}: 0.149 vs. 0.066, $p < 0.0001$). L-fucose supported significantly

greater growth in watery diarrhea-associated strains at both temperatures (**Figure 2B**). These

results indicate that L-fucose metabolism is a consistent and temperature-independent trait among

watery diarrhea-associated strains. To assess whether differences in mucosal substrate metabolism

extended beyond individual monosaccharides, we next tested the ability of each strain to grow in

medium supplemented with bovine submaxillary gland mucin at 37°C. Although there was strain-

level variability, no significant difference in mucin utilization was observed between watery and

bloody/inflammatory strains ($p = 0.67$), (**Figure 3, Supplementary Figure 3**).

L-glutamine utilization supported higher levels of growth among all strains compared to L-fucose,

with several significant differences between different strains at 37°C or 42°C (**Figure 4A**). Among

all strains tested, only A13a showed significantly higher growth at 42°C compared to 37°C, while

81-176 and A9a both showed increased growth at 37°C compared to 42°C. However, when

grouped by diarrheal manifestation, bloody/inflammatory associated strains exhibited enhanced

growth at 37°C in L-glutamine-supplemented medium compared to watery diarrhea associated

strains ($p = 0.0005$) (**Figure 4B**). There was no difference in growth potential at 42°C between

watery and bloody/inflammatory strains, ($p = 0.68$), suggesting a temperature or host dependent

role for glutamine utilization among *C. jejuni* strains. Together, these results demonstrate that

carbon source utilization differs consistently between strain groups and that these traits may be

modulated by host temperature. The enhanced L-fucose metabolism of watery diarrhea associated

strains may facilitate rapid growth and colonization in both poultry and human hosts, while the

slower growth and utilization of L-glutamine by bloody/inflammatory diarrhea associated strains

contributes to persistence within the human host.

Motility varies widely among C. jejuni strains but does not distinguish diarrheal phenotype

Motility is critically important for *C. jejuni* colonization and pathogenesis, aiding in both adhesion

as well as invasion of host cells[53-55]. To assess whether motility differs between strains

associated with watery versus bloody/inflammatory diarrhea, we performed swarming motility

assays in soft agar at both 37°C and 42°C. A wide range of motility phenotypes was observed

among *C. jejuni* strains: bloody/inflammatory strains NCTC11168, D33a, and 81-176, and watery

strains D42a and RM1221, were highly motile (>10 mm radius), while bloody/inflammatory strain

A9a and watery strain S3 were weakly motile or non-motile (<10 mm) (**Figure 5**). While no

significant difference in motility was observed between the two diarrheal manifestations at either

temperature, a definite trend toward reduced motility among bloody/inflammatory diarrhea

associated strains was noted at 37°C ($p = 0.052$). No such trend was observed at 42°C, where the

difference remained non-significant ($p = 0.66$; **Supplementary Figure 4**). There was no

significant difference in motility at different temperatures for the bloody/inflammatory ($p = 0.18$)

or watery ($p = 0.073$) diarrhea associated strains, but watery diarrhea associated strains trended

towards significance.

Biofilm formation differs by diarrheal phenotype

Biofilm formation contributes to the persistence, and transmission of *C. jejuni*[56]. Therefore, to

assess the potential differences in these traits among our *C. jejuni* strains, we quantified biofilm

formation at 37°C using crystal violet staining. Strong biofilm producers ($A_{570} > 0.30$) included

bloody/inflammatory diarrhea associated strain M129 and watery diarrhea associated strains A13a

and D42a. Medium biofilm formation (A_{570} between 0.20–0.30) was observed in

bloody/inflammatory diarrhea associated strains 81-176 and NCTC11168 and watery diarrhea

associated strains A14a and S3, while poor biofilm formation ($A_{570} < 0.20$) was noted for A9a,

D33a, and RM1221 (**Figure 6A**).Although strains varied in their individual capacities, biofilm

formation at 37°C was found to significantly differ between diarrheal manifestation groups (p =

0.019; **Figure 6B**).

Discussion

This study investigated the variation between strains of *C. jejuni* with defined diarrheal phenotypes

in the neonatal piglet model, we assessed growth rates, carbon source utilization, motility, and

biofilm formation at human and avian body temperatures. In human and poultry hosts, *C. jejuni*

must pass through the mucosal layer of the intestines to reach the intestinal epithelial surface and

establish an infection. Here, *C. jejuni* strains showed significant strain to strain variation, including

differences in the utilization of L-fucose, L-glutamine, generation times, and biofilm formation,

but did not display significantly different abilities to utilize bovine submaxillary gland mucin when

grouped by diarrheal manifestation, findings that were not surprising based on *C. jejuni's* limited

ability to metabolize carbohydrates[35]. It is known that *C. jejuni* growth is enhanced when co-

cultured with commensal *Bacteroides vulgatus*, a response suggested to be due to the breakdown

of mucin by *B. vulgatus*[57]. There are several types of bacteria that are known mucin degraders

that have been identified in humans, calves and goats, and include *Bifidobacterium*[58; 59],

Bacteroides[59] and *Akkermansia muciniphila*[60]. Mucin degradation by *A. muciniphila* is

known to support intestinal barrier function[61] and provide physiological benefits that prevent

disease[61-63]. The composition of mucin degraders in chickens is less well defined, but

Bacteroides has been found to be one of the most predominating genera of bacteria in the chicken

intestinal microbiome[62; 64]. The growth of these commensal bacteria is supported by the highly

glycosylated mucins within the intestinal mucus[65], during which active fructosidases are

secreted[66; 67]. *C. jejuni* lacks the ability to release L-fucose from glycans and instead is known

to utilize free fucose from mucin degraders such as *B. vulgatus*[57].

The mucus layer of the intestinal tract is where *C. jejuni* encounters L-fucose, the only known

carbohydrate chemoattractant for *C. jejuni*[68; 69]. In the human gut, 80% of the total dry weight

of intestinal mucins is made up of *O*-linked glycans[70; 71], however in poultry L-fucose is

suggested to comprise an estimated 7% of the *O*-glycans in the intestinal tract[72]. The *fuc* locus

for L-fucose catabolism has been identified in >50% of sequenced *C. jejuni* isolates[73] and has

been observed among a high percentage of *C. jejuni* isolates from humans[74]. Here we found

that strains associated with watery diarrhea exhibited enhanced utilization of L-fucose at 37°C and

also grew significantly faster than those linked to bloody/inflammatory disease at 37°C and 42°C.

We hypothesis that this combination of rapid growth and efficient metabolism may support early

colonization and transmission, contributing to the acute nature of watery diarrhea.

The concentration of L-fucose is particularly high among breastfed infants due to the secretion of

L-fucose from fucosylated human milk oligosaccharides by intestinal commensals such as

Bifidobacterium spp and *Bacteriodes* spp[57; 67; 75; 76]. The amount of available L- fucose is

known to be as high as 4-5 mg/g in the feces of breastfed infants and in low to middle income

countries, higher proportions of *C. jejuni* have been found among these infants compared to those

that were not breastfed[77; 78]. Despite the high levels of available L-fucose, breastfed infants

with campylobacteriosis had significantly higher proportions of non-L-fucose metabolizing

Campylobacter than non-breastfed infants[77], a finding suggested to be driven in part by L-fucose

chemotaxis[79].

Chemotaxis towards L-fucose in the intestinal lumen through the upregulation of L-fucose

dehydrogenase, FucX, is suggested to contribute to its clearance while non-L-fucose metabolizing

C. jejuni strains move towards the intestinal mucosa to establish disease[79]. These findings

support the hypothesis that watery diarrhea associated strains utilize L-fucose in the mucus layer

where they are maintained further away from the intestinal epithelial cells while

bloody/inflammatory strains move to establish themselves at the enterocyte surface. Indeed, we

have recently found that bloody/inflammatory diarrhea associated *C. jejuni* strains are significantly

more invasive than watery diarrhea associated strains (Bosquez et al 2025). However, in contrast,

L-fucose utilization has also been observed *in vitro* to enhance invasion of Caco-2 cells and

significantly increase fibronectin binding efficacy in strain NCTC11168[44]. Another study found

similar results with invasion of Caco-2 increased with the addition of 10 mM L-fucose under

microaerophilic conditions using *C. jejuni* strain 108[67]. The same study observed growth

enhancement of *C. jejuni* strain 108 in the presence of L-fucose, a finding similar to ours,

suggesting that watery diarrhea associated strains may alter their invasiveness if tested under the

same conditions. In addition, we found that the increased growth potential of watery diarrhea

associated strains was maintained at both host-relevant temperatures, suggesting adaptation to

colonization across multiple hosts with enhanced growth in humans. The ability to utilize L-fucose

provided a colonization advantage *in vivo* in piglets but had no effect on the colonization of

chickens when inoculated at a low dose[42]. Since piglets are a model for human disease, whereas

chickens are the natural reservoir for *C. jejuni*, this suggests L-fucose utilization at 37°C (human

body temperature) versus 42°C (chicken body temperature) may only be relevant to human

disease.

In contrast, bloody/inflammatory diarrhea-associated strains grew more slowly particularly at

37°C, a factor that potentially contributes to prolonged colonization and disease development.

Slower growth may allow for slower establishment of an infection, which allows for delayed

immune recognition of the pathogen until the bacteria more established, potentially resulting in a

more damaging infection. Initial BIOLOG screening found that some strains preferentially utilized

L-glutamine, we observed a significant increase in growth potential in the presence of L-glutamine

by bloody/inflammatory associated strains compared to watery diarrhea associated strains at 37°C

but not at 42°C. In humans, glutamine is known to support epithelial barrier function and modulate

host immune responses, and its depletion may promote tissue damage and inflammation[80-82].

Because L-glutamine is a key fuel source for enterocytes and supports tight junction integrity[80;

83], *C. jejuni* utilization of L-glutamine may not only provide a metabolic advantage but also

actively disrupt enterocyte homeostasis. In the context of inflammation, where glutamine demand

is heightened due to increased epithelial turnover and immune cell activation, its depletion could

contribute to impaired repair to tissue damage as well as reduced immune function[84]. Other

studies have found that *C. jejuni* strains have considerable variation in their ability to utilize L-

glutamine, with 31% of strains[40] containing the enzyme gamma-glutamyl transpepidase (GGT)

necessary for glutamine and glutathione metabolism. One study found a higher percentage of *C.*

jejuni strains from human isolates (37.93%) to contain the *ggt* gene compared to 14.29% among

strains from non-human sources[51], suggesting an advantage for these strains in the human host.

Furthermore, a study of 166 *C. jejuni* strains from Finnish patients found an association between

bloody diarrhea and GGT production[85], while another study found purified *C. jejuni* GGT

impaired intestinal cell and lymphocyte proliferation[86]. The complicated and variable response

by *C. jejuni* strains containing *ggt* suggests that individual strains may have complex metabolic

requirements or can compensate with alternative pathways not yet characterized. Overall, the

metabolic diversity observed among strains highlights the importance of considering host-specific

and strain-specific adaptations when investigating L-glutamine utilization among *C. jejuni* strains,

especially among those associated with bloody/inflammatory diarrhea.

This study found that motility, a critical factor associated with colonization/pathogenicity[87-90],

had no significant differences between the different diarrheal manifestation groups. In terms of

motility*, Campylobacter* species utilize a single or bipolar flagella to move[91]. Upon entering a

host intestinal tract, the ability to move is important for navigating the thick layer of mucus to

reach the epithelial surface and colonize[91]. In this study, motility was variable at the strain level

and did not correlate strongly with diarrheal manifestations at either 37°C or 42°C in this study.

All strains were motile except strain S3, which was originally able to colonize poultry[92; 93], but

minimal laboratory passages rendered it defective for colonization[94]. Previously, similar results

were demonstrated for nonmotile *C. jejuni* strain CS that caused few pathological changes in the

piglet model, but grew to a greater maximum cell density in MH medium compared to a

genotypically similar but motile strain[88]. Watery diarrhea associated strains with limited to no

motility, such as S3, may be able to overcome the mucus layer via the more rapid growth that we

observed in this study, but additional studies are needed to confirm this hypothesis. Our results

suggest that motility is essential for the pathogenesis of *C. jejuni* regardless of the diarrheal

manifestation associated with the strain.

Biofilm formation is known to differ among *C. jejuni* strains[95] and has been suggested to

promote survival, transmission and colonization of hosts[96; 97]. The present study revealed

significantly higher biofilm formation among watery diarrhea associated strains at 37°C,

suggesting that these strains may have a colonization advantage in humans. In addition to

temperature, biofilm formation by bacteria can be triggered by various environmental stressors

such as nutrient availability, varying oxygen tensions and osmotic changes[98-100], for example,

one study showed that *C. jejuni* biofilm formation was increased under low-oxygen tensions as

well as in minimal nutrient medium[101]. An enhanced capacity to form biofilms would be

particularly useful in surviving the harsh conditions of the intestinal tract such as stomach acid,

bile salts, low-oxygen tension, oxidative stress or even immune defenses. When nutrients are

available, such as L-fucose, it is known that biofilm formation is reduced, and growth is enhanced

in vitro[73]. Future studies may investigate the ability of *C. jejuni* strains associated with watery

diarrhea to form biofilms when L-fucose is provided in the medium, particularly given their

enhanced growth with L-fucose. Stronger biofilm formation by strains associated with watery

diarrhea may be connected to their enhanced growth rate, which in combination would allow these

strains to quickly colonize the intestinal tract and contribute to the rapid production of watery

diarrhea.

It is important to note that our study revealed substantial phenotypic variability within each

diarrheal manifestation group. This is not unexpected given the known genomic and functional

diversity of *C. jejuni*, but it reinforces the need to study multiple strains to identify group-level

trends. Within the watery diarrhea group, strain D42a had a longer generation time at 37°C than

the other watery associated strains. S3, RM1221 and A14a had higher growth potential at 42°C in

the presence of L-fucose, while A13a showed significantly higher growth at 42°C in the presence

of L-glutamine. Among bloody/inflammatory strains, NCTC11168, A9a, D33a and M129 all had

slower generation times at 37°C than at 42°C. 81-176, D33a and NCTC-11168 showed similar

levels of motility, but A9a and M129 were only weakly. The presence of these outliers highlights

the spectrum of metabolic and growth capabilities within each pathotype and suggests that these

diarrheal manifestations may require several phenotypes to be present and may also rely on host

factors.

Another consideration is that strain classification in this study was based on diarrheal outcomes

observed in the neonatal piglet model, rather than direct human clinical data. While this may limit

immediate generalizability, the piglet model is the only experimental systems that reliably

reproduces both watery and bloody/inflammatory diarrheal manifestations caused by *C. jejuni.* It

provides a controlled environment to directly associate specific bacterial traits with clinical

outcomes, eliminating many of the confounding variables inherent in human observational studies.

Additionally, clinical strains are likely biased towards bloody/inflammatory diarrhea associated

strains, as people suffering from the less severe watery diarrhea are not likely to seek medical

attention. Nonetheless, validation of these findings using well-characterized human isolates with

known clinical presentations will be important for confirming their broader relevance.

Despite these limitations, our findings provide evidence that clinical outcome in *C. jejuni* infection

may be shaped by fundamental differences in bacterial growth and metabolism. Strains associated

with bloody/inflammatory diarrhea are slower growers, utilizing L-glutamine for enhanced growth

at 37°C, while those linked to watery diarrhea appear optimized for rapid expansion, biofilm

formation, and mucus-derived sugar utilization. These strategies may reflect differing host

adaptation trajectories and suggest that metabolic profiling could provide new insight into the

pathogenic potential of *C. jejuni* lineages.

Contributors

JB: Conceptualization, methodology, experimentation, formal analysis, writing – original draft, and review & editing. **KKC:** Conceptualization, methodology, resources, funding acquisition, project administration, supervision, writing, and review & editing. **CP:** Writing, and review & editing. **BP:** Writing, and review & editing.

Conflicts of Interest

The authors have no conflicts of interest.

Funding information

This work was supported in part by Start-up funds provided to Kerry Cooper by the University of Arizona, and Technology and Research Initiative Fund (TRIF) award CALS_ACBC_Cooper_2101696 provided to Kerry Cooper by the University of Arizona. No funding agency had any role in the study design, data collection, data analysis, or preparation of the manuscript.

Acknowledgments

The authors thank all members of the Cooper laboratory for providing critical technical assistance in sample processing and other aspects of the study. The authors thank Bryan Roxas for assistance with Supplementary Figure 2 and thank Dr. Jared Smith for his contributions to editing the manuscript.

492 **Table 1.** *Campylobacter jejuni* strains utilized in the study and associated diarrheal manifestation

Strain	Source	ST	Clonal Complex	Diarrheal Manifestation
NCTC11168	Clinical	ST43	ST-21	Bloody and/or Inflammatory
81-176	Clinical	ST604	ST-42	Bloody and/or Inflammatory
M129	Clinical	ST353	ST-353	Bloody and/or Inflammatory
A9a	Poultry	ST2827	ST-460	Bloody and/or Inflammatory
D33a	Poultry	ST459	ST-42	Bloody and/or Inflammatory
RM1221	Poultry	ST354	ST-354	Watery
S3	Poultry	ST354	ST-354	Watery
D42a	Poultry	ST21	ST-21	Watery
A14a	Poultry	ST1212	ST-607	Watery
A13a	Poultry	ST1212	ST-607	Watery

Figure Legends

Figure 1. Generation time of *C. jejuni* strains at 37°C and 42°C. (A) Generation Time in minutes at 37°C and 42°C for all *C. jejuni* strains. Average generation time calculated from logarithmic phase from three biologically independent assays with three technical replicates at both 37°C and 42°C (n = 3). Compact letter displays (CLDs) indicate statistically significant differences between groups (p < 0.05) as determined by Kruskal-Wallis test followed by Dunn's post hoc test. Bars labeled with different letters indicate statistically significant differences (p < 0.05) between those strains and/or temperatures, whereas shared letters denote groups that are not significantly different from each other. (B) Generation time in minutes of *C. jejuni* strains grouped by diarrheal manifestation (n = 15). There was a significant difference between bloody/inflammatory and watery diarrhea associated strains at 37°C (*, p = 0.018) but not at 42°C (p = 0.38). There was a significant difference between 37°C and 42°C among Bloody/inflammatory diarrhea associated strains (**, p = 0.0057), but among watery diarrhea associated strains (p = 0.12).

Figure 2. Utilization of L-fucose at 37°C and 42°C. (A) L-fucose utilization at 37°C and 42°C between *C. jejuni* strains. Compact letter displays (CLDs) indicate statistically significant differences between groups (p < 0.05) as determined by Kruskal-Wallis test followed by Dunn's post hoc test (n = 3). Bars labeled with different letters indicate statistically significant difference (p < 0.05) between those strains and/or temperatures, whereas shared letters denote groups that are not significantly different from each other. (B) L-fucose utilization at 37°C and 42°C between *C. jejuni* strains grouped by diarrheal manifestation. There was a significant difference between the growth of bloody/inflammatory and watery diarrhea associated strains when grown in minimal

media containing L-fucose at both 37°C (****, $p < 0.0001$) and at 42°C (****, $p < 0.0001$). The

average highest OD_{600} was calculated from three biologically independent assays with three

technical replicates (n = 15).

Figure 3. Utilization of mucin at 37°C. Bovine submucosally gland mucin utilization at 37°C by

C. jejuni strains. Compact letter displays (CLDs) indicate statistically significant differences

between groups ($p < 0.05$) as determined by Kruskal-Wallis test followed by Dunn's post hoc test.

Bars labeled with different letters indicate statistically significant difference ($p < 0.05$) between

those strains, whereas shared letters denote groups that are not significantly different from each

other. The average highest OD_{600} was calculated from three biologically independent assays with

three technical replicates (n = 3).

Figure 4. Utilization of L-glutamine at 37°C and 42°C. (A) Average highest OD_{600} L-glutamine

utilization at 37°C and 42°C across all *C. jejuni* strains. Compact letter displays (CLDs) indicate

statistically significant differences between groups ($p < 0.05$) as determined by Kruskal-Wallis

test followed by Dunn's post hoc test (n = 3). Bars labeled with different letters indicate statistically

significant difference ($p < 0.05$) between those strains and/or temperatures, whereas shared letters

denote groups that are not significantly different from each other. (B) Average highest OD_{600} L-

glutamine utilization at 37°C and 42°C across all *C. jejuni* strains grouped by diarrheal

manifestation. There was a significant difference between the growth of bloody/inflammatory and

watery diarrhea associated strains when grown at 37°C in minimal media containing L- glutamine

(***, $p = 0.0005$) but not at 42°C ($p = 0.68$). The average highest OD_{600} was calculated from three

biologically independent assays with three technical replicates (n = 15).

Figure 5. Motility at 37°C and 42°C. Average motility of *C. jejuni* strains after 48-hour incubation at both 37°C and 42°C. Motility was measured as the radius of the halo of growth in millimeters after 48-hour incubation. Average motility was calculated from three biologically independent assays with two technical replicates (n = 3). Compact letter displays (CLDs) indicate statistically significant differences between groups (p < 0.05) as determined by Kruskal-Wallis test followed by Dunn's post hoc test (n = 3). Bars labeled with different letters indicate statistically significant difference (p < 0.05) between those strains and/or temperatures, whereas shared letters denote groups that are not significantly different from each other.

Figure 6. Biofilm production at 37°C. (A) Formation of biofilm after 72 hours at 37°C among *C. jejuni* strains. Biofilm formation was quantified using spectrometer readings at 570 nm following a 72-hour incubation at 37°C. Compact letter displays (CLDs) indicate statistically significant differences between groups (p < 0.05) as determined by Kruskal-Wallis test followed by Dunn's post hoc test. Bars labeled with different letters indicate statistically significant difference (p < 0.05) between those strains, whereas shared letters denote groups that are not significantly different from each other. Average biofilm formation was determined from three biologically independent assays at 37°C with two technical replicates (n = 3) (B) Average biofilm production of bloody/inflammatory and watery diarrhea associated strains after 72 hours at 37°C. There was a significant difference between the biofilm production of watery and bloody/inflammatory strains at 37°C (**, p = 0.019). Average biofilm production at 37°C was calculated from three biologically independent assays with two technical replicates (n = 15).

References

[1] N. O. Kaakoush, N. Castaño-Rodríguez, H. M. Mitchell and S. M. Man, "Global Epidemiology

of *Campylobacter* Infection," *Clin Microbiol Rev,* Vol. 28, No. 3, 2015, pp. 687-720.

[2] M. B. Batz, E. Henke and B. Kowalcyk, "Long-term consequences of foodborne infections,"

Infect Dis Clin North Am, Vol. 27, No. 3, 2013, pp. 599-616.

[3] R. E. Black, M. M. Levine, M. L. Clements, T. P. Hughes and M. J. Blaser, "Experimental

Campylobacter jejuni infection in humans," *J Infect Dis,* Vol. 157, No. 3, 1988, pp. 472-479.

[4] J. P. van Spreeuwel, G. C. Duursma, C. J. Meijer, R. Bax, P. C. Rosekrans and J. Lindeman,

"*Campylobacter* colitis: histological immunohistochemical and ultrastructural findings," *Gut,*

Vol. 26, No. 9, 1985, pp. 945-951.

[5] F. A. Klipstein, R. F. Engert, H. Short and E. A. Schenk, "Pathogenic properties of

Campylobacter jejuni: assay and correlation with clinical manifestations," *Infect Immun,* Vol. 50,

No. 1, 1985, pp. 43-49.

[6] J. L. Fauchere, A. Rosenau, M. Veron, E. N. Moyen, S. Richard and A. Pfister, "Association

with HeLa cells of *Campylobacter jejuni* and *Campylobacter coli* isolated from human feces,"

Infect Immun, Vol. 54, No. 2, 1986, pp. 283-287.

[7] C. Centers for Disease and Prevention. "Campylobacteriosis 2015 Case Definition." National

Notifiable Diseases Surveillance System (NNDSS), 2015. Print.

[8] K. K. Nyati and R. Nyati, "Role of *Campylobacter jejuni* infection in the pathogenesis of

Guillain-Barré syndrome: an update," *Biomed Res Int,* Vol. 2013, 2013, p. 852195.

[9] E. Schorling, S. Knorr, S. Lick, P. Steinberg and D. A. Brüggemann, "Probability of sequelae

following *Campylobacter* spp. infections: Update of systematic reviews and meta-analyses,"

Public Health Challenges, Vol. 2, No. 4, 2023, p. e145.

[10] P. Harvey, T. Battle and S. Leach, "Different invasion phenotypes of *Campylobacter* isolates

in Caco-2 cell monolayers," *J Med Microbiol,* Vol. 48, No. 5, 1999, pp. 461-469.

[11] D. M. Tracz, M. Keelan, J. Ahmed-Bentley, A. Gibreel, K. Kowalewska-Grochowska and D.

E. Taylor, "pVir and bloody diarrhea in *Campylobacter jejuni* enteritis," *Emerg Infect Dis,* Vol.

11, No. 6, 2005, pp. 838-843.

[12] R. P. L. Louwen, A. van Belkum, J. A. Wagenaar, Y. Doorduyn, R. Achterberg and H. P.

Endtz, "Lack of Association between the Presence of the pVir Plasmid and Bloody Diarrhea in

Campylobacter jejuni Enteritis," *Journal of Clinical Microbiology,* Vol. 44, No. 5, 2006, pp. 1867-

1868.

[13] W. M. Johnson and H. Lior, "Toxins produced by *Campylobacter jejuni* and *Campylobacter*

coli," *Lancet,* Vol. 1, No. 8370, 1984, pp. 229-230.

[14] W. P. Yeen, S. D. Puthucheary and T. Pang, "Demonstration of a cytotoxin from

Campylobacter jejuni," *J Clin Pathol,* Vol. 36, No. 11, 1983, pp. 1237-1240.

[15] R. L. Guerrant, C. A. Wanke, R. A. Pennie, L. J. Barrett, A. A. Lima and A. D. O'Brien,

"Production of a unique cytotoxin by *Campylobacter jejuni,*" *Infect Immun,* Vol. 55, No. 10, 1987,

pp. 2526-2530.

[16] H. Goossens, J.-P. Butzler and Y. Takeda, "Demonstration of cholera-like enterotoxin

production by *Campylobacter jejuni*," *FEMS Microbiology Letters,* Vol. 29, No. 1-2, 1985, pp.

73-76.

[17] G. I. Perez-Perez, D. L. Cohn, R. L. Guerrant, C. M. Patton, L. B. Reller and M. J. Blaser,

"Clinical and immunologic significance of cholera-like toxin and cytotoxin production by

Campylobacter species in patients with acute inflammatory diarrhea in the USA," *J Infect Dis,*

Vol. 160, No. 3, 1989, pp. 460-468.

[18] G. M. Ruiz-Palacios, Y. López-Vidal, J. Torres and N. Torres, "Serum antibodies to heat-

labile enterotoxin of *Campylobacter jejuni*," *J Infect Dis,* Vol. 152, No. 2, 1985, pp. 413-416.

[19] F. A. Klipstein and R. F. Engert, "Properties of crude *Campylobacter jejuni* heat-labile

enterotoxin," *Infect Immun,* Vol. 45, No. 2, 1984, pp. 314-319.

[20] B. Wysok, J. Wojtacka, A. Wiszniewska-Łaszczych, M. Sołtysiuk and A. Kobuszewska, "The

Enterotoxin Production and Antimicrobial Resistance of *Campylobacter* Strains Originating from

Slaughter Animals," *Pathogens,* Vol. 11, No. 10, 2022.

[21] X. Zang, H. Lv, H. Tang, X. Jiao and J. Huang, "Capsular Genotype and Lipooligosaccharide

Class Associated Genomic Characterizations of *Campylobacter jejuni* Isolates From Food

Animals in China," *Frontiers in Microbiology,* Vol. Volume 12 - 2021, 2021.

[22] A. Culebro, J. Revez, B. Pascoe, Y. Friedmann, M. D. Hitchings, J. Stupak, S. K. Sheppard,

J. Li and M. Rossi, "Large Sequence Diversity within the Biosynthesis Locus and Common

Biochemical Features of *Campylobacter coli* Lipooligosaccharides," *J Bacteriol,* Vol. 198, No.

20, 2016, pp. 2829-2840.

[23] S. F. Mouftah, J. F. Cobo-Díaz, A. Álvarez-Ordóñez, A. Mousa, J. K. Calland, B. Pascoe, S.

K. Sheppard and M. Elhadidy, "Stress resistance associated with multi-host transmission and

enhanced biofilm formation at 42 °C among hyper-aerotolerant generalist *Campylobacter jejuni*,"

Food Microbiology, Vol. 95, 2021, p. 103706.

[24] S. Peters, B. Pascoe, Z. Wu, S. C. Bayliss, X. Zeng, A. Edwinson, S. Veerabadhran-

Gurunathan, S. Jawahir, J. K. Calland, E. Mourkas, R. Patel, T. Wiens, M. Decuir, D. Boxrud, K.

Smith, C. T. Parker, G. Farrugia, Q. Zhang, S. K. Sheppard and M. Grover, "*Campylobacter jejuni*

genotypes are associated with post-infection irritable bowel syndrome in humans," *Commun Biol,*

Vol. 4, No. 1, 2021, p. 1015.

[25] D. A. John, L. K. Williams, V. Kanamarlapudi, T. J. Humphrey and T. S. Wilkinson, "The

Bacterial Species *Campylobacter jejuni* Induce Diverse Innate Immune Responses in Human and

Avian Intestinal Epithelial Cells," *Frontiers in Microbiology,* Vol. 8, 2017.

[26] C. Cayrou, N. A. Barratt, J. M. Ketley and C. D. Bayliss, "Phase Variation During Host

Colonization and Invasion by *Campylobacter jejuni* and Other *Campylobacter* Species," *Front*

Microbiol, Vol. 12, 2021, p. 705139.

[27] R. L. Wallace, D. M. Cribb, D. M. Bulach, D. J. Ingle, K. G. Joensen, E. M. Nielsen, P.

Leekitcharoenphon, K. Stingl and M. D. Kirk, "*Campylobacter jejuni* ST50, a pathogen of global

importance: A comparative genomic analysis of isolates from Australia, Europe and North

America," *Zoonoses Public Health,* Vol. 68, No. 6, 2021, pp. 638-649.

[28] S. F. Mouftah, B. Pascoe, J. K. Calland, E. Mourkas, N. Tonkin, C. Lefevre, D. Deuker, S.

Smith, H. Wickenden, M. D. Hitchings, S. K. Sheppard and M. Elhadidy, "Local accessory gene

sharing among Egyptian *Campylobacter* potentially promotes the spread of antimicrobial

resistance," *Microbial Genomics,* Vol. 8, No. 6, 2022.

[29] E. Mourkas, A. J. Taylor, G. Méric, S. C. Bayliss, B. Pascoe, L. Mageiros, J. K. Calland, M.

D. Hitchings, A. Ridley, A. Vidal, K. J. Forbes, N. J. C. Strachan, C. T. Parker, J. Parkhill, K. A.

Jolley, A. J. Cody, M. C. J. Maiden, D. J. Kelly and S. K. Sheppard, "Agricultural intensification

and the evolution of host specialism in the enteric pathogen *Campylobacter jejuni,*" *Proceedings*

of the National Academy of Sciences, Vol. 117, No. 20, 2020, pp. 11018-11028.

[30] B. Pascoe, L. K. Williams, J. K. Calland, G. Meric, M. D. Hitchings, M. Dyer, J. Ryder, S.

Shaw, B. S. Lopes, C. Chintoan-Uta, E. Allan, A. Vidal, C. Fearnley, P. Everest, J. A. Pachebat,

T. A. Cogan, M. P. Stevens, T. J. Humphrey, T. S. Wilkinson, A. J. Cody, F. M. Colles, K. A.

Jolley, M. C. J. Maiden, N. Strachan, B. M. Pearson, D. Linton, B. W. Wren, J. Parkhill, D. J.

Kelly, A. H. M. van Vliet, K. J. Forbes and S. K. Sheppard, "Domestication of *Campylobacter jejuni* NCTC 11168," *Microbial Genomics,* Vol. 5, No. 7, 2019.

[31] J. Parkhill, B. W. Wren, K. Mungall, J. M. Ketley, C. Churcher, D. Basham, T. Chillingworth, R. M. Davies, T. Feltwell, S. Holroyd, K. Jagels, A. V. Karlyshev, S. Moule, M. J. Pallen, C. W. Penn, M. A. Quail, M. A. Rajandream, K. M. Rutherford, A. H. van Vliet, S. Whitehead and B. G. Barrell, "The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences," *Nature,* Vol. 403, No. 6770, 2000, pp. 665-668.

[32] O. Gundogdu, S. D. Bentley, M. T. Holden, J. Parkhill, N. Dorrell and B. W. Wren, "Re-annotation and re-analysis of the *Campylobacter jejuni* NCTC11168 genome sequence," *BMC genomics,* Vol. 8, No. 1, 2007, p. 162.

[33] K. K. Cooper, M. A. Cooper, A. Zuccolo and L. A. Joens, "Re-sequencing of a virulent strain of *Campylobacter jejuni* NCTC11168 reveals potential virulence factors," *Res Microbiol,* Vol. 164, No. 1, 2013, pp. 6-11.

[34] C. Gaynor Erin, S. Cawthraw, G. Manning, K. MacKichan Joanna, S. Falkow and G. Newell Diane, "The Genome-Sequenced Variant of *Campylobacter jejuni* NCTC 11168 and the Original Clonal Clinical Isolate Differ Markedly in Colonization, Gene Expression, and Virulence-Associated Phenotypes," *Journal of Bacteriology,* Vol. 186, No. 2, 2004, pp. 503-517.

[35] M. Stahl, J. Butcher and A. Stintzi, "Nutrient acquisition and metabolism by *Campylobacter jejuni,*" *Front Cell Infect Microbiol,* Vol. 2, 2012, p. 5.

[36] Y. He, S. Reed and T. P. Strobaugh, Jr., "Complete Genome Sequence and Annotation of *Campylobacter jejuni* YH003, Isolated from Retail Chicken," *Microbiol Resour Announc,* Vol. 9, No. 4, 2020.

[37] M. McClelland, K. E. Sanderson, J. Spieth, S. W. Clifton, P. Latreille, L. Courtney, S.

Porwollik, J. Ali, M. Dante, F. Du, S. Hou, D. Layman, S. Leonard, C. Nguyen, K. Scott, A.

Holmes, N. Grewal, E. Mulvaney, E. Ryan, H. Sun, L. Florea, W. Miller, T. Stoneking, M. Nhan,

R. Waterston and R. K. Wilson, "Complete genome sequence of *Salmonella enterica* serovar

Typhimurium LT2," *Nature,* Vol. 413, No. 6858, 2001, pp. 852-856.

[38] D. Hofreuter, V. Novik and J. E. Galán, "Metabolic diversity in *Campylobacter jejuni*

enhances specific tissue colonization," *Cell Host Microbe,* Vol. 4, No. 5, 2008, pp. 425-433.

[39] A. X. van der Stel, A. van Mourik, P. Łaniewski, J. P. van Putten, E. K. Jagusztyn-Krynicka

and M. M. Wösten, "The *Campylobacter jejuni* RacRS two-component system activates the

glutamate synthesis by directly upregulating γ-glutamyltranspeptidase (GGT)," *Front Microbiol,*

Vol. 6, 2015, p. 567.

[40] C. P. de Haan, A. K. Llarena, J. Revez and M. L. Hänninen, "Association of *Campylobacter*

jejuni metabolic traits with multilocus sequence types," *Appl Environ Microbiol,* Vol. 78, No. 16,

2012, pp. 5550-5554.

[41] D. Hofreuter, J. Tsai, R. O. Watson, V. Novik, B. Altman, M. Benitez, C. Clark, C. Perbost,

T. Jarvie, L. Du and J. E. Galán, "Unique features of a highly pathogenic *Campylobacter jejuni*

strain," *Infect Immun,* Vol. 74, No. 8, 2006, pp. 4694-4707.

[42] M. Stahl, L. M. Friis, H. Nothaft, X. Liu, J. Li, C. M. Szymanski and A. Stintzi, "L-fucose

utilization provides *Campylobacter jejuni* with a competitive advantage," *Proc Natl Acad Sci U S*

A, Vol. 108, No. 17, 2011, pp. 7194-7199.

[43] W. T. Muraoka and Q. Zhang, "Phenotypic and genotypic evidence for L-fucose utilization

by *Campylobacter jejuni,*" *J Bacteriol,* Vol. 193, No. 5, 2011, pp. 1065-1075.

[44] P. S. Middendorf, L. M. Wijnands, S. Boeren, A. L. Zomer, W. F. Jacobs-Reitsma, H. M. W.

den Besten and T. Abee, "Activation of the l-fucose utilization cluster in *Campylobacter jejuni*

induces proteomic changes and enhances Caco-2 cell invasion and fibronectin binding," *Heliyon,*

Vol. 10, No. 16, 2024, p. e34996.

[45] F. Meurens, A. Summerfield, H. Nauwynck, L. Saif and V. Gerdts, "The pig: a model for

human infectious diseases," *Trends Microbiol,* Vol. 20, No. 1, 2012, pp. 50-57.

[46] J. Vítovec, B. Koudela, J. Štěrba, I. Tomancová, Z. Matyáš and P. Vladík, "The gnotobiotic

piglet as a model for the pathogenesis of *Campylobacter jejuni* infection," *Zentralblatt für*

Bakteriologie, Vol. 271, No. 1, 1989, pp. 91-103.

[47] J. E. Line, K. L. Hiett, J. Guard-Bouldin and B. S. Seal, "Differential carbon source utilization

by *Campylobacter jejuni* 11168 in response to growth temperature variation," *J Microbiol*

Methods, Vol. 80, No. 2, 2010, pp. 198-202.

[48] M. Reuter, A. Mallett, B. M. Pearson and A. H. van Vliet, "Biofilm formation by

Campylobacter jejuni is increased under aerobic conditions," *Appl Environ Microbiol,* Vol. 76,

No. 7, 2010, pp. 2122-2128.

[49] R. J. Reeser, , R. T. Medler, , S. J. Billington, , B. H. Jost, , L. A. Joens and "Characterization

of *Campylobacter jejuni* biofilms under defined growth conditions." 2007/01/26 ed: American

Society for Microbiology, 2007. 1908-1913. Vol. 73. Print.

[50] D. Hermans, A. Martel, K. Van Deun, M. Verlinden, F. Van Immerseel, A. Garmyn, W.

Messens, M. Heyndrickx, F. Haesebrouck and F. Pasmans, "Intestinal mucus protects

Campylobacter jejuni in the ceca of colonized broiler chickens against the bactericidal effects of

medium-chain fatty acids," *Poultry Science,* Vol. 89, No. 6, 2010, pp. 1144-1155.

[51] I. H. Barnes, M. C. Bagnall, D. D. Browning, S. A. Thompson, G. Manning and D. G. Newell,

"Gamma-glutamyl transpeptidase has a role in the persistent colonization of the avian gut by

Campylobacter jejuni," *Microb Pathog,* Vol. 43, No. 5-6, 2007, pp. 198-207.

[52] M. Stahl and B. A. Vallance, "Insights into *Campylobacter jejuni* colonization of the

mammalian intestinal tract using a novel mouse model of infection," *Gut Microbes,* Vol. 6, No.

2, 2015, pp. 143-148.

[53] D. Hermans, K. Van Deun, A. Martel, F. Van Immerseel, W. Messens, M. Heyndrickx, F.

Haesebrouck and F. Pasmans, "Colonization factors of *Campylobacter jejuni* in the chicken gut,"

Veterinary Research, Vol. 42, No. 1, 2011, p. 82.

[54] T. M. Wassenaar, B. A. van der Zeijst, R. Ayling and D. G. Newell, "Colonization of chicks

by motility mutants of *Campylobacter jejuni* demonstrates the importance of flagellin A

expression," *J Gen Microbiol,* Vol. 139 Pt 6, 1993, pp. 1171-1175.

[55] P. Guerry, "*Campylobacter* flagella: not just for motility," *Trends Microbiol,* Vol. 15, No.

10, 2007, pp. 456-461.

[56] M. Zimmer, H. Barnhart, U. Idris and M. D. Lee, "Detection of *Campylobacter jejuni* strains

in the water lines of a commercial broiler house and their relationship to the strains that colonized

the chickens," *Avian Dis,* Vol. 47, No. 1, 2003, pp. 101-107.

[57] J. M. Garber, H. Nothaft, B. Pluvinage, M. Stahl, X. Bian, S. Porfirio, A. Enriquez, J. Butcher,

H. Huang, J. Glushka, E. Line, J. A. Gerlt, P. Azadi, A. Stintzi, A. B. Boraston and C. M.

Szymanski, "The gastrointestinal pathogen *Campylobacter jejuni* metabolizes sugars with

potential help from commensal *Bacteroides vulgatus,*" *Communications Biology,* Vol. 3, No. 1,

2020, p. 2.

[58] J. Killer and M. Marounek, "Fermentation of mucin by bifidobacteria from rectal samples of

humans and rectal and intestinal samples of animals," *Folia Microbiol (Praha),* Vol. 56, No. 2,

2011, pp. 85-89.

[59] P. Ruas-Madiedo, M. Gueimonde, M. Fernández-García, C. G. de los Reyes-Gavilán and A.

Margolles, "Mucin degradation by *Bifidobacterium* strains isolated from the human intestinal

microbiota," *Appl Environ Microbiol,* Vol. 74, No. 6, 2008, pp. 1936-1940.

[60] M. Derrien, M. C. Collado, K. Ben-Amor, S. Salminen and W. M. de Vos, "The Mucin

degrader *Akkermansia muciniphila* is an abundant resident of the human intestinal tract," *Appl*

Environ Microbiol, Vol. 74, No. 5, 2008, pp. 1646-1648.

[61] B. P. Ganesh, R. Klopfleisch, G. Loh and M. Blaut, "Commensal *Akkermansia muciniphila*

Exacerbates Gut Inflammation in *Salmonella* Typhimurium-Infected Gnotobiotic Mice," *PLoS*

One, Vol. 8, No. 9, 2013, p. e74963.

[62] I. Biasato, I. Ferrocino, E. Grego, S. Dabbou, F. Gai, L. Gasco, L. Cocolin, M. T. Capucchio

and A. Schiavone, "Gut Microbiota and Mucin Composition in Female Broiler Chickens Fed Diets

including Yellow Mealworm (*Tenebrio molitor*, L.)," *Animals,* Vol. 9, No. 5, 2019, p. 213.

[63] A. Smirnov, R. Perez, E. Amit-Romach, D. Sklan and Z. Uni, "Mucin Dynamics and

Microbial Populations in Chicken Small Intestine Are Changed by Dietary Probiotic and

Antibiotic Growth Promoter Supplementation," *The Journal of Nutrition,* Vol. 135, No. 2, 2005,

pp. 187-192.

[64] I. Biasato, I. Ferrocino, E. Biasibetti, E. Grego, S. Dabbou, A. Sereno, F. Gai, L. Gasco, A.

Schiavone, L. Cocolin and M. T. Capucchio, "Modulation of intestinal microbiota, morphology

and mucin composition by dietary insect meal inclusion in free-range chickens," *BMC Veterinary*

Research, Vol. 14, No. 1, 2018, p. 383.

[65] J. A. Ferreyra, K. M. Ng and J. L. Sonnenburg, "The Enteric Two-Step: nutritional strategies

of bacterial pathogens within the gut," *Cell Microbiol,* Vol. 16, No. 7, 2014, pp. 993-1003.

[66] B. Shuoker, M. J. Pichler, C. Jin, H. Sakanaka, H. Wu, A. M. Gascueña, J. Liu, T. S. Nielsen,

J. Holgersson, E. Nordberg Karlsson, N. Juge, S. Meier, J. P. Morth, N. G. Karlsson and M. Abou

Hachem, "Sialidases and fucosidases of *Akkermansia muciniphila* are crucial for growth on mucin

and nutrient sharing with mucus-associated gut bacteria," *Nature Communications,* Vol. 14, No.

1, 2023, p. 1833.

[67] Y. M. C. A. Luijkx, N. M. C. Bleumink, J. Jiang, H. S. Overkleeft, M. M. S. M. Wösten, K.

Strijbis and T. Wennekes, "Fucosidases facilitate growth and invasion of *Campylobacter jejuni* in

the presence of mucins," *Cellular microbiology,* Vol. 22, No. 12, 2020, p. e13252.

[68] J. T. LAMONT, "Mucus: The Front Line of Intestinal Mucosal Defense," *Annals of the New*

York Academy of Sciences, Vol. 664, No. 1, 1992, pp. 190-201.

[69] M. B. Hugdahl, J. T. Beery and M. P. Doyle, "Chemotactic behavior of *Campylobacter*

jejuni," *Infect Immun,* Vol. 56, No. 6, 1988, pp. 1560-1566.

[70] L. E. Tailford, E. H. Crost, D. Kavanaugh and N. Juge, "Mucin glycan foraging in the human

gut microbiome," *Front Genet,* Vol. 6, 2015, p. 81.

[71] K. S. Bergstrom and L. Xia, "Mucin-type O-glycans and their roles in intestinal homeostasis,"

Glycobiology, Vol. 23, No. 9, 2013, pp. 1026-1037.

[72] P. Tsirtsikos, K. Fegeros, C. Balaskas, A. Kominakis and K. C. Mountzouris, "Dietary

probiotic inclusion level modulates intestinal mucin composition and mucosal morphology in

broilers," *Poultry Science,* Vol. 91, No. 8, 2012, pp. 1860-1868.

[73] R. Dwivedi, H. Nothaft, J. Garber, L. Xin Kin, M. Stahl, A. Flint, A. H. van Vliet, A. Stintzi

and C. M. Szymanski, "L-fucose influences chemotaxis and biofilm formation in *Campylobacter*

jejuni," *Mol Microbiol,* Vol. 101, No. 4, 2016, pp. 575-589.

[74] P. S. Middendorf, A. L. Zomer, I. L. Bergval, W. F. Jacobs-Reitsma, H. M. W. den Besten

and T. Abee, "Host associations of *Campylobacter jejuni* and *Campylobacter coli* isolates carrying

the L-fucose or d-glucose utilization cluster," *International Journal of Food Microbiology,* Vol.

425, 2024, p. 110855.

[75] G. Shani, L. Hoeflinger Jennifer, E. Heiss Britta, F. Masarweh Chad, A. Larke Jules, M.

Jensen Nick, S. Wickramasinghe, C. Davis Jasmine, E. Goonatilleke, A. El-Hawiet, L. Nguyen, S.

Klassen John, M. Slupsky Carolyn, B. Lebrilla Carlito and A. Mills David, "Fucosylated Human

Milk Oligosaccharide Foraging within the species *Bifidobacterium pseudocatenulatum* Is Driven

by Glycosyl Hydrolase Content and Specificity," *Applied and Environmental Microbiology,* Vol.

88, No. 2, 2022, pp. e01707-01721.

[76] M. Moya-Gonzálvez Eva, N. Peña-Gil, A. Rubio-del-Campo, M. Coll-Marqués José, R.

Gozalbo-Rovira, V. Monedero, J. Rodríguez-Díaz and J. Yebra María, "Infant Gut Microbial

Metagenome Mining of α-l-Fucosidases with Activity on Fucosylated Human Milk

Oligosaccharides and Glycoconjugates," *Microbiology Spectrum,* Vol. 10, No. 4, 2022, pp.

e01775-01722.

[77] X. Bian, M. Garber Jolene, K. Cooper Kerry, S. Huynh, J. Jones, K. Mills Michael, D. Rafala,

D. Nasrin, L. Kotloff Karen, T. Parker Craig, M. Tennant Sharon, G. Miller William and M.

Szymanski Christine, "*Campylobacter* Abundance in Breastfed Infants and Identification of a New

Species in the Global Enterics Multicenter Study," *mSphere,* Vol. 5, No. 1, 2020, pp.

10.1128/msphere.00735-00719.

[78] D. Chen, S. L. McKune, N. Singh, J. Yousuf Hassen, W. Gebreyes, M. J. Manary, K. Bardosh,

Y. Yang, N. Diaz, A. Mohammed, Y. Terefe, K. T. Roba, M. Ketema, N. Ameha, N. Assefa, G.

Rajashekara, L. Deblais, M. Ghanem, G. Yimer and A. H. Havelaar, "*Campylobacter*

Colonization, Environmental Enteric Dysfunction, Stunting, and Associated Risk Factors Among

Young Children in Rural Ethiopia: A Cross-Sectional Study From the *Campylobacter* Genomics

and Environmental Enteric Dysfunction (CAGED) Project," *Frontiers in Public Health,* Vol.

Volume 8 - 2020, 2021.

[79] B. Zhou, J. M. Garber, J. Vlach, P. Azadi, K. K. S. Ng, J. C. Escalante-Semerena and C. M.

Szymanski, "*Campylobacter jejuni* uses energy taxis and a dehydrogenase enzyme for l-fucose

chemotaxis," *mBio,* Vol. 14, No. 6, 2023, p. e0273223.

[80] N. Li, P. Lewis, D. Samuelson, K. Liboni and J. Neu, "Glutamine regulates Caco-2 cell tight

junction proteins," *Am J Physiol Gastrointest Liver Physiol,* Vol. 287, No. 3, 2004, pp. G726-733.

[81] M.-H. Kim and H. Kim, "The Roles of Glutamine in the Intestine and Its Implication in

Intestinal Diseases," *International Journal of Molecular Sciences,* Vol. 18, No. 5, 2017, p. 1051.

[82] B. J. Deters and M. Saleem, "The role of glutamine in supporting gut health and

neuropsychiatric factors," *Food Science and Human Wellness,* Vol. 10, No. 2, 2021, pp. 149-154.

[83] B. Wang, Z. Wu, Y. Ji, K. Sun, Z. Dai and G. Wu, "L-Glutamine Enhances Tight Junction

Integrity by Activating CaMK Kinase 2–AMP-Activated Protein Kinase Signaling in Intestinal

Porcine Epithelial Cells12," *The Journal of Nutrition,* Vol. 146, No. 3, 2016, pp. 501-508.

[84] V. Cruzat, M. Macedo Rogero, K. Noel Keane, R. Curi and P. Newsholme, "Glutamine:

Metabolism and Immune Function, Supplementation and Clinical Translation," *Nutrients,* Vol.

10, No. 11, 2018.

[85] B. Feodoroff, P. Ellström, H. Hyytiäinen, S. Sarna, M. L. Hänninen and H. Rautelin,

"*Campylobacter jejuni* isolates in Finnish patients differ according to the origin of infection," *Gut*

Pathog, Vol. 2, No. 1, 2010, p. 22.

[86] P. Floch, V. Pey, M. Castroviejo, J. W. Dupuy, M. Bonneu, A. H. de la Guardia, V. Pitard, F.

Mégraud and P. Lehours, "Role of *Campylobacter jejuni* gamma-glutamyl transpeptidase on

epithelial cell apoptosis and lymphocyte proliferation," *Gut Pathog,* Vol. 6, 2014, p. 20.

[87] C. C. Grant, M. E. Konkel, W. Cieplak, Jr. and L. S. Tompkins, "Role of flagella in adherence,

internalization, and translocation of *Campylobacter jejuni* in nonpolarized and polarized epithelial

cell cultures," *Infect Immun,* Vol. 61, No. 5, 1993, pp. 1764-1771.

[88] P. Malik-Kale, B. H. Raphael, C. T. Parker, L. A. Joens, J. D. Klena, B. Quiñones, A. M.

Keech and M. E. Konkel, "Characterization of genetically matched isolates of *Campylobacter*

jejuni reveals that mutations in genes involved in flagellar biosynthesis alter the organism's

virulence potential," *Appl Environ Microbiol,* Vol. 73, No. 10, 2007, pp. 3123-3136.

[89] I. Nachamkin, X. H. Yang and N. J. Stern, "Role of *Campylobacter jejuni* flagella as

colonization factors for three-day-old chicks: analysis with flagellar mutants," *Appl Environ*

Microbiol, Vol. 59, No. 5, 1993, pp. 1269-1273.

[90] D. G. Newell, H. McBride and J. M. Dolby, "Investigations on the role of flagella in the

colonization of infant mice with *Campylobacter jejuni* and attachment of *Campylobacter jejuni* to

human epithelial cell lines," *J Hyg (Lond),* Vol. 95, No. 2, 1985, pp. 217-227.

[91] P. M. Burnham and D. R. Hendrixson, "*Campylobacter jejuni*: collective components

promoting a successful enteric lifestyle," *Nat Rev Microbiol,* Vol. 16, No. 9, 2018, pp. 551-565.

[92] N. Luo, O. Sahin, J. Lin, L. O. Michel and Q. Zhang, "In vivo selection of *Campylobacter*

isolates with high levels of fluoroquinolone resistance associated with gyrA mutations and the

function of the CmeABC efflux pump," *Antimicrob Agents Chemother,* Vol. 47, No. 1, 2003, pp. 390-394.

[93] O. Sahin, N. Luo, S. Huang and Q. Zhang, "Effect of *Campylobacter*-specific maternal antibodies on *Campylobacter jejuni* colonization in young chickens," *Appl Environ Microbiol,* Vol. 69, No. 9, 2003, pp. 5372-5379.

[94] K. K. Cooper, M. A. Cooper, A. Zuccolo, B. Law and L. A. Joens, "Complete genome sequence of *Campylobacter jejuni* strain S3," *J Bacteriol,* Vol. 193, No. 6, 2011, pp. 1491-1492.

[95] S. K. Sheppard, J. F. Dallas, N. J. Strachan, M. MacRae, N. D. McCarthy, D. J. Wilson, F. J. Gormley, D. Falush, I. D. Ogden, M. C. Maiden and K. J. Forbes, "*Campylobacter* genotyping to determine the source of human infection," *Clin Infect Dis,* Vol. 48, No. 8, 2009, pp. 1072-1078.

[96] P. M. Araújo, E. Batista, M. H. Fernandes, M. J. Fernandes, L. T. Gama and M. J. Fraqueza, "Assessment of biofilm formation by *Campylobacter* spp. isolates mimicking poultry slaughterhouse conditions," *Poult Sci,* Vol. 101, No. 2, 2022, p. 101586.

[97] B. Pascoe, G. Méric, S. Murray, K. Yahara, L. Mageiros, R. Bowen, N. H. Jones, R. E. Jeeves, H. M. Lappin-Scott, H. Asakura and S. K. Sheppard, "Enhanced biofilm formation and multi-host transmission evolve from divergent genetic backgrounds in *Campylobacter jejuni,*" *Environ Microbiol,* Vol. 17, No. 11, 2015, pp. 4779-4789.

[98] G. O'Toole, H. B. Kaplan and R. Kolter, "Biofilm formation as microbial development," *Annu Rev Microbiol,* Vol. 54, 2000, pp. 49-79.

[99] J. H. Pringle and M. Fletcher, "Influence of substratum hydration and adsorbed macromolecules on bacterial attachment to surfaces," *Appl Environ Microbiol,* Vol. 51, No. 6, 1986, pp. 1321-1325.

[100] W. Wang, Y. Cao, J. Li, S. Lu, H. Ge, S. Pan, X. Pan and L. Wang, "The impact of osmotic

stresses on the biofilm formation, immunodetection, and morphology of *Aeromonas hydrophila,*"

Microbiological Research, Vol. 269, 2023, p. 127301.

[101] J. Reeser Ryan, T. Medler Robert, J. Billington Stephen, B. H. Jost and A. Joens Lynn,

"Characterization of *Campylobacter jejuni* Biofilms under Defined Growth Conditions," *Applied*

and Environmental Microbiology, Vol. 73, No. 6, 2007, pp. 1908-1913.

Figure 1

Figure 2

Figure 3

A

B

Figure 5

Figure 6