1 **Title:** More running causes more ocular dominance plasticity in mouse primary visual cortex:
2 new gated running wheel setup allows to quantify individual running behaviour of group-
3 housed mice.
4
5 **Abbreviated title:** More running, more ocular dominance plasticity
6

7 **Authors:** Cornelia Schöne[1,2], Jaya Sowkyadha Sathiyamani[1,2,3], Mihaela Guranda[1], Siegrid
8 Löwel[1,2]
9
Affiliation:
[1] University of Göttingen, Department of Systems Neuroscience, 37075 Göttingen,
Germany
[2] Göttingen Campus Institute for Dynamics of Biological Networks, 37075 Göttingen,
Germany
[3] International Max Planck Research School for Neurosciences, 37075 Göttingen,
Germany
17
18 **Corresponding author:**
19 Cornelia Schöne
20 email: cornelia.schoene@gwdg.de
21 orcid: https://orcid.org/0000-0003-2060-1084
22
23

24 **Graphical abstract:**

With our newly developed gated running wheel setup, we observed a striking correlation between individual running activity, and a measure of experience dependent plasticity in mouse primary visual cortex. More running caused more plasticity: running speed, running distance, total running time, number of running bouts and bout duration all correlated with a measure of visual cortical plasticity, the ocular dominance index. Thus, our observations add to the growing body of evidence that individual behavioural choices strongly affect individual brain plasticity.

45 **Acknowledgements:** We thank DFG (SFB889, Project B05 to S.L.) and the Dorothea
46 Schlözer-Program for funding. We thank Tobias Mühmer and Jan Hoffmann, for constructing
47 the gated running wheel cage, and Christina Ahlbrecht for technical assistance and animal
48 care.

Title: More running causes more ocular dominance plasticity in mouse primary visual cortex: new gated running wheel setup enables individual tracking of wheel running in group-housed mice.

Abstract

Environmental enrichment boosts neuronal plasticity of standard-cage raised (SC) mice. Since it becomes increasingly more important to track individual mouse behaviours and its influence on brain plasticity, we designed a gated running wheel (gRW) setup allowing to correlate wheel running with neuronal plasticity, using the established paradigm of ocular dominance (OD)-plasticity after monocular deprivation (MD).

After SC-rearing until adulthood (>P110), group-housed mice were transferred to gRW cages, that provided an additional running wheel compartment for tracking individual wheel activity via implanted RFID chips. Notably, individual running parameters varied enormously: mice ran from close to 0 to ~20 km across the 7 days of gRW experience, with on average running 0-3.96 km in 0-3.85 h/d and running bouts lasting from <1 up to 10 min, while running at a speed of 6-26 cm/s. OD-plasticity in V1 after 7 days of MD in the gRW was visualized using intrinsic signal optical imaging, and compared to control gRW-mice without MD via calculation of an OD-index. Most, notably - while wheel running enabled OD-plasticity - *individual* running parameters correlated with *individual* OD-indices after MD: Mice running longer distances, for longer time, at higher speeds and with longer and more frequent bouts displayed more experience-dependent V1-plasticity. In turn, a composite measure of overall running wheel activity derived from principal component analysis of running parameters accounted for 65% of inter-individual variability of OD-index following MD. Together our study demonstrates that interindividual variability of running behaviour is high, and mice intrinsically motivated to run more show enhanced V1-plasticity, underscoring the huge importance of analysing individual behavioural parameters together with any measure of brain plasticity. End

Keywords: Voluntary wheel running, ocular dominance plasticity, primary visual cortex, inter-individual variability, behavioural tracking

Introduction

While the natural environment of humans and mice is complex, that of laboratory rodents has been simplified and standardised in the past decades to an extent that deprives the animals of essential behaviours (Löwel et al., 2017; Cait et al., 2024). For instance, standard cage rearing (SC) leaves minimal room for the expression of individual traits and both behavioural and physiological interindividual variability is further reduced by using inbred strains lacking genetic variability (Wahlsten et al., 2006; Körholz et al., 2018). Recently, a number of laboratories demonstrated that raising animals under less deprived rearing conditions – in so-called "enriched environments", i.e. housing in larger social groups and/or larger cages, with regularly changed mazes to navigate through and running wheels for voluntary physical exercise – elicits remarkable effects on brain wiring and plasticity across molecular, anatomical, and functional levels when compared to animals raised in an SC environment (Sale et al., 2007; Fabel et al., 2009; Baroncelli et al., 2010; Di Garbo et al., 2011; Greifzu et al., 2014; Kalogeraki et al., 2014; Löwel et al., 2017; Stryker and Löwel, 2018; Bogado Lopes et al., 2023).

Both environmental enrichment, but also just voluntary wheel running have been shown to boost experience-dependent changes in rodent primary visual cortex (V1) (Sale et al., 2007; Baroncelli et al., 2010; Greifzu et al., 2014; Kalogeraki et al., 2014; Kaneko and Stryker, 2014), using the established model of ocular dominance (OD) plasticity after monocular deprivation (MD)(Gordon and Stryker, 1996; Cang et al., 2005): when one eye is closed for few days, the relative V1-activation strength is shifted towards the open eye (Cang et al., 2005; Gordon and Stryker, 1996). SC-mice beyond P110 do no longer show OD-plasticity with 7 days of MD (Lehmann and Löwel, 2008; Sato and Stryker, 2008), and need ~7 weeks for V1-activation changes (Hosang et al., 2018)

While it is clearly established that running increases OD-plasticity in mice (Kalogeraki et al., 2014; Kaneko and Stryker, 2014) the consequences of interindividual variability of wheel running on experience-dependent V1-plasticity has not yet been studied. We therefore aimed to analyse the effects of individual running behaviour on OD-plasticity of adult (>P110) group housed SC-mice. As tracking individual wheel running activity of group housed mice is still challenging (Mayr et al., 2020; Reuser et al., 2022), we designed a custom-built gated running wheel (gRW) setup consisting of a rat-sized home cage connected to a compartment with a surveyed running wheel that allowed individual tracking of mice implanted with an RFID sensor. We quantified running distance, time, speed, bout number and bout length of running activity and tested whether they correlated with the OD-index, which compares visual stimulus evoked V1-activation via the ipsi- and contralateral eye and thus quantifies the magnitude of OD-plasticity (Cang et al., 2005).

Running parameters varied extensively between individual mice, with some mice running only few meters while others ran >20 km during their 7 days of running wheel exposure. Most notably, correlating the macro (running distance, time and speed) and micro (bout number and duration) architecture of running wheel activity of individual mice with their individual OD-index, quantifying OD-plasticity, revealed strong correlations with Pearson correlation coefficients ranging from -0.77 to -0.83: more active runners displayed stronger OD-plasticity. Our results underscore the importance of taking individual behavioural choices into account when analysing influences on brain plasticity.

Methods:

Animals and rearing conditions

In order to avoid conflict between group housed males, we limited our experiments to groups of 2-5 adult female C57Bl6/J mice (age range P127-225, called P160), growing up in Tecniplast 1284 standard mouse cages (SC; size: 17x32x19cm) until the day an MD/noMD was performed. After the MD/noMD, animals were cohoused in the gRW-setup, provided with *ad libitum* food and water, and maintained on a 12 h light dark cycle.. All experimental procedures complied with the National Institutes of Health guidelines for the use of laboratory animals and were approved by the local government of Lower Saxony, Germany (Niedersächsisches Landesamt für Verbraucherschutz und Lebensmittelsicherheit).

Optomotry, monocular deprivation and implantation of RFID tags

To ensure that all experimental mice had regular vision before testing OD-plasticity with MD, we measured the spatial frequency threshold (SFT) of the optomotor reflex of both eyes using an optomotor system (Prusky et al. 2004; Lehmann and Löwel, 2008). SFT values were within published values and differed by less than 0.02 cyc/deg between the two eyes in all mice.

Thereafter, the right eye was sutured shut (MD) for 7 d according to published protocols (Gordon and Stryker, 1996; Cang et al, 2005; Greifzu et al., 2014). In brief, mice were anaesthetized using 2.5 % isoflurane in 1:1 O_2/N_2, and kept at 1-1.5% isoflurane for stable anaesthesia. Body temperature was maintained at 37°C using a heat pad and rectal probe. After subcutaneous injection of Rimadyl (Carprofen, 5 mg/kg) and covering of eyes with Bepanthen eye cream, a local analgesic (Lidocaine, xylocaine gel 2 %, Aspen Pharma Trading Limited, Ireland) was applied to the eyelid of the right eye before trimming and closing of the eye with 2 mattress stiches (suture material: 7-0 Perma-Hand silk, Ethicon, 8.0mm diameter). The eyes of the animals were examined daily to ensure that the MD-eye stayed closed. Additionally, optomotor reflex thresholds (SFT) of the open eye were tested daily after MD, as a measure for functional MD, including the day of intrinsic signal optical imaging.

Cylindrical glass-covered radio frequency identification devices (RFID; length: 12.5 mm; diameter: 1.93 mm, Sparkfun Electronics, Colorado, USA) were implanted during the MD or noMD (control without MD) surgery to minimize mouse discomfort. First, a patch of fur was shaved in the animal's neck using small surgical scissors, then a local analgesic gel (Lidocaine, xylocaine gel 2 %, Aspen Pharma Trading Limited, Ireland) was applied to the exposed skin. A small incision was made and the chip carefully placed subcutaneously in the scruff to make it difficult for the mice to pick at and thereby reducing the risk of an infection. The incision was sealed with a simple interrupted stitch (suture material: 7-0 Perma-Hand silk, Ethicon, 8.0mm diameter). After MD/RFID implantation and complete recovery from anaesthesia, mice were moved to a gRW setup.

Gated running wheel setup (gRW)

In order to assess individual running wheel behaviour, we developed an open source gated running wheel (gRW) cage, which served as the home cage for the duration of the experiment (Figure 1A, Table 1). It consisted of a standard size rat cage (43cm x 27cm) connected to a separate gRW-compartment, equipped with the same RW as used in the enrichment cages in Greifzu et al., 2014 (Figure 1A). Wheel diameter was 12.5 cm, so that the calculated RW-circumference was 39.3 cm. RW-turns were registered using a hall sensor for detecting motion of two magnets attached at opposing sites of the RW. The mice needed to pass through a seesaw in order to reach the RW-compartment. Entering would flip the seesaw blocking other mice from entering (see video 1 and 2). To ensure the seesaw remained closed while a mouse was inside the RW-compartment, the seesaw was actively blocked from flipping back. When activity was detected by beam break sensors outside the RW-compartment, an electromagnet was activated to hold the seesaw closed. Attempts of mice to exit the RW-compartment were detected using beam break sensors inside the RW-compartment, which would deactivate the seesaw magnet. Seesaw position was registered through a roller switch.

Mice were chipped with radio frequency identification devices (RFID). The RFID-tags transmitted the encrypted individual ID to a USB-driven reading device (ID-20LA SparkFun Electronics, Colorado, USA) at 125 kHz via a custom-made RFID coil. Thus, each time a mouse passed underneath the RFID coil into the gRW compartment, RFID and time stamps were saved. All sensors were connected to the GPIO pins of a raspberry pi 4B+. In addition, a raspberry pi noir camera was used to record movement inside the RW-compartment at 40 Hz; camera data were used to confirm sensor data and to extract RW-turns when the Hall sensor failed. All data was recorded using custom written python scripts running on the raspberry pi. A 3D .pdf file illustrating dimensions of cage components and electronic assembly is provided under supplementary material (Model 1 and Figure S1, respectively).

Analysis of gated wheel running

In order to assign each RW-turn to individual mice, we first defined running bouts as running at a speed above 0.1 Hz, then the most recently detected RFID was assigned to all wheel turns within the respective running bout. When more than one RFID was detected since the last gate flip, this indicated that more than one mouse had entered the gRW compartment. This occasionally happened, due to two mice squeezing through the gate simultaneously, or due to failure of the magnetic seesaw lock. In this case, the RW-turns were labelled as ambiguous. Mice with ambiguous RW-turns of more than 7% were excluded from the RW-analysis, but included for analysis of OD-plasticity in Figure 2. For the remaining mice, on average, we quantified 1.7±0.5% ambiguous RW-turns (n=20 mice), confirming that the self-locking seesaw of the gRW was well suited to restrict access to the wheel compartment to individual mice.

Optical imaging of intrinsic signals and visual stimuli.

As a final step of the experiment, OD-plasticity was assessed after 7 days of MD/noMD in the gRW-setup, using optical imaging of intrinsic signals (Cang et al., 2005; Greifzu et al., 2014).

Surgery. Briefly, mice were box-anaesthetised with 2.5% halothane in O_2 and N_2O (1:1) and
injected with atropine (5mg/kg, s.c.; Franz Köhler Chemie), dexamethasone (8mg/kg s.c.;
Merck), and chlorprothixene (8mg/kg, i.m.; Sigma-Aldrich). After placing animals in a
stereotaxic frame, anaesthesia was maintained with 0.8% halothane in a 1:1 mixture of O_2
and N_2O.

Data acquisition and visual stimulation (Kalatsky and Stryker, 2003; Cang et al, 2005).
Mouse V1 responses were recorded through the skull using the "Fourier" imaging method of
Kalatsky and Stryker (2003) and optimised for the assessment of OD-plasticity (Cang et al.,
2005). V1-signals were visualised with a CCD camera (Dalsa 1M30) using a 135x50 mm
tandem lens configuration (Nikon) with red illumination light (610±10 nm). Active brain
regions absorb more red light and appear darker in the images. Frames were acquired at a
rate of 30 Hz, temporally binned to 7.5 Hz, and stored as 512x512-pixel images after spatial
binning of the camera image. Visual stimuli were presented on a 60 Hz refresh rate monitor
(21 inches; Accuvue HM-4921-D, Hitachi) positioned 25 cm from the eyes. Stimuli consisted
of white drifting horizontal bars (2° wide) limited to the contralateral binocular visual field (-5°
to 15°) as described previously (Greifzu et al., 2014). The amplitude component of the
optical signal represents the intensity of neuronal activation (expressed as fractional change
in reflectance times 10^{-4}) and was used to calculate an ocular dominance (OD) index. At
least three maps per animal were averaged by an experimenter blinded to the experimental
conditions to compute the OD-index as (C+I)/(C-I), with C and I representing the response
magnitudes of each pixel to visual stimulation of the contralateral (C) and ipsilateral (I) eye.

EXPERIMENTAL DESIGN AND STATISTICAL ANALYSES

Principal component analysis

Principal component analysis was performed using the MATLAB (R2022b) pca function on
wheel running parameters (distance, time, speed, bout duration and bout number) obtained
from combined noMD and MD mice in order to eliminate multicollinear variables which
effectively reduces dimensions of descriptive variables with the first principal component
(PC1) capturing the maximum variance of the data (86-90% of variance in wheel running
parameters, depending on whether average or individual data across days was used,
respectively). In both cases, PC1 had the biggest loading for bout no., with smaller loadings
for all other wheel parameters (Figure S2). Together this suggests, that PC1 is well suited to
represent overall wheel activity capturing maximal variance of individual running wheel
performance.

Statistical analysis

Normality of data was assessed using the Kolmogorov-Smirnov test, and non-parametric
versus parametric tests were chosen accordingly as stated in the text. Inter-group
comparisons between two groups were done by Student's t-test or Mann Whitney test, for 3
or more groups, we used one or two-way analysis of variance (ANOVA), as detailed. In
analyses in which a within-subject factor was present (i.e. eye), ANOVA with repeated
measurements was performed. Post hoc multiple comparison tests were corrected by
applying Sidak correction to p-values. The levels of significance were set as *: $p < 0.05$; **:
$p < 0.01$; ***: $p < 0.001$. Data are represented as means ± standard error of mean (s.e.m.).

246 Software accessibility

247 Any software and code used for running the gRW setup and for processing obtained data
248 are available upon request.

249 Results

250 The gated running wheel setup allows for detailed readout of
251 individual mouse running behaviour.

Given the importance of social interactions for the well-being of mice, we specifically
designed a gated running wheel setup (gRW), so that we could track individual running
activity of group-housed mice in a home cage setting. The gRW setup consists of a standard
rat cage connected to a separate compartment with a running wheel for voluntary physical
exercise (Figure 1A). Upon entering the gRW-compartment, mice are individually registered
using an implanted RFID sensor allowing to allocate running bouts to individual animals. This
enables us to quantify individual wheel running parameters and to correlate individual
running behaviours with individual V1-plasticity parameters assessed using intrinsic signal
optical imaging.

In order to provide a detailed analysis of wheel activity of mice, we first compared running
wheel activity between animals undergoing monocular deprivation (MD, n=12) and control
mice (noMD, n=8): wheel running was not affected by MD (Figure S2). Next, we evaluated
whether the separate gRW-compartment would impair access and reduce overall running
activity compared to a previous publication using identical running wheels located inside the
home cage. This did not seem to be the case: mice in the gRW setup ran 3307±763 RW
turns per day, corresponding to 1.3±0.3 km/d (Figure 1B), rather similar to the 3991±445 RW
turns/day, corresponding to 1.6±0.2 km/d of the animals of a previous study (Kalogeraki et
al., 2014).

Since, typically, only a single mouse entered the RW compartment at a time, the activity of
individual mice could be well separated by assigning running bouts to individual mice for
assessing interindividual variability in wheel running (Figure 1C-G). Across mice, running
distance increased over days as revealed by linear regression analysis (Figure 1D,
$F(1,138)=4.98$, p=0.027). However, running was not consistent across days for individual
mice (Figure 1D), and could range from close to 0 to above 10 km on individual days and for
individual mice (Figure 1D). Average cumulative running distances reached 8.2±1.8 km after
7 days, ranging from close to 0 to above 20 km of total distance travelled by individuals
(Figure 1D). Mice exhibited a significant increase in running distance from day 1 to day 7
(Figure 1E , one-sample t-test p=0.001), whereas the difference between day 4 and day 1
was not significant (Figure 1E). This was paralleled by a larger proportion of mice running
more than one bout per day, which increased from 50% on day 1 to 75% on day 7 (Figure
1F).

As expected from nocturnal animals, mice ran more during night time compared to day time,
with approximately 50% of mice showing a clear circadian pattern and running predominantly
during night time (Figure 1F), while the other half of the animals did not run sufficiently to
detect circadian activity preferences, and few individuals also occasionally ran during day
time.

On average, mice achieved running times of 1.5±0.3 h/d consisting of 17±3 bouts/d with
individual running bouts lasting on average 5.0±0.5 min, while running at a speed of 18±1
cm/s (Figure 1G). Of note, mice covered a wide range of running wheel parameters, running
0-3.96 km/d in 0-3.85 hours with running bouts lasting from <1 up to 10 min, while running at
a speed of 6-26 cm/s. Interestingly, mice running longer distances typically also spent more
time running, at faster speeds, which was also reflected in a larger number of running bouts,
with few exceptions, as illustrated by the colour coding of individuals in figures 1E,G. A
correlation matrix was computed to examine the interrelationships between the 5 quantified
running parameters more precisely (Figure H): The strongest correlation was observed
between distance and time (R=0.97, p<0.0001), i.e. mice that ran for longer time periods also
travelled a larger total distance. Speed and bout number also strongly correlated with running
distance (speed: R=0.81, p<0.0001; bout no: R=0.72, p<0.0001), i.e. mice running faster and
more often, also reached a higher final distance. In turn, bout duration did not correlate with
distance, time or number of bouts (p>0.05), however there was a significant correlation with
running speed (R=0.63, p<0.01), i.e. faster mice also ran longer bouts.

Together, these data demonstrate i) substantial interindividual variability in multiple running
wheel parameters of our group-housed mice which may influence cortical plasticity, and ii)
that our new gRW-setup is ideally suited to analyse this question in detail.

Gated wheel running restores OD-plasticity to adult standard cage raised mice

We had previously shown that running can boost OD-plasticity in SC-raised adult mice, even
if running was possible only during 7 days of MD (Kalogeraki et al., 2014). Here, we tested
whether i) the gRW setup also boosts OD-plasticity in SC-mice and ii) individual running
parameters, i.e. running more, would influence individual OD-plasticity. gRW-housing started
immediately after MD, and V1-activity maps were visualized after 7 days of MD. We
recorded intrinsic signal optical imaging responses of binocular V1 to visual stimulation of
the left and right eye with horizontal drifting bars (Cang et al., 2005).
Confirming previous imaging data in mice without MD (Lehmann and Löwel, 2008; Sato and
Stryker, 2008), visual stimulation of the contralateral eye induces a stronger V1-activation
compared to ipsilateral eye stimulation, visible as darker patches in the V1-activity maps
(Figure 2A), and reflected in a higher V1-activation strength (Figure 2C; contra/ipsi:
2.0±0.2/1.3±0.1, n=10, p=0.0003). In contrast, in gRW mice, MD induced a clear OD-shift:
After MD, the contralateral eye no longer activated V1 more strongly than the ipsilateral eye
(contra/ipsi: 1.7±0.2/1.5±0.0.2, n=9; p=0.7). Calculating the OD-index, quantifying OD-
plasticity by comparing ipsi- and contralateral eye induced V1-activity (Fig. 2B), confirms the
gRW boosted OD-shift: in noMD mice, the OD-index was 0.24±0.04 (n=10), indicating
contralateral eye dominance, whereas following MD, the OD-index was reduced to
0.05±0.04 (n=9, Mann-Whitney p-value 0.001), demonstrating a clear OD-shift in P160 mice.
Thus, our new gRW setup boosts OD-plasticity in group-housed mice.
327

Optomotor results

To test animals' basic visual abilities and to confirm the effectiveness of MD, we used the
virtual reality optomotor system, and measured the spatial frequency threshold (SFT) of the
optomotor reflex before and after MD (Prusky et al., 2004): vertical drifting gratings of

various spatial frequencies evoke small horizontal head and neck movements following the
stimulus. MD typically leads to an enhancement of optomotor reflex thresholds through the
open eye (citation!). Confirming effective MD, the SFT of our group-housed mice increased
from 0.39±0.00 cyc/deg on day 0 to 0.51±0.00 cyc/deg on day 7 after MD (n=10/9; 2-way
ANOVA for effect of MD p< 0.0001; Figure 2H), while values remained stable in noMD mice
(day 0/7: 0.39±0.00/0.39±0.00 cyc/deg, 2-way ANOVA for effect of MD: p<0.0001).

Gated running wheel parameters correlate with a measure of ocular dominance (OD) plasticity, the OD-index

Correlation of overall running wheel activity with OD-index: In order to investigate how
individual behavioural choices shape brain plasticity, we tested whether individual running
wheel parameters were correlated with individual OD-indices, which quantify the magnitude
of OD-plasticity. For this, we first obtained a measure of overall gRW activity for individual
mice by performing a principal component analysis (PCA) on the quantified running wheel
parameters (distance, time, speed, running bout numbers, bout duration) of pooled no MD
and MD mice, averaged across the 7 days of wheel exposure. The first principal component
explained 87% of interindividual variability in wheel running and hence was a suitable
measure of overall gRW activity of individual mice (see methods and figure S2 for details).
Strikingly, interindividual variability in gRW activity explained 65% of variability of the OD-
index (r^2=0.65, p=0.017, Figure 3A,B): Animals with high gRW performance showed a lower
OD-index and thus increased plasticity (Figure 3A,B). In contrast, correlation analyses in no
MD mice revealed no significant relationship between gRW activity and OD-index (r^2=0.02,
p=0.77, Figure 3B).
Correlation of individual running wheel parameters with OD-index: In addition, we also
observed striking correlations of individual running wheel parameters with individual OD-
index after MD (Figure 3C). In MD mice, bout duration, speed, bout number, running
distance and running time explained close to or above 60% (e.g. r^2 ≥ 0.60) of the
interindividual variability in OD-index of MD mice (n=9; bout duration: r^2=0.69, p=0.0102;
speed: r^2=0.63, p=0.0175; bout no: r^2=0.58, p=0.0271; running distance: r^2=0.60, p=0.0243;
running time r^2=0.62, p=0.0192). Importantly, no significant correlation between running
wheel parameters and OD-index was found in the no MD control group (p>0.05 for all
parameters), suggesting that baseline OD-index is not affected by wheel running.
Correlation of wheel activity on specific days with OD-index: Previous experimental evidence
has demonstrated that running wheel activity correlates with cortical gamma activity in V1
(Niell & Stryker, 2010), and that MD is associated with increased gamma activity in V1 for
several hours after MD, at least in juvenile mice aged 24-27 days (Quast et al., 2023).
Hence, we wondered whether running wheel activity on a specific day after MD was
correlated more strongly with the OD-index. For this we used PCA for dimensionality
reduction of running wheel parameters measured for *each day separately* and correlated the
first principal component (explaining 90% of variability of running wheel parameters across
days and animals) with the OD-index: gRW activity significantly correlated with the OD-index
on days 6 and 7 after MD (day6/7: r^2=0.52/0.53, p=0.045/0.041), while this relationship was
not significant on days 1-5 (p>0.05 for all correlations), suggesting that predominantly
running on days 6 and 7 affects the OD-index. Nevertheless, gRW activity on the first days
still contributed to individual variability in OD-index, since including gRW activity of
cumulative numbers of days in the analysis already produced significant correlations when

only data from days 1 and 2 was included (p>0.05 for day 1 alone, p<0.05 for data from >1
days).
Overall these data document that group housed mice display huge interindividual variability
in voluntary physical exercise when given the possibility to use a running wheel: individuals
differ in bout number, bout duration, running speed, running distance and running time. Most
notably, the amount of wheel running was predictive of the magnitude of OD-plasticity: Mice
that ran more had stronger experience-dependent changes in V1-activation after MD. This
directly highlights the importance of inter-individual behavioural variability for brain plasticity
and suggests that individual behavioural choices and their influence on brain physiology and
plasticity should be an integral part of future studies.

Discussion

Using our newly developed gated running wheel setup (gRW) that allows to track individual
running parameters of group-housed mice, we observed an enormous variability of individual
behaviours of the tested animals: running speed, running distance, total running time,
number of running bouts and bout duration varied by many orders of magnitude between
individuals. Furthermore, we revealed a striking correlation between individual running
parameters and experience-dependent plasticity in mouse V1. More running caused more
ocular dominance (OD) plasticity: all quantified running parameters significantly correlated
with a measure of visual cortical plasticity, the OD-index. Thus, our observations add to the
growing body of evidence that individual behavioural choices can strongly affect individual
brain plasticity, and should therefore be considered when analysing neuronal plasticity.

399
Mice housed in the gRW for one week showed huge interdividual differences in running
wheel performance, running 0-3.96 km/d in 0-3.85 hours with running bouts lasting between
0 and 10 min, while running at a speed of 6-26 cm/s, suggesting diverse intrinsic motivation
to run. Wheel running is distinct from home cage running because the distances and speeds
reached per day are drastically increased when mice have access to a running wheel. Mice
in their home cages only achieve speeds of up to 1 cm/s and distances of 0.1-0.2 km/d
(Iannello, 2019), while for running wheels allow speeds of up to 15-120 cm/s and distances
of 1-20 km/d have been reported (Koteja et al., 1999; Kopp, 2001; Manzanares et al., 2018).
While our data spans the lower end of the reported range - with individual mice running up to
10 km/d on single days – this is expected from the specific conditions used here: i) group
housing reduces running distances by ~50% (Plenz and Kanold, 2021) and ii) the short-term
wheel access provided here for only 7 days likely was not sufficient for mice to reach a
stable performance, which has been reported to take ~2 weeks (De Bono et al., 2006).

413
It is widely accepted that OD-plasticity in SC-raised mice is age-dependent, with clearly
decreasing plasticity in adult animals beyond P110, which requires an extended MD for
observable OD-shifts (Sawtell et al., 2003; Pham, 2004; Hofer et al., 2006; Lehmann and
Löwel, 2008; Sato and Stryker, 2008; Hosang et al., 2018). A number of environmental and
behavioural interventions have been established by now for restoring plasticity in adult SC-
raised rodents or for sustaining the plastic potential of adult rodent V1 beyond that age
(Espinosa and Stryker, 2012; Hübener and Bonhoeffer, 2014). This includes previous
episodes of MD (Hofer et al., 2006), forced visual stimulation (Matthies et al., 2013),
combined visual stimulation and running (Kaneko and Stryker, 2014), and dark rearing (He

et al., 2006; Stodieck et al., 2014), all of which promoted visual cortical plasticity in adult
animals. Notably, mice and rats with access to a running wheel or housing in enriched
environment retained OD-plasticity into late adulthood (Sale et al., 2007; Baroncelli et al.,
2010; Greifzu et al., 2014; Kalogeraki et al., 2014), demonstrating that voluntary physical
exercise can sustain the brain's capacity for adaptive modification to environmental changes.
428
While OD-shifts obtained in the present gRW-mice were similar to previously published data
after 7 days of RW-enrichment (Kalogeraki et al., 2014), OD-shifts were smaller compared to
animals experiencing lifelong enrichment in even larger 2-floor cages with a regularly
changed maze (Greifzu et al., 2014). Thus, while short-term running can restore OD-
plasticity in adult mice, more complex enrichment of their immediate environment for longer
periods of time, enables stronger experience-dependent V1-activity changes. Alternatively,
extension of MD-duration can also boost OD-plasticity in adult SC mice (Hosang et al.,
2018). Thus, there seems to be a trade-off between age and MD duration in SC-raised
animals: in younger mice, shorter MDs are sufficient to induce significant OD shifts. In
contrast, older SC-mice need considerably extended MD duration to display OD plasticity,
but these long MD times can be shortened by specific environmental and behavioural
interventions such as, e.g., raising animals in an enriched environment (Sale et al., 2007;
Greifzu et al., 2014) or providing access to a running wheel (Kalogeraki et al., 2014), like in
the present study.
443
Using C57Bl/6J inbred mice housed in the gRW-setup allowed us to test how individual
behavioural trajectories affect brain plasticity with minimal influence of genetics on the
observed phenotype. In line with the model of Kempermann (2019), we observed a linear
relationship between running parameters and OD-plasticity quantified by the OD-index, with
individual running performance predicting up to 65% of phenotypic variability in OD-index
between mice. Thus, individual behavioural choices exert a strong influence on experience-
dependent plasticity. Interestingly, all wheel parameters correlated similarly strong with
individual OD-index, with p-values of correlations only varying slightly between 0.010 and
0.027 between bout duration, bout number, running speed, running distance and running
time. As wheel running comprises a rewarding behaviour that even mice in the wild pursue
voluntarily (Sherwin, 1998), it remains unclear why some mice showed very little wheel
activity. Multiple factors affecting wheel running have been identified, including mouse strain,
sex, group housing, social hierarchy, age and circadian rhythms (Kopp, 2001; De Bono et
al., 2006; Basterfield et al., 2009; Bartling et al., 2017; Bains et al., 2018; Balog et al., 2019;
Plenz and Kanold, 2021). Importantly, social hierarchy has been shown to affect both wheel
running and OD-plasticity in male mice, with dominant males exhibiting stronger OD-shifts
compared to subordinate males (Balog et al., 2019). While social hierarchies are less
prominent in female mice (Williamson et al., 2019) used here, we cannot exclude that
subordinate social rank may have caused reduced access to the gRW in some of our mice,
and thus might have contributed to both the amount of wheel running and variability of OD-
index.
465
Both high running speed and bout duration require high physical fitness and might be
associated with better sensory-motor coordination for efficient running that may improve with
practice (De Bono et al., 2006), suggesting that pre-training or rearing under less deprived
conditions could also benefit OD-plasticity by enabling more efficient running and higher
fitness. Interestingly, as intermitted access to running wheels has been shown to specifically

increase adult neurogenesis in the hippocampus as opposed to continuous long term access
(Nguemeni et al., 2018), gRW housing might provide additional benefits as wheel access
was limited to one mouse at a time.
474

In addition, variability in the activity of specific neural networks has been linked to mouse
wheel activity (Rhodes et al., 2003): lateral hypothalamic orexin/hypocretin and GAD65
networks have emerged as drivers of locomotion initiation (Kosse et al., 2017; Karnani et al.,
2020), suggesting that interindividual differences in network activity might explain variability
in wheel running. In addition, other factors which were not captured in our experiment, might
also have contributed to variability in OD-plasticity. This includes exploratory behaviours not
captured by our setup, a higher intrinsic plastic ability of the brain enabling OD-plasticity
more independently of external factors, which might be related to overall higher fitness levels
as also suggested by younger mice running more than older mice (Bartling et al., 2017).
484

Together, our newly developed gRW-setup allowing to correlate individual mouse running
behaviour with individual measures of experience-dependent V1-plasticity has demonstrated
a striking correlation between mouse running activity and OD-plasticity, highlighting the
importance of individual behavioural tracking for explaining experimental data. Thus, our
observations add to the growing body of evidence that individual behavioural choices can
strongly affect individual brain plasticity and thus should be analysed more carefully in future
studies.
492

Author contributions: Study design by C.S. Data collection and initial processing by C.S.,
J.S and M.G. Formal analysis by C.S. Data interpretation by C.S. and S.L. Original
manuscript draft by C.S. Reviewing and editing of manuscript draft C.S. and S.L.

Conflict of Interest Statement:
The authors whose names are listed above certify that they have NO affiliations with or
involvement in any organization or entity with any financial interest (such as honoraria;
educational grants; participation in speakers' bureaus; membership, employment,
consultancies, stock ownership, or other equity interest; and expert testimony or patent-
licensing arrangements), or non-financial interest (such as personal or professional
relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed
in this manuscript.

Data Availability Statement:
The data supporting the findings of this research are available on request to the
corresponding author, pending a formal data-sharing agreement and approval from the local
ethics committee.

Acknowledgements: We thank DFG (SFB889, Project B05 to S.L.) and the Dorothea
Schlözer-Program for funding. We thank Tobias Mühmer and Jan Hoffmann, for constructing
the gated running wheel cage, and Christina Ahlbrecht for technical assistance and animal
care.

References

Bains RS, Wells S, Sillito RR, Armstrong JD, Cater HL, Banks G, Nolan PM (2018) Assessing mouse behaviour throughout the light/dark cycle using automated in-cage analysis tools. J Neurosci Methods 300:37–47.

Balog J, Hintz F, Isstas M, Teichert M, Winter C, Lehmann K (2019) Social hierarchy regulates ocular dominance plasticity in adult male mice. Brain Struct Funct 224:3183–3199.

Baroncelli L, Sale A, Viegi A, Maya Vetencourt JF, De Pasquale R, Baldini S, Maffei L (2010) Experience-dependent reactivation of ocular dominance plasticity in the adult visual cortex. Exp Neurol 226:100–109.

Bartling B, Al-Robaiy S, Lehnich H, Binder L, Hiebl B, Simm A (2017) Sex-related differences in the wheel-running activity of mice decline with increasing age. Exp Gerontol 87:139–147.

Basterfield L, Lumley LK, Mathers JC (2009) Wheel running in female C57BL/6J mice: impact of oestrus and dietary fat and effects on sleep and body mass. Int J Obes 33:212–218.

Bogado Lopes J, Senko AN, Bahnsen K, Geisler D, Kim E, Bernanos M, Cash D, Ehrlich S, Vernon AC, Kempermann G (2023) Individual behavioral trajectories shape whole-brain connectivity in mice Lerch JP, Behrens TE, Lerch JP, Yee Y, eds. eLife 12:e80379.

Cait J, Winder CB, Mason GJ (2024) How much "enrichment" is enough for laboratory rodents? A systematic review and meta-analysis re-assessing the impact of well-resourced cages on morbidity and mortality. Appl Anim Behav Sci 278:106361.

Cang J, Kalatsky VA, Löwel S, Stryker MP (2005) Optical imaging of the intrinsic signal as a measure of cortical plasticity in the mouse. Vis Neurosci 22:685–691.

De Bono JP, Adlam D, Paterson DJ, Channon KM (2006) Novel quantitative phenotypes of exercise training in mouse models. Am J Physiol-Regul Integr Comp Physiol 290:R926–R934.

Di Garbo A, Mainardi M, Chillemi S, Maffei L, Caleo M (2011) Environmental enrichment modulates cortico-cortical interactions in the mouse. PLoS ONE 6.

Espinosa JS, Stryker MP (2012) Development and Plasticity of the Primary Visual Cortex. Neuron 75:230–249.

Fabel K, Wolf S, Ehninger D, Babu H, Galicia P, Kempermann G (2009) Additive effects of physical exercise and environmental enrichment on adult hippocampal neurogenesis in mice. Front Neurosci 3.

Gordon JA, Stryker MP (1996) Experience-Dependent Plasticity of Binocular Responses in the Primary Visual Cortex of the Mouse. J Neurosci 16:3274–3286.

Greifzu F, Pielecka-Fortuna J, Kalogeraki E, Krempler K, Favaro PD, Schlüter OM, Löwel S (2014) Environmental enrichment extends ocular dominance plasticity into adulthood

and protects from stroke-induced impairments of plasticity. Proc Natl Acad Sci U S A
111:1150–1155.

He H-Y, Hodos W, Quinlan EM (2006) Visual Deprivation Reactivates Rapid Ocular
Dominance Plasticity in Adult Visual Cortex. J Neurosci 26:2951–2955.

Hofer SB, Mrsic-Flogel TD, Bonhoeffer T, Hübener M (2006) Lifelong learning: ocular
dominance plasticity in mouse visual cortex. Curr Opin Neurobiol 16:451–459.

Hosang L, Yusifov R, Löwel S (2018) Long-Term Visual Training Increases Visual Acuity and
Long-Term Monocular Deprivation Promotes Ocular Dominance Plasticity in Adult
Standard Cage-Raised Mice. eneuro 5:ENEURO.0289-17.2017.

Hübener M, Bonhoeffer T (2014) Neuronal Plasticity: Beyond the Critical Period. Cell
159:727–737.

Iannello F (2019) Non-intrusive high throughput automated data collection from the home
cage. Heliyon 5:e01454.

Kalogeraki E, Greifzu F, Haack F, Löwel S (2014) Voluntary physical exercise promotes
ocular dominance plasticity in adult mouse primary visual cortex. J Neurosci
34:15476–15481.

Kaneko M, Stryker MP (2014) Sensory experience during locomotion promotes recovery of
function in adult visual cortex Nelson SB, ed. eLife 3:e02798.

Karnani MM, Schöne C, Bracey EF, González JA, Viskaitis P, Li H-T, Adamantidis A,
Burdakov D (2020) Role of spontaneous and sensory orexin network dynamics in
rapid locomotion initiation. Prog Neurobiol:101771.

Kempermann G (2019) Environmental enrichment, new neurons and the neurobiology of
individuality. Nat Rev Neurosci 20:235–245.

Kopp C (2001) Locomotor activity rhythm in inbred strains of mice: implications for
behavioural studies. Behav Brain Res 125:93–96.

Körholz JC, Zocher S, Grzyb AN, Morisse B, Poetzsch A, Ehret F, Schmied C, Kempermann
G (2018) Selective increases in inter-individual variability in response to
environmental enrichment in female mice. eLife 7:e35690.

Kosse C, Schöne C, Bracey E, Burdakov D (2017) Orexin-driven GAD65 network of the
lateral hypothalamus sets physical activity in mice. Proc Natl Acad Sci 114:4525–
4530.

Koteja P, Swallow JG, Carter PA, Garland Jr Theodore (1999) Energy Cost of Wheel
Running in House Mice: Implications for Coadaptation of Locomotion and Energy
Budgets. Physiol Biochem Zool Ecol Evol Approaches 72:238–249.

Lehmann K, Löwel S (2008) Age-dependent ocular dominance plasticity in adult mice. PLoS
ONE 3.

Löwel S, Dehmel S, Makowiecki K, Kalogeraki E (2017) Environmental conditions strongly
affect brain plasticity. Neuroforum 24:A19–A29.

Manzanares G, Brito-da-Silva G, Gandra PG (2018) Voluntary wheel running: patterns and
physiological effects in mice. Braz J Med Biol Res 52:e7830.

Matthies U, Balog J, Lehmann K (2013) Temporally Coherent Visual Stimuli Boost Ocular
Dominance Plasticity. J Neurosci 33:11774–11778.

Mayr KA, Young L, Molina LA, Tran MA, Whelan PJ (2020) An economical solution to record
and control wheel-running for group-housed mice. J Neurosci Methods 331:108482.

Nguemeni C, McDonald MW, Jeffers MS, Livingston-Thomas J, Lagace D, Corbett D (2018)
Short- and Long-term Exposure to Low and High Dose Running Produce Differential
Effects on Hippocampal Neurogenesis. Neuroscience 369:202–211.

Pham TA (2004) A semi-persistent adult ocular dominance plasticity in visual cortex is
stabilized by activated CREB. Learn Mem 11:738–747.

Plenz UT, Kanold PO (2021) Differences in running performance of single- and group-
housed mice. :2021.12.29.474296.

Prusky GT, Alam NM, Beekman S, Douglas RM (2004) Rapid Quantification of Adult and
Developing Mouse Spatial Vision Using a Virtual Optomotor System. Investig
Opthalmology Vis Sci 45:4611.

Reuser A, Wenzel K, Felix SB, Dörr M, Bahls M, Könemann S (2022) Simultaneous
assessment of spontaneous cage activity and voluntary wheel running in group-
housed mice. Sci Rep 12:4444.

Rhodes JS, Garland T, Gammie SC (2003) Patterns of Brain Activity Associated With
Variation in Voluntary Wheel-Running Behavior. Behav Neurosci 117:1243–1256.

Sale A, Maya Vetencourt JF, Medini P, Cenni MC, Baroncelli L, De Pasquale R, Maffei L
(2007) Environmental enrichment in adulthood promotes amblyopia recovery through
a reduction of intracortical inhibition. Nat Neurosci 10:679–681.

Sato M, Stryker MP (2008) Distinctive features of adult ocular dominance plasticity. J
Neurosci Off J Soc Neurosci 28:10278–10286.

Sawtell NB, Frenkel MY, Philpot BD, Nakazawa K, Tonegawa S, Bear MF (2003) NMDA
Receptor-Dependent Ocular Dominance Plasticity in Adult Visual Cortex. Neuron
38:977–985.

Sherwin CM (1998) Voluntary wheel running: a review and novel interpretation. Anim Behav
56:11–27.

Stodieck SK, Greifzu F, Goetze B, Schmidt K-F, Löwel S (2014) Brief dark exposure
restored ocular dominance plasticity in aging mice and after a cortical stroke. Exp
Gerontol 60:1–11.

Stryker MP, Löwel S (2018) Amblyopia: New molecular/pharmacological and environmental
approaches. Vis Neurosci 35:E018.

Wahlsten D, Bachmanov A, Finn DA, Crabbe JC (2006) Stability of inbred mouse strain
differences in behavior and brain size between laboratories and across decades.
Proc Natl Acad Sci 103:16364–16369.

Williamson CM, Lee W, DeCasien AR, Lanham A, Romeo RD, Curley JP (2019) Social
hierarchy position in female mice is associated with plasma corticosterone levels and
hypothalamic gene expression. Sci Rep 9:7324.

Tables

Table 1: Hardware essentials for building a gated running wheel setup.

Item	Catalog No.	Company
Raspberry pi 4B+ 4GB, 32 GB SD	RP-4B-4GB	Raspberry Pi Foundation, Cambridge, England
Raspberry Pi Noir Camera 2.0 8MP	RB-CAMERAV2IR	Raspberry Pi Foundation, Cambridge, England
Raspberry Pi® RB-LCD-7 Display-Modul 17.8 cm (7 Zoll) 800 x 480 Pixel	RB-LCD-7	Raspberry Pi Foundation, Cambridge, England
RFID reader	ID-20LA	SparkFun Electronics, Colorado, USA
RFID implants	SEN-09416	Sparkfun Electronics, Colorado, USA
roller switch for seesaw	190.072.013	Marquardt, Germany
phototransistor for beam break sensors	BPV11F	Vishay, Germany
reed contact for Hall sensor	MS-213-3	PIC Proximity Instrument Controls GmbH, Germany

Figure legends

Figure 1. Gated running wheel setup allows to track individual mouse running
wheel activity.
A. Standard-sized rat cage (left, 43 cm x 27 cm) with an additional compartment (right)
containing beam break sensors (1,3), a seesaw (2), RFID sensor (4), hall sensor (5),
magnet (6), and a running wheel (7). All the sensors are connected to a raspberry pi
4B+ (8). (9) Illustrates the size of the implanted RFID.
B. Comparison of average daily running activity of gRW housed mice (orange) with
published data from Kalogeraki et al. 2014 (black).
C. Example of quantified gRW-activity of two co-housed mice illustrating allocation of
running bouts to individual mice (orange/black line) for 2 consecutive days and nights
(grey background) (top) and 60 minutes during night time (bottom), with detected
running bouts highlighted in grey.
D. Daily (left) and cumulative (right) running trajectories of individual mice housed in
gRW enrichment for 7 days, document a large range of individual running behaviour.

While some mice achieve high distances from the first day, for most mice daily running
distances only achieved high distances torwards the end of the 7 days. Colours of
traces and dots are matched for individual mice across figures E-G
E. Overall there was a significant increase in daily running between day 1 and day 7,
but not day 1 and 4.
F. Running distances of all animals across 7 days binned per hour with top: heat map
of indivdual mice and bottom: average distances across all animals, with overlaid
proportion of mice running on each day (defined as at least 1 bin above 0 distance
travelled). Note that animals with sufficient running activity predominantly run during
night time (higher average values and darker patches in heatmap), with few
expceptions of running also during the light period.
G. Running wheel parameters in individual mice. Note the huge interindividual
variability across parameters. Mice that ran longer distances also spent more time
running at higher speeds with larger bout numbers.
H. Correlation matrix illustrating interdependencies of 5 running wheel parameters
displayed in G.
671
Figure 2. gRW enrichment boosts ocular dominance plasticity in group-housed
mice.
A. Intrinsic signal optical imaging examples of grey-scale coded activity maps after
visual stimulation of the contralateral and ipsilateral eye in binocular V1 of C57Bl/6J
gRW housed mice after 7 days of monocular deprivation (MD) and without MD (no
MD). The magnitude of V1-activation is expressed as the fractional change in
reflectance $x10^{-4}$ and indicated in the top right. Warm colors in the color-coded OD-
maps indicate contralateral eye dominance, colder colors indicate an OD-shift towards
to ipsilateral eye. Histograms in A,E represent distribution of pixel wise OD-scores.
Scale bar: 1mm.
B. Quantification of OD-index. Symbols represent values of individual mice, means
are marked by horizontal lines.
C. V1-activation elicited by stimulation of the contralateral (C) or ipsilateral (I) eye
(filled circle indicates MD eye). Note that MD did not reduce V1-activation through the
deprived eye in C.
D. Spatial frequency threshold of the optomotor reflex for noMD and MD mice, plotted
vs. days (x-axis). Note that thresholds increase after MD, indicating successfully
induced MD (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ and ****$p < 0.0001$ in Mann-Whitney
or 2-way ANOVA with Sidak's post hoc test).
691
Figure 3. Individual running wheel (RW) activity correlates with ocular
dominance (OD) plasticity, quantified via the OD-index after monocular
deprivation (MD) in P160 mice.
A. Example intrinsic signal optical imaging recordings from noMD and MD P160
animals, with low (gRW activity <0) and high (gRW activity >0) RW performance (gRW
activity was defined as the first principal component (PC1) of RW parameters
displayed in C, and shown in B). Data displayed as in figure 2A. Note that OD-
plasticity after MD is only boosted in P160 mice, when mice run sufficiently in the gRW
setup.
B. The OD-index of individual P160 mice is correlated with their individual gRW
activity after MD (orange), but not in no MD control mice (grey). Lines represent linear
regression fitted to MD mice and no MD mice, respectively. Pearson r-values and p-
values for MD mice are in the figure.

C. The OD-index of individual mice after MD is correlated with all quantified running
wheel parameters. Data is displayed as in B. Pearson correlation between OD-index
and individual wheel running activity was significant for all tested running wheel
parameters in P160 mice after MD, but not no MD control mice.
709
Video 1. Video footage of a mouse entering the gated running wheel compartment via
flipping a seesaw.
Video 2. Video footage of a mouse leaving the gated running wheel compartment via
flipping a seesaw. Note that its cage mate is blocked from entering the compartment.
Model 1. 3D model of gated running wheel components
Figure S1. Electronic circuit of gated running wheel setup
Figure S2. llustration of PCA analysis of wheel running parameters averaged across 7
days (A-C) and from individual days (D-F). A,D: Data transformed into coordinate
system of principal component (PC) 1 and PC 2 and graphical representation of
loadings. B,E: Loadings of all 5 PC C,F: Variability explained by each PC.
Figure S3. Comparison of wheel running parameters between no MD and MD mice.
No significant difference was observed.

Figure 1 Gated running wheel setup allows for detailed readout of wheel running behavior in group housed mice

Figure 2 Gated running wheel housing promotes ODP in group housed mice

Figure 3 Running wheel activity correlates with ODI