1   **Title:** More running causes more ocular dominance plasticity in mouse primary visual cortex:
2   new gated running wheel setup allows to quantify individual running behaviour of group-
3   housed mice.
4
5   **Abbreviated title:** More running, more ocular dominance plasticity
6

7   **Authors:** Cornelia Schöne[1,2], Jaya Sowkyadha Sathiyamani[1,2,3], Mihaela Guranda[1], Nele
8   Schnurbusch[1], Siegrid Löwel[1,2]
9
10 **Affiliation:**
11 [1] University of Göttingen, Department of Systems Neuroscience, 37075 Göttingen, Germany
12 [2] Göttingen Campus Institute for Dynamics of Biological Networks, 37075 Göttingen,
13 Germany
14 [3] International Max Planck Research School for Neurosciences, 37075 Göttingen, Germany
15
16 Corresponding authors email: cornelia.schoene@gwdg.de

17
Number of pages: 19
Number of figures: 5
Number of multimedia: 2
Number of 3D models: 1
Number of words:
Abstract: 250
Introduction: 649
Discussion: 1500
26
27
28
Conflict of Interest Statement:
The authors whose names are listed above certify that they have NO affiliations with or
involvement in any organization or entity with any financial interest (such as honoraria;
educational grants; participation in speakers' bureaus; membership, employment,
consultancies, stock ownership, or other equity interest; and expert testimony or patent-
licensing arrangements), or non-financial interest (such as personal or professional
relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in
this manuscript.
37
38

39 **Acknowledgements:** We thank DFG (SFB889, Project B05 to S.L.) and the Dorothea
40 Schlözer-Program for funding. We thank Tobias Mühmer and Jan Hoffmann, for constructing
41 the gated running wheel cage, Dr Subhodeep Bhattacharya for assisting with analysis and
42 Christina Ahlbrecht for technical assistance and animal care.

Abstract

Environmental enrichment boosts neuronal plasticity of standard-cage raised (SC) mice. We previously showed that wheel running during 7 days of monocular deprivation (MD) can restore ocular dominance (OD)-plasticity in the primary visual cortex (V1) of adult SC-mice beyond postnatal day (P) 110 (P160). It remained – however – unknown whether individual wheel running parameters boost OD-plasticity differently and whether running for just 4 days would also boost OD-plasticity in younger mice aged 50-100 days (P70).

To this end, we designed a gated running wheel setup (gRW) attached to the home cage, and tracked individual running wheel activity of group-housed mice. Mouse vision was checked by optometry and OD-plasticity in V1 was visualized using intrinsic signal optical imaging. As hypothesized, wheel running during MD boosted OD-plasticity in both P160 and P70 mice: after 4d/7d of MD (P70/P160), the OD-index, comparing ipsi- and contralateral eye induced V1-activation, dropped from 0.25±0.04 without MD to 0.12±0.03 with MD in P70 (t-test p<0.01), and from 0.24±0.04 to 0.05±0.04 in P160 mice (t-test p<0.01). Quantitative comparison of running parameters yielded that average running distance, running time and speed were similar in P70 and P160 mice, but P70 mice ran more bouts (P70/P160: 19.5±3.3/17.0±3.3 bouts/d, t-test p<0.001) of shorter durations (P70/P160: 2.9±0.5/5.0±0.5 min, t-test p<0.05). Most notably, individual running parameters correlated with individual OD-indices after MD: mice running longer distances, for longer time, at higher velocity and with longer bouts displayed a stronger OD-shift, i.e. animals showed more experience-dependent V1-plasticity (Pearson r²=0.42, p=0.022).

Significance statement

The link between individual choices and brain physiology has become a rising topic in systems neuroscience. With our newly developed gated running wheel setup, we observed a striking correlation between individual running activity, and experience-dependent plasticity in mouse primary visual cortex. More running caused more plasticity: *running speed, running distance, total running time, number of running bouts and bout duration all significantly correlated with a measure of visual cortical plasticity, the ocular dominance index.* Thus, our observations add to the growing body of evidence that individual behavioural choices can strongly affect individual brain plasticity.

Introduction

The link between individual choices and brain physiology has become a rising topic in systems neuroscience, and reflects the growing importance to study individuals, rather than populations by giving meaning to variability in biological data. While the natural environment of humans and mice is complex, that of laboratory rodents has been simplified and standardised in the past decades to an extent that deprives the animals of essential behaviours (Löwel et al., 2017). Furthermore, standard cage rearing (SC) leaves minimal room for the expression of individual traits and both behavioural and physiological inter-individual variability is further reduced by using inbred strains lacking genetic variability. Recently, a number of laboratories has started to systematically investigate the impact of less deprived rearing conditions on brain wiring and plasticity, demonstrating that raising animals in a so-called "enriched environment" elicits remarkable effects on the brain across molecular, anatomical, and functional levels when compared to animals raised in an SC environment (Sale et al., 2007; Fabel et al., 2009; Baroncelli et al., 2010; Di Garbo et al., 2011; Greifzu et al., 2014; Kalogeraki et al., 2014; Löwel et al., 2017; Stryker and Löwel, 2018; Bogado Lopes et al., 2023).

Both environmental enrichment and voluntary wheel running boost ocular dominance (OD) plasticity in mouse primary visual cortex (V1) (Greifzu et al., 2014; Kalogeraki et al., 2014). OD-plasticity is typically induced after monocular deprivation (MD), which shifts V1-activation strength towards the open eye (Cang et al., 2005; Gordon and Stryker, 1996). During the critical period of visual system development of SC mice, a 4-day MD is sufficient to induce an OD-shift in V1 intrinsic signal optical imaging responses to drifting bars, which is quantified using an OD-index (Cang et al., 2005). The required MD duration to induce OD-plasticity in SC-animals increases with age: While SC-mice between postnatal day 50 to 90 (P70) require 7 days of MD for significant changes in OD-index (Lehmann and Löwel, 2008; Sato and Stryker, 2008), this reaches ~7 weeks in mice older than postnatal day 110 (Hosang et al., 2018). Importantly, providing a running wheel to P160 SC-mice was sufficient to re-enable OD-plasticity after just one week of MD (Kalogeraki et al., 2014). While it is clearly established that running increases OD-plasticity in P160 mice (Kalogeraki et al., 2014; Kaneko and Stryker, 2014) the consequences of behavioural inter-individual variability of wheel running on experience-dependent V1-plasticity has not yet been studied. Moreover, it is unknown whether younger animals would also benefit from short-term running during MD.

The aim of this study was therefore to analyse the effects of individual running wheel activity on OD-plasticity in group housed P70 and P160 mice. As tracking individual wheel running activity of group housed mice is still challenging (Mayr et al., 2020; Reuser et al., 2022), we designed a custom-built gated running wheel (gRW) setup that allowed individual tracking of mice implanted with an RFID sensor while using a running wheel accessible from their home cage. We quantified various parameters of running activity and tested whether they correlated with the OD-index of individual P70 and P160 mice. The OD-index compares V1-activation via the ipsi- and contralateral eye and thus quantifies the magnitude of OD-shift (Cang et al., 2005). As hypothesized, gRW housing boosted OD-plasticity in both P70 and P160 animals. Average running distance in the gRW setup was similar to published values of group housed mice with running wheels provided directly in the home-cage (Kalogeraki et al., 2014; Plenz and Kanold, 2021). Importantly, our newly designed experimental gRW setup allowed us to correlate the macro (running distance, time and speed) and micro (bout number and duration) architecture of running wheel activity of individual mice with their individual OD-index: our data

123 clearly show a strong correlation between running wheel parameters of individual mice and
124 their OD-index quantifying OD-plasticity after MD that was absent in no MD control animals:
125 mice who ran more and longer distances displayed stronger OD-shifts.

126 Methods:

127 Animals and rearing conditions

Female C57Bl6/J mice were group-housed on a 12 h light dark cycle and lived in standard
cages (SC) until the day an MD was performed. Mice were provided with ad libitum food and
water. Two age groups were used for this study: P70 mice (age range P46-86) and P160 mice
(age range P127-225). All experimental procedures complied with the National Institutes of
Health guidelines for the use of laboratory animals and were approved by the local government
of lower saxony, Germany (Niedersächsisches Landesamt für Verbraucherschutz und
Lebensmittelsicherheit).

135 Monocular deprivation and implantation of RFID tags

To ensure that experimental mice had regular vision before induction of OD-plasticity, we
tested the spatial frequency threshold (SFT) of the optomotor reflex of both eyes before MD
(Prusky et al. 2004; Lehmann and Löwel, 2008). SFT values were within published values and
differed by less than 0.02 cyc/deg between the two eyes in all mice. The right eye was deprived
for 4 days (d) (P70) or 7 d (P160) according to published protocols (Gordon and Stryker, 1996;
Cang et al, 2005; Greifzu et al., 2014). The eyes of the animals were examined daily to ensure
that eyes remained healthy and the MD-eye stayed closed. Optomotor reflexes of the open
eye were tested daily after MD including the day of intrinsic signal optical imaging.
Cylindrical glass-covered radio frequency identification devices (RFID; length: 12.5 mm;
diameter: 1.93 mm, Sparkfun Electronics, Colorado, USA) were implanted during the MD or
noMD (control without MD) surgery to minimize mouse discomfort. First, a patch of fur was
shaved in the nape of the animal's neck using small surgical scissors, then a local anaesthetic
gel (Lidocaine, xylocaine gel 2 %, Aspen Pharma Trading Limited, Ireland) was applied to the
exposed skin. A small incision was made and the chip carefully placed subcutaneously in the
scruff to make it difficult for the mice to pick at and thereby reducing the risk of an infection.
The incision was sealed with a simple interrupted stitch (suture material: 7-0 Perma-Hand silk,
Ethicon, 8.0mm diameter). After MD/RFID implantation and complete recovery from
anaesthesia, mice were moved to a gRW setup.

154 Optical imaging of intrinsic signals and visual stimuli.

Surgery. Surgical preparations for optical imaging were performed as published (Cang et al.,
2005; Greifzu et al., 2014). Briefly, mice were box-anaesthetised with 2% halothane in O_2 and
N_2O (1:1) and injected with atropine (5mg/kg, s.c.; Franz Köhler Chemie), dexamethasone
(8mg/kg s.c.; Merck), and chlorprothixene (8mg/kg, i.m.; Sigma-Aldrich). After placing animals
in a stereotaxic frame, anaesthesia was maintained with 0.8% halothane in a 1:1 mixture of
O_2 and N_2O.
161

Data acquisition and visual stimulation (Kalatsky and Stryker, 2003; Cang et al, 2005). Mouse
V1 responses were recorded through the skull using the "Fourier" imaging method of Kalatsky
and Stryker (2003) and optimised for the assessment of OD-plasticity (Cang et al., 2005). V1-
signals were visualised with a CCD camera (Dalsa 1M30) using a 135x50 mm tandem lens
configuration (Nikon) with red illumination light (610±10 nm). Active brain regions absorb more
red light and appear darker in the images. Frames were acquired at a rate of 30 Hz, temporally
binned to 7.5 Hz, and stored as 512x512-pixel images after spatial binning of the camera
image. Visual stimuli were presented on a 60 Hz refresh rate monitor (21 inches; Accuvue
HM-4921-D, Hitachi) positioned 25 cm from the eyes. Stimuli consisted of white drifting
horizontal bars (2° wide) as described previously (Greifzu et al., 2014). The amplitude
component of the optical signal represents the intensity of neuronal activation (expressed as
fractional change in reflectance times 10^{-4}) and was used to calculate an ocular dominance
(OD) index. At least three maps per animal were averaged to compute the OD-index as
(C+I)/(C-I), with C and I representing the response magnitudes of each pixel to visual
stimulation of the contralateral (C) and ipsilateral (I) eye.

Gated running wheel setup (gRW)

In order to assess individual running wheel (RW) behaviour, we developed an open source
gated running wheel cage, which served as the home cage for the duration of the experiment.
It consisted of a standard size rat cage (43cm x 27cm) connected to a separate gRW-
compartment, equipped with the same RW as used in the enrichment cages in Greifzu et al.,
2014 (Figure 1A). Wheel diameter is 12.5 cm, so that the calculated RW-circumference was
39.3 cm. RW-turns were registered using a hall sensor for detecting motion of two magnets
attached at opposing sites of the RW. The mice needed to pass through a seesaw in order to
reach the RW-compartment. Entering would flip the seesaw blocking other mice from entering
(see video 1 and 2). To ensure the seesaw remained closed while a mouse was inside the
RW-compartment, the seesaw was actively blocked from flipping back. When activity was
detected by beam break sensors outside the RW-compartment, an electromagnet was
activated to hold the seesaw closed. Attempts of mice to exit the RW-compartment were
detected using beam break sensors inside the RW-compartment, which would deactivate the
seesaw magnet. Seesaw position was registered through a roller switch.
Mice were chipped with radio frequency identification devices (RFID). The RFID-tags
transmitted the encrypted individual ID to a USB-driven reading device (ID-20LA SparkFun
Electronics, Colorado, USA) at 125 kHz via a custom-made RFID coil. Thus, each time a
mouse passed underneath the RFID coil into the gRW compartment, RFID and time stamps
were saved. All sensors were connected to the GPIO pins of a raspberry pi 4B+. In addition,
a raspberry pi noir camera was used to record movement inside the RW-compartment at 40
Hz; camera data were used to confirm sensor data and to extract RW-turns when the Hall
sensor failed. All data was recorded using custom written python scripts running on the
raspberry pi. A 3D .pdf file illustrating dimensions of cage components and electronic
assembly is provided upon request.
202
203 Hardware essentials

Item	Catalog No.	Company
Raspberry pi 4B+ 4GB, 32	RP-4B-4GB	Raspberry Pi Foundation,

GB SD		Cambridge, England
Raspberry Pi Noir Camera 2.0 8MP	RB-CAMERAV2IR	Raspberry Pi Foundation, Cambridge, England
Raspberry Pi® RB-LCD-7 Display-Modul 17.8 cm (7 Zoll) 800 x 480 Pixel	RB-LCD-7	Raspberry Pi Foundation, Cambridge, England
RFID reader	ID-20LA	SparkFun Electronics, Colorado, USA
RFID implants	SEN-09416	Sparkfun Electronics, Colorado, USA
roller switch for seesaw	190.072.013	Marquardt, Germany
phototransistor for beam break sensors	BPV11F	Vishay, Germany
reed contact for Hall sensor	MS-213-3	PIC Proximity Instrument Controls GmbH, Germany

Analysis of gated wheel running

In order to assign each RW-turn to individual mice, we first defined running bouts as running at a speed above 0.1 Hz, then the most recently detected RFID was assigned to all wheel turns within the respective running bout. When more than one RFID was detected since the last gate flip, this indicated that more than one mouse had entered the gRW compartment. This occasionally happened, due to two mice squeezing through the gate simultaneously, or due to failure of the magnetic seesaw lock. In this case, the RW-turns were labelled as ambiguous. Mice with ambiguous RW-turns of more than 9% were excluded from the RW-analysis, but included for analysis of OD-plasticity in Fig. 2. For the remaining mice, on average, we quantified 2.7±0.8% ambiguous RW-turns for P70 mice (n=10 mice) and 1.7±0.5% ambiguous RW-turns for P160 mice (n=20 mice).

EXPERIMENTAL DESIGN AND STATISTICAL ANALYSES

Statistical analysis

Inter-group comparisons between two groups were done by Student's t-test, for 3 or more groups, we used one or two-way analysis of variance (ANOVA), as detailed. In analyses in which a within-subject factor was present (i.e. eye), ANOVA with repeated measurements was performed. Post hoc multiple comparison tests were corrected by applying Sidak correction to p-values. The levels of significance were set as *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$. Data are represented as means ± standard error of mean (s.e.m.).

225 Software accessibility

226 Any software and code used for running the gRW setup and for processing obtained data
227 are available upon request.

228 Results

229 The gated running wheel setup allows for detailed readout of
230 individual mouse running behaviour.

Given the importance of social interactions for the well-being of mice, we specifically designed
a gated running wheel setup (gRW) so that we could track individual running activity of group-
housed mice in a home cage setting. The gRW setup consists of standard rat cage connected
to a separate compartment with a running wheel for voluntary physical exercise (Figure 1A).
Upon entering the gRW-compartment, mice are individually registered using an implanted
RFID sensor allowing to allocate running bouts to individual mice (Figure 1C) and to correlate
individual running behaviors with individual V1-plasticity parameters assessed using intrinsic
signal optical imaging. In order to avoid conflict between group housed males, we limited our
experiments to groups of two to six female mice belonging to two age groups, termed P70 and
P160, whose gRW activity was tracked for 4 or 7 days respectively (Figure 1) and occurred
predominantly during night time (Figure 1C). As typically only single mice entered the RW
compartment the activity of individual mice could be well separated highlighting interindividual
variability for example in running bout length and speeds achieved by mice (Figure 1C,E).
P160 gRW mice ran on average 3307±763 RW turns per day, corresponding to 1.3±0.3 km/d
(Figure 1B). These values are similar to previously published data showing 3991±445 RW
turns/day, corresponding to 1.6±0.2 km/d for seven days of RW enrichment in group housed
P160 C57Bl/6J mice using identical running wheels (Kalogeraki et al., 2014). The total average
distance individual mice ran per day was highly variable and ranged from 0 to 3.7 km/d in P70,
and from 0 to 3.5 km/d in P160 mice. Cumulative running distances consistently increased
across days (Figure 1D), reaching 6.7±1.2 km after 4 days in P70 animals and 8.2±1.8 km
after 7 days in P160 animals. We could not identify significant differences in the trajectories of
cumulative wheel running between the age groups: linear regression lines fitted to cumulative
running distances of each age group revealed neither difference in slopes (F=1.21, p=0.272)
nor in intercepts (F=3.54, p=0.062) of the fitted curves.

We next compared several wheel running parameters between between P70 and P160 mice
(Figure 1E). Mice of both age groups reached similar average daily running distances of
1.7±0.3 km/d and 1.3±0.3 km/d (p=0.31), corresponding to daily running times of 1.8±0.3 h/d
and 1.5±0.3 h/d for P70 and P160 age groups, respectively (p=0.37). The average running
speed of P70 and P160 was 18±2 cm/s and 18±1 cm/s, again not significantly different
(p=0.84). However, P160 mice showed longer average running bouts of 5.0±0.5 min,
compared to the 2.9±0.5 min of P70 mice (p=0.009). P70 mice also ran 19.5±3.3 bouts/d, and
thus more than P160 mice, which only ran 17.0±3.3 bouts/d (p=0.01). Together, these data
demonstrate that voluntary physical exercise in group-housed mice shows huge interindividual
differences in various parameters, that may influence cortical plasticity, and that our new gRW-
setup is ideally suited to quantify these parameters of individual running activity of animals at
different ages.

gRW re-enables OD-plasticity in P70 and P160 mice

It was previously shown that 4 days of MD in P70 mice and 7 days of MD in P160 is not sufficient to induce OD-plasticity in standard cage (SC) raised rodents measured using intrinsic signal optical imaging (Lehmann and Löwel, 2008; Sato and Stryker, 2008). Here, we tested whether running in the gRW setup might also boost OD-plasticity in adult mice as previously shown for standard running wheels after 7 days of MD (Kalogeraki et al., 2014). gRW-housing started immediately after MD, and V1-activity maps were visualized after 4 (P70) or 7 days MD (P160), respectively. We recorded intrinsic signal optical imaging responses of binocular V1 to horizontal drifting bars (Cang et al., 2005). Confirming previous imaging data in mice without MD (Lehmann and Löwel, 2008; Sato and Stryker, 2008), visual stimulation of the contralateral eye induces a stronger V1-activation compared to ipsilateral eye stimulation, visible as the darker patches in the V1-activity maps (Figure 2A,E), and reflected in the V1-activation strength (Figure 2C,G): V1 was dominated by input from the contralateral eye (P70 contra/ipsi: $2.4\pm0.2/1.5\pm0.0.2$, n=9, p<0.0001; P160: $2.0\pm0.2/1.3\pm0.0.1$, n=10, p=0.0003). In contrast, in gRW mice of both age groups, MD induced a clear OD-shift: after MD, both eyes activated V1 more equally strong, i.e. V1-activity patches induced by ipsi- and contralateral eye stimulation show more similar grey values. Quantification of V1-activity (Fig. 2C,G) and the OD-index, comparing ipsi- and contralateral eye induced V1-activation (Fig. 2B,F) confirms the gRW boosted OD-shifts: while in noMD mice, the OD-index was positive, indicating contralateral eye dominance, after MD, the OD-index was lower compared to the noMD control groups (P70 noMD/MD: $0.25\pm0.04/0.12\pm0.03$, n=9/11, t-test p-value 0.007; P160 noMD/MD: $0.24\pm0.04/0.05\pm0.04$, n=10/9, t-test p-value 0.003). Comparing V1-activation values thus shows that in P70 animals, contralateral eye dominance was diminished after MD (P70 contra/ipsi: $2.6\pm0.03/2.2\pm0.3$, n=11, p=0.016). In P160 animals after MD, the contralateral eye no longer activated V1 more strongly than the ipsilateral eye (MD P160 contra/ipsi: $1.7\pm0.2/1.5\pm0.0.2$, n=9; p=0.7). Furthermore, V1-activation values in P70 mice indicate that the MD-induced OD-shifts do not result from a reduction of V1-activation through the previously deprived eye, like observed in juvenile OD-plasticity (Sato and Stryker, 2008), but rather from an increase in V1-activation through the open eye (Fig. 2C), as previously shown for OD-plasticity in young enriched mice (Kalogeraki et al., 2017).

Together our data highlight that even short-term gRW enrichment is able to boost OD-plasticity in group-housed mice of both age groups.

Optomotor results in gRW mice

To test animals' basic visual abilities and to confirm the effectiveness of MD, we used the virtual reality optomotor system to measure the spatial frequency threshold (SFT) of the optomotor reflex before and after MD (Prusky et al., 2004): vertical drifting gratings of various spatial frequencies evoke small horizontal head movements following the stimulus. MD typically leads to an enhancement of optomotor reflex thresholds through the open eye. As expected, and confirming effective MD, in P70 mice, 4 days of MD led to an increase of the SFT from 0.38 ± 0.00 to 0.43 ± 0.01 cyc/deg, while SFT remained stable in noMD mice (P70 day 0/4: $0.38\pm0.00/0.39\pm0.00$ cyc/deg; n(noMD/MD)=9/10; 2-way ANOVA for effect of MD: p=0.002; Figure 2D). Similarly, in P160 mice, 7 days of MD led to a significant increase of the SFT from 0.39 ± 0.00 (P160, day 0) to 0.51 ± 0.00 cyc/deg (n=10/9; 2-way ANOVA for effect of MD p< 0.0001; Figure 2H), while values remained stable in noMD mice (P 160, day 0/7): $0.39\pm0.00/0.39\pm0.00$ cyc/deg).

gRW running wheel parameters correlate with ocular dominance index

In order to investigate how individual behavioural choices feedback to shape brain plasticity, we tested whether individual running wheel parameters were correlated with individual OD-indices which quantify the magnitude of OD-plasticity in both age groups. For this we performed a principal component analysis (PCA) to reduce the dimensions of *average* running wheel parameters (daily running distance, daily running time, average daily running speed, average daily bout number, average daily bout duration) to the first principal component (explaining 96 % of interindividual variability in wheel running of pooled P70 and P160 data) as a measure of overall gRW activity. Mice showing less than median gRW activity (median gRW activity = 2.3) had weaker or absent OD-plasticity than mice showing higher than median gRW activity: pooling data from P70/P160 (Figure 3B), the OD-index of mice with high gRW activity was -0.03±0.02 compared to 0.12±0.04 for low gRW activity (n=6/6, p=0.007).

Confirming this result, inter-individual variability in gRW activity of P160 mice explained 67% in variability of the OD-index (r^2=0.67, p=0.013). Correlation analyses in P70 mice closely followed the same distribution as data from P160 mice such that the pooled correlations were also significant (r^2=0.48, p=0.012, Figure 3B). Since we could obtain only 4 wheel-parameter/OD-index pairs for P70 mice we refrained from testing correlations in this dataset separately.

In addition, we also observed striking correlations of individual running wheel parameters with individual OD-index (Figure 3C). In P160 mice, bout duration, speed, bout no, running distance and running time explained close to or above 60% (e.g. $r^2 \geq 0.60$) of the interindividual variability in OD-index (P160, n=9; bout duration: r^2=0.69, p=0.0102; speed: r^2=0.63, p=0.0175; bout no: r^2=0.58, p=0.0271; running distance: r^2=0.60, p=0.0243; running time r^2=0.62, p=0.0192). Correlation analyses in P70 mice again closely followed the same distribution such that the pooled correlations were significant for all quantified parameters except bout number (bout duration: r^2=0.61, p=0.0025; speed: r^2=0.45, p=0.0163; bout no: r^2=0.20, p=0.14; running distance: r^2=0.42, p=0.0219; running time r^2=0.45, p=0.0150). Importantly, no significant correlation between running wheel parameters and OD-index was found in the no MD control group (p>0.05 for all correlations).

Previous experimental evidence has demonstrated that running wheel activity correlates with cortical gamma activity in V1 (Niell & Stryker, 2010), and that MD is associated with increased gamma activity in V1 for several hours after MD, at least in juvenile mice aged 24-27 days (Quast et al., 2023). Hence, we wondered whether running wheel activity was correlated more strongly with the OD-index on a specific day after MD. For this we used PCA for dimensionality reduction of running wheel parameters measured for *each day separately* and correlated the first principal component (explaining 90 % of variability of running wheel parameters across days and animals) with the OD-index: gRW activity significantly correlated with the OD-index only on days 6 and 7 after MD (day6/7: r^2=0.52/0.53, p=0.045/0.041), while this relationship was not significant on days 1-5 (p>0.05 for all correlations). Nevertheless, gRW activity on the first day still contributed to variability in OD-index, since averaging gRW activity of increasing numbers of days, already produced significant correlations when only data from days 1 and 2 was included (p>0.05 for day 1 alone, p<0.05 for data from >1 days).

Overall these data document that even short-term voluntary physical exercise strongly boosts experience-dependent changes in V1-activation and highlight that more running leads to stronger OD-plasticity in mouse V1. Since interindividual variability in running behaviour is

359   very high, individual behavioural choices and their influence on brain physiology and plasticity
360   should be more strongly considered in future studies.

Discussion

Using our newly developed gated running wheel setup (gRW) that allows to track individual
running activity in group-housed mice, we observed a striking correlation between individual
running parameters and experience-dependent plasticity in mouse primary visual cortex. More
running caused more plasticity: *running speed, running distance, total running time, number*
of running bouts and bout duration all significantly correlated with a measure of visual cortical
plasticity, the ocular dominance index. Thus, our observations add to the growing body of
evidence that individual behavioural choices can strongly affect individual brain plasticity.
Confirming previous studies with conventional running wheels (Kalogeraki et al. 2014), 7 days
of gRW-running was sufficient to boost OD-plasticity in P160 mice. Likewise, running for only
4 days also restored OD-plasticity in P70 mice. While OD-shifts obtained in P160 were similar
to previously published data after 7 days of RW-enrichment, OD-shifts were smaller compared
to animals experiencing lifelong enrichment (Kalogeraki et al. 2017). Similarly, the magnitude
of OD-shifts for P70 mice was smaller compared to lifelong environmental enrichment
(Kalogeraki et al. 2017). Thus, while short-term running can restore OD-plasticity to adult mice
more enrichment, like adding a maze, for longer periods of time, can induce stronger
experience-dependent V1-activity changes. Alternatively, extension of MD-duration can also
boost OD-plasticity in adult SC-mice (Hosang et al., 2018), indicating that there is a trade-off
between age and MD duration in SC-raised animals and that longer MD times can be
shortened by specific environmental interventions, as shown here.
381
Using C57Bl/6J inbred mice housed in the gRW-setup allowed to test how individual
behavioural trajectories affect brain plasticity with minimal influence of genetics on the
observed phenotype. In line with the model of Kempermann (2019), we observed a linear
relationship between running parameters and OD-plasticity quantified by the OD-index:
individual running performance predicted up to 67% of phenotypic variability between mice.
Thus, individual behavioural choices exert a strong influence on experience-dependent
plasticity. Interestingly, bout duration was the best predictor of individual OD-index than bout
number, running speed, running distance and running time, as suggested by the obtained p-
values. Both high running speed and bout duration require high physical fitness and might be
associated with better sensory-motor coordination for efficient running that may improve with
practice (De Bono et al., 2006), suggesting that pre-training could also benefit OD-plasticity
by enabling more efficient running.
394
Wheel running comprises a rewarding behaviour that even mice in the wild pursue voluntarily
(Sherwin, 1998). Multiple factors affecting wheel running have been identified, including
mouse strain, sex, group housing, social hierarchy, age, circadian rhythms (Kopp, 2001; De
Bono et al., 2006; Basterfield et al., 2009; Bains et al., 2018; Balog et al., 2019; Plenz and
Kanold, 2021). In addition, variability in the activity of specific neural networks has been linked
to mouse wheel activity (Rhodes et al., 2003): lateral hypothalamic orexin/hypocretin and
GAD65 networks have emerged as drivers of locomotion initiation (Kosse et al., 2017; Karnani
et al., 2020), suggesting that inter-individual differences in network activity might explain
variability in wheel running. Importantly, social hierarchy has been shown to affect both wheel

running and OD-plasticity in male mice, with dominant males exhibiting stronger OD-shifts
compared to subordinate males (Balog et al., 2019). While social hierarchies are less
prominent in female mice (Williamson et al., 2019) used here, this may have contributed to
variability of OD-index and wheel running in our data.
408
Mechanisms underlying V1-activity changes differ between juvenile and adult mice: In
juveniles, OD-plasticity is initially mediated through depression of V1-responses through the
previously closed (contralateral) eye, while adult mice show OD-shifts initiated by increased
open (ipsilateral) eye V1-responses (Lehmann and Löwel, 2008; Sato and Stryker, 2008).
While we have not examined this in detail, our P70 data suggest that gRW-enrichment enables
OD-plasticity mainly through potentiation of ipsi-eye V1-responses, without contra-eye V1-
response depression. Our data are thus in line with published data showing that, both short-
term running and life-long environmental enrichment restore the *adult* form of OD-plasticity
but can*not* reinstate juvenile-like OD-plasticity (Kalogeraki et al 2017).
418
Together, our newly developed gRW-setup allowing to correlate individual mouse running
behaviour with individual measures of experience-dependent plasticity has demonstrated a
striking correlation between mouse running activity and OD-plasticity, highlighting the
importance of individual behavioural tracking for explaining experimental data. Thus, our
observations add to the growing body of evidence that individual behavioural choices can
strongly affect individual brain plasticity, and thus should be analysed more carefully in future
studies.

References

427
Bains RS, Wells S, Sillito RR, Armstrong JD, Cater HL, Banks G, Nolan PM (2018)
Assessing mouse behaviour throughout the light/dark cycle using automated in-cage
analysis tools. J Neurosci Methods 300:37–47.

Balog J, Hintz F, Isstas M, Teichert M, Winter C, Lehmann K (2019) Social hierarchy
regulates ocular dominance plasticity in adult male mice. Brain Struct Funct
224:3183–3199.

Baroncelli L, Sale A, Viegi A, Maya Vetencourt JF, De Pasquale R, Baldini S, Maffei L (2010)
Experience-dependent reactivation of ocular dominance plasticity in the adult visual
cortex. Exp Neurol 226:100–109.

Basterfield L, Lumley LK, Mathers JC (2009) Wheel running in female C57BL/6J mice:
impact of oestrus and dietary fat and effects on sleep and body mass. Int J Obes
33:212–218.

Bogado Lopes J, Senko AN, Bahnsen K, Geisler D, Kim E, Bernanos M, Cash D, Ehrlich S,
Vernon AC, Kempermann G (2023) Individual behavioral trajectories shape whole-
brain connectivity in mice Lerch JP, Behrens TE, Lerch JP, Yee Y, eds. eLife
12:e80379.

Cang J, Kalatsky VA, Löwel S, Stryker MP (2005) Optical imaging of the intrinsic signal as a
measure of cortical plasticity in the mouse. Vis Neurosci 22:685–691.

De Bono JP, Adlam D, Paterson DJ, Channon KM (2006) Novel quantitative phenotypes of
exercise training in mouse models. Am J Physiol-Regul Integr Comp Physiol
290:R926–R934.

Di Garbo A, Mainardi M, Chillemi S, Maffei L, Caleo M (2011) Environmental enrichment
modulates cortico-cortical interactions in the mouse. PLoS ONE 6.

Fabel K, Wolf S, Ehninger D, Babu H, Galicia P, Kempermann G (2009) Additive effects of
physical exercise and environmental enrichment on adult hippocampal neurogenesis
in mice. Front Neurosci 3.

Greifzu F, Pielecka-Fortuna J, Kalogeraki E, Krempler K, Favaro PD, Schlüter OM, Löwel S
(2014) Environmental enrichment extends ocular dominance plasticity into adulthood
and protects from stroke-induced impairments of plasticity. Proc Natl Acad Sci U S A
111:1150–1155.

Hosang L, Yusifov R, Löwel S (2018) Long-Term Visual Training Increases Visual Acuity and
Long-Term Monocular Deprivation Promotes Ocular Dominance Plasticity in Adult
Standard Cage-Raised Mice. eneuro 5:ENEURO.0289-17.2017.

Kalatsky VA, Stryker MP (2003) New paradigm for optical imaging: Temporally encoded
maps of intrinsic signal. Neuron 38:529–545.

Kalogeraki E, Greifzu F, Haack F, Löwel S (2014) Voluntary physical exercise promotes
ocular dominance plasticity in adult mouse primary visual cortex. J Neurosci
34:15476–15481.

Kalogeraki E, Pielecka-Fortuna J, Löwel S (2017) Environmental enrichment accelerates
ocular dominance plasticity in mouse visual cortex whereas transfer to standard
cages resulted in a rapid loss of increased plasticity. PLOS ONE 12:e0186999.

Kaneko M, Stryker MP (2014) Sensory experience during locomotion promotes recovery of
function in adult visual cortex Nelson SB, ed. eLife 3:e02798.

Karnani MM, Schöne C, Bracey EF, González JA, Viskaitis P, Li H-T, Adamantidis A,
Burdakov D (2020) Role of spontaneous and sensory orexin network dynamics in
rapid locomotion initiation. Prog Neurobiol:101771.

Kempermann G (2019) Environmental enrichment, new neurons and the neurobiology of
individuality. Nat Rev Neurosci 20:235–245.

Kopp C (2001) Locomotor activity rhythm in inbred strains of mice: implications for
behavioural studies. Behav Brain Res 125:93–96.

Kosse C, Schöne C, Bracey E, Burdakov D (2017) Orexin-driven GAD65 network of the
lateral hypothalamus sets physical activity in mice. Proc Natl Acad Sci 114:4525–
4530.

Lehmann K, Löwel S (2008) Age-dependent ocular dominance plasticity in adult mice. PLoS
ONE 3.

Löwel S, Dehmel S, Makowiecki K, Kalogeraki E (2017) Environmental conditions strongly
affect brain plasticity. Neuroforum 24:A19–A29.

Mayr KA, Young L, Molina LA, Tran MA, Whelan PJ (2020) An economical solution to record
and control wheel-running for group-housed mice. J Neurosci Methods 331:108482.

Plenz UT, Kanold PO (2021) Differences in running performance of single- and group-
housed mice. :2021.12.29.474296 Available at:
https://www.biorxiv.org/content/10.1101/2021.12.29.474296v1.

Prusky GT, Alam NM, Beekman S, Douglas RM (2004) Rapid Quantification of Adult and
Developing Mouse Spatial Vision Using a Virtual Optomotor System. Investig
Opthalmology Vis Sci 45:4611.

Reuser A, Wenzel K, Felix SB, Dörr M, Bahls M, Könemann S (2022) Simultaneous
assessment of spontaneous cage activity and voluntary wheel running in group-
housed mice. Sci Rep 12:4444.

Rhodes JS, Garland T, Gammie SC (2003) Patterns of Brain Activity Associated With
Variation in Voluntary Wheel-Running Behavior. Behav Neurosci 117:1243–1256.

Sale A, Maya Vetencourt JF, Medini P, Cenni MC, Baroncelli L, De Pasquale R, Maffei L
(2007) Environmental enrichment in adulthood promotes amblyopia recovery through
a reduction of intracortical inhibition. Nat Neurosci 10:679–681.

Sato M, Stryker MP (2008) Distinctive features of adult ocular dominance plasticity. J
Neurosci Off J Soc Neurosci 28:10278–10286.

Sherwin CM (1998) Voluntary wheel running: a review and novel interpretation. Anim Behav
56:11–27.

Stryker MP, Löwel S (2018) Amblyopia: New molecular/pharmacological and environmental
approaches. Vis Neurosci 35:E018.

Williamson CM, Lee W, DeCasien AR, Lanham A, Romeo RD, Curley JP (2019) Social
hierarchy position in female mice is associated with plasma corticosterone levels and
hypothalamic gene expression. Sci Rep 9:7324.

Figure legends

Figure 1. Gated running wheel setup allows to track individual mouse running wheel activity.

A. Standard-sized rat cage (left, 43 cm x 27 cm) with an additional compartment (right) containing beam break sensors (1,3), a seesaw (2), RFID sensor (4), hall sensor (5), magnet (6), and a running wheel (7). All the sensors are connected to a raspberry pi 4B+ (8). (9) Illustrates the size of the implanted RFID.

B. Comparison of average daily running activity of gRW housed mice (orange) with published data from Kalogeraki et al. 2014 (black).

C. Example of quantified gRW-activity of two co-housed mice illustrating allocation of running bouts to individual mice (blue/black line for P70, orange/black line for P160) for 3 consecutive days and nights (grey background) (top) and 90 minutes during night time (bottom) for P70 (left) and P160 mice (right), with detected running bouts highlighted in gray. Note that animals predominantly run during night time.

526   **D.** Cumulative running trajectories of individual P70/P160 (left/right) mice housed in
527   gRW enrichment for 4/7 days, document a large range of individual running behaviour.
528   **E.** Comparison of running wheel parameters in individual P70 (blue dots) and P160 mice
529   (orange squares); $*p < 0.05$, $**p < 0.01$ in t-test. Note the huge interindividual variability.
530

Figure 2. gRW enrichment boosts ocular dominance plasticity in group-housed
P70 (A,B,C,D) and P160 (E,F,G,H) mice.
A,E. Intrinsic signal optical imaging examples of grey-scale coded activity maps after
visual stimulation of the contralateral and ipsilateral eye in binocular V1 of C57Bl/6J
gRW housed mice after 4 days (P70; A-D) or 7 days (P160, E-H) of monocular
deprivation (MD) and without MD (no MD). The magnitude of V1-activation is expressed
as the fractional change in reflectance $x10^{-4}$ and indicated in the top right. Warm colors
in the color-coded OD-maps indicate contralateral eye dominance, colder colors
indicate an OD-shift towards to ipsilateral eye. Histograms in A,E represent distribution
of pixel wise OD-scores. Scale bar: 1mm.
B,F. Quantification of OD-index. Symbols represent values of individual mice, means
are marked by horizontal lines.
C,G. V1-activation elicited by stimulation of the contralateral (C) or ipsilateral (I) eye
(filled circle indicates MD eye). Note that MD did not reduce V1-activation through the
deprived eye in C.
D,H. Spatial frequency threshold of the optomotor reflex for noMD and MD mice, plotted
vs. days (x-axis). Note that thresholds increase after MD, indicating successfully
induced MD ($*p < 0.05$, $**p < 0.01$, $***p < 0.001$ and $****p < 0.0001$ in t-test or 2-way
ANOVA with Sidak's post hoc test).
550

Figure 3. Individual running wheel (RW) activity correlates with ocular dominance
(OD) plasticity, quantified via the OD-index after monocular deprivation (MD) in
P160 mice.
A. Example intrinsic signal optical imaging recordings from noMD and MD animals, with
low (gRW activity <2.3) and high (gRW activity >2.4) RW performance (gRW activity
was defined as PCA1 of RW parameters displayed in C, and displayed in B). Data
displayed as in figure 2A,D. Note that OD-plasticity after MD is only boosted when mice
run sufficiently in the gRW setup.
B. The OD-index of individual P70 /P160 mice after MD (blue/orange squares) is
correlated with their individual gRW activity. Lines represent linear regression of P160
mice (orange) or pooled data from P70/P160 groups (black). Pearson r-values and p-
values for P160 and pooled P70/P160 in figure.
C. The OD-index of individual mice P160 after MD is correlated with all quantified
running wheel parameter. Data is displayed as in B. For the P160 mice, Pearson
correlation between OD-index and individual wheel running activity was significant for
all tested running wheel parameters. Pearson correlation for pooled P70/P110 data was
significant for all wheel parameters except bout no.
568

Video 1. Video footage of a mouse entering the gated running wheel compartment via
flipping a seesaw.
Video 2. Video footage of a mouse leaving the gated running wheel compartment via
flipping a seesaw. Note that its cage mate is blocked from entering the compartment.
Model 1. 3D model of gated running wheel components
Figure S1. Electronic circuit of gated running wheel setup

Figure 1 Gated running wheel setup allows for detailed readout of wheel running behavior in group housed mice

Figure 2 Gated running wheel housing promotes ODP in group housed mice

Figure 3 Running wheel activity correlates with ODI in P110 animals