Flow cytometry-based isolation combined with ultra-sensitive proteomics of *Salmonella*-containing phagosomes reveals novel insights into host-pathogen interactions

Ritika Chatterjee[1], José Luis Marin Rubio[1,2], Francesca Romana Cianfanelli[1], Andrew Frey[1], Benjamin Bernard Armando Raymond[1], Meihan Meng[1], Abeer Dannoura[1], Tiaan Heunis[1], Mengchun Li[1], Andrew Filby[1], Matthias Trost[1]*

[1] *Biosciences Institute, Newcastle University, Newcastle upon Tyne, UK*
[2] *Current address: : Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, USA*

Correspondence: matthias.trost@ncl.ac.uk

Keywords: Phagosome, flow cytometry, proteomics, macrophage, Salmonella, infection

ABSTRACT

Macrophages engulf pathogens into dynamic phagosomes, which many bacteria manipulate for survival. However, isolating pure pathogen-containing phagosomes remains challenging. Here, we developed a novel flow cytometry-based isolation and ultrasensitive proteomics approach to analyse phagosomal and bacterial proteomes from macrophages infected with wild-type (WT) *Salmonella enterica* serovar Typhimurium (STM), a *ΔphoP* mutant, or dead WT bacteria at 30 min and 4 hrs post-infection. Our approach provides higher throughput, requires lower cell numbers and quantifies more proteins than previous techniques. Our data reveals key host-pathogen interactions, showing induction of PhoP-dependent virulence factors and novel effectors to shape STM's intracellular niche. Notably, phagosomes containing live WT STM resemble in some parts more those with dead bacteria than those with the PhoP mutant. Moreover, our data indicates that bacteria-containing phagosomes recruit mitochondrial membrane for production of reactive oxygen species. These findings provide new insights into *Salmonella*'s manipulation of phagosomal maturation and intracellular niche formation.

Introduction

Phagocytosis is an evolutionary conserved cell function in which professional phagocytes such as macrophages, neutrophils, and dendritic cells engulf and remove particles like dead cells, foreign particles and invading pathogens.[1,2] Upon binding to a wide range of receptors, pathogens are internalised by cytoskeleton rearrangements, leading to the formation of a *de novo* organelle, the phagosome. The phagosome undergoes a series of maturation steps, sequentially fusing with early and late endosomes and ultimately with lysosomes, forming the phagolysosome, within which the ingested particles are degraded. This degradation process generates antigens that are presented to the cell surface, thereby linking innate and adaptive immunity. However, numerous intracellular pathogens, including *Salmonella enterica* serovar Typhimurium (STM), have evolved mechanisms to manipulate phagosome maturation, altering its composition to establish an intracellular replicative niche within macrophages.

Salmonella infection can cause a spectrum of diseases, ranging from self-limiting gastroenteritis to systemic typhoidal fevers, affecting a wide array of hosts. After breaching the intestinal barrier, the bacteria disseminate to secondary infection site through the lymphatic system. During this dissemination macrophages phagocytose the invading bacteria, however, *Salmonella* has evolved a two-component system PhoP/PhoQ[3] which senses reduced Mg^{2+} concentrations, cationic antimicrobial peptides, and acidic pH within phagosomes. The PhoP system activates the expression of virulence genes, including proteins of the *Salmonella* pathogenicity island-2 (SPI-2) encoded type 3 secretion system which enables the injection of effector proteins into the host cytoplasm, modulating host trafficking pathways to support bacterial survival and replication[4].

To comprehensively investigate global changes in phagosome composition, researchers have developed various methods to isolate pathogen-containing phagosomes for proteomic analysis. However, while latex or polystyrene bead phagosome isolation followed by proteomics has been a hugely successful technology to understand the molecular composition of phagosomes of inert beads[4–6], isolating highly pure pathogen-containing vacuoles and phagosomes (for simplicity from now on called phagosomes) remains a technical challenge. This difficulty arises primarily due to the similar densities of mitochondria and bacterial phagosomes, as well as the complex tubular structures of certain pathogen-containing vacuoles, such as *Salmonella*-containing vacuoles (SCVs), which hinder fractionation-based isolation.[5,6] Nonetheless, since the early 2000s, researchers have successfully enriched bacteria-containing phagosomes for proteomics analysis using a variety of methods, including density gradient centrifugation, magnetic bead affinity purification, and biotin affinity purification[7–9].

Here, we utilized an innovative antibody-free technique to isolate *Salmonella*-containing phagosomes from infected human THP-1 macrophages using flow-cytometry that we termed PhagoCyt. Utilising highly sensitive proteomic analyses, we achieve unprecedented depth in characterizing the proteomes of both the phagosome and intra-phagosomal *Salmonella*, revealing the intricate recruitment of host protein complexes and organellar membranes to STM-containing phagosomes.

Results

Isolation of bacteria-containing phagosomes using antibody-free fluorescence-activated cell sorting (FACS).

We have developed a robust flow cytometry-based method, for isolating pathogen-containing phagosomes, which we termed PhagoCyt. This powerful approach enables the isolation of highly pure phagosome populations, which can be analysed using ultrasensitive proteomic techniques to characterise the proteomes of both phagosomes and intraphagosomal pathogens (**Fig.1A**).

Microbes can be labelled either by expressing fluorescent proteins or using live-cell stains such as CellTrace Violet (CTV), allowing for the study of non-genetically modified clinical isolates (**Suppl. Fig.1A**). Importantly, CTV staining does not affect phagocytic uptake (**Suppl. Fig 1B**). We optimised the gating strategy for bacterial phagosomes first on size (**Fig. 1B; Suppl. Fig 1D**). To visualise phagosomal membranes in macrophage lysates, we identified 4 µM CellVue Claret Far Red as optimal, enabling the exclusion of bacteria merely attached to membrane fragments (**Suppl. Fig. 1C**). Importantly, CellVue does not stain bacterial membranes, ensuring specific labelling of host-derived phagosomal structures (**Fig. 1B; Suppl. Fig 1E**).

Human THP-1 cells were differentiated into macrophages using 10 ng/ml Phorbol-12-myristate-13-acetate (PMA), resulting in high expression levels of macrophage markers CD11b, CD11c and CD14 (**Suppl Fig 2A-E**). These macrophages were then infected with BFP-expressing *Salmonella* Typhimurium at a multiplicity of infection (MOI) of 1:100 for 4 hrs.

Figure 1: PhagoCyt workflow and mass spectrometry optimisation.
(**A**) PhagoCyt workflow to isolate *Salmonella*-containing phagosomes by flow cytometry. (**B**) Flow
cytometry strategy and control experiments. Bacterial phagosomes were first selected sorted based
on forward scatter height (FSC-H) and forward scatter area (FSC-A), and then only double positives
for BFP (bacteria) and Cellvue Claret Far Red were selected for sorting. Unstained bacteria (black),
BFP-tagged *Salmonella* alone (blue), BFP-tagged *Salmonella* treated with CellVue Claret Far Red
(purple) and cell lysates containing phagosomes with *Salmonella* not expressing BFP (red) do not
generate any double positive hits. Only BFP-expressing *Salmonella* within phagosomes are double
positive (green). (**C**) Imaging flow cytometry picture of a selected phagosome using ImageStream X
Mark II. (**D**) Mass spectrometry parameters and specifically PASEF windows were optimised for the
timsTOF HT. (**E**) Protein identifications on an EvoSep One/timsTOF HT platform for HeLa digests
using these optimised parameters.

BFP-expression in STM does not affect bacterial growth and survival (**Suppl. Fig 1F,G**). As STM infection induces macrophage cell death at time-points past 4 hours[10], centrifugation steps were included to ensure that only live macrophages were pelleted.

Live, infected macrophages were mechanically lysed using a dounce homogenizer, and unbroken cells and nuclei were removed by centrifugation. Phagosomes were then pelleted by centrifugation from this post-nuclear supernatant, membranes stained with CellVue and analysed by flow cytometry. This revealed double-positive phagosomes containing bacteria and CellVue-stained membranes. Control experiments (**Fig. 1B**) showed that unstained bacteria (black), BFP-tagged *Salmonella* alone (blue), BFP-tagged *Salmonella* treated with CellVue Claret Far Red (purple) and cell lysates containing phagosomes with *Salmonella* not expressing BFP (red) do not generate any double positive signals. Only BFP-expressing *Salmonella* within phagosomes exhibited a double positive signal (green). Imaging cytometry confirmed >95% purity and intact phagosomal membranes (**Fig. 1C**).

Assuming that a single phagosome contains about 0.5% of a cell's content (~0.25 pg), 10^5 phagosomes contain an estimated 25 ng of protein. We optimised sample preparation for these low-input samples, employing the high-recovery S-Trap method for cell lysis and protein digestion. To enhance proteomics sensitivity, we also optimised diaPASEF parameters on a timsTOF HT instrument (**Fig. 1D**). Specifically, we designed a number of variable-width diaPASEF acquisition schemes using pydiAID[11] and tested them with optimal and low loads of tryptic peptides. Testing 500 ng of the protein digests with a 60-min gradient yielded >8000 protein group IDs and >120,000 precursors with the optimal 16 window (W) diaPASEF acquisition scheme. In high-sensitivity mode, the timsTOF HT identified 6000–7000

protein groups and ~60,000 precursors from as little as 25–50 ng of HeLa lysate digest (**Fig. 1E**).

Using this workflow, we performed a proteomics analysis of phagosomes isolated from THP-1 macrophages that were infected with BFP-expressing wild-type (WT) *Salmonella* Typhimurium (STM), a *ΔphoP* mutant, which is unable to respond to the hostile phagosomal environment[3], and dead WT STM, killed by incubation with 4% paraformaldehyde (PFA) for 1 h. For each condition, macrophages were challenged with an MOI of 1:100 for 30 min, followed gentamicin treatment to eliminate extracellular bacteria. Macrophages were then chased for 30 min or 4 h to generate phagosome samples at early/medium and late/matured timepoints.

Maturation of WT STM phagosomes shows intricate interactions with organelles and protein complexes

Phagosome maturation involves dynamic interaction with host organelles and protein complexes[1]. Across six biological replicates, PhagoCyt showed very high reproducibility despite being isolated on different days (**Suppl. Fig 2A-C**) and clear separation of individual conditions (**Suppl. Fig. 2D**). Overall, we identified 5800 proteins, of which were 4625 human and 1175 STM proteins (<1% FDR, minimum of 2 peptides per protein). Stringent filtering, i.e. only taking proteins that have been quantified in 4/6 replicates, reduced that to 4761 quantified proteins of which 3900 were human and 861 bacterial (**Fig. 2A, Suppl. Tables 1-7**). A total of 1331 proteins significantly changed in abundance during phagosome maturation in STM WT phagosomes (q < 0.05, fold-change > 1.5), with 858 human and 444 *Salmonella* proteins increasing, while 557 human and 15 *Salmonella* proteins were decreasing (**Fig. 2A**).

170 Phagosomal proteomes were highly enriched for vesicular, endosomal and lysosomal proteins, along with proteins from the endoplasmic reticulum, mitochondria and nuclear envelope proteins (**Fig. 2B**). ER proteins and some mitochondrial proteins have also been identified on immunologically silent polystyrene-bead phagosomes[12]. However, while recruitment of ER membrane is

175 known for STM phagosomes[11], the identification of mitochondrial proteins in bacteria-containing phagosomes was in previous publications considered a contaminant, specifically when isolated by ultracentrifugation[9], due to the similar density of mitochondria and bacteria. However, our FACS-based isolation does not rely on density gradients. Furthermore, mitochondrial protein abundances varied

180 between conditions (**Fig. 2C**), suggesting that mitochondria may be actively recruited to phagosomes rather than present as passive contaminants.

STM secretes several effector proteins that modify mitochondria[13,14]. Moreover, it has been shown that reactive oxygen species (ROS) generated by mitochondria contribute to killing of phagocytosed bacteria[15] and induction of mitochondrial

185 superoxide by antimycin A improves killing[16]. Moreover, mitochondrial ROS-producing membranes have been shown to be delivered to bacteria-containing phagosomes via mitochondria-derived vesicles (MDVs)[17], indicating that recruitment of mitochondrial membranes to the phagosome represents a host defence mechanism.

Figure 2: Proteomics results show intricate interaction of phagosomes with host organelles.
(**A**) Volcano plot of the proteomics data of STM WT 4h vs 30 min. (**B**) GO-term analysis of organelles identified in STM phagosomes. (**C**) Heatmap of identified mitochondrial proteins shows significant changes of mitochondrial proteins over time and within bacterial strains tested.

Several proteins were highly enriched in 4-hour STM phagosomes, including matrix metalloproteinase-9 (MMP9)[18], which plays a key role in extracellular matrix remodelling, and during bacterial infections it recruits leukocytes and facilitates

200 cytokine, chemokine and defensin activation[19]. Interestingly its inhibitor, metalloproteinase inhibitor 3 (TIMP3)[20] is equally strong more abundant. Other strongly upregulated proteins included IFI30/GILT, a lysosomal thiol reductase involved in disulfide bond reduction; Galectin-3 (LGALS3), which is crucial for phagocytosis and phagosomal membrane repair[21,22]; and Gasdermin-A, a pore-

205 forming protein, that can induce pyroptosis but appears to preferentially target mitochondrial membranes[23]. Additionally, alpha-L-iduronidase IDUA which degrades glycosaminoglycans, was upregulated nearly fourfold, raising the possibility that it may contribute to the sugar degradation of the bacterial surface (**Fig. 3B**).

As expected, most plasma membrane receptors such as the Fc- and cytokine

210 receptors significantly decreased with maturation, either through recycling or degradation (**Fig. 3C**). Interestingly, integrins exhibited differential regulation: some decreased, while others increased over time. Notably, ITGAL and ITGB2, which are critical for *Salmonella* clearance[24,25], were among those increasing. Given their role in actin cytoskeleton regulation, integrin recruitment may contribute to regulating the

215 highly abundant actin cytoskeleton machinery around STM phagosomes (**Fig. 3H**). A particularly striking observation was the significant increase in Toll-like receptor 2 (TLR2) over time (**Fig. 3D**), which contrasts with the behaviour of TLR2 on normally maturing bead phagosomes[26]. This suggests that STM phagosomes may engage distinct signalling pathways during maturation.

Figure 3: Selection of significantly changing proteins and protein complexes in WT STM phagosomes.
(**A-E**) Normalised protein log2 intensities of selected proteins in WT 30 min (early, light blue) and WT 4 hr (late, dark blue) STM phagosomes. **(F-J)** Mean normalised protein log2 intensities of protein
225 complex members of selected protein complexes. Mean log2 fold-changes (FC) were calculated from the log2 differences of the individual proteins. Numbers in the figure represent log2 fold-changes.*: p<0.05; **: p<0.01; ***: p<0.001; ****: p<0.0001, unpaired t-test (FDR controlled).

In a simplified model of phagosome maturation, lysosomal proteins such as
230 cathepsins, increase with maturation. Indeed, these key lysosomal proteases mostly

increase with maturation, but similarly to integrins, not all members behave as expected, with four cathepsins reducing with time (**Fig. 3E**), suggesting that trafficking to the STM phagosome from cellular cathepsin reservoirs is complex and potentially remodelled by STM.

235 Several protein complexes exhibited decreasing abundance with maturation, including septins that are essential for phagosome formation but have also roles in recognising bacterial cell division within phagosomes[27,28] (**Fig. 3F**), the coatomer complex (COP proteins) that transports proteins from the Golgi (**Fig. 3G**) and the aforementioned actin cytoskeleton (**Fig. 3H**). Conversely, protein complexes that

240 increased during maturation included the proteasome (**Fig. 3J**), which has just been shown to generate anti-bacterial peptides[29] as well as peptides for antigen presentation. Fittingly, MHC molecules increase as well during maturation (**Fig. 3K**). These findings highlight the intricate regulation of phagosome maturation, revealing both expected and unexpected protein dynamics that shape the phagosomal

245 environment during *Salmonella* infection.

***Salmonella* actively modulates phagosomal protein recruitment in a PhoP-dependent manner**

Salmonella Typhimurium actively modifies its phagosomal environment by secreting

250 effector proteins, primarily through the *Salmonella* Pathogenicity Islands I and II (SPI-1/SPI-2)[4] . The ability to recognise the acidic phagosome environment through PhoP is essential for the induction of SPI-2 effector proteins, responsible for intracellular survival and replication[3].

Reactive oxygen species (ROS) are essential in killing intracellular pathogens. The

255 cell counteracts oxidative stress using superoxide dismutases which convert

superoxide into H_2O_2. Interestingly, the cytoplasmic superoxide dismutase 1 (SOD1) does not change in the measured conditions. However, the mitochondrial SOD2 is not only significantly more abundant on the phagosome, but also increased hugely with maturation, which is an indication of very high ROS levels at the maturing phagosome (**Fig. 4A, Suppl. Table 7**). This is also supported by the induction of bacterial superoxide dismutases such as SodCA (**Fig. 5B**). Interestingly, while the NADPH oxidase (NOX) complex decreased in WT STM phagosomes, mitochondrial ROS production may compensate for this decline, specifically as interferon-induced nitric oxide synthases (iNOS) were undetectable at the analysed time points (**Fig. 4A**).

Glycolysis is essential for macrophages to clear *Salmonella* Typhimurium. Despite previous reports suggesting that STM reduces overall glycolytic activity in infected macrophages[30], we observed an increased association of glycolytic proteins with maturing phagosomes. However, this pattern was reversed in the PhoP mutant, suggesting that PhoP regulates metabolic rewiring at the phagosome (**Fig. 4B**).

Notably, several proteins that increase with maturation in STM WT phagosomes—including MMP9, TIMP3, Galectin-3, and Gasdermin-A—exhibited opposite trends in the PhoP mutant (**Fig. 4C**). A similar reversal was observed for interferon-induced proteins, such as ISG15, its ligase HERC5, and IFI30/GILT, which decreased with maturation in PhoP mutant phagosomes. Overall, the data suggests that live and dead WT STM phagosomes are more similar to each other than the phagosomes of the live STM PhoP-mutant (**Fig. 4D**). This suggests that PhoP-dependent effectors actively remodel the phagosomal environment, influencing key immune and metabolic processes.

Figure 4: Selected protein changes in phagosomes of the PhoP-mutant and dead WT STM.
(A) Normalised protein log2 intensities of selected proteins in superoxide conversion and reactive oxygen species generation in early (30 min) and late (4 h) phagosomes of (live) WT STM, PhoP mutant and dead WT STM. **(B)** Mean normalised protein log2 intensities of proteins involved in
285 glycolysis and glucogenesis. Mean log2 fold-changes (FC) were calculated from the log2 differences of the individual proteins. **(C)** Normalised protein log2 intensities of selected significantly changing proteins. Numbers in the figure represent log2 fold-changes. **(D)** Principal component analysis (PCA) of all proteomes shows separation of individual sample groups. Circles represent 95% confidence interval.
290 *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$; ****: $p < 0.0001$, unpaired t-test (FDR controlled).

PhagoCyt identifies novel putative *Salmonella* effector proteins

STM utilises the PhoP/Q system to sense vacuolar low pH, which induces the expression of effector proteins secreted through the SPI-2 T3SS. These effectors are critical for bacterial survival and replication within the host[31].

295 Consistent with previous studies[3], we observed a strong induction of PhoP-regulated genes in STM phagosomes at the 4-hour time point, compared to 30-minute phagosomes (**Fig. 5A**). This included key PhoP/Q system components (PhoP, PhoQ, PhoN, PhoU, PhoB) as well as the phosphate transporter PstS, part of the ABC transporter complex PstSACB, which is activated by PhoB in response to

300 phosphate starvation[32]. Additionally, we observed a significant increase in the outer membrane adhesin and invasin protein PagN, which facilitates adaptation to intracellular host niches[33], as well as the PhoP-dependent upregulation of PagC, which has been implicated in bacterial outer membrane vesicle (OMV) formation[34,35]. Finally, we detected a marked increase in PdgL (PcgL), a periplasmic D-Ala-D-Ala

305 dipeptidase. The identification of these well-documented effector proteins validates our approach.

Beyond known effectors, PhagoCyt identified several putative effectors. Among the most highly upregulated proteins at 4 hours were SL1344_1263 (lipid A deacylase LpxR family protein)[36] which is regulated by HilD[37], SL1344_1563, a predicted-cell

310 envelope protein, and SL1344_3560, a predicted- phosphatidic acid phosphatase (**Fig. 5A**). The latter two have been shown to be essential for establishing infection in the intestinal milieu in a transposon screen[38].

To counter high levels of ROS generated by the host cell, STM significantly induces the abundance of the ROS-scavenging proteins, including the superoxide dismutases SodCa and SodA, the methionine sulfoxide reductase A (MsrA), and the alkyl hydroperoxide reductase AhpC[39]. We also observed increased levels of the Iron-Sulfur (Fe-S) cluster assembly proteins (SufC, SufD, SufI, and SufS) which support survival in macrophages[40] and act as potent negative regulators of SPI-1 thereby helping STM establish a competent niche inside host cells[41] (**Fig. 5B**).

While a large number of bacterial proteins increased in abundance during phagosome maturation, only a smaller subset showed decreased levels. These include FlgK, a flagellar gene which is downregulated upon entering the cell to evade the host innate-immune pathways of recognition of flagellar components as PAMPs[42]. Furthermore, we observed a reduction in the SPI-1 injectisome complex proteins such as SctB1[43] and also effectors delivered by SPI-1 such as SipB, SopB and SopE, validating the existing literature. However, we also identified novel potential effectors of early infection, including the mostly uncharacterised Copper resistance protein YobA and lipoprotein YbaY, which is involved in biofilm formation, both of which are about 2-fold more abundant in the early phagosomes (**Fig. 5C**). This data suggests that PhagoCyt can serve as a powerful tool to identify novel microbial effector proteins of intracellular pathogens.

Figure 5: Selected bacterial protein changes in WT STM within phagosomes. (A) Normalised protein log2 intensities of selected bacterial proteins induced in late (4 hr) Salmonella phagosomes. **(B)** Normalised protein log2 intensities of selected bacterial proteins of redox and iron-sulfur cluster assembly proteins. **(C)** Normalised protein log2 intensities of selected bacterial proteins reduced in abundance in late (4 hr) *Salmonella* phagosomes Numbers in the figure represent log2 fold-changes. *: $p<0.05$; **: $p<0.01$; ***: $p<0.001$; ****: $p<0.0001$, unpaired t-test (FDR controlled).

Discussion

Phagosomes are highly dynamic organelles that interact with practically all cellular membrane trafficking pathways[1]. Due to this dynamic nature, proteomics of isolated phagosomes has been a powerful tool to identify unbiasedly novel biological insights. Polystyrene bead phagosome can be isolated to high purity by floatation in ultracentrifugation gradients and have therefore been extensively used to study these membrane dynamics and how macrophage activation affects phagosomal maturation[12,26,44,45]. The isolation of pathogen-containing phagosomes, however, has been much more difficult as many intracellular pathogens, and specifically bacteria, have a similar density to host mitochondria and are therefore difficult to isolate through centrifugation alone. Additional steps such as magnetic bead affinity purification or biotin affinity purification[7–9,46] were used in attempts to overcome these problems. Nonetheless, nuclear and mitochondrial proteins were often considered "contaminants" as their presence in phagosomes was considered a side effect of the isolation procedures. The here presented approach was carefully optimised to obtain a phagosome-enriched cellular supernatant from which phagosomes were then isolated using a highly sensitive flow cytometry approach. Our approach is not only relatively fast compared to other approaches but – when combined with low-input proteomics – also requires fewer cells. We only injected the digests of the equivalent of 100,000 phagosomes (~25 ng) onto the mass spectrometer.

While flow cytometry (phagoFACS) has been used to analyse phagosomal subpopulations[47,48] and isolate intracellular bacteria[49], this is the first time where flow cytometry has been used to isolate intact phagosomes. The power of this approach is that not only the phagosomal host proteome, but also the bacterial proteome is available through the proteomics analysis. This data showed the induction of many

365 known SPI-2 effector proteins, validating our approach. Additionally, it identified several novel putative effectors that will require further study. Furthermore, this study is the first to isolate *Salmonella*-containing vacuoles (SCV)/phagosomes from macrophages as previous studies were performed in HeLa cells[50] and *Dictyostelium discoideum*[51].

370 Overall, this method will enhance the ability to isolate pathogen containing phagosome and allow the unbiased molecular characterisation of host-pathogen interaction at this important organelle.

Online Methods

Antibodies

- BUV737 Rat Anti-CD11b (#612801, BD Biosciences).

- BUV737 Rat IgG2b, κ Isotype Control (#612762, BD Biosciences).

- CD11c Monoclonal Antibody (N418), APC (#17-0114-82, Invitrogen).

- Armenian Hamster IgG Isotype Control (eBio299Arm), APC (#17-4888-82,

380   Invitrogen).

- CD14 Monoclonal Antibody (61D3), Alexa Fluor 700 (#56-0149-42, Invitrogen).

- Mouse IgG1 kappa Isotype Control (P3.6.2.8.1), Alexa Fluor 700 (#56-4714-80, Invitrogen).

- FITC anti-human CD44 Antibody (#338803, BioLegend).

- FITC Mouse IgG1, κ Isotype Control Antibody (#400107, BioLegend).

- Alexa Fluor 488 (AF488) Goat anti-Rabbit IgG antibody (Invitrogen).

THP-1 cell culture and differentiation into macrophages.

390   The human monocytic leukaemia cell line THP-1 (TIB-202, ATCC) was cultured in RPMI-1640 medium (Gibco) supplemented with 10 % FBS and 4 mM L-glutamine at 37°C in a humidified 5 % CO_2 atmosphere. ATCC routinely performs cell line authentication using short tandem repeat profiling. Cell experimentation was always performed within a period not exceeding 20 passages after resuscitation and cells

395   were regularly tested for mycoplasma.

To differentiate THP-1 cells into macrophages, we cultured THP-1 cells at a concentration of 5×10^5 cells/mL with 10 ng/mL Phorbol 12-myristate 13-acetate (PMA) (Sigma-Aldrich) for 48 h, followed by 24 h with fresh medium without PMA.

These culture conditions were optimised by evaluating cell viability and immunophenotype using CD11b, CD11c, CD14, and CD44 (**Supplementary Fig. 1**). We assessed cell viability by propidium iodide staining.

Bacterial strains and culture conditions

Salmonella enterica serovar Typhimurium SL1344 and genetically modified *Salmonella* (**Supplementary Table 1**) were kindly provided by Dirk Bumann (Biozentrum Basel) and grown in LB broth at 37⁰C with constant rotation.

Table 1. *Salmonella* strain list.

Salmonella strain	Deletion (Δ) / Induced promoter (P)	Reporter	Constitutive expression
Wild-type (WT) SL1344	-	-	BFP
ΔPhoP	*ΔPhoP*	-	BFP
Acidity response	P_{Asr}	GFP	BFP

BFP, blue fluorescent protein; GFP, green fluorescent protein.

Bacterial infection

For each experimental replicate, 4×10^7 THP-1 cells were differentiated into macrophages, seeding 1×10^7 cells per 150 mm tissue culture dish. Before infection, the cell medium was refreshed with 10 mL of complete medium (washed three times with warm PBS before refreshing the medium). Overnight bacteria cultures (optical density (OD600 nm) of 1.0 – 1.2 , where OD= 1 is equals to approximately 1×10^9 bacteria/mL), were washed in warm PBS twice at 4,000 rpm for 15 min and, subsequently, resuspended in warm cell culture medium at the desired MOI of 100 at a concentration of 1×10^9/mL. Bacterial input was placed on LB agar plates to count

colony-forming units (CFUs). Then bacteria were added at an MOI of 1:100 and incubated for 30 min for uptake. This produced the 30 min time-point. After 30 min, we removed the bacteria-containing medium, washed twice with warm PBS and refreshed the medium with 50 µg/mL gentamicin. After one hour, cells were rewashed twice with warm PBS and incubated with fresh medium supplemented with 15 µg/mL gentamicin up to 4 h (4 h post-infection or p.i.). CFUs were determined by lysing cells in 0.1 % Triton-X 100 in PBS at indicated time points.

Bacterial staining with CellTrace Violet (CTV)

Before the infection, untagged bacteria were stained with different concentrations of CellTrace Violet (CTV) dye (Thermo-Fisher Scientific), following the manufacturer's protocol. Higher concentration of CTV dye tested, 500 µM, stained 97 % of bacteria, and its fluorescence intensity was compatible with mTagBFP. Moreover, CTV dye did not affect bacteria uptake (**Supplementary Fig. 2B**). Briefly, the OD_{600} was measured for overnight culture and about 10^9 bacteria were incubated in 500 µl of warm PBS with 500 µM of CTV stain (resuspend immediately to achieve homogenous staining) for 20 min at 37°C. Then, the bacteria were washed twice with PBS (at 37°C) for 5 min and centrifugated at 4,000 rpm for 15 min. Finally, bacteria were resuspended in a warm cell culture medium at a MOI of 100.

Cell lysis for phagosome isolation.

After the indicated time point (30 minutes for uptake and 4 h p.i.), the medium was removed, and ice-cold PBS was added. Then, cells were scraped from four dishes, transferred into 15 mL tubes, and centrifuged at 300 xg at 4°C for 10 min. Cell pellets were washed twice with ice-cold PBS. Afterwards, pellets were re-suspended in 1 mL hypotonic buffer (250 mM sucrose, 3 mM imidazole, pH 7.4) with protease and phosphatase inhibitors and 0.1μl (25 U) DNase (Pierce Universal Nuclease for Cell Lysis, Thermo-Fisher Scientific). To lyse the cells, we used a Dounce homogeniser in a class II biological safety cabinet (between 5 – 7 strokes until about 70% of cells were broken). We monitored the lysis using trypan blue dye under the microscope. Cell lysates were centrifuged at 1,200 xg at 4°C for 10 min to pellet nuclei. The post-nuclear supernatant I was then transferred into a 2 mL tube, and again treated with DNase for 5mins at 37°C and the nuclei were pelleted again at 1,200 x g for 5 mins. The post-nuclear supernatant II was then transferred into a fresh 2 mL tube and diluted with the same volume of ice-cold PBS and centrifuged at 2,500 xg at 4°C for 5 min to pellet phagosomes.

CellVue Claret Far Red staining

We used the CellVue Claret Far Red Membrane Label (Sigma-Aldrich) for phagosomal membrane staining as per manufacturer's protocol. In brief, we resuspended the phagosome pellet in 50 μl of diluent C. Then, 50 μl with 4 μM dye was added, mixed and incubated for 5 min at room temperature. The staining was quenched using 100 μL of 1% BSA in PBS and washed twice with ice-cold PBS by centrifugation at 2,500 xg at 4°C for 5 min. Samples were fixed with 0.5 % PFA in

PBS at 4°C for 15 min, followed by one washed with PBS. Finally, we resuspended the samples in 1 mL of PBS with 1 mM EDTA for flow cytometry.

Flow cytometry

For immunophenotype analysis, cells were blocked with flow cytometry staining buffer at 4°C for 20 min, then incubated with the corresponding antibody at a 1:100 dilution at 4°C for 20 min, washed with PBS, and analysed using a BD Symphony A5

470   flow cytometer (BD Biosciences).

The samples were then flow sorted for double positives (BFP+ and Claret Far Red+) on a BD FACS Aria III. The BFP excitation was set to 399 nm and emission to 454 nm and for Cell Vue Claret Far Red excitation to 655 nm and emission to 675 nm. The initial gating strategy involved the selection for bacterial phagosome size

475   followed by gating based on single channel controls. All samples were sorted based on a double positive gating channel and the analysis was performed using FlowJo software.

Sample preparation for LC-MS/MS

480   After sorting, phagosomes were pelleted by centrifugation at 16,000 xg for 15 min at 4°C, resuspended in PBS supplemented with protease inhibitors, transferred to low protein binding microcentrifuge tubes. Next, the samples were resuspended in a 50 µL lysis buffer (5% SDS and 50 mM triethylammonium bicarbonate (TEAB) in HPLC water, pH 7.4) supplemented with protease inhibitors (cOmplete, EDTA-free, Roche).

485   To prepare the samples for mass spectrometry, S-Trap micro columns (Protifi) were used according to the manufacturer's recommended protocol for high recovery. Proteins were reduced with 20 mM tris(2-carboxyethyl)phosphine at 37ºC for 30 min

and alkylated with 20 mM iodoacetamide in the dark for 30 min. The proteins were then acidified with 5 μL 55 % phosphoric acid and 330 μL S-Trap binding buffer. 1

490   micrograms of Sequencing Grade Modified Trypsin (Promega) were added to the acidified samples before being transferred into the S-Trap column. The columns were washed five times with 150 μL S-Trap binding buffer before incubating them with 0.5 μg of Sequencing Grade Modified Trypsin in 25 μl of 50 mM TEAB, pH 8 for 2 h at 47°C. Finally, the samples were sequentially eluted in 40 μl 50 mM TEAB, pH

495   8, 40 μl 0.2 % formic acid and 35 μl 50 % acetonitrile with 0.2 % formic acid. The eluted peptides were dried in a vacuum concentrator and stored in -80°C until further use.

Data-independent acquisition mass spectrometry (DIA-MS)

500   Liquid chromatography (LC) was performed using an Evosep One system with a 15 cm Aurora Elite C18 column with integrated captive spray emitter (IonOpticks), at 50 °C. Buffer A was 0.1 % formic acid in HPLC water, buffer B was 0.1 % formic acid in acetonitrile. Immediately prior to LC-MS, peptides were resuspended in 30 μl of buffer A and a volume of peptides equivalent to 20 % of the sample (100,000 events

505   isolated during the FACS) was loaded onto the LC system-specific C18 EvoTips, according to manufacturer instructions, and subjected to the predefined Whisper100 or Whisper-Zoom 20 SPD protocol (where the gradient is 0-35 % buffer B, 100 nl/min or 200 nl/min, for 58 minutes). The Evosep One (EvoSep) was used in line with a timsTOF-HT mass spectrometer (Bruker) operated in DIA-PASEF mode with high

510   sensitivity on. Mass and IM ranges were 300-1200 m/z and 0.6-1.45 1/K_0, DIA-PASEF was performed using variable width IM-m/z windows without overlap, designed using py_diAID[11], and are provided (see data sharing section). TIMS ramp

and accumulation times were 100 ms, total cycle time was ~1.8 seconds. Collision

energy was applied in a linear fashion, where ion mobility = 0.6-1.6 $1/K_0$, and

515   collision energy = 20 - 59 eV.

MS data analysis

Raw data files were searched using DIA-NN 1.9,[52] using its *in silico* generated

spectral library from the *Homo sapiens* (downloaded from Uniprot on the 14th of

520   September 2023, containing 42437 entries) and *Salmonella* Typhimurium

(downloaded from Uniprot on the 14th of September 2023 containing 4659 entries)

proteomes and a common contaminants list[53]. Carbamidomethylation (C) was set as

a fixed modification, and oxidation (M) to variable. Precursor masses from 300 to

1800 were included, with charge 1 to 4. Mass accuracy for MS1 and MS2 was set to

525   15, Match between runs was performed and normalisation set to 'Global'. Trypsin

specificity was set to one missed cleavage and a protein and PSM false discovery

rate of <1%, respectively. Stringent filtering was applied: Proteins required intensities

in a minimum of 3/6 in one group, and 0,3-6 in the other and proteins required to

have a minimum of 2 peptides.

530   Proteomics data analysis was performed on a R software pipeline (v4.2.1). Raw

intensities were log2-transformed. Protein groups detected in at least four samples

within any experimental group were kept for downstream analysis. For some data,

missing values were imputed by v2-mnar method implemented in msImpute

package[54] (v1.7.0). Differential expression between groups was assessed by fitting a

535   linear model and applying empirical Bayes moderated t-statistics in limma package[55]

(v3.60.4). Resulting P-values were adjusted using the Benjamini–Hochberg FDR

correction. Proteins with a log2-fold change ≥ ±0.5 and adjusted p-value ≤ 0.05 were considered as differentially expressed.

540 **Data availability statement**

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository[56] with the data set identifier PXD062621.

Acknowledgements

This research was funded by a Wellcome Trust Investigator Award (215542/Z/19/Z) to MT, a Marie Skłodowska-Curie and a Swiss National Science Foundation Fellowship to FRC, and a BBSRC studentship to MM. The mass spectrometer used in this work was funded by a Wellcome Trust multi-user equipment grant (212947/Z/18/Z). We thank Dr Camila Valenzuela-Montenegro and Prof Jost Enninga for critical proof reading of the manuscript.

References

1.   Pauwels, A.-M., Trost, M., Beyaert, R. & Hoffmann, E. Patterns, Receptors, and Signals:
Regulation of Phagosome Maturation. *Trends Immunol.* **38**, 407–422 (2017).

2.   Boulais, J. *et al.* Molecular characterization of the evolution of phagosomes. *Mol. Syst. Biol.*
6, 423 (2010).

3.   Groisman, E. A., Duprey, A. & Choi, J. How the PhoP/PhoQ System Controls Virulence and
Mg2+ Homeostasis: Lessons in Signal Transduction, Pathogenesis, Physiology, and Evolution.
Microbiol. Mol. Biol. Rev. MMBR **85**, e0017620 (2021).

4.   Jennings, E., Thurston, T. L. M. & Holden, D. W. Salmonella SPI-2 Type III Secretion System
Effectors: Molecular Mechanisms And Physiological Consequences. *Cell Host Microbe* **22**, 217–231
(2017).

5.   Vorwerk, S., Krieger, V., Deiwick, J., Hensel, M. & Hansmeier, N. Proteomes of host cell
membranes modified by intracellular activities of Salmonella enterica. *Mol. Cell. Proteomics MCP* **14**,
81–92 (2015).

6.   Herweg, J.-A. *et al.* Purification and proteomics of pathogen-modified vacuoles and
membranes. *Front. Cell. Infect. Microbiol.* **5**, 48 (2015).

7.   Lönnbro, P., Nordenfelt, P. & Tapper, H. Isolation of bacteria-containing phagosomes by
magnetic selection. *BMC Cell Biol.* **9**, 35 (2008).

8.   Lührmann, A. & Haas, A. A method to purify bacteria-containing phagosomes from infected
macrophages. *Methods Cell Sci. Off. J. Soc. Vitro Biol.* **22**, 329–341 (2000).

9.   Stévenin, V. *et al.* Dynamic Growth and Shrinkage of the Salmonella-Containing Vacuole
Determines the Intracellular Pathogen Niche. *Cell Rep.* **29**, 3958-3973.e7 (2019).

10.   Starr, T., Bauler, T. J., Malik-Kale, P. & Steele-Mortimer, O. The phorbol 12-myristate-13-
acetate differentiation protocol is critical to the interaction of THP-1 macrophages with Salmonella
Typhimurium. *PloS One* **13**, e0193601 (2018).

11.   Skowronek, P. *et al.* Rapid and In-Depth Coverage of the (Phospho-)Proteome With Deep
Libraries and Optimal Window Design for dia-PASEF. *Mol. Cell. Proteomics MCP* **21**, 100279 (2022).

12.   Bilkei-Gorzo, O. *et al.* The E3 ubiquitin ligase RNF115 regulates phagosome maturation and
host response to bacterial infection. *EMBO J.* **41**, e108970 (2022).

13.   Achi, S. C. *et al.* Proteome profiling identifies a link between the mitochondrial pathways and
host-microbial sensor ELMO1 following Salmonella infection. *bioRxiv* 2024.05.03.592405
(2024) doi:10.1101/2024.05.03.592405.

14.   Chen, H.-Y. *et al.* Mitochondrial injury induced by a Salmonella genotoxin triggers the
proinflammatory senescence-associated secretory phenotype. *Nat. Commun.* **15**, 2778 (2024).

15.   West, A. P. *et al.* TLR signalling augments macrophage bactericidal activity through
mitochondrial ROS. *Nature* **472**, 476–480 (2011).

16.   Krause, K. *et al.* Caspase-11 counteracts mitochondrial ROS-mediated clearance of
Staphylococcus aureus in macrophages. *EMBO Rep.* **20**, e48109 (2019).

17.   Abuaita, B. H., Schultz, T. L. & O'Riordan, M. X. Mitochondria-Derived Vesicles Deliver Antimicrobial Reactive Oxygen Species to Control Phagosome-Localized Staphylococcus aureus. *Cell Host Microbe* **24**, 625-636.e5 (2018).

18.   Lee, H. S. & Kim, W. J. The Role of Matrix Metalloproteinase in Inflammation with a Focus on Infectious Diseases. *Int. J. Mol. Sci.* **23**, 10546 (2022).

19.   Elkington, P. T. G., O'Kane, C. M. & Friedland, J. S. The paradox of matrix metalloproteinases in infectious disease. *Clin. Exp. Immunol.* **142**, 12–20 (2005).

20.   Fan, D. & Kassiri, Z. Biology of Tissue Inhibitor of Metalloproteinase 3 (TIMP3), and Its Therapeutic Implications in Cardiovascular Pathology. *Front. Physiol.* **11**, 661 (2020).

21.   Jia, J. *et al.* Galectin-3 Coordinates a Cellular System for Lysosomal Repair and Removal. *Dev. Cell* **52**, 69-87.e8 (2020).

22.   Sano, H. *et al.* Critical role of galectin-3 in phagocytosis by macrophages. *J. Clin. Invest.* **112**, 389–397 (2003).

23.   Kondolf, H., Da, D., Gr, D. & Dw, A. Protein engineering reveals that gasdermin A preferentially targets mitochondrial membranes over the plasma membrane during pyroptosis. *J. Biol. Chem.* **299**, (2023).

24.   Nascimento, D. de O. *et al.* Integrin αDβ2 (CD11d/CD18) Modulates Leukocyte Accumulation, Pathogen Clearance, and Pyroptosis in Experimental Salmonella Typhimurium Infection. *Front. Immunol.* **9**, 1128 (2018).

25.   Zhang, J. *et al.* A Loss-of-Function Mutation in the Integrin Alpha L (Itgal) Gene Contributes to Susceptibility to Salmonella enterica Serovar Typhimurium Infection in Collaborative Cross Strain CC042. *Infect. Immun.* **88**, e00656-19 (2019).

26.   Dill, B. D. *et al.* Quantitative proteome analysis of temporally resolved phagosomes following uptake via key phagocytic receptors. *Mol. Cell. Proteomics MCP* **14**, 1334–1349 (2015).

27.   Huang, Y.-W. *et al.* Mammalian septins are required for phagosome formation. *Mol. Biol. Cell* **19**, 1717–1726 (2008).

28.   Robertin, S. & Mostowy, S. The history of septin biology and bacterial infection. *Cell. Microbiol.* **22**, e13173 (2020).

29.   Goldberg, K. *et al.* Cell-autonomous innate immunity by proteasome-derived defence peptides. *Nature* **639**, 1032–1041 (2025).

30.   Gutiérrez, S. *et al.* Salmonella Typhimurium impairs glycolysis-mediated acidification of phagosomes to evade macrophage defense. *PLoS Pathog.* **17**, e1009943 (2021).

31.   Garvis, S. G., Beuzón, C. R. & Holden, D. W. A role for the PhoP/Q regulon in inhibition of fusion between lysosomes and Salmonella-containing vacuoles in macrophages. *Cell. Microbiol.* **3**, 731–744 (2001).

32.   Bruna, R. E., Kendra, C. G. & Pontes, M. H. An intracellular phosphorus-starvation signal activates the PhoB/PhoR two-component system in Salmonella enterica. *mBio* **15**, e0164224 (2024).

33.   Lambert, M. A. & Smith, S. G. J. The PagN protein of Salmonella enterica serovar Typhimurium is an adhesin and invasin. *BMC Microbiol.* **8**, 142 (2008).

630   34.   Dehinwal, R. *et al.* Increased Production of Outer Membrane Vesicles by Salmonella Interferes with Complement-Mediated Innate Immune Attack. *mBio* **12**, e0086921 (2021).

35.   Kitagawa, R. *et al.* Biogenesis of Salmonella enterica serovar typhimurium membrane vesicles provoked by induction of PagC. *J. Bacteriol.* **192**, 5645–5656 (2010).

36.   Reynolds, C. M. *et al.* An outer membrane enzyme encoded by Salmonella typhimurium lpxR
635   that removes the 3'-acyloxyacyl moiety of lipid A. *J. Biol. Chem.* **281**, 21974–21987 (2006).

37.   Petrone, B. L., Stringer, A. M. & Wade, J. T. Identification of HilD-regulated genes in Salmonella enterica serovar Typhimurium. *J. Bacteriol.* **196**, 1094–1101 (2014).

38.   Chaudhuri, R. R. *et al.* Comprehensive assignment of roles for Salmonella typhimurium genes in intestinal colonization of food-producing animals. *PLoS Genet.* **9**, e1003456 (2013).

640   39.   De Groote, M. A. *et al.* Periplasmic superoxide dismutase protects Salmonella from products of phagocyte NADPH-oxidase and nitric oxide synthase. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 13997–14001 (1997).

40.   Wang, B. X. *et al.* High-throughput fitness experiments reveal specific vulnerabilities of human-adapted Salmonella during stress and infection. *Nat. Genet.* **56**, 1288–1299 (2024).

645   41.   Fontecave, M., Py, B., Ollagnier de Choudens, S. & Barras, F. From Iron and Cysteine to Iron-Sulfur Clusters: the Biogenesis Protein Machineries. *EcoSal Plus* **3**, (2008).

42.   Li, B. *et al.* Salmonella STM1697 coordinates flagella biogenesis and virulence by restricting flagellar master protein FlhD4C2 from recruiting RNA polymerase. *Nucleic Acids Res.* **45**, 9976–9989 (2017).

650   43.   Godlee, C. & Holden, D. W. Transmembrane substrates of type three secretion system injectisomes. *Microbiol. Read. Engl.* **169**, 001292 (2023).

44.   Trost, M. *et al.* The phagosomal proteome in interferon-gamma-activated macrophages. *Immunity* **30**, 143–154 (2009).

45.   Guo, M. *et al.* Triggering MSR1 promotes JNK-mediated inflammation in IL-4-activated
655   macrophages. *EMBO J.* **38**, (2019).

46.   Steinhäuser, C. *et al.* Lipid-labeling facilitates a novel magnetic isolation procedure to characterize pathogen-containing phagosomes. *Traffic Cph. Den.* **14**, 321–336 (2013).

47.   E, H. *et al.* Autonomous phagosomal degradation and antigen presentation in dendritic cells. *Proc. Natl. Acad. Sci. U. S. A.* **109**, (2012).

660   48.   Hoffmann, E. *et al.* Autonomous phagosomal degradation and antigen presentation in dendritic cells. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 14556–14561 (2012).

49.   Becker, D. *et al.* Robust Salmonella metabolism limits possibilities for new antimicrobials. *Nature* **440**, (2006).

50.   Stévenin, V. *et al.* Dynamic Growth and Shrinkage of the Salmonella-Containing Vacuole
665   Determines the Intracellular Pathogen Niche. *Cell Rep.* **29**, (2019).

51.   Valenzuela, C. *et al.* SopB- and SifA-dependent shaping of the Salmonella-containing vacuole proteome in the social amoeba Dictyostelium discoideum. *Cell. Microbiol.* **23**, (2021).

52.   Demichev, V., Messner, C. B., Vernardis, S. I., Lilley, K. S. & Ralser, M. DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. *Nat.*
670   *Methods* **17**, 41–44 (2020).

53.   Frankenfield, A. M., Ni, J., Ahmed, M. & Hao, L. Protein Contaminants Matter: Building Universal Protein Contaminant Libraries for DDA and DIA Proteomics. *J. Proteome Res.* **21**, 2104–2113 (2022).

54.   Hediyeh-Zadeh, S., Webb, A. I. & Davis, M. J. MsImpute: Estimation of Missing Peptide
675   Intensity Data in Label-Free Quantitative Mass Spectrometry. *Mol. Cell. Proteomics MCP* **22**, 100558 (2023).

55.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

56.   Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019:
680   improving support for quantification data. *Nucleic Acids Res.* **47**, D442–D450 (2019).