Engineering NIR-Sighted Bacteria

Stefanie S. M. Meier[1,‡], Michael Hörzing[1,‡], Cornelia Böhm[2,3,‡], Emma L. R. Düthorn[1,‡], Heikki Takala[2,‡], René Uebe[4,5,‡], Andreas Möglich[1,5,6,*,‡]

[1] Department of Biochemistry, University of Bayreuth, 95447 Bayreuth, Germany.
[2] Department of Biological and Environmental Science, Nanoscience Center, University of Jyvaskyla, Jyvaskyla 40014, Finland.
[3] Institute of Biochemistry, Graz University of Technology, 8010 Graz, Austria
[4] Department of Microbiology, University of Bayreuth, 95447 Bayreuth, Germany.
[5] Bayreuth Center for Biochemistry & Molecular Biology, Universität Bayreuth, 95447 Bayreuth, Germany.
[6] North-Bavarian NMR Center, Universität Bayreuth, 95447 Bayreuth, Germany.

* for correspondence: andreas.moeglich@uni-bayreuth.de
‡ ORCID identifiers: S.S.M.M. 0009-0002-6028-5223; M.H. 0009-0008-2301-3783; C.B. 0000-0002-3552-2349; E.L.R.D. 0009-0000-4156-5428; H.T. 0000-0003-2518-8583; R.U. 0000-0003-2357-1589; A.M. 0000-0002-7382-2772

Abstract

Spatially and temporally orchestrated gene expression underpins organismal development, physiology, and adaptation. In bacteria, two-component systems (TCS) translate environmental cues into inducible expression outputs. Inducible expression also serves as a versatile instrument in both basic and applied science. Here, we harness the photosensors of rhizobial bathy-phytochromes to construct synthetic TCSs for stringent activation of gene expression by near-infrared (NIR) light in laboratory and probiotic *Escherichia coli* strains, and in *Agrobacterium tumefaciens*. Orthogonal TCSs afford the multiplexed expression control of several genes by NIR and visible light. Notwithstanding substantial photochemical activation of bathy-phytochromes by visible radiation, the NIR-light-responsive systems hardly responded to red light. Evidently, light signals can be processed by TCSs into highly nonlinear responses at the physiological relevant level of gene expression. These fundamental aspects likely extend to naturally occurring TCSs. Depending on their photosensor traits

and environmental conditions, bathy-phytochromes may thus either be NIR-specific or function as

colorblind receptors of light vs. darkness.

Keywords

bacterial phytochrome; gene expression; near-infrared light; optogenetics; sensor histidine kinase;

sensory photoreceptor; two-component system

Introduction

Proportionate to its essential roles in organismal physiology, development, replication, and

adaptation, gene expression is stringently and precisely regulated. Various cellular circuits integrate

cognate input signals and, in turn, ramp up or down the expression of target genes. Molecular

analyses of these circuits not only benefit fundamental research, but also, they underlie applications

in biotechnology and synthetic biology for the precise control of gene expression. Two-component

systems (TCS) are the predominant signal-transduction pathways by which bacteria and several

other organisms process environmental stimuli and convert them into intracellular gene-expression

responses (Buschiazzo and Trajtenberg, 2019; Parkinson and Kofoid, 1992). In their canonical form,

TCSs comprise as one component a membrane-spanning, homodimeric sensor histidine kinase (SHK)

that relays signals to the cell interior. SHKs generally consist of N-terminal extracellular/periplasmic

sensor and C-terminal cytosolic effector modules, the latter of which subdivide into the DHp

(dimerization and phospho-accepting histidine) and CA (catalytic) domains. In signal-dependent

manner, SHKs undergo autophosphorylation at a conserved, eponymous histidine residue to enable

subsequent phosphoryl-group transfer to an aspartic acid residue within the response regulator

(RR), i.e., the second TCS component. Once phosphorylated, the RR commonly dimerizes and is

thereby rendered capable of sequence-specific DNA binding and transcription initiation at target

promoters (Gao et al., 2019). As it is the overall levels of phosphorylated RR that govern the

physiological output of the TCS, many SHKs are bifunctional and also exert phosphatase in addition

to their kinase activities (Möglich et al., 2009; Russo and Silhavy, 1993). Exceedingly stringent and

steep regulatory responses can be achieved by finely balancing the opposing RR phosphorylation

and dephosphorylation reactions. Depending on the signal, one or the other of the two elementary

activities prevails, and the SHK consequently acts as either net kinase or net phosphatase. SHKs are

specific for their respective RRs, thus allowing the coexistence in a single cell of multiple TCSs with

minimal crosstalk (Laub and Goulian, 2007).

Among the plethora of known TCSs, certain representatives respond to light as their cognate

stimulus (Möglich, 2019; Ohlendorf and Möglich, 2022). Pertinent TCSs comprise soluble

photoreceptor SHKs that reside in the cytosol and absorb light at specific wavelengths within the

near-ultraviolet (UV) to near-infrared (NIR) range of the electromagnetic spectrum (Krell et al., 2010;

Möglich, 2019). Beyond their pivotal physiological role, these TCSs are of prime interest in

biotechnology as they unlock the control of gene expression by light, an approach belonging to

optogenetics (Deisseroth et al., 2006). As an inducer of gene expression, light brings several benefits

in that it affords superior spatial and temporal precision; can be applied, dosed, and withdrawn with

ease and in automatable fashion; and penetrates living tissue to a wavelength-dependent extent

(Weissleder, 2001). Several light-responsive TCSs have become available and support strong

regulatory responses covering the electromagnetic spectrum from the near-UV to the NIR (Lazar and

Tabor, 2021; Ohlendorf and Möglich, 2022). Apart from little-modified naturally occurring setups

(Tabor et al., 2011), additional photoreceptor SHKs have been engineered by modular recombination

of sensor and effector entities (Levskaya et al., 2005; Möglich et al., 2010, 2009). As a case in point,

the widely deployed pDusk and pDawn plasmids resort to the blue-light-inhibited SHK YF1

originating from the fusion of a light-oxygen-voltage (LOV) photosensor to the DHp/CA effector of

the *Bradyrhizobium japonicum* FixL SHK (Möglich et al., 2009; Ohlendorf et al., 2012). Irrespective

of the wealth of setups for the optical control of bacterial gene expression, there is a dearth of

circuits affording stringent responses to NIR light (Ong et al., 2018).

We previously expanded the pDusk plasmid family by exchanging said LOV photosensor for

photosensory core modules (PCM) of bacterial phytochromes (BphP) (Meier et al., 2024a;

Multamäki et al., 2022). The resultant pREDusk systems respond to red rather than blue light which

facilitates multiplexing (Meier et al., 2024b; Multamäki et al., 2022) with scores of bacterial

optogenetic circuits that mostly sense blue light (Ohlendorf and Möglich, 2022). As an added benefit,

red light penetrates mammalian tissue to a higher extent than blue light owing to reduced scattering

and absorption by cellular constituents, hemoglobin, and other pigments (Ash et al., 2017;

Weissleder, 2001). BphPs react to red and NIR light via biliverdin (BV) chromophores covalently

attached within their PCMs (Davis et al., 1999; Hughes et al., 1997; Takala et al., 2020). The linear

tetrapyrrole BV traverses between the Z and E isomers of its C15=C16 double bond which underpin

the red-absorbing Pr and the far-red-absorbing Pfr states of BphPs, respectively. Red and NIR light

triggers the photochromic Z/E isomerization of BV and thereby drives the photoconversion in the

Pr→Pfr and Pfr→Pr directions, respectively. In conventional BphPs, as in those underpinning the

pREDusk systems, the Pr state is the thermodynamically more stable form prevailing in darkness.

After red-light exposure, these BphPs revert to their Pr resting state thermally within the dark-

recovery reaction which is generally protracted and plays out over many hours to days. Conventional

BphPs therefore effectively act as ratiometric red/NIR-light sensors. In common with many natural

BphP-SHKs (Huber et al., 2024; Karniol and Vierstra, 2003; Lamparter et al., 2002; Multamäki et al.,

2021), the engineered ones in the pREDusk circuits exhibit net kinase activity towards their RR within

their Pr states and net phosphatase activity within their Pfr states. In pREDusk, red light therefore

downregulates target gene expression by several hundred-fold (Meier et al., 2024a; Multamäki et

al., 2022). More recently, we inverted the light response of the underlying BphP-SHKs by changing

the composition of the linkers through which their sensor and catalytic modules transduce signals

(Meier et al., 2024a). Doing so yielded BphP-SHKs with net kinase activity in their Pfr states and net

phosphatase activity in their Pr states, thus supporting pronounced increases of gene expression

under red light in the resultant pDERusk systems.

A second group of bacterial phytochromes, denoted bathy-BphPs, adopt the Pfr rather than

the Pr state in darkness (Karniol and Vierstra, 2003). Consequently, NIR light promotes the Pfr→Pr

conversion, as does visible light, if to a lesser degree. To the extent it has been studied, the thermal

dark recovery in bathy-BphPs is generally fast and complete within seconds to minutes. Due to these

aspects, bathy-BphPs are considered to primarily serve as sensors of light vs. darkness rather than

of a specific light color (Huber et al., 2024).

Informed by spectroscopic analyses, we here harness the PCMs of select bathy-BphPs for

stringently controlling TCS output and bacterial gene expression by NIR light. The resultant plasmid

systems, dubbed pNIRusk, prompt stringent upregulation of gene expression under NIR light. Equally

advantageously and unexpectedly, the engineered pNIRusk circuits show only minimal activation by

red light. To enable multiplexed applications, we also advance light-regulated setups based on

orthogonal TCSs. The pNIRusk plasmids not only afford pronounced NIR-light responses in

Escherichia coli laboratory strains but also in the probiotic Nissle bacteria and in *Agrobacterium*

tumefaciens. Taken together, we deliver versatile implements for the precision control of gene

expression by NIR light in both fundamental and applied research.

Results

Bathy-Bacteriophytochromes as Sensors of Light and pH

To subject bacterial gene expression to NIR light, we considered the PCMs of previously reported

bathy-BphPs as input modules (Rottwinkel et al., 2010; Tasler et al., 2005). To better inform our

choice, we expressed, purified, and spectroscopically characterized BphP PCMs from *Azorhizobium*

caulinodans (*Ac*PCM), *Agrobacterium vitis* (*Av*PCM), and *Pseudomonas aeruginosa* (*Pa*PCM).

Consistent with earlier reports, all three PCMs fully populated the Pfr state in darkness with distinct

Soret- and Q-band absorbance peaks at 415 nm and (755 ± 3) nm, respectively (Fig. 1a-c). Exposure

to NIR light drove the complete conversion to the Pr state characterized by maxima of the Soret band

at (393 ± 2) nm and of the Q band at around (698 ± 3) nm. Illumination with blue and red light

prompted partial population of the Pr state to around 65-80% and 40-70%, respectively. Strikingly,

at pH 8.0, the Q-band intensity of the pure Pr state differed markedly across the PCMs, with *Pa*PCM

showing the strongest signal, and *Ac*PCM the weakest. Based on earlier reports, these data may

principally reflect differences in BV protonation at the pyrrole C ring (Borucki et al., 2005; Zienicke

et al., 2013). To put this notion to the test, we recorded absorbance spectra for the three PCMs at

pH values between 6 and 11 (Fig. 1d-f). While the Q-band intensity remained invariant within the

Pfr state, in the Pr state it increased with decreasing pH. By evaluating the corresponding absorbance

peak heights as a function of pH, we determined apparent pK_a values of 7.2 ± 0.1, 8.0 ± 0.1, and 9.9

± 0.2 for *Ac*PCM, *Av*PCM, and *Pa*PCM, respectively (Fig. 1g). Following previous studies (Zienicke et

al., 2013), we tentatively ascribe these pK_a values to (de-)protonation of the nitrogen atom within

the pyrrole C ring in BV. In addition, pH also affects the photostationary Pr:Pfr ratio under red light

which increases with rising alkalinity (Fig. 1h, Suppl. Fig. 1a-c).

Having ascertained the bathy-character of the candidate PCMs and their Pr⇄Pfr

interconversion, we next assessed the kinetics of reversion to their dark-adapted Pfr state at pH 8.0

following NIR illumination (Fig. 1i). All three PCMs fully recovered to the Pfr state in single-

exponential fashion, but the time constants τ for *Ac*PCM and *Av*PCM of (140 ± 1) s and (80 ± 1) s

were markedly lower than the value of (610 ± 1) s for *Pa*PCM. Closely similar recovery time constants

resulted after exposing the samples to red instead of NIR light (Suppl. Fig. 1d-e). With rising alkalinity,

we observed slower and less complete recovery after NIR-light exposure (Suppl. Fig. 1g-l).

Potentially, this deceleration directly owes to BV deprotonation evidenced in the absorbance

spectra. That is because BV remains protonated in the Pfr state across all investigated pH values, but

within the Pr state it is progressively deprotonated at higher pH values. The return to the Pfr state

hence requires re-protonation at elevated pH values which may slow down the dark recovery.

Subjecting Bacterial Gene Expression to NIR-Light Control

To achieve control of bacterial expression by NIR light, we adapted the previous *Dm*REDusk plasmid

which encodes a red-light-responsive chimeric SHK originating from the fusion of the PCM of

Deinococcus maricopensis BphP (*Dm*BphP) to the DHp/CA fragment of *B. japonicum* FixL (Meier et

al., 2024a). As seen not least in the above spectroscopic experiments, bathy-BphPs also respond to

visible besides NIR light, which stands to cause undesired crosstalk with genetic circuits that react

to, e.g., blue or red light. We reasoned that this adverse effect may be ameliorated by deliberately

rendering the NIR-responsive systems comparatively insensitive to light. Guided by the above

spectroscopic analyses, we selected *Ac*PCM and *Av*PCM as they exhibit a faster Pr→Pfr dark recovery

than *Pa*PCM, which translates into a lower effective light sensitivity at photostationary state

(Ohlendorf and Möglich, 2022; Ziegler and Möglich, 2015).

With *Ds*Red as a fluorescence reporter gene, we replaced *Dm*PCM within *Dm*REDusk by

either *Ac*PCM or *Av*PCM, while maintaining the same register for the linker between the PCM and

DHp/CA moieties (Suppl. Fig. 2). In the resultant plasmids, referred to as pNIRusk, the heme

oxygenase enzyme for intracellular BV provision, the BphP-SHK, and the matching RR FixJ thus form

a constitutively expressed tricistronic operon (Fig. 2a). The initial pNIRusk designs, denoted '0',

showed considerable reporter fluorescence in darkness that only slightly elevated under both red

and NIR light, despite their BphP-SHKs adhering to the same design as in *Dm*REDusk (Fig. 2b).

Spurred by our recent construction of the pDERusk setups (Meier et al., 2024a), we

hypothesized that light responsiveness may be installed by varying the composition of the linker

between the PCM and DHp/CA moieties. That is because this linker is expected to adopt continuous

α-helical coiled-coil conformation and serves as the conduit for conformational signals traveling from

the PCM to the effector module (Diensthuber et al., 2013). Using the PATCHY approach (Ohlendorf

et al., 2016), we generated libraries of the *Ac*NIRusk and *Av*NIRusk setups that differ in the length

and sequence of said linker. In brief, the PATCHY protocol achieves linker variation by randomly

recombining different portions of the linkers deriving from the parental receptors, i.e., the bathy-

BphP PCM and the FixL SHK, respectively (see Suppl. Fig. 3). By assessing the reporter fluorescence

in darkness and under NIR and red light, we identified five light-switchable variants for *Ac*NIRusk and

six for *Av*NIRusk (Fig. 2b, Suppl. Fig. 3). Variants were considered light-switchable if they exhibited at

least a 2-fold difference in reporter fluorescence in darkness vs. illumination with either red or NIR

light. We refer to individual variants by their linker lengths relative to the initial constructs. Although

the light-responsive variants all rely on the same PCMs and DHp/CA modules, they drastically

differed in their response to NIR light, with some elevating and others lowering reporter-gene

expression compared to darkness. Earlier application of PATCHY to SHKs responding to blue and red

light (Meier et al., 2024a; Ohlendorf et al., 2016) had pinpointed sets of light-regulated receptor

variants that exhibited systematic trends in the distribution of the underlying linkers. Specifically,

receptor variants differing in their linker length by multiples of seven residues tended to show similar

responses to light, which could be rationalized by the coiled-coil nature of said linkers (Diensthuber

et al., 2013; Ohlendorf et al., 2016). At present, we only identified a handful of light-responsive SHK

variants, thus largely precluding systematic analyses of the linker properties.

We subsequently focused on variants inducing gene expression under NIR light, albeit with

intriguing variation in their response to red light. For instance, *Ac*NIRusk-1b with a linker one residue

shorter than in *Dm*REDusk showed sizeable induction under red light. By contrast, the variant *Ac*NIRusk-1a with the same linker length but different sequence did much less so. For an eventual application, we deem a reduction of red-light-induced activation advantageous. We thus continued with *Ac*NIRusk-1a and *Av*NIRusk+17a, both of which strongly induce expression under NIR light compared to darkness but only show small responses to red light. For simplicity, we refer to these setups as *Ac*NIRusk and *Av*NIRusk in the following.

Flow cytometry revealed homogeneous single-cell fluorescence distributions for *E. coli* bacteria harboring *Ac*NIRusk or *Av*NIRusk (Fig. 2c). When incubated in darkness, the median fluorescence was only 2-fold above the background fluorescence. Saturating NIR light (800 nm, 250 μW cm^{-2}) induced largely uniform shifts of the entire cell population to several 10-fold higher median fluorescence for *Ac*NIRusk and *Av*NIRusk, respectively, an effect similar in magnitude to the red-light response observed previously for the pDERusk systems (Meier et al., 2024a). At sub-saturating light doses, the single-cell fluorescence of bacteria containing *Ac*NIRusk and *Av*NIRusk uniformly rose to intermediate levels with increasing NIR-light intensity (Suppl. Figs. 4 and 5a-b). Red (100 μW cm^{-2}) and blue (70 μW cm^{-2}) light partially activated *Ac*NIRusk 11-fold and 35-fold, respectively (Fig. 2c).

At the ensemble level, the average fluorescence per cell density ramped up sigmoidally with NIR light by up to around 70- and 150-fold for *Ac*NIRusk and *Av*NIRusk, respectively (Fig. 2d-e). The two systems exhibited half-maximal activation at relatively high NIR-light intensities of (29 ± 1) μW cm^{-2} and (43 ± 1) μW cm^{-2}, i.e., about an order of magnitude above the half-maximal red-light doses previously determined for induction of the pREDusk and pDERusk systems (Meier et al., 2024a; Multamäki et al., 2022). Arguably, these differences reflect at least in part the respective dark-recovery times of the underlying bathy-BphPs which are in the seconds to minutes range rather than in the hours range for conventional BphPs. Bacterial cell growth was unaffected by NIR light (Suppl. Fig. 5c-e).

Red light also induced *Ac*NIRusk and *Av*NIRusk but only around 2- and 6-fold relative to

darkness (Fig. 2d-e). Blue light elicited a stronger 16- and 25-fold induction, respectively, at the

highest probed light intensity of around 120 μW cm^{-2}. Higher blue-light intensities were not accessed

owing to instrumental limitations and onsetting phototoxicity. From the data, we determined half-

maximal blue-light doses of (46 ± 3) μW cm^{-2} and (100 ± 20) μW cm^{-2} for *Ac*NIRusk and *Av*NIRusk,

respectively. Given the around 1.7-fold higher energy of blue compared to NIR light, the circuits

therefore trigger at similar half-maximal photon fluencies for either light color. At least qualitatively,

the degree of induction by the individual light colors (red, blue, NIR) follows the same trend as the

photochemical responses evidenced in the underlying *Ac*PCM and *Av*PCM (see Fig. 1a-c) where NIR

light elicited the most pronounced Pr formation, followed by blue and red light.

Engineering Orthogonal Light-responsive Two-Component Systems

In common with earlier members of the pDusk pedigree, the pNIRusk systems are based on the FixLJ

TCS from *B. japonicum* (Anthamatten and Hennecke, 1991). Combinations of several of these setups

would hence experience crosstalk at the level of RR phosphorylation which, in all but a few cases, is

undesired. To overcome this limitation, we set out to engineer light-responsive TCSs orthogonal to

those used in pDusk, pNIRusk, and related setups. As recently surveyed, several TCSs have been

characterized to the point that they have found application in synthetic biology (Lazar and Tabor,

2021). Based on their reported stringency of regulation, we selected as candidate TCSs TtrSR from

Shewanella baltica that responds to tetrathionate and TodST from *Pseudomonas putida* that can

detect toluene compounds (Daeffler et al., 2017; Lau et al., 1997).

Starting from *Dm*REDusk, we substituted the original FixL DHp/CA domains for those from

TtrS, while also exchanging FixJ for the TtrR RR and putting the expression of a *Ds*Red reporter under

control of the Pttr promoter that is subject to regulation by TtrSR (Fig. 3a). Within this framework,

we then varied the linker interconnecting the *Dm*PCM and the TtrS DHp/CA module via PATCHY

(Ohlendorf et al., 2016). Screening of the resultant linker library for differential *Ds*Red expression

under darkness, and NIR and red light identified several variants, termed *Dm*Ttr, with substantial

light responses (Fig. 3a, Suppl. Fig. 6a). Notably, the overall expression levels within the identified

*Dm*Ttr systems were markedly higher than those in the pREDusk and pNIRusk setups.

Next, we extended the strategy to *Av*NIRusk by introducing TodST. Of note, TodS is a complex

SHK with integrated phosphorelay that comprises in sequential order PAS-1, DHp-1, CA-1, REC, PAS-

2, DHp-2, and CA-2 domains (Busch et al., 2009). Toluene signals, detected by PAS-1, modulate the

activity of the DHp-1/CA-1 unit which phosphorylates the REC domain that in turn governs the

activity of the DHp-2/CA-2 module and eventual phosphorylation of the TodT RR. Earlier reports

showed that a miniaturized TodS variant only comprising the PAS-1, DHp-2, and CA-2 domains,

dubbed Min-TodS, retains toluene sensitivity and stringent regulation of TodT (Silva-Jiménez et al.,

2012). Consequently, we replaced the FixLJ TCS in *Av*NIRusk by the TodS DHp-2/CA-2 fragment, while

also introducing the TodT RR and its cognate PtodX promoter (Fig. 3b). Linker variation by PATCHY

and screening for *Ds*Red reporter fluorescence pinpointed several variants, termed *Av*Tod, which

either elevated or lowered gene expression in response to NIR light (Fig. 3b, Suppl. Fig. 6b).

For further analyses, we selected the variants *Dm*Ttr+7b, *Av*Tod+16, and *Av*Tod+21 with PCM-

DHp/CA linkers 7, 16, and 21 residues, respectively, longer than that in the parental *Dm*REDusk

circuit. Exposed to red light, *Dm*Ttr+7b elicited an about 15-fold decrease of gene expression

compared to darkness (Suppl. Fig. 7a). Whereas *Av*Tod+16 supported a 20-fold reporter-

fluorescence decrease under NIR light, *Av*Tod+21 promoted an 8-fold increase of gene expression

(Suppl. Fig. 7b-c). Although the attained regulatory efficiencies pale in comparison to those of

*Dm*REDusk (Meier et al., 2024a) and *Av*NIRusk, the data show that the principal design and screening

strategies readily translate to other TCSs.

To allow multiplexing of several light-responsive circuits in a single bacterial cell, we modified

the *Dm*DERusk and *Av*NIRusk plasmids by replacing their kanamycin resistance marker, their ColE1

origin of replication (ori), and their *Ds*Red reporter gene by a streptomycin marker, the CloDF13 ori,

and the yellow-fluorescent YPet reporter, respectively. The resultant *Dm*DERusk-Str-YPet and

*Av*NIRusk-Str-YPet circuits showed similar responses to red and NIR light as the parental plasmids

they derive from (Suppl. Fig. 7d-e). Pairs of compatible plasmids with different TCSs, oris, resistance

markers, and fluorescent reporters were transformed into bacteria, followed by cultivation in

darkness, under red (100 μW cm^{-2}) or NIR light (200 μW cm^{-2}), or under both light colors. By

employing individual plasmid combinations, the expression of the *Ds*Red and YPet reporters could

be individually controlled by the two light colors (Fig. 3c, Suppl. Fig. 7f). For instance, when

combining *Dm*DERusk-Str-YPet with *Av*Tod+21-*Ds*Red, YPet and *Ds*Red expression rose under red and

NIR light, respectively, whereas the joint application of both light colors only induced the YPet

reporter gene. Taken together, well-chosen plasmid sets enable the orthogonal expression control

of two target genes by red and NIR light. Notably, the plasmids can be deployed as is with no further

modifications required.

Although not probed here, the overall performance and relative response in multiplexed

applications may be optimized by variation of the copy number and expression strength of the

individual plasmids. Likewise, rather than using saturating light intensities as in the above

experiments, yet finer-grained expression control could be exerted by adjusting the red- and NIR-

light intensities. In a similar vein, different wavelengths, currently determined by the available

instrumental setups, may be chosen to better separate the individual light responses of two (or,

several) multiplexed systems.

Applications in the Laboratory and Beyond

As TCSs widely recur across bacterial phyla, they often retain functionality upon transfer between

different bacterial strains. Earlier reports not least demonstrated that representatives of the pDusk

family support light-regulated gene expression across several prokaryotes (Ohlendorf and Möglich,

2022). For instance, pDusk and its derivative pDawn (Ohlendorf et al., 2012) have been successfully

deployed in *E. coli* Nissle 1917 (*Ec*N) and other probiotic bacteria (Cui et al., 2021; Magaraci et al.,

2014; Pan et al., 2021; Sankaran and del Campo, 2019; Yang et al., 2020). These experiments augur

the application of light-responsive bacteria in the gut of mammals as programmable, living

therapeutics (Ohlendorf and Möglich, 2022). To date, however, pertinent applications are limited by

the shallow tissue penetration of blue light required for triggering pDusk and pDawn. Although

upconverting nanoparticles can be administered jointly with the light-responsive bacteria to allow

actuation by NIR light, this approach introduces complexity and departs from a fully self-contained,

optogenetic strategy (Yang et al., 2020). To assess whether the presently generated light-responsive

gene-expression systems may overcome this potential limitation, we introduced *Ac*NIRusk and

*Av*NIRusk into *Ec*N. Both setups mediated a more than 100-fold upregulation of reporter-gene

expression under NIR light with half-maximal doses of (27 ± 1) and (28 ± 2) $\mu W\ cm^{-2}$, respectively

(Fig. 4a, Suppl. Fig. 8). Of advantage, the response to blue and NIR light was even lower than

observed above for the *E. coli* laboratory strain.

We next probed the performance of *Av*NIRusk and *Dm*REDusk in the biotechnologically

relevant *Agrobacterium tumefaciens*. To this end, we constructed derivative plasmids, denoted

*Av*NIRlux and *Dm*REDlux, that subject the expression of a luciferase operon under the control of the

light-responsive TCSs (Fig. 4b). After conjugational transfection into *A. tumefaciens*, the light

response was assessed by spotting the bacteria on agar plates (Fig. 4c). As visualized by

luminescence, the *Av*NIRlux system strongly induced expression upon incubation under NIR light

compared to darkness. Different from *E. coli* (see Fig. 2e, Suppl. Fig. 9a), red light activated to similar

extents as NIR light, whereas blue light induced expression only weakly. In the case of *Dm*REDlux,

the expression of the *lux* reporter was strongly reduced under both blue and red light, but not under

NIR light or in darkness. Although *A. tumefaciens* harbors two endogenous BphPs (Karniol and

Vierstra, 2003), our experiments did not detect any differences in growth or phenotype between

cultivation under NIR and red light. We next characterized the response of *A. tumefaciens* cultured

in liquid medium. Once reaching the stationary phase after around 20 h (Suppl. Fig. 9b), the cultures

bearing *Av*NIRlux exhibited a 7-fold activation of *lux* expression by NIR light compared to darkness.

For *Dm*REDlux, the *lux* expression was around 300-fold higher in darkness than under red light (Fig.

4d, Suppl. Fig. 9c-d).

Discussion

Bathy-Bacteriophytochromes as Tunable Sensors of Light Intensity and Quality

As conventional bacteriophytochromes do, bathy-BphPs transition between their Pr and Pfr states

with 15*Z*- and 15*E*-configured biliverdin, respectively. In marked contrast to the conventional

representatives, bathy-BphPs assume their 15*E*-configured Pfr state as the thermodynamically most

stable form in darkness, rather than their Pr state. Despite this principal difference, the overall PCM

architecture and the residues lining the bilin binding pocket are remarkably conserved across known

BphPs. Notwithstanding limited successes in reprogramming certain BphPs (Böhm et al., 2025; Xu et

al., 2024), the molecular determinants underlying bathy-character remain elusive and their design

demanding.

Owing to their Pfr dark-adapted state, bathy-BphPs undergo net Pfr→Pr conversion when

irradiated with light within the near-UV to NIR region of the electromagnetic spectrum. Put another

way, there is no light quality that would fully populate the dark-adapted Pfr state. Bathy-BphPs

surmount this challenge by a fast thermal Pr→Pfr recovery that restores the Pfr state within seconds

to minutes. The primary physiological role of bathy-BphPs may hence be that of sensors of light

quantity rather than quality (i.e., color) (Huber et al., 2024).

Intriguingly, we uncovered a profound pH influence on photoreception in bathy-BphPs. With

rising alkalinity, the absorption cross-section of the Q band strongly decreases within the Pr state

but remains unchanged for Pfr. Based on earlier studies (Borucki et al., 2005; Velazquez Escobar et

al., 2015; Zienicke et al., 2013), we attribute these spectral changes to (de-)protonation of the

nitrogen atom of the BV pyrrole C within the Pr state. The pertinent pK_a values range between 7-8

for *Ac*PCM and *Av*PCM, similar to the value of 7.6 for the bathy-BphP Agp2 from *A. tumefaciens*

(Zienicke et al., 2013), but substantially lower than the value of about 10 for *Pa*PCM. Inspection of

the experimentally determined *Pa*PCM and Agp2PCM structures in their Pfr states (Schmidt et al.,

2018; Yang et al., 2008) and the corresponding Pfr models for *Ac*PCM and *Av*PCM (Abramson et al.,

2024) reveals highly conserved chromophore-binding pockets with most residues adjacent to BV

identical in all four proteins (Suppl. Fig. 10). However, *Pa*PCM features tyrosine and serine residues

at positions 190 and 275 rather than phenylalanine and alanine, respectively, in the equivalent

places of the other three PCMs. Although speculative at present, these systematic differences may

render the *Pa*PCM binding pocket comparatively hydrophilic and thereby account for the divergent

protonation equilibrium of the pyrrole C ring.

BV deprotonation within the Pr state at elevated pH incurs at least three principal effects that

under constant light all promote population of the Pr state at the cost of the Pfr state. First, the

Pr→Pfr dark recovery slows down between pH 7 and 10 for all three investigated PCMs. At least in

part, this may reflect the need for BV re-protonation upon return from the Pr to the Pfr state which

remains fully protonated at all investigated pH values. Second, as noted above, BV deprotonation

leads to a drastic reduction of the Q-band absorbance within the Pr state (see Fig. 1). For a given

intensity of visible light, photoconversion to the Pfr state thus becomes less effective, thereby again

favoring the Pr state at photostationary state. Third, BV deprotonation also goes along with a

dramatic decrease in the apparent photoconversion quantum yields in the Pr→Pfr vs. the Pfr→Pr

directions (Suppl. Fig. 1f). As best exemplified for *Ac*PCM, at pH 6.5, the respective quantum yields

for the photoreversible Pr⇌Pfr interconversion are about equal, but at pH 10.5, the quantum yield

in the Pr→Pfr direction is less than 1% that in the Pfr→Pr direction. Together with the pH-dependent

reduction in the Pr-state Q-band absorbance, this mechanism leads to vastly different red-light-

induced photostationary states as a function of pH (see Fig. 1h). Whereas at low pH, saturating red

light results in about 60-70% Pfr population, at high pH values almost no Pfr state remains (Fig. 5).

As a direct consequence, at pH 10.5, red light counter-intuitively drives the complete conversion to

the Pr state of *Ac*PCM. Put another way, in its deprotonated form this bathy-BphP becomes

essentially colorblind and acts as a pure light-intensity sensor. Although less pronounced, *Av*PCM

and *Pa*PCM follow the same trend with pH.

While it is debatable whether the rhizobia *Ac*PCM and *Av*PCM derive from ever encounter

such extreme pH swings, it has been proposed that the resultant changes in photochemical

properties could serve a sensing function (Zienicke et al., 2013). Even independent of this aspect,

our present analyses show how pH impacts bathy-BphP photoreception at multiple levels. While it

remains true that all light colors net promote the Pfr→Pr transition, they do so to different extents

which can have a profound impact on downstream signal transduction as discussed below. We

propose that modulation of the BV protonation midpoint has been harnessed during evolution to

sculpt the light response of extant bathy-BphPs according to the individual needs of the respective

organism of origin. In line with this hypothesis, we presently evidence a considerable span of pK_a

values for BV protonation across several bathy-BphP PCMs.

Engineering of NIR-Light-Responsive Proteins

As best exemplified by sensor histidine kinases, PCMs of bathy-BphPs can occur in conjunction with

the same type of effector as those of conventional BphPs (Blum et al., 2025). However, it has been

unclear whether a single, specific output protein can be regulated in its activity by both PCM types.

The successful recombination of bathy-BphP PCMs with the same DHp/CA module earlier controlled

by conventional BphP PCMs (Meier et al., 2024a; Multamäki et al., 2022) now demonstrates the

principal validity of this idea. At least to some extent, the PCMs of conventional and bathy-BphPs are

hence interchangeable, a finding that indicates shared or compatible signal-transduction

mechanisms, as also suggested by the conserved structure and photochemistry across the BphP

class (Möglich, 2019; Takala et al., 2020). As one important ramification, numerous previously

engineered red-light-responsive receptors may thus be subjugated to NIR-light control by swapping

out their underlying PCMs from conventional BphPs for ones from bathy-BphPs (Chernov et al.,

2017; Etzl et al., 2018; Leopold et al., 2022, 2019; Levskaya et al., 2005; Schmidl et al., 2014; Stabel

et al., 2019; Stüven et al., 2019; Tang et al., 2021). Compared to the challenging approach of

imparting bathy-character via limited PCM modifications (Böhm et al., 2025; Xu et al., 2024), the

design task thus considerably simplifies.

We identify several caveats that likely hold for future design endeavors, too. Notably, the

connection between PCM and effector of choice needs to be conducive to transmitting light signals.

When at present employing the same linker register that supports red-light-inhibited gene

expression in *Dm*REDusk, we initially derived *Ac*PCM- and *Av*PCM-based systems giving rise to

constitutive target gene expression without pronounced light responses. Adjusting the length and

sequence of the linker intervening the PCM and the DHp/CA domains then yielded variants with

robust gene-expression responses to NIR light. Based on structural and biochemical evidence, said

linker is expected to form a continuous parallel α-helical coiled coil within the homodimeric SHK

(Bódizs et al., 2024; Burgie et al., 2024; Diensthuber et al., 2013; Jacob-Dubuisson et al., 2018; Malla

et al., 2024; Möglich, 2019; Möglich et al., 2009). Although other BphP-based photoreceptor designs

may involve linkers adopting different structures, the variation and probing of length and sequence

variants generally represents a worthwhile strategy that may eventually bear fruit. As, depending on

the use case, sizeable variant libraries may have to be assessed, two principal aspects emerge that

benefit the implementation of this strategy. First, the construction of libraries covering the desired

diversity in linker composition should be efficient, which can, e.g., be achieved by resorting to the

PATCHY protocol (Ohlendorf et al., 2016) as done here. Second, the screening for light-regulated

function should afford sufficient throughput, e.g., by coupling photoreceptor function to a

fluorescent readout or cell survival.

Tunability of NIR-Light-Activated Two-Component Systems

The recombination of bathy-BphP PCMs with a SHK effector module led to the pNIRusk plasmids for

stringent activation by NIR light of bacterial gene expression. As the underlying PCMs absorb light

over the entire near-UV to NIR region of the electromagnetic spectrum, the pNIRusk systems are

expected to also activate when exposed to other light colors. To dampen the commonly undesirable

activation by visible light and thereby facilitate the multiplexing with other photoresponsive circuits,

we deliberately chose *Ac*PCM and *Av*PCM because of their comparatively fast Pr→Pfr thermal

recovery. At photostationary state under constant illumination, a faster recovery results in an

effective desensitization against light, be it of blue, red, or NIR color (Ziegler and Möglich, 2015).

Consistent with this design rationale, the *Ac*NIRusk and *Av*NIRusk setups indeed exhibit half-maximal

light doses around ten times higher than for, e.g., *Dm*DERusk which relies on a conventional BphP

PCM (Meier et al., 2024a). Given its low phototoxicity, NIR light may be applied at even high powers

without harming the biological system under study. We note that NIR light around 800 nm is

commonly used in a therapeutic setting for photobiomodulation at intensities exceeding the ones

presently used by up to several orders of magnitude (Hamblin, 2016).

Perplexingly, the *Ac*NIRusk and *Av*NIRusk circuits respond moderately or very weakly to blue

and red light, respectively (see Fig. 2d, e). Although the strength of the gene-expression response

monotonically increases with the extent of Pfr→Pr photoconversion elicited by the individual light

colors within the underlying PCMs (see Fig. 1a-c), there is no direct proportionality. For instance,

*Ac*PCM roughly assumes a 0.7:0.3 photostationary Pr:Pfr equilibrium under red light (see Fig. 1a),

but at the level of gene expression, the response of the *Ac*NIRusk circuit to red light is only 2% that

of the NIR-light response. Initially unexpected, this baffling effect therefore further benefits the

application of the pNIRusk circuits in combination with other light-responsive entities, be it optogenetic circuits or fluorescent reporters (Meier et al., 2024b; Tabor et al., 2011).

Our work provides pointers to the potential molecular underpinnings in that we isolated additional pNIRusk variants with similar NIR-light responses but substantially higher expression under red light (see Fig. 2b). Notably, these variants only differ in the composition of their linkers connecting PCM to effector. Informed by our recent investigation of linker variants that culminated in the generation of *Dm*DERusk (Meier et al., 2024a), we hypothesize that the different red-light responses find their origin in an altered balance between the counteracting elementary histidine kinase and phosphatase activities of SHKs. At the ensemble level, even small perturbations, e.g., caused by linker variation or changes in the Pr:Pfr proportion, can utterly tilt the net activity in either the kinase or phosphatase direction (Landry et al., 2018; Möglich, 2019; Möglich et al., 2009; Russo and Silhavy, 1993). The histidine kinase-phosphatase duality of TCSs thus provides a means of rendering the signal response to light highly malleable and adaptable. At a practical level, when engineering light-responsive TCSs, we advise to screen multiple candidate variants that may well differ substantially in their signal response. As noted previously (Meier et al., 2024a), similar concepts may also operate for cyclases (GGDEF) and phosphodiesterases (EAL) enzymes that frequently occur in conjunction with BphP PCMs (Gourinchas et al., 2017) and that antagonistically catalyze the making and breaking, respectively, of the bacterial second messenger cyclic-di-guanosine monophosphate (Jenal et al., 2017).

Applications in the NIR Future

The advent of stringently controlled NIR-responsive bacterial gene-expression platforms now facilitates a number of innovative applications. First, the currently available optogenetic setups for expression regulation in bacteria predominantly respond to blue and red light but are often blind to NIR light (Ohlendorf and Möglich, 2022). Given their fully intended relative insensitivity and their

unexpectedly weak response to visible light, the pNIRusk systems lend themselves to combinations

with pertinent setups. For example, applied at low to moderate intensity, blue light would trigger

one setup, e.g., pAurora2 or EL222-based systems (Jayaraman et al., 2016; Ranzani et al., 2024), but

leave pNIRusk largely untouched. By contrast, NIR light would selectively activate pNIRusk as the

LOV receptors underlying the blue-light-responsive setups are insensitive to this light quality. To

facilitate multiplexed applications of pNIRusk, we supply derivatives reliant on alternative TCSs that

can be readily combined in single bacterial cells and mediate the expression of several target genes

orchestrated by two light colors.

Second, NIR light penetrates mammalian tissue much more deeply than light of shorter

wavelengths (Lehtinen et al., 2022; Weissleder, 2001), owing to reduced absorption and scattering

which are, e.g., 5-10 lower for near-infrared light than for blue light. The exponential attenuation of

light intensity with tissue depth is therefore much less pronounced for NIR light than it is for UV or

visible light. Given the low phototoxicity of NIR light and its established use in photobiomodulation,

noted above, the pNIRusk setups stand to be activatable through substantial tissue layers of at least

several millimeters. Although the pNIRusk setups are (deliberately) less light-sensitive than, e.g., the

earlier pREDusk and pDERusk systems, they can be nonetheless actuated by light fluences in the

range of several tens of μW per cm^2 which is orders of magnitudes less than the light intensities

commonly used in neuronal optogenetics. These aspects open up intriguing applications involving

programmable, light-responsive bacteria as agents inside the bodies of mammals, e.g., as living

therapeutics (Cubillos-Ruiz et al., 2021; Hartsough et al., 2020). The pDawn system has already been

applied in this capacity within probiotic bacteria, but the approach has been hampered by delivering

blue light *in situ* (Cui et al., 2021; Magaraci et al., 2014; Pan et al., 2021; Sankaran and del Campo,

2019; Yang et al., 2020). One remedy has been the use of upconverting nanoparticles (UCNP)

administered alongside the light-responsive bacteria (Pan et al., 2021; Yang et al., 2020). Notably,

the UCNPs can be charged by absorption of several photons in the NIR range to then emit a photon

in the blue spectral region. Though feasible, pertinent applications introduce complexity and depart

from a purely genetically encoded optogenetic strategy. The new pNIRusk setups directly address

this challenge as they can be activated by wavelengths closely similar to those previously used for

charging the UCNPs but at around 10-fold lower light intensity. Towards this goal, we demonstrate

at present that both *Ac*NIRusk and *Av*NIRusk operate in the probiotic *E. coli* Nissle strain out of the

box, without need for optimization. Used in place of pDawn, the pNIRusk systems thus obviate the

need for the UCNPs and thereby broaden the application scope of living therapeutics considerably.

Third, we demonstrate functionality of pNIRusk and pREDusk in the biotechnologically

relevant *A. tumefaciens* (Fig. 4c-d) which is widely used as a transfection vehicle in plant

biotechnology (Liu et al., 2013). Prospectively, both pNIRusk and pREDusk may achieve

spatiotemporal control of *A. tumefaciens* infectivity, e.g., to confine plant transfection to specific

time windows or organs. Of particular advantage, the bathochromically shifted absorbance spectra

of BphPs compared to plant Phys (Hughes and Winkler, 2024; Yi et al., 2025) stand to enable NIR-

light control without interfering with plant photoreception and photoautotrophic growth.

Methods

Molecular Biology

Oligonucleotides and plasmids used in this study are listed in Suppl. Tables 1 and 2. All plasmid

constructs were confirmed by Sanger sequencing (Microsynth Seqlab, Göttingen, Germany). To

produce bathy-BphP PCMs, the respective genes were introduced into a pCDF-Duet plasmid

(Novagen) already containing heme oxygenase 1 (HO1) from *Synechocystis* sp. for biliverdin supply

(Mukougawa et al., 2006; Stüven et al., 2019). To this end, the genes encoding the PCMs from *A.*

caulinodans (Uniprot A8HU76, residues 1-501), *A. vitis* (Uniprot B9JR96, residues 1-499), and *P.*

aeruginosa (Uniprot Q9HWR3, residues 1-497) were amplified by PCR from the template plasmids

pQX091, pQX086, and pQX037, respectively (Xu et al., 2024), followed by Gibson assembly (Gibson

et al., 2009).

The initial *Ac*NIRusk and *Av*NIRusk constructs (denoted '0') were created by exchanging

*Dm*PCM in *Dm*REDusk (Meier et al., 2024a) for *Ac*PCM and *Av*PCM, respectively (Suppl. Fig. 2), via

PCR amplification and Gibson assembly. The second amino acid in *Ac*PCM and *Av*PCM was changed

from proline to threonine to maintain the polycistronic structure of the operon comprising HO1, the

BphP-SHK, and the RR FixJ. To apply the PATCHY strategy (Ohlendorf et al., 2016), start constructs

were generated by elongating the PCM-DHp/CA linkers in the initial *Ac*NIRusk and *Av*NIRusk plasmids

as reported before (Meier et al., 2024a). The resulting constructs thus comprise the N-terminal

portions of the parental BphPs from A. *caulinodans* (Uniprot A8HU76) and A. *vitis* (Uniprot B9JR96)

up to and including residues 520 and 518, respectively, connected to the C-terminal fragment of *B.*

japonicum FixL starting from residue 257 (Uniprot P23222). At the junction, a restriction site for

*Ksp*AI and a one-nucleotide frameshift were inserted (Suppl. Fig. 3). Cloning was performed by two

consecutive overhang-PCRs and blunt-end ligation.

The TtrSR TCS from *Shewanella baltica* was introduced into *Dm*REDusk (Meier et al., 2024a)

by first replacing the FixK2 promoter by the 85-bp minimal promoter sequence PttrB185-269

(Daeffler et al., 2017) via PCR amplification and restriction cloning using *Bgl*II and *Xba*I sites. Next,

the *B. japonicum* FixLJ TCS was exchanged for the *S. baltica* TtrSR (Genbank CP000891.1, loci

Sbal195_3858 and Sbal195_3859) via Gibson assembly using as a PCR template a synthetic gene

(Gene Art) with codons optimized for expression in *E. coli*. For conducting PATCHY, a start construct

was made that includes residues 1-539 of the *Dm*BphP (Uniprot E8U3T3) and the *S. baltica* TtrSR

TCS (Uniprot A9L3S5) starting at residue 407. Cloning was performed as described above except that

an *Eco*32I restriction site and a two-nucleotide frameshift were placed at the junction between the

two gene fragments (Suppl. Fig. 6a).

The cognate TodX promoter of the TodST TCS from *Pseudomonas putida* was amplified by

PCR from pMIR77 (Ramos-González et al., 2002) and inserted into pDusk (Ohlendorf et al., 2012) by

restriction cloning with *Bgl*II and *Xba*I. Next, the TodST TCS was amplified from pMIR66 (Ramos-

González et al., 2002) and fused to the *Bacillus subtilis* YtvA LOV domain within pDusk via Gibson

cloning. The *Av*Tod-NIRusk PATCHY start construct was obtained by replacing YtvA-LOV with HO1 and

and *Av*PCM (residues 1-518) via Gibson assembly. The resultant construct comprised the TodS

starting from residue 726 (Uniprot E0X9C7) and TodT. At the junction, a *KspA*I restriction site and a

one-nucleotide frameshift were inserted (Suppl. Fig. 6b).

For multiplexing experiments, the *Ds*Red Express2 reporter gene (Strack et al., 2008) in

*Dm*DERusk and *Av*NIRusk was replaced by the YPet reporter gene (Nguyen and Daugherty, 2005).

Additionally, the kanamycin (Kan) resistance marker and the ColE1 origin of replication (ori) were

exchanged for streptomycin (Str) and a CloDF13 ori via Gibson assembly. A version of *Av*NIRusk

containing a multiple-cloning site rather than a fluorescent reporter was cloned by Gibson assembly

and deposited at Addgene for distribution (accession number 235084).

To apply *Dm*REDusk and *Av*NIRusk in *Agrobacterium tumefaciens*, the *Photorhabdus*

luminescence luxABCDE operon from pBAM2-luxAE (Dziuba et al., 2021) was cloned into the miniTn5

delivery plasmid pBAM2 (Uebe, unpublished). The sequence was amplified via PCR, digested with

*Nde*I/*Sac*I, and ligated into a similarly digested and Fast-AP-dephosphorylated pBAM2 vector to

generate pBAM2-luxAE. Subsequently, the light-regulated TCS cassettes of *Dm*REDusk (*DrhemO-

Dmpcm-BjfixL*) and *Av*NIRusk (*DrhemO-Avpcm-BjfixL*) were amplified. These genes were set under

the control of the strong constitutive *M. gryphiswaldense* PmamDC promoter. The *B. japonicum*

FixK2 promoter was also amplified by PCR. The amplified fragments were fused by overlap extension

PCR and digested with *Nde*I/*Not*I as was the pBAM2-luxAE. Ligation resulted in the reporter plasmids

*Dm*REDlux and *Av*NIRlux in which the *luxABCDE* operon is under the control of the FixK2 promoter.

Protein Purification

For production of *Ac*PCM, *Av*PCM, and *Pa*PCM, *E. coli* LOBSTR cells (Andersen et al., 2013) were transformed with the respective pCDF-Duet-PCM/HO1 constructs. Gene expression and protein purification were performed as before (Meier et al., 2024a). Briefly, two flasks with 800 mL lysogeny broth (LB) medium each supplemented with 100 µg mL^{-1} Str were inoculated with a 5-mL overnight culture, and bacteria were grown in baffled Erlenmeyer flasks at 37°C and 225 rpm agitation. At an optical density at 600 nm (OD$_{600}$) of around 0.7, the expression was induced by 1 mM isopropyl-β-thiogalactopyranoside (IPTG), and 0.5 mM δ-aminolevulinic acid was added to support biliverdin incorporation. The expression was performed at 16°C and 225 rpm agitation in darkness for around 65 h. All following steps were performed in green safe light. After harvesting the cells by centrifugation, they were resuspended in lysis buffer [50 mM tris(hydroxy-methyl)-amino-methane (Tris)/HCl, 20 mM NaCl, 10 mM imidazole, 1 mM β-mercaptoethanol, pH 8.0, protease inhibitor completeOne (Roche), 1 mg mL^{-1} lysozyme]. Cells were lysed by sonification, and the supernatant of the centrifuged lysate was applied to a HiTrap TALON crude column loaded with cobalt-nitrilotriacetic resin (Cytiva). The column was washed with buffer [50 mM Tris/HCl, 20 mM NaCl, 1 mM β-mercaptoethanol, pH 8.0], and the His-tagged proteins were eluted by an imidazole gradient from 0 to 0.5 M. Fractions were analyzed by denaturing SDS polyacrylamide gel electrophoresis (PAGE) in the presence of 1 mM Zn^{2+} to visualize biliverdin incorporation (Berkelman and Lagarias, 1986) and pooled based on protein content. During an overnight dialysis at 4°C into buffer [50 mM Tris/HCl, 20 mM NaCl, 1 mM β-mercaptoethanol, pH 8.0], the His-tag was cleaved off by His$_6$-tagged TEV protease. The sample was applied twice to a gravity-flow IMAC column with HisPur cobalt resin (ThermoFisher) using the same buffers as before. Cleavage of the His-tag and protein purity were analyzed by Zn-SDS-PAGE. Fractions were pooled, dialyzed overnight at 4°C against storage buffer

[20 mM Tris/HCl, 20 mM NaCl, 10% (w/v) glycerol, 1 mM dithiothreitol, pH 8.0], concentrated via

spin filtration, and stored at -80°C.

UV/vis Absorption Spectroscopy

Absorption spectroscopy was carried out on a Cary60 UV/vis spectrophotometer (Agilent

Technologies) at 25°C under green safe light. Protein samples were diluted into buffers with pH

ranging from 6.0 to 11.0 (30 mM 2-(N-morpholino)ethanesulfonic acid/HCl, 30 mM NaCl, pH

6.0/6.5/7.0; 30 mM Tris/HCl, 30 mM NaCl, pH 7.5/8.0/8.5/9.0; 30 mM glycine/HCl, 30 mM NaCl, pH

10.0/10.5/11.0] (Rumfeldt et al., 2019). Samples were illuminated for 120 s with LEDs emitting in

the NIR [(800 ± 18) nm], red [(650 ± 10) nm] and blue [(470 ± 5) nm]. Data were normalized by the

Soret absorbance at 403 nm. Pr→Pfr dark recovery kinetics were monitored every 5 s at 760 nm and

fitted to single-exponential functions using the Fit-o-mat software (Möglich, 2018). Pr:Pfr ratios after

red or blue illumination were calculated via linear combination of the basis spectra as described

previously (Yi et al., 2025). The variation of photochemical parameters with pH was evaluated by

fitting to the Henderson-Hasselbalch equation. The ratio of the quantum yields Φ_p and Φ_q for the

red-light-driven photoconversion in the Pr→Pfr and Pfr→Pr directions, respectively (see Suppl. Fig.

1f), can be determined as follows (Yi et al., 2025):

$$\Phi_q/\Phi_p = P_R/P_{FR} \times (1 - 10^{-A_{Pr}})/(1 - 10^{-A_{Pfr}})$$

P_R and P_{FR} are the Pr and Pfr proportions at red-light-induced photostationary state, and A_{Pr} and A_{Pfr}

denote the absorbance in the Pr and Pfr states at the peak emission wavelength of the red-light

source.

Generation and Screening of Linker Variants

*Ac*NIRusk, *Av*NIRusk, *Dm*Ttr-REDusk and *Av*Tod-NIRusk linker variants were created with the PATCHY

strategy (Meier et al., 2024a; Ohlendorf et al., 2016). Sets of forward and reverse primers to generate

linker variants were designed using a custom Python script (https://github.com/vrylr/PATCHY). In brief, PCR amplification was carried out using the primer sets in a final concentration of 0.25 µM. After purifying the PCR product by gel extraction, the start constructs were digested with *Ksp*AI or *Eco*32I (Suppl. Fig. 3, 6), followed by phosphorylation with T4 polynucleotide kinase and ligation with T4 DNA ligase. The reaction mixes were transformed into DH10b cells and plated on LB plates supplemented with kanamycin (LB/Kan). Plates were incubated overnight at 37°C in darkness, or under red light [100 µW cm^{-2}, (650 ± 11) nm] or NIR light [200 µW cm^{-2}, (800 ± 18) nm]. Applied light intensities were calibrated with a power meter (model 842-PE, equipped with a 918D-UV-OD3 silicon photodiode, Newport, Darmstadt, Germany). Bacterial colonies developing red color upon overnight incubation, indicative of *Ds*Red production, were used to inoculate 200 µL LB/Kan in a clear 96-well microtiter plate (MTP) (Nunc, ThermoFisher). Bacteria were incubated for 21-24 h at 30°C and 750 rpm agitation. Next, the cultures were diluted 100-fold into fresh LB/Kan medium and split into three separate cultures which were then incubated at 37°C and 750 rpm agitation in either darkness, red light [100 µW cm^{-2}, (650 ± 11) nm], or NIR light [200 µW cm^{-2}, (800 ± 18) nm]. After 18 h, the optical density at (600 ± 9) nm (OD_{600}) and the *Ds*Red fluorescence [(554 ± 9) nm excitation and (591 ± 20) nm emission] were measured with a Tecan Infinite M200 Pro MTP reader (Tecan Group, Ltd., Männedorf, Switzerland). The *Ds*Red fluorescence was normalized by OD_{600} and represent the mean ± s.d. of three biologically independent replicates.

Light-Dose-Response Assays in *Escherichia coli*

Analyses were performed as before (Meier et al., 2024a; Multamäki et al., 2022). *E. coli* CmpX13 (Mathes et al., 2009) or *E. coli* Nissle (*Ec*N, DSMZ no. 115365) bacteria containing the respective construct were used to inoculate 5 mL LB medium supplemented with 50 µg mL^{-1} Kan (LB/Kan) or 100 µg mL^{-1} Str (LB/Str). Cultures were incubated for 21-24 h at 30°C and 225 rpm agitation under non-inducing light conditions. Under green safe light, the bacteria were diluted 100-fold into fresh

LB/Kan or LB/Str, before distributing 200 µL each into 64 wells of a black-walled MTP with clear

bottom (µClear, Greiner BioOne, Frickenhausen, Germany). As a control, bacteria harboring an

empty plasmid variant with a multiple-cloning site (MCS) were also placed on the MTP. After sealing

the MTP with a gas-permeable membrane, it was placed on an Arduino-controlled illumination

device. For the application of red and blue light, a previous setup was used (Hennemann et al., 2018)

with LED peak wavelengths of (624 ± 8) nm and (463 ± 12) nm, respectively. To apply NIR light, we

devised a new matrix of 8-by-8 LEDs emitting at (800 ± 18) nm (LED800-01AU, Roithner Lasertechnik)

which is controlled by an Arduino circuit board (Suppl. Fig. 4). The bacteria inside the MTPs were

incubated for 18 h at 37°C and 750 rpm agitation. The illumination was applied in 1:2 or 1:10 duty

cycles (i.e., 1 s light and 1 s darkness, or 20 s light and 180 s darkness). After incubation, the OD_{600}

and *Ds*Red fluorescence were determined as above. For measurements of YPet fluorescence,

excitation and emission wavelengths of (500 ± 9) nm and (554 ± 9) nm were used. Following

normalization to OD_{600}, the MCS background was subtracted. Data represent mean ± s.d. of three

biological replicates and were evaluated as a function of the light intensity averaged over the duty

cycle according to Hill isotherms using Fit-o-mat.

Flow Cytometry

E. coli CmpX13 cells (Mathes et al., 2009) containing *Ac*NIRusk, *Av*NIRusk, or an empty-vector control

(MCS) were analyzed by flow cytometry (Meier et al., 2024a). After overnight incubation for 21-24

h at 30°C and 750 rpm agitation, bacterial cultures were diluted 100-fold into fresh LB/Kan medium.

For each variant, the cultures were split and incubated for 18 h at 37°C and 750 rpm agitation in

either darkness, red light [100 µW cm^{-2}, (624 ± 8) nm], blue light [70 µW cm^{-2}, (463 ± 12) nm], or

different intensities (18-270 µW cm^{-2}) of NIR light [(800 ± 18) nm]. Following incubation, the cultures

were diluted 10-fold into phosphate-buffered saline (1x sheath fluid, Bio-Rad) and analyzed on a S3e

cell sorter (Bio-Rad). The *Ds*Red fluorescence of at least 200,000 events was detected using

excitation lasers at 488 nm and 561 nm and an emission channel at (585 ± 15) nm. Data were fitted to skewed Gaussian distributions using Fit-o-mat. Experiments were repeated three times with similar results.

Growth and Expression Kinetics

The growth and reporter-gene expression of *E. coli* CmpX13 cells (Mathes et al., 2009) containing *Ac*NIRusk, *Av*NIRusk, or an empty-vector control (MCS) were analyzed over time as described before (Multamäki et al., 2022). For each variant, two 5-mL LB/Kan cultures were inoculated and incubated in darkness and under NIR light [200 µW cm^{-2}, (800 ± 18) nm] at 37°C and 600 rpm agitation. At discrete times, samples of 200 µL were supplemented with 3.5 mg mL^{-1} chloramphenicol and 0.4 mg mL^{-1} tetracycline to arrest cell growth and translation. After an incubation of at least 2 h, the OD_{600} and *Ds*Red fluorescence were determined as above. Data represent mean ± s.d. of three biological replicates and were fitted to logistic functions.

Multiplexing of Optogenetic Circuits

Pairs of plasmids harboring light-responsive TCSs were serially transformed into chemically competent *E. coli* CmpX13. 200 µL LB supplemented with 50 µg mL^{-1} Kan and 100 µg mL^{-1} Str (LB/Kan+Str) were inoculated with single bacterial clones and incubated for 21-24 h inside MTP wells at 30°C and 750 rpm agitation in darkness. The cultures were then diluted 100-fold into LB/Kan+Str before dispensation of 200 µL each into individual wells of four µClear MTPs. The MTPs were incubated for 18 h at 37°C and 750 rpm agitation in either darkness, under red light [100 µW cm^{-2}, (624 ± 8) nm], NIR light [200 µW cm^{-2}, (800 ± 18) nm], or both light colors combined. Where applicable, red light was applied continuously from below using the Arduino-controlled RGB LED matrix, whereas NIR light shone from above. After 18 h incubation, the OD_{600}, and *Ds*Red and YPet fluorescence were measured as above. Data represent mean ± s.d. of three biological replicates.

Luciferase Assays in *Agrobacterium tumefaciens*

The *Dm*REDlux and *Av*NIRlux plasmids were transferred to *Agrobacterium tumefaciens* C58 (also

known as *Agrobacterium fabrum* C58, Lab collection, ATCC 33970) by biparental conjugation using

E. coli WM3064 as a diaminopimelic acid auxotrophic donor strain (William Metcalf, UIUC,

unpublished; *thrB1004 pro thi rpsL hsdS lacZΔM15 RP4-1360 Δ(araBAD)567 ΔdapA1341::[erm pir*

(wt)]). Unless specified otherwise, C58 was routinely cultivated in LB medium at 28°C with 170 rpm

agitation. Selection for mini-Tn5 insertion mutants carrying the *lux* reporter constructs was carried

out on solid LB with 1.5% (w/v) agar and 50 µg mL^{-1} Kan. *E. coli* WM3064 strains carrying plasmids

were cultivated at 37°C with 200 rpm shaking in LB supplemented with 0.1 mM DL-a,e-

diaminopimelic acid (DAP) and 25 µg mL^{-1} Kan or 50 µg mL^{-1} ampicillin (Amp).

The protocol for biparental conjugation was described before (Dziuba et al., 2021). Briefly,

cultures of the C58 receptor and plasmid-containing *E. coli* strains were cultivated in LB or LB/Amp

medium overnight. Then, 2×10^9 cells of each strain were mixed in 15-mL falcon tubes and pelleted

at 5,421 g for 10 min. The supernatant was discarded while the cell pellet was resuspended in 100-

200 µL LB medium and spotted onto LB agar plates. After overnight incubation at 28°C, cells were

resuspended in 1 mL LB and transferred to 15-mL falcon tubes containing 9 mL LB medium. After

incubation for 2 h at 28°C and 120 rpm agitation, the cell suspension was pelleted by centrifugation,

and the supernatant was discarded. Following resuspension in 1 mL LB medium, 100 µL of 10- and

100-fold diluted cell suspensions were plated on LB-Kan agar plates. After incubation for three days

at 28°C, Kan-resistance plasmid insertion mutants were transferred to fresh LB/Kan agar plates and

screened by PCR.

At least three randomly selected transconjugants containing the luciferase reporter

constructs were analyzed for luminescence. Each experiment was performed in duplicate.

Luminescence signals, measured as arbitrary light units (ALU), were detected using a Tecan Infinite

M200 Pro MTP reader equipped with a luminometer module (Tecan Group, Ltd., Männedorf, Switzerland) during culture growth in LB medium at 23°C and 280 rpm. Measurements were taken every 20 minutes over 300 cycles (~90 h). ALU readings were normalized by the optical density at 650 nm (OD_{650}) to calculate the relative light units (RLU). To this end, tested clones were inoculated into 600 μL LB/Kan medium in 24-well plates and incubated for ~ 20 h under non-inducing conditions (i.e., red light for *Dm*REDlux and darkness for *Av*NIRlux, respectively) at 28°C and 140 rpm. Cultures were then diluted 10-fold into 1 mL LB/Kan and incubated under identical conditions for an additional 4-5 h. Subsequently, the OD_{650} was adjusted to 0.1 by adding fresh LB/Kan medium, and 100 μL of the normalized cell suspension was transferred into two 96-well MTPs. Cultivation of both MTPs was initiated simultaneously, with one incubated under dark conditions and the other one under red [(650 ± 11) nm] or NIR light [(800 ± 18) nm].

Alternatively, luminescence was assessed using a spotting assay. Precultures were prepared as above, and 5 μL of the OD_{650}-normalized cell suspension were spotted onto LB/Kan agar plates. The plates were incubated for 44 h at 28°C under dark conditions or under blue [(470 ± 13) nm], red [(650 ± 11) nm], or NIR light [(800 ± 18) nm]. Luminescence signals were documented as cumulative signal intensities using the Bio-Rad ChemiDoc XRS+ System (Bio Rad Laboratories Inc., Herkules, USA), with images captured every 0.6 s over a 1-min period. In addition, colorimetric images were acquired to visualize the outlines of the spotted cell suspensions. Luminescence was quantified from the image time series by calculating the slope of the mean pixel intensities within the outlined cell suspension areas using the Image J Fiji package (v1.54f) (Schindelin et al., 2012). Background correction was performed by subtracting the mean pixel intensity of an uninoculated region of the agar plate.

References

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung C-C, O'Neill M, Reiman D, Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli O, Bridgland A, Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB, Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A, Savy P, Singh S, Stecula A, Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Žídek A, Bapst V, Kohli P, Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**:493–500. doi:10.1038/s41586-024-07487-w

Andersen KR, Leksa NC, Schwartz TU. 2013. Optimized E. coli expression strain LOBSTR eliminates common contaminants from His-tag purification. *Prot Struct Funct Bioinf* **81**:1857–1861. doi:10.1002/prot.24364

Anthamatten D, Hennecke H. 1991. The regulatory status of the fixL- and fixJ-like genes in Bradyrhizobium japonicum may be different from that in Rhizobium meliloti. *Mol Gen Genet* **225**:38–48.

Ash C, Dubec M, Donne K, Bashford T. 2017. Effect of wavelength and beam width on penetration in light-tissue interaction using computational methods. *Lasers Med Sci* **32**:1909–1918. doi:10.1007/s10103-017-2317-4

Berkelman TR, Lagarias JC. 1986. Visualization of bilin-linked peptides and proteins in polyacrylamide

gels. *Anal Biochem* **156**:194–201.

Blum M, Andreeva A, Florentino LC, Chuguransky SR, Grego T, Hobbs E, Pinto BL, Orr A, Paysan-

Lafosse T, Ponamareva I, Salazar GA, Bordin N, Bork P, Bridge A, Colwell L, Gough J, Haft DH,

Letunic I, Llinares-López F, Marchler-Bauer A, Meng-Papaxanthos L, Mi H, Natale DA, Orengo CA,

Pandurangan AP, Piovesan D, Rivoire C, Sigrist CJA, Thanki N, Thibaud-Nissen F, Thomas PD,

Tosatto SCE, Wu CH, Bateman A. 2025. InterPro: the protein sequence classification resource in

2025. *Nucl Acids Res* **53**:D444–D456. doi:10.1093/nar/gkae1082

Bódizs S, Mészáros P, Grunewald L, Takala H, Westenhoff S. 2024. Cryo-EM structures of a bathy

phytochrome histidine kinase reveal a unique light-dependent activation mechanism. *Structure*

32:1952-1962.e3. doi:10.1016/j.str.2024.08.008

Böhm C, Lehtinen K, Multamäki E, Vanhatalo R, Brander O, Meier SSM, Rumfeldt JA, Möglich A,

Takala H. 2025. Traits and application of bathy phytochromes to bacterial optogenetics.

submitted.

Borucki B, von Stetten D, Seibeck S, Lamparter T, Michael N, Mroginski MA, Otto H, Murgida DH,

Heyn MP, Hildebrandt P. 2005. Light-induced proton release of phytochrome is coupled to the

transient deprotonation of the tetrapyrrole chromophore. *J Biol Chem* **280**:34358–64.

Burgie ES, Basore K, Rau MJ, Summers B, Mickles AJ, Grigura V, Fitzpatrick JAJ, Vierstra RD. 2024.

Signaling by a bacterial phytochrome histidine kinase involves a conformational cascade

reorganizing the dimeric photoreceptor. *Nat Commun* **15**:6853. doi:10.1038/s41467-024-50412-

y

Busch A, Guazzaroni M-E, Lacal J, Ramos JL, Krell T. 2009. The Sensor Kinase TodS Operates by a

Multiple Step Phosphorelay Mechanism Involving Two Autokinase Domains*. *J Biol Chem*

284:10353–10360. doi:10.1074/jbc.M900521200

Buschiazzo A, Trajtenberg F. 2019. Two-Component Sensing and Regulation: How Do Histidine

Kinases Talk with Response Regulators at the Molecular Level? *Annu Rev Micro* **73**:507–528.

doi:10.1146/annurev-micro-091018-054627

Chernov KG, Redchuk TA, Omelina ES, Verkhusha VV. 2017. Near-Infrared Fluorescent Proteins,

Biosensors, and Optogenetic Tools Engineered from Phytochromes. *Chem Rev* **117**:6423–6446.

doi:10.1021/acs.chemrev.6b00700

Cubillos-Ruiz A, Guo T, Sokolovska A, Miller PF, Collins JJ, Lu TK, Lora JM. 2021. Engineering living

therapeutics with synthetic biology. *Nat Rev Drug Discov* **20**:941–960. doi:10.1038/s41573-021-

00285-3

Cui M, Sun T, Li Shubin, Pan H, Liu J, Zhang X, Li L, Li Shanshan, Wei C, Yu C, Yang C, Ma N, Ma B, Lu

S, Chang J, Zhang W, Wang H. 2021. NIR light-responsive bacteria with live bio-glue coatings for

precise colonization in the gut. *Cell Rep* **36**:109690. doi:10.1016/j.celrep.2021.109690

Daeffler KN, Galley JD, Sheth RU, Ortiz-Velez LC, Bibb CO, Shroyer NF, Britton RA, Tabor JJ. 2017.

Engineering bacterial thiosulfate and tetrathionate sensors for detecting gut inflammation. *Mol*

Syst Biol **13**:923. doi:10.15252/msb.20167416

Davis SJ, Vener AV, Vierstra RD. 1999. Bacteriophytochromes: Phytochrome-Like Photoreceptors

from   Nonphotosynthetic   Eubacteria.   *Science*   **286**:2517–2520.

doi:10.1126/science.286.5449.2517

Deisseroth K, Feng G, Majewska AK, Miesenböck G, Ting A, Schnitzer MJ. 2006. Next-generation

optical technologies for illuminating genetically targeted brain circuits. *J Neurosci* **26**:10380–

10386. doi:10.1523/JNEUROSCI.3863-06.2006

Diensthuber RP, Bommer M, Gleichmann T, Möglich A. 2013. Full-length structure of a sensor

histidine kinase pinpoints coaxial coiled coils as signal transducers and modulators. *Structure*

21:1127–1136. doi:10.1016/j.str.2013.04.024

Dziuba M, Riese CN, Borgert L, Wittchen M, Busche T, Kalinowski J, Uebe R, Schüler D. 2021. The

Complex Transcriptional Landscape of Magnetosome Gene Clusters in Magnetospirillum

gryphiswaldense. *mSystems* **6**:10.1128/msystems.00893-21. doi:10.1128/msystems.00893-21

Etzl S, Lindner R, Nelson MD, Winkler A. 2018. Structure-guided design and functional

characterization of an artificial red light–regulated guanylate/adenylate cyclase for optogenetic

applications. *J Biol Chem* **293**:9078–9089. doi:10.1074/jbc.RA118.003069

Gao R, Bouillet S, Stock AM. 2019. Structural Basis of Response Regulator Function. *Annu Rev Micro*

73:175–197. doi:10.1146/annurev-micro-020518-115931

Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic assembly of

DNA molecules up to several hundred kilobases. *Nat Methods* **6**:343–345.

doi:10.1038/nmeth.1318

Gourinchas G, Etzl S, Göbl C, Vide U, Madl T, Winkler A. 2017. Long-range allosteric signaling in red

light-regulated diguanylyl cyclases. *Sci Adv* **3**:e1602498. doi:10.1126/sciadv.1602498

Hamblin MR. 2016. Shining light on the head: Photobiomodulation for brain disorders. *BBA Clinical*

6:113–124. doi:10.1016/j.bbacli.2016.09.002

Hartsough LA, Park M, Kotlajich MV, Lazar JT, Han B, Lin C-CJ, Musteata E, Gambill L, Wang MC, Tabor

JJ. 2020. Optogenetic control of gut bacterial metabolism to promote longevity. *Elife* **9**:e56849.

doi:10.7554/eLife.56849

Hennemann J, Iwasaki RS, Grund TN, Diensthuber RP, Richter F, Möglich A. 2018. Optogenetic Control

by Pulsed Illumination. *Chembiochem* **19**:1296–1304. doi:10.1002/cbic.201800030

Huber C, Strack M, Schultheiß I, Pielage J, Mechler X, Hornbogen J, Diller R, Frankenberg-Dinkel N.

2024. Darkness inhibits autokinase activity of bacterial bathy phytochromes. *J Biol Chem*

300:107148. doi:10.1016/j.jbc.2024.107148

Hughes J, Lamparter T, Mittmann F, Hartmann E, Gärtner W, Wilde A, Börner T. 1997. A prokaryotic

phytochrome. *Nature* **386**:663. doi:10.1038/386663a0

Hughes J, Winkler A. 2024. New Insight Into Phytochromes: Connecting Structure to Function. *Annu*

Rev Plant Biol **75**:153–183. doi:10.1146/annurev-arplant-070623-110636

Jacob-Dubuisson F, Mechaly A, Betton J-M, Antoine R. 2018. Structural insights into the signalling

mechanisms of two-component systems. *Nat Rev Micro* **16**:585. doi:10.1038/s41579-018-0055-

7

Jayaraman P, Devarajan K, Chua TK, Zhang H, Gunawan E, Poh CL. 2016. Blue light-mediated

transcriptional activation and repression of gene expression in bacteria. *Nucl Acids Res* **44**:6994–

7005. doi:10.1093/nar/gkw548

Jenal U, Reinders A, Lori C. 2017. Cyclic di-GMP: second messenger extraordinaire. *Nat Rev Microbiol*

15:271–284. doi:10.1038/nrmicro.2016.190

Karniol B, Vierstra RD. 2003. The pair of bacteriophytochromes from Agrobacterium tumefaciens are

histidine kinases with opposing photobiological properties. *Proc Natl Acad Sci U S A* **100**:2807–

2812. doi:10.1073/pnas.0437914100

Krell T, Lacal J, Busch A, Silva-Jiménez H, Guazzaroni M-E, Ramos JL. 2010. Bacterial Sensor Kinases:

Diversity in the Recognition of Environmental Signals. *Annu Rev Micro* **64**:539–559.

doi:10.1146/annurev.micro.112408.134054

Lamparter T, Michael N, Mittmann F, Esteban B. 2002. Phytochrome from Agrobacterium

tumefaciens has unusual spectral properties and reveals an N-terminal chromophore

attachment site. *Proc Natl Acad Sci U S A* **99**:11628–11633. doi:10.1073/pnas.152263999

Landry BP, Palanki R, Dyulgyarov N, Hartsough LA, Tabor JJ. 2018. Phosphatase activity tunes two-

component system sensor detection threshold. *Nat Commun* **9**:1433. doi:10.1038/s41467-018-

03929-y

Lau PCK, Wang Y, Patel A, Labbé D, Bergeron H, Brousseau R, Konishi Y, Rawlings M. 1997. A bacterial

basic region leucine zipper histidine kinase regulating toluene degradation. *Proc Natl Acad Sci U*

S A **94**:1453–1458. doi:10.1073/pnas.94.4.1453

Laub MT, Goulian M. 2007. Specificity in Two-Component Signal Transduction Pathways. *Annu Rev*

Genet **41**:121–145. doi:10.1146/annurev.genet.41.042007.170548

Lazar JT, Tabor JJ. 2021. Bacterial two-component systems as sensors for synthetic biology

applications. *Curr Opin Syst Biol* **28**:100398. doi:10.1016/j.coisb.2021.100398

Lehtinen K, Nokia MS, Takala H. 2022. Red Light Optogenetics in Neuroscience. *Front Cell Neurosci*

15.

Leopold AV, Chernov KG, Shemetov AA, Verkhusha VV. 2019. Neurotrophin receptor tyrosine kinases

regulated with near-infrared light. *Nat Commun* **10**:1–13. doi:10.1038/s41467-019-08988-3

Leopold AV, Thankachan S, Yang C, Gerashchenko D, Verkhusha VV. 2022. A general approach for

engineering RTKs optically controlled with far-red light. *Nat Methods* **19**:871–880.

doi:10.1038/s41592-022-01517-z

Levskaya A, Chevalier AA, Tabor JJ, Simpson ZB, Lavery LA, Levy M, Davidson EA, Scouras A, Ellington

AD, Marcotte EM, Voigt CA. 2005. Synthetic biology: Engineering Escherichia coli to see light.

Nature **438**:441–442. doi:10.1038/nature04405

Liu W, Yuan JS, Stewart Jr CN. 2013. Advanced genetic tools for plant biotechnology. *Nat Rev Genet*

14:781–793. doi:10.1038/nrg3583

Magaraci MS, Veerakumar A, Qiao P, Amurthur A, Lee JY, Miller JS, Goulian M, Sarkar CA. 2014.

Engineering Escherichia coli for Light-Activated Cytolysis of Mammalian Cells. *ACS Synth Biol*

3:944–948. doi:10.1021/sb400174s

Malla TN, Hernandez C, Muniyappan S, Menendez D, Bizhga D, Mendez JH, Schwander P, Stojković

EA, Schmidt M. 2024. Photoreception and signaling in bacterial phytochrome revealed by single-

particle cryo-EM. *Sci Adv* **10**:eadq0653. doi:10.1126/sciadv.adq0653

Mathes T, Vogl C, Stolz J, Hegemann P. 2009. In vivo generation of flavoproteins with modified

cofactors. *J Mol Biol* **385**:1511–1518. doi:10.1016/j.jmb.2008.11.001

Meier SSM, Multamäki E, Ranzani AT, Takala H, Möglich A. 2024a. Leveraging the histidine kinase-

phosphatase duality to sculpt two-component signaling. *Nat Commun* **15**:4876.

doi:10.1038/s41467-024-49251-8

Meier SSM, Multamäki E, Ranzani AT, Takala H, Möglich A. 2024b. Multimodal Control of Bacterial

Gene Expression by Red and Blue Light. *Meth Mol Biol* **2760**:463–477. doi:10.1007/978-1-0716-

3658-9_26

Möglich A. 2019. Signal transduction in photoreceptor histidine kinases. *Prot Sci* **28**:1923–1946.

doi:10.1002/pro.3705

Möglich A. 2018. An Open-Source, Cross-Platform Resource for Nonlinear Least-Squares Curve

Fitting. *J Chem Educ* **95**:2273–2278. doi:10.1021/acs.jchemed.8b00649

Möglich A, Ayers RA, Moffat K. 2010. Addition at the molecular level: signal integration in designed

Per-ARNT-Sim receptor proteins. *J Mol Biol* **400**:477–486. doi:10.1016/j.jmb.2010.05.019

Möglich A, Ayers RA, Moffat K. 2009. Design and signaling mechanism of light-regulated histidine

kinases. *J Mol Biol* **385**:1433–1444. doi:10.1016/j.jmb.2008.12.017

Mukougawa K, Kanamoto H, Kobayashi T, Yokota A, Kohchi T. 2006. Metabolic engineering to produce

phytochromes with phytochromobilin, phycocyanobilin, or phycoerythrobilin chromophore in

Escherichia coli. *FEBS Lett* **580**:1333–1338. doi:10.1016/j.febslet.2006.01.051

Multamäki E, García de Fuentes A, Sieryi O, Bykov A, Gerken U, Ranzani AT, Köhler J, Meglinski I,

Möglich A, Takala H. 2022. Optogenetic Control of Bacterial Expression by Red Light. *ACS Synth*

Biol **11**:3354–3367. doi:10.1021/acssynbio.2c00259

Multamäki E, Nanekar R, Morozov D, Lievonen T, Golonka D, Wahlgren WY, Stucki-Buchli B, Rossi J,

Hytönen VP, Westenhoff S, Ihalainen JA, Möglich A, Takala H. 2021. Comparative analysis of two

paradigm bacteriophytochromes reveals opposite functionalities in two-component signaling.

Nat Commun **12**:4394. doi:10.1038/s41467-021-24676-7

Nguyen AW, Daugherty PS. 2005. Evolutionary optimization of fluorescent proteins for intracellular

FRET. *Nat Biotechnol* **23**:355–360. doi:10.1038/nbt1066

Ohlendorf R, Möglich A. 2022. Light-regulated gene expression in Bacteria: Fundamentals, advances,

and perspectives. *Front Bioeng Biotech* **10**:1029403.

Ohlendorf R, Schumacher CH, Richter F, Möglich A. 2016. Library-Aided Probing of Linker

Determinants in Hybrid Photoreceptors. *ACS Synth Biol* **5**:1117–1126.

doi:10.1021/acssynbio.6b00028

Ohlendorf R, Vidavski RR, Eldar A, Moffat K, Möglich A. 2012. From dusk till dawn: one-plasmid

systems for light-regulated gene expression. *J Mol Biol* **416**:534–542.

doi:10.1016/j.jmb.2012.01.001

Ong NT, Olson EJ, Tabor JJ. 2018. Engineering an E. coli Near-Infrared Light Sensor. *ACS Synth Biol*

7:240–248. doi:10.1021/acssynbio.7b00289

Pan H, Li L, Pang G, Han C, Liu B, Zhang Y, Shen Y, Sun T, Liu J, Chang J, Wang H. 2021. Engineered NIR

light-responsive bacteria as anti-tumor agent for targeted and precise cancer therapy. *Chem Eng*

J **426**:130842. doi:10.1016/j.cej.2021.130842

Parkinson JS, Kofoid EC. 1992. Communication Modules in Bacterial Signaling Proteins. *Annu Rev*

Genet **26**:71–112. doi:10.1146/annurev.ge.26.120192.000443

Ramos-González M-I, Olson M, Gatenby AA, Mosqueda G, Manzanera M, Campos MJ, Víchez S,

Ramos JL. 2002. Cross-Regulation between a Novel Two-Component Signal Transduction System

for Catabolism of Toluene in Pseudomonas mendocina and the TodST System from Pseudomonas

putida. *J Bact* **184**:7062–7067. doi:10.1128/jb.184.24.7062-7067.2002

Ranzani AT, Buchholz K, Blackholm M, Kopkin H, Möglich A. 2024. Induction of bacterial expression

at the mRNA level by light. *Nucl Acids Res* **52**:10017–10028. doi:10.1093/nar/gkae678

Rottwinkel G, Oberpichler I, Lamparter T. 2010. Bathy Phytochromes in Rhizobial Soil Bacteria. *J Bact*

192:5124–5133. doi:10.1128/JB.00672-10

Rumfeldt JA, Takala H, Liukkonen A, Ihalainen JA. 2019. UV-Vis Spectroscopy Reveals a Correlation

Between Y263 and BV Protonation States in Bacteriophytochromes. *Photochem Photobiol*

95:969–979. doi:https://doi.org/10.1111/php.13095

Russo FD, Silhavy TJ. 1993. The essential tension: opposed reactions in bacterial two-component

regulatory systems. *Trends Microbiol* **1**:306–310.

Sankaran S, del Campo A. 2019. Optoregulated Protein Release from an Engineered Living Material.

Adv Biosyst **3**:1800312. doi:10.1002/adbi.201800312

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012.

Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682.

doi:10.1038/nmeth.2019

Schmidl SR, Sheth RU, Wu A, Tabor JJ. 2014. Refactoring and Optimization of Light-Switchable

Escherichia coli Two-Component Systems. *ACS Synth Biol* **3**:820–831. doi:10.1021/sb500273n

Schmidt A, Sauthof L, Szczepek M, Lopez MF, Escobar FV, Qureshi BM, Michael N, Buhrke D, Stevens

T, Kwiatkowski D, von Stetten D, Mroginski MA, Krauß N, Lamparter T, Hildebrandt P, Scheerer P.

2018. Structural snapshot of a bacterial phytochrome in its functional intermediate state. *Nat*

Commun **9**:4912. doi:10.1038/s41467-018-07392-7

Silva-Jiménez H, Ramos JL, Krell T. 2012. Construction of a prototype two-component system from

the phosphorelay system TodS/TodT. *Protein Eng Des Sel* **25**:159–169.

doi:10.1093/protein/gzs001

Stabel R, Stüven B, Hansen JN, Körschen HG, Wachten D, Möglich A. 2019. Revisiting and Redesigning

Light-Activated Cyclic-Mononucleotide Phosphodiesterases. *J Mol Biol* **431**:3029–3045.

doi:10.1016/j.jmb.2019.07.011

Strack RL, Strongin DE, Bhattacharyya D, Tao W, Berman A, Broxmeyer HE, Keenan RJ, Glick BS. 2008.

A noncytotoxic DsRed variant for whole-cell labeling. *Nat Methods* **5**:955–957.

doi:10.1038/nmeth.1264

Stüven B, Stabel R, Ohlendorf R, Beck J, Schubert R, Möglich A. 2019. Characterization and

engineering of photoactivated adenylyl cyclases. *Biol Chem* **400**:429–441. doi:10.1515/hsz-

2018-0375

Tabor JJ, Levskaya A, Voigt CA. 2011. Multichromatic control of gene expression in Escherichia coli. *J*

Mol Biol **405**:315–324. doi:10.1016/j.jmb.2010.10.038

Takala H, Edlund P, Ihalainen JA, Westenhoff S. 2020. Tips and turns of bacteriophytochrome

photoactivation. *Photochem Photobiol Sci* **19**:1488–1510. doi:10.1039/D0PP00117A

Tang K, Beyer HM, Zurbriggen MD, Gärtner W. 2021. The Red Edge: Bilin-Binding Photoreceptors as

Optogenetic Tools and Fluorescence Reporters. *Chem Rev* **121**:14906–14956.

doi:10.1021/acs.chemrev.1c00194

Tasler R, Moises T, Frankenberg-Dinkel N. 2005. Biochemical and spectroscopic characterization of

the bacterial phytochrome of Pseudomonas aeruginosa. *FEBS J* **272**:1927–1936.

doi:10.1111/j.1742-4658.2005.04623.x

Velazquez Escobar F, Piwowarski P, Salewski J, Michael N, Fernandez Lopez M, Rupp A, Qureshi BM,

Scheerer P, Bartl F, Frankenberg-Dinkel N, Siebert F, Andrea Mroginski M, Hildebrandt P. 2015. A

protonation-coupled feedback mechanism controls the signalling process in bathy

phytochromes. *Nature Chem* **7**:423–430. doi:10.1038/nchem.2225

Weissleder R. 2001. A clearer vision for *in vivo* imaging. *Nat Biotech* **19**:316–317. doi:10.1038/86684

Xu Q, Vogt A, Frechen F, Yi C, Küçükerden M, Ngum N, Sitjà-Roqueta L, Greiner A, Parri R, Masana M,

Wenger N, Wachten D, Möglich A. 2024. Engineering Bacteriophytochrome-coupled

Photoactivated Adenylyl Cyclases for Enhanced Optogenetic cAMP Modulation. *J Mol Biol*,

Photosensory Receptors-Mechanisms and Effects **436**:168257. doi:10.1016/j.jmb.2023.168257

Yang C, Cui M, Zhang Y, Pan H, Liu J, Wang S, Ma N, Chang J, Sun T, Wang H. 2020. Upconversion

optogenetic micro-nanosystem optically controls the secretion of light-responsive bacteria for

systemic immunity regulation. *Commun Biol* **3**:1–14. doi:10.1038/s42003-020-01287-4

Yang X, Kuk J, Moffat K. 2008. Crystal structure of Pseudomonas aeruginosa bacteriophytochrome:

Photoconversion and signal transduction. *Proc Natl Acad Sci U S A* **105**:14715–14720.

Yi C, Meier SSM, Kehr M, Möglich A. 2025. Spectroscopic Analyses of Photoconversion in

Phytochromes. *Meth Mol Biol* **submitted**.

Ziegler T, Möglich A. 2015. Photoreceptor engineering. *Front Mol Biosci* **2**:30.

doi:10.3389/fmolb.2015.00030

Zienicke B, Molina I, Glenz R, Singer P, Ehmer D, Escobar FV, Hildebrandt P, Diller R, Lamparter T.

2013. Unusual Spectral Properties of Bacteriophytochrome Agp2 Result from a Deprotonation

of the Chromophore in the Red-absorbing Form Pr*. *J Biol Chem* **288**:31738–31751.

doi:10.1074/jbc.M113.479535

Acknowledgements

We thank our colleagues for discussion; Dr. Roman S. Iwasaki, Dr. Ralph P. Diensthuber, and Tobias

von Basse for preliminary experiments; Tino Krell for generously providing the pMIR66 and pMIR77

plasmids; and Mr. Braun and the electronical workshop at University Bayreuth for support. Funding

was provided by the European Commission (FET Open NEUROPA grant 863214 to A.M.); the German

Research Foundation (grants MO2192/4-2 to A.M. and UE200/1-1 to R.U.); and the Research Council

of Finland (grant 330678 to H.T.).

Author Contributions

S.S.M.M. and A.M. conceived of the project; H.T., R.U., and A.M. acquired funding; S.S.M.M., M.H.,

C.B., R.U., and A.M. designed research; S.S.M.M., M.H., C.B., E.L.R.D., R.U., and A.M. conducted

research; S.S.M.M., M.H., C.B., E.L.R.D., R.U., and A.M. analyzed data; S.S.M.M. and A.M. wrote the

manuscript; all authors edited the manuscript.

Competing Interests

987 The authors declare no competing interests.

Figure Legends

Figure 1

Spectroscopic characterization of bacteriophytochrome photosensory core modules (PCM) from

Azorhizobium caulinodans (*Ac*PCM), *Agrobacterium vitis* (*Av*PCM), and *Pseudomonas aeruginosa*

(*Pa*PCM). (a) UV/vis absorbance spectra of *Ac*PCM at pH 8 in darkness (black line), and after

illumination with near-infrared (NIR, purple line), red (red), and blue light (blue). Data are normalized

by the absorbance at 403 nm within the Soret band. The lower panel shows the NIR-dark difference

absorbance spectrum with maxima and minima marked. (b) and (c) as in (a) but for *Av*PCM and

*Pa*PCM. (d) Absorbance spectra of *Ac*PCM in darkness (solid lines) and after NIR illumination (dashed

lines) at different pH values as indicated by color. (e) and (f) as in (d) but for *Av*PCM and *Pa*PCM. (g)

Absorbance at 700 nm for *Ac*PCM (light purple circles), *Av*PCM (dark green squares), and *Pa*PCM

(orange triangles) upon NIR-light exposure at different pH values. Data were fitted to the Henderson-

Hasselbalch equation to determine apparent pK_a values (dashed lines, *Ac*PCM: 7.2 ± 0.1, *Av*PCM: 8.0

± 0.1, *Pa*PCM: 9.9 ± 0.2). (h) Calculated Pfr fractions of *Ac*PCM (light purple circles), *Av*PCM (dark

green square), and *Pa*PCM (orange triangles) after red-light exposure at different pH values. The

underlying absorbance spectra are shown in Suppl. Fig. 1a-c. (i) Dark-recovery kinetics at 25°C and

pH 8 after NIR-illumination as followed by absorbance at 760 nm. Time constants for *Ac*PCM (light

purple), *Av*PCM (dark green), and *Pa*PCM (orange) were determined by single-exponential fits.

Figure 2

Engineering of light-regulated sensor histidine kinases (SHK) based on photosensory core modules

(PCM) from bathy-bacteriophytochromes (BphP). (a) Schematic of the pNIRusk plasmids to control

gene expression in bacteria by NIR light. The LacI^q promoter constitutively expresses a tricistronic

operon consisting of heme oxygenase (HO1), the light-regulated SHK (chimera of *Ac/Av*PCM and

FixL), and the response regulator FixJ. When phosphorylated, FixJ binds to the FixK2 promoter and

triggers the expression of a gene of interest (GOI). (b) Response of select *Ac*NIRusk (left) and

*Av*NIRusk (right) systems to different light conditions (grey: darkness; purple: NIR light; red: red

light). *Ds*Red reporter fluorescence was normalized by the optical density of the bacterial cultures.

Individual pNIRusk variants are denoted by the relative lengths of the linkers within their underlying

SHKs (see Suppl. Fig. 3), with the variants chosen for further analyses highlighted in bold. As control,

an empty-vector construct (MCS) is shown, and data are normalized by the reference fluorescence

value determined for the *Av*+17a variant. (c) Flow-cytometric analyses of bacteria containing

*Ac*NIRusk-1a (top) or *Av*NIRusk+17a (bottom) upon incubation in darkness (black), blue (blue, 70 μW

cm^{-2}), red (red, 100 μW cm^{-2}), or NIR light (purple, 200 μW cm^{-2}). An empty-vector control (MCS) is

shown in green. (d) Light-dose response of bacteria containing *Ac*NIRusk-1a incubated in NIR (purple

circles), red (red squares), and blue light (blue pentagons). (e) as in (d) but for *Av*NIRusk+17a.

Figure 3

Engineering of orthogonal light-regulated two-component systems (TCS). (a) Schematic of the

*Dm*Ttr-REDusk circuit based on the TtrSR TCS from *Shewanella baltica* combined with the

photosensory core module (PCM) from the *Deinococcus maricopensis* bacteriophytochrome (BphP)

(top). Reporter fluorescence of bacteria bearing circuits with different linker lengths of their

underlying sensor histidine kinases (SHK) upon cultivation in darkness (grey), NIR (purple), and red

light (red) (bottom). An empty-vector control is denoted as 'MCS', and data are normalized as in Fig.

2. Variants are marked by the relative lengths of the linkers within their underlying SHKs compared

to pREDusk (see Suppl. Fig. 6). Variants characterized further are highlighted in bold. (b) as in (a) but

for the *Av*Tod-NIRusk circuit based on the TodST TCS from *Pseudomonas putida* combined with the

BphP PCM from *Agrobacterium vitis*. (c) Multiplexing of bacteria containing two optogenetic circuits,

as noted below the graph, to separately control the expression of *Ds*Red (top) and YPet (bottom)

reporters. After incubation in darkness (gray), NIR (purple), red (red), and combined NIR and red

light (red and purple stripes), fluorescence was measured and, in the case of *Ds*Red, normalized as

in Fig. 2. YPet fluorescence was normalized by the reference value determined for *Dm*REDusk-YPet

under red light (this panel).

Figure 4

Light-regulated gene expression in *Escherichia coli* Nissle (*Ec*N) and *Agrobacterium tumefaciens*. (a)

Light-dose response of *Ec*N harboring *Av*NIRusk after incubation at varying NIR-light (purple circles),

red-light (red squares), and blue-light (blue pentagons) intensities. Corresponding data for *Ac*NIRusk

are shown in Suppl. Fig. 8. (b) Schematic of the *Av*NIRlux and *Dm*REDlux circuits based on *Av*PCM

and *Dm*PCM. The tricistronic cassette comprising heme oxygenase (HO1), the sensor histidine kinase

(chimera of *Av/Dm*PCM and FixL), and FixJ is controlled by the constitutive PmamDC promoter from

Magnetospirillum gryphiswaldense, whereas the FixK2 promoter regulates the expression of the

luxABCDE operon. (c) *A. tumefaciens* bacteria bearing *Av*NIRlux (left) and *Dm*REDlux (right) were

spotted on plates and incubated at different light conditions (darkness, NIR, red, or blue light),

followed by luminescence (Lumi.) measurements (see Suppl. Fig. 9a). (d) *A. tumefaciens* bacteria

carrying *Av*NIRlux and *Dm*REDlux were incubated in liquid culture for 20 h under different light

conditions (gray: darkness, purple: NIR light, red: red light), followed by recording of reporter

luminescence (see Suppl. Fig. 9c-d).

Figure 5

Photoreception by select bathy-bacteriophytochromes (BphP) at different pH values. Throughout

the entire tested pH range, the photosensory core modules (PCM) of the bathy-BphPs from

Azorhizobium caulinodans (*Ac*PCM), *Agrobacterium vitis* (*Av*PCM), and *Pseudomonas aeruginosa*

(*Pa*PCM) adopt a pure Pfr state in darkness with fully protonated biliverdin (BV) chromophore. NIR

light drives the complete conversion to the Pr state in which BV can be either protonated or

deprotonated as governed by pH and the respective pK_a value of the bathy-BphP. Red light populates

a photostationary mixture with Pr and Pfr proportions strongly depending on pH. At high pH values,

red light can drive the bathy-BphP nearly completely to the Pr state for *Ac*PCM and *Av*PCM. Owing

to a higher pK_a value for BV protonation, *Pa*PCM is less affected but follows the same trend.

Figure 1

Figure 2

Figure 3

a

*Av*NIRusk (*Ec*N)

Red
Blue
NIR

Norm. Fluorescence [a. u.]

<Light Intensity> [µW cm⁻²]

b

*Av*NIRlux/*Dm*REDlux

P$_{mamDC}$ FixK2

HO | PCM | FixL | FixJ // luxA | B | C | D | E

Av
Dm

c

	*Av*NIRlux		*Dm*REDlux	
	Bright field	Lumi.	Bright field	Lumi.
Dark				
NIR				
Red				
Blue				

d

Dark   NIR   Red

Norm. RLU [a. u.]

*Av*NIRlux   *Dm*REDlux

Figure 4

Figure 5

Engineering NIR-Sighted Bacteria

Stefanie S. M. Meier[1,‡], Michael Hörzing[1,‡], Cornelia Böhm[2,3,‡], Emma L. R. Düthorn[1,‡], Heikki Takala[2,‡], René Uebe[4,5,‡], Andreas Möglich[1,5,6,*,‡]

[1] Department of Biochemistry, University of Bayreuth, 95447 Bayreuth, Germany.

[2] Department of Biological and Environmental Science, Nanoscience Center, University of Jyvaskyla, Jyvaskyla 40014, Finland.

[3] Institute of Biochemistry, Graz University of Technology, 8010 Graz, Austria

[4] Department of Microbiology, University of Bayreuth, 95447 Bayreuth, Germany.

[5] Bayreuth Center for Biochemistry & Molecular Biology, Universität Bayreuth, 95447 Bayreuth, Germany.

[6] North-Bavarian NMR Center, Universität Bayreuth, 95447 Bayreuth, Germany.

* for correspondence: andreas.moeglich@uni-bayreuth.de

‡ ORCID identifiers: S.S.M.M. 0009-0002-6028-5223; M.H. 0009-0008-2301-3783; C.B. 0000-0002-3552-2349; E.L.R.D. 0009-0000-4156-5428; H.T. 0000-0003-2518-8583; R.U. 0000-0003-2357-1589; A.M. 0000-0002-7382-2772

Contents

Supplementary Figure 1

Additional spectroscopic characterization of bacteriophytochrome photosensory core modules

(PCM) from *Azorhizobium caulinodans* (*Ac*PCM), *Agrobacterium vitis* (*Av*PCM), and *Pseudomonas*

aeruginosa (*Pa*PCM). (a) Absorbance spectra of *Ac*PCM at different pH values as indicated below the

graph in darkness (solid line) and after red-light-illumination (dotted line). Data are normalized by the absorbance at 403 nm within the Soret band. (b) and (c) as in (a) but for *Av*PCM and *Pa*PCM. (d) Dark-recovery kinetics of *Ac*PCM (light purple circles), *Av*PCM (dark green circles), and *Pa*PCM (orange circles) monitored at 760 nm, 25°C, and pH 8 after illumination with red light. Time constants determined by single-exponential fits are (147 ± 1) s, (71 ± 1) s and (599 ± 2) s for *Ac*PCM, *Av*PCM, and *Pa*PCM, respectively. (e) Absorbance spectra recorded in darkness (solid lines) and after dark recovery (dotted lines) for *Ac*PCM (light purple), *Av*PCM (dark green), and *Pa*PCM (orange) at pH 8. (f) Ratio of the quantum yields for Pr→Pfr over Pfr→Pr photoconversion as a function of pH. (g) Dark-recovery kinetics of *Ac*PCM recorded at 760 nm after illuminating with NIR light at 25°C and different pH as indicated below the graph. Time constants were determined by single-exponential fits, with the results shown in the table. (h) and (i) as in (g) but for *Av*PCM and *Pa*PCM. (j) Absorbance spectra of *Ac*PCM recorded in darkness (solid lines) and after dark recovery (dotted lines) at different pH as indicated below the graph. (k) and (l) as in (j) but for *Av*PCM and *Pa*PCM.

Supplementary Figure 2

Multiple sequence alignment of the sensor histidine kinases within *Dm*REDusk, *Ac*NIRusk+0, and *Av*NIRusk+0. The different shading marks the fusion of the photosensory core modules (PCM) of the bacteriophytochromes from *Deinococcus maricopensis* (*Dm*PCM), *Azorhizobium caulinodans* (*Ac*PCM), and *Agrobacterium vitis* (*Av*PCM) with the catalytic domains of the sensor histidine kinase FixL from *Bradyrhizobium japonicum*. The linker region is shown in pink, and the active-site histidine is marked with a red triangle.

Supplementary Figure 3

a

```
Plasmid Label    501                        257                               291              Expression
MH013   sc    VLRHN ELLADERGKADVRQRLLNE   TEHQQTQARLQELQSELVHVSRLSAMGEMASALA HELN      No
MH021   −4    VLRHN ELLADERG                             RLSAMGEMASALA HELN      NIR/Red
MH019   −1a   VLRHN ELLAD                          ELVHVSRLSAMGEMASALA HELN      NIR
MH034   −1b   VLRHN ELL                         QSELVHVSRLSAMGEMASALA HELN      NIR/Red
MH005   0     VLRHN                          LQELQSELVHVSRLSAMGEMASALA HELN      Dark/NIR/Red
MH017   +16a  VLRHN ELLADER              TEHQQTQARLQELQSELVHVSRLSAMGEMASALA HELN      Dark/Red
MH018   +16b  VLRHN ELLADERG              EHQQTQARLQELQSELVHVSRLSAMGEMASALA HELN      Dark
```

b

```
Plasmid Label    499                        257                               291              Expression
MH014   sc    VLRHN ELMQEERNKADIRQRVLNE   TEHQQTQARLQELQSELVHVSRLSAMGEMASALA HELN      No
MH030   −5    VLRHN ELMQEER                             RLSAMGEMASALA HELN      NIR/Red
MH023   −1    VLRHN ELMQE                          ELVHVSRLSAMGEMASALA HELN      NIR
MH006   0     VLRHN                          LQELQSELVHVSRLSAMGEMASALA HELN      Dark/NIR/Red
MH029   +16a  VLRHN ELMQEERN             EHQQTQARLQELQSELVHVSRLSAMGEMASALA HELN      Dark/Red
MH020   +16b  VLRHN ELMQEER             TEHQQTQARLQELQSELVHVSRLSAMGEMASALA HELN      Dark/Red
MH028   +17a  VLRHN ELMQEERK             TEHQQTQARLQELQSELVHVSRLSAMGEMASALA HELN      NIR
MH038   +17b  VLRHN ELMQEERN             TEHQQTQARLQELQSELVHVSRLSAMGEMASALA HELN      NIR/Red
```

*Ac*NIRusk (a) and *Av*NIRusk (b) PATCHY start constructs (sc) and derived linker variants. The linker between the *Ac*/*Av*PCM and FixL was adopted from the parental proteins and is defined as the region between the PCM (residues VLRHN) and the active-site histidine at position 291. At the junction, a *Ksp*AI restriction site and a frameshift were inserted. Underlined amino acids originate from a cloning artifact. PATCHY primers are listed in Suppl. Table 1.

Supplementary Figure 4

Setup for illumination with near-infrared (NIR) light at different intensities in a microtiter-plate (MTP) format. (a) The programmable 8-by-8 matrix of NIR light-emitting diodes (LED) is based on a previous setup (Hennemann et al., 2018; Stüven et al., 2019) and integrated into a 3D-printed adapter. NIR-light intensities can be adjusted by an Arduino microcontroller. (b) Emission spectrum of the NIR LED matrix with a maximum at 800 nm and a full width at half-maximum of 18 nm.

Supplementary Figure 5

(a-b) Analyses by flow cytometry of *Ds*Red production in bacteria containing *Ac*NIRusk (a) or *Av*NIRusk (b) when incubated in darkness (black), or under different NIR-light intensities. An empty-vector control (MCS) is shown in green. (c) Growth kinetics of bacteria bearing *Ac*NIRusk-*Ds*Red (circles) or empty-vector controls (MCS, squares) in darkness (gray) or NIR light (purple). (d) as in (c) but for *Av*NIRusk. (e) *Ds*Red production induced by NIR light in bacteria containing *Ac*NIRusk (light purple circles) or *Av*NIRusk (dark green circles) over time. The background fluorescence of bacteria carrying the empty-vector control is indicated by the dashed line. Fluorescence readings are normalized as in Fig. 2b. Data were fitted by logistic functions.

Supplementary Figure 6

a

Plasmid	Label	510		407	441	Expression
SM123	sc	GARL AALQDLNARLERANRELEGRNAELRRFAF	RELINTQRQLNENRALLEHAQRIAIAGELGASLS HELN			No
SM146	−13	GARL A	AIAGELGASLS HELN			Red
SM147	0	GARL	LNENRALLEHAQRIAIAGELGASLS HELN			Dark/NIR
SM148	+5	GARL AA	QRQLNENRALLEHAQRIAIAGELGASLS HELN			Red
SM149	+7a	GARL A	INTQRQLNENRALLEHAQRIAIAGELGASLS HELN			Dark/NIR
SM150	+7b	GARL AALQDLNARLERANRELEGRNAELRR	LGASLS HELN			Dark/NIR
SM153	+14	GARL AALQDLNARLERANRELEGRNAELRRFAF<u>D</u>	AGELGASLS HELN			Dark/NIR

b

Plasmid	Label	499		726	760	Expression
SM128	sc	VLRHN ELMQEERNKADIRQRVLNE	TEKKQAQENLNQLQQQLVYVSRSATMGEFAAYIA HEIN			No
SM160	+3	VLRHN ELMQEERNKADIRQRVLNE	MGEFAAYIA HEIN			NIR
SM161	+16	VLRHN ELMQEERNKADIRQRVLNE<u>V</u>	QQQLVYVSRSATMGEFAAYIA HEIN			Dark/Red
SM162	+19	VLRHN ELMQEERNKADIRQRVLNE<u>V</u>	NQLQQQLVYVSRSATMGEFAAYIA HEIN			Dark
SM163	+21	VLRHN ELMQEERNKADIRQR	KQAQENLNQLQQQLVYVSRSATMGEFAAYIA HEIN			NIR/Red
SM164	+26	VLRHN ELMQEERNKADIRQRVL	TEKKQAQENLNQLQQQLVYVSRSATMGEFAAYIA HEIN			Dark/Red

*Dm*Ttr-REDusk (a) and *Av*Tod-NIRusk (b) PATCHY start constructs (sc) and derived linker variants. The

linker between the *Dm*/*Av*PCM and the TtrS/TodS two-component system was adopted from the

natural proteins. At the junction, a *Eco*32I or *Ksp*AI restriction site and a frameshift were inserted.

Underlined amino acids highlighted derive from a cloning artifact. PATCHY primers are listed in Suppl.

Table 1.

Supplementary Figure 7

Characterization of constructs used in multiplexing experiments. (a) Schematic of *Dm*Ttr+7b (top) and associated light-dose response (bottom). (b) and (c) as in (a) but for *Av*Tod+16 and *Av*Tod+21. Data in panels a-c are normalized as in Fig. 2. (d) Schematic of *Dm*DERusk-StrR-YPet (top) and the associated light-dose response (bottom). (e) as in (d) but for *Av*NIRusk-StrR-YPet. Data in panels d-f are normalized as in Fig. 3. (f) The schematic summarizes the results of the multiplexing experiments from Fig. 3c. The production of *Ds*Red and YPet under different light conditions are shown by pink pentagons and green heptagons, respectively.

Supplementary Figure 8

Light-regulated gene expression in *Escherichia coli* Nissle (*Ec*N) containing *Ac*NIRusk after incubation at varying NIR-light (purple circles), red-light (red squares), and blue-light (blue pentagons) intensities.

Supplementary Figure 9

Gene expression regulated by *Av*NIRlux and *Dm*REDlux in *Agrobacterium tumefaciens*. (a) Quantified

luminescence from the spotting assay in Fig. 4b. Bar colors represent incubation in darkness (gray),

NIR (purple), red (red), or blue (blue) light. (b) Optical density at 650 nm of *A. tumefaciens* bacteria

bearing *Av*NIRlux or *Dm*REDlux grown in liquid culture under different light conditions. (c)

Normalized reporter luminescence of the bacteria from panel (b) harboring *Av*NIRlux and grown in

darkness (grey) or under NIR light (purple). The vertical black line denotes a time point of 20 h on

which the data in Fig. 4d are based. (d) as for (c) but for *A. tumefaciens* bacteria containing *Dm*REDlux

and illumination with red instead of NIR light (red).

Supplementary Figure 10

Biliverdin binding pocket in the photosensory core modules (PCM) of (a) *Azorhizobium caulinodans* bacteriophytochrome (BphP), (b) *Agrobacterium vitis* BphP, (c) *Pseudomonas aerugionsa* BphP, and (d) *Agrobacterium tumefaciens* Agp2 BphP. Whereas experimental structures are available for the PCMs of AgP2 [PDB entry 6g1y (Schmidt et al., 2018)] and *Pa*BphP [3c2w (Yang et al., 2008)], for *Ac*BphP and *Av*BphP, PCM models were calculated with AlphaFold3 (Abramson et al., 2024). Tyrosine 190 and serine 275 are highlighted in the *Pa*PCM structure by magenta color. The other three PCMs feature less polar phenylalanine and alanine residues, respectively, in the structurally equivalent positions.

Supplementary Table 1

Oligonucleotides used in this study. Overhang regions are indicated in bold, and restriction enzyme

recognition sites are underlined.

Name	Sequence 5'-3'
Primers for cloning of bathy-PCM expression constructs	
M001_Ac_Bb_for	CCTGGTTGAAATTGTGCTGCGTCATAATGAGAATCTTTATTTTCAGCATCACCA
M002_Ac_Bb_rev	GATCAACTGCTTCTGCATTCGGCATGGTATATCTCCTTATTAAAGTTAAACAAAA
M003_Ac_Ins_for	TTTTGTTTAACTTTAATAAGGAGATATACCATGCCGAATGCAGAAGCAGTTGATC
M004_Ac_Ins_rev	TGGTGATGCTGAAAATAAAGATTCTCATTATGACGCAGCACAATTTCAACCAGG
M005_Av_Bb_for	ACCGTTAAAGTGGTTCTGCGCCATAATGAGAATCTTTATTTTCAGCATCACCA
M006_Av_Bb_rev	ATCAACGGTCTGGGTTGCAGGCATGGTATATCTCCTTATTAAAGTTAAACAAAA
M007_Av_Ins_for	TTTTGTTTAACTTTAATAAGGAGATATACCATGCCTGCAACCCAGACCGTTGAT
M008_Av_Ins_rev	TGGTGATGCTGAAAATAAAGATTCTCATTATGGCGCAGAACCACTTTAACGGT
M069_Pa-Ins-fw	TTTTGTTTAACTTTAATAAGGAGATATACCATGACGAGCATCACCCCGGTTACCC
M070_Pa-Ins-rev	TGGTGATGGCCCTGAAAATAAAGATTCTCCGCATGGTTGAGGCACAGTTCCATC
M071_Pa-Bb-fw	GATGGAACTGTGCCTCAACCATGCGGAGAATCTTTATTTTCAGGGCCATCACCA
M072_Pa-Bb-rev	GGGTAACCGGGGTGATGCTCGTCATGGTATATCTCCTTATTAAAGTTAAACAAAA
M017_AcAv_TEVrepair_fw	GGCCATCACCATCACCATCACCATCAC
M018_Ac_TEVrepair_rev	CTGAAAATAAAGATTCTCATTATGACGC
M019_Av_TEVrepair_rev	CTGAAAATAAAGATTCTCATTATGGCGC
Primers for Gibson cloning of bathy-PCMs into pREDusk	
M009_Ac_Bb_for	CCTGGTTGAAATTGTGCTGCGTCATAATCTCCAGGAACTGCAATCCGAGCTCGTC
M010_Ac_Bb_rev	TTGTCAGATCAACTGCTTCTGCATTCGTCATGCGTGGGCGACCTCAGGC
M011_Ac_Ins_for	GCCTGAGGTCGCCCACGCATGACGAATGCAGAAGCAGTTGATCTGACAA
M012_Ac_Ins_rev	GACGAGCTCGGATTGCAGTTCCTGGAGATTATGACGCAGCACAATTTCAACC

	AGG
M013_Av_Bb_for	CCGTTAAAGTGGTTCTGCGCCATAATCTCCAGGAACTGCAATCCGAGCTCGTC
M014_Av_Bb_rev	GTCAGATCAACGGTCTGGGTTGCAGTCATGCGTGGGCGACCTCAGGC
M015_Av_Ins_for	GCCTGAGGTCGCCCACGCATGACTGCAACCCAGACCGTTGATCTGAC
M016_Av_Ins_rev	GACGAGCTCGGATTGCAGTTCCTGGAGATTATGGCGCAGAACCACTTTAACGG

Primers for cloning of PATCHY *Ac*/*Av*NIRusk start constructs

A07_P_I_II_Step1_for	TTAACCGAGCATCAGCAGACACAAGCACGTCTCCAGGAACTGCAATCCGAGCTCG
M020_AcREDusk_PATCHY_rev1	CGCTCGTCAGCCAGTAATTCATTATGACGCAGCACAATTTCAACC
M021_AcREDusk_PATCHY_rev2	GACGCACGTCTGCTTTCCCGCGCTCGTCAGCCAGTAATTCATTAT
M022_AcREDusk_PATCHY_fw2	AGCGTCTTCTTAATGAAGTTAACCGAGCATCAGCAGACACAAG
M023_AvREDusk_PATCHY_rev1	CGCTCCTCTTGCATCAGTTCATTATGGCGCAGAACCACTTTAACG
M025_AvREDusk_PATCHY_fw2	AGCGTGTCTTAAACGAGGTTAACCGAGCATCAGCAGACACAAG
M028_AvREDusk_PATCHY_rev2	GACGAATGTCTGCTTTGTTACGCTCCTCTTGCATCAGTTCATTAT

PATCHY forward primers for amplification of FixL

A14_P_I_II_for_1	ACCGAGCATCAGCAGAC
A15_P_I_II_for_2	GAGCATCAGCAGACACAAG
A16_P_I_II_for_3	CATCAGCAGACACAAGC
A17_P_I_II_for_4	CAGCAGACACAAGCACG
A18_P_I_II_for_5	CAGACACAAGCACGTCTC
A19_P_I_II_for_6	ACACAAGCACGTCTCC
A20_P_I_II_for_7	CAAGCACGTCTCCAGG
A21_P_I_II_for_8	GCACGTCTCCAGGAAC
A22_P_I_II_for_9	CGTCTCCAGGAACTGC
A23_P_I_II_for_10	CTCCAGGAACTGCAATCC
A24_P_I_II_for_11	CAGGAACTGCAATCCG
A25_P_I_II_for_12	GAACTGCAATCCGAGC
A26_P_I_II_for_13	CTGCAATCCGAGCTCG
A27_P_I_II_for_14	CAATCCGAGCTCGTCC
A28_P_I_II_for_15	TCCGAGCTCGTCCACG

A29_P_I_II_for_16	GAGCTCGTCCACGTCTC
A30_P_I_II_for_17	CTCGTCCACGTCTCCAG
A31_P_I_II_for_18	GTCCACGTCTCCAGGC
A32_P_I_II_for_19	CACGTCTCCAGGCTGAG
A33_P_I_II_for_20	GTCTCCAGGCTGAGCG
A34_P_I_II_for_21	TCCAGGCTGAGCGCCATG
A35_P_I_II_for_22	AGGCTGAGCGCCATGG
A36_P_I_II_for_23	CTGAGCGCCATGGGCG
A37_P_I_II_for_24	AGCGCCATGGGCGAAATG
A38_P_I_II_for_25	GCCATGGGCGAAATGG
A39_P_I_II_for_26	ATGGGCGAAATGGCGTC
A40_P_I_II_for_27	GGCGAAATGGCGTCCG
A41_P_I_II_for_28	GAAATGGCGTCCGCGC
A42_P_I_II_for_29	ATGGCGTCCGCGCTCG

PATCHY forward primers for amplification of TtrS

B92_P_Ttr_for_01	CGCGAACTGATTAATACCC
B93_P_Ttr_for_02	GAACTGATTAATACCCAGCG
B94_P_Ttr_for_03	CTGATTAATACCCAGCGTCAG
B95_P_Ttr_for_04	ATTAATACCCAGCGTCAGC
B96_P_Ttr_for_05	AATACCCAGCGTCAGC
B97_P_Ttr_for_06	ACCCAGCGTCAGCTGAATG
B98_P_Ttr_for_07	CAGCGTCAGCTGAATG
B99_P_Ttr_for_08	CGTCAGCTGAATGAAAATCG
C01_P_Ttr_for_09	CAGCTGAATGAAAATCGTG
C02_P_Ttr_for_10	CTGAATGAAAATCGTGCACTG
C03_P_Ttr_for_11	AATGAAAATCGTGCACTGC
C04_P_Ttr_for_12	GAAAATCGTGCACTGCTG
C05_P_Ttr_for_13	AATCGTGCACTGCTGG
C06_P_Ttr_for_14	CGTGCACTGCTGGAAC
C07_P_Ttr_for_15	GCACTGCTGGAACATGC
C08_P_Ttr_for_16	CTGCTGGAACATGCACAG
C09_P_Ttr_for_17	CTGGAACATGCACAGC
C10_P_Ttr_for_18	GAACATGCACAGCGTATTG
C11_P_Ttr_for_19	CATGCACAGCGTATTGC
C12_P_Ttr_for_20	GCACAGCGTATTGCCATTG

C13_P_Ttr_for_21	CAGCGTATTGCCATTGC
C14_P_Ttr_for_22	CGTATTGCCATTGCCG
C15_P_Ttr_for_23	ATTGCCATTGCCGGTG
C16_P_Ttr_for_24	GCCATTGCCGGTGAAC
C17_P_Ttr_for_25	ATTGCCGGTGAACTGG
C18_P_Ttr_for_26	GCCGGTGAACTGGGTG
C19_P_Ttr_for_27	GGTGAACTGGGTGCAAG
C20_P_Ttr_for_28	GAACTGGGTGCAAGCC
C21_P_Ttr_for_29	CTGGGTGCAAGCCTGAG

PATCHY forward primers for amplification of TodS

C28_P_Tod_for_01	ACTGAGAAGAAACAAGCACAGG
C29_P_Tod_for_02	GAGAAGAAACAAGCACAGG
C30_P_Tod_for_03	AAGAAACAAGCACAGGAAAATC
C31_P_Tod_for_04	AAACAAGCACAGGAAAATCTTAACC
C32_P_Tod_for_05	CAAGCACAGGAAAATCTTAACC
C33_P_Tod_for_06	GCACAGGAAAATCTTAACCAG
C34_P_Tod_for_07	CAGGAAAATCTTAACCAGTTGC
C35_P_Tod_for_08	GAAAATCTTAACCAGTTGCAGC
C36_P_Tod_for_09	AATCTTAACCAGTTGCAGCAAC
C37_P_Tod_for_10	CTTAACCAGTTGCAGCAACAAC
C38_P_Tod_for_11	AACCAGTTGCAGCAACAAC
C39_P_Tod_for_12	CAGTTGCAGCAACAACTTG
C40_P_Tod_for_13	TTGCAGCAACAACTTGTG
C41_P_Tod_for_14	CAGCAACAACTTGTGTACG
C42_P_Tod_for_15	CAACAACTTGTGTACGTTTCC
C43_P_Tod_for_16	CAACTTGTGTACGTTTCCC
C44_P_Tod_for_17	CTTGTGTACGTTTCCCGATC
C45_P_Tod_for_18	GTGTACGTTTCCCGATCAG
C46_P_Tod_for_19	TACGTTTCCCGATCAGC
C47_P_Tod_for_20	GTTTCCCGATCAGCTACG
C48_P_Tod_for_21	TCCCGATCAGCTACGATG
C49_P_Tod_for_22	CGATCAGCTACGATGGG
C50_P_Tod_for_23	TCAGCTACGATGGGTGAATTTG
C51_P_Tod_for_24	GCTACGATGGGTGAATTTG
C52_P_Tod_for_25	ACGATGGGTGAATTTGC

C53_P_Tod_for_26	ATGGGTGAATTTGCAGC
C54_P_Tod_for_27	GGTGAATTTGCAGCCTATATTG
C55_P_Tod_for_28	GAATTTGCAGCCTATATTGC
C56_P_Tod_for_29	TTTGCAGCCTATATTGCACAC

PATCHY reverse primers for amplification of *Ac*PCM

M029_Ac_rev1	TTCATTAAGAAGACGCTGACG
M030_Ac_rev2	ATTAAGAAGACGCTGACGC
M031_Ac_rev3	AAGAAGACGCTGACGC
M032_Ac_rev4	AAGACGCTGACGCACG
M033_Ac_rev5	ACGCTGACGCACGTCTG
M034_Ac_rev6	CTGACGCACGTCTGCTTTC
M035_Ac_rev7	ACGCACGTCTGCTTTCC
M036_Ac_rev8	CACGTCTGCTTTCCCG
M037_Ac_rev9	GTCTGCTTTCCCGCGC
M038_Ac_rev10	TGCTTTCCCGCGCTCG
M039_Ac_rev11	TTTCCCGCGCTCGTCAG
M040_Ac_rev12	CCCGCGCTCGTCAGCC
M041_Ac_rev13	GCGCTCGTCAGCCAGTAATTC
M042_Ac_rev14	CTCGTCAGCCAGTAATTCATTATG
M043_Ac_rev15	GTCAGCCAGTAATTCATTATGACG
M044_Ac_rev16	AGCCAGTAATTCATTATGACG
M045_Ac_rev17	CAGTAATTCATTATGACGCAGC
M046_Ac_rev18	TAATTCATTATGACGCAGCAC
M047_Ac_rev19	TTCATTATGACGCAGCAC
M048_Ac_rev20	ATTATGACGCAGCACAATTTC

PATCHY reverse primers for amplification of *Av*PCM

M049_Av_rev1	CTCGTTTAAGACACGCTGAC
M050_Av_rev2	GTTTAAGACACGCTGACGAATG
M051_Av_rev3	TAAGACACGCTGACGAATG
M052_Av_rev4	GACACGCTGACGAATGTC
M053_Av_rev5	ACGCTGACGAATGTCTGC
M054_Av_rev6	CTGACGAATGTCTGCTTTG
M055_Av_rev7	ACGAATGTCTGCTTTGTTACG
M056_Av_rev8	AATGTCTGCTTTGTTACGC
M057_Av_rev9	GTCTGCTTTGTTACGCTCC

M058_Av_rev10	TGCTTTGTTACGCTCCTC
M059_Av_rev11	TTTGTTACGCTCCTCTTGC
M060_Av_rev12	GTTACGCTCCTCTTGCATC
M061_Av_rev13	ACGCTCCTCTTGCATCAG
M062_Av_rev14	CTCCTCTTGCATCAGTTCATTATG
M063_Av_rev15	CTCTTGCATCAGTTCATTATGG
M064_Av_rev16	TTGCATCAGTTCATTATGGC
M065_Av_rev17	CATCAGTTCATTATGGCGC
M066_Av_rev18	CAGTTCATTATGGCGCAG
M067_Av_rev19	TTCATTATGGCGCAGAACC
M068_Av_rev20	ATTATGGCGCAGAACC

Primers for cloning of orthogonal TtrSR TCS

B55_Pttr_BglII_for	CGTTAAAGATCTATATTTGTTGCGCTAGA
B56_Pttr_XbaI_rev	TGCCAGTCTAGAACAGGCATTC
B57_Bb_for_DmTtrSR	ACGTCTGCGTCTGGAAGATAATCGTTAACAATTGATGTAAGTTAGCTCACTCA
B58_Bb_rev_DmTtrSR	CATTCAGCTGACGCTGGGTATTAATCAGCTCACGATTAGCACGCTCTAAG
B59_Ins_for_DmTtrSR	CTTAGAGCGTGCTAATCGTGAGCTGATTAATACCCAGCGTCAGCTGAATG
B60_Ins_rev_DmTtrSR	TGAGTGAGCTAACTTACATCAATTGTTAACGATTATCTTCCAGACGCAGACGT
B84_P_V_sc_for	GGCGTTTCGCCTTTGATATCGCGAACTGATTAATACCCAGCGTCAG
B85_P_V_sc_rev	GCAACTCTGCGTTACGCCCTTCAAGCTCACGATTAGCACGCTCTA

Primers for cloning of orthogonal TodST TCS

TodX_fw	GCACAAGATCTGGTCTGAGGTTTTCATCGAC
TodX_rv	CGTGCTCTAGAAATTACAATCCTTCCAC
0_pDusk_fw	GAGTCAATTGAGGGTGGTGAATGTGGCTAG
1_TodS_fw	GATATCACCGAGAAGAAACAAGCACAGGAAAATC
2_NoMfeI_rv	CGTACACAAGTTGTTGCTGCAACTGGTTAAGATTTTCCTGTGCTTGTTTC
3_TodS_fw	GCAGCAACAACTTGTGTACGTTTCCCGATC
4_TodT_rv	GCATCAATTGCTATTCCAGGCTATCCTTGAG
5_pDusk_rv	GCTTGTTTCTTCTCGGTGATATCATTCTGAATTC
M014_Av_Bb_rev	GTCAGATCAACGGTCTGGGTTGCAGTCATGCGTGGGCGACCTCAGGC
M015_Av_Ins_for	GCCTGAGGTCGCCCACGCATGACTGCAACCCAGACCGTTGATCTGAC
C26_Bb_for_P_X_sc	ATTCGTCAGCGTGTCTTAAACGAGGTTAACTGAGAAGAAACAAGCACAG
C27_Ins_rev_P_X_sc	CTGTGCTTGTTTCTTCTCAGTTAACCTCGTTTAAGACACGCTGACGAAT

Primers for cloning StrR variants

B73_Bb_for_pStrepDusk	GATCACCAAGGTAGTCGGCAAATAAGAATTAATTCATGAGCGGATACATATT
B74_Bb_rev_pStrepDusk	AAGGATCTCAAGAAGATCCTTTGATCTGATGCCTCCGTGTAAGGGG
B75_Ins_for_pStrepDusk	CCCCTTACACGGAGGCATCAGATCAAAGGATCTTCTTGAGATCCTT
B76_Ins_rev_pStrepDusk	AATATGTATCCGCTCATGAATTAATTCTTATTTGCCGACTACCTTGGTGATC
B26_Ins_rev	CGAGATGATAGGAGGTCTAGCATGACGACCAAGGGACATATCTACG
B27_Bb_for	CGTAGATATGTCCCTTGGTCGTCATGCTAGACCTCCTATCATCTCG
M014_Av_Bb_rev	GTCAGATCAACGGTCTGGGTTGCAGTCATGCGTGGGCGACCTCAGGC
M015_Av_Ins_for	GCCTGAGGTCGCCCACGCATGACTGCAACCCAGACCGTTGATCTGAC

Primers for cloning MCS variant

C76_Bb_for_MCS	ATCAGCAACTACATGAAGGGCTCGCGGCGGCTGCTTGCCG
C77_Ins_rev_MCS	CGGCAAGCAGCCGCCGCGAGCCCTTCATGTAGTTGCTGAT
M014_Av_Bb_rev	GTCAGATCAACGGTCTGGGTTGCAGTCATGCGTGGGCGACCTCAGGC
M015_Av_Ins_for	GCCTGAGGTCGCCCACGCATGACTGCAACCCAGACCGTTGATCTGAC

Primers for cloning lux variants

lux-op_Nde_for	**GGATCC<u>CATATG</u>** AAATTTGGAAACTTTTTGCTTACATACC
lux-op_Sac_rev	**TGAC<u>GAGCTC</u>** TCAACTATCAAACGCTTCG
PfixK2_NdeI_for	**TGAC<u>CATATG</u>** ATATCTCCTTCTTAAAGTTAAACAAAATT
PfixK2_rev_OLPmamG	**ACTAAGAGCTAGTAAAGCGAAAAAG** CGGGATCTCGACGCTCTC
DrHo_OLPmamG_for	**CTTTTTCGCTTTACTAGCTCTTAGTTCTCCAATAAATTCCCTGCGAAGCTTAGGAGATCAGTATATG** CTCCGACTCATGATCATGAAGC
FixJ_NotI_rev	**TGAC<u>GCGGCCGC</u>** TCAATCGTTGAGCATGCC

Supplementary Table 2

Plasmids used in this study.

Id	Construct	Reference
pBAM2	*p15A* ori, miniTn5, Amp^R, Kan^R	Uebe, unpublished
pBAM2-luxAE	*p15A* ori, miniTn5, Amp^R, Kan^R, *luxABCDE*	(Dziuba et al., 2021)
p*Dm*REDlux	*p15A* ori, miniTn5, Amp^R, Kan^R, PfixK2-*luxABCDE*, PmamDC-*DrhemO-Dmpcm-fixL*	(this study)
p*Av*NIRlux	*p15A* ori, miniTn5, Amp^R, Kan^R, PfixK2-*luxABCDE*, PmamDC-*DrhemO-Avpcm-fixL*	(this study)
pMH005	*Ac*NIRusk-0	(this study)
pMH006	*Av*NIRusk-0	(this study)
pMH008	pCDF-Duet-*Ac*PCM/HO	(this study)
pMH009	pCDF-Duet-*Av*PCM/HO	(this study)
pMH013	*Ac*NIRusk-PATCHY-start construct	(this study)
pMH014	*Av*NIRusk-PATCHY-start construct	(this study)
pMH017	*Ac*NIRusk+16a	(this study)
pMH018	*Ac*NIRusk+16b	(this study)
pMH019	*Ac*NIRusk-1a (*Ac*NIRusk)	(this study)
pMH020	*Av*NIRusk+16b	(this study)
pMH021	*Ac*NIRusk-4	(this study)
pMH023	*Av*NIRusk-1	(this study)
pMH026	pCDF-Duet-*Pa*PCM/HO	(this study)
pMH028	*Av*NIRusk+17a (*Av*NIRusk)	(this study)
pMH029	*Av*NIRusk+16a	(this study)
pMH030	*Av*NIRusk-5	(this study)
pMH034	*Ac*NIRusk-1b	(this study)
pMH038	*Av*NIRusk+17b	(this study)
pMIR66	TodS and TodT proteins	(Ramos-González et al., 2002)
pMIR77	PtodX promoter	(Ramos-González et al., 2002)

pQX037	*Pa*PAC	(Xu et al., 2024)
pQX086	*Av*PAC	(Xu et al., 2024)
pQX091	*Ac*PAC	(Xu et al., 2024)
pRD349	pDusk-YT1-X	(unpublished)
pSM001	*Dr*REDusk-MCS	(Multamäki et al., 2022)
pSM003	*Dm*REDusk	(Meier et al., 2024a)
pSM024	*Dm*REDusk-MCS	(Meier et al., 2024a)
pSM087	*Dm*DERusk	(Meier et al., 2024a)
pSM090	*Dm*DERusk-YPet	(Meier et al., 2024a)
pSM091	*Dm*DERusk-MCS	(Meier et al., 2024a)
pSM122	*Dm*DERusk-StrR-YPet	(this study)
pSM123	*Dm*Ttr-REDusk-PATCHY-start construct	(this study)
pSM128	*Av*Tod-NIRusk PATCHY-start construct	(this study)
pSM136	*Av*NIRusk-StrR-YPet	(this study)
pSM146	*Dm*Ttr-13	(this study)
pSM147	*Dm*Ttr-0	(this study)
pSM148	*Dm*Ttr+5	(this study)
pSM149	*Dm*Ttr+7a	(this study)
pSM150	*Dm*Ttr+7b	(this study)
pSM153	*Dm*Ttr+14	(this study)
pSM160	*Av*Tod+3	(this study)
pSM161	*Av*Tod+16	(this study)
pSM162	*Av*Tod+19	(this study)
pSM163	*Av*Tod+21	(this study)
pSM164	*Av*Tod+26	(this study)
pSM182	*Av*NIRusk-MCS	(this study) Addgene: 235084