eRNAkit: Expanding the Functional Atlas of human Enhancer RNAs Beyond the Nucleus

Natalia Benova[1*], Rene Kuklinkova[1*], Mahmoud K. Eldahshoury[1], Chinedu A. Anene[1 #]

1. Centre for Biomedical Science Research, School of Health, Leeds Beckett University,
Leeds, LS1 3HE, United Kingdom
* These authors contributed equally.
# Corresponding author. Centre for Biomedical Science Research, School of Health, Leeds
Beckett University, Leeds LS1 3HE, United Kingdom. E-mail: c.a.anene@leedsbeckett.ac.uk

8
9
Abstract

Enhancer RNAs (eRNAs) are a class of non-coding RNAs transcribed from active enhancers
that regulate various aspects of transcription. Although traditionally viewed as nuclear-
localised and unstable transcripts, large transcriptomic studies reveal they vary widely in
localisation and biochemical properties, including detectable accumulation in cytoplasmic
compartments. These observations suggest potential non-nuclear functions for eRNAs, yet
existing databases remain focused on nuclear, cis-regulatory roles, limiting systematic
exploration of their broader regulatory repertoire. Here we present **eRNAkit**, a
comprehensive and accessible resource designed to address this gap by integrating
subcellular localisation, RNA-RNA interaction and expression data for annotated eRNAs.
Leveraging fractionation-based RNA-Seq datasets, eRNAkit profiles eRNA distribution across
nuclear and cytoplasmic compartments. It incorporates gene expression data spanning
major human organs and primary cell types, enabling tissue-specific analysis of eRNA
function. Crucially, eRNAkit includes experimentally derived RNA-RNA interaction data from
RIC-Seq, PARIS, and KARR-Seq, supporting exploration of trans-acting and cytoplasmic roles
for eRNAs grounded in physical interaction evidence. eRNAkit expands current eRNA
resources beyond the enhancer-promoter paradigm, offering a robust platform for
dissecting non-canonical functions of eRNAs and advancing our understanding of their full
regulatory potential in human biology. The eRNAkit resource is available to download at
https://github.com/AneneLab/eRNAkit.

30
Introduction
eRNAs are a class of non-coding RNAs transcribed from active enhancer regions [1]. They
regulate gene expression by activating transcription, bridging enhancer-promoter
interactions, and remodelling chromatin [2]. Although initially considered nuclear, unstable
and non-polyadenylated, transcriptomic data — including those from the ENCODE
consortium, reveal that eRNAs vary widely in both biochemical properties and cellular
localisation [3,4]. While most studies report nuclear enrichment based on relative abundance,
they also detect eRNAs in cytoplasmic fractions, raising the question of what functions these
cytoplasmic eRNAs might perform — and which molecular targets they may regulate
outside the nucleus.
41
Existing eRNA databases predominantly emphasise *cis*-regulatory mechanisms, reflecting
the prevailing model in which enhancers influence the expression of nearby genes through

spatial proximity. Resources such as eRNAbase, HeRA, Anima-eRNAdb, eRic and GPIeR [5–9]
systematically catalogue eRNA-target relationships based on genomic co-localisation,
chromatin conformation, or correlative expression patterns. While these approaches have
illuminated many aspects of enhancer biology, their design inherently favours local
interactions and overlook more distal, cytoplasmic effects. Furthermore, these databases
largely ignore the localisation of eRNAs, an essential factor in understanding their functional
potential. Nevertheless, mounting evidence suggests that certain eRNAs can function
beyond their sites of origin, modulating the expression of distant genes or engaging in post-
transcriptional regulation. Some studies have uncovered eRNAs that act in *trans*, regulating
gene expression far from their sites of origin. For example, a distal regulatory region eRNA
transcribed from the MYOD1 locus mediates cohesion recruitment and Myogenin gene
expression in trans to control myogenic differentiation [10]. Similarly, an eRNA transcribed
from the KLK3 locus enhance androgen receptor-dependent gene expression in human
prostate cancer [11]. In both examples, the eRNAs were found to be polyadenylated, which is
contrary to the standard model of eRNA processing. In a different study, knockdown of two
p53-induced enhancer RNAs, p53BER2 and p53BER4, impaired the induction of distant
target genes in response to p53 activation [12]. This regulation occurred despite the absence
of detectable p53 binding at the promoters of these target genes, suggesting that the eRNAs
act independently of local enhancer–promoter contact. Moreover, given that siRNA
knockdowns used to validate most of these observations are more effective for cytoplasm
localised RNAs [13], these observations raise the possibility that some of these effects is
through post-transcriptional mechanisms, such as mRNA stabilisation or miRNA sponging,
rather than classical nuclear transcriptional activation.
67
Therefore, these findings may support a broader model in which cytoplasmic functions may
contribute significantly to gene regulation and function. However, a comprehensive
understanding of their subcellular distribution remains incomplete, and the functional
significance of eRNAs beyond the nuclear compartment, particularly in the cytoplasm, has
yet to be systematically explored. A centralised and accessible resource cataloguing eRNA
localisation across subcellular compartments and mapping potential eRNA-eRNA
interactions is currently lacking. This gap hampers efforts to distinguish nuclear-restricted
enhancer activity from putative post-transcriptional functions and limits our ability to
interrogate the full functional repertoire of eRNAs in gene expression control.

To address this gap, we developed **eRNAkit**, a comprehensive and user-friendly resource for
exploring eRNA subcellular localisation, expression, and cytoplasmic functions. eRNAkit
integrates publicly available fractionation-based transcriptomic datasets to systematically
profile eRNA distribution across nuclear and cytoplasmic compartments. It also incorporates
expression data across major human organs and primary cell types, allowing users to assess
the tissue and cell-type specificity of individual eRNAs. Additionally, eRNAkit includes
experimentally derived RNA-RNA interaction data from technologies such as RIC-Seq, PARIS,
and KARR-Seq, enabling users to explore putative eRNA-eRNA interactions grounded in
physical proximity and hybridisation evidence. By combining localisation, expression, and
interaction data, eRNAkit offers a powerful framework to interrogate the full regulatory
potential of eRNAs beyond the classical enhancer–promoter paradigm.

Results

eRNAkit Overview

eRNAkit represents the first comprehensive resource specifically designed to investigate the
cytoplasmic functions of eRNAs (Figure 1). It integrates a suite of analytical workflows for
processing and analysing eRNA-related data, facilitating tasks such as data manipulation,
statistical analysis, and visualisation. To assist users, eRNAkit provides a transcriptome-wide
landscape of eRNA subcellular localisation, eRNA-mRNA interactions and eRNA-ribosome
associations, thereby elucidating their function beyond the nucleus. Compared to existing
resources, eRNAkit offers the most extensive collection of cytoplasmic eRNA-specific
functional information and analytical methods (Table 1), enabling a comprehensive
interrogation of eRNA functions. To facilitate exploration of the database, it includes a user-
friendly Rshiny interface.

Table 1: Leading eRNA resources for comparison to eRNAkit

	Subcellular localisation	mRNA interaction	Ribosome interaction	Target assignment
eRNAkit	Nucleus, Cytosol, Nucleolus, Chromatin, Nucleoplasm, Membrane, Insoluble cytoplasmic fraction	KARR-Seq, RIC-Seq, PARIS	Ribo-Seq, Polysome-Seq	Physical eRNA-mRNA (TAD used to differentiate nuclear from cytoplasmic)
eRNAbase	None	RIC-Seq	None	Genomic distance (multiple thresholds)
HeRA	None	None	None	Genomic distance (1MB) and Correlation (Rho > 0.3)
Anima — eRNAdb	None	None	None	Genomic distance (1MB) and Correlation (Rho > 0.3)
eRic	None	None	None	Genomic distance (1MB)
GPIeR	None	None	None	Genomic distance (1MB)

eRNAkitApp

Extracting relevant information for a few eRNAs from the transcriptome-wide database can
be tedious due to the complexity of navigating multiple interlinked tables. Thus, we build an
interactive Rshiny UI called eRNAkitApp. The interface includes two core components:
search panel and interactive visualisation modules. The search panel allows users to query
the database by eRNA ID, genomic coordinates, or Ensemble gene ID, returning all relevant
records. The visualisation modules include subcellular localisation (Supplementary Figure
S1b), tissue/cell expression and specificity (Supplementary Figure S1c), eRNA-mRNA
interactions (Supplementary Figure S1a) and eRNA-ribosome associations. Finally, the
application offers straightforward results download option, allowing users to export search
results tables. Additionally, the full database is also available for download for integration
into other workflows. Collectively, this tool serves as a powerful platform for researchers to
interactively explore and analyse eRNA-related data.

Data Summary
We obtained 46,650 consensus eRNAs by integrating enhancer annotations from five large
databases (Methods, Figure 1a). The median length of these eRNAs was 992 bp, with an
interquartile range of 238–3,241 bp, and the longest extending up to 3,198,781 bp. We
found 11 extremely large regions (> 300, 000 bp), which are derived from overlapping super-
enhancers in the source databases. Across tissues and primary cells, 81.19% of the eRNAs
were detected at a >1 CPM threshold (Supplemental Figure 1a-b). The number of detectable
eRNAs ranged from 5,209 in the liver to 13,444 in the spinal cord for tissues, and from 1,957
in astrocytes to 20,688 in bladder smooth muscle cells (SMCs) for primary cells. Notably,
SMCs and fibroblast populations accounted for the majority of the top 10 cell types. All
tissues and cell types had at least one specific eRNA, with the vagina showing the highest
number among organs (2,383) and mononuclear cells having the most among cell types
(1,202).
130
Unlike protein-coding genes, the functional output of eRNAs is the RNA molecule itself. As
such, their regulatory potential is closely linked to their subcellular localisation and physical
interactions with other RNAs. Defining these spatial and molecular contexts is essential for
understanding the mechanisms by which eRNAs exert their functions. As expected, more
eRNAs are detectable in the nucleus, regardless of polyadenylation status (Figure 2b).
However, polyadenylation influences the number of eRNAs detectable in the cytoplasm,
potentially through controlling eRNA stability and transport. Analysis of the cytoplasm-
nucleus REI distribution across cell types and RNA classes reveals a trimodal pattern, with
subsets of eRNAs enriched in the nucleus, in the cytoplasm, or showing no strong bias
(Figure 2c). While nuclear enrichment is more common, a substantial proportion of eRNAs
are cytoplasm enriched. In the two cell lines with subcytoplasmic fractionation data, we
observe widespread localisation of eRNAs in both the insoluble (organelle-associated) and
membrane fractions (Figure 2d).
144
Our curated dataset of eRNA-mRNA interactions identified 82,587 pairs present in at least
two samples, encompassing 25.28% (11,793 out of 46,650) of the annotated consensus
eRNAs (Figure 3a). Given the conservative nature of our filtering, this number is expected to
increase with broader dataset inclusion. Surprisingly, correlation analysis between eRNA
length and the number of interactions revealed a significant weak negative correlation (rho
$= -0.207$, $p < 2.2 \times 10^{-16}$; $S = 3.30 \times 10$, Figure 2a). This suggests that longer eRNAs may be
slightly less likely to engage in RNA-RNA interactions. However, the low correlation
coefficient indicates that factors beyond transcript size-such as sequence features,
localisation, or binding affinity-likely contribute to their interaction potential. Indeed,
stratified analysis of these interactions across diverse cellular perturbations reveals that
some are dependent on active translation or RNA-binding proteins (Figure 3b). Notably,
similar perturbations-such as translation inhibition by cycloheximide or harringtonine, and
viral infection by RSV or VSV-lead to consistent alterations in the RNA-RNA interaction
landscape (Figure 3b). Additionally, only 28.7% (23, 703 out of 82, 587) of the mRNA were
located within the same TAD or within 5kb of their corresponding eRNA pair (Methods).
Interestingly, 30% of all the interactions pairs occur between eRNAs and mRNAs located on
different chromosomes. These observations suggest that most eRNA-mRNA interactions do
not adhere to the canonical *cis*-regulatory function of eRNAs.
163

Case study

Potential miRNA sponging by eRNAs

An established cytoplasmic function of several classes of non-coding RNAs is the ability to act as miRNA sponges — sequestering miRNAs and preventing them from repressing their target mRNAs. This mechanism has been widely described for circular RNAs and long non-coding RNAs [14,15]. Given this context, it is plausible that eRNAs, especially those with stable cytoplasmic presence — may also function as miRNA sponges. This possibility has not been systematically explored, but the pervasive localisation of eRNAs hints at such roles. To illustrate this, we examined the eRNA targets of miR130b and miR96, both of which have been implicated in diverse cellular processes, including metabolism and tissue development. miR-130B has been shown to regulate adipogenesis [16], while miR-96 is involved in neuronal cell differentiation [17]. Using eRNAkitApp, we first input their Ensemble IDs "ENSG00000283871, ENSG00000199158" and selected **TARGET** (Figure S1a). The results showed two distinct clusters of target eRNAs, indicating potential differences in function. Then we selected the **LOCALISATION** tab, which revealed that most of the associated eRNAs are detectable in the cytoplasm and sub-cytoplasmic fractions (Figure S1b). Given the strong cytoplasmic localisation of these eRNAs in this context, and the precedence of miRNA sponging among other cytoplasmic non-coding RNAs, we proposed that miRNA sequestration may be a functional mechanism for these eRNAs. To investigate this possibility, we examined changes in eRNA expression in response to miR-130 overexpression in mesenchymal stem cells (GSE202135) and miR-96 knockdown in prostate epithelial cells (GSE255227). Following miR-130 overexpression (by mimics), 60% (12 out of 20) of the detectable (CPM > 1 in > 2 samples) eRNA targets were downregulated (log2FC < -0.5), a proportion markedly higher than expected by chance (empirical p = 0.0, Figure 3c). By contrast, all detectable eRNA targets of miR-96 (6 out of 6) were upregulated (log2FC > 0.5) upon miR-96 knockdown (by siRNA), also surpassing random expectation (empirical p = 0.0019, Figure 3d). These findings suggest that, like other cytoplasmic ncRNAs, eRNAs may contribute to post-transcriptional regulation by modulating miRNA availability and illustrate the biological value of the eRNAkit.

Inter-organelle communication through eRNAs

Bi-directional communication between the nucleus and mitochondria is critical for maintaining cellular function [18]. Emerging research suggests that RNA-mediated nuclear signal can influence mitochondrial biology [19]. We identified an eRNA, "en135763" located on chromosome 5, which is altered under various cellular stress conditions (data not shown), and searched eRNAkitApp using the eRNA ID option (Figure S3a). In the **LOCALISATION** tab, results showed that this eRNA exhibits comparable expression levels in the cytoplasm and nucleus. It is also enriched in the organelle-associated fraction in K562 cells and has eRNA-RNA interactions with mitochondrial encoded genes (Figure S3b), indicating that eRNAkit could support research into RNA mediated inter-organelle communication. The **EXPRESSION** tab revealed that this eRNA is widely expressed across organs and primary cell types, with notably higher levels in T-cell populations, consistent with the involvement of immune cells in the stress response. These findings highlight the underappreciated value of eRNA subcellular localisation and physical interactions.

Discussion

eRNAs have long been considered nuclear-localised and unstable byproducts of enhancer
activation, with functions largely confined to cis-regulatory roles at nearby promoters.
However, emerging data challenge this restricted view, revealing that eRNAs are more
biochemically and functionally diverse than previously appreciated. Unlike protein-coding
genes, the final product of eRNAs is RNA itself. Consequently, eRNA function relies on the
physical properties and interactions of the RNA molecule, which are inherently dependent
on spatial proximity. Here, we present eRNAkit, a comprehensive resource that consolidates
subcellular localisation, expression specificity, and RNA-RNA interaction data to
systematically map the full regulatory repertoire of eRNAs, including potential trans-acting
and post-transcriptional functions. Compared to existing databases, eRNAkit offers several
advances. Most notably, it is the first to integrate larger scale subcellular localisation data
alongside physical RNA-RNA interactions, thereby facilitating functional hypotheses that
extend beyond proximity-based enhancer-promoter models.
223
A key component of eRNAkit is the transcriptome-wide landscape of eRNA localisation
across subcellular compartments. Our REI-based analyses demonstrate that while nuclear
enrichment is prevalent, a considerable proportion of eRNAs are detectable — and in some
cases, enriched in cytoplasmic and subcytoplasmic fractions. This observation supports prior
reports suggesting the presence of cytoplasmic eRNAs [3] and offers a transcriptome-wide
perspective that challenges the long-standing assumption of nuclear restriction. Notably,
the influence of polyadenylation on cytoplasmic localisation suggests that RNA processing
features may be critical determinants of eRNA stability and trafficking, potentially mirroring
the behaviour of other non-coding RNAs with post-transcriptional roles, such as lncRNAs
and circRNAs. eRNAkit also highlights strong cell- and tissue-specific expression patterns of
eRNAs, consistent with the known context-dependence of enhancer activity [1,20]. The
presence of highly specific eRNAs in reproductive organs and in immune cells highlight their
potential roles in tissue-specialised gene regulatory programs. Such specificity may be
leveraged in future studies to prioritise candidate eRNAs for functional validation or
biomarker development.
239
The inclusion of tens of thousands of robust eRNA-mRNA interactions further expands the
functional landscape of eRNAs. These interactions, derived from multiple RNA-RNA
interactome technologies (RIC-Seq, PARIS, KARR-Seq), provide direct evidence of physical
proximity between eRNAs and potential target transcripts. Importantly, a substantial
fraction of these interactions is translation or RBP-dependent, suggesting that eRNAs may
engage in post-transcriptional regulation through mechanisms such as mRNA stabilisation,
translational control, or miRNA competition. The weak negative correlation between eRNA
length and interaction count also points to a nuanced interaction landscape that may be
shaped more by structure, binding motifs, or localisation than by transcript size alone.
249
Despite its utility, several limitations should be noted. First, while the use of multiple
datasets enhances confidence in reported interactions and expression patterns, coverage
across all cell types and compartments remains incomplete. Second, although physical
interactions provide strong evidence for potential function, mechanistic validation is
essential to determine whether these interactions are regulatory, structural, or stochastic.
Lastly, the dynamic regulation of eRNA expression and localisation — particularly under
stress, development, or disease contexts — remains to be fully characterised and could

provide further insight into the functional adaptability of eRNAs. To address these
limitations, we plan to update eRNAkit regularly by incorporating additional subcellular
localisation data and functional eRNA interaction profiles as more high-throughput datasets
become available. Due to our work in cellular senescence, we are also building an extension
to characterise the expression profiles of these eRNAs in cellular stress response. We are
confident that eRNAkit provides a critical first step toward uncovering the full regulatory
potential of eRNAs and will serve as a foundational tool for advancing our understanding of
their diverse roles in human health and disease.

Methods

eRNAkit Design

eRNAkit comprises three main layers: analytics, database and Rshiny application (Figure 1).
The analytics layer includes a suite of functions designed to characterise the cytoplasmic
functions of eRNAs. Key routines encompass rna2rna for identifying eRNA-mRNA
interactions from STAR aligner chimeric junctions, entropyTS for assessing tissue or cell
specificity and getREI for calculating the relative expression index. The database layer,
referred to as emi, contains four structured tables that store consensus eRNA annotations,
transcriptome-wide eRNA-mRNA interactions, subcellular localisation data of eRNA classes
and their association with ribosomes, as well as expression levels of eRNAs across major
human organs and primary cell types, including tissue and cell-type specificity metrics.
Additionally, the database integrates genome-wide enhancer-target gene information,
facilitating the differentiation between nuclear and cytoplasmic functions of eRNAs. The R
Shiny application layer provides an interactive interface for users to explore and visualise
the database, enabling dynamic querying and presentation of results.
280
Consensus human eRNA annotation
To build consensus eRNAs annotation, we collected previously published human enhancer
elements across multiple tissues and cell types from EnhancerAtlas 2.0 [21], Ensemble
regulatory build [22], eRNAbase [5], neuronal transcribed enhancers [23], and cancer associated
eRNAs [24]. We manually downloaded 2, 492, 442 coordinates in bed format and converted
those in hg19 genome build to hg38 coordinates using liftOver tool [25]. Then, overlapping
elements across the five datasets were merged to generate a consensus file using BEDTools
[26]. To ensure robust genomic coordinates, we validated all annotations against the hg38
chromosome sizes, excluding any entries with start or end positions outside of chromosome
bounds. To ensure that eRNA quantification is not confounded by normal gene expression,
we removed elements overlapping with the Ensemble gene annotation v112. We extended
the gene regions by 500 bp at both sides for this analysis to remove possible unknown
variants. Finally, we removed elements with a size < 10 bp, which are within the range of
largely unstable nuclear eRNAs.
295
eRNA subcellular localisation
Subcellular RNA-Seq data representing two RNA classes (polyA− and polyA+) across seven
cellular fractions (nucleus, cytosol, nucleolus, chromatin, nucleoplasm, membrane, insoluble
cytoplasmic fraction) were obtained from ENCODE [3]. BAM files aligned to the hg38 human
genome assembly for 13 cell lines (SK-N-SH, K562, A549, H1, HeLa-S3, MCF-7, HepG2, IMR-

90, GM12878, SK-N-DZ, SK-MEL-5, NCI-H460, and HT1080) were downloaded from the
ENCODE Data portal (Supplemental Table S2). Read counts were quantified using HTSeq
(v0.11.1)[27] on the consensus eRNA GTF file. Each RNA class and subcellular fraction had
biological replicates, and read counts were summarised by the median count per million
(CPM) across replicates.

To assess relative compartment expression of eRNAs across RNA classes and cell types, we
calculated the relative expression index (REI) across predefined pairs of subcellular
compartments for each RNA class (polyA−, polyA+ or total RNA) and cell line. REI was
defined as:

$$310 \qquad REI = log_2\left(\frac{C_1}{C_2}\right) \qquad Eq.1$$

where C_1 and C_2 represent the median of a given eRNA in two compartments. A higher REI
value indicates greater expression in C_1 relative to C_2.

Compartment pairs were defined such that finer cytoplasmic subfractions (e.g., insoluble
cytoplasmic fraction, membrane) were compared to the cytosol, and nuclear subfractions
(e.g., nucleoplasm, nucleolus, chromatin) were compared to the nucleus. The complete set
of compartment comparisons were cytosol-nucleus, insoluble cytoplasmic fraction-cytosol,
membrane-cytosol, nucleolus-nucleus, nucleoplasm-nucleus, chromatin-nucleus. For each
RNA class and cell line, REIs were computed only when expression values were available for
at least one compartments being compared. To investigate eRNA processing, a second REI
was calculated between polyA− and polyA+ samples within each compartment and cell
using the same filtering strategy.

Calculation of tissue and cell specificity for eRNAs
To assess tissue and cell specificity, we downloaded and processed total RNA-Seq BAM files
(hg38) for 32 major tissues and 77 primary cell types from the ENCODE data portal.
Biological replicates were summarised by the median CPM across replicates. eRNAs that
were lowly expressed (<1 CPM) in all tissues were removed. We call an eRNA specific (tissue
or cell) when its maximum median CPM was more than double the second largest median
CPM and the difference between the logarithm of the total number of tissues/cells and the
Shannon entropy of the median CPMs is greater that 1 [28].
330
eRNA-mRNA interaction
The eRNA-mRNA interactions in eRNAkit were identified using three high-throughput
approaches, including KARR-Seq [29], RIC-Seq [30] and PARIS [31] (Table 2). These methods differ
in their underlying chemistries, interaction specificity, and compartmental bias, together
providing a more comprehensive and nuanced map of eRNA-mRNA contacts. By integrating
datasets with differing sensitivities to nuclear versus cytoplasmic interactions, eRNAkit
enables exploration of potential post-transcriptional regulatory roles for eRNAs that might
be overlooked by any single approach.

Table 2: Comparison of RNA–RNA Interaction Mapping Methods

Feature	KARR-Seq	RIC-Seq	PARIS
Chemistry	Kethoxal-modified guanine bases (selective to ssRNA)	Formaldehyde (reversible; targets protein and RNA interactions)	Psoralen (intercalates in double-stranded RNA)
Interaction	Base-paired duplexes, especially transient or regulatory pairs	Both direct and indirect (via RNA-binding proteins) interactions	Base-paired duplexes with high confidence
Specificity	Prefers G-rich and dynamic regions	Protein-mediated proximity bias	Enriches for stable RNA duplexes
Compartment Bias (Whole Cell Lysate)	**Cytoplasmic-enriched** — favours mature, exported RNAs	**Balanced** — captures both nuclear and cytoplasmic interactions	**Nuclear-biased** — stronger detection of pre-mRNA and nuclear lncRNA
Advantages	Sensitive to dynamic, potentially regulatory interactions	Captures a broad spectrum of RNA interactions	Provides structural information; precise mapping of duplexes
Limitations	May miss highly structured or non-G-rich regions	Less resolution for direct base-pairing	Bias toward stable, long helices

We manually downloaded 78 raw sequencing datasets from 9 cells lines via the NCBI GEO
database (Supplemental Table S1). To identify eRNA-mRNA interactions, reads were quality
filtered (Q < 20) and adapter trimmed using Trimmomatic (v0.39) [32]. Cleaned reads were
aligned to the human reference genome (hg38) using STAR [33] with chimeric alignment
options enabled (--runMode alignReads --chimOutType Junctions --
chimJunctionOverhangMin 8 --chimSegmentMin 15 --outFilterMultimapNmax 100 --
alignIntronMin 1 --scoreGapNoncan -4 --scoreGapATAC -4 --alignSJoverhangMin 15 --
alignSJDBoverhangMin 10 --alignSJstitchMismatchNmax 5 -1 5 5).

Chimeric junctions were then processed using a custom R-based workflow implemented in
eRNAkit (rna2rna). Briefly, chimeric alignments were parsed into a pseudo-BEDPE format by
splitting each read into two genomic intervals (n1 and n2) and extracting positions and
strand information using the CIGAR string. These were intersected with Ensemble v111 gene
annotations and the consensus eRNA annotations. Only junctions connecting an eRNA to a
known gene were retained as putative eRNA-mRNA interactions. To enhance robustness,
only eRNA-mRNA pairs detected in at least two independent samples were included in the
final interaction set. This conservative filtering strategy prioritises reproducibility across
biological contexts and offers a high-confidence, global view of eRNA interactions.

Some of the samples analysed by KARR-Seq included cellular perturbations such as
translation inhibition, RNA-binding protein knockdowns and viral infections (Supplemental
Table S1). These experimental contexts provided an opportunity to explore potential

mechanisms driving eRNA-mRNA interactions. To assess perturbation responsiveness, we focused on binary changes-specifically, the complete gain or loss of individual interactions in treated versus untreated conditions. Interactions were considered perturbation-sensitive if they were entirely absent in control samples but appeared in treated samples, or vice versa. This stringent approach highlights interactions potentially under regulatory control in response to environmental or cellular cues.

eRNA genomic targets

To identify potential genomic targets to facilitate differentiation of cytoplasm exclusive interactions, we used two strategies (1) distal targets, based on co-localisation within the same topologically associating domains (TADs), and (2) proximal targets, based on linear distance to the eRNA loci. For the distal approach, we obtained human TAD annotations across multiple cell types and tissues from the ENCODE database [34]. Using BEDTools (v2.31.1) [26] and human gene annotation (hg38, Ensemble v111), we identified genes residing in the same TAD as each eRNA locus. For the proximal approach, genes located within ±5 kb of an eRNA locus were annotated as their proximal targets. Note that both approaches mirror what is used by existing tools to call eRNA targets (Table 1).

RNA-Seq and Ribo-Seq analysis

Standard RNA-Seq and Ribo-Seq (Supplemental Table S3) analysis was performed using our existing pipeline [35]. Briefly, the raw reads were quality filtered (Q < 20) and adapter trimmed using Trimmomatic (v0.39) [32]. Cleaned reads were aligned to the human reference genome (hg38) using HISAT2 (v2.1.0) in default settings [36]. Read counts were quantified using HTSeq (v0.11.1)[27] on the consensus eRNA annotation. The expression levels were normalised by CPM.

Permutation test

To determine whether the number of differentially expressed targets associated with a given miRNA was significantly greater than expected by chance, we performed a non-parametric permutation test. The observed count was defined as the number of eRNA targets in a specific threshold (Log$_2$ fold change (Log2FC) > 1 or < -1) among the set of detected eRNA targets of miR-130b or miR-96. To generate the null distribution, we randomly sampled the same number of eRNA (x10,000) from all the detected eRNA and counted the number matching the threshold. The empirical p-value was computed as the proportion of permutations in which the number of random eRNAs in the specified threshold was greater than or equal to the observed count. This one-tailed test estimates the probability of obtaining the observed level of enrichment under the null hypothesis of random association.

Data availability

The research community can access eRNAkit freely from the GitHub repository https://github.com/AneneLab/eRNAkit.

Author contributions

Project was conceived of by C.A.A. Data curation and processing were done by N.B, R.K, M.E, and C.A.A. Software and database creation was done by C.A.A, N.B and R.K. C.A.A supervised the project and wrote the manuscript. All authors edited the final manuscript.

Disclosure and declaration

We declare no conflict of interest.

Figure Legends

Figure 1: Overview of the main components of eRNAkit.
eRNAkit consists of three layers: **Analytics**, **Database**, and **eRNAkitApp**. The **Analytics** layer provides workflows for processing eRNA-related data and supports the design of validation experiments. The **Database** contains data on eRNA expression across major tissues and primary cells, subcellular localisation, and physical eRNA–mRNA interactions. The **eRNAkitApp** offers a user-friendly interface for querying the database and includes modules for exploring eRNA function.

Figure 2: Summary statistics from the database layer.
(a) Distribution of consensus eRNAs across chromosomes.
(b) Number of detected eRNAs (CPM > 1) in both the nucleus and cytoplasm across multiple cell types. polyA+ indicates selection of polyadenylated RNAs prior to sequencing, while polyA− indicates depletion of polyadenylated RNAs.
(c) Distribution of the relative expression index (REI) between cytoplasm (C1) and nucleus (C2) across RNA classes (see Methods). Positive values indicate enrichment in the cytoplasm, and negative values indicate enrichment in the nucleus.
(d) Number of detected eRNAs (CPM > 1) in cytoplasmic sub-fractions across RNA classes in HepG2 and K562 cells.

Figure 3: eRNA–mRNA interactions reveal potential cytoplasmic functions.
(a) Distribution of eRNA–mRNA interactions across mRNA biotypes.
(b) Similarity of eRNA–mRNA interactions (gain or loss) under different cellular perturbations. SRSF1 and DDX3X represent knockdowns of RNA-binding proteins. Harringtonine and cycloheximide indicate translational inhibition via drug treatment. RSV and VSV represent infection with respiratory syncytial virus or vesicular stomatitis virus, respectively. Arsenite denotes induction of cellular stress.
(c–d) Scatter plots showing the median expression of eRNAs in control cells versus the $\log_2$ fold change in expression upon perturbation with miR-130 **(c)** or miR-96 **(d)**.

Figure S1: Main interface and usage of eRNAkitApp.
(a) Navigation bar with the TARGET module activated.
(b) View of the LOCALISATION module.
(c) View of the EXPRESSION module.
Note: Graphs shown in S1 relate to searching for "ENSG00000283871, ENSG00000199158" (MIR130 and MIR96).

Figure S2: eRNAs are widely expressed across major tissues and primary cells.
(a–b) Number of detected eRNAs (CPM > 1) across tissues **(a)** and primary cells **(b)**.

Figure S3: Searching eRNAkitApp using eRNA ID.
(a) Navigation bar with the LOCALISATION module activated.
(b) View of the TARGET module.

(c) View of the EXPRESSION module.

Note: Graphs shown in S3 relate to searching for "en135763".

References

1. Andersson, R. *et al.* An atlas of active enhancers across human cell types and tissues.

Nature **507**, 455–461 (2014).

2. Li, Q. *et al.* Enhancer RNAs: mechanisms in transcriptional regulation and functions in

diseases. *Cell Communication and Signaling* **21**, 191 (2023).

3. Djebali, S. *et al.* Landscape of transcription in human cells. *Nature* **489**, 101–108 (2012).

4. Sartorelli, V. & Lauberth, S. M. Enhancer RNAs are an important regulatory layer of the

epigenome. *Nature structural & molecular biology* **27**, 521–528 (2020).

5. Song, C. *et al.* eRNAbase: a comprehensive database for decoding the regulatory eRNAs in

human and mouse. *Nucleic Acids Research* **52**, D81–D91 (2024).

6. Zhang, Z. *et al.* HeRA: an atlas of enhancer RNAs across human tissues. *Nucleic acids*

research **49**, D932–D938 (2021).

7. Jin, W. *et al.* Animal-eRNAdb: a comprehensive animal enhancer RNA database. *Nucleic*

acids research **50**, D46–D53 (2022).

8. Zhang, Z. *et al.* Transcriptional landscape and clinical utility of enhancer RNAs for eRNA-

targeted therapy in cancer. *Nature communications* **10**, 4562 (2019).

9. Zhang, Z. *et al.* Genetic, pharmacogenomic, and immune landscapes of enhancer RNAs

across human cancers. *Cancer Research* **82**, 785–790 (2022).

10.   Tsai, P.-F. *et al.* A muscle-specific enhancer RNA mediates cohesin recruitment and

regulates transcription in trans. *Molecular cell* **71**, 129–141 (2018).

11.   Hsieh, C.-L. *et al.* Enhancer RNAs participate in androgen receptor-driven looping

that selectively enhances gene activation. *Proceedings of the National Academy of*

Sciences **111**, 7319–7324 (2014).

12.   Melo, C. A. *et al.* eRNAs are required for p53-dependent enhancer activity and gene

transcription. *Molecular cell* **49**, 524–535 (2013).

13.   Lennox, K. A. & Behlke, M. A. Cellular localization of long non-coding RNAs affects

silencing by RNAi more than by antisense oligonucleotides. *Nucleic acids research* **44**,

863–877 (2016).

14.   Singh, S., Sinha, T. & Panda, A. C. Regulation of microRNA by circular RNA. *Wiley*

Interdisciplinary Reviews: RNA **15**, e1820 (2024).

15.   Paci, P., Colombo, T. & Farina, L. Computational analysis identifies a sponge

interaction network between long non-coding RNAs and messenger RNAs in human

breast cancer. *BMC systems biology* **8**, 1–15 (2014).

16.   Wang, Y. *et al.* Circulating miR-130b mediates metabolic crosstalk between fat and

muscle in overweight/obesity. *Diabetologia* **56**, 2275–2285 (2013).

17.   Kuhn, S. *et al.* miR-96 regulates the progression of differentiation in mammalian

cochlear inner and outer hair cells. *Proceedings of the National Academy of Sciences* **108**,

2355–2360 (2011).

18.   Soledad, R. B., Charles, S. & Samarjit, D. The secret messages between mitochondria

and nucleus in muscle cell biology. *Archives of biochemistry and biophysics* **666**, 52–62

(2019).

19.   Vendramin, R., Marine, J. & Leucci, E. Non-coding RNA s: the dark side of nuclear–

mitochondrial communication. *The EMBO journal* **36**, 1123–1133 (2017).

20.   Bonn, S. *et al.* Tissue-specific analysis of chromatin state identifies temporal

signatures of enhancer activity during embryonic development. *Nature genetics* **44**, 148–

156 (2012).

21.   Gao, T. & Qian, J. EnhancerAtlas 2.0: an updated resource with enhancer annotation

in 586 tissue/cell types across nine species. *Nucleic acids research* **48**, D58–D64 (2020).

22.   Zerbino, D. R., Wilder, S. P., Johnson, N., Juettemann, T. & Flicek, P. R. The ensembl

regulatory build. *Genome biology* **16**, 1–8 (2015).

23.   Dong, P. *et al.* Population-level variation in enhancer expression identifies disease

mechanisms in the human brain. *Nature genetics* **54**, 1493–1503 (2022).

24.   Chen, H. & Liang, H. A high-resolution map of human enhancer RNA loci

characterizes super-enhancer activities in cancer. *Cancer cell* **38**, 701–715 (2020).

25.   Hinrichs, A. S. *et al.* The UCSC genome browser database: update 2006. *Nucleic acids*

research **34**, D590–D598 (2006).

26.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**, 841–842 (2010).

27.   Anders, S., Pyl, P. T. & Huber, W. HTSeq—a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

28.   Li, S. *et al.* exoRBase: a database of circRNA, lncRNA and mRNA in human blood

exosomes. *Nucleic acids research* **46**, D106–D112 (2017).

29.   Wu, T. *et al.* KARR-seq reveals cellular higher-order RNA structures and RNA–RNA

interactions. *Nature Biotechnology* **42**, 1909–1920 (2024).

30.   Cai, Z. *et al.* RIC-seq for global in situ profiling of RNA–RNA spatial interactions.

Nature **582**, 432–437 (2020).

31.   Lu, Z. *et al.* RNA duplex map in living cells reveals higher-order transcriptome

structure. *Cell* **165**, 1267–1279 (2016).

32.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

33.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21

(2013).

34.   Hammal, F., De Langen, P., Bergon, A., Lopez, F. & Ballester, B. ReMap 2022: a

database of Human, Mouse, Drosophila and Arabidopsis regulatory regions from an

integrative analysis of DNA-binding sequencing experiments. *Nucleic acids research* **50**,

D316–D325 (2022).

35.   Cogan, J. A., Benova, N., Kuklinkova, R., Boyne, J. R. & Anene, C. A. Meta-analysis of

RNA interaction profiles of RNA-binding protein using the RBPInper tool. *Bioinformatics*

Advances **4**, vbae127 (2024).

36.   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory

requirements. *Nature methods* **12**, 357 (2015).

540
541
542

Figure 1

Figure 2

Figure 3

a

b

c

Figure S1

a

b

Figure S2

Figure S3