VGLL1 contributes to both the transcriptome and epigenome of the developing trophoblast compartment

Ruben I. Calderon[1,2,3], Nirvay Sah[1,3], Molly Huang[4], Ryan H. Kittle[1,7], Walee B. Shaik[1,3], Jennifer N. Chousal[1], Sampada Kallol[1,3], Tony Bui[1], Robert Morey[1,3], Alejandra Mitre[1], Norah M.E. Fogarty[5,8], Claudia Gerri[5,9,10], Claire Zheng[1,2], Peter DeHoff [3,4], Pratik Home[6,11], Kathy K. Niakan[5,12,13,14], Heidi Cook-Andersen[2,3,4], Kathleen M. Fisch[4], Soumen Paul[6], Francesca Soncin[1,3,*]

Affiliations:

1. Department of Pathology, School of Medicine, University of California San Diego, La Jolla, CA 92093, USA

2. School of Biological Sciences, University of California San Diego, La Jolla, CA 92093, USA

3. Sanford Consortium for Regenerative Medicine, La Jolla, CA, 92093, USA

4. Department of Obstetrics, Gynecology, and Reproductive Sciences, School of Medicine, University of California San Diego, La Jolla, CA, 92093, USA.

5. Human Embryo and Stem Cell Laboratory, The Francis Crick Institute, London, NW1 1AT, UK.

6. Department of Pathology & Laboratory Medicine and Institute for Reproductive and Developmental Sciences, University of Kansa Medical Center, Kansas City, KS 66160.

Current addresses:

7. Department of Genetics and Development, Columbia University Irving Medical Center, New York, NY 10032, USA.

8. Centre for Gene Therapy and Regenerative Medicine, King's College London, London, SE1 9RT, UK.

9. Max Planck Institute of Molecular Cell Biology and Genetics, Pfotenhauerstrasse 108, 01307, Dresden, Germany.

10. Center for Systems Biology Dresden, Pfotenhauerstrasse 108, 01307, Dresden, Germany.

11. BioIVT, 1101 W Cambridge Circle Dr. Kansas City, KS 66103.

12. The Loke Centre for Trophoblast Research, Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge CB2 3EG, UK.

13. Wellcome Trust – Medical Research Council Stem Cell Institute, University of Cambridge, Jeffrey Cheah Biomedical Centre, Puddicombe Way, Cambridge CB2 0AW, UK.

14. Epigenetics Programme, Babraham Institute, Cambridge CB22 3AT, UK.

15. Lead contributor   * Corresponding author: fsoncin@health.ucsd.edu

Abstract

The trophectoderm (TE), the first lineage specified during mammalian development, initiates implantation and gives rise to placental trophoblasts. While animal models have elucidated key conserved signaling pathways involved in early TE specification, including BMP, WNT, and HIPPO, species-specific differences during early development emphasize the need for human-specific models. We previously identified VGLL1, a coactivator of TEAD transcription factors, as a human-specific placental marker. In this study, we employed a pluripotent stem cell (PSC)-based model of TE induction by BMP4 to investigate chromatin remodeling and transcriptional dynamics during TE specification. BMP4-induced chromatin accessibility changes promoted a trophoblast gene expression program, while mesoderm lineage markers were only transiently expressed upon canonical WNT activation. We found that VGLL1 was expressed downstream of key TE transcription factors (GATA2/3, TFAP2A/C) but was essential for full trophoblast differentiation by up-regulating EGFR and reinforcing GATA3 expression through positive feedback. Notably, VGLL1 enhanced canonical WNT signaling via direct regulation of WNT receptors and effectors. We also identified KDM6B, a histone demethylase that removes H3K27me3 repressive marks, as a direct VGLL1 target. KDM6B facilitated activation of bivalent promoters associated with TE markers, linking epigenetic regulation to lineage specification. Our findings establish a mechanistic framework positioning VGLL1 as a central regulator that integrates HIPPO, BMP, and WNT signaling pathways to drive human TE and trophoblast development.

Statement of Significance

This research explores human-specific mechanisms for the induction of the trophectoderm (TE) during early development, which is crucial for placental function and pregnancy establishment. Using a pluripotent stem cell system, we showed that BMP4 treatment induced TE specification by a combination of TE-specific gene expression activation and chromatin changes. We showed that the transcription co-factor VGLL1 played a critical role in directing cells toward a TE fate via three mechanisms: (1) regulating the expression of TE-specific genes; (2) enhancing activation of canonical WNT pathway; and (3) activating the expression of KDM6B, which acts on the chromatin to promote TE-specific gene accessibility.

Introduction

The trophectoderm (TE) is the first lineage specified during mammalian development and is important for embryo cavitation, implantation, and pregnancy initiation (1). After implantation, the TE gives rise to the trophoblast compartment of the placenta, contributing to the formation of the maternal/fetal interface (2). Early pregnancy losses and several placental-associated disorders are linked to abnormalities in the establishment and differentiation of the TE compartment during early development (2). However, ethical and technical considerations have limited our understanding of the mechanisms of TE specification and trophoblast development in humans. Studies in animal models have determined that tight spatio-temporal regulation of highly conserved signaling pathways, including BMP, WNT, and HIPPO, are required to establish the TE in the pre-implantation embryos and for early placental development (3–8). However, increasing evidence points to species-specific differences in the composition of regulatory networks that contribute to the development of the extra-embryonic ectoderm compartment (3, 9–11). Therefore, species-specific models are better suited to investigate TE specification and early placental development (3, 9, 12).

The HIPPO signaling pathway is a conserved cascade that initiates the specification of the TE lineage following activation of GATA3 and CDX2 expression by the TEAD4/YAP1 complex (10, 13, 14). In early mouse embryos, differential HIPPO pathway activation between the inner and outer cells leads to the expression of CDX2, which initiates a TE expression program, including induction of GATA3 and ELF5 (13, 15, 16). EOMES is expressed down-stream of CDX2 and, while not required for TE specification, it is fundamental for the maintenance of the TE transcriptional program and maintenance of mouse trophoblast stem cells (TSCs) (17, 18). In humans, four transcription factors, GATA2, GATA3, TFAP2A, and TFAP2C, form a transcriptional network required for TE initiation (TEtra factors) downstream of the HIPPO signaling pathway (19). This network drives the establishment of an extra-embryonic ectoderm signature, which includes mature markers such as VGLL1, TP63, and EGFR, but not EOMES, which is dispensable for human TE and trophoblast development (9, 20). In particular, we have previously identified VGLL1 as a human-specific marker

expressed in the trophoblast progenitor compartments of the placenta (9). VGLL1 is a
conserved binding partner of the TEAD family of transcription factors and competes with
YAP1 to form a transcriptional complex with TEAD4, therefore potentially modulating HIPPO
signaling in human placenta (21).

Canonical WNT signaling is a well-characterized developmental pathway mediated
by nuclear localization of β-catenin, where it binds to TCF/LEF complexes and drives
expression of WNT-specific target genes (22). This pathway is required for maintenance and
expansion of various human epithelial stem cells, including intestinal and skin stem cells
(23, 24). In human pre-implantation embryos, the TE compartment is enriched in WNT
components, including ligands and receptors (25–29). Moreover, compared to mouse, both
human and non-human primate embryos show higher canonical WNT-related gene
expression in the TE (30). WNT genes are also highly expressed in the trophoblast
compartment of the placenta. *In vitro*, both human TSCs and trophoblast organoids require
activation of canonical WNT signaling for proliferation (31–33) while its transient inhibition is
required to initiate differentiation of extravillous trophoblasts (EVT) (32, 34).

Bone Morphogenetic Protein (BMP) signaling has well described roles in gastrulation,
including specification of both the embryonic mesoderm lineage and primordial germ cells
(5). However, recent evidence demonstrated BMP activation in the TE of human, but not
murine, blastocysts (5, 12, 35), though autocrine BMP signaling has been reported in both
mouse and human TSCs (36). *In vitro*, exogenous BMP4 has been shown to drive
differentiation of human pluripotent stem cells (PSCs) into different lineages, namely
embryonic mesoderm, primordial germ cells, or extra-embryonic trophoblast lineage
progenitors, based on the cellular context and the activation of other signaling pathways (5).
In particular, embryonic mesoderm differentiation requires BMP-induced WNT activation,
while inhibition of β-catenin-dependent WNT signaling downstream of BMP effectively
blocks mesoderm differentiation, instead promoting trophoblast lineage (37). However, the
mechanisms underlying WNT activation by BMP4 and the switch from mesoderm to
trophoblast lineage in PSC-derived TE-like cells have not been elucidated yet.

Regulatory networks drive establishment and maintenance of cellular identity by acting on a lineage-specific epigenetic landscape downstream of signaling pathways (38). DNA methylation and histone marks surrounding lineage-specific genes determine chromatin accessibility for specific transcription factor (TF) binding (38). Recent studies in pre-implantation embryos have described epigenome changes across early development from zygote to blastocyst, including segregation of TE and ICM (39–41). However, there is little mechanistic understanding on how epigenetic marks are laid out in a lineage-specific manner during the specification of the TE and in trophoblast development. Moreover, it is largely unknown how multiple signaling pathways like HIPPO, WNT, and BMP converge towards a lineage-specific outcome and drive trophoblast development.

PSCs represent a unique resource to study species-specific genetic and epigenetic mechanisms of TE and trophoblast specification as they are tractable models for genetic and chemical manipulation. Multiple protocols have been published to study TE and trophoblast specification from human PSCs (42–46). We have established a 2-step protocol based on BMP4 treatment of PSCs to first differentiate them into TE-like cells, followed by either further differentiation into mature trophoblasts or derivation of *bona fide* hTSCs (47, 48). Based on literature, we further optimized the first step by adding a WNT inhibitor, IWP2, to block concurrent activation of a mesoderm program and obtain a more homogeneous population of TE-like cells (37, 45). We previously reported that VGLL1 is one of the most highly upregulated genes in the BMP4-directed differentiation of human embryonic stem cells (hESCs) into TE-like cells (9). Here, we use this model to first investigate the BMP4-dependent chromatin modifications that occur during the transition from PSC to TE-like cells in relation to gene expression changes and WNT activation status. Then, we present our findings on how VGLL1 directly and indirectly drives a trophoblast-specific gene expression and epigenomic program in this PSC-derived trophoblast model.

Results

BMP4 mediates chromatin accessibility to stabilize a trophoblast-specific expression program

We have previously optimized a protocol for differentiation of PSCs into TE-like cells
by treatment with 10ng/mL BMP4 and the WNT inhibitor IWP2 over 4 days, which
successfully blocks mesoderm differentiation and promotes a uniform EGFR+ population
(45). To start dissecting the transcriptional changes towards TE-like cells during this
differentiation protocol, we first performed bulk RNA-seq analysis of H9 hESCs at day 0, day
2, day 4 following treatment with BMP4 or BMP4/IWP2 (**Suppl. Fig. S1A**) and identified
differentially expressed genes for each timepoint and treatment compared to d0 (**Suppl.
Table S1**). Genes downregulated in both treatments included pluripotency and embryonic
lineage markers, i.e. *NANOG, POU5F1,* and *SOX2* (**Suppl. Table S1**). Upregulated genes
common across timepoints and treatments included the TEtra factors *GATA2, GATA3,*
TFAP2A, and *TFAP2C*. By day 4, common upregulated genes included *ENPEP*, and
trophoblast markers *VGLL1* and *TP63 (***Suppl. Fig. S1B and Suppl. Table S1**). Genes
uniquely upregulated by BMP4 treatment on day 2 were associated with various signaling
pathways, including WNT signaling (**Fig. 1A, Suppl. Table S1**). As expected, secondary to
WNT signaling activation in hESCs, we detected upregulation of mesoderm markers *EOMES,*
TBXT, and *TBX6* at day 2 of BMP4 treatment (**Fig. 1B**). However, by day 4, the expression of
these markers decreased to levels comparable to the treatment with the addition of IWP2
(**Fig. 1B**). Accordingly, the WNT signaling pathway was no longer one of the top significantly
enriched pathways at day 4 of BMP4 alone treatment (**Suppl. Fig. S1C**). To confirm the
activation timeframe of the WNT pathway during BMP4 treatment, we used a TCF/LEF-eGFP
reporter hESC line (HUES9) where detection of GFP fluorescence is dependent on canonical
WNT activation (49). We detected WNT signaling activation starting at day 2, a peak at day 3,
and decreased activity at day 4 of BMP4 treatment (**Suppl. Fig. S1D**). The WNT inhibitor IWP2
completely blocked eGFP expression, while another inhibitor, IWR1, showed only partial
suppression of WNT activity at the dose used (**Suppl. Fig. S1D**). Accordingly, we detected a
dose-dependent inhibition of both WNT targets, *LEF1* and *WNT5B,* as well as mesoderm
markers *TBXT* and *EOMES* between the two inhibitors (**Suppl. Fig. S1E**). Overall, these data
suggest that the WNT pathway is transiently activated during treatment of hESCs with low

dose BMP4, but that both WNT activity and mesoderm gene expression decrease over time

as the TE identity is established in this model.

To investigate chromatin accessibility changes associated with the differential

establishment of the mesoderm and TE lineage during BMP4 treatment of PSCs, we

performed Assay for Transposase-Accessible Chromatin (ATAC) followed by sequencing in

hESCs treated with BMP4 or BMP4/IWP2 at days 0, 2, and 4. We found that peaks resided at

promoter (31%), intergenic (24%), and intronic (23%) regions (**Suppl. Fig. S1F**). Following

differential peak analysis, both at day 2 and day 4 of treatment, we found open chromatin

regions specific to BMP4 or BMP4/IWP2, as well as regions common to both treatment

protocols (**Fig. 1C, Suppl. Table S2**). We performed Hypergeometric Optimization of Motif

EnRichment (HOMER) analysis on common and day/treatment-specific regions to identify

TF binding motifs. Commons regions at both days 2 and 4 contained binding motifs for well-

known TE and trophoblast TFs, including TEAD and AP1-complex members (i.e. JUN and

FOS) (**Suppl. Fig S1G**). Binding motifs of both TEtra factors GATA3 and TFAP2C appeared in

both common and treatment-specific regions. At day 2 of BMP4-only treatment, we found

open chromatin at LEF-1 and TCF3, TFs downstream of the WNT pathway. We then looked at

the chromatin regions around the transcription start sites of well-known trophoblast and

mesoderm genes (**Fig.1D and Suppl. Fig. S1H**). The open chromatin region across

treatments/days showed a sharp peak and remained consistent around trophoblast markers

(**Fig. 1D**), including GATA3 and EGFR (**Suppl. Fig. S1H**). Conversely, open chromatin around

mesoderm genes showed a broader distribution that decreased between day 2 and day 4 in

response to both treatments (**Fig. 1D**), including at EOMES and TBXT (**Suppl. Fig. S1H**) These

chromatin accessibility data are largely consistent with the gene expression patterns

observed above.

Finally, we looked at the differential expression of known epigenetic modifiers that

could account for these changes in chromatin accessibility. We found multiple lineage-

specific chromatin modifiers differentially expressed during TE-specification. These

included down-regulation of *de novo* DNA methyltransferases *DNMT3A* at day 2, a switch

from *TET1* to *TET2* (50) expression, and up-regulation of histone demethylases like *KDM6B*,

206 *KDM4B,* and *KDM1B* (**Fig. 1E**). These data suggest that low levels of BMP4 preferentially

207 enable a trophoblast transcriptional signature via a combination of lineage-specific gene

208 expression activation and finely tuned regulation of the epigenetic landscape.

VGLL1 regulates a TE-specific gene expression program

210  In the context of BMP4 treatment of human PSCs, the TEtra factors GATA2, GATA3,

211 TFAP2A, and TFAP2C have been identified as initiators of the TE/trophoblast transcriptional

212 program, upstream of more mature trophoblast markers like TP63, EGFR, ELF5, and

213 VGLL1(9, 19, 20). To further tease apart the timeframe of activation of these TE/trophoblast

214 genes in this BMP4-based model, we investigated the expression of TEtra factors, and

215 mature trophoblast markers by qPCR at short intervals within the first 72 hours of

216 BMP4/IWP2 treatment. As previously reported, TEtra factors *GATA3* and *TFAP2C* were the

217 first to be activated, with upregulation as early as 6h after the start of the treatment and

218 continuous increase over time (**Fig. 2A**). *VGLL1* expression started to rise between 12 hours

219 and 24 hours of BMP4/IWP2 treatment (**Fig. 2A**) while other mature TE/trophoblast markers

220 were observed at 48 hours (*TP63*) and 72 hours (*ELF5* and *EGFR*), respectively (**Fig. 2A**). By

221 immunofluorescence, VGLL1 protein was detected in the nucleus after 24 hours of

222 BMP4/IWP2 treatment, where it co-localized with TEAD4 and YAP1 throughout TE induction

223 (**Suppl. Fig. S2A**). While previous literature considered VGLL1 a mature trophoblast marker

224 (like TP63), these data suggest that VGLL1 might play an earlier role downstream of TEtra

225 factors but before other trophoblast markers.

226  To understand the role of VGLL1 in this context, we established hESC lines that

227 constitutively express VGLL1-specific shRNA for targeted gene knock-down (or shScramble

228 as control, hereafter referred to as Scramble). Compared to Scramble, treatment of shVGLL1

229 hESCs with BMP4/IWP2 led to blunted up-regulation of late trophoblast markers, *TP63*,

230 *ASCL2*, and *GCM1* (**Fig. 2B**). Interestingly, TEtra factor *GATA3* and the early TE marker *CDX2*

231 also showed lower up-regulation. In addition, surface levels of EGFR levels were also lower

232 in the BMP4/IWP2-treated shVGLL1-hESCs, compared to similarly-treated Scramble-hESCs

233 (**Suppl. Fig. S2B**), suggesting an incomplete TE differentiation. Therefore, to further explore

the effect of VGLL1 knock-down in TE induction, we performed bulk RNA-seq analysis on two

independent shVGLL1 clones and one Scramble control at days 0 and 4 of BMP4/IWP2

treatment (**Suppl. Fig. S2C**). Overall, shVGLL1 clones showed a marked delay in

differentiation compared to Scramble (**Suppl. Fig. S2C**). We found 7,284 genes that were

differentially dysregulated in BMP4/IWP2-treated shVGLL1 hESCs compared to Scramble

(adj. p value<0.05), organized in 4 distinct cluster based on pattern of expression (Suppl.

Table S3). Specifically, 3,056 genes showed blunted down-regulation (cluster #2) compared

to Scramble and were associated with transcriptional regulation of pluripotent stem cells

and embryo development (**Suppl. Fig. S2D**), including pluripotency markers *POU5F1*, *SOX2*,

and *SALL2*. The 4,192 up-regulated genes that showed misregulation in shVGLL1 hESCs

compared to Scramble (**Fig. 2C**) organized in 3 clusters of expression. Cluster #3 contained

genes that failed to be up-regulated, including TE/trophoblast markers *GATA2*, *NR2F2* (51),

and *CCR7* (52), while cluster #4 included genes with blunted expression compared to

Scramble, including TEtra factors *GATA3* and *TFAP2C*, TE markers *ENPEP* (53), and

trophoblast markers *EGFR*, *ARID3A*, and *ARID5B* (54). Gene ontology analysis confirmed

that genes that failed to be fully up-regulated were associated with response to hormones

and cell migration and were highly enriched for placental genes (**Suppl. Fig. S2E**).

Interestingly, the remaining cluster #1 contained genes that were more highly up-regulated

in shVGLL1 cells compared to Scramble and included TFAP2A, a potential compensatory

response to lack of VGLL1 and lower expression of other TEtra factors. These data suggest

VGLL1 acts through a positive feedback mechanism downstream of the TEtra factors to

complete a TE/trophoblast-specific program.

We then investigated if VGLL1 was required for TSC derivation. Recently, we

demonstrated that hESC-derived TE-like cells (hESCs following 4 days of BMP4/IWP2

treatment) can adapt to hTSC culture conditions, resulting in cells that show characteristics

of *bona fide* primary hTSCs (48). hTSC culture medium includes a canonical WNT agonist

and EGF signaling activation combined with TGFβ and ROCK signaling inhibition. Therefore,

we attempted to derive hTSC-like cells from both Scramble and two shVGLL1 hESC clones.

Similar to our initial report with wild type PSCs, both Scramble and shVGLL1 hESCs rapidly

lost both epithelial morphology and cell surface expression of EGFR during the early stages of adaptation (**Suppl. Fig. S3A-B**). However, while Scramble cells acquired a hTSC-like colony morphology and regained expression of surface EGFR by passage 6, shVGLL1 clones maintained a single cell morphology and failed to regain EGFR (**Fig. 2D-E**). Moreover, shVGLL1 clones expressed lower levels of TEtra factors *GATA2*, *GATA3*, and *TFAP2C*, compared to Scramble hTSCs (**Fig. 2F, Suppl. Fig. S3D**). Interestingly, while the TE marker *ENPEP* also decreased in shVGLL1 cells, *CDX2* increased compared to Scramble (**Suppl. Fig. S3C-D**). These data suggest that VGLL1 knock-down PSCs failed to form TSCs and that VGLL1 is required for the maintenance of a trophoblast signature, in particular related to the expression of some TEtra factors and EGFR.

We further evaluated VGLL1 activity, uncoupled from BMP4 treatment, to investigate the function of VGLL1 in promoting the TE-specific transcriptional program. We leveraged a system composed of a doxycycline (DOX)-inducible dCas9-VPR and VGLL1-specific guide RNAs for CRISPR-mediated gene activation in undifferentiated hESCs (iVGLL1a) (**Fig. 3A**). In the absence of exogenous BMP4, ectopic expression of VGLL1 in hESCs initiated changes in cell morphology with cells taking on a cobblestone shape, typical of TE/trophoblast cells (**Fig. 3B**), and increased surface expression of EGFR (**Fig. 3C**). Moreover, we observed up-regulation of the TEtra factor *GATA3* and trophoblast marker *EGFR* accompanied by down-regulation of pluripotency marker *NANOG* by qPCR (**Fig. 3D**). We then performed bulk RNA-seq analysis on three independent iVGLL1a clones treated with DOX (or DMSO as carrier control) (**Suppl. Fig. S4A**). We found 6,530 differentially expressed genes (DEGs) in DOX-treated cells compared to DMSO (adj. p-value 0.05), of which 3,053 were up-regulated and 3,477 down-regulated (**Fig. 3E, Suppl. Table S4**). Significantly down-regulated genes were enriched in pluripotency genes, including *MYC*, *NANOG*, *POU5F1*, and *SOX2* (**Fig. 3E and Suppl. Fig. S4B**). Up-regulated DEGs included TE/trophoblast markers *ARID5B*, *EGFR*, and *GATA3* (**Fig. 3E**). Moreover, gene ontology analysis of pathways and biological processes revealed the activation of signaling pathways important for TE and trophoblast specification and function, including TGFβ, MAPK, and PI3K-Akt pathways (**Suppl. Fig. S4C**). Interestingly, iVGLL1a cells could not be adapted to TSC culture conditions. These data point to a specific

292 role of VGLL1 in controlling a TE/trophoblast transcriptional program but suggest that

293 ectopic expression of VGLL1 alone is insufficient for TSC establishment from hPSCs.

VGLL1 regulates canonical WNT pathway effectors

Gene ontology analysis of up-regulated genes in response to DOX treatment of

iVGLL1a hESCs suggested activation of WNT signaling pathway (**Suppl. Fig. S4C**). Indeed,

VGLL1 expression induced up-regulation of WNT pathway components and mediators,

including *EPHA2, FZD2, LEF1, LRP4, WLS, WNT4/5A/5B/6/10A/11*, as well as WNT signaling

targets like *FST* (**Fig. 4A and Suppl. Fig S4D**). We also confirmed β-catenin dependent WNT

activation using a reporter expressing mCherry under constitutive promoter and GFP under

TCF/LEF promoter (55). We observed GFP signal in a subset of mCherry expressing cells

following DOX treatment compared to DMSO control (**Fig. 4B**). Interestingly, we observed up-

regulation of mesoderm markers on DOX-treated iVGLL1a hESCs by qPCR, including EOMES

and TBXT (**Fig. 4C**). During BMP4-induced hESC differentiation, co-treatment with WNT

inhibitor IWP2 effectively suppressed mesoderm marker expression. Therefore, we tested

the dependence of mesoderm marker expression down-stream of VGLL1 activation in

iVGLL1a hESCs on activation of the WNT signaling pathway. Addition of IWP2 during

doxycycline induction of VGLL1 effectively reduced the expression of WNT genes *FST, LEF1*,

and *LGR5* (**Suppl. Fig. S4D**). Moreover, this resulted in reduced expression of mesoderm

markers *EOMES* and *TBXT* (**Fig. 4C**) and improved expression levels of *GATA3* and *EGFR* by

qPCR and flow cytometry (**Fig. 4D-E**). We compared gene expression profiles of DOX and

DOX/IWP2 treated cells (**Fig. 4F, Suppl. Table S4**) and found that DOX/IWP2 treated cells

showed significant loss of mesoderm-associated genes, including *EOMES* and *TBXT* (**Suppl.**

Table S4). These data support a role for VGLL1 in modulating β-catenin-dependent WNT

signaling, an important pathway in trophoblast development and placenta function.

VGLL1 directly regulates trophoblast-associated genes at placenta-specific distal regulatory regions

We compared the differentially expressed genes in both the constitutive knock-down
and gene activation systems to further narrow down VGLL1 contributions to the TE
transcriptional program (**Fig. 5A**). We found 528 genes that were concordantly regulated by
VGLL1 expression, specifically down-regulated in VGLL1sh hESCs (relative to Scramble)
after TE induction, and up-regulated in the iVGLL1a hESCs treated with DOX/IWP2 (**Suppl.**
Table S4). These genes were involved in cancer pathways, focal adhesion, TGFβ and HIPPO
signaling pathways (**Suppl. Fig. S4E**) and were highly associated with placental gene
expression (**Suppl. Fig. S5A**). We then set out to investigate whether VGLL1 directly regulates
the transcription of a subset of these genes. VGLL1 lacks a DNA binding domain but is a
transcriptional co-factor for TEAD family members. TEAD4 is expressed in both hESCs and
hESC-derived TE-like cells and plays a role in trophoblast differentiation of hESCs as well as
in maintenance of stemness in primary hTSCs (56, 57). Therefore, we predicted that VGLL1
binding to regulatory regions might be mediated through its association with TEAD4. To test
this, we performed immunoprecipitation with either an anti-TEAD4 or anti-VGLL1 antibody
on the choriocarcinoma cell line JEG3 and confirmed co-precipitation of VGLL1 and TEAD4,
respectively (**Suppl. Fig. S5B**). We also confirmed binding of VGLL1 and TEAD4 to a validated
5'UTR region of *GATA3* by ChIP-qPCR (**Suppl. Fig. S5C**) (58).

Having validated the interaction between VGLL1 and TEAD4 in a trophoblast cell line,
we next set out to identify binding regions of VGLL1 in complex with TEAD4 in hESC-derived
TE-like cells. Therefore, we performed ChIP-seq using both anti-VGLL1 and anti-TEAD4
antibodies on BMP4/IWP2-treated hESCs at day 4. VGLL1 ChIP peaks were distributed
mainly between intronic (44.4%), intergenic (30.6%), and promoter (20%) regions (**Suppl. Fig**
S5D), suggesting that VGLL1 might regulate target genes through transcription activation at
promoters as well as through interactions at distal enhancer regions. In line with this, we
found evidence of VGLL1 occupancy at *EGFR* and *ENPEP* promoter regions (**Fig. 5B)**, as well
as in distal regulatory regions of *GATA3* and *EGFR* identified as putative active enhancers in
the human placenta (59) (**Suppl. Fig. 5E, Suppl. Table S5**). VGLL1 peaks were enriched in
TEAD family member motifs, including TEAD4 and TEAD1 (**Fig. 5C**). In our ChIP-seq data,
VGLL1 and TEAD4 co-occupied regions at trophoblast-associated genes *EGFR*, *ARID3A*,

CDH1, *ENPEP*, *KRT7*, *KRT8*, *PTPN14*, and *TEAD1* (**Fig. 5D**, **Suppl. Fig. S5F, Suppl. Table S5**). In addition, we identified VGLL1-specific binding at TE/trophoblast regulators *ASCL2*, *ARID5B*, *CCR7*, CITED2, *NODAL*, *TP63,* and WNT signaling effectors, such as the long non-coding *LEF-AS1*, and components *EPHA2*, *FZD2*, *ITGB1, LGR4*, *LRP5/6*, *WLS*, *WNT5A*, and *WNT5B* (**Fig. 5D and Suppl. Table S5**). The genes bound by VGLL1 were highly associated with the human trophectoderm transcriptional signature and not with mesoderm-derived cell types (**Suppl. Fig. S5G**). Together, these data demonstrate that VGLL1 directly regulates a TE/trophoblast-specific program at promoter regions in complex with TEAD4 and at distal regulatory regions potentially with a novel partner.

VGLL1 contributes to BMP4-dependent epigenetic changes by regulating the histone demethylation enzyme KDM6B

To investigate if VGLL1 might contribute to the epigenetic changes for TE induction of hESC downstream of BMP4-treatment, we first looked at chromatin modifiers whose expression patterns followed VGLL1 manipulation (**Fig. 5A, Suppl. Table S4**). Among the 528 genes concordantly regulated, we identified *TET2*, involved in DNA demethylation, and *KDM6B*, an enzyme responsible for demethylation of histone 3 lysine residue 27 (H3K27me3/2/1). Furthermore, we detected VGLL1 occupancy at *KDM4B* and *KDM6B* loci in BMP4/IWP2-treated hESCs (**Fig. 5D**) as well as in hTSCs (data not shown), indicating these lysine demethylases are direct targets of VGLL1. KDM6B has been reported to regulate the demethylation of H3K27me3 at promoter regions of *Gata3* and *Cdx2*, thus promoting their expression in the TE of pre-implantation murine embryos (60). Therefore, we treated hESCs with a KDM6B demethylase activity inhibitor, GSKJ4, during BMP4/IWP2-mediated differentiation, and, similar to VGLL1 knock-down, we found decreased levels of TE markers *GATA3* and *CDX2*, compared to vehicle control (**Fig. 5F**). Collectively, these data suggest that VGLL1 further contributes to the establishment of a TE/trophoblast transcriptional signature by regulating the expression of chromatin remodeler enzymes.

VGLL1 and KDM6B are co-expressed in the TE compartment of pre-implantation embryos

To extend our findings beyond PSC-based models, we examined VGLL1 expression in
human embryos. Using published datasets, we investigated its temporal and spatial
expression during early embryonic development. Yan *et al.* previously performed RNA
sequencing (RNA-seq) of individual early human embryos from the oocyte to pre-
implantation blastocyst (27). Analysis of this data revealed that *VGLL1* was expressed at very
low levels until the blastocyst stage, at which point its expression began to increase (**Suppl.**
Fig. S6A). Blakeley *et al.* investigated the transcriptional signature of the three main cell
types found in the pre-implantation blastocyst: TE, epiblast (EPI), and primitive endoderm
(PE) with single-cell RNA-seq (3). Examination of datasets from these cell types revealed that
VGLL1 was highly expressed in TE, with some expression in the PE but not in EPI (**Suppl. Fig.**
S6B). To further distinguish between polar TE (adjacent to the inner cell mass) and mural TE
(adjacent to the blastocoele) clusters, we re-analyzed single-cell RNA-seq data of
dissociated human embryo cells from another study that performed extended embryo
culture (61). We found that *VGLL1* expression was increased in the TE relative to the EPI and
PE, with slightly higher expression in the polar TE compared to the mural TE. By contrast,
CDX2, another TE marker central to the TE program in humans and mice, was similarly
increased in TE compared to EPI and PE cells but was higher in the mural compared to the
polar compartment (**Fig 6A**). To assess if protein expression correlated with our
transcriptomic results, we co-stained pre-implantation embryos with VGLL1 and both extra-
embryonic (GATA3, TP63) and embryonic (NANOG) markers. We observed VGLL1 positive
cells at the morula stage in both inside and outside cells and co-staining with GATA3 in a
subset of outside cells (**Fig 6B**). At the blastocyst stage, VGLL1 expression was localized in
the nuclei of TE cells (**Fig 6B, Suppl. Fig. S6C**). These data confirm VGLL1 activity in the TE
compartment of human embryos, further suggesting a role for this transcriptional co-factor
in establishment/maintenance of this lineage.

Finally, we examined KDM6B expression in both human embryos and first trimester
placentas and found that its expression correlated with VGLL1 in the TE/trophoblast
compartment. In human pre-implantation embryos, *KDM6B* was highly expressed in the TE,
with increased expression in the polar compared to mural TE (**Fig. 6C**). In the first trimester

placenta, KDM6B was expressed in the cytotrophoblast (CTB) compartment and proximal
cell column as well as in the syncytiotrophoblast compartment (STB) (**Fig. 6D**). Taken
together with our ChIPseq, these data point to a direct role for VGLL1 in regulating KDM6B
expression in the progenitor compartment of first trimester placenta. Overall, this study
demonstrates that VGLL1 regulates both genetic and epigenetic features to support a
trophoblast lineage-specific transcriptional program.

Discussion

Correct establishment of the TE lineage is fundamental for placental development
and a successful pregnancy outcome. Because of the ethical and technical limitations of
working with human embryos, various protocols have been reported to differentiate human
pluripotent stem cells into trophoblast in order to study species-specific mechanisms of
development. In this work, we used a BMP4-model for TE induction from primed PSCs
previously established in our lab, which includes addition of the WNT inhibitor IWP2 to block
mesoderm differentiation, followed by TSC culture adaptation (48). First, we further explore
the transcriptional programs and lineages elicited by BMP4 in the presence and absence of
IWP2. We demonstrated that activation of the canonical WNT pathway and consequent up-
regulation of mesoderm markers was limited to the first 2 days of treatment, with
transcription directed towards a TE-like program by day 4. Universal TE markers including
GATA3, ENPEP, and *TACSTD2* were all induced by BMP4 treatment. Our ATAC-seq data
comparing BMP4-driven differentiation with and without WNT signaling inhibition indicates
that, in primed hPSCs, the chromatin surrounding trophoblast lineage genes was accessible
for transcription throughout differentiation. The changes in chromatin reported here
accompany the temporal increase of TE/trophoblast markers that we observed in our RNA-
seq data. On the other hand, we showed that, at these concentrations of BMP4, chromatin
near mesoderm lineage genes closed over time even in the absence of WNT inhibition.

Temporal activation of TE/trophoblast lineage in our model was similar to previously
published studies of BMP4-dependent induction of the trophoblast lineage, with TEtra
factors GATA2/3 and TFAP2A/C expressed early during differentiation followed by more

mature markers (19). However, in their study Krendl and colleagues defined VGLL1 as a late

marker. In this study, we determined that *VGLL1* induction is downstream of TEtra factors

but upstream of other mature markers like *EGFR* and *TP63*. Knock-down of VGLL1 prevented

a complete exit from pluripotency with maintenance of factors like *POU5F1* and *SOX2*, and

blunted expression TE markers including *EGFR*, *ENPEP*, and *NR2F2*. Interestingly, we also

observed reduced expression of TEtra factors *GATA3*, *GATA2*, and *TFAP2C*. These data

suggest that while VGLL1 is not required for the initiation of the TE/trophoblast program, it

belongs to a second wave of factors necessary for the completion of this program in part via

a positive feedback loop onto the TEtra factors.

Such a role is not unique to VGLL1. In the mouse, Eomesodermin (Eomes),

downstream of Cdx2, is not required to initiate the TE program (18, 62) but instead plays a

key role as regulator of TE differentiation and maintenance of TSCs in the extra-embryonic

compartment (17). Embryos lacking Eomes are arrested in the early post-implantation

period, initiate formation of TE (e.g. by cavitation), but fail to form a proper chorion (17). While

EOMES is not involved in human TE/trophoblast development (3, 9), our data suggest that

VGLL1 might play a similar function in reinforcing the TE/TSC identity, downstream of early

key TE lineage regulators and allowing hTSC expansion. In fact, VGLL1 knock-down TE-like

cells adapted to hTSC culture conditions showed decreased expression of GATA2 and

GATA3, and failed to regain cell surface expression of EGFR, demonstrating lack of *bona fide*

hTSC derivation.

Canonical WNT signaling is a crucial pathway to establish and maintain various

epithelial stem cells (23). As mentioned, canonical WNT signaling supports mesoderm

differentiation of BMP4-treated hPSCs (37). VGLL1 is uniquely expressed in the

TE/trophoblast lineage during early human development, thus the up-regulation of

mesoderm genes *EOMES*, *MIXL1,* and *TBXT* in hPSCs as a consequence of ectopic

expression of VGLL1 was unexpected. As in our BMP4-based differentiation protocol,

addition of IWP2 significantly decreased the expression of mesoderm genes in iVGLL1a

hESCs, suggesting that canonical WNT activation, but not mesoderm genes, may be directly

regulated by VGLL1. In fact, temporal activation of the canonical WNT reporter at 48-72h of

BMP4-treatment correlated with the timeline of VGLL1 induction. We also detected active

WNT signaling in DOX-treated iVGLL1a hESCs. However, our data also suggest that VGLL1

alone cannot sustain long-term autocrine activation of the canonical WNT pathway. We

detected VGLL1 occupancy at WNT coreceptors *LRP5/6* as well as the WNT agonist R-

Spondin receptor *LGR4*. Canonical WNT pathway activation is necessary for hTSC and

trophoblast organoid derivation from placental tissue. Our data therefore suggest that

VGLL1 is key in supporting and maintaining the trophoblast stem cell state, potentially via

up-regulation of canonical WNT signaling pathway receptors and coreceptors.

Recently, Yang and colleagues explored the role of VGLL1 and TEAD4 in TE

specification, starting from naïve PSCs, which represent cells of the pre-implantation

embryonic compartment (63). Similar to our results, they demonstrated that loss of VGLL1

led to maintenance of pluripotency genes, blunted expression of TE genes, including *GATA3*,

ENPEP and *NR2F2*, and inability to derive hTSCs. They demonstrated that the VGLL1/TEAD4

complex directly regulates expression of markers such as *GATA3* and *ENPEP* at promoter and

intergenic regions. In contrast, our model utilizes primed PSCs, which represent a post-

implantation epiblast state. Despite the different initial states, our data largely corroborate

their findings with VGLL1 directly regulating markers such as *GATA3* and *ENPEP*.

Interestingly, we showed VGLL1 occupancy at distal enhancer regions of *GATA3* and *EGFR*.

These regions were recently mapped to predicted enhancers active in the human placenta

across gestation (59), suggesting a role for VGLL1 in trophoblast-specific distal regulatory

regions. In fact, when we looked at the presence of these distal enhancer peaks in the Yang

et al data, we found similar peaks, corroborating that they might represent key regulatory

regions for TE and trophoblast identity.

The TE and ICM lineages segregate early during mammalian development and are

distinguished by unique transcriptomes and epigenomes. Both naïve and primed hPSCs

contain bivalent promoters at developmental genes, which maintain them in a

transcriptionally poised state (19, 64, 65). Such bivalent regions are characterized by

H3K4me3, a post-translational histone modification mark associated with active promoters,

and H3K27me3, a marker correlated with transcriptional repression. Global distribution of

H3K27me3 differs between primed and naive hPSCs. However, many of the genes marked

by H3K27me3 and shared between these two pluripotency states are trophoblast regulators,

including GATA3 (64, 65). In their study, Yang and colleagues found a correlation between

loss of VGLL1 and lower levels of H3K27ac, a histone mark associated with active promoters

and enhancers, at TE-specific genes (63). However, the authors did not show a direct link

between VGLL1 expression and H3K27ac levels at TE genes. Moreover, before H3K27 can be

modified by acetylation, the methyl groups need to be removed from the bivalent promoter

containing H3K27me3. We have also observed decreased levels of global H3K27ac in our

primed model (data not shown) and identified KDM6B, one of two H3K27me3 specific

demethylases in mammals, as a potential mediator of VGLL1-dependent chromatin

remodeling. The transcript levels of *KDM6B* were largely dependent on VGLL1 expression

and we observed direct VGLL1 binding on *KDM6B* in our ChIP-seq data. In the murine

blastocyst, KDM6B removes the repressive H3K27me3 at the promoter regions of *Gata3* and

Cdx2 specifically in the TE (60). Loss of *Kdm6b* results in maintenance of H3K27me3 at TE-

specific gene regulatory regions, lower expression of Gata3 and Cdx2 in the TE, and an

impaired ability for the embryos to implant into the uterus. In humans, *VGLL1* and *KDM6B*

are highly expressed in the polar TE of the human blastocyst, which attaches to the uterine

wall during implantation. It remains unclear whether KDM6B is involved in human

implantation, but our results show that chemical inhibition of KDM6B demethylase activity

phenocopies loss of VGLL1 and results in the blunted expression of *GATA3* and *CDX2*, *in*

vitro. Thus, we posit a model whereby VGLL1 directly up-regulates the expression of KDM6B,

which contributes to demethylation of H3K27me3 at bivalent/poised promoters prior to

deposition of the acetyl group by acetyltransferases at trophoblast-specific regulatory

regions to mark a transcriptionally permissive chromatin state (**Fig.7**). Evidence suggests

that, while mouse TE is fully specified in the pre-implantation blastocyst, human TE

commitment might not be completed until post-implantation (51). VGLL1-dependent

regulation of chromatin remodelers like KDM6B might provide a mechanism explaining the

higher plasticity of human post-implantation pluripotent cells.

Overall, our data extend previous findings and add a substantial mechanistic hypothesis on
the role of VGLL1 in TE and trophoblast lineage specification. We also demonstrated that
primed hPSCs and the BMP4-dependent differentiation protocol remain a functional and
practical *in vitro* model to study fundamental aspects of genetic and epigenetic regulation
of TE and trophoblast cell identity in humans (66).

Material and Methods

Human embryo and Placental Tissue Collection. Human embryos were donated to the
research project by informed consent under the UK Human Fertilisation and Authority
Licence number R0162 and independently reviewed by the Health Research Authority's
Research Ethics Committee IRAS projects 308099. Human placental tissues were collected
under a protocol approved by the Human Research Protections Program Committee of the
University of California, San Diego Institutional Review Board; all patients gave informed
consent for the collection and use of these tissues.

Culture of human embryonic stem cells. Human embryonic stem cells H9/WA09 (H9) and
HUES9TOPGFP were cultured and passaged in mTeSR Plus (StemCell Technologies cat. no.
100-1130) on Geltrex (Gibco cat. no. A14133-02)-coated plates as previously described (67).
shRNA H9 hESC lines were previously derived and maintained in 10 µg/ml puromycin (Sigma
Aldrich cat. no. P8833-25MG)(9).

Derivation of dCas9-VPR VGLL1 gRNAs (iVGLL1a) H9 hESCs. Five RNA guides for VGLL1
activation were designed using the sgRNA Designer tool by the Broad Institute (68) and
cloned into the Golden Gate pMA Cas9 Guide vector system (69), modified to incorporate an
Eif1a-Bsd antibiotic resistance cassette for selectable incorporation into a mammalian
cells. Each of the guides were cloned into the U6 promoter driven cloning site on the
plasmid, and sequence verified. Plasmids containing DOX-inducible dCas9-VPR (gift from
Dr. Church) (70), VGLL1-specific guide RNAs, and TOPGFP.mC (Addgene cat. no. 35491) (55)
were introduced in H9 ESCs using the Amaxa Nucleofector II electroporation system
following manufacturer instructions (setting B-016). Briefly, 1 million cells were collected,
resuspended and mixed with 2 µg of plasmid DNA before nucleofection using the Lonza

Human Stem Cell Nucleofector Kit 1 (Lonza cat. no. VPH-5012). Cells were immediately re-

plated, and selection started 48 hours later. iVGLL1a cell lines were maintained with 2 µg/mL

puromycin and 8 µg/mL blasticidin S (Gibco cat. no. A1113903). Ectopic expression of VGLL1

was induced with 2 µg/mL doxycycline (Millipore Sigma cat. no. D3447-500MG) for 4 days.

Trophoblast differentiation of hESC. hESCs were first differentiated into TE-like cells

followed by adaptation to hTSC culture conditions according to a previously published

protocol (67). Briefly, hESCs were dissociated with TrypLE Express (Gibco cat. no. 12604021)

and plated at a density of 10,000 cells per cm^2 on Geltrex-coated plates in mTeSR Plus with

5µM Y27632 (Selleck Chemicals cat. no. S1049). After 24 hours, media was changed to

differentiation media composed of DMEM/F12 (Corning, cat. no. 10-092-CV), 1x ITS

(MilliporeSigma, cat. no. I1884), L-ascorbic acid 64 µg/mL (MilliporeSigma, cat. no. A8960),

sodium bicarbonate 543 µg/mL (ThermoFisher, cat. no. 25080094), 2% BSA (Gemini Bio-

Products, cat. no. 700-110), 10 ng/mL recombinant human BMP4 (R&D Systems cat. no. 314-

BP-010), and 2 µM IWP2 (Selleck Chemicals cat. no. S7085). Media was changed daily. At

day 4, cells were either collected for analysis or re-plated on collagen IV (5µg/mL)-coated

plates (MilliporeSigma cat. no. C0543-1VL) and media changed to iCTB media, composed of

Advanced DMEM/F12 (ThermoFisher, cat. no. 12634028), 1X N2 (ThermoFisher, cat. no.

17502001), 1X B27 (ThermoFisher, cat. no. 17504001), 1X GlutMax (ThermoFisher, cat. no.

35050061), 150 µM 1-thioglycerol (MilliporeSigma, cat. no. M6145), 0.05% BSA (Gemini Bio,

cat. no. 700-110-100), 1% knockout serum replacement (ThermoFisher, cat. no. 10828028),

2 µM CHIR99021 (MilliporeSigma SML1046-25MG), 0.5 µM A83-01 (Tocris, cat. no. 2939), 1

µM SB431542 (Millipore Sigma cat. no. 616464-5MG), 5 µM Y27632 (Selleck Chemicals, cat.

no. S1049), 130 µg/mL VPA sodium salt (MilliporeSigma, cat. no. 676380), 100 ng/mL

recombinant human FGF2 (BioPioneer, cat. no. HRP-0011), 50 ng/mL recombinant human

EGF (R&D Systems, cat. no. 236-EG), 50 ng/mL recombinant human HGF (STEMCELL

Technologies, cat. no. 78019.2), and 20 ng/mL Noggin (R&D Systems cat. no. 6057-NG).

IWP2 was withheld in the BMP4 only treatment. 2 µM IWR1 (MilliporeSigma, cat. no. I0161)

was used as an alternative WNT inhibitor for HUES9TOPGFP. 1 µM GSK-J4 (MedChem

Express cat. no. HY-15648B) was used for KDM6B activity inhibition.

Immunohistochemistry. Formalin-fixed, paraffin-embedded placental tissue sections were stained with anti-KDM6B (Abcam cat. no. ab38113) and Goat anti-Rabbit IgG secondary antibody conjugated with AlexaFluor 488 (Invitrogen cat. no. A11008) following previously published protocol (20). Embryos were thawed according to recommendations from Bourn Hall Clinic, the *in vitro* fertilization clinic coordinating donations. hESCs and embryos were fixed in 4% paraformaldehyde (Electron Microscopy Sciences cat. no. 15710) and stained as previously described(3). The following antibodies were used: anti-VGLL1 (Sigma Aldrich cat. no. HPA042403-100UL), anti-TEAD4 (Abcam cat. no. ab58310), anti-YAP1 (Abcam cat. no. ab52771), anti-NANOG (Company), anti-GATA3, mouse anti-p40 (ΔNp63-specific antibody, clone BC28; Biocare Medical). Embryos were imaged on a Leica SP5 inverted confocal microscope (Leica Microsystems). Cells and placentas were imaged on an ECHO Rebel microscope (ECHO).

Flow cytometry. Flow cytometry was conducted using live cells as previously described(67). Briefly, single cell suspensions were stained with EGFR-APC (Biolegend cat. no. 352906), or APC-conjugated mouse IgG (BioLegend cat. no. 400120) antibodies, in flow cytometry buffer (1% FBS, 2% BSA, and 0.03% sodium azide) for 1 hour at 4°C. Cells were then washed three times in PBS and fixed with 2% PFA (Electron Microscopy Sciences cat. no. 15710). Flow cytometry analysis was performed on a Fortessa X20 flow cytometer (BD Biosciences). Data were visualized using FlowJo Software (v10).

RNA isolation, cDNA preparation, and quantitative real-time PCR. Total RNA was purified using NucleoSpin isolation kit (Macherey-Nagel cat. no. 740955.250) and RNA concentration measured using NanoDrop 2000C spectrophotometer (ThermoScientific). cDNA was prepared from total RNA using the PrimeScript RT-Kit (Takara Bio cat. no. RR037A). Quantitative real-time PCR (qPCR) was performed using TB Green Premix Ex Taq (Takara bio cat. no. RR420A) on a QuantStudio5 thermocycler (AppliedBiosystems). The primer sequences used are listed in **Supplemental Table 2**. Relative expression of each transcript was calculated using ΔΔCT method, normalized to 18S rRNA. Statistical analysis and graphs were generated using GraphPad Prism 10. Bar chart data display mean fold change and standard deviation. Student's t tests were performed to determine significance between two

groups. Two-way ANOVA was performed when analyzing significant differences between
three or more groups. All qRT-PCR experiments were done using three replicates and
statistical tests were performed using log transformed fold change. Level of significance is
represented with asterisks.

RNA sequencing and analysis. Total RNA extraction with mirVana (Ambion cat. no. AM1561)
and library preparation for RNA-sequencing were performed as previously described (71).
Quality control was performed using FastQC (v.0.12.1) and multiQC (v.1.19). Two samples
with a lower signal quality than their replicates were eliminated. Reads were mapped to
GRCh38.p14 (GENCODE release 45) using STAR (v.2.7.9a) (72) and annotated using
featureCounts (subread v.2.0.6). Ensembl genes without at least three samples with 10 or
more reads were removed from analysis. Normalization and differential expression analysis
was performed using the R (v.4.4.0) package DESeq2 (v.1.46.0) (73). Genes with an adjusted
p-value < 0.05 and $\log_2$ fold change > 1 were considered differentially expressed. In iVGLL1a
models, adjusted p-value < 0.05 and $\log_2$ fold change > 0.5 were used to include subtle but
biologically relevant changes. Heatmaps and PCA were generated with Qlucore. Statistical
analysis was performed using a two-group comparison or multi-group comparison with q-
value <0.05. Gene list enrichment analysis was performed with Enrichr on multiple database
as specified in the text (74).

Assay for Transposase-Accessible Chromatin (ATAC)-sequencing. Nuclei extraction and
ATAC-seq library prep ware performed according to previously published protocol (75). The
bcbio-nextgen (v1.2.8) ATAC-seq pipeline was used to process the ATAC-seq reads. The
nucleosome free fraction was normalized using deeptools (v 3.5.1) (76) bamCoverage by
CPM. One sample with a lower signal quality than its replicates, as determined by limma-
voom multi-dimensional scaling plots, was eliminated. Bigwigs for replicates were
combined using wiggletools (v 1.2.11) (77) mean to obtain one average track from biological
replicates. Genomic tracks were visualized using IGV Viewer (v2.11.7). Heatmaps were
generated using deeptools computeMatrix and plotHeatmap. Binding profiles were
generated deeptools computeMatrix and plotProfile. Consensus peaks were annotated
using ChIPpeakAnno (v 3.32.0)(78) with EnsDb.Hsapiens.v86 (v 2.99.0). Chromosome

regions were annotated using ChIPpeakAnno assignChromosomeRegion with
TxDb.Hsapiens.UCSC.hg38.knownGene (v 3.16.0). Differential expression analysis was
carried out using edgeR (v 3.40.2)(79) with limma-voom normalization (80). HOMER (v
5.1)(81) findMotifsGenome.pl was then used to identify transcription factor binding motifs.

Chromatin Immunoprecipitation (ChIP)-sequencing. Chromatin immunoprecipitation
was performed following a previously published protocol(58). Briefly, cells were crosslinked
with 1% formaldehyde (Pierce cat. no. 28906) for 10 minutes at room temperature on a
horizontal rocker. Crosslinking was stopped with 2.5 M glycine. Cells were collected into
15mL conical tubes and centrifuged 1200rpm for 5 minutes at 4°C. Cell pellets were washed
in PBS two times and cell pellets were stored in -80°C until ChIP. Frozen cell pellets were
resuspended in cell lysis buffer with protease inhibitors and incubated at 4°C for 10 minutes.
Following centrifugation, nuclear pellets were resuspended in nuclei lysis buffer with
protease inhibitors and incubated at 4°C for 10 minutes. Covaris shearing buffer was added
and chromatin was sheared in microtube-500 AFA (Covaris cat. no. 520185) using Covaris
E220 Focused-Ultrasonicator with settings 5% duty, PiP 140, 15 minutes in 30 second bursts,
cycles 200. Sodium chloride concentration was adjusted to 150mM and Triton-X to 1%
following shearing. Pre-clearing was done with agarose beads (Cell Signaling Technology
cat. no. 9863S) for 30 minutes at 4°C on a rocker. Input was collected and stored in -20°C.
Cleared lysates were incubated with 6 µg of either anti-TEAD4 (Abcam cat. no. ab58310) or
anti-VGLL1 antibody (Sigma Aldrich cat. no. HPA042403-100UL) overnight at 4°C on a rocker.
Agarose beads (50 µL) were added and incubated at 4°C for 1 hour on a rocker. Lysates were
centrifuged at 14000rpm for 5 minutes at 4°C, and washed five times. DNA-protein-antibody
complexes were eluted in two steps with 0.1 M sodium bicarbonate, 1% SDS, vortexed
briefly, and centrifuged at 7500rpm for 2 minutes at room temperature. Reverse crosslinking
for eluted and input DNA was done with 3 µg/mL RNAse A (ThermoScientific cat. no.
EN0531) and 0.3 M sodium chloride (Invitrogen cat. no. AM9759) on a heat block at 67°C for
5 hours. Proteinase K (Invitrogen cat. no. 100005393) was added (0.2 mg/mL) and incubated
at 45°C overnight. DNA purification was performed with Monarch PCR and DNA Cleanup kit

(NEB #T1030) and submitted for library preparation and sequencing conducted at the IGM Genomics Center, University of California, San Diego, La Jolla, CA.

Adapters were trimmed from reads using Trim Galore (v0.6.10) -q 20 --fastqc --paired. Trimmed reads were aligned to GRCh38 using bowtie2 (v2.4.1) with setting -p 24. samtools (v1.10) were used to sort and index bam files. Peaks were called using MACS2 (v2.2.6) callpeak with settings -f BAMPE -g 2.7e+9 -q 0.05 --keep-dup all --call-summits. DiffBind (v3.16.0) was used to find consensus peaks among replicates using minOverlapmembers = 2. BigWig files were generated with deeptools (v3.5.5) bamCoverage with settings --binSize 20 --smoothLength 60 --effectiveGenomeSize 2913022398 –normalizeUsing CPM --extendReads 150 --centerReads --scaleFactorsMethod None --minMappingQuality 30 --ignoreForNormalization chrM, then replicates were averaged using bigwigAverage with settings -bs 20. Heatmaps were generated with deeptools computeMatrix reference-point --referencePoint center -b 1000 -a 1000 -missingDataAsZero -refPointLabel 0 and visualized using plotHeatmap --kmeans 2. ChIPpeakAnno (v3.14.2) was used to annotate peaksets. Motif analysis was performed using HOMER (v5.1) findMotifsGenome.pl with -size 200. Pie charts of genomic annotations were generated with ChIPpeakAnno pie1 function and genomic element distribution gED. Genomic tracks were visualized using IGV Viewer (v2.11.7). Computationally intensive sequencing analyses were done on the San Diego Supercomputer Center (2025): Expanse through the University of California San Diego(82).

Data availability

All datasets are available in Geo: GSE295121, GSE295122, GSE295124, GSE295126.

Acknowledgments

All sequencing was conducted at the IGM Genomics Center, University of California, San Diego, La Jolla, CA. We thank you Karl Willert for the generous donation of the TOPGFP HUES9 cell line, and Marisa Korody for the IWR1. We are grateful to Dr. Mana Parast for the continuous encouragement and support. The contents of this publication are solely the

687 responsibility of the authors and do not necessarily represent the official views of CIRM or
688 any other agency of the State of California.

Fundings

FS and RC were supported by the Eunice Kennedy Shriver National Institute of Child Health
and Human Development (NICHD) grant R01HD096260 to FS and Diversity supplement to
RC. This research was also supported by the California Institute for Regenerative Medicine
(CIRM) awards DISC0-13757 to FS and DISC0-13816 to HCA. NS was funded by CIRM EDUC4
– 12804 Interdisciplinary Stem Cell Training Grant. KF was partially supported by NIH Grant
UL1TR001442 and MH was partially supported by T32 GM145427. AM was funded by the
CIRM Bridges program. PDH was supported by the UC San Diego OB/GYN and Reproductive
Sciences. Work in the laboratory of K.K.N. was supported by the Wellcome Trust
(221856/Z/20/Z). Work in the laboratory of KKN was also supported by the Francis Crick
Institute, which receives its core funding from Cancer Research UK CC2074, the Medical
Research Council CC2074, and Wellcome Trust CC2074. This research was supported by
NIH grants HD062546, HD0098880 and HD079363 to SP. This publication includes data
generated at the UC San Diego IGM Genomics Center utilizing an Illumina NovaSeq 6000
that was purchased with funding from a National Institutes of Health SIG grant (#S10
OD026929).

Bibliography

1. V. B. Alarcon, Y. Marikawa, Trophectoderm formation: regulation of morphogenesis and gene expressions by RHO, ROCK, cell polarity, and HIPPO signaling HHS Public Access. *Reproduction* **164**, 75–86 (2022).

2. M. Y. Turco, A. Moffett, Development of the human placenta. (2019). https://doi.org/10.1242/dev.163428.

3. P. Blakeley, *et al.*, Defining the three cell lineages of the human blastocyst by single-cell RNA-seq (Development, (2015) 142, 3151-3165). *Dev.* **142**, 3613 (2015).

4. C. Gerri, *et al.*, Initiation of a conserved trophectoderm program in human, cow and mouse embryos. *Nature* **587**, 443–447 (2020).

5. R. M. Roberts, *et al.*, The role of BMP4 signaling in trophoblast emergence from pluripotency. *Cell. Mol. Life Sci.* **79** (2022).

6. N. Nishioka, *et al.*, The Hippo Signaling Pathway Components Lats and Yap Pattern Tead4 Activity to Distinguish Mouse Trophectoderm from Inner Cell Mass. *Dev. Cell* **16**, 398–410 (2009).

7. A. C. Denicol, *et al.*, Canonical WNT signaling regulates development of bovine embryos to the blastocyst stage. *Sci. Reports 2013 31* **3**, 1–7 (2013).

8. K. B. Lee, J. K. Folger, S. K. Rajput, G. W. Smith, Temporal regulation of mRNAs for select bone morphogenetic proteins (BMP), BMP receptors and their associated SMAD proteins during bovine early embryonic development: Effects of exogenous BMP2 on embryo developmental progression. *Reprod. Biol. Endocrinol.* **12**, 1–10 (2014).

9. F. Soncin, *et al.*, Comparative analysis of mouse and human placentae across gestation reveals species-specific regulators of placental development. *Dev.* **145** (2018).

10. C. Gerri, *et al.*, A conserved role of the Hippo signalling pathway in initiation of the first lineage specification event across mammals. *Dev.* **150** (2023).

11. M. Regin, *et al.*, Lineage segregation in human pre-implantation embryos is specified by YAP1 and TEAD1. *Hum. Reprod.* **38**, 1484–1498 (2023).

12. B. A. T. Weatherbee, *et al.*, Distinct pathways drive anterior hypoblast specification in the implanting human embryo. *Nat. Cell Biol.* **26**, 353–365 (2024).

13. H. Sasaki, Roles and regulations of Hippo signaling during preimplantation mouse development. *Dev. Growth Differ.* **59**, 12–20 (2017).

14. M. Azagury, Y. Buganim, Unlocking trophectoderm mysteries: In vivo and in vitro perspectives on human and mouse trophoblast fate induction. *Dev. Cell* **59**, 941–960 (2024).

15. A. Ralston, *et al.*, Gata3 regulates trophoblast development downstream of Tead4
and in parallel to Cdx2. *Development* **137**, 395–403 (2010).

16. H. Sasaki, Position- and polarity-dependent Hippo signaling regulates cell fates in
preimplantation mouse embryos. *Semin. Cell Dev. Biol.* **47–48**, 80–87 (2015).

17. A. P. Russ, *et al.*, Eomesodermin is required for mouse trophoblast development and
mesoderm formation. *Nature* **404**, 95–99 (2000).

18. D. Strumpf, *et al.*, Cdx2 is required for correct cell fate specification and
differentiation of trophectoderm in the mouse blastocyst. *Development* **132**, 2093–
2102 (2005).

19. C. Krendl, *et al.*, GATA2/3-TFAP2A/C transcription factor network couples human
pluripotent stem cell differentiation to trophectoderm with repression of
pluripotency. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E9579–E9588 (2017).

20. Y. Li, *et al.*, BMP4-directed trophoblast differentiation of human embryonic stem
cells is mediated through a ΔNp63+ cytotrophoblast stem cell state. *Dev.* **140**, 3965–
3976 (2013).

21. H. M. Sonnemann, B. Pazdrak, D. A. Antunes, J. Roszik, G. Lizée, Vestigial-like 1
(VGLL1): An ancient co-transcriptional activator linking wing, placenta, and tumor
development. *Biochim. Biophys. Acta - Rev. Cancer* **1878**, 188892 (2023).

22. K. Qin, *et al.*, Canonical and noncanonical Wnt signaling: Multilayered mediators,
signaling mechanisms and major signaling crosstalk. *Genes Dis.* **11**, 103–134 (2024).

23. A. Haegebarth, H. Clevers, Wnt signaling, lgr5, and stem cells in the intestine and
skin. *Am. J. Pathol.* **174**, 715–721 (2009).

24. D. Pinto, H. Clevers, Wnt control of stem cells and differentiation in the intestinal
epithelium. *Exp. Cell Res.* **306**, 357–363 (2005).

25. J. Adjaye, *et al.*, Primary Differentiation in the Human Blastocyst: Comparative
Molecular Portraits of Inner Cell Mass and Trophectoderm Cells. *Stem Cells* **23**,
1514–1525 (2005).

26. L. Aghajanova, *et al.*, Comparative transcriptome analysis of human trophectoderm
and embryonic stem cell-derived trophoblasts reveal key participants in early
implantation. *Biol. Reprod.* **86**, 11–12 (2012).

27. L. Yan, *et al.*, Single-cell RNA-Seq profiling of human preimplantation embryos and
embryonic stem cells. *Nat. Publ. Gr.* (2013). https://doi.org/10.1038/nsmb.2660.

28. M. Krivega, W. Essahib, H. Van de Velde, WNT3 and membrane-associated β-catenin
regulate trophectoderm lineage differentiation in human blastocysts. *Mol. Hum.*
Reprod. **21**, 711–722 (2015).

29. S. Sonderegger, J. Pollheimer, M. Knöfler, Wnt signalling in implantation,

decidualisation and placental differentiation--review. *Placenta* **31**, 839–847 (2010).

30.   Y. Hu, *et al.*, Single-cell analysis of nonhuman primate preimplantation development
in comparison to humans and mice. *Dev. Dyn.* **250**, 974–985 (2021).

31.   H. Okae, *et al.*, Derivation of Human Trophoblast Stem Cells. *Cell Stem Cell* **22**, 50-
63.e6 (2018).

32.   S. Haider, *et al.*, Self-Renewing Trophoblast Organoids Recapitulate the
Developmental Program of the Early Human Placenta. *Stem Cell Reports* **11**, 537–
551 (2018).

33.   M. Y. Turco, *et al.*, Trophoblast organoids as a model for maternal–fetal interactions
during human placentation. *Nature* **564**, 263–281 (2018).

34.   B. Dietrich, S. Haider, G. Meinhardt, J. Pollheimer, · Martin Knöfler, WNT and NOTCH
signaling in human trophoblast development and differentiation. **79**, 292 (2022).

35.   A. S. Brumm, *et al.*, Initiation and maintenance of the pluripotent epiblast in pre-
implantation human development is independent of NODAL signaling. *Dev. Cell* **60**
(2025).

36.   J. Au, *et al.*, Role of autocrine bone morphogenetic protein signaling in trophoblast
stem cells. *Biol. Reprod.* (2021). https://doi.org/10.1093/biolre/ioab213.

37.   D. Kurek, *et al.*, Endogenous WNT signals mediate BMP-induced and spontaneous
differentiation of epiblast stem cells and human embryonic stem cells. *Stem Cell*
Reports **4**, 114–128 (2015).

38.   N. Sah, F. Soncin, Conserved and divergent features of trophoblast stem cells. *J.*
Mol. Endocrinol. **72** (2024).

39.   L. Li, *et al.*, Single-cell multi-omics sequencing of human early embryos. *Nat. Cell*
Biol. 2018 207 **20**, 847–858 (2018).

40.   L. Liu, *et al.*, An integrated chromatin accessibility and transcriptome landscape of
human pre-implantation embryos. *Nat. Commun. 2019 101* **10**, 1–11 (2019).

41.   A. L. Wilkinson, I. Zorzan, P. J. Rugg-Gunn, Epigenetic regulation of early human
embryo development. *Cell Stem Cell* **30**, 1569–1584 (2023).

42.   S. Io, *et al.*, Capturing human trophoblast development with naive pluripotent stem
cells in vitro. *Cell Stem Cell* **28**, 1023-1039.e13 (2021).

43.   G. Guo, *et al.*, Human naive epiblast cells possess unrestricted lineage potential.
Cell Stem Cell **28**, 1040-1056.e6 (2021).

44.   J. K. Cinkornpumin, *et al.*, Naive Human Embryonic Stem Cells Can Give Rise to
Cells with a Trophoblast-like Transcriptome and Methylome. *Stem Cell Reports* **15**,
198–213 (2020).

45.   M. Horii, T. Bui, O. Touma, H. Y. Cho, M. M. Parast, An Improved Two-Step Protocol
for Trophoblast Differentiation of Human Pluripotent Stem Cells. *Curr. Protoc. Stem*
Cell Biol. **50**, 1–21 (2019).

46.   M. Amita, *et al.*, Complete and unidirectional conversion of human embryonic stem
cells to trophoblast by BMP4. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 1–10 (2013).

47.   M. Horii, *et al.*, Human pluripotent stem cells as a model of trophoblast
differentiation in both normal development and disease. *Proc. Natl. Acad. Sci. U. S.*
A. **113**, E3882–E3891 (2016).

48.   F. Soncin, *et al.*, Derivation of functional trophoblast stem cells from primed human
pluripotent stem cells. *Stem Cell Reports* **17**, 1303–1317 (2022).

49.   N. Moya, J. Cutts, T. Gaasterland, K. Willert, D. A. Brafman, Endogenous WNT
signaling regulates hPSC-derived neural progenitor cell heterogeneity and specifies
their regional identity. *Stem Cell Reports* **3**, 1015–1028 (2014).

50.   W. Huang, *et al.*, Uncovering the role of TET2-mediated ENPEP activation in
trophoblast cell fate determination. *Cell. Mol. Life Sci.* **81** (2024).

51.   D. Meistermann, *et al.*, Integrated pseudotime analysis of human pre-implantation
embryo single-cell transcriptomes reveals the dynamics of lineage specification.
Cell Stem Cell **28**, 1625-1640.e6 (2021).

52.   X. Luan, *et al.*, Down-regulation of CCR7 via AKT pathway and GATA2 inactivation
suppressed trophoblast migration and invasion in recurrent spontaneous abortion†.
Biol. Reprod. **102**, 424–433 (2020).

53.   M. Drukker, *et al.*, Isolation of primitive endoderm, mesoderm, vascular endothelial
and trophoblast progenitors from human pluripotent stem cells. *Nat. Biotechnol.* **30**,
531–542 (2012).

54.   C. Dong, *et al.*, A genome-wide CRISPR-Cas9 knockout screen identifies essential
and growth-restricting genes in human trophoblast stem cells. *Nat. Commun. 2022*
13 1 **13**, 1–16 (2022).

55.   D. Horst, *et al.*, Differential WNT activity in colorectal cancer confers limited
tumorigenic potential and is regulated by MAPK signaling. *Cancer Res.* **72**, 1547–
1556 (2012).

56.   B. Saha, *et al.*, TEAD4 ensures postimplantation development by promoting
trophoblast self-renewal: An implication in early human pregnancy loss. *Proc. Natl.*
Acad. Sci. U. S. A. **117**, 17864–17875 (2020).

57.   G. Meinhardt, *et al.*, Pivotal role of the transcriptional co-activator YAP in trophoblast
stemness of the developing human placenta. *Proc. Natl. Acad. Sci. U. S. A.* **117**,
13562–13570 (2020).

58.   P. Home, *et al.*, Altered subcellular localization of transcription factor TEAD4

regulates first mammalian cell lineage commitment. *Proc. Natl. Acad. Sci. U. S. A.*
109, 7362–7367 (2012).

59.   D. M. Owen, *et al.*, Genome-wide identification of transcriptional enhancers during
human placental development and association with function, differentiation, and
disease†. *Biol. Reprod.* **109**, 965–981 (2023).

60.   B. Saha, *et al.*, EED and KDM6B coordinate the first mammalian cell lineage
commitment to ensure embryo implantation. *Mol. Cell. Biol.* **33**, 2691–2705 (2013).

61.   F. Zhou, *et al.*, Reconstituting the transcriptome and DNA methylome landscapes of
human implantation. *Nature* **572**, 660–664 (2019).

62.   H. Niwa, *et al.*, Interaction between Oct3/4 and Cdx2 determines trophectoderm
differentiation. *Cell* **123**, 917–929 (2005).

63.   Y. Yang, *et al.*, VGLL1 cooperates with TEAD4 to control human trophectoderm
lineage specification. *Nat. Commun.* **15** (2024).

64.   B. Kumar, *et al.*, Polycomb repressive complex 2 shields naïve human pluripotent
cells from trophectoderm differentiation. *Nat. Cell Biol. 2022 246* **24**, 845–857
(2022).

65.   D. W. Zijlmans, *et al.*, Integrated multi-omics reveal polycomb repressive complex 2
restricts human trophoblast induction. **24**, 858–871 (2022).

66.   A. S. Seetharam, *et al.*, The product of BMP-directed differentiation protocols for
human primed pluripotent stem cells is placental trophoblast and not amnion. *Stem*
Cell Reports **17. 10.**, /j.stemcr.2022.04.014 (2022).

67.   V. C. Cheung, *et al.*, Current Strategies of Modeling Human Trophoblast Using
Human Pluripotent Stem Cells in vitro. (2023). https://doi.org/10.1002/cpz1.875.

68.   H. K. Kim, *et al.*, Deep learning improves prediction of CRISPR–Cpf1 guide RNA
activity. *Nat. Biotechnol. 2018 363* **36**, 239–241 (2018).

69.   J. Vad-Nielsen, L. Lin, L. Bolund, A. L. Nielsen, Y. Luo, Golden Gate Assembly of
CRISPR gRNA expression array for simultaneously targeting multiple genes. *Cell.*
Mol. Life Sci. **73**, 4315–4325 (2016).

70.   A. Bertero, *et al.*, Optimized inducible shRNA and CRISPR/Cas9 platforms for in vitro
studies of human development using hPSCs. *Development* **143**, 4405–4418 (2016).

71.   R. Morey, *et al.*, Transcriptomic Drivers of Differentiation, Maturation, and Polyploidy
in Human Extravillous Trophoblast. *Front. Cell Dev. Biol.* **9** (2021).

72.   A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

73.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15** (2014).

74. E. Y. Chen, *et al.*, Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. *BMC Bioinformatics* **14** (2013).

75. F. C. Grandi, H. Modi, L. Kampman, M. R. Corces, Chromatin accessibility profiling
by ATAC-seq. https://doi.org/10.1038/s41596-022-00692-9.

76. F. Ramírez, *et al.*, deepTools2: a next generation web server for deep-sequencing
data analysis. *Nucleic Acids Res.* **44**, W160–W165 (2016).

77. D. R. Zerbino, N. Johnson, T. Juettemann, S. P. Wilder, P. Flicek, WiggleTools:
parallel processing of large collections of genome-wide datasets for visualization
and statistical analysis. *Bioinformatics* **30**, 1008–1009 (2014).

78. L. J. Zhu, *et al.*, ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and
ChIP-chip data. *BMC Bioinformatics* **11** (2010).

79. Y. Chen, L. Chen, A. T. L. Lun, P. L. Baldoni, G. K. Smyth, edgeR v4: powerful
differential analysis of sequencing data with expanded functionality and improved
support for small counts and larger datasets. *Nucleic Acids Res.* **53**, 13–14 (2025).

80. C. W. Law, Y. Chen, W. Shi, G. K. Smyth, Voom: Precision weights unlock linear
model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, 1–17 (2014).

81. S. Heinz, *et al.*, Simple combinations of lineage-determining transcription factors
prime cis-regulatory elements required for macrophage and B cell identities. *Mol.*
Cell **38**, 576–589 (2010).

82. S. Strande, *et al.*, Expanse: Computing without Boundaries: Architecture,
Deployment, and Early Operations Experiences of a Supercomputer Designed for the
Rapid Evolution in Science and Engineering. *ACM Int. Conf. Proceeding Ser.* (2021).
https://doi.org/10.1145/3437359.3465588.

 Figure Legends

Figure 1. BMP4 treatment of hESCs induces chromatin remodeling. A. Venn diagram of up-regulated genes following treatment of hESCs with BMP4 with and without IWP2 for 2 days. Gene list enrichment analysis on the KEGG pathway database (2021) for genes uniquely up-regulated with BMP4 alone treatment is presented, a=WNT-mediated b=FGF-mediated c=TGFβ-mediated. **B.** Gene expression levels of markers for the mesoderm lineage across treatment/days by bulk RNA-seq. **C.** Heatmaps illustrating open chromatin loci by ATAC-seq. **D.** Signal intensity of open chromatin at mesoderm- and TE-specific genes. **E.** Heatmap of differentially expressed chromatin modifiers genes across treatment/days.

Figure 2. Loss of VGLL1 prevents completion of the TE/trophoblast program. A. Gene expression levels of early and late trophoblast markers in the first 72h of BMP4/IWP2 treatment of hESCs by qPCR. Statistical analysis relative to 0h **B.** Gene expression levels of early and late trophoblast markers in shVGLL1 compared to Scramble hESCs during BMP4/IWP2 treatment by qPCR. Asterisks represent statistical differences within cell lines compared to d0. Letters represent statistical significance of shVGLL1 compared to same timepoint on Scramble. **C.** Heatmaps of differentially expressed genes between shVGLL1 and Scramble hESCs at d0 and d4 of BMP4/IWP2 treatment by RNA-seq organized by gene cluster analysis. **D.** Bright field images of Scramble and shVGLL1 hESC-derived TSCs. **E.** Cell surface expression of EGFR in hESC-derived TSCs by flow cytometry. **F.** Gene expression levels of TEtra factors and EGFR in hESC-derived TSCs by qPCR. Statistical analysis compared to Scramble. * p≤0.05, ** p≤0.01, *** p≤0.001, **** p≤0.0001.

Figure 3. VGLL1 regulates a TE/trophoblast program independently of BMP4 treatment. A. Schematic representation of the inducible CRISPR-mediated gene activation system, which includes a doxycycline (DOX)-inducible dCas9-VPR and VGLL1-specific guide RNAs. **B.** Bright field images of DMSO (control) and DOX-treated hESCs at day 4. **C.** Cell surface expression of EGFR in DOX-treated iVGLL1a hESCs at day 4 by flow cytometry. **D.** Gene expression levels of lineage markers in DOX-treated iVGLL1a hESCs at day 4 by qPCR. Statistical analysis compared to DMSO. **E.** Heatmap of differentially expressed genes in iVGLL1a hESCs at day 4 of DMSO or DOX treatment. * p≤0.05, ** p≤0.01, *** p≤0.001, **** p≤0.0001

Figure 4. VGLL1 regulates expression of canonical WNT effectors. A. Heatmap of WNT target genes differentially expressed in iVGLL1a hESCs at day 4 of DMSO or DOX treatment. **B.** β-catenin-dependent WNT activation in DOX-treated hESC using a TCF/LEF-GFP reporter plasmid. **C.** Gene expression levels of mesoderm markers in DOX-treated iVGLL1a hESCs with and without IWP2 by qPCR. **D.** Gene expression levels of early and late trophoblast markers in DOX-treated iVGLL1a hESCs with and without IWP2 by qPCR. **E.** Cell surface

expression of EGFR in DOX-treated iVGLL1a hESCs with and without IWP2 at day 4 by flow
cytometry. **F.** Heatmap of differentially expressed genes in iVGLL1a hESCs at day 4 of DMSO
or DOX treatment, with and without IWP2. * p≤0.05, ** p≤0.01, *** p≤0.001, **** p≤0.0001.

Figure 5. VGLL1 directly regulates chromatin modifier gene KDM6B. A. Venn diagram of
concordantly regulated genes downregulated in shVGLL1 clones and up-regulated in DOX-
treated iVGLL1a hESCs. **B.** Genome browser tracks showing VGLL1 and TEAD4 genomic
enrichment peaks (ChIP-seq) and chromatin accessibility peaks (ATAC-seq) for
representative TE gene loci of BMP4/IWP2-treated hESCs at day 4. **C.** Motif analysis using
HOMER showing the significantly enriched DNA-binding motifs at VGLL1-binding sites in
BMP4/IWP2-treated hESCs at day 4. **D.** Co-occupancy analysis by signal density pileups of
VGLL1 and TEAD4 peaks in ChIP-seq of BMP4/IWP2-treated hESCs at day 4. **E.** Genome
browser tracks showing VGLL1 genomic enrichment peaks (ChIP-seq) and chromatin
accessibility peaks (ATAC-seq) for chromatin modifiers, KDM6B and KDM4B loci of
BMP4/IWP2-treated hESCs at day 4. **F.** Gene expression levels of early trophoblast markers
after 4 days of treatment of hESCs with BMP4/IWP2 in the presence of KDM6B inhibitor
GSKJ4 (or DMSO as control). * p≤0.05, ns = non-significant

Figure 6. VGLL1 and KDM6B are highly expressed in the polar TE of pre-implantation
blastocysts. A. Violin plots showing expression levels of VGLL1 and CDX2 in the mural TE,
polar TE, epiblast, and primitive endoderm (PE) cells of d6 blastocysts (data from Zhou *et al*.
2019). **B.** Immunofluorescence staining of human morula and day 5 blastocyst with VGLL1,
TP63, and GATA3 or NANOG. **C.** Violin plots showing expression levels of KDM6B in the mural
TE, polar TE, epiblast, and primitive endoderm (PE) cells of d6 blastocysts (data from Zhou
et al. 2019). **D.** Immunofluorescence staining of KDM6B in week 6 placenta. CTB=
cytotrophoblast, STB=syncytiotrophoblast, CC=cell column

Figure 7. Graphical abstract.

A Upregulated genes_Day 2

Term	Adj. P-value
Gastric cancer[a,b,c]	2.96E-04
Hippo signaling pathway	6.77E-04
Wnt signaling pathway	1.56E-03
Basal cell carcinoma[a]	1.56E-03
Breast cancer[a,b]	6.95E-03
Regulation of actin cytoskeleton	7.32E-03

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7