A SNARE-like mechanism mediates bacterial outer membrane exchange

Carolina Basurto De Santiago[1][¶], Elias J. Topo[1][¶], Pengbo Cao[2,3], Daniel Wall[2][#]

and Beiyan Nan[1][#]

[1]Department of Biology, Texas A&M University, College Station TX 77843, USA

[2]Department of Molecular Biology, University of Wyoming, Laramie, WY 82071, USA

[3]Present address: School of Medicine, Westlake University, Hangzhou, China

Running title: OME molecular mechanism
8

9   [¶]C. B. D. S. and E, J. T. contributed equally to this work

10   #Address correspondence to Beiyan Nan, bnan@tamu.edu; and Daniel Wall,

11   dwall2@uwyo.edu

12   **Keywords:** *Myxococcus xanthus*, single-particle microscopy, cell surface protein, outer

13   membrane exchange, kin recognition, membrane fusion, fusogen

Abstract

Kin recognition, the ability to distinguish self from nonself at the cellular level is critical to multicellular life. *Myxococcus xanthus* is a multicellular bacterium that cooperates among genetically-related cells and reduces exploitation by nonkin through outer membrane exchange (OME) of common goods and toxins. The polymorphic cell surface receptor called TraA and its partner protein TraB mediate kin recognition by OME, but its molecular mechanism remains unknown. Here we show that TraAB induce OME by overcoming the repulsive interaction between cells. Using quantitative microscopy techniques, we determined that TraA and TraB form complexes at a 1:1 ratio and that as few as one intercellular TraAB-TraAB dimer is sufficient to induce OME. We visualized the OME of single protein particles between cells and revealed that OME depends on the free diffusion of outer membrane (OM) contents. Our findings suggest a mechanism that shows analogies to the eukaryotic soluble N-ethylmaleimide-sensitive factor attachment protein receptors (SNAREs), which mediate plasma membrane fusion. *M. xanthus* OME provides a novel pathway that leads to an underlying conserved mechanism for membrane fusion that is a foundation process for multicellularity.

Introduction

The ability to distinguish self from nonself at the cellular level is critical to multicellular life. This process ensures that highly related or clonal cells interact cooperatively while excluding unrelated cells, which are often detrimental to multicellularity. For organisms that build multicellularity by aggregating cells from their environment, self/nonself-recognition, or kin recognition, is of paramount importance for coalescing a cooperative kin collective.

Myxobacteria are a group of aggregative multicellular organisms that possess known systems for distinguishing relatedness at the genus and species levels (Kroos *et al.*, 2025). Specifically, *Myxococcus xanthus* uses the type VI secretion system (T6SS) and outer membrane exchange (OME) for this purpose (Vassallo & Wall, 2019, Vassallo *et al.*, 2020, Gong *et al.*, 2018). The T6SS is used by many Gram-negative bacteria for various types of social conflicts and virulence and is relatively well understood (Cherrak *et al.*, 2019). In contrast, OME is restricted to myxobacteria and is not as well studied.

OME self-recognition occurs through homotypic binding between adjacent cells, mediated by their polymorphic cell surface receptor called TraA and its operonic partner, TraB (Pathak *et al.*, 2013, Cao & Wall, 2017). Following cell-cell recognition, TraAB catalyzes OME, which involves the bidirectional transfer of different outer membrane proteins (OMPs) and lipids between cells (Nudleman *et al.*, 2005, Wall, 2014, Pathak *et al.*, 2012). This process is thought to occur by transient outer membrane (OM) fusion, where TraAB acts as a fusogen on each interacting cell (Cao *et al.*, 2015, Cao & Wall, 2019a). The TraA protein contains a signal peptide (SP), a variable domain (VD) responsible for recognition specificity, a cysteine-rich repeat region that likely functions as a presentation stalk for the VD, and a MYXO-CTERM sorting tag that is likely lipidated to anchor it to the cell surface (Sah *et al.*, 2020, Guo *et al.*, 2025). TraB contains an SP, an OM β-barrel, and an OmpA cell-wall-binding domain. Forward genetic screens suggest that TraAB are the only proteins required for OME (Dey & Wall, 2014, Vassallo *et al.*, 2021).

OME serves two broad functions: mediating cooperation and discriminating

between cells (Sah & Wall, 2020). For cooperation, OME can rescue genetic defects in

recipient cells by transferring a wild-type protein from donor cells (Hodgkin & Kaiser,

1977, Pathak *et al.*, 2012). OME can also rescue lethal OM defects by transferring

lipopolysaccharides (LPS) from healthy donors (Vassallo & Wall, 2016). Additionally,

when individual cells coalesce into tissue-like assemblages, genetically identical cells can

have distinct life histories that result in different adaptations depending on their age

and/or microenvironment exposures. Here, OME facilitates the transition to

homogeneous and synchronized cell populations with respect to their OM components

(Cao & Wall, 2019b), thus facilitating cooperation. Finally, adaptive traits suited to

particular environments can be transferred to naïve siblings via OME (Subedi *et al.*, 2024),

thus preparing or adapting them against future stresses.

OME also functions to discriminate against nonkin. This occurs by the exchange of

dozens of polymorphic lipoprotein toxins that belong to six different families (Dey *et al.*,

2016, Vassallo *et al.*, 2017, Vassallo & Wall, 2019, Weltzer *et al.*, 2025). Therefore, when

partnering cells are clonal, they are not intoxicated because they express a cognate set of

immunity proteins, which themselves are not exchanged. In contrast, when nonclonal

cells fortuitously possess compatible TraA receptors, a second discrimination step occurs,

wherein cells are reciprocally poisoned because they lack a complete set of immunity

proteins. These steps result in an extremely high level of specificity for self-recognition,

ensuring that highly related collectives form for multicellular functions, such as fruiting

body development.

Membrane fusion plays a central role in many biological processes in eukaryotic

cells, but in bacteria there are few examples and little is known. Here, we sought to

address key questions about the molecular mechanism of OME. First, we sought to

understand the nature of TraAB multimeric complexes that form upon cell-cell contact

and OME engagement. Following the coalescing of receptors, these complexes are readily

visualized as foci when using functional fluorescent protein fusions to TraA or TraB.

However, the stoichiometry of the intercellular TraA junction and the intracellular TraAB

complexes remains unknown. Second, since client OME cargos do not contain specificity

factors for transfer, but must be diffusible in a fluid OM, we aimed to measure the fluidity

of *M. xanthus* OM. Third, do these foci junctions serve as conduits for the transfer of client

proteins, and what trajectories do these proteins follow upon OME? Using

photobleaching and single molecule tracking techniques, we addressed these questions

and proposed an updated model of OME in myxobacteria.

Fig. 1. TraAB facilitate OME by overcoming the repulsive interactions between cells. A) cryoEM images show that juxtaposed *pilA⁻* cells do not make direct OM-OM contacts. In the same cell, the OM is relaxed at non-contact sites while stressed at cell-cell contact sites. **B)** Cell-cell contact induces stress compression of periplasms at contact sites. Periplasmic thickness was measured as the distance between the centers of the OM and IM densities from seven contacting cell pairs, 30 measurements/pair. The *p* value was calculated using a one-way ANOVA test between two unweighted, independent samples. **C)** Juxtaposed *pilA⁻* cells that over produce TraAB frequently form direct OM-OM contacts that lead to OM fusion.

Results

Repulsive interaction between cells obstructs OME

Cells lacking functional TraAB or expressing incompatible TraA variants do not exchange

their OM contents (Wei *et al.*, 2011). This raises the question; why are TraAB required and

physical cell-cell contacts alone are insufficient for OM fusion? To answer this, we investigate

cell-cell contacts using cryogenic electron microscopy (cryoEM). To avoid the potential

interference from extracellular polysaccharides (EPS), we grew *pilA⁻* cells that do not

produce EPS (Yang *et al.*, 2010, Zhou & Nan, 2017) in liquid culture, concentrated cells by

centrifugation, and captured micrographs of *M. xanthus* cells in close contact, where we

found cell groups of various sizes. As OME does not occur in liquid, these cells displayed the

character of OMs before fusion (**Fig. 1A**).

Analyzing these micrographs, we found two remarkable features. First, regardless of

group size, physical interaction between OMs was not observed (**Fig. 1A**). The distances

between the centers of the OM densities at side-to-side cell contact sites varied between 20.4

nm and 34.5 nm, with an average distance of 28.8 ± 3.4 nm (measure from three pairs of cells,

30 measurements/pair). Therefore, the interaction between the OMs is under the regime of

electrostatic repulsion, likely driven by the strong negative charge on lipopolysaccharide

molecules, and beyond the range of van der Waals attraction, $0.3 - 0.5$ nm (Batsanov, 2001).

Such repulsion is stronger than the one between weakly charged liposomes $(3 - 10$ nm

distance) that prevents their fusion in aqueous media (Jahn *et al.*, 2024). Hence juxtaposed

cells are always separated by a thin water layer and the repulsion between cells physically

obstructs OM fusion. Second, the periplasms appeared compressed at cell-cell contact sites

(**Fig. 1A**). Strikingly, in the cells that contact others only on one side, only the contact sides

displayed compressed periplasms with the thickness (the distance between the centers of the

OM and IM densities) of 30.4 ± 3.8 nm, while the non-contact sides retained relaxed

periplasms of 37.5 ± 5.9 nm thickness (measured from the same seven pairs of cells, 30

measurements/pair, for both measurements, **Fig. 1B**). This result shows that cell-cell contacts

increase the surface tension of OMs. Therefore, similar to plasma membranes that do not fuse

spontaneously (Rand & Parsegian, 1984), repulsion must be overcome to bring OMs closer,

which poses a major energy barrier in OM fusion.

TraAB induce OME by overcoming the repulsive interaction between cells

To visualize how TraAB facilitate OME, we subjected a $pilA^-$ strain that expressed $traAB$

ectopically by the strong $pilA$ promoter ($traAB^{OE}$) (Pathak $et\ al.$, 2012), to cryoEM imaging. In

stark contrast to its parent strain that maintained a nearly constant OM spacing (**Fig. 1A**), the

OM-OM distance between juxtaposed $traAB^{OE}$ cells varied considerably. Notably, these cells

frequently made close OM contacts that often culminated in an apparent OM fusion-like

configuration (**Fig. 1C**). Hence, TraAB-TraAB complexes between cells overcomes repulsive

interactions between juxtaposed OMs and thus induces OME.

Stoichiometry of the intercellular TraA foci

TraAB are thought to function as fusogens to catalyze OM fusion between cells. By using

fluorescent fusion reporters to TraA or TraB they were found to coalesce into foci aggregates

upon cell-cell contact (Cao & Wall, 2019a). Here we sought to address (i) how many TraA

molecules are present in these foci? (ii) Do all the TraA foci contain the same stoichiometry,

and (iii) do interacting cells contribute the same number of TraA proteins in each focus? To

answer these questions, we used a functional monomeric super folder green fluorescent

protein (GFP)-labeled TraA (TraA-GFP) expressed from the heterologous $pilA$ promoter and

placed it in a $\Delta traA$ strain (Cao & Wall, 2019a). We mixed this strain with wild-type

(unlabeled) cells at a 1:2 ratio and spotted the cell mixture on a 1.5% agar surface. As reported

(Cao & Wall, 2019a), TraA-GFP distributed uniformly throughout the OM of isolated cells.

However, upon cell-cell contact, we observed two types of TraA foci at contact junctions,

bright foci occurred between two labeled cells and dim foci between labeled and unlabeled

cells (**Fig. 2A**).

The intensities of TraA-GFP foci appeared heterogeneous, regardless of whether only

one or both cells were labeled (**Fig. 2A**), suggesting that individual TraA foci do not adopt

the same stoichiometry. We then subjected the TraA-GFP foci to photobleaching using a 488-

nm laser (0.5 kW/cm²) and monitored the decrease of fluorescent signals continuously at 67

Hz (15 ms/frame). The intensity of these foci decreased in discrete steps of the same size,

which we define as Δi that reflected the photobleach of one GFP molecule (**Fig. 2B**).

Occasionally we saw larger bleach steps where the intensity decreased by $2\Delta i$, indicating that

two molecules were bleached simultaneously (**Fig. 2B**). Therefore, the number of Δi bleach

steps equal the number of TraA-GFP molecules within the focus (**Fig. 2B, Movie S1**).

Fig. 2. TraA forms heterogeneous foci at cell-cell contact sites. **A)** Representative images of the TraA-GFP foci. The Δ*traA traA-gfp* cells were mixed with wild-type (unlabeled) at a 1:2 ratio. White arrows indicate foci formed between labeled cells and red arrows between labeled and unlabeled cells. **B)** Examples of the photobleached foci that contain 4 and 6 TraA-GFP molecules, respectively. The decrease step of fluorescence corresponds to one GFP molecule being photobleached (*Δi*). **C)** The number of TraA-GFP molecules in each focus between labeled cells (left, n = 174) and between labeled and unlabeled ones (right, n = 204).

Consistent with the variation of their fluorescent intensities, the 174 foci between labeled

cells lost fluorescence after 2 to 10 steps of photobleaching, indicating that these foci

contained 2 to 10 TraA-GFP molecules (**Fig. 2B, Movie S1**). The numbers of TraA-GFP

molecules in each focus displayed a typical Gaussian distribution. Importantly, we never

observed any foci that was bleached in a single step of Δi, suggesting that at least one TraA

molecule in each juxtaposed cell was required to establish the intercellular junction. This

result was consistent with the observation that TraA-GFP expressing cells cannot form TraA

foci when it contacts a $\Delta traA$ cell (Cao & Wall, 2019a). In addition, 46% of these foci contained

odd numbers of TraA molecules, indicating that the formation of these foci does not

necessarily require the same number of TraA molecules from two juxtaposed cells.

Among the 204 foci found between labeled and unlabeled cells, the number of TraA-GFP

molecules also showed a nearly symmetric distribution between 1 and 6, roughly half as

many as detected in the foci between two labeled cells (**Fig. 2C**). The detection of foci that

contained only one TraA-GFP molecule further confirmed that a single TraA molecule from

each contacting cell was sufficient to establish an intercellular TraA junction. Taken together,

TraA foci at cell-cell contact sites contain molecules from both juxtaposed cells where each

cell may contribute different numbers of TraA molecules.

TraA and TraB form complexes at a 1:1 ratio

TraA requires its partner OMP TraB to form functional complexes (Pathak *et al.*, 2012, Cao &

Wall, 2017). Although TraB does not play a specificity role in kin recognition, TraA and TraB

colocalize into the foci between adjacent cells in an interdependent manner, suggesting that

these proteins form intracellular complexes (Cao & Wall, 2019a). However, the stoichiometry

of these complexes is unknown. To address this question, we used a strain that expressed

both TraA-mCherry and TraB-GFP and photobleached both mCherry and GFP

simultaneously using Hamamatsu W-VIEW GEMINI™ beam-splitter. TraA and TraB

colocalized in the foci between juxtaposed cells (**Fig. 3A**). To simplify analysis, we recorded

the photobleach of both fluorophores at 5 Hz (200 ms/frame, **Movie S2**). Using this method,

we found that the numbers of TraA-mCherry in each focus varied between 2 and 12 in the 59

analyzed foci (**Fig. 3B**), consistent with the photobleach result from TraA-GFP (**Fig. 2C**).

Importantly, the TraB-GFP molecules displayed a similar distribution pattern (**Fig. 3B**). We

further investigated the correlation between TraA and TraB in each focus and found that each

TraA molecule correlated with 0.98 TraB molecules, indicating that TraA and TraB form

complexes at a 1:1 ratio (**Fig. 3C**).

Fig. 3. TraA and TraB form intracellular complexes at a 1:1 ratio. **A)** TraA-mCherry and TraB-GFP colocalize in the foci between adjacent cells. **B)** From 59 foci analyzed, the number of TraA and TraB proteins in each focus show similar distribution patterns. **C)** The correlation between the numbers of TraA and TraB in individual foci suggests a 1:1 ratio in the intracellular TraAB complexes. In each focus, the number of TraB is plotted against that of TraA. The numbers of foci that have the same TraA:TraB ratio are indicated by color.

M. xanthus **contains a highly fluid OM**

The model bacterium *Escherichia coli* has a rigid OM (Rojas *et al.*, 2018, Sun *et al.*, 2022). As a

consequence, the lateral mobility of OMPs is restricted due to the formation of large OMP

islands and the selective presence of bulky LPS molecules in the outer leaflet (Rassam *et al.*,

2015, Benn *et al.*, 2021). In contrast, *M. xanthus* cells are flexible and can exchange a broad

range of OM molecules efficiently, including OMPs, LPS, and dyes (Ducret *et al.*, 2013,

Vassallo *et al.*, 2015, Pathak *et al.*, 2012). Since heterologous proteins, like a lipidated GFP, are

readily exchanged as cargos, OME does not require other *cis* specificity sequences for

transfer. Instead, these cargos are likely transferred between cells by diffusion, which

requires a fluid OM (Cao & Wall, 2020, Cao & Wall, 2019a). However, the fluidity of *M.*

xanthus OM has not been measured.

Fig. 4. SS$_{OM}$-PAmCherry diffuses rapidly in the *M. xanthus* OM. **A)** Single PAmCherry particles diffuse freely in both the OM and IM. The overall distribution of SS$_{OM}$- and SS$_{IM}$-PAmCherry is displayed using the composite of 100 consecutive frames taken at 15-ms intervals. Single-particle trajectories of PAmCherry were generated from the same frames. Individual trajectories are distinguished by colors. **B)** The linear relationship between MSD and Δt indicates that PAmCherry diffuses freely in both OM (n = 1518) and IM (n = 2848). **C)** PAmCherry shows similar diffusion coefficients in OM (n = 1518) and IM (n = 2848).

To measure membrane fluidity, we used the diffusion rate of OMPs as a proxy. Here, we

used a lipoprotein where a type II signal sequence was fused to the fluorescent protein

mCherry and expressed as a recombinant SS$_{OM}$-mCherry reporter in *M. xanthus.* This simple

lipoprotein tag was sufficient for the OM localization of mCherry and enabled it to be

transferred by OME (Wei *et al.*, 2011). We reengineered this strain by replacing mCherry with

the photoactivatable mCherry (PAmCherry). mCherry and PAmCherry are of the same size

and share 95.8% protein sequence identity. We used a 405-nm excitation laser (0.2 kW/cm^2,

0.1 s) to activate the fluorescence of a few labeled SS_{OM}-PAmCherry particles randomly (here

we refer isolated fluorescence spots as single particles rather than "single molecules" because

each spot could contain more than one molecule, albeit only the fluorescence of one molecule

was activated). We imaged the activated particles at 67 Hz using a 561-nm laser with total

internal reflection fluorescence (TIRF) microscopy (Fu *et al.*, 2018, Nan *et al.*, 2015, Nan *et al.*,

2013, Pogue *et al.*, 2018, Ramirez Carbo *et al.*, 2024, Zhang *et al.*, 2020). Using this setting, only

a thin section of each cell close to the coverslip was illuminated, which included the cell

surface. To exclude the potential influence of cell-cell contact, we only analyzed the dynamics

of SS_{OM}-PAmCherry particles in isolated cells. Using single particle tracking photo-activated

localization microscopy (sptPALM), we identified 1,518 fluorescent particles that remained

in focus for 4 - 12 frames (0.4 - 1.2 s) and plotted their mean squared displacements (MSDs)

against time lag (Δt). The linear relationship between MSD and Δt indicated that single SS_{OM}-

PAmCherry particles moved by typical diffusion (**Fig. 4A, B, Movie S3**). The diffusion

coefficient (*D*) of the entire population was 0.19 ± 0.04 μm^2/s (n = 1518, **Fig. 4C**), in the same

order of magnitude as the unrestricted diffusion of proteins in the cytoplasm, IM, and

periplasm in other bacteria, such as *E. coli* and *Bacillus subtilis* (Parry *et al.*, 2014, Lucena *et al.*,

2018, Tran *et al.*, 2024). Importantly, the diffusion of SS_{OM}-PAmCherry was about 10-fold

faster than the restricted diffusion of *M. xanthus* proteins in the IM and periplasm (Fu *et al.*,

2018, Nan *et al.*, 2013, Zhang *et al.*, 2023b, Ramirez Carbo *et al.*, 2024), suggesting that SS_{OM}-

PAmCherry diffuses with little restriction in the OM.

The bacterial IM is known to be fluid, where many proteins diffuse freely (Lucena *et al.*,

2018). As a control, we engineered a strain that localizes mCherry to the IM (SS_{IM}-mCherry)

(Wei *et al.*, 2011) by replacing mCherry with PAmCherry. As expected, we found that the

recombinant SS_{IM}-PAmCherry also diffused freely, with a *D* value of 0.22 ± 0.02 μm^2/s (n =

2848), close to that of SS$_{OM}$-PAmCherry (**Fig. 4, Move S4**). Taken together, like the IM, the

OM of *M. xanthus* is highly fluid and such fluidity provides the mechanistic foundation for

OME.

OME depends on the diffusion of OM components

OME between *M. xanthus* cells has been imaged in real-time (Ducret *et al.*, 2013, Cao & Wall,

2019a). However, the mechanism of OME is not fully understood. For instance, does OME

depend on the diffusion of the client cargos, or the exchange of large OM fragments? To

investigate the dynamics and trajectories of client proteins during OME, we visualized and

tracked single SS$_{OM}$-PAmCherry particles. If OME requires diffusion, then SS$_{OM}$-PAmCherry

particles should quickly travel between cells. Alternatively, if cells exchange bulk OM

fragments, the exchange of SS$_{OM}$-PAmCherry particles should be slow due to the size of such

fragments.

To test these possibilities, we used cells that expressed SS$_{OM}$-PAmCherry as donors and

Δ*traA* P$_{pilA}$-*traA-gfp* cells as recipients. We mixed these strains at a 1:1 ratio on a 1.5% agar

surface. To increase the chance of capturing OME, we first exposed cells under the 405-nm

laser for 2 s, where the majority of PAmCherry was photoactivated (Fu *et al.*, 2018). We then

excited the fluorescence of PAmCherry and GFP using the 561-nm and 488-nm lasers,

respectively, and imaged both channels simultaneously using the beam-splitter. As SS$_{OM}$-

PAmCherry particles moved rapidly (**Fig. 4**), we had to track their movements at a high frame

rate (67 Hz). For this reason, we could only image each field for a few seconds before the

reporters' photobleach (**Fig. 2**). In these short time windows, OME was rarely visualized.

Nevertheless, we captured OME of 39 SS$_{OM}$-PAmCherry particles (**Fig. 5A-C, Movie S5-S7**).

Their MSD showed a linear relationship with Δ*t*, with a *D* value of 0.21 ± 0.07 μm^2/s (n = 39),

close to the intracellular diffusion rate of SS$_{OM}$-PAmCherry particles. These results indicate

that cell-cell contact does not hinder OMP diffusion, and that OME depends on the diffusion

of OM content, rather than the transfer of large OM fragments (**Fig. 5D**).

By closely tracking particle trajectories during OME, we found six cases where SS$_{OM}$-

PAmCherry particles transferred to recipient cells and then returned to the donors (**Fig. 5A,**

Movie S5). This observation was consistent with OME occurring bidirectionally, where

'recipients' can also act as donors. Strikingly, we identified five SS$_{OM}$-PAmCherry particles

being transferred between more than two cells. For instance, the particle shown in **Fig. 5B**

Fig. 5. OME requires cargo diffusion. **A - C)** Example trajectories of single SS$_{OM}$-PAmCherry particles during OME. The cells expressing SS$_{OM}$-PAmCherry (magenta) and TraA-GFP (green) were mixed at a 1:1 ratio. Arrows point to the exchanged particles. On the right side of each time-lapse series, the outlines of the donors and recipients are marked with purple and green dotted lines, respectively. The trajectories of particles are marked with black segmented lines. Their start and end positions are marked with purple and blue circles, respectively. Scale bars, 1 µm. The uncropped videos for these OME events are Movies S5, S6, and S7. In A) and B), the initial positions of the donor cell poles are marked with white dashed lines in the time-lapse images. In C), the positions of two TraA foci are marked with green circles on the right panel. **D)** The linear relationship between MSD and Δt of the 39 transferred particles indicates they diffuse freely during OME.

and **Movie S6** passed three cell-cell boundaries. Thus, through TraAB-mediated kin

recognition, client proteins can be serially transferred between clonemates to achieve long-

range OME among multiple adjacent cells.

TraA foci do not correlate as the channels for OME

Here we sought to investigate the role of TraAB complexes in OME and if they serve as

transfer portals. To answer these questions, we followed the trajectories of single-particle

OME events between cells that made lateral contacts. Such juxtaposed side-by-side cells

frequently formed multiple TraA foci, which allowed us to ask whether these foci correlated

as the sites where SS_{OM}-PAmCherry particles crossed the cell-cell boundaries. However, out

of the seven SS_{OM}-PAmCherry particles that transferred through lateral side-by-side cells,

none of them passed through the TraA foci (**Fig. 5C, Movie S7**). These observations strongly

suggest that TraAB complexes do not form transfer channels for OME.

A SNAREs-like mechanism for TraAB-mediated OM fusion

To understand how one intercellular TraAB-TraAB dimer induces OM fusion, we used

AlphaFold (Jumper *et al.*, 2021) to predict the structures of TraA and TraB. Using the

sequences of the processed proteins (amino acids 37 – 684 for TraA, excluding its SP and the

cleaved MYXO-CTERM (Sah *et al.*, 2020) and 23 – 543 for TraB, excluding its SP), AlphaFold

generated highly-confident structure models for both proteins in their monomeric states (**Fig.**

6A). TraA monomer forms a dumbbell-like structure where the N-terminal VD, containing

at least eight β-sheets, forms the first dumbbell head. The shaft is an extended cysteine-rich

region, which makes a 180° turn at the end, forming the second dumbbell head (**Fig. 6A**). This

region, containing more than 70 cysteines, forms at least eight β-hairpins that each consists

of two antiparallel β-strand connected by a short loop (**Fig. 6A**). We predict that at cell

surfaces where thiol groups are oxidized, this region is stiffened by dozens of disulfide

bonds. Consistent with a previous finding that the VD determines the recognition between

TraA variants (Cao & Wall, 2017), the AlphaFold structure showed two TraA molecules

interacting through their VDs, forming a head-to-head dimer (**Fig. 6A**). Strikingly, the length

of a TraA dimer is ~ 28 nm, matching the distance between OMs in tightly packed cell groups

(**Fig. 1A**).

Fig. 6. Predicted structures of TraA, TraB, and their complexes suggest a mechanism for OM fusion comparable to the eukaryotic SNARE system. **A)** The predicted structures of TraA, TraB, and their complex. Structure elements are colored according to their pLDDT (predicted local-distance difference test) scores, which indicate the prediction confidence. **B)** A proposed bulge-and-slide mechanism for TraAB-mediated OM fusion. When two cells with compatible TraA molecules make contact, TraA molecules form head-to-head dimers, first through their VDs, and then TraB anchors such TraA dimers to PG. As the juxtaposed cells slide against each other, the two TraAB complexes move in opposite directions, pulling the OMs. Such mechanical stress will bring the OMs of adjacent cells closer and thus trigger OME. **C)** A model of SNARE-mediated plasma membrane fusion in eukaryotic cells. Assembly of SNAREs requires four different SNARE motifs on three or four separate proteins. Once a SNARE complex forms, the four coiled-coil domains transform into a four-helix bundle that "zips" the two membranes together.

The TraB protein consists of a N-terminal β-barrel and a C-terminal OmpA-like domain,

linked together by short thrombospondin type 3 (TSP3) repeats predicted to bind calcium

(Dai *et al.*, 2017) (**Fig. 6A**). Consistent with their colocalization and 1:1 stoichiometry (**Fig. 3**),

AlphaFold also predicted a 1:1 TraAB complex in which the second dumbbell head of TraA

is positioned on top of the β-barrel in TraB (**Fig. 6A**). However, AlphaFold did not predict

significant conformational changes in TraA after its dimerization and interaction with TraB

(**Fig. 6A**). Nevertheless, because TraB is diffusive in isolated cells but colocalizes with TraA

in the relatively stationary intercellular foci (Cao & Wall, 2019a), TraA dimerization could

introduce conformational changes in TraB, which then promotes binding to peptidoglycan

(PG) and thus restrains the diffusion of the TraAB-TraAB dimers.

Based on these predictions and our findings, we propose a model for TraAB-mediated

OM fusion that has similarities to the eukaryotic soluble N-ethylmaleimide-sensitive factor

attachment protein receptors (SNAREs) that mediate vesicle membrane fusion. To facilitate

fusion, membranes must be brought close together, and one way to do this is to minimize

their contact area. Eukaryotes utilize membrane-anchored SNARE proteins to establish

point-to-point contacts that perturb the bilayer structure and thereby lower the energy barrier

for fusion. Once SNARE proteins from juxtaposed membranes establish contact, the SNARE

complex tighten their coiled-coil domains into bundles, mimicking a zipper that pulls the

membranes together until they merge (Jahn *et al.*, 2024) (**Fig. 6D**). Through this mechanism,

one to three SNARE complexes are sufficient to catalyze membrane fusion (Jahn *et al.*, 2024,

Sinha *et al.*, 2011, Shi *et al.*, 2012, van den Bogaart *et al.*, 2010). Similarly, we propose that when

two cells that express compatible TraA variants approach each other, TraA molecules

establish point-to-point contacts between them. Diffusing TraA molecules can then form

intercellular dimers in a head-to-head orientation, which is consistent with TraA foci

formation between topologically opposed OMs. The TraAB complexes could anchor to PG

through the OmpA domain in TraB and thus forms relatively stationary junction foci like

SNARE complexes (**Fig. 6B**). While the TraA molecules in a dimer are not likely to form

SNARE-like zippers (**Fig. 6D**), when juxtaposed cells slide against each other, the movements

of cells pull the PG-anchored TraAB complexes toward opposite directions that tugs on the

OM and generates bulges at the base of each complex (**Fig. 6B**). As the cells continue to slide,

due to the rigidity of TraA, such stressed OM bulges are primed for membrane fusion as they

are forced into juxtaposed OMs, resulting in OM fusion between cells (**Fig. 6B**). In agreement

with our model, transmission electron microscopy (TEM) of cryo-fixed thin sections showed

that cells overexpressing TraAB contacted each other through OM bulges (Vassallo *et al.*, 2015)

(**Fig. 6C**).

Discussion

Multicellular organisms independently evolved from unicellular ancestors at least 25 times

during evolution (Grosberg & Strathmann, 2007). Myxobacteria achieve multicellularity

through the aggregation of single cells (Kroos *et al.*, 2025). While forming aggregates is

relatively straightforward, it is less understood how individual cells from genetically mixed

populations establish stable multicellular structures. For instance, how do individual cells

maintain cooperative cell-cell interactions while keeping cheaters at bay (Velicer *et al.*, 2000)?

Myxobacteria achieve the transition from solitary, single-cell life into cooperative,

multicellular tissues through kin recognition and the exchange of cellular contents (Sah &

Wall, 2020). Specifically, the homotypic binding by their polymorphic cell surface TraA

protein and its cohort protein TraB, leads to OM content sharing and further discrimination

against nonself by OME of polymorphic toxins. To better understand the mechanism of OME

we used quantitative microscopy and single molecule tracking. We showed that the ratio of

TraA and TraB in cell-cell junctions is 1:1. Moreover, we showed that overexpression of the

chromosomal copy of the *traAB* operon generates TraA foci of various stoichiometry, ranging

from 2 to 12. Our results confirmed that each partnering cell contributes at least one TraAB

complex to form a functional junction. Because the native, low expression of *traAB* is

sufficient for OME (Wei *et al.*, 2011), we postulate that the minimum number of TraAB

complexes in junctions for OME is two. Indeed, the foci that contained two TraAB complexes

only accounted for ~5% of all the foci we detected, which explains why such foci were hard

to image when TraAB were expressed with the native promoter (Cao & Wall, 2019a, Vassallo

et al., 2015). Thus, overexpression increased both the number and size of such foci. The wide

range of stoichiometry and the fact that two partnering cells do not necessarily contribute

equal numbers of TraAB proteins in foci junctions, suggest that TraAB do not form

intercellular protein channels with defined structures.

Instead, our results invoke TraAB serving as fusogens to transiently catalyze the OM

fusion between cells and thus allow rapid and facile cargo exchange. This model presumes

that the OM of *M. xanthu*s is fluid, distinct from the rigid OMs from other Gram-negative

bacteria such as *E. coli* (Cao & Wall, 2020). By tracking single molecules of the lipidated SS$_{OM}$-

PAmCherry, we showed this reporter rapidly diffuses in the OM. This analysis thus

confirmed *M. xanthus* OM fluidity, showing that diffusion rates in the OM were similar to

unrestricted protein diffusion rates in the cytoplasm, IM, and periplasm in other bacteria. It

is unclear how widely such fluid OMs exist in Gram-negative bacteria. Nevertheless, our

method can be easily applied to measure OM fluidity in other organisms.

Further support of the membrane fusion comes from tracking OME at single-particle

resolution. Here SS$_{OM}$-PAmCherry particles were observed to transfer from donor into

recipient cells at rates that are nearly identical to their diffusion rates. Strikingly, particles

were also observed to enter recipient cells and then return to the original donor, supporting

prior observations of bidirectional transfer mediated by OM fusion (Cao & Wall, 2019a).

Finally, in tracking experiments, particles also traveled across multiple cell boundaries when

those cells were adjacent to one another. This result indicates that multiple cells can

simultaneously form fusion junctions, which are in a tissue-like state. This latter point is also

supported by prior findings of serial cell-cell transfer of SitA toxins (Vassallo *et al.*, 2017).

The last question we addressed is whether TraAB foci serve as transfer portals for

OME. However, our observations did not find a correlation between TraA foci and single

particle transfer of the SS$_{OM}$-PAmCherry reporter. Instead, transfer occurs at various sites

along juxtaposed cells, which are not directly correlated with TraAB foci. Given these

findings, we propose TraAB foci serve as sites to nucleate membrane fusion. Once

"handshake" recognition occurs between TraAB complexes of juxtaposed cells, a small slide

or movement between cells is sufficient to bring the OMs together by pulling forces that

generate bulges on OMs, as observed in TEM micrographs of cell cross-sections (Vassallo *et*

al., 2015). Additionally, fluorescent time-lapse microscopy of gliding *M. xanthus* cells engaged

in OME shows the functional TraA-mCherry reporter dynamically coalescing in a zipper-like

fashion along the lengths of the interacting cells (Cao & Wall, 2019a). Taken together, we

envision TraAB initiating an OM fusion junction that widens along the cell length. Fission

occurs when cells move apart from one another.

The outer leaflet of the *M. xanthus* OM contains LPS, consisting of the lipid A membrane

anchor and extended aqueous exposed polysaccharide O-antigen repeats (Perez-Burgos *et al.*,

2019). Additional types of lipids may also reside in their outer leaflet, but that is not fully

understood (Lorenzen *et al.*, 2014). Since TraAB has no known connection to the IM or

cytoplasmic energy sources, we propose that when cells slide against each other, the

mechanical force of cell movements, an essential requirement for OME (Wei *et al.*, 2011),

could pull OMs outward through the rigid TraAB-TraAB dimer and eventually cause OM

fusion (**Fig. 6B**). Our model shows similarities to the eukaryotic SNARE mechanism in how

membranes are brought together to overcome the energy barrier that obstructs membrane

fusion (**Fig. 6D**). However, in contrast to the SNARE complexes that require four SNARE

motifs from three to four proteins, the TraAB-TraAB dimer is simpler but contains the same

essential structural elements: the interaction domains in TraA and the membrane anchor in

TraA and TraB (**Fig. 6A**). Different from the SNARE complexes, which pull membranes

together through large-scale conformational changes (**Fig. 6D**), the energy for bringing

membranes in proximity comes from sliding movements between cells by either motility or

cell growth (**Fig. 6B**). Nevertheless, we do not rule out the possibility that conformational

changes in TraA could also play important roles in OM fusion. In this regard, a candidate

conformational switch within the diverse TraA family is the invariant Ser-Cys-Asn-Cys-Cys-

Pro motif found in the C-terminal stalk region (Cao *et al.*, 2019). Here the adjacent Cys

residues may form a sterically constrained disulfide bond that brakes upon tension by TraA-
TraA receptors adhering as cells move in opposite directions.

Our model explains two important observations. First, whereas TraA and TraB do not
depend on each other to localize to OMs, their function as cell-cell adhesins and for the
formation of junction foci requires both proteins. In this case, TraA is required for kin
recognition through its VD to initiate intercellular dimer formation, while TraB is essential to
anchor such dimers to PG and to restrain their diffusion. This latter point is supported by the
finding that deletion of the OmpA domain from TraB results in a truncated protein able to
form foci and act as an adhesion with TraA, but is not functional for OME (Balagam *et al.*,
2021). Second, OME requires cell motility (Wall *et al.*, 1998, Wall & Kaiser, 1998, Wei *et al.*,
2011). Our model predicts that the relative movements pull on OMs through the rigid TraA-
TraA dimers, and such mechanical stress will produce pronounced OM bulges that forces
OMs together, while also perturbing the bilayer structure, and thereby lowering the energy
barrier to the point where fusion occurs spontaneously (**Fig. 6B**). Our model further predicts
that after fusion, fission occurs when cells move apart from each other, where long narrow
OM tubes are sometimes seen between separating cells (Ducret *et al.*, 2013, Wei *et al.*, 2014).
In future studies, it will be interesting to test if exogenous expression of TraAB in other
bacteria with fluid OMs facilitates OME. However, there are technical issues reconstituting
the MYXO-CTERM sorting pathway for TraA cell surface localization in a heterologous host
(Sah *et al.*, 2020, Guo *et al.*, 2025). Nevertheless, if successful, this method can be used in
synthetic biology for generating bacterial multicellular structures.

Materials and methods
Bacterial strains, plasmids and growth conditions
Vegetative *M. xanthus* cells were grown in liquid CYE medium (10 mM MOPS pH 7.6, 1%
(w/v) Bacto™ casitone (BD Biosciences), 0.5% yeast extract and 8 mM $MgSO_4$) at 32 °C, in
125-ml flasks with vigorous shaking, or on CYE plates that contains 1.5% agar. Strains,

plasmids, and primers used in this study are listed in Supplementary **Table 1**. To create
P$_{pilA}$-SS$_{OM}$-PAmCherry (pPC60), fragments of P$_{pilA}$-SS$_{OM}$ and *PAmCherry* were PCR
amplified (Q5, New England Biolabs), and the resulting amplicons containing 25 bp
overhangs were ligated into the pSWU19 vector (linearized with EcoRI and XbaI) in
Gibson Assembly Master Mix (New England Biolabs). Construction of the SS$_{IM}$–
PAmCherry fusion was done following the published method (Wei *et al.*, 2011). This
construct was made by PCR amplification of genomic DNA with primers P$_{pilA}$-F2 and
SS$_{IM}$–mCherry-R (**Table 1**). The 5′ tail of the latter primer encodes 81 bp from MXAN_1176
that encode the N-terminal IM localization signal plus the downstream 10 amino acids
(CKDSDKKESM) for IM retention (Wei *et al.*, 2011). This PCR product was cloned to the
5′ end of PAmCherry at the XbaI site. Plasmids were verified by PCR, restriction enzyme
digestion, and if necessary, DNA sequencing. Verified plasmids were then electroporated
into *M. xanthus* cells and selected with appropriate antibiotics. The strains, plasmids and
primers used in this study are listed in **Table 1**.

Table 1. Strains, plasmids, and primers used in this study.

Strain name	Description	Source
DZ4469	DZ2 *pilA::tet,* Tc^R	(Vlamakis *et al.*, 2004)
DW1463	DK8601 (*aglQ1 pilA::tet*) *PpilA-traAB*, Tc^R	(Pathak *et al.*, 2012)
DW1483	DK8601 (*aglQ1 pilA::tet*) Δ*traAB*, Tc^R	(Cao & Wall, 2017)
DW2279	DW1483 pPC44, Km^R, Tc^R	(Cao & Wall, 2019a)
DW2285	DW1483 pPC49, Km^R, Tc^R	(Cao & Wall, 2019a)
DW2302	DW8601 pPC60 Km^R, Tc^R	This study
BN360	DW8601 pBN-IMPAC Km^R, Tc^R	This study

Plasmid name	Description	Source
pPC60	P$_{pilA}$-SS$_{OM}$-PAmCherry in pSWU19, Km^R	This study
pPC41	P$_{pilA}$-traA-gfp-traB in pSWU19, Km^R	(Cao & Wall, 2019a)
pPC44	P$_{pilA}$-traA-mCherry-traB in pSWU19, Km^R	(Cao & Wall, 2019a)
pPC49	P$_{pilA}$-traA-traB-gfp in pSWU19, Km^R	(Cao & Wall, 2019a)
pBN-IMPAC	P$_{pilA}$-SS$_{IM}$-PAmCherry in pSWU19, Km^R	This study

Primer name	Sequence (from 5' to 3')	Use
pSWU19-EcoRI-P$_{pilA}$-F	AGGAAACAGCTATGACCATGATTACGAATTCCGTCATGTTGGACGAGGT	pPC60
P$_{pilA}$-SS$_{OM}$-PAmCherry-R	GCCCTTGCTCACTCTAGAAGGAAACGAGCAACCG	pPC60
P$_{pilA}$-SS$_{OM}$-PAmCherry-F	GTTTCCTTCTAGAGTGAGCAAGGGCGAGGAG	pPC60
PAmCherry-XbaI-R	TGCATGCCTGCAGGTCGACTCTAGAAAGCTTTTACTTGTACAGCTCGTCCATGC	pPC60

P_{pilA}-F2	GGATCTAGATGCATGCGGCCGCGTCATGTTGGACGAGGTC	pBN-IMPAC
SS_{IM}-mCherry-R	CACATCTAGACATGGATTCCTTCTTGTCGGAGTCCTTGCAGG CGGTGAGGGTAAGGCAGAGGAGGACAGTCGTCAGGAATTG GGTCCGCATGGTGACTCTCCTCCGGGGGTCCTCAGAGAAG GTTG	pBN-IMPAC

Light microscopy and image analysis

For all imaging experiments, 5 ml cells or cell mixtures were spotted on a 1.5% (w/v) agar pad containing 10 mM MOPS pH 7.6 and 8 mM MgSO₄. All microscopy images were captured using an Andor iXon Ultra 897 EMCCD camera (effective pixel size 160 nm) on an inverted Nikon Eclipse-Ti™ microscope with a 100× 1.49 NA TIRF objective. GFP and mCherry were photobleached using the 488-nm and 561-nm lasers (0.5 kW/cm^2), respectively at the stated frame rates. sptPALM was performed as previously described (Fu *et al.*, 2018). *M. xanthus* cells were grown in CYE to OD$_{600}$ around 1. PAmCherry was activated using a 405-nm laser (0.3 kW/cm^2), excited and imaged using a 561-nm laser (0.2 kW/cm^2). Images were acquired at 67 Hz (15 ms/frame). For each sptPALM experiment, single PAmCherry particles were localized in at least 100 individual cells from three biological replicates. sptPALM and cell morphology data were analyzed as previously described using a MATLAB (MathWorks) script (Fu *et al.*, 2018, Zhang *et al.*, 2023a, Zhang *et al.*, 2023b). Briefly, cells were identified using differential interference contrast images. Single PAmCherry particles inside of cells were fitted by a symmetric 2D Gaussian function; its center was assumed to be the particle's position (Fu *et al.*, 2018). Particles in consecutive frames were considered to belong to the same trajectory when they were within a user-defined distance of 320 nm (two pixels). Sample trajectories were generated using the TrackMate plugin (Ershov *et al.*, 2022) in the ImageJ suite (https://imagej.net). For simplicity, we considered all the mobile particles of these proteins displaying typical diffusion and determined their *D* from a linear fit to the first four points of the MSD using a simpler formula MSD = 4*DΔt* (Fu *et al.*, 2018, Lee *et al.*, 2016). Error bars were the standard derivation of 1,000 bootstrap samples using the published method (Morgenstein *et al.*, 2015).

Cryo-EM

Bacterial strains were grown overnight in CYE media with appropriate antibiotics,

incubated with shaking at 32 °C and 250 rpm to an optical density of OD_{600} 0.6. Cells were

collected by centrifugation at 5,000 rpm and room temperature for 3 min and

resuspended in CYE media to a final OD_{600} of 12. The cell suspension (3 μl) was applied

to C Flat-1.2/1.3 200 mesh copper grids (Electron Microscopy Sciences) that were glow

discharged for 30 s at 15 mA. Grids were plunge-frozen in liquid ethane with an FEI

Vitrobot Mark IV (Thermo Fisher Scientific) at 4 °C, 100% humidity with a waiting time

of 30 s, two-side blotting time of 2.5 - 3 s, and blotting force of 0. All subsequent grid

handling and transfers were performed in liquid nitrogen. Grids were clipped onto cryo-

FIB autogrids (Thermo Fisher Scientific). We used a Titan Krios G4 transmission electron

microscope (Thermo Fisher Scientific), equipped with a Gatan K3 direct electron detector

and a Gatan BioContinuum energy filter. Imaging was performed using a 15 eV slit width

on the energy filter to enhance image contrast. Micrographs were recorded in counted

mode at a nominal magnification of 30,000 x, using a dose rate of 14.65

electrons/pixel/second over a 2.6-second exposure, resulting in a total accumulated dose

of 50 electrons/$Å^2$. Low magnification images were taken at 6500 x with a pixel size of

1.39 nm.

Acknowledgments

This work was supported by the National Institutes of Health grants GM129000 to B. N. and GM140886 to D.W.

Movie captions

Movie S1. TraA-GFP foci being photobleached. Time-lapse images were recorded at 67 Hz and played at the same frame rate (real time).

Movie S2. Simultaneous photobleaching of TraA-mCherry (right) and TraB-GFP (left) in foci between adjacent cells. Time-lapse images were recorded at 5 Hz and played at the same frame rate (real time).

Movie S3. Diffusion of OM_{SS}-PAmCherry particles. Time-lapse images were recorded at 67 Hz and played at the same frame rate (real time).

Movie S4. Diffusion of IM_{SS}-PAmCherry particles. Time-lapse images were recorded at 67 Hz and played at the same frame rate (real time).

Movie S5. OME is bidirectional. The donor cell expresses OM_{SS}-PAmCherry (magenta) and the recipient TraA-GFP (green). The particle was first transferred to the recipient and then returned to the donor. Time-lapse images were recorded at 67 Hz and played at 7 frames/s, about 1/10 speed of real time.

Movie S6. Serial cell-cell transfer of OM_{SS}-PAmCherry particle by OME. Donor cell expresses OM_{SS}-PAmCherry (magenta) and the recipients TraA-GFP (green). Particle was serially transferred, crossing three cell-cell boundaries. Time-lapse images were recorded at 67 Hz and played at 7 frames/s, about 1/10 speed of real time.

Movie S6. TraAB foci do not function as OME conduits. Donor cell expresses OM_{SS}-PAmCherry (magenta) and the recipients TraA-GFP (green). Two OM_{SS}-PAmCherry particles transferred across cell boundaries, but not through the TraAB foci. Time-lapse images were recorded at 67 Hz and played at 7 frames/s, about 1/10 speed of real time.

References

Balagam, R., Cao, P., Sah, G.P., Zhang, Z., Subedi, K., Wall, D., and Igoshin, O.A. (2021) Emergent myxobacterial behaviors arise from reversal suppression induced by kin contacts. *mSystems* **6**: e0072021.

Batsanov, S.S. (2001) Van der Waals Radii of Elements. *Inorganic Materials* **37**: 871-885.

Benn, G., Mikheyeva, I.V., Inns, P.G., Forster, J.C., Ojkic, N., Bortolini, C., Ryadnov, M.G., Kleanthous, C., Silhavy, T.J., and Hoogenboom, B.W. (2021) Phase separation in the outer membrane of *Escherichia coli*. *Proc Natl Acad Sci U S A* **118**.

Cao, P., Dey, A., Vassallo, C.N., and Wall, D. (2015) How myxobacteria cooperate. *J Mol Biol* **427**: 3709-3721.

Cao, P., and Wall, D. (2017) Self-identity reprogrammed by a single residue switch in a cell surface receptor of a social bacterium. *Proc Natl Acad Sci U S A* **114**: 3732-3737.

Cao, P., and Wall, D. (2019a) Direct visualization of a molecular handshake that governs kin recognition and tissue formation in myxobacteria. *Nature communications* **10**: 3073.

Cao, P., and Wall, D. (2020) The Fluidity of the Bacterial Outer Membrane Is Species Specific: Bacterial Lifestyles and the Emergence of a Fluid Outer Membrane. *Bioessays* **42**: e1900246.

Cao, P., Wei, X., Awal, R.P., Muller, R., and Wall, D. (2019) A highly polymorphic receptor governs many distinct self-recognition types within the *Myxococcales* order. *mBio* **10**: 10.1128/mbio.02751-02718.

Cao, P.B., and Wall, D. (2019b) Direct visualization of a molecular handshake that governs kin recognition and tissue formation in myxobacteria. *Nat Commun* **10**.

Cherrak, Y., Flaugnatti, N., Durand, E., Journet, L., and Cascales, E. (2019) Structure and Activity of the Type VI Secretion System. *Microbiol Spectr* **7**.

Dai, S., Sun, C., Tan, K., Ye, S., and Zhang, R. (2017) Structure of thrombospondin type 3 repeats in bacterial outer membrane protein A reveals its intra-repeat disulfide bond-dependent calcium-binding capability. *Cell Calcium* **66**: 78-89.

Dey, A., Vassallo, C.N., Conklin, A.C., Pathak, D.T., Troselj, V., and Wall, D. (2016) Sibling Rivalry in *Myxococcus xanthus* Is Mediated by Kin Recognition and a Polyploid Prophage. *Journal of Bacteriology* **198**: 994-1004.

Dey, A., and Wall, D. (2014) A genetic screen in *Myxococcus xanthus* identifies mutants that uncouple outer membrane exchange from a downstream cellular response. *J Bacteriol* **196**: 4324-4332.

Ducret, A., Fleuchot, B., Bergam, P., and Mignot, T. (2013) Direct live imaging of cell-cell protein transfer by transient outer membrane fusion in *Myxococcus xanthus*. *Elife* **2**: e00868.

Ershov, D., Phan, M.S., Pylvanainen, J.W., Rigaud, S.U., Le Blanc, L., Charles-Orszag, A., Conway, J.R.W., Laine, R.F., Roy, N.H., Bonazzi, D., Dumenil, G., Jacquemet, G., and Tinevez, J.Y. (2022) TrackMate 7: integrating state-of-the-art segmentation algorithms into tracking pipelines. *Nat Methods* **19**: 829-832.

Fu, G., Bandaria, J.N., Le Gall, A.V., Fan, X., Yildiz, A., Mignot, T., Zusman, D.R., and Nan, B. (2018) MotAB-like machinery drives the movement of MreB filaments during bacterial gliding motility. *Proc Natl Acad Sci U S A* **115**: 2484-2489.

Gong, Y., Zhang, Z., Zhou, X.W., Anwar, M.N., Hu, X.Z., Li, Z.S., Chen, X.J., and Li, Y.Z. (2018) Competitive Interactions Between Incompatible Mutants of the Social Bacterium Myxococcus xanthus DK1622. *Front Microbiol* **9**: 1200.

Grosberg, R.K., and Strathmann, R.R. (2007) The evolution of multicellularity: A minor major

transition? *Annu Rev Ecol Evol S* **38**: 621-654.

Guo, T., Haft, D.H., and Wall, D. (2025) Myxosortase: an intramembrane protease that sorts
MYXO-CTERM proteins to the cell surface. *mBio*: e0406724.

Hodgkin, J., and Kaiser, D. (1977) Cell-to-cell stimulation of motility in nonmotile mutants of
Myxococcus. *Proc. Natl. Acad. Sci. USA* **74**: 2938-2942.

Jahn, R., Cafiso, D.C., and Tamm, L.K. (2024) Mechanisms of SNARE proteins in membrane
fusion. *Nat Rev Mol Cell Biol* **25**: 101-118.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Zidek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J.,
Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S.,
Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T.,
Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W., Kavukcuoglu, K., Kohli, P., and
Hassabis, D. (2021) Highly accurate protein structure prediction with AlphaFold. *Nature*
596: 583-589.

Kroos, L., Wall, D., Islam, S.T., Whitworth, D.E., Munoz-Dorado, J., Higgs, P.I., Singer, M.,
Mauriello, E.M., Treuner-Lange, A., Sogaard-Andersen, L., Kaimer, C., Elias-Arnanz, M.,
Stojkovic, E.A., Muller, R., Volz, C., Velicer, G.J., and Nan, B. (2025) Milestones in the
development of *Myxococcus xanthus* as a model multicellular bacterium. *J Bacteriol* **207**:
e0007125.

Lee, T.K., Meng, K., Shi, H., and Huang, K.C. (2016) Single-molecule imaging reveals modulation
of cell wall synthesis dynamics in live bacterial cells. *Nature communications* **7**: 13170.

Lorenzen, W., Bozhuyuk, K.A., Cortina, N.S., and Bode, H.B. (2014) A comprehensive insight
into the lipid composition of *Myxococcus xanthus* by UPLC-ESI-MS. *J Lipid Res* **55**: 2620-
2633.

Lucena, D., Mauri, M., Schmidt, F., Eckhardt, B., and Graumann, P.L. (2018) Microdomain
formation is a general property of bacterial membrane proteins and induces heterogeneity
of diffusion patterns. *BMC Biol* **16**: 97.

Morgenstein, R.M., Bratton, B.P., Nguyen, J.P., Ouzounov, N., Shaevitz, J.W., and Gitai, Z. (2015)
RodZ links MreB to cell wall synthesis to mediate MreB rotation and robust morphogenesis.
Proc Natl Acad Sci U S A.

Nan, B., Bandaria, J.N., Guo, K.Y., Fan, X., Moghtaderi, A., Yildiz, A., and Zusman, D.R. (2015)
The polarity of myxobacterial gliding is regulated by direct interactions between the
gliding motors and the Ras homolog MglA. *Proc Natl Acad Sci U S A* **112**: E186-193.

Nan, B., Bandaria, J.N., Moghtaderi, A., Sun, I.H., Yildiz, A., and Zusman, D.R. (2013) Flagella
stator homologs function as motors for myxobacterial gliding motility by moving in helical
trajectories. *Proc Natl Acad Sci U S A* **110**: E1508-1513.

Nudleman, E., Wall, D., and Kaiser, D. (2005) Cell-to-cell transfer of bacterial outer membrane
lipoproteins. *Science* **309**: 125-127.

Parry, B.R., Surovtsev, I.V., Cabeen, M.T., O'Hern, C.S., Dufresne, E.R., and Jacobs-Wagner, C.
(2014) The bacterial cytoplasm has glass-like properties and is fluidized by metabolic
activity. *Cell* **156**: 183-194.

Pathak, D.T., Wei, X., Bucuvalas, A., Haft, D.H., Gerloff, D.L., and Wall, D. (2012) Cell contact-
dependent outer membrane exchange in myxobacteria: genetic determinants and
mechanism. *PLoS Genet* **8**: e1002626.

Pathak, D.T., Wei, X., Dey, A., and Wall, D. (2013) Molecular recognition by a polymorphic cell
surface receptor governs cooperative behaviors in bacteria. *PLoS Genet.* **9**: e1003891.

Perez-Burgos, M., Garcia-Romero, I., Jung, J., Valvano, M.A., and Sogaard-Andersen, L. (2019) Identification of the lipopolysaccharide O-antigen biosynthesis priming enzyme and the O-antigen ligase in *Myxococcus xanthus*: critical role of LPS O-antigen in motility and development. *Mol Microbiol* **112**: 1178-1198.

Pogue, C.B., Zhou, T., and Nan, B. (2018) PlpA, a PilZ-like protein, regulates directed motility of the bacterium *Myxococcus xanthus*. *Mol Microbiol* **107**: 214-228.

Ramirez Carbo, C.A., Faromiki, O.G., and Nan, B. (2024) A lytic transglycosylase connects bacterial focal adhesion complexes to the peptidoglycan cell wall. *Elife* **13**.

Rand, R.P., and Parsegian, V.A. (1984) Physical force considerations in model and biological membranes. *Can J Biochem Cell Biol* **62**: 752-759.

Rassam, P., Copeland, N.A., Birkholz, O., Toth, C., Chavent, M., Duncan, A.L., Cross, S.J., Housden, N.G., Kaminska, R., Seger, U., Quinn, D.M., Garrod, T.J., Sansom, M.S., Piehler, J., Baumann, C.G., and Kleanthous, C. (2015) Supramolecular assemblies underpin turnover of outer membrane proteins in bacteria. *Nature* **523**: 333-336.

Rojas, E.R., Billings, G., Odermatt, P.D., Auer, G.K., Zhu, L., Miguel, A., Chang, F., Weibel, D.B., Theriot, J.A., and Huang, K.C. (2018) The outer membrane is an essential load-bearing element in Gram-negative bacteria. *Nature* **559**: 617-621.

Sah, G.P., Cao, P., and Wall, D. (2020) MYXO-CTERM sorting tag directs proteins to the cell surface via the type II secretion system. *Mol Microbiol* **113**: 1038-1051.

Sah, G.P., and Wall, D. (2020) Kin recognition and outer membrane exchange (OME) in myxobacteria. *Curr Opin Microbiol* **56**: 81-88.

Shi, L., Shen, Q.T., Kiel, A., Wang, J., Wang, H.W., Melia, T.J., Rothman, J.E., and Pincet, F. (2012) SNARE proteins: one to fuse and three to keep the nascent fusion pore open. *Science* **335**: 1355-1359.

Sinha, R., Ahmed, S., Jahn, R., and Klingauf, J. (2011) Two synaptobrevin molecules are sufficient for vesicle fusion in central nervous system synapses. *Proc Natl Acad Sci U S A* **108**: 14318-14323.

Subedi, K., Roy, P.C., Saiz, B., Basile, F., and Wall, D. (2024) Cell-cell transfer of adaptation traits benefits kin and actor in a cooperative microbe. *Proc Natl Acad Sci U S A* **121**: e2402559121.

Sun, J., Rutherford, S.T., Silhavy, T.J., and Huang, K.C. (2022) Physical properties of the bacterial outer membrane. *Nat Rev Microbiol* **20**: 236-248.

Tran, B.M., Punter, C.M., Linnik, D., Iyer, A., and Poolman, B. (2024) Single-protein Diffusion in the Periplasm of *Escherichia coli*. *J Mol Biol* **436**: 168420.

van den Bogaart, G., Holt, M.G., Bunt, G., Riedel, D., Wouters, F.S., and Jahn, R. (2010) One SNARE complex is sufficient for membrane fusion. *Nat Struct Mol Biol* **17**: 358-364.

Vassallo, C., Pathak, D.T., Cao, P., Zuckerman, D.M., Hoiczyk, E., and Wall, D. (2015) Cell rejuvenation and social behaviors promoted by LPS exchange in myxobacteria. *Proc Natl Acad Sci U S A* **112**: E2939-2946.

Vassallo, C.N., Cao, P., Conklin, A., Finkelstein, H., Hayes, C.S., and Wall, D. (2017) Infectious polymorphic toxins delivered by outer membrane exchange discriminate kin in myxobacteria. *Elife* **6**.

Vassallo, C.N., Sah, G.P., Weltzer, M.L., and Wall, D. (2021) Modular Lipoprotein Toxins Transferred by Outer Membrane Exchange Target Discrete Cell Entry Pathways. *mBio* **12**: e0238821.

Vassallo, C.N., Troselj, V., Weltzer, M.L., and Wall, D. (2020) Rapid diversification of wild social

groups driven by toxin-immunity loci on mobile genetic elements. *ISME J* **14**: 2474-2487.

Vassallo, C.N., and Wall, D. (2016) Tissue repair in myxobacteria: A cooperative strategy to heal
cellular damage. *BioEssays* **38**: 306-315.

Vassallo, C.N., and Wall, D. (2019) Self-identity barcodes encoded by six expansive polymorphic
toxin families discriminate kin in myxobacteria. *Proc. Natl. Acad. Sci. USA* **116**: 24808-
24818.

Velicer, G.J., Kroos, L., and Lenski, R.E. (2000) Developmental cheating in the social bacterium
Myxococcus xanthus. *Nature* **404**: 598-601.

Vlamakis, H.C., Kirby, J.R., and Zusman, D.R. (2004) The Che4 pathway of *Myxococcus xanthus*
regulates type IV pilus-mediated motility. *Mol Microbiol* **52**: 1799-1811.

Wall, D. (2014) Molecular recognition in myxobacterial outer membrane exchange: functional,
social and evolutionary implications. *Mol Microbiol* **91**: 209-220.

Wall, D., and Kaiser, D. (1998) Alignment enhances the cell-to-cell transfer of pilus phenotype.
Proc Natl Acad Sci U S A **95**: 3054-3058.

Wall, D., Wu, S.S., and Kaiser, D. (1998) Contact stimulation of Tgl and type IV pili in
Myxococcus xanthus. *J Bacteriol* **180**: 759-761.

Wei, X., Pathak, D.T., and Wall, D. (2011) Heterologous protein transfer within structured
myxobacteria biofilms. *Mol Microbiol* **81**: 315-326.

Wei, X., Vassallo, C.N., Pathak, D.T., and Wall, D. (2014) Myxobacteria produce outer membrane-
enclosed tubes in unstructured environments. *J Bacteriol* **196**: 1807-1814.

Weltzer, M.L., Guo, T., and Wall, D. (2025) Crosstalk between polymorphic toxin-immunity
systems involved in kin discrimination. *mBio* **0**: e00468-00425.

Yang, Z., Lux, R., Hu, W., Hu, C., and Shi, W. (2010) PilA localization affects extracellular
polysaccharide production and fruiting body formation in *Myxococcus xanthus*. *Mol*
Microbiol **76**: 1500-1513.

Zhang, H., Mulholland, G.A., Seef, S., Zhu, S., Liu, J., Mignot, T., and Nan, B. (2020) Establishing
rod shape from spherical, peptidoglycan-deficient bacterial spores. *Proc Natl Acad Sci U*
S A **117**: 14444-14452.

Zhang, H., Venkatesan, S., Ng, E., and Nan, B. (2023a) Coordinated peptidoglycan synthases and
hydrolases stabilize the bacterial cell wall. *GitHub*.

Zhang, H., Venkatesan, S., Ng, E., and Nan, B. (2023b) Coordinated peptidoglycan synthases and
hydrolases stabilize the bacterial cell wall. *Nature communications* **14**: 5357.

Zhou, T., and Nan, B. (2017) Exopolysaccharides promote *Myxococcus xanthus* social motility by
inhibiting cellular reversals. *Mol Microbiol* **103**: 729-743.