# Immunogenicity and Structural Features of a Stabilized Clade-C HIV-1 Env Derived from a Pediatric Elite-neutralizer

Swarandeep Singh[1,†], Sanjeev Kumar[1,2,†], Arnab Chatterjee[3,†], Lakshay Malhotra[4], Abdul Rahman[1], Ifrah Andrabi[1], Bapan Mondal[3], Sanket Katpara[1], Shaifali Sharma[1], Harsh Bhakri[1], Anjli Gaur[1] ,Tanu Bansal[1], Muneeb Pervez[1], Vikram Iyer[1] , Abdul Wahid Hussain[1], Shubbir Ahmed[6], Rajesh Kumar[5], Ethayathulla Abdul Samath[4], Rakesh Lodha[7], Somnath Dutta[3*], Kalpana Luthra[1*]

[1]Department of Biochemistry, All India Institute of Medical Sciences, New Delhi, India

[2]ICGEB-Emory Vaccine Center, International Center for Genetic Engineering and Biotechnology, New Delhi, India

[3]Molecular Biophysics Unit, Indian Institute of Science, Bangalore, India

[4]Department of Biophysics, All India Institute of Medical Sciences, New Delhi, India

[5]Indian Institute of Technology, Roorkee, India

[6] Centralized Core Research Facility, All India Institute of Medical Sciences, New Delhi, India

[7] Department of Pediatrics, All India Institute of Medical Sciences, New Delhi, India

[†]These authors contributed equally

[*]Correspondence**:** somnath@iisc.ac.in (S.D.), kalpanaluthra@gmail.com (K.L.)

**Keywords:** HIV-1 vaccine, Immunogenicity**,** Neutralizing Antibodies, Cryo-EM structure

## ABSTRACT

The envelope (Env) glycoprotein from viruses circulating in elite-neutralizers-individuals who naturally develop broadly neutralizing antibody (bNAb) responses, serves as potential template for rational HIV-1 vaccine design. Here, we engineered a soluble clade-C Env trimer (330) derived from a pediatric elite-neutralizer (AIIMS_330) using SOSIP, NFL, and self-assembling ferritin-nanoparticles (NP) platforms. Both Env SOSIP and NP-displayed Envs exhibited high thermostability and nanomolar binding affinity to diverse HIV-1 bNAbs, with minimal reactivity to non-neutralizing antibodies, indicative of a native-like antigenic profile. A cryo-EM structure of the 330-SOSIP trimer in complex with the autologous bNAb 44m, revealed recognition of the GDIR motif and N332 supersite with additional contacts at T144, and T413. Rabbits immunized with the SOSIP and NP elicited potent autologous and other tier-1 clade-C viruses neutralizing

responses, supporting the utility of multivalent vaccine. These findings broadens our understanding of the structure and immunogenicity of the HIV-1 Env from pediatric ENs and contributes valuable information towards a rational immunogen design.

**INTRODUCTION**

A critical goal in HIV vaccine development is to elicit broadly neutralizing antibodies (bNAbs) in uninfected and high-risk individuals[1]. Elite neutralizers (ENs), a rare group of HIV-1-infected individuals who develop exceptionally potent bNAb responses, provide valuable blueprints for rational HIV-1 immunogen design, aimed at inducing similar bNAb responses upon vaccination[1–3]. Studies involving simian-human immunodeficiency virus (SHIV) challenges in nonhuman primates (NHPs) have indicated that passive immunization with bNAbs can effectively prevent infection, supporting the development of an HIV-1 vaccine that induces such protective immune responses in humans upon vaccination[4–8]. Stabilized HIV-1 envelope (Env) glycoproteins have emerged as promising candidates for immunogen design, with the potential to elicit protective immune responses through vaccination, based on insights from the RV144 Thai trial[9,10]. Native like HIV-1 Env trimers, particularly the clade-A BG505 Env have shown the ability to induce neutralizing antibody (nAb) responses in animal models and are currently under evaluation in clinical trials[11–18]. HIV-1 clade C, predominates global HIV-1 infection with more than 50% infections[19,20]. A limited number of clade-C Envs has been stabilized; with low expression levels and instability, posing hurdles to develop clade C Env-based vaccines[21–24].

Recent investigations highlight distinct differences in surface antigen presentation between HIV-1 and HPV, contributing to HIV-1's limited ability to elicit a strong immune response compared to HPV[25]. In contrast, currently approved multivalent vaccines for HPV and HBV generate robust immune responses with >95% efficacy[26–28]. Recent studies suggest that single Env-based vaccine regimens may not be the most effective approach for HIV-1 prevention. Notably, presenting HIV-1 Env trimers on self-assembling ferritin nanoparticles has shown promise in eliciting robust immune response in rabbits, suggesting improved immunogenicity[29,30]. Studies also suggest that HIV-infected children who often acquire the infection through vertical transmission, are inherently better at generating bNAbs than adults[31–36]. The relative immaturity of the immune system in children may allow for more effective

protective immune responses, as founder virus encounters minimal selection pressure,
providing the  evolving B-cells an advantage in generating NAbs that effectively neutralize
contemporary viruses[31–33,37]. Env trimers from contemporaneous viruses can serve as effective
tools for immunogen design, and  structural characterization of Env in complex with autologous
bNAbs from the same donor can provide valuable insights for vaccine development[36].

We previously reported on a pair of HIV-1-infected monozygotic twins (AIIMS_330 and
AIIMS_329) demonstrating elite neutralizing activity, with longitudinal sampling of AIIMS_330
revealing sustained elite neutralizing activity. The Env from this EN exhibited multiple broad
bNAb epitope specificities, making it an ideal template for designing and characterizing HIV-1
Env mimics[33,35]. In this study, we designed and characterized an HIV-1 Clade-C 330 Env trimer
derived from a pediatric EN AIIMS_330, as soluble SOSIP & NFL Env trimers, alongside tailoring
the Env to display a repetitive array of HIV-1 trimers on self-assembling ferritin nanoparticles
(NPs)[11,29,38]. Designed Env constructs, stabilized by trimer-stabilizing mutations, exhibited
enhanced expression and a greater propensity to assemble into native-like HIV-1 Env
mimics[12,15,39,40]. The 330 Env constructs exposed multiple epitopes and demonstrated strong
affinity to bNAbs while occluding the immunodominant epitopes of non-neutralizing antibodies
(non-NAbs). Ferritin NPs exhibited strong immunogenicity and tier-1 autologous neutralizing
activity, upon immunization in rabbits compared to soluble counterparts. Negative stain-
electron microscopy (NS-EM) showed that both soluble Envs and multivalent NP adopted native-
like morphology. A cryo-EM structure, revealed a stable, well-organized Envelope structure in
the prefusion closed conformation of the 330 SOSIP Env trimer. Structural insights from the
cryo-EM model of the 330 SOSIP Env trimer in complex with the autologous bNAb 44m, a V3-
glycan-specific potent bNAb identified from the AIIMS_330 EN[36], provide information on the
antigenic determinants. Collectively, this study broadens our understanding of the structure and
immunogenicity of the HIV-1 Env from pediatric ENs. The 330 Env could serve as valuable
antigenic probes for HIV-1 bNAb discovery and potential Env for a multivalent vaccine design
and vaccination strategies.

## RESULTS

### Design & biophysical characterization of 330 Env trimers and ferritin nanoparticles

Previously we reported a pair of HIV-1 chronically infected monozygotic twins AIIMS_330 & AIIMS_329 exhibiting elite plasma neutralizing activity[35]. Over five years of longitudinal sampling, the AIIMS_330 EN showed sustained evolution of plasma bNAbs targeting major neutralizing determinants on the HIV-1 Env glycoprotein[33,35,41]. The pseudoviral Env clone 330.16.E6, derived from the AIIMS_330 EN, exhibited high susceptibility to neutralization by a broad range of HIV-1 bNAbs with multiple epitope specificity, along with autologous contemporaneous plasma antibodies, and resistance to the panel of non-nAbs tested[33,35]. Therefore, we engineered this Env in membrane soluble forms, as SOSIP.664 gp120-gp41 cleaved Env trimers, NFL.664 gp140 single-chain uncleaved Env trimers and SOSIP Env trimers on self-assembling ferritin NPs to achieve multivalency (Figure 1 A-C). The 330 Env trimers were stabilized by incorporating mutations for increased expression and trimerization, as illustrated in Figure 1A-D, Table S1. The SOSIP Env was designed by introducing a SOS disulfide bond (A501C-T605C)[46], an I559P mutation[42], and an optimized furin cleavage site (RRRRRR)[47] . We designed the NFL Env construct, that carried the  I559P mutation, by linking the gp120 and gp41 using a flexible glycine-serine linker (GGGGSGGGGS)[38]. Both Env constructs were human codon-optimized to enhance expression in mammalian systems.

The stabilized and optimized 330 Envs were expressed in HEK293F cells, followed by a two-step purification viz. HIV-1 Env trimer specific bNAb PGT145-Immunoaffinity chromatography followed by Size Exclusion Chromatography (SEC). The SEC purification of 330 Env trimers (SOSIP and NFL) yielded a peak at an elution volume of 10 mL, which shifted further upward for ferritin nanoparticles, eluting at 8.5 mL on a Superdex 200 10/300 column (Figure 1E). The purification yield of 330 Env trimers following SEC was around ~1 mg/L for SOSIP and ~1.45 mg/L for NFL (Figure S2), which was higher or comparable to the Indian and South African Clade-C strains (CZA97: ~0.4-0.6 mg/mL, 329 SOSIP: ~0.6 mg/mL, 16055 NFL: ~2 mg/mL)[23,48–50]. The NFL trimer of Env 330 had better expression yields than the SOSIP version, likely due to the increased stability and functionality provided by the covalent linkage between the gp120 and gp41 subunits, resulting in more functional gp140 molecules[38,50]. However, low yield of Ferritin NPs was observed, around 0.05 mg/L, likely due to the complex assembly of Env trimers into synthesized ferritin nanocages as reported previously[51,52]. The 330 Env trimers

displayed single band on Blue Native-PAGE (BN-PAGE) stained with Coomassie, whereas the 330

ferritin NPs were too large to efficiently enter the gels and visible at the top of the lanes,

indicating the high purity of Env trimers and NPs preparations (Figure S2-Left). The SDS-PAGE

analysis revealed intact trimeric nature of the 330 Env proteins under non-reducing conditions

that separated as monomers in the presence of DTT under reducing conditions (Figure S2-

Right). Negative-Stain Electron Microscopy (NS-EM) reference-free two-dimensional (2D) class

averages demonstrated that over 99% of the gp140 proteins adopted a native-like trimeric

morphology, with multiple copies of the Env spikes efficiently displayed on the self-assembling

ferritin NPs (Figure 1F, Supplementary S3).  The 3D model of soluble gp140 trimers reveals a

trilobular structure, while ferritin NPs display multiple copies of Env spikes (Figure 1G).

Thermal stability & secondary structures of 330 Envs were evaluated using sypro-orange

Differential Scanning Fluorimetry (DSF) and by Circular Dichroism (CD) spectroscopy. The 330

SOSIP Env exhibited a Tm of ~74°C indicating its ability to retain its structural integrity even at

elevated temperatures (Figure 1H, S4 A-B). The CD spectra of 330 SOSIP Env reveal ~25% alpha-

helical and ~25% antiparallel beta-sheet content, contributing to the stability and rigidity of the

Env, while random coils impart the flexibility required for molecular recognition, collectively

suggesting a thermodynamically stable secondary structure for the Envelope protein (Figure S4

C-D). Overall, biophysical characterization demonstrated that the 330 Envs are well-stabilized,

and effectively displayed on ferritin NP cages.

**Antigenicity of 330 Env trimers and ferritin nanoparticles indicated strong binding affinity to HIV-1 bNAbs**

Next, we assessed the antigenicity of the 330 Envs using binding ELISA with an extensive

panel of HIV-1 bNAbs and non-nAbs. The 330 Env trimers exhibited well-exposed epitopes

recognized by potent bNAbs, including the V1V2 apex (PGT145, PGDM1400, CAP256.25), the V3-

glycan supersite (2G12, PGT121, 10-1074, AIIMS-P01), the CD4 binding site (VRC01, N6), and

the gp120-gp41 interface (PGT151), while showing minimal reactivity to non-NAbs (F105, A32,

48d, 17b+sCD4), suggesting that the 330 Env trimers adopt native-like conformations (Figure 2).

The antigenic profile of ferritin NPs was found to resemble that of SOSIP, though with less

saturation to tested monoclonal antibodies (mAbs) concentration and allowing all the

conformational epitopes to be accessible to HIV-1 neutralizing mAbs targeting to the Envs

presented on the nanoparticle. A minor degree of binding of the non-NAbs was observed with

NFL and ferritin particles, with negligible to SOSIP. Notably, the 330 Env trimers and particles
exhibited preferential affinity for neutralizing mAbs over non-neutralizing mAbs.

Subsequently, we assessed the binding affinity of 330 Envs to neutralizing & non-
neutralizing mAbs with distinct specificities was then assessed by BLI-Octet analysis. To assess
binding affinity and kinetics, we selected conformation-specific bNAbs (PGT145, PGDM1400, and
PGT151) as well as non-neutralizing antibodies (F105, 17b, and sCD4). The gp140 Envs (affinity)
& ferritin NPs (avidity) exhibited strong binding affinity to the tested bNAbs (Figure 3). The
SOSIP variants (Ferritin NPs) (KD 1.07 nm) & NFL (KD 1.39 nm) showed a high affinity of ~1nM
to the V1V2 directed bNAb PGT145. A similar trend in binding of PGDM1400 with SOSIP (KD
1.34 nm) & NFL Env (KD 3.32 nm) was observed respectively. The interface-directed bNAb
PGT151 demonstrated an exceptionally high affinity to SOSIP variants (Ferritin NPs), with an
affinity of ~5 pM, indicating that the PGT151 epitopes on the Env proteins displayed on the
nanoparticle were highly accessible (Table S2). We observed that PGT145-purified SOSIP
trimers were efficiently cleaved at the gp120-gp41 junction, as indicated by strong binding to the
cleavage-specific bNAb PGT151, while no binding was detected with NFL trimer (Figure 2 & 3).
The  introduction of intraprotomer locking 201C –433C disulfide bond as described previously[50],
in these constructs prevented the CD4-induced Env rearrangements as assessed by low
reactivity with sCD4 & 17b non-nAbs, indicating the prefusion conformation of the Envelopes. As
expected, and consistent with the ELISA binding results, the 330 Envs exhibited minimal
reactivity to the non-NAbs F105 and 17b+sCD4 (Figure 2 & 3, Table S2). Collectively, these
findings suggest that both the soluble Envs and multivalent NPs adopt a native-like
conformation, exposing multiple key bNAb epitopes while shielding the immunodominant
epitopes targeted by non-NAbs. Based on the negligible antigenic reactivity of the SOSIP Env
with non-nAbs, we next performed the structural analysis of the SOSIP Env.

**Cryo-EM structure of 330 SOSIP Env in complex with autologous bNAb 44m**

The structure of the natively glycosylated 330 SOSIP Env was solved using single-particle
cryo-electron microscopy (cryo-EM). A total 109004 number of particles were analyzed to
determine the 2D class averages and reconstruct the trilobular 3D model of 330 SOSIP Env in its
prefusion closed conformations (Figure S5 A-C). In a recent study, we reported heavy chain
matured HIV-1 lineage bNAb 44m, which effectively neutralizes its autologous pseudovirus,
330.16.E6, from which the 330 Env trimer is derived[36]. We aimed to investigate the key

interacting residues of bNAb 44m to its autologous Envelope 330. A cryo-EM structure of 330

SOSIP Env in complex with 44m bNAb was solved (Figure 5, S6). The architecture of 330 Env

displayed the trilobular shapes of trimers with three extra densities of corresponding bNAb 44m

Fab bound moieties (Figure 5 A-B, Supply S6A). The model showed that three 44m FAbs bind to

each gp120 monomeric subunit, enabling effective neutralization (Figure 5C-F). The interacting

surface area covered by the mainly by heavy chain of 44m and 330 Envs are 522 Å², and 554 Å²

respectively (Figure 5B). The model further reveals that the 44m CDRH1 is positioned near the

330 Env V1V2 loop, while CDRH2 and CDRH3 are located near the V3 region (Figure 5C). Herein ,

we observed  that 44m bNAb  showed dependence on V3-glycan N332-supersite and GDIR motif

along with distinct key residues identified in case of autologous Env 330 as compared to the

heterologous clade-A Env BG505 as reported previously[36]. The key interacting residues include

H31 in the CDRH1 loop, which forms a contact site with T144 in the V1 loop; Y52 and Y53 in the

CDRH2 loop, which interact with D325 of the GDIR motif in the V3 region; R109 in the CDRH3

loop, which engages with glycan N332 in the V3 region; and T413 in the V4 loop of gp120 (Figure

5D). These interactions are primarily stabilized through hydrogen bonding and van der Waals

forces. The glycan interaction at the 330 Env-44m Fab junction involves N146 from the V1 loop

and N301 and N332 from the V3 loop (Figure 5E). The D235 residue was shielded within the

hydrophobic cleft formed by the V3 loop of the gp120 region (Figure 5F). Next, we performed

rigid-body docking of the 44m-BG505 atomic model (EMDB-36815) into our cryo-EM map

(Figure S7). This analysis allowed us to investigate the binding modes of 44m[36] and other V3-

glycan specific bNAbs 10-1074[53], PGT121[54], 438-B11[55], BG18[56], mature BG18[57], DH270.6[58], and

BF520.1[59] to a 330 Env superimposed BG505 model. Our findings suggest that 44m interacts

with the V3 loop via a slightly declining shift, similar to BF520 and BG18, but exhibits a distinct

binding mode compared to PGT121 and 35022 (Figure S7).

**Structural and glycan homology of 330 Env to BG505**

To assess structural similarity, The HIV-1 Clade C 330 SOSIP trimer was superimposed

onto a clade C Env 16055 NFL[60] (PDB: 5UM8) and a Clade-A Env BG505 SOSIP[61] (PDB: 4TVP). We

observed that the 330 Env trimer exhibited greater structural similarity to BG505 Env, with

lower root-mean-square deviations (RMSDs) of 0.685 Å for gp120 and 0.478 Å for gp41

compared to modelled 16055 Env, which had RMSDs of 0.798 Å for gp120 and 0.739 Å for gp41

(Figure 4A-D). Upon glycan prediction analysis by Los Alamos HIV-1 sequence database

(LANGT), we observed that 330 Env lacked conserved PNGS at positions 295 and 339 in the

V3/C4 region and at position 130 in the V1V2 apex, in contrast to BG505 Env, which lacks PNGS

at positions 241 and 289 (Figure 4E). Since BG505-mediated immune responses are directed

against positions 241 and 289 in the V3 loop[62–64], the glycan hole in the V1V2 and V3/C4 region

of the 330 Env may favor the elicitation of immune responses focused on these regions. Next, we

calculated the total glycan hole area in the 330 Env using Los Alamos glycan shield mapping tool

(LAGSM), suggesting that the optimal protein accessible area was 967 $Å^2$, compared to BG505,

B41, C9Z7A & Du422 counterparts (Figure 4F). The potential N-glycosylation sites (PNGS)

conserved in the 330 Env were determined by Los Alamos HIV-1 sequence database (LANGT)

using N-Glycosite tool. Conserved PNGS analysis revealed that 330 Env comprises of 28 PNGS

comparable to that reported in the BG505[65] & 16055[66] Envs (Figure 4F, Table S3). Targeting

exposed protein surfaces within glycan holes can be an effective strategy to boost the immune

response against these vulnerable sites. Overall, our structural homology modelling and glycan

analysis demonstrate that the stabilized clade-C Env 330 shares numerous features with the

well-studied BG505 Env, suggesting its capacity to adopt well-ordered structures.

**330 Ferritin nanoparticles and SOSIP trimer exhibit enhanced immunogenicity compared**
**to NFL in rabbits**

Following the establishment of native-like prefusion-closed Env structures, we next

evaluated and compared the immunogenicity of 330 Env trimers (SOSIP & NFL) and multivalent

ferritin NPs in New Zealand White (NZW) rabbits. Four groups of female rabbits were

immunized, primed at week 0, and boosted at weeks 4 and 20 and administered a dose of 20 μg

of 330 gp140 Env trimers or an equimolar amount (25 μg) of ferritin nanoparticles, formulated

with AddaVax™ adjuvant (Figure 6A). The AddaVax™ adjuvant is a squalene-based oil-in-water

nano-emulsion and has been shown not to impact the structural integrity or antigenicity of the

Env trimers, as demonstrated in previous studies[67,68]. Rabbits were bled at weeks 0, 4, 6, 16, 20,

and 22 and sera samples were subsequently evaluated for binding reactivity with the Env trimer

using capture ELISA and for neutralizing activity against Env pseudotyped viruses. First, we

evaluated the binding titers of the collected immune sera from different timepoints against the

immunogens tested and measured as median titers across each group. The binding titers of

immune sera exhibited a significant increase between 4 to 8 weeks and then again between 20 to

22 weeks, immediately after the second and third boosters (Figure 6B). The polyclonal immune

sera demonstrated sustained binding reactivity to the autologous Envelope trimer and with the

cross-clade Envelopes, though with a lower extent to heterologous Envs. The half-lives ($t1/2$) of

the immune sera titers against the autologous Env were significantly longer (p value: 0.007) as compared to that of the heterologous Envs (SF162, 6240, 1086) tested. The overall immune sera binding titers were comparable across the test groups; however, rabbits immunized with 330 SOSIP and ferritin NPs exhibited statistically significantly antibody titers (p value: 0.01) compared to those immunized with NFL counterparts. Peak serum antibody binding to 330 Env was observed at week 22, prompting further analysis of week 22 serum samples for their neutralization activity.

Neutralization breadth was further evaluated against autologous (330.16.E6), closely related (329.14.B1), and heterologous clade C (MW965.26 and 25925), Tier 1 clade B (SF162), and Tier 2 clade A (BG505.T332N) pseudoviruses. No immune sera response was detected against murine leukemia virus (MLV) used as a negative control and all animals exhibited neutralizing antibody responses following immunization, except for one in the NFL group (NZW5273) (Table S4). Multivalent 330 ferritin NPs and SOSIP elicited a strong neutralizing response against autologous (330.16.E6) and closely related Envs (329.14.B1) as compared to NFL counterpart. The Ferritin NP group elicited the most robust autologous neutralizing antibody response, with a median $ID_{50}$ titer of 1160 compared to the soluble gp140 trimer groups (SOSIP and NFL) exhibited median $ID_{50}$ titers of 830 and 375, respectively (Figure 6C). The Ferritin NP group demonstrated a statistically significant increase in neutralizing antibody titers compared to both the SOSIP (p = 0.014691) and NFL (p = 0.010323) groups, with fold differences of 2.5-fold and 3.5-fold, respectively. These findings are consistent with the results from the capture ELISA, where the NP and SOSIP groups exhibited sustained binding reactivity, whereas the NFL group showed a decline in response after boosters. The NP and SOSIP groups demonstrated moderate heterologous neutralizing responses against the clade C Tier 1 viruses (MW965.26 and 25925) and the Tier 1 clade B virus (SF162), with responses comparable to each other and higher than the NFL group although the differences were not statistically significant. None of the 330 immunogens induced neutralizing responses against the Tier 2 BG505 virus, which aligns with the low BG505 nAb activity observed in the plasma of the AIIMS_330 HIV-1-infected elite neutralizer[35]. Overall, we observed that both 330 SOSIP Env trimer and ferritin NPs elicited strong autologous NAb responses, as well as cross-clade neutralization, though with a limited magnitude. Notably, the presentation of these antigens on ferritin NPs was associated with a higher trend in NAb response induction.

## DISCUSSION

HIV-1 clade-C accounts for more than half of global HIV-1 infections, with particularly high prevalence in South Africa and India[19,20]. Comparative analyses of clade-C viruses from these regions reveal the significant differences in their sequences, highlighting distinct evolutionary trajectories[69]. These regional variations in intra-clade-C Env sequences likely contribute to the differential sensitivity of circulating strains to HIV-1 nAbs. HIV-1 Clade-C characterized by unique features such as enhanced transmission efficiency, extensive genetic diversity within the V1-V3 region of the Env gene, and a distinct profile of drug resistance mutations compared to other clades[69–72]. Understanding these differences is crucial for tailoring the prevention and treatment strategies in clade C prevalent regions[19,20]. Structural information on the HIV-1 Env trimer from clade-C is crucial for structure-guided immunogen design. However, only a limited number of high-resolution structures are available for native-like clade-C Env trimers, such as 16055[50,73], DU172.17[74], 426c[75], and CZA97.12[76]. Although many HIV-1 Env immunogens have been stabilized, most studies have extensively characterized BG505, a Clade-A transmitted founder Env derived from a 6-week-old infant with elite neutralizing activity[11]. Such elite individuals from the pediatric group could provide valuable Env immunogen templates, as HIV-1 infection and disease progression in children differs from that in adults[31,34,35]. Notably, neutralizing antibodies tend to arise early in the course of infection in pediatric individuals, with the ability to neutralize contemporaneous viral Envelopes[33,37]. We previously reported a pair of monozygotic twins (AIIMS_329 and AIIMS_330) with elite neutralizing activity, whose plasma demonstrated exceptionally potent bNAb responses with multi-epitope specificities against a diverse panel of multi-clade heterologous Env pseudoviruses[33,35]. These Envs hold significant potential as templates for vaccine design and as antigenic baits to isolate HIV-1 bNAbs for immunotherapeutic applications[49]. Recent studies demonstrated that, multimeric presentation of viral class I fusion viral glycoproteins, such as , Ebola virus Glycoprotein (GP)[77,78], influenza hemagglutinin (HA)[79,80], Respiratory Syncytial Virus (RSV) F protein[81,82], SARS-CoV-2 S protein[83–86] and HIV-1 Env[29, 30] on ferritin NPs has enabled the efficient production of well-ordered NP-based vaccine candidates with a capacity to induce robust immune response.

Herein, we designed and characterized a soluble HIV-1 clade-C 330 Env trimer on SOSIP and NFL platform, derived from a circulating virus in an Indian pediatric EN AIIMS_330[35], by stabilizing the Env using version 7 mutations and displaying the SOSIP Env on self-assembling ferritin NPs[29,30]. The purification yield of Env trimers post SEC was around ~1 mg/L for SOSIP

and ~1.45 mg/L for NFL, which was higher or comparable to the Indian and South African Clade-
C strains (CZA97: ~0.4-0.6 mg/mL, 329 SOSIP: ~0.6 mg/mL, 16055 NFL: ~2 mg/mL, 1PGE-
THIVC SOSIP)[23,48–50]. Using Negative stain Electron Microscopy, we demonstrated that SOSIP and
NFL Envs displaying key features of native-like trimeric conformation and Env trimers were
presented efficiently on ferritin NPs as observed in stabilized BG505 Env-particles[29]. The
structure of the 330 SOSIP, resolved by single-particle cryo-EM, revealed trilobular Env
structures in their prefusion, closed conformation. The generated Env trimer and particles
displays a desirable antigenic profile, with nanomolar affinity to conformation specific bNAbs
(PGT145 & PGT151), with negligible reactivity to non-nAbs (F105 & 17b+sCD4), shielding the
Env immunodominant regions. Biophysical characterization showed  that the 330 Env exhibits a
balanced thermodynamic profile and high degree of thermal stability with a Tm of ~74°C which
is notably higher as compared to other reported clade C Envs ZM197 (69°C)[12] , IPGE-THIVC
(62°C)[23] and 16055 (68°C)[87]. We observed greater structural similarity of 330 Env to clade-A
BG505 Env (RMSD gp120-0.685 Å & gp41-0.478 Å) compared to clade-B AMC009 Env reported
(RMSD gp120-2.623 Å & gp41-1.624 Å),  suggesting well-ordered structures[14]. Cryo-EM
structure of 330 Env in complex with autologous bNAb 44m revealed the well-exposed epitopes;
the three trilobular structures occupied by the three Fab arms as observed with BG505-44m
structure. We  identified herein, additional key antigenic determinants T144 and T413 in the
autologous 330 Env, that interacted with 44m; , in addition to  D325 and N322 dependency, as
observed with heterologous BG505 Env[36]. Our findings offer crucial insights into the antigenic
determinants in the clade C virus Env, that interact with the autologous antibody 44m, that can
inform a clade-based vaccine design. Furthermore, using a reverse vaccinology approach, these
insights can guide the structure-based engineering of neutralizing antibodies[88].

Rabbit immunizations demonstrated that the 330 Env immunogens elicited robust
immune responses against autologous and other tier-1 viruses, with limited cross-clade
neutralization. This may be attributed to distinct antigenic determinants on heterologous Envs,
adjuvant used or by varied glycan compositions[89,90]. Further insight can be achieved by
investigating the specific epitopes targeted by NAbs in the immune sera, however it has not been
addressed herein and is a limitation of this study. The multivalent presentation of HIV-1 antigens
on ferritin NPs correlated with enhanced neutralization potency, likely attributed to augmented
B-cell activation signals[91]. This observation aligns with findings from germline-targeting eOD-
GT8 NPs, suggesting a potential mechanism for eliciting more robust and coordinated immune

responses[91]. These findings emphasize the need for further exploration of the role of glycans and

structure guided mutations in shaping immunogenicity.

In summary, we designed and characterized soluble HIV-1 Env trimers derived from a

clade-C-infected pediatric elite-neutralizer from India. Structural analysis of the 330 Env trimer

in complex with autologous the bNAb 44m-a heavy chain matured version of the first bNAb

identified from AIIMS_330 donor, revealed molecular features of 330 Env-44m bNAb interaction

which could inform structure-based vaccine design. Our findings indicate that the 330 SOSIP Env

trimer can serve as a valuable antigenic probe for bNAb discovery and that both the SOSIP and

ferritin nanoparticle versions represent promising candidates for multivalent HIV-1 vaccine

design and development.

**MATERIALS AND METHODS**

**Institutional Review Board & Ethics approval**

This study was conducted after obtaining approval from the institutional ethics committee, All

India Institute of Medical Sciences (AIIMS), New Delhi, India (250/IAEC-1/2020). Rabbit

immunization was outsourced to a contract research organization (CRO) Liveon Biolabs Private

Limited, Bengaluru, Karnataka, India with IAEC (IAEC-approved protocol No.: LBPL-IAEC-008-

01/2021; study number: LBPL/NG-1736 (EF)). The immunization studies described here were

carried out on female naive New Zealand White rabbits of 2.0–2.5 kg and aged 4 months.  All

animal experiments adhered to the guidelines for the care and use of laboratory animals as

outlined by the Committee for Control and Supervision of Experiments on Animals (CCSEA),

Government of India[92].

**Construct design**

The 330 Env gene was derived from a previously identified Indian clade C HIV-1 Env (330.16.E6,

GenBank: MK076720) obtained from a pediatric elite-neutralizer AIIMS_330, as previously

described[35]. The 330 Env construct were design in soluble forms as SOSIP.664[11] gp120-gp41

cleaved Env trimers and NFL.664[38] gp140 single-chain Env trimers along with tailoring of SOSIP

Env trimers on self-assembling ferritin nanoparticles[29] to achieve multivalency. SOSIP & NFL

constructs were stabilize by incorporating I559P & ΔMPER (Env prefusion stabilizing

mutation)[42,43], R315Q (reduced V3 non-nAb immunodominant exposure)[44], I201C-A433C

(intraprotomer locking disulfide bond, reduced V3 exposure)[12], and clade C stabilizing B505-

Trimer Derived (TD8) mutations (E47D, T49E, V65K, I165L, K429R, K432Q, K500R)[39]. The SOSIP construct was designed by introducing an SOS disulfide bond (A501C-T605C)[46], an I559P mutation[42], and an optimized furin cleavage site (RRRRRR)[47]. In contrast, the NFL (Native Flexible Linker) construct retained the I559P mutation and linked gp120 and gp41 using a flexible glycine-serine linker (GGGGSGGGGS)[38]. Native signal sequence is replaced by the CD5 signal peptide (MGSLQPLATLYLLGMLVASVLA) for enhanced secretion of Envelope proteins, constructs were codon optimized for mammalian expression and synthesized commercially by GeneArt (Thermo Fischer Scientific). For binding assays, 330 Env constructs were tagged with either a His-tag (GSGSGGSGHHHHHHHH) or a twin strep tag (tst) (WSHPQFEKGGGSGGGSWSHPQFEK) at the C-terminus of gp41ECTO, preceding the stop codon. These constructs were subsequently cloned into pPI4 expression vector using Pst1 and Not1 restriction sites. The 330 ferritin NP's construct was generated by fusing the N-terminus from *Helicobacter pylori* ferritin (GenBank: WP_000949190.1), starting from Asp5 to Ser167, to the SOSIP.664 C-terminus (truncated at position 664), separated by a Gly-Ser (GSG) linker, as described previously[29].

**Envelope Protein Expression**

The 330 Env encoding plasmids were transiently expressed in 293F suspension cells (Gibco, R790-07). 293F cells were thawed and maintained in Freestyle Expression Medium (Gibco, K9000-01) in a Shaker incubator at 37°C, with 120 r.p.m. and 8% CO2. For transfection, HIV-1 Env and furin protease encoding plasmids mixed in a 3:1 ratio (w/w) with NFL (500 µg), SOSIP Env (333.3 µg), to furin (166.6 µg) incubated with transfection reagent PEImax (Polysciences, 24765) in a 3:1 ratio (w/ w) PEI max to DNA for 20 min at room temperature (RT) and then added to 1L culture at density of 1 million cells/mL. Seven days post-transfection, the supernatants were harvested, centrifuged, and filtered using 0.45-mm filters (Nalgene, 2954545).

**Envelope Protein Purification**

330 SOSIP Env and SOSIP Env-NP fusion proteins were purified by PGT145 bNAb immunoaffinity chromatography, as described previously[48]. Briefly, HIV-1 pre-fusion closed trimer-specific bNAb PGT145[93] was covalently coupled to the CNBr-activated Sepharose 4B beads (GE Healthcare, 17043001) according to the manufacturer's protocol for antibody coupling. The PGT145-coupled CNBr Sepharose 4B bead column was used to purify HIV-1 Env

trimers or Env trimer NPs. Env proteins contained in Expi293F filtered supernatants were captured on PGT145-coupled CNBr-activated sepharose beads by overnight rolling incubation at 4 °C. Subsequently, the mixture of supernatants and beads was passed over Econo-columns (Biorad, 7371512). The columns were then washed with three column volumes of 0.5 M NaCl and 20 mM Tris HCl pH 8.0 solution. After elution with 3 M $MgCl_2$ pH 7.5, the proteins were buffer-exchanged into TN75 (75 mM NaCl, 20 mM Tris HCl pH 8.0) or PBS buffer by ultrafiltration with Vivaspin concentrators MWCO 100 kDa (Cytivia, 28932363). Protein concentrations were determined from the A280 values measured using a NanoDrop One (Thermo Fisher Scientific, 29018226) and the molecular weight and extinction coefficient values were calculated by the ProtParam Expasy webtool. Env proteins used for immunization were first purified using PGT145 affinity chromatography, followed by size-exclusion chromatography (SEC) on a Superdex 200 Increase 10/300 GL column (28990944, GE Healthcare Life Sciences) using an Äkta Pure 25M system (Cytiva, 29018226). Envs containing fractions were pooled, concentrated, filter-sterilized, snap-frozen in liquid nitrogen, and stored at -80°C until further use.

**SDS-PAGE and BN-PAGE Analyses**

For SDS-PAGE and blue native polyacrylamide gel electrophoresis (BN-PAGE) analyses, 2 μg of 330 gp140 Env trimers, or equimolar amounts of ferritin NP's (2.5 μg) was ran over precast native gel 4–20 % Bis-Tris gradient gel (Biorad, 4561094). The degree gp120-gp41 cleavage was examined by incubating the SOSIP & NFL protein with 0.1M dithiothreitol (DTT) and analysed by SDS-PAGE under reducing conditions. Subsequently, gels were stained with the Colloidal Blue Staining Kit (Invitrogen, LC6025), respectively.

**Enzyme linked Immunosorbent assay (ELISA)**

ELISAs were performed as previously described[94] using tst-tagged Env constructs on StrepTactinXT-coated microplates (IBA Lifesciences, 2-4101-001). Purified proteins (1 μg/mL) were diluted in NaHCO3 buffer and immobilized on 96-well Streptactin plates by O/N incubation at 4°C. Following a three wash step with PBST (0.05% Tween20) to remove unbound trimers, plates were blocked by 1% BSA in PBST for 2 hours at RT. Further the serial dilutions of test antibodies ranging from 10ug – 0.004 ug/ml in PBST/1% BSA solution were added and incubated for 2 h. After 3 washes with PBST, HRP-labelled goat anti-human IgG (Jackson ImmunoResearch, AB_2337578) diluted 1:3000 in PBST/1% BSA was added and incubated for

30 mins, followed by 3 washes with PBST and 3 washes of PBS. A developing solution (1% TMB

(Mabtech, 3652-F10), allowed the colorimetric reaction, which was stopped by the addition of 2

N H2SO4. Finally, colour development (absorption at 450 nm, OD450) was measured at

NanoDrop One (Thermo Fisher Scientific, 29018226) to obtain the different binding curves.

**Biolayer Interferometry (BLI)**

Binding kinetics were measured using an Octet K2 (ForteBio, RED96e) device at RT with 1000

rpm agitation. Anti-human Fc sensors (Sartorius AHC, 18-5060) were used to capture mAbs on

the Octet K2, were equilibrated in kinetics buffer for 60 s to obtain a baseline prior to protein

association. 330 Envelope protein was used as analyte in concentrations ranging from 210 to 2.6

nM in the HEPES buffer (pH 7.2) background supplemented with 0.02% Tween 20 and 0.1%

BSA. Binding was assessed by incubating mAb-captured sensors in wells containing 10 μg/ml

Env for 120 seconds with 1000 rpm agitation. Association was recorded for 150 s followed by

dissociation for 450 s. Data were analysed using the ForteBio Data analysis software (Forte- Bio

Inc, Version 9.0) and a global fit was performed using a 1:1 binding model to fit the association

and dissociation curves.

**Sypro orange differential scanning fluorimetry (DSF)**

330 SOSIP protein were prepared at a concentration of 0.2 mg/mL in 1XPBS and transferred into

0.1 mL polypropylene PCR tube strips (Tarsons, 611020) with a final volume of 25 μL. Sypro

Orange protein stain (Invitrogen, 4461146) was added at a 1:200 dilution. The samples were

analyzed using a Rotor-Gene Q system (Qiagen, 9001862) with a high-resolution melt (HRM)

protocol, starting at 25°C and increasing to 90°C in 1°C increments, with each step lasting 4

seconds. Melt curve data were processed using Rotor-Gene Q Series Software.

**Circular dichroism (CD) spectroscopy**

The Far-UV spectrum was recorded in the wavelength range of 200–250 nm using a CD

Spectrophotometer (JASCO, J-1500) with a 0.1 cm path length Quartz cuvette (Sigma Aldrich,

I0285). Measurements were conducted for the 330 protein in a 20 mM phosphate buffer (pH 7.5)

with100 mM NaCl concentration at 4°C for 30 minutes. Each spectral scan was averaged over 2–3

measurements. The CD spectra were visualized using SigmaPlot 10.0, and the secondary

structure was analyzed through the BESTSEL online server. For thermal studies, measurements

were performed between 20℃ and 80℃ using the same Quartz cuvette, and the melting curve

was generated using SigmaPlot 10.0.

**Negative Stain Microscopy (NS-EM)**

Negative-stain electron microscopy (NS-EM) was performed essentially as described

elsewhere[95]. Briefly, 3 μl of purified 330 gp140 Env trimers or ferritin NPs (~0.03 mg/mL) were

applied to glow discharged (20 mA for 30 s) carbon-coated 400 Cu mesh grid. After 5 s, grids

were negatively stained with 2% (w/v) uranyl formate for 60 s. Imaging was performed using a

120 kV Talos L120C electron microscope (TALOSL120CMS) equipped with a 4k x 4k Ceta camera

at a magnification of 73kx (3.84 Å/pixel). The collected images were processed in EMAN 2.1[96].

Particle extraction and reference-free 2D class averages were generated using simple_prime2D

in SIMPLE 2.1 software[97] with a mask diameter of 30 pixels at 3.84 Å/pix.

**Preparation of 44m Fab fragment**

Fab fragments were generated from 4 mg of AIIMS-44m IgG using a Fab fragmentation kit (G

Biosciences, 786-273) following the manufacturer's protocol. Briefly, 0.5 ml of 44m IgG was

desalted using a SpinOUT™ GT-600 column. Cysteine.HCl was added to the Fab Digestion Buffer

(pH) to a final concentration of 20 mM before papain digestion. The 0.5 ml IgG sample was then

incubated with Immobilized Papain in the spin column at 37°C for 6 hours with end-over-end

mixing. After incubation, the digested antibody solution was collected by centrifugation at 5,000

x g for 1 minute. The flow-through was further incubated with Protein A resin for 10-15 minutes

at room temperature (RT) with end-over-end mixing. Subsequently, the Fab fragments were

collected in the flow-through while Fc fragments and undigested IgG remained bound to the

resin. The purity and size of the Fab fragments were assessed by SDS-PAGE and the purified Fab

fragments were stored at -80°C.

**Cryo-EM sample preparation, data acquisition, data analysis and model fitting**

330 Env trimer (1.5 mg/ml) incubated with 44m Fab at 1:3 molar ratio. 3 μl of sample were

applied to glow discharged R1.2/R1.3 300 mesh copper grids and immediately blotted for 8 secs

without any blotting force using FEI Vitrobot Mark IV plunger at 22°C. The grid was immediately

plunge-frozen into liquid ethane at liquid nitrogen temperature. Cryo-EM data were collected

using 200 kV Talos Arctica transmission electron microscope (Thermo Scientific,

TALOSARCTICA) equipped with Gatan K2 Summit Direct Electron Detector. Movies were

recorded automatically using Latitude-S (DigitalMicrograph - GMS 3.5) at a nominal magnification of 45,000x at the effective pixel size of 1.17 Å[97]. Micrographs were acquired in counting mode with a total dose of 60 e$^-$/Å$^2$, with an exposure time of 8 sec distributed for 20 frames. A total of 5215 movies were acquired for the 330 SOSIP Env trimer and 44m Fab complexes, respectively. Single-Particle Analysis (SPA) was performed for the acquired Cryo-EM movies using the Relion version 3.1[98] and CryoSPARC v4.2.1[99]. At first, drift and gain corrections of the individual movies were performed with MotionCorr2[100] and estimated Contrast transfer function (CTF) parameters using CTFFIND 4.1.13 in Relion[101]. Subsequently, CTF-estimated micrographs were subjected to analysis to eliminate bad micrographs using cisTEM[102], and also to remove poor resolution micrographs with a fit resolution threshold of 7 Å. The particles from the best micrographs were chosen for automated picking using 2D reference in Relion and extracted with the box sizes of 320 pixels for the 44m Fab bound 330 SOSIP Env trimer complexes. The extracted particles were imported to CryoSPARC v4.2.1, and the rest of the processing was performed there. After four rounds of rigorous 2D classification, 879,956 particles were selected from the best 2D classes with high-resolution features of the Ab-bound complex. This dataset was divided into 6 classes for Ab initio reconstruction with C3 symmetry. The class04, with 109004 particles, had the best Ab bound features among all the classes (Figure S6). Therefore, we considered the class04 for non-uniform refinement by imposing C3 symmetry. The class04 was further refined by local refinement, and the resolution of the 3D map was estimated at FSC 0.143. For model fitting, the Cryo-EM maps of 330 Env trimers were docked on ferritin structure (PDB: 3BVE) symmetrically on four sites. Overviews of Cryo-EM data processing are shown in S. Table S3.

**Immunizations**

The rabbit immunization studies were conducted at a contract research organization (CRO), Liveon Biolabs Private Limited, Karnataka, India, under GLP-compliant conditions. Liveon Biolabs is an AAALAC International-accredited facility and is registered with the CCSEA, Government of India. Female, naive New Zealand White rabbits, aged 4 months and weighing between 2.0–2.5 kg, were used for the study. The protocol for animal use was approved by the Institutional Animal Ethics Committee (IAEC) of Liveon Biolabs (Approval No.: LBPL-IAEC-008-01/2021; Study No.: LBPL/NG-1736 (EF)). Prior to housing, the experimental room underwent fumigation for decontamination, and microbial load was assessed using the settle plate method. The rabbits were kept in a temperature-controlled Environment (20 ± 3 °C) with relative

humidity ranging from 30–70%, a 12-hour light/dark cycle, 12–15 air changes per hour, and a noise level below 80 dBs. Mortality and morbidity were monitored twice daily, while clinical observations were conducted at least once per day throughout the study. Body weight measurements were recorded for all animals prior to the administration of the test item on Day 1 and subsequently on a weekly basis (±1 day) during the treatment period. The gp140 SOSIP and NFL Env immunogens (G1 and G2) were administered at a dose of 20 µg of trimers, or an equivalent molar amount SOSIP-ferritin nanoparticles (25 µg) (G3) formulated with the AddaVax adjuvant (1:1 v/v) (vac-adx-10, Invivogen). The placebo group (G4) received only adjuvant with sterile 1XPBS. Doses were calculated based on the molecular weight of the protein peptides, excluding glycans, as described elsewhere[66,66]. Blood samples were collected from rabbits at weeks 0, 4, 6, 16, 20, and 22. Before blood collection from the marginal ear vein, a local anaesthetic cream containing 4% lidocaine was applied for 5 minutes. The immunization process, including bleeding protocols, caused minimal and brief pain or distress, with no significant impact on animal health. At the end of the experimental period, the rabbits were euthanized using a lethal dose of sodium thiopental. Complement-deactivated serum samples were subsequently analysed for their binding reactivity and neutralization potential.

**HIV-1 pseudovirus generation**

The HIV-1 pseudoviruses were produced in HEK293T cells as described previously[103] by co-transfecting the corresponding full HIV-1 gp160 Envelope plasmid and a pSG3ΔEnv backbone plasmid. Briefly, $1 \times 10^5$ cells in 2 mL complete DMEM (10% fetal bovine serum (FBS) (Gibco, 26140079 and 1% penicillin and streptomycin (Gibco, 15140-122) were seeded per well of a 6-well cell culture plate (Costar, 3506) the day before transfection. For transfection, Envelope (1.25 µg) and delta Envelope (pSG3ΔEnv backbone) plasmids (2.50 µg) were mixed in a 1:2 ratio in Opti-MEM (Gibco, 31985062), with a final volume of 200 µL per well, and incubated for 5 min at room temperature. Next, 3 µL of PEI-Max transfection reagent (Polysciences, 24765) (1 mg/mL) was added to this mixture before further incubation for 15 min at room temperature. This mixture was then added dropwise to the HEK 293T cells supplemented with fresh complete DMEM growth media and incubated at 37 °C for 48 h. Pseudoviruses were then harvested by filtering cell supernatants with 0.45 µm sterile filters (mdi), aliquoted, and stored at −80 °C until usage.

**HIV-1 Neutralization assay**

Neutralization assays were carried out using TZM-bl reporter cells which is described elsewhere[104]. The autologous and heterologous NAb response of end point sera was measured in TZM-bl cell neutralization assays. Neutralization was measured as a reduction in luciferase gene expression after a single round of infection of TZM-bl cells (NIH AIDS Reagent Program) with HIV-1 Env pseudoviruses. The $TCID_{50}$ of the HIV-1 pseudoviruses was calculated and 200 $TCID_{50}$ of the virus was used in the neutralization assays through incubation with three-fold serially diluted rabbit sera starting at a 1:20 dilution. Next, freshly trypsinized TZM-bl cells in a complete DMEM (10% fetal bovine serum (FBS) (Gibco, 26140079 and 1% penicillin and streptomycin (Gibco, 15140-122) containing 50 μg/mL DEAE-dextran (Sigma, D9885) at $10^5$ cells/well were added and the plates were incubated at 37 °C for 48 h. Virus controls (cells with HIV-1 virus only) and cell controls (cells without virus and antibody) were included. MuLV was used as a negative control. After the incubation of the plates for 48 h, luciferase activity was measured using the Bright-Glow Luciferase Assay System (Promega, E2610). $ID_{50}$ for antibodies were calculated from a dose–response curve fit with a non-linear function using GraphPad Prism 8 software (San Diego, CA, USA). All neutralization assays were repeated at least 2 times, and the data shown are from representative experiments.

**Statistical analysis**

Statistical significance between groups and subgroups was assessed using the Mann–Whitney U test, Student's t-test, and one-way analysis of variance for continuous variables. For representative results, experiments was done in replicates or repeated twice. All statistical analyses and plotting were performed using GraphPad Prism (version 8.0; La Jolla, CA, USA)

**FUNDING**

This study was funded by the Department of Biotechnology (DBT), India (BT/PR39156/DRUG/134/91/2021, BT/PR30120/MED/29/1339/2018) grants awarded to K.L. This work was also supported by Department of Science & Technology-FIST (SR/FST/LSII-039/2015) and DBT-BUILDER Program (BT/INF/22/SP22844/2017) grants awarded to S.D. S.K. is supported by the DBT-Wellcome Trust India Alliance Early Career Fellowship Grant IA/E/18/1/504307. The funders played no role in the study design, data collection, analysis, or interpretation, nor in the writing of the manuscript or the decision to publish the results.

**ACKNOWLEDGMENTS**

The authors are grateful to the NIH AIDS reagent program for providing the HIV-1 research reagents and the Neutralizing Antibody Consortium (NAC), IAVI, USA for providing the HIV-1 neutralizing antibodies donated by Michel Nussenzweig, Hermann Katinger, Mark Connors, James Robinson, Dennis Burton, John Mascola, Peter Kwong, and William Olson. We express gratitude to Prof. R.W. Sanders for generously providing the PPI4 expression vector and the training provided to S.D.S as a part of his HIV-1 Trust Fellowship. We appreciate the support of Liveon Biolabs for conducting the immunization studies. We also extend our appreciation for the IAVI-Leader Development Program awarded to S.D.S.

**AUTHOR CONTRIBUTIONS**

Conceptualization and implementation; S.D.S., S.Ku, and K.L., EM work & data acquisition; S.D.S., A.C., Formal analysis; S.D.S., S.Ku, K.L., S.D., Biophysical analysis; A.C., S.D.S., L.M., B.M., E.A.S., Expression and purifications: S.D.S., A.R., I.A., S.S., V.I., A.G., T.B., & R.Ku., ELISA & neutralization assays; S.D.S., S.S, A.R., S.Ka, A.W.H., Animal studies; S.D.S., S.Ka, H.B., A.G., & A.W.H. Octet experimentation; S.D.S., R.Ku., S.A., and M.P., Original manuscript writing; S.D.S., S.Ku, A.C., & K.L., Supervision; S.D., & K.L., Project administration; S.D.S. & S.Ku, funding acquisition; S.D., & K.L. All authors have read and agreed to the published version of the manuscript.

**CONFLICT OF INTERESTS**

An Indian patent application (202211004612) has been filed on the 330 SOSIP Env trimer described in this study. The following authors: K.L., S.D.S., S.Ku. and S.S. are listed as inventors on the patent. The other authors declare no conflicts of interest.

**DATA AVAILABILITY**

The modified 330 Envelope protomer sequence been deposited in GenBank with accession number XXXX. Cryo-EM maps & atomic coordinates of reported structures are deposited into the Electron Microscopy Data Bank (EMDB) with accession code EMD: XXXX for 330 SOSIP Env trimer in complex with bNAb 44m. Any additional data are available upon reasonable request from the corresponding authors.

**REFERENCES**

1. Burton, D. R. *et al.* A Blueprint for HIV Vaccine Discovery. *Cell Host Microbe* **12**, 396–407 (2012).

2. Simek, M. D. *et al.* Human Immunodeficiency Virus Type 1 Elite Neutralizers: Individuals with Broad and Potent Neutralizing Activity Identified by Using a High-Throughput Neutralization Assay together with an Analytical Selection Algorithm. *J. Virol.* **83**, 7337–7348 (2009).

3. van den Kerkhof, T. L. G. M. *et al.* Early development of broadly reactive HIV-1 neutralizing activity in elite neutralizers. *AIDS* **28**, 1237 (2014).

4. Mascola, J. R. *et al.* Protection of macaques against vaginal transmission of a pathogenic HIV-1/SIV chimeric virus by passive infusion of neutralizing antibodies. *Nat. Med.* **6**, 207–210 (2000).

5. Mascola, J. R. *et al.* Protection of Macaques against pathogenic simian/human immunodeficiency virus 89.6PD by passive transfer of neutralizing antibodies. *J. Virol.* **73**, 4009–4018 (1999).

6. Parren, P. W. *et al.* Antibody protects macaques against vaginal challenge with a pathogenic R5 simian/human immunodeficiency virus at serum levels giving complete neutralization in vitro. *J. Virol.* **75**, 8340–8347 (2001).

7. Moldt, B. *et al.* Highly potent HIV-specific antibody neutralization in vitro translates into effective protection against mucosal SHIV challenge in vivo. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 18921–18925 (2012).

8. Hessell, A. J. *et al.* Broadly neutralizing human anti-HIV antibody 2G12 is effective in protection against mucosal SHIV challenge even at low serum neutralizing titers. *PLoS Pathog.* **5**, e1000433 (2009).

9. Karasavvas, N. *et al.* The Thai Phase III HIV Type 1 Vaccine Trial (RV144) Regimen Induces Antibodies That Target Conserved Regions Within the V2 Loop of gp120. *AIDS Res. Hum. Retroviruses* **28**, 1444–1457 (2012).

10. Kim, J., Excler, J.-L. & Michael, N. Lessons from the RV144 Thai Phase III HIV-1 Vaccine Trial and the search for correlates of protection. *Annu. Rev. Med.* **66**, (2014).

11. Sanders, R. W. *et al.* A Next-Generation Cleaved, Soluble HIV-1 Env Trimer, BG505 SOSIP.664 gp140, Expresses Multiple Epitopes for Broadly Neutralizing but Not Non-Neutralizing Antibodies. *PLOS Pathog.* **9**, e1003618 (2013).

12. Torrents de la Peña, A. *et al.* Improving the Immunogenicity of Native-like HIV-1 Envelope
Trimers by Hyperstabilization. *Cell Rep.* **20**, 1805–1817 (2017).

13. Torrents de la Peña, A. *et al.* Immunogenicity in Rabbits of HIV-1 SOSIP Trimers from Clades
A, B, and C, Given Individually, Sequentially, or in Combination. *J. Virol.* **92**, (2018).

14. Schorcht, A. *et al.* Neutralizing Antibody Responses Induced by HIV-1 Envelope Glycoprotein
SOSIP Trimers Derived from Elite Neutralizers. *J. Virol.* **94**, 10.1128/jvi.01214-20 (2020).

15. de Taeye, S. W. *et al.* Immunogenicity of Stabilized HIV-1 Envelope Trimers with Reduced
Exposure of Non-neutralizing Epitopes. *Cell* **163**, 1702–1715 (2015).

16. Sok, D. *et al.* Rapid elicitation of broadly neutralizing antibodies to HIV by immunization in
cows. *Nature* **548**, 108–111 (2017).

17. Sanders, R. W. *et al.* HIV-1 VACCINES. HIV-1 neutralizing antibodies induced by native-like
Envelope trimers. *Science* **349**, aac4223 (2015).

18. Houser, K. V. *et al.* Safety and immunogenicity of an HIV-1 prefusion-stabilized Envelope
trimer (Trimer 4571) vaccine in healthy adults: A first-in-human open-label, randomized,
dose-escalation, phase 1 clinical trial. *eClinicalMedicine* **48**, (2022).

19. Gartner, M. J., Roche, M., Churchill, M. J., Gorry, P. R. & Flynn, J. K. Understanding the
mechanisms driving the spread of subtype C HIV-1. *EBioMedicine* **53**, 102682 (2020).

20. Hemelaar, J., Gouws, E., Ghys, P. D. & Osmanov, S. Global and regional distribution of HIV-1
genetic subtypes and recombinants in 2004. *AIDS Lond. Engl.* **20**, W13-23 (2006).

21. Ahmed, S. *et al.* Stabilization of a soluble, native-like trimeric form of an efficiently cleaved
Indian HIV-1 clade C Envelope glycoprotein. *J. Biol. Chem.* **292**, 8236–8243 (2017).

22. Rutten, L. *et al.* A Universal Approach to Optimize the Folding and Stability of Prefusion-
Closed HIV-1 Envelope Trimers. *Cell Rep.* **23**, 584–595 (2018).

23. Kumar, R. *et al.* Elicitation of potent serum neutralizing antibody responses in rabbits by
immunization with an HIV-1 clade C trimeric Env derived from an Indian elite neutralizer.
*PLOS Pathog.* **17**, e1008977 (2021).

24. Kumar, S. *et al.* Design and immunogenicity of an HIV-1 clade C pediatric Envelope
glycoprotein stabilized by multiple platforms. *BioRxiv Prepr. Serv. Biol.* 2024.09.14.613016
(2024) doi:10.1101/2024.09.14.613016.

25. Schiller, J. & Chackerian, B. Why HIV Virions Have Low Numbers of Envelope Spikes:
Implications for Vaccine Development. *PLOS Pathog.* **10**, e1004254 (2014).

26. Zhao, Q., Li, S., Yu, H., Xia, N. & Modis, Y. Virus-like particle-based human vaccines: quality
assessment based on structural and functional properties. *Trends Biotechnol.* **31**, 654–663
(2013).
27. Ludwig, C. & Wagner, R. Virus-like particles-universal molecular toolboxes. *Curr. Opin.*
*Biotechnol.* **18**, 537–545 (2007).
28. Jennings, G. T. & Bachmann, M. F. The coming of age of virus-like particle vaccines. *Biol. Chem.*
**389**, 521–536 (2008).
29. Sliepen, K. *et al.* Presenting native-like HIV-1 Envelope trimers on ferritin nanoparticles
improves their immunogenicity. *Retrovirology* **12**, 82 (2015).
30. He, L. *et al.* Presenting native-like trimeric HIV-1 antigens with self-assembling nanoparticles.
*Nat. Commun.* **7**, 12041 (2016).
31. Goo, L., Chohan, V., Nduati, R. & Overbaugh, J. Early development of broadly neutralizing
antibodies in HIV-1–infected infants. *Nat. Med.* **20**, 655–658 (2014).
32. Ditse, Z. *et al.* HIV-1 Subtype C-Infected Children with Exceptional Neutralization Breadth
Exhibit Polyclonal Responses Targeting Known Epitopes. *J. Virol.* **92**, (2018).
33. Kumar, S. *et al.* An HIV-1 Broadly Neutralizing Antibody from a Clade C-Infected Pediatric
Elite Neutralizer Potently Neutralizes the Contemporaneous and Autologous Evolving
Viruses. *J. Virol.* **93**, e01495-18 (2019).
34. Makhdoomi, M. A. *et al.* Neutralization resistant HIV-1 primary isolates from antiretroviral
naïve chronically infected children in India. *Virology* **499**, 105–113 (2016).
35. Mishra, N. *et al.* Viral Characteristics Associated with Maintenance of Elite Neutralizing
Activity in Chronically HIV-1 Clade C-Infected Monozygotic Pediatric Twins. *J. Virol.* **93**,
e00654-19, /jvi/93/17/JVI.00654-19.atom (2019).
36. Kumar, S. *et al.* Recognition determinants of improved HIV-1 neutralization by a heavy chain
matured pediatric antibody. *iScience* **26**, 107579 (2023).
37. Simonich, C. A. *et al.* HIV-1 Neutralizing Antibodies with Limited Hypermutation from an
Infant. *Cell* **166**, 77–87 (2016).
38. Sharma, S. K. *et al.* Cleavage-Independent HIV-1 Env Trimers Engineered as Soluble Native
Spike Mimetics for Vaccine Design. *Cell Rep.* **11**, 539–550 (2015).
39. Guenaga, J. *et al.* Structure-Guided Redesign Increases the Propensity of HIV Env To Generate
Highly Stable Soluble Trimers. *J. Virol.* **90**, 2806–2817 (2015).
40. Joyce, M. G. *et al.* Soluble Prefusion Closed DS-SOSIP.664-Env Trimers of Diverse HIV-1
Strains. *Cell Rep.* **21**, 2992–3002 (2017).

41. Kumar, S. *et al.* B cell repertoire sequencing of HIV-1 pediatric elite-neutralizers identifies
multiple broadly neutralizing antibody clonotypes. *Front. Immunol.* **15**, 1272493 (2024).
42. Sanders, R. W. *et al.* Stabilization of the soluble, cleaved, trimeric form of the Envelope
glycoprotein complex of human immunodeficiency virus type 1. *J. Virol.* **76**, 8875–8889
(2002).
43. Klasse, P. J. *et al.* Influences on trimerization and aggregation of soluble, cleaved HIV-1 SOSIP
Envelope glycoprotein. *J. Virol.* **87**, 9873–9885 (2013).
44. de Taeye, S. W. *et al.* Immunogenicity of Stabilized HIV-1 Envelope Trimers with Reduced
Exposure of Non-neutralizing Epitopes. *Cell* **163**, 1702–1715 (2015).
45. Andrabi, R. *et al.* The Chimpanzee SIV Envelope Trimer: Structure and Deployment as an HIV
Vaccine Template. *Cell Rep.* **27**, 2426-2441.e6 (2019).
46. A Recombinant Human Immunodeficiency Virus Type 1 Envelope Glycoprotein Complex
Stabilized by an Intermolecular Disulfide Bond between the gp120 and gp41 Subunits Is an
Antigenic Mimic of the Trimeric Virion-Associated Structure | Journal of Virology.
https://journals.asm.org/doi/full/10.1128/JVI.74.2.627-643.2000.
47. Binley, J. M. *et al.* Enhancing the proteolytic maturation of human immunodeficiency virus
type 1 Envelope glycoproteins. *J. Virol.* **76**, 2606–2616 (2002).
48. Ringe, R. P. *et al.* Improving the Expression and Purification of Soluble, Recombinant Native-
Like HIV-1 Envelope Glycoprotein Trimers by Targeted Sequence Changes. *J. Virol.* **91**,
(2017).
49. The Design and Immunogenicity of an HIV-1 Clade C Pediatric Envelope Glycoprotein
Stabilized by Multiple Platforms. https://www.mdpi.com/2076-393X/13/2/110.
50. Guenaga, J. *et al.* Glycine Substitution at Helix-to-Coil Transitions Facilitates the Structural
Determination of a Stabilized Subtype C HIV Envelope Glycoprotein. *Immunity* **46**, 792-
803.e3 (2017).
51. Rodrigues, M. Q., Alves, P. M. & Roldão, A. Functionalizing Ferritin Nanoparticles for Vaccine
Development. *Pharmaceutics* **13**, 1621 (2021).
52. López-Sagaseta, J., Malito, E., Rappuoli, R. & Bottomley, M. J. Self-assembling protein
nanoparticles in the design of vaccines. *Comput. Struct. Biotechnol. J.* **14**, 58–68 (2016).
53. Schommers, P. *et al.* Restriction of HIV-1 Escape by a Highly Broad and Potent Neutralizing
Antibody. *Cell* **180**, 471-489.e22 (2020).

54. Julien, J.-P. *et al.* Broadly Neutralizing Antibody PGT121 Allosterically Modulates CD4 Binding

via Recognition of the HIV-1 gp120 V3 Base and Multiple Surrounding Glycans. *PLoS Pathog.*

**9**, e1003342 (2013).

55. Kumar, S. *et al.* A V $_H$ 1-69 antibody lineage from an infected Chinese donor potently

neutralizes HIV-1 by targeting the V3 glycan supersite. *Sci. Adv.* **6**, eabb1328 (2020).

56. Barnes, C. O. *et al.* Structural characterization of a highly-potent V3-glycan broadly

neutralizing antibody bound to natively-glycosylated HIV-1 Envelope. *Nat. Commun.* **9**, 1251

(2018).

57. Steichen, J. M. *et al.* A generalized HIV vaccine design strategy for priming of broadly

neutralizing antibody responses. *Science* **366**, eaax4380 (2019).

58. Saunders, K. O. *et al.* Targeted selection of HIV-specific antibody mutations by engineering B

cell maturation. *Science* **366**, eaay7199 (2019).

59. Simonich, C. A. *et al.* Kappa chain maturation helps drive rapid development of an infant HIV-

1 broadly neutralizing antibody lineage. *Nat. Commun.* **10**, 2190 (2019).

60. Guenaga, J. *et al.* Glycine Substitution at Helix-to-Coil Transitions Facilitates the Structural

Determination of a Stabilized Subtype C HIV Envelope Glycoprotein. *Immunity* **46**, 792-

803.e3 (2017).

61. Pancera, M. *et al.* Structure and immune recognition of trimeric pre-fusion HIV-1 Env. *Nature*

**514**, 455–461 (2014).

62. Bianchi, M. *et al.* Electron-Microscopy-Based Epitope Mapping Defines Specificities of

Polyclonal Antibodies Elicited during HIV-1 BG505 Envelope Trimer Immunization.

*Immunity* **49**, 288-300.e8 (2018).

63. Klasse, P. J. *et al.* Epitopes for neutralizing antibodies induced by HIV-1 Envelope

glycoprotein BG505 SOSIP trimers in rabbits and macaques. *PLOS Pathog.* **14**, e1006913

(2018).

64. McCoy, L. E. *et al.* Holes in the Glycan Shield of the Native HIV Envelope Are a Target of

Trimer-Elicited Neutralizing Antibodies. *Cell Rep.* **16**, 2327–2338 (2016).

65. Behrens, A.-J. *et al.* Composition and Antigenic Effects of Individual Glycan Sites of a Trimeric

HIV-1 Envelope Glycoprotein. *Cell Rep.* **14**, 2695–2706 (2016).

66. Brouwer, P. J. M. *et al.* Immunofocusing and enhancing autologous Tier-2 HIV-1

neutralization by displaying Env trimers on two-component protein nanoparticles. *Npj*

*Vaccines* **6**, 1–14 (2021).

67. Ozorowski, G. *et al.* Effects of Adjuvants on HIV-1 Envelope Glycoprotein SOSIP Trimers *In Vitro*. *J. Virol.* **92**, (2018).

68. Ahmed, S. *et al.* Enhancing anti-viral neutralization response to immunization with HIV-1 Envelope glycoprotein immunogens. *Npj Vaccines* **8**, 1–10 (2023).

69. Sutar, J. *et al.* Geospatial HIV-1 subtype C gp120 sequence diversity and its predicted impact on broadly neutralizing antibody sensitivity. *PLOS ONE* **16**, e0251969 (2021).

70. Ariën, K. K., Vanham, G. & Arts, E. J. Is HIV-1 evolving to a less virulent form in humans? *Nat. Rev. Microbiol.* **5**, 141–151 (2007).

71. Taylor, B. S., Sobieszczyk, M. E., McCutchan, F. E. & Hammer, S. M. The challenge of HIV-1 subtype diversity. *N. Engl. J. Med.* **358**, 1590–1602 (2008).

72. Huang, A. *et al.* Global Comparison of Drug Resistance Mutations After First-Line Antiretroviral Therapy Across Human Immunodeficiency Virus-1 Subtypes. *Open Forum Infect. Dis.* **3**, ofv158 (2016).

73. Julien, J.-P. *et al.* Design and structure of two HIV-1 clade C SOSIP.664 trimers that increase the arsenal of native-like Env immunogens. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 11947–11952 (2015).

74. Kumar, S. *et al.* Neutralizing Antibodies Induced by First-Generation gp41-Stabilized HIV-1 Envelope Trimers and Nanoparticles. *mBio* **12**, 10.1128/mbio.00429-21 (2021).

75. Borst, A. J. *et al.* Germline VRC01 antibody recognition of a modified clade C HIV-1 Envelope trimer and a glycosylated HIV-1 gp120 core. *eLife* **7**, e37688 (2018).

76. Ringe, R. P. *et al.* Glycan heterogeneity as a cause of the persistent fraction in HIV-1 neutralization. *PLOS Pathog.* **19**, e1011601 (2023).

77. Xu, D. *et al.* Design of universal Ebola virus vaccine candidates via immunofocusing. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2316960121.

78. Powell, A. E. *et al.* Multimerization of Ebola GPΔmucin on protein nanoparticle vaccines has minimal effect on elicitation of neutralizing antibodies. *Front. Immunol.* **13**, 942897 (2022).

79. Casazza, J. P. *et al.* Phase 1 dose-escalation trial evaluating a group 2 influenza hemagglutinin stabilized stem nanoparticle vaccine. *Npj Vaccines* **9**, 1–10 (2024).

80. Tang, P. *et al.* A ferritin nanoparticle vaccine based on the hemagglutinin extracellular domain of swine influenza A (H1N1) virus elicits protective immune responses in mice and pigs. *Front. Immunol.* **15**, (2024).

81. Stephens, L. M. & Varga, S. M. Nanoparticle vaccines against respiratory syncytial virus. *Future Virol.* **15**, 763–778.

82. Swanson, K. A. *et al.* A respiratory syncytial virus (RSV) F protein nanoparticle vaccine
focuses antibody responses to a conserved neutralization domain. *Sci. Immunol.* **5**, eaba6466
(2020).

83. Joyce, M. G. *et al.* SARS-CoV-2 ferritin nanoparticle vaccines elicit broad SARS coronavirus
immunogenicity. *bioRxiv* 2021.05.09.443331 (2021) doi:10.1101/2021.05.09.443331.

84. Weidenbacher, P. A.-B. *et al.* A ferritin-based COVID-19 nanoparticle vaccine that elicits
robust, durable, broad-spectrum neutralizing antisera in non-human primates. *Nat. Commun.*
**14**, 2149 (2023).

85. Yu, J. *et al.* Ad26.COV2.S and SARS-CoV-2 spike protein ferritin nanoparticle vaccine protect
against SARS-CoV-2 Omicron BA.5 challenge in macaques. *Cell Rep. Med.* **4**, 101018 (2023).

86. Johnston, S. C. *et al.* A SARS-CoV-2 Spike Ferritin Nanoparticle Vaccine Is Protective and
Promotes a Strong Immunological Response in the Cynomolgus Macaque Coronavirus
Disease 2019 (COVID-19) Model. *Vaccines* **10**, 717 (2022).

87. Guenaga, J. *et al.* Structure-Guided Redesign Increases the Propensity of HIV Env To Generate
Highly Stable Soluble Trimers. *J. Virol.* **90**, 2806–2817 (2016).

88. Rappuoli, R., Bottomley, M., D'Oro, U., Finco, O. & Gregorio, E. Reverse vaccinology 2.0:
Human immunology instructs vaccine antigen design. *J. Exp. Med.* **213**, jem.20151960 (2016).

89. Klasse, P. J., Ozorowski, G., Sanders, R. W. & Moore, J. P. Env Exceptionalism: Why Are HIV-1
Env Glycoproteins Atypical Immunogens? *Cell Host Microbe* **27**, 507–518 (2020).

90. Seabright, G. E., Doores, K. J., Burton, D. R. & Crispin, M. Protein and Glycan Mimicry in HIV
Vaccine Design. *J. Mol. Biol.* **431**, 2223–2247 (2019).

91. Veneziano, R. *et al.* Role of nanoscale antigen organization on B-cell activation probed using
DNA origami. *Nat. Nanotechnol.* **15**, 716–723 (2020).

92. ACTS, RULES AND GUIDELINES: Committee for the Purpose of Control And Supervision of
Experiments on Animals.
https://ccsea.gov.in/Content/54_1_ACTS,RULESANDGUIDELINES.aspx.

93. Sok, D. *et al.* Recombinant HIV Envelope trimer selects for quaternary-dependent antibodies
targeting the trimer apex. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 17624–17629 (2014).

94. del Moral-Sánchez, I. *et al.* Triple tandem trimer immunogens for HIV-1 and influenza nucleic
acid-based vaccines. *Npj Vaccines* **9**, 1–18 (2024).

95. Kumar, S. *et al.* Recognition determinants of improved HIV-1 neutralization by a heavy chain
matured pediatric antibody. *iScience* **26**, 107579 (2023).

96. Tang, G. *et al.* EMAN2: An extensible image processing suite for electron microscopy. *J. Struct.*
*Biol.* **157**, 38–46 (2007).

97. Kumar, A., P., S., Gulati, S. & Dutta, S. User-friendly, High-throughput, and Fully Automated
Data Acquisition Software for Single-particle Cryo-electron Microscopy. *J. Vis. Exp.* 62832
(2021) doi:10.3791/62832.

98. Scheres, S. H. W. RELION: Implementation of a Bayesian approach to cryo-EM structure
determination. *J. Struct. Biol.* **180**, 519–530 (2012).

99. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

100.   Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat. Methods* **14**, 331–332 (2017).

101.   Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J. Struct. Biol.* **192**, 216–221 (2015).

102.   Grant, T., Rohou, A. & Grigorieff, N. cisTEM, user-friendly software for single-particle
image processing. *eLife* **7**, e35383 (2018).

103.   Mishra, N. *et al.* Broadly neutralizing plasma antibodies effective against autologous
circulating viruses in infants with multivariant HIV-1 infection. *Nat. Commun.* **11**, 4409
(2020).

104.   Montefiori, D. C. Measuring HIV Neutralization in a Luciferase Reporter Gene Assay. in *HIV*
*Protocols* (eds. Prasad, V. R. & Kalpana, G. V.) 395–405 (Humana Press, Totowa, NJ, 2009).
doi:10.1007/978-1-59745-170-3_26.

**FIGURE LEGENDS:**

**Figure 1: Design & biochemical characterization of 330 gp140 Env trimers and ferritin**
**nanoparticles.** Linear representation of three 330 Envelope variants **(A)** Design of SOSIP with
the SOS-bond (A605-T605C), IP (I559P) stabilizing mutation along with TD8 mutations, a
intraprotomer disulfide bond (I201C-A433C), an improved furin cleavage (R6) with CD5 Signal
Peptide (SP) and Twin Strep-Tag (tst) tag **(B)** Design of NFL covalently linking gp120 to gp41
with flexible glycine serine (GGGGSGGGGS) linker **(C)** Design of H. pylori ferritin nanoparticle
linked glycine serine linker to SOSIP construct **(D)** A surface structure highlighting the TD8
stabilizing mutations (right) in the 330 Envelope construct. Structure was illustrated using
PyMOL and based on the BG505 SOSIP.664 crystal structure (PDB ID: 4ZMJ) **(E)** Size-exclusion

chromatography (SEC) profiles of AIIMS 330 Env trimers (SOSIP and NFL) and Ferritin

nanoparticles (NPs), eluted between 8–12 mL on a Superdex 200 Increase 10/300 GL column **(F)**

Negative Stain-EM 2-D class averages of NFL & SOSIP gp140 Env trimers and ferritin

nanoparticles **(G)** 3D reconstruction of SOSIP gp140 Env trimers and multivalent ferritin

nanoparticles using chimera X **(H)** Melting curve around 74°C of AIIMS 330 Envelope

**Figure 2: Antigenicity of 330 gp140 Env trimers and ferritin nanoparticles to HIV-1 bNAbs**

**and non-NAbs.** Representative AIIMS 330 Env binding curves with conformational specific

bNAbs (PGT145, CAP256.25, PGDM1400, PGT151), CD4 specific bNAbs (VRC01, N6), V3

Supersite (10.1074, AIIMS P01) along with non-nAbs (17b+sCD4, F105, A32, 48d) assessed by

ELISA.

**Figure 3: Octet sensograms kinetics of 330 Env to HIV-1 bNAbs and non-NAbs. (B)** Octet

Sensograms showing response shift (nm) over time (sec) of AIIMS 330 Env trimers and particles

with varied concentrations (500 nM, 166.6 nM, 55.5 nM, 18.5 nM, 6.1 nM) against the quaternary

specific bNAbs PGT145, PGD1400, cleavage dependent bNAb PGT151 and the non-nAbs F105 &

17b+sCD4.

**Figure 4: Structural and glycan homology of 330 SOSIP with BG505 & 16055 trimers (A-B)**

Atomic model comparison of 330 SOSIP Env trimer with BG505 SOSIP (PDB: 4TVP) (RMSD

gp120: 0.685 $A^0$, gp41: 0.478 $A^0$) **(C-D)** 16055 NFL (PDB: 5UM8) (RMSD gp120: 0.798 $A^0$, gp41:

0.739 $A^0$) SOSIP trimer. Colour coding of SOSIP trimers corresponding to segmented EM

densities are: dodger blue, AIIMS 330 SOSIP, deep pink, BG505 SOSIP and 16055. Within each

protomer, gp120 regions are highlighted in blue and pink, while gp41 regions are shown in

yellow and green **(E)** Presence of glycan hole in 330 Env at position 130, 295, 339 in HXB2

reference coordinates showed in top, front & side view **(F)** Figure depicting the total protein

accessible area across cross-clade Envelopes **(G)** Presence of total 28 PNGS in AIIMS 330 &

BG505 Envelope. An N indicates the presence of a PNGS, if NXT/S motif in a.a sequence whereas

X should not be proline. The glycan structure and alignment created by N-Glycosite tool from Los

Alamos HIV-1 sequence database.

**Figure 5: Cryo-EM structure of 330 SOSIP Env trimer in complex with autologous bNAb**

**AIIMS 44m (A)** Top and side views of Cryo-EM reconstructed map of AIIMS 330 SOSIP Env

trimer in complex with 44m bNAb. Colour coding corresponding to segmented EM densities are:

magenta, 44m bNAb; dodger blue, gp120; gold, gp41 **(B)** The corresponding atomic model fitted

in the EM map of AIIMS 330 SOSIP trimer in complex with 44m bNAb. The model illustrates that three 44m bNAbs bind to each gp120 monomeric subunit, facilitating effective neutralization **(C)** Location of CDRH and FR3 regions are shown here in different colours (CDRH1- cyan, CDRH2- deep green, CDRH3- light green, FR3- light brown) **(D)** Key interacting residues AIIMS 44m bNAb - H31, Y52, Y53, and R109 highlighted in cyan colour, and corresponding interacting residues of gp120 loops - T144, D325, N332, and T413 highlighted in yellow colour **(E)** The N-glycans N146, N301, and N332 located at the junction of AIIMS 44m bNAb and AIIMS 330 gp120 protomers **(F)** The D325 residue from the Env V3 loop oriented toward a hydrophobic cleft in the mAb CDRH2 region.

**Figure 6: Immunogenicity & Neutralizing potential of 330 SOSIP, NFL Env trimers & Ferritin NP's in NZW rabbits. (A)** Rabbit immunization Schedule. Four groups of rabbits were immunized at week 0 (prime), 4 & 20 (boost), gp140 SOSIP and NFL Env immunogens (G1 and G2) were administered at a dose of 20 μg of trimers, or an equivalent molar amount SOSIP-ferritin nanoparticles (25 μg) (G3) formulated with the AddaVax adjuvant (1:1 v/v). Sera was obtained at indicated intervals of week 4, 6, 8, 12, 16, 20 & 22 (red arrows). **(B)** Binding endpoint titers (OD at A450) over time against autologous virus (330.16.E6) Dots and error bars represent the median binding titers and standard deviations. The assay cut-off is marked with a dotted line **(C)** Midpoint neutralization titers ($ID_{50}$) for week 22 sera of the immunized rabbits against autologous (330.16.E6) and closely related (329.14.B1), and heterologous clade C (MW965.26 and 25925), Tier 1 clade B (SF162), and Tier 2 clade A (BG505.T332N) pseudoviruses. Median ID50 titers NFL, SOSIP & Ferritin NP group was 375, 830 & 1160. Statistical analysis was done using an unpaired two-tailed Student's t-test and the assay cut-off is marked with a dotted line.

A

330 SOSIP

CD5 SP | C1 | V1 | V2 | C2 | V3 | C3 | V4 | C4 | V5 | C5 | | HR1 | HR2 | tst

R6          .664

E47D     EI06T          I165L     I201C          R315Q          A433C     K429R     A501C     I559P     T605C
T49E                                                        K432Q     K500R
V65K

B

330 NFL

CD5 SP | C1 | V1 | V2 | C2 | V3 | C3 | V4 | C4 | V5 | C5 | GS Linker | HR1 | HR2 | tst

.664

E47D     EI06T          I165L     I201C          R315Q          A433C     K429R     K500R     I559P
T49E                                                        K432Q
V65K

C

330 SOSIP Ferritin NP

CD5 SP | C1 | V1 | V2 | C2 | V3 | C3 | V4 | C4 | V5 | C5 | | HR1 | HR2 | GS Linker | Ferritin

R6          .664

E47D     EI06T          I165L     I201C          R315Q          A433C     K429R     A501C     I559P     T605C
T49E                                                        K432Q     K500R
V65K

D

I165L
I201C
R315Q          K432Q
A433C          K429R
V65K
E106T
T49E
E47D
K500R

E

SEC-Superdex 200 10/300 GL

nanoparticles    aggregates    trimers    dimers/monomers

330 SOSIP
330 NFL
330 Ferritin NPs

F

NS- EM 2D Class Averages

NFL Env Trimers          SOSIP Env Trimers          Ferritin Nanoparticles

G

3D Reconstruction

Soluble gp140 Trimers          Ferritin Nanoparticles

H

Tm 74° C

330 SOSIP
Control

**A**

**B**

330 SOSIP
BG505 SOSIP

RMSD    gp120 : 0.685 A°
        gp41 : 0.478 A°

**C**

**D**

330 SOSIP
16055 NFL

RMSD    gp120 : 0.798 A°
        gp41 : 0.739 A°

**E**

Front view                Top View                Side view

**F**

**G**