**Title: Persistent effects of dietary selection and inbreeding on microbiome composition**

**and longevity in Drosophila**

ⱳPeyton K. Warren, ORCID 0009-0003-6100-1381

ⱳGarrison Miller,

‡Gaurav S. Kandlikar, ORCID 0000-0003-3043-6780

ⱳEnoch Ng'oma ORCID 0000-0002-6741-7922

ⱳDivision of Biological Sciences, University of Missouri-Columbia, MO, USA

‡Department of Biological Sciences, Louisiana State University, LA

18 Peyton K. Warren and Garrison Miller contributed equally to this work.

*Correspondence: Enoch Ng'oma

ngomae@missouri.edu

**Abstract**

Background

Dietary environments can shape host-microbe co-evolution by imposing selection pressures that

favor host and microbial genotypes with enhanced fitness under specific nutritional conditions.

However, the long-term evolutionary dynamics of these interactions after selection pressures

are relaxed remain poorly understood. Here, we investigated the microbiome of *Drosophila*

*melanogaster* populations that underwent long-term experimental evolution under three

temporally variable nutritional regimes (constant high sugar, progressively decreasing protein,

and fluctuating protein), and compared them to unselected controls. Following inbreeding of

evolved populations, we examined how persistent evolutionary effects interact with aging to

shape host-microbe relationships and survival.

Results

All selected regimes exhibited reduced survival relative to controls, indicating lasting

physiological costs associated with historical selection and inbreeding. Survival differed among

regimes: the deteriorating-protein regime was closest to controls, the fluctuating-protein regime

was intermediate, and the high-sugar regime showed the shortest lifespan. Survival effects were

sex-specific: relative to control females, those from the fluctuating-protein regime exhibited

reduced early-life survival, whereas females from the deteriorating-protein regime experienced

greater late-life mortality. Microbiome composition varied with both selection regime and age.

Although Firmicutes and Proteobacteria dominated across groups, selected lines showed

increased Firmicutes and reduced Proteobacteria, especially early in life, suggesting early-life

taxonomic restructuring. The decreasing-protein regime maintained more stable microbial

diversity over time, whereas high-sugar and fluctuating-protein diets were associated with

progressive microbiome instability with age. Core *Acetobacter* species (*A. aceti*, *A.*

*oryzifermentans*) declined in abundance in selected flies, indicating persistent disruption of

microbiome integrity. A random forest model predicted fly age from microbiome composition

with 78.8% accuracy, reinforcing links between microbial dynamics and host aging.

Conclusion

Historical dietary selection and inbreeding can leave lasting signatures on survival and

microbiome composition in *Drosophila*. Protein restriction promoted late-life longevity and

microbial stability, whereas high-sugar and fluctuating diets were associated with early-life

effects followed by later-life shifts in dominant taxa. Together, these findings illustrate how

nutritional history and its microbial legacies influence lifespan and aging through persistent host-

microbe interactions.

**Keywords:** experimental evolution, nutritional history, longevity, microbiome legacies, dietary

stress, persistent effects.

**Introduction**

Microorganisms are deeply integrated into host biology across taxonomic and functional scales, playing critical roles in organismal performance and fitness (Ley et al. 2008; Muegge et al. 2011; Gould et al. 2018). Rapidly changing environmental conditions increase the pressure for hosts to cope with novel conditions faster than their genomes can evolve. Microbial genomes, which evolve more rapidly than host genomes, represent an important source of ecological and functional flexibility, enabling hosts to respond to environmental stressors through shifts in symbiotic composition and activity. Microbiotas, as part of the functional hologenome, can mediate developmental timing, immune defense, energy allocation, and other traits central to host survival and reproduction [4, 5]. In many species, the microbiome is a key modulator of host fitness [6], development [7, 8], reproduction [9, 10], stress tolerance [11], pathogenic defense [12, 13], survival, and aging [13–15]. Disruptions to microbial composition can increase gut permeability and systemic inflammation, accelerating physiological decline [16], whereas beneficial microbes can attenuate immune overactivation and extend lifespan [14]. These findings position inflammation as a central axis through which microbiotas mediate life-history trade-offs, particularly under dietary stress [14, 16], reviewed in [17, 18].

Diet is a dominant ecological determinant of host-associated microbiomes. In *Drosophila melanogaster*, while variation in microbiome composition has been linked to host allele frequency shifts in genes associated with local adaptation [4], microbial composition is often more strongly influenced by the nutritional environment than host genotype [19, 20]. For example, *Lactobacillus plantarum* enhances larval growth on poor diets by promoting insulin signaling [7], while acetic acid bacteria regulate sugar metabolism and immune function [21]. Nutrient availability thus shapes both microbiota composition and downstream host traits including immune investment, somatic maintenance, and fecundity. Many studies have shown that reduced protein-to-carbohydrate (P:C) ratios extend lifespan at the cost of reproductive

output, whereas protein-rich diets promote early-life fecundity but reduce survival [22–24].

These physiological responses are mediated through nutrient-sensing pathways such as IIS and

mTOR [25, 26], which are also modulated by gut microbes [27]. Given this complex interplay,

sustained dietary stress is expected to shape host physiology and microbiome composition in

concert.

Intrinsic host factors such as age also interact with microbial dynamics. In *D. melanogaster*,

aging is associated with shifts in gut microbiota composition, including reduced diversity and

expansion of inflammatory taxa [14, 28]. These changes are not simply passive consequences

of aging but actively contribute to age-related physiological decline, suggesting that the

microbiome is both a target and mediator of aging processes.

Here, we examine a set of experimentally evolved *D. melanogaster* populations that were

maintained for 65 generations under distinct dietary selection regimes and subsequently inbred

for >30 generations due to COVID pandemic-related disruptions. This design enables us to

investigate whether long-term dietary selection has left lasting genomic or microbiome

signatures that persist after extended inbreeding and rearing under a common diet. Selection

treatments included: 1) deteriorating availability (DA), representing high-to-low protein provision

within each generation; 2) fluctuating availability (FA), reflecting predictable high-low-high

protein cycles; and 3) constant high sugar (CH), modeling persistent metabolic excess. A

control group (C) was maintained without dietary stress.

Our primary goal was to determine whether historical dietary selection is reflected in

contemporary microbiome composition and host survival patterns. We therefore refer to any

observed differences as *legacies of dietary history* rather than direct evidence of ongoing

adaptation. We hypothesized that hosts derived from contrasting nutritional histories would

retain distinct microbiome compositions reflecting their ancestral environments, potentially

influencing lifespan through pathways linked to inflammation and metabolic homeostasis.

Specifically, we predicted that the DA regime would show signatures consistent with investment

in somatic maintenance and longevity, potentially mediated by microbes that mitigate

inflammatory processes. In the FA regime, evolved under predictable cycles of abundance and

scarcity, we anticipated intermediate survival outcomes and dynamically shifting microbial

compositions reflecting metabolic flexibility. Chronic high-sugar exposure (CH regime) was

expected to produce signatures of metabolic stress and reduced microbial diversity. Finally, we

anticipated that age-related microbiota succession would interact with these dietary legacies,

producing distinct temporal patterns of microbial resilience or instability.

Collectively, these predictions provide a framework for testing how long-term nutritional history,

filtered through both host and microbial evolution, can leave enduring effects on host–

microbiome interactions and life-history dynamics, even after prolonged inbreeding and

common-garden rearing.

**Materials and methods**

Experimental evolution

As part of a larger study, we experimentally evolved a population of *D. melanogaster* under

several stressful nutritional conditions. We generated the initial population by mass-

intercrossing 835 recombinant inbred lines from the B population (pB) of the *Drosophila*

Synthetic Population Resource, DSPR [29, 30] for five generations under common garden

conditions (Fig. 1a) [31]. Several Drosophila community panels in North America, including

*Drosophila* Genetic Reference Panel (DGRP) and the *D. melanogaster* Reference Genome,

exhibit mosaic genomic ancestry (~20% African, ~80% European) shaped by natural selection

against incompatible allele combinations [32]. This can also be expected in DSPR populations,

highlighting the need for a genetically diverse population in experimental evolution, as such

expanded genotypic diversity would provide a broad substrate for selection facilitating the study

of complex traits and adaptation. The properties of the DSPR are described elsewhere [29, 30].

From this genetically diverse base population, 36 lines were independently subjected to one of

three dietary selection regimes for 65 generations with each regime replicated 12 times in large

population cages (N > 4,000 flies; Fig. 1a). One population (N > 4,000) was maintained without

selection exposure for the duration of selection. Selection in these populations acted primarily

on reproductive success and early-life survival under dietary stress, rather than directly on

lifespan. Consequently, differences in longevity observed in the inbred lines likely reflect

correlated responses to selection on nutrient-dependent fitness rather than direct selection for

extended lifespan.

**Figure 1:** Study design. **a.** A total of 835 recombinant inbred lines of the DSPR rared on

standard maintenance diet were interbred into a genetically diverse population and subjected to

experimental evolution in four diet treatments for 65 generations. Selection was terminated after

65 generations, and flies in each regime inbred in vial replicates, here denoted, *selection inbred*

*lines*, or SILs and maintained on common NutriFly diet. Control flies were maintained on

standard diet throughout selection in one cage at N > 4,000 and also moved to NutriFly at

selection termination. Six SILs per regime were each expanded into 6 replicates, alongside

pseudo replicate vials of C flies. Microbiome composition was measured at 15, 35 and 60 days

post-eclosion in 3 vial replicates. Survival was measured in a separate set of 3 vials. **b.**

Selection regimes: 1) fluctuating (FA), 2) deteriorating (DA) and 3) constant high (CH) nutrient

availability were set up in 12 replicate cages (N > 4000 flies per cage) and fed sequences of low

protein (LP) and control (C) diet alternations or constant high sugar (HS) diet over a 21-day

cycle post-oviposition (ovp) per generation (see methods). Note: DA and FA received temporal

alternations of contro and low protein levels; CH received an oversupply of sugar; C was

retained on standard diet for the DSPR (no deliberate selection).

*Nutritional composition of diets constituting selection regimes.* Based on previously published

formulations [25, 31, 33], we used three diets that systematically varied in the ratio of protein

(yeast) to carbohydrate (sucrose) to impose distinct nutritional selection pressures. Each diet

was prepared per liter of water with the following composition: Control diet - 10 g agar, 50 g

sucrose, 200 g yeast, and 2.7 g tegosept (dissolved in 11 mL ethanol); Low-protein diet - 10 g

agar, 50 g sucrose, 100 g yeast, and 2.7 g tegosept; High-sugar diet - 10 g agar, 342 g sucrose,

200 g yeast, and 2.7 g tegosept. Yeast provided the principal protein source, and sucrose

supplied carbohydrates. The control diet yielded an intermediate protein: carbohydrate ratio

(P:C ≈ 1 : 0.5) typical of standard laboratory diets that support optimal growth and fecundity.

The low-protein diet reduced yeast content by half while maintaining sucrose levels, generating

a protein-limited condition (P:C ≈ 1 : 1). The high-sugar diet increased sucrose concentration

roughly sevenfold, creating a carbohydrate-rich environment (P:C ≈ 1 : 3). Percent

macronutrient content by energy was approximately 45 - 53% protein and 47–52% carbohydrate

for the control, 36–41% protein and 59–64% carbohydrate for the low protein diet, and 17–19%

protein and 81–83% carbohydrate for high sugar diet.

*Maintenance of the control population.* Prior to the onset of experimental evolution, the

recombinant inbred lines used to generate the base population were maintained on a standard

cornmeal–dextrose maintenance diet (40.8 g cornmeal, 21.6 g yeast, 86.3 g dextrose, 6.25 g

agar per liter of water, with 1.8 g tegosept). This base diet contained lower protein and

carbohydrate levels than any of the experimental diets, ensuring that the defined sugar–protein

diets represented clear nutritional shifts.

To preserve an ancestral baseline for comparison, the unselected control population was

maintained throughout the experiment on the cornmeal–dextrose maintenance diet described

above - the same diet on which the DSPR founder population was reared prior to the start of

selection. We note that this "maintenance" diet differs from the "control (C)" diet used as one of

the experimental selection treatments (see above). Using the ancestral diet for the unselected

population ensured that these flies represented the pre-selection reference state, avoiding

inadvertent selection under any of the experimental diets. The control diet was considered

optimal for growth and reproduction and provided a reference for evaluating nutritionally

stressful regimes (low protein or high sugar) that impose either limitation or oversupply of key

nutrients (previously characterized in Ng'oma et al. [34]).

*Construction of selection treatment regimes*. From these diets, we designed three selection

conditions (Fig. 1b) by alternating the sequence in which the diets were presented to flies in

each selection generation. The Fluctuating Availability (FA) treatment alternated nutrient

provision as follows: flies received a control until day 14 post-oviposition (po), low protein diet

from day 14 - 18 po, and control diet again until day 21 po when eggs were collected to seed

the next generation. Flies in the Deteriorating Availability treatment (DA) received a standard

diet until day 18 po, followed by a low protein diet until day 21 po, at which eggs were collected.

Files in the Constant High sugar treatment (CH) received the high sugar diet throughout, until

day 21 po. Alongside selection cages, we maintained one population (N > 4000) that was not

exposed to experimental selection and likely remained outbred at the time of this assay, with no

observed population bottlenecks and assuming weak genetic drift. Because eggs were collected

from survivors at day 21 po each generation, selection acted primarily through differences in

fecundity and viability during early adulthood under each dietary regime. Longevity beyond this

window was therefore not a direct target of selection but could evolve indirectly as a correlated

response to nutrient-dependent reproductive success and stress tolerance. These dietary

regimes therefore varied in both macronutrient composition and temporal availability during

selection. This may invoke distinct evolutionary responses in the reproduction–survival trade-off.

We interpret results with this fact in mind.

*Selection inbred lines (SILs).* Experimental evolution was performed for 65 generations, after

which populations were inbred in replicated vials for each of 36 selection populations at the

onset of the Covid pandemic for 38 generations (~19 months). At the end of experimental

evolution, all flies were of similar age and sampled from the actively reproducing cohort,

ensuring that alleles influencing fecundity and early-life survival were those most consistently

transmitted to the next generation. We refer to inbred lines derived from experimental selection

as Selection Inbred Lines (SILs, Fig. 1a). SILs were maintained in common garden conditions

on a molasses-based Nutri-Fly® diet (Genesee Scientific) until assaying, and the unselected

group was held at high population in a cage until assay.

*Sample collection.* For this study, we selected six random SILs from each selection regime (FA,

DA, and CH) along with 12 pseudo replicated vials of the non-selected population. Each

selected SIL was expanded over three generations, and three vials each were assigned to

microbiome and lifespan estimation assays. Each assay vial was seeded with 24 females and 6

males to reduce male mating pressure, which is known to reduce female lifespan in *Drosophila*

[34, 35]. This set up minimizes mating-induced costs and allows clearer inference on intrinsic

female lifespan differences among selection treatments. Our design therefore yields a reduced

male sample in lifespan analysis which we consider in the interpretation of results (i.e., 1,670

deaths, 176 censored events for females; 572 deaths, 9 censored events for males). Death

events were counted three times per week (Monday, Wednesday, Friday) at which time also a

fresh food vial was provided. For the microbiome portion, we assayed only females. Based on

our prior study indicating key variability in lifespan and fecundity responses to diet across life

phases [36], we collected whole flies at three time points (15, 35, and 60 days po, Fig. 1). At

each time point, four females were flash-frozen and stored at -80°C for analysis.

*Diet sources.* Diets were prepared with inactivated SAFPro Relax + YF 73050 yeast (Lesaffre

Yeast Corp., Milwaukee, USA), stored at 4°C, and used within 14 days. Evolved lines were held

on a common Nutri-fly molasses formulation (Genesee Scientific) based on a Bloomington

Drosophila Stock Center Nutri-fly molasses protocol (12.4 g/L yeast, 61.3 g/L yellow cornmeal,

and 75.2 ml/L molasses). This change was necessitated by shortages in supplies during and

after the pandemic. Across all assays, flies were maintained in a temperature-controlled

chamber at 23°C under continuous light, mirroring conditions under which the DSPR was

originally generated and maintained [29]. Flies were supplied with fresh food three times per

week (Monday, Wednesday, Friday).

16S rRNA amplicon sequencing

Whole-fly 16S rRNA sequencing without surface sterilization captures both internal and surface-

associated microbes and thus represents the total host-associated microbiome. This approach

aligns with our goal of characterizing microbiome responses to long-term dietary selection at the

whole-organism level. All samples were reared on a common diet medium prepared in bulk and

distributed across all selection lines, ensuring a shared environmental microbial background.

Therefore, observed differences across selection treatments likely reflect evolved host-

microbiome associations rather than environmental contamination. DNA was extracted from

pools of two to four flies using the Qiagen DNeasy Blood & Tissue Kit (Cat. No 69504). Samples

were manually homogenized with a glass pestle in ATL buffer, incubated for 4 hours, and

processed following the kit manufacturer's protocol. DNA yield and quality were assessed using

a nanodrop spectrophotometer (ND-1000) and Qubit 4 Fluorometer with the Qubit 1X dsDNA

HS Assay Kit (Cat. No Q33231). Samples were normalized as needed with elution buffer and

sequenced on a MiSeq instrument (PE250-v2 Nano flow cell, 25 cycles) targeting the V4 region

of the bacterial 16S rRNA gene with primers 515F (5'-GTGCCAGCMGCCGCGGTAA-3') and

806R (5'-GGACTACHVGGGTWTCTAAT-3') [37, 38]. Sequencing was performed at the

University of Missouri Genomics Technology Core.

Data Analysis

*Lifespan*. We first explored and compared survival patterns among selection regimes in a total

of 2,427 individuals (2,242 deaths, 185 censored), using survival models that included sex

either as a covariate or in sex-stratified analyses. Kaplan-Meier survival curves indicated

marked sex differences in survival probability (Additional File 2 Table S1) and changes in the

relative survival rankings of selection regimes over time. Because each selection regime (CH,

DA, FA) consisted of multiple unique selection inbred lines (SILs) replicated across vials, our

design required modeling fixed effects of regime, sex, and their interaction, while accounting for

random effects of SIL and vial.

To evaluate the effects of dietary regime and sex on survival, we fitted mixed-effects Cox

proportional hazards models with a random intercept for SIL nested within vial. We compared

an additive model (`event ~ regime + sex + (1 | SIL/vial)`) and an interaction model

(`event ~ regime * sex + (1 | SIL/vial)`) using ΔAIC as the comparison criterion.

Proportional hazards assumptions were tested using Schoenfeld residuals. All fixed effects

(regime, sex, and their interaction) significantly violated this assumption ($\chi^2$ test, $p < 0.01$),

indicating time-varying hazard ratios. A time-transformed Cox model (`regime + tt(sex)`,

where `tt(x) = x * log(t)`) improved fit (ΔAIC = 15.6), confirming time-dependent effects of

sex.

Given these violations and the hierarchical structure of our data, we proceeded with a discrete-

time survival framework. We implemented a generalized linear mixed model (GLMM) with a

complementary log–log (cloglog) link using **lme4**, specifying event ~ regime * sex + (1 | SIL).

This model structure best fit the data (ΔAIC = 17, $p = 2.7 \times 10^{-5}$), while additional vial-level

random effects did not improve fit (ΔAIC = +2).

To confirm robustness, we re-estimated the same model using a fully Bayesian hierarchical

approach implemented in **brms**, specifying identical fixed and random effects and a cloglog link.

Weakly informative priors were applied (normal(0, 2) for fixed effects; Student-t(3, 0, 2.5) for

random effects). Results from both approaches were concordant, and the Bayesian model was

used for inference and visualization. Full model specifications and diagnostics are provided in

Additional File 1.

Our recent study testing effects of larval nutrition on adult phenotypes in this population [36]

showed that patterns of survival can vary significantly across shorter periods of time in adult

lifespan. To understand evolved patterns of survival in each selection treatment, we examined

percentile survival differences in ten-day intervals in which each of our sampling time points for

the microbiome portion (described next) formed the median: 10-20, 30-40, and 55-65 days, post

eclosion. We tested for significant differences between treatments in each time interval I1, I2

and I3 using either analysis of variance (Anova) or Kruskall-Wallis tests followed by an

appropriate post-hoc test (Turkey's HSD or Dunn's Test).

*Microbiome analysis*: We processed Illumina reads using the R library DADA2 pipeline [39],

leveraging its use of Amplicon Sequence Variants (ASV), an alternative to Operational

Taxonomic Units (OTUs) that enables the detection of single-nucleotide differences in amplicon

sequencing and avoids the limitations of traditional OTU clustering by using exact biological

sequences. Reads were quality-filtered and trimmed at positions 225 (forward) and 200

(reverse), and chimeras were removed. Taxonomy was assigned using the Silva database

(v138.1). Samples with fewer than 200 reads after sequence merging were excluded. We

investigated the proportion of reads identified as *Wolbachia* and found that these constituted

less than 1% reads in the entire data set (Additional File 3, Fig. S7). Prior to analysis, these

reads were removed as common practice [40]. We performed mainstream analysis and

visualization using the R library **phyloseq** [41]. Read counts were normalized using a log-

transformation: `log(1 + x)/sum(log(1 + x))`. This approach combines two commonly used

normalization equations, log(1 + x) and (x)/sum(x), into one step and effectively scales data to

relative abundances (i.e., the proportion of specific microbial groups within samples) while

revealing differences in rare taxa through its log transformation [42–44]. This method was used

instead of rarefication because rarefication is known to unnecessarily discard samples and to

have a higher false positive rate (McMurdie and Holmes 2014).

We assessed microbiome diversity using both alpha and beta diversity metrics. For alpha

diversity, we calculated Shannon and observed species indices from non-normalized sequence

data, as standard practice to preserve true richness and evenness without introducing artifacts

from normalization. Beta diversity was also evaluated on a full data set using principal

coordinate analysis (PCoA) based on UniFrac distances, with phylogenetic trees generated via

the *ape* package [45]. To test for overall differences in microbial community structure across

treatments and time points, we applied permutational multivariate analysis of variance

(PERMANOVA) using the **vegan** R package [46], applying Bonferroni correction to control for

false positives. We further used PERMDISP2, a multivariate analogue of Levene's test [47], to

determine whether observed differences were influenced by dispersion within groups.

Collectively, these analyses provided a robust framework for understanding the drivers of

microbial community composition in our study.

To identify taxa associated with specific selection treatments or age groups, we performed

Indicator Species Analysis using the **indicspecies** R package [48]. Based on the ASV table, we

assessed associations between microbial taxa and either selection treatment or fly age.

Indicator values were calculated using the `multipatt()` function, which evaluates both

specificity and fidelity of taxa to groups, and statistical significance was assessed using 999

permutations.

To assess whether microbiome composition could predict fly age, dietary regime, or their

interaction, we implemented a predictive random forest model adapted from [49]. Using the

**caret** R package, we trained models to classify samples based on time point, treatment, and

their combination, accounting for each unique vial and treatment-age group. Model performance

was evaluated using repeated five-fold cross-validation, repeated five times. To account for

variability in accuracy estimates, we ran 100 model iterations and calculated the mean

predictive accuracy for each classification task. Additionally, we extracted the top 15 most

important taxa contributing to each model, based on feature importance scores from the trained

random forest classifiers.

**Results**

Overall longevity following diet selection

Lifespan performance in the non-selected base population (C) served as a benchmark for

evaluating selection treatments (Fig. 2). Summary survival patterns are presented in (Additional

file 2, Table S1). Overall, a log-rank test comparing survival across selection regimes revealed

highly significant differences in lifespan ($\chi^2$ = 274, df = 3, $p$ < 2e–16). CH (i.e. constant high

sugar) flies showed markedly lower survival (Fig. 2a-c), with far more deaths than expected

under the null hypothesis of equal survival curves (median 27 days vs 42 days in C; Additional

File 1, Table S2). In contrast, DA and FA (i.e. deteriorating- and fluctuating-availability) flies

exhibited higher-than-expected survival (median 41 and 37 days, respectively). The control

group lived the longest, significantly exceeding both CH (p = 5.91e-09) and FA (p = 0.0273)

regimes (Fig. 2a). These patterns are reflected when survival is analyzed within 10-day bins

around microbiome sampling time points, with the CH showing steeper mortality (Fig. 2c).

**Figure 2:** Survival performance: **a.** overall with both sexes together, **b.** median survival in each selection group of SILs, **c.** survival by sex based on bins of 10 days in which the sampling time point (of the microbiome) is the median (only bins containing assayed time points of 15, 35 and 60 days are shown (I1=10-20 days, I2=30-40 days, I3=55-65 days); **d-g**, comparison of sex performance averaged across SILs in each selection treatment.

Sex-specific survival patterns

Sex differences in lifespan were apparent across all selection regimes. A Cox proportional hazards model with a time-varying effect of sex revealed significant differences in survival across regimes ($\chi^2$ = 304.8, *df* = 4, *p* < 2e-16). Flies from the constant high-sugar regime (CH) had the highest risk of death (HR = 2.68, 95% CI: 2.28–3.15, *p* < 2e-16), followed by the fluctuating-availability (FA) regime (HR = 1.49, 95% CI: 1.27–1.75, *p* = 9.9e-07). The deteriorating-availability (DA) regime showed a modest, borderline effect (HR = 1.17, 95% CI: 1.00–1.38, *p* = 0.055). The time-varying term for sex, *tt(Sex_binary)*, was also significant (HR = 1.10, 95% CI: 1.06–1.13, *p* = 2.9e-10), indicating that the sex effect on mortality increased with age (Additional File 3 Fig. S1c). Males thus faced progressively higher mortality risk than

females over time. Together, these results demonstrate that long-term selection under stressful

dietary regimes, particularly constant high sugar, led to pronounced lifespan reductions, with

sex-specific differences in mortality risk that intensified with age.

Although sex differences were clear, their magnitude and direction varied (Fig. 2d-g). In the

control population, females outlived males, with median lifespans of 46 and 40.5 days,

respectively (95% CI for females: 42 - 49; males: 35 - 44). A similar, though smaller, sex gap

was observed under the DA regime (median: females 42, males 39 days). In the FA regime,

survival differences between sexes were minimal, with overlapping confidence intervals and

near-identical medians (females: 37 days, males: 36 days). Interestingly, under the CH regime,

males lived slightly longer than females, with median lifespans of 32 vs 26 days. Generally, CH

and FA treatments favored male survival earlier, reversing later in life with higher female

survival (Fig. 2c, d, f). Conversely, DA and C regimes showed greater female survival through

most of life (Fig. 2c, Fig. 2e,g, Additional file 3 Fig. S1 - S2), suggesting adjustment to dietary

stress through reduced early mortality (DA, FA) or combined early and late-life mortality

reduction (DA, C). These results indicate that selection regime influences not only overall

lifespan but also modulates sex-specific survival patterns, with a marked sex-based lifespan

reversal occurring under constant high sugar (CH). Results further suggest that dietary stress

adaptation might involve distinct survival strategies in this population, with DA and FA regimes

prioritizing early-life resilience and DA further extending late-life survival.

Interactive impacts of sex and diet regime

Relative to C flies, the DA and FA regimes showed reduced early mortality (in I1 and I2, Fig. 2c)

favoring longevity, but diverged later (in I3), with FA experiencing slightly higher late mortality

while DA maintained slower rates. Control flies exhibited the lowest mortality in the first half of

lifespan (I1 and I2). In contrast, CH flies suffered heavy mortality across the entire curve.

Underlying these differences are apparent rank changes in female vs male curves suggesting

strong regime by sex interactions in survival (Fig. 2d-g). To evaluate the effects of selection

regime, sex, and their interaction on mortality risk over time, we fit both Bayesian and

frequentist generalized linear mixed models (GLMMs) using a complementary log-log link.

Estimates from both approaches were highly consistent, showing strong effects of time and the

CH selection regime on increased mortality probability, while the DA and FA regimes had

weaker or non-significant effects (Table 1, Fig. 3, Additional file 3 Fig. S3-S5). Male flies

experienced a modestly elevated risk overall, and these differences were most pronounced in

the CH treatment (Fig. 3). Bayesian posterior direction (PD) values further supported the

robustness of key effects (e.g., time, CH, FA, Male, and CH : Male), with >96% of posterior

samples in the direction of the mean. Full model estimates are provided in Table 1.

Predicted probabilities of death varied by selection regime and sex in both Bayesian and

frequentist GLMMs (Additional File 3 Fig. S3, S4, S6). In C flies, females showed the lowest

mortality probability (~1%), while CH females had the highest (~3–4%), with overlapping

estimates between models. Males consistently exhibited higher predicted risks than females

within most regimes, particularly under DA and FA. Both models revealed minimal sex

differences in CH, where female and male predictions were comparably elevated. These

adjusted predictions support the significant regime and sex effects identified in the full models.

**Table 1.** Comparison of Bayesian and frequentist estimates from generalized linear mixed

models (GLMMs) assessing the effects of selection regime, sex, and time (standardized week)

on the probability of mortality. Bayesian estimates are posterior means with 95% credible

intervals and posterior direction (PD), while frequentist GLMM estimates are shown with

standard errors (SE), z-values, and associated p-values. The reference levels are the base

population (C) and females.

| Effect | Bayesian estimate (95% CI) | PD direction (%) | glmer estimate (se) | z-value | p-value |
|---|---|---|---|---|---|
| Intercept | -4.53 (-4.83, -4.23) | 100 | -4.54 (0.14) | -32.23 | < 2e-16 |
| Week (z) | 0.89 (0.85, 0.93) | 100 | 0.89 (0.02) | 40.39 | **< 2e-16** |
| CH | 1.21 (0.75, 1.67) | 100 | 1.23 (0.21) | 5.82 | **5.9e-09** |
| DA | 0.09 (-0.37, 0.56) | 65.4 | 0.11 (0.21) | 0.51 | 0.6084 |
| FA | 0.45 (-0.01, 0.93) | 97.2 | 0.47 (0.21) | 2.21 | **0.0273** |
| Male | 0.33 (-0.01, 0.67) | 96.7 | 0.35 (0.17) | 2.04 | **0.0416** |
| CH : Male | -0.47 (-0.84, -0.09) | 99.0 | -0.49 (0.19) | -2.55 | **0.0109** |
| DA : Male | 0.12 (-0.25, 0.53) | 73.4 | 0.10 (0.20) | 0.52 | 0.6007 |
| FA : Male | -0.14 (-0.51, 0.26) | 75.4 | -0.16 (0.20) | -0.81 | 0.4210 |

**Figure 3:** Fixed effects from hierarchical modeling of survival. Posterior distributions and 95%

credible intervals (CIs) for fixed effects from the Bayesian model. Each ridge represents the

estimated posterior distribution of a model coefficient. Points indicate posterior medians, and

horizontal bars show 95% CIs. Vertical dashed line denotes the null effect (0). Parameters

include treatment effects (CH, DA, FA), sex (Male), and their interactions, relative to the

reference group (females in the control C regime). Estimates are on the scale of the linear

predictor.

Microbial composition across selection regimes and time points

To assess microbial composition, we first analyzed the average relative abundance of phyla

across time points and selection regimes (Fig. 4). Overall, all selection regimes (CH, DA, FA)

displayed broadly similar microbial compositions that were distinct from the control, primarily

due to lower Proteobacteria abundance and higher Firmicutes at the initial time point. Over time,

Proteobacteria steadily increased across all treatments, ultimately converging between 45–65 %

by T2. Other phyla, including Actinobacteria, Bacteroidota, and Fusobacteriota, were more

abundant in the selected regimes than in controls but generally declined with age (Additional file

3: Fig. S8c–e).

**Figure 4:** Phylum-level relative abundance of bacterial communities across timepoints and

dietary treatments. Stacked bar plots show the log-transformed relative abundance of bacterial

phyla at three timepoints t1, t2 and t3 (i.e. 15, 35, and 60 days post-eclosion) within each dietary

treatment: **a.** constant high sugar (CH), **b.** decreasing nutrient availability (DA), **c.** fluctuating

nutrient availablity (FA), and **d.** control non-selected group (C). Each bar represents the mean

relative abundance of samples collected at a given timepoint (n indicated above each bar).

Colors correspond to bacterial phyla, as shown in the shared legend.

To explore compositional significance of observed relative abundance dynamics, we performed

an indicator species analysis (ISA) at both the genus and species level. At the species level, an

indicator species analysis (ISA) identified 35 species differing across time points, 5 across

selection regimes, and 18 across both factors. Most time-related species (81 %) were enriched

at the earliest time point (T1), highlighting pronounced early-life microbiome restructuring.

Control samples were enriched in three *Acetobacter* species (*A. aceti*, *A. oryzifermentans*, and

*A. tropicalis*), whereas *Corynebacterium amycolatum* and *Mobiluncus curtisii* (Actinobacteria)

were associated with CH and FA, respectively (Table 2). Several species were linked to specific

combinations of diet and time. Notably, the genus of core Drosophila gut colonizers,

*Acetobacter,* were depleted in the selected lines but present across all time points in controls,

indicating that long-term dietary selection disrupted the *Acetobacter*-dominated microbiome

characteristic of non-selected flies. The full output of the ISA looking across time point and

selection regime can be found through (Additional file 4).

At the genus level, ISA identified 39 genera differing across time points, 8 across selection

regimes, and 28 across both factors (Additional file 2 Table S2). Among the diet-associated

genera, *Fructilactobacillus* (Firmicutes) was enriched in CH; *Mobiluncus* (Actinobacteria) and

*Pelomonas* (Proteobacteria) in FA; *Leptotrichia* (Proteobacteria) in CH + FA; *Acinetobacter*

(Proteobacteria) in DA + FA; and *Actinomyces* (Actinobacteria), *Neisseria* (Proteobacteria), and

*Lacticaseibacillus* (Firmicutes) in CH + DA + FA. Together, these results indicate that dietary

selection history reshaped both phylum- and genus-level composition, resulting in distinct,

convergent microbiomes among selected lines that diverged from the control.

Based on prior studies linking microbial composition to immune function [50, 51], we

hypothesized that Proteobacteria and Firmicutes could influence hosts positively or negatively

depending on their relative proportions. We calculated phyla ratios as age-specific indicators of

treatment effects (Additional file 3 Fig. S8f-g). In the selection regimes, the Proteobacteria-

Firmicute ratio generally increased over time (Additional file 3 Fig. S8f), while the Bacteriodota-

Firmicute ratio decreased between T1 and T2 but then increased in DA between T2 to T3

(Additional file 3 Fig. S8g). The control group began with higher ratios than the treatments and

exhibited a general decline, suggesting an age-related increase in Firmicutes relative to

Proteobacteria and Bacteriodota. These patterns reveal contrasting dynamics between

Proteobacteria and Bacteroidota that are influenced by Firmicutes.

Table 2: Species-level enrichment across time and selection regime.

| Species | Time point | Regime | Remark |
|---|---|---|---|
| *Acetobacter aceti* | T1, T2, T3 (C) | C only | Core gut colonizer; disrupted in treatments |
| *Acetobacter oryzifermentans* | T3 (DA), all time points (C) | C, DA | Found in fermented substrates |
| *Acetobacter tropicalis* | T3 (C) | C only | Common hindgut colonizer |
| *Acetobacter persici* | T1 (C, CH, DA), T2 (all), T3 (C, DA) | Mixed | Broad presence but absent in FA at later time points |

| | | | |
|---|---|---|---|
| *Corynebacterium amycolatum* | - | CH | Treatment-associated species |
| *Mobiluncus curtisii* | - | FA | Treatment-associated species |

## Microbial diversity dynamics and predictive modeling

We assessed microbial diversity across time points and dietary selection treatments using both

alpha and beta diversity metrics. At the ASV level, both observed richness (Fig. 5a) and

Shannon index (Fig. 5b), shifted significantly between T1 and T2 in all selection regimes,

particularly in DA and FA indicating substantial early-life microbial restructuring. Some

differences persisted into T3 in DA, although CH and FA lacked T3 data due to shortened

lifespans. Similar patterns were observed at the genus- evel (Fig. 5c-d). Beta-diversity analyses

(PCoA and PERMANOVA) revealed significant effects of time point, selection regime, and their

interaction (p = 0.001), with pairwise contrasts showing pronounced T1 to T2 shifts in DA and

FA (Fig. 6b-c). Dispersion tests further supported temporal changes within treatments,

especially in DA and CH, and early compositional divergence between control and selected

lines. Overall, age exerted a stronger influence than dietary regime on microbial community

structure.

**Figure 5:** Temporal changes in microbial alpha diversity across dietary treatments(CH, DA, FA and C) at ASV (a-b) and genus (c-d) levels. **a&c** are observed diversity while **b&d** are Shannon-Weiner index. Significance between timepoints is from pairwise Wilcoxon rank-sum tests ($p <$ 0.05), using raw p-values. Timepoints t1, t2 and t3 are 15, 35, and 60 days post-eclosion.

To evaluate the predictive value of microbial profiles, we trained random forest models to classify age, selection regime, and their interaction (Additional file 3 Fig. S10). The model predicting time point achieved the highest accuracy (78.8 %), identifying *Acetobacter persici*, *Cutibacterium acnes*, and *A. oryzifermentans* as key predictors. Models predicting dietary regime or the time x regime interaction performed less accurately (40.1 % and 32.2 %,

respectively), likely reflecting limited sample sizes in the control and T3. Across all models, *A.*

*persici* consistently emerged as a strong age-associated taxon, increasing in abundance in

older flies across regimes.

**Figure 6:** PCoA showing species diversity of dietary regimes over time (a - d) and across

regimes at each timepoint (e - g). **a.** constant high sugar (CH), **b.** decreasing nutrient availability

(DA), **c.** fluctuating nutrient availability (FA), **d.** control non-selected group (C), **e.** 15 days post

eclosion (dpe) (T1), **f.** 35 dpe (T2), and **g.** 60 dpe (T3). CH and FA groups did not reach t3,

hence no comparisons for T3 for a, c, and g. UniFrac distances are used as beta-diversity

metric.

Given its prominence in both random forest and indicator species analyses, we examined

*Acetobacter* abundance in greater detail (Fig. 7). *A. persici* was the most abundant species

across regimes, followed by *A. aceti* and *A. oryzifermentans*. From T1 to T2, *A. persici*

increased significantly across all selected regimes but not in controls, while *A. aceti* rose in DA

and FA. Between treatments, *A. aceti* differed significantly from controls in both T1 and T2, and

*A. oryzifermentans* showed broader regime-specific differences. Together, these results indicate

that age-related microbial shifts, dominated by *Acetobacter* taxa, are more consistent and

predictable than those driven by dietary selection history. However, persistent differences in

these core taxa between control and selected regimes suggest that long-term dietary selection

also left distinct microbial legacies shaping community composition across ages.

**Figure 7:** Prevalence of various Acetobacter species across samples: **a.** overview of the

average relative abundance of four different acetobacter species. **b.** change in abundance of

each species between 15 dpe (T1) and 35 dpe (T2) of each dietary regime. **c-e**. differences

between dietary regimes in **c.** *A. aceti*, **d.** *A. persici*, and **e.** *A. oryzifermentans* at both T1 and

T2. **b-e**. p-values from Wilcoxon ranked-sum tests.

**Discussion**

We investigated how long-term dietary selection and inbreeding histories affect survival and

microbial community dynamics in *Drosophila melanogaster*. To do so, we used individual diets

we have characterized earlier in the DSPR (i.e. high sugar, reduced yeast, and standard diet)

[31, 52, 53] to construct selection conditions, DA, FA and CH. While none of the selected lines

outperformed the control in lifespan, DA (decreasing protein) selection, representing repeated

exposure to a high-to-low  dietary environment, produced lifespan performance comparable to

the control group after 65 generations. In contrast, flies from FA and CH selection regimes

exhibited reduced lifespan when maintained on the standard diet. This pattern indicates that

chronic exposure to specific dietary stressors can leave lasting physiological and microbial

legacies on lifespan, even after long-term rearing in a common environment. These findings

extend our prior work and highlight that historical nutritional stress can generate enduring, and

sometimes maladaptive, consequences for longevity.

**Longevity and sex-specific responses to dietary selection regimes**

Among selection regimes, the constant high-sugar (CH) lines showed the strongest survival

cost, particularly in females. Rather than indicating incomplete adaptation, this likely reflects a

persistent metabolic legacy of chronic sugar exposure, which may influence host–microbe

interactions. Prolonged selection under carbohydrate-rich conditions may have reinforced

physiological priorities toward nutrient storage and reproduction at the expense of somatic

maintenance, leading to reduced lifespan when flies are maintained on a more balanced diet

[54, 55]. The pronounced early mortality in CH females suggests carryover effects of metabolic

dysregulation during peak reproductive activity, consistent with sugar-induced insulin resistance

observed in experimental evolution studies (e.g., [56]).

By contrast, lines exposed to fluctuating or deteriorating availability (FA and DA) exhibited more

moderate declines, potentially reflecting historical exposure that promoted metabolic flexibility or

resilience to nutrient variation rather than chronic abundance. In these lines, survival during

early adulthood (a key reproductive window) was maintained, whereas later mortality increased,

suggesting trade-offs in resource allocation between reproduction and maintenance.

Notably, sex-specific patterns emerged: FA males lived longer than females early in life,

possibly facilitating sustained mating at a cost to female survival, while in DA a mildly opposite

pattern was observed, consistent with maternal investment being favored under limited protein

supply. Given that our data focus on longevity patterns rather than a complete set of life-history

traits, we interpret these findings as reflecting evolved differences in physiological maintenance

rather than comprehensive life-history dynamics. Further, although mutation accumulation is a

theoretical factor influencing survival differences, it is unlikely to explain our results. The 65

generations of directional selection in large populations would have efficiently purged

deleterious alleles. Thus, the observed survival differences more plausibly reflect evolved

physiological adjustments to historical nutritional regimes rather than stochastic accumulation of

mutations. Overall, the results point to divergent physiological trajectories rooted in past dietary

environments, with lasting effects on survival even under a standardized diet.

Microbiome responses to selection

Selection regimes also produced distinct microbiome profiles, representing the primary and

most robust evidence of persistent effects of dietary selection. Compared to the control group,

selected lines exhibited increased microbial richness and altered taxonomic composition,

particularly early in life. The control flies maintained lower Firmicutes and higher Proteobacteria

in early adulthood, consistent with a more beneficial, stable microbiome previously associated

with greater lifespan in *Drosophila* [57, 58]. Selected lines experienced early colonization by

taxa such as *Corynebacterium* and *Mobiluncus*, which in other systems include opportunistic or

inflammation-associated species [16, 59, 60]. Studies in both flies and mammals show that early

microbiota disruptions can impair host immunity and longevity [12]. These findings suggest that

historical dietary stress can imprint persistent effects on microbiome assembly, potentially

compromising host survival persistently after the selective pressure is withdrawn. The DA (and

to a lesser degree, FA) lines, which maintained more survivors into later life, also showed

similar dynamics in microbial composition across age. This is consistent with work showing that

early adult microbiome establishment is critical for later-life health in flies [59, 61]. The ability to

maintain beneficial taxa may therefore represent a retained trait from prior selection rather than

ongoing adaptation in stressful nutritional environments.

We also detected consistent reductions in beneficial microbial taxa between the control and

selection treatments in both T1 and T2, particularly in two core *Acetobacter* species: *A. aceti*

and *A. oryzifermentans*. As members of the acetic acid bacteria group, *Acetobacter* species are

integral to the fruit fly microbiome and have been linked to critical host functions, including

growth and development, insulin signaling, lipid metabolism, and stem cell activity [62, 63]. In

our study, flies from the selection regimes exhibited lower relative abundances of these

microbes, suggesting that nutritional stress and inbreeding histories may have disrupted the

core microbiota. Such disruptions could impair physiological processes and contribute to

microbial imbalance and reduced survival. Interestingly, although *A. aceti* has been previously

associated with shorter lifespan in *D. melanogaster* [64], we observed the highest levels of *A.*

*aceti* and the longest lifespan in control flies. This contrast may reflect context-dependent

consequences rather than directional adaptation, underscoring how dietary stress can interfere

with host–microbiome relationships in ways that persist beyond the period of exposure.

Although flies were maintained for multiple generations on a standardized diet after the

selection phase, microbiome differences among lines remained evident, suggesting that

evolutionary history of selection constrained microbial assembly long after the environmental

stress was removed.

Patterns in dominant bacterial phyla

Patterns in dominant bacterial phyla further underscore the link between microbiome dynamics

and host physiology. The ratio of Bacteroidota to Firmicutes decreased with age in all lines,

probably suggesting an age-related shift in immune potential. Previous work associates higher

Bacteroidota with immune activation [50] and lower Firmicutes with gut homeostasis [58]. The

control flies, which had higher early Proteobacteria and lower Firmicutes, may have benefited

from this composition. Conversely, historically stressed lines with elevated Firmicutes may carry

immune and metabolic imbalances linked to prior selection, leading to greater community

reorganization. We also observed an inverse relationship between Proteobacteria and

Bacteroidota in selected lines, a pattern often linked to inflammation and metabolic stress [65,

66]. This microbial imbalance is well-documented in human and mouse studies under high-fat or

sugar diets [67], suggesting deep evolutionary parallels in how dietary stress leaves microbial

legacies across taxa.

Microbiome succession and indicator taxa

Indicator species analysis showed rapid microbiome restructuring early in adulthood. Most

microbial turnover occurred at T1, reinforcing the importance of early life as a window for

microbiome establishment. While many groups declined over time, only a few were enriched in

specific treatments, suggesting that historical exposure to dietary stress shaped which taxa

persisted, rather than implying active adaptation of the microbial community. The presence of

potentially pro-inflammatory species such as *Porphyromonas bennonis* and *Cutibacterium*

*acnes* in some selection treatments raises further concern. These taxa are linked with chronic

inflammation in mammals [68], and their persistence in flies may indicate long-term

consequences for immune homeostasis. The apparent protective role of *Lactobacillus* in some

groups aligns with studies highlighting its antagonistic effect on pathogens [69]. Thus, persistent

survival differences among selection lines may result from retained microbial assemblages

capable of modulating immune and metabolic balance, rather than ongoing adaptive co-

evolution.

Limitations and future directions

Our experimental design imposed a shared long-term post-selection environment, which could

mask environment-specific adaptive responses. Moreover, fewer flies from the CH and FA

groups survived to late age points, limiting effective comparisons at T3. Another limitation is that

we did not characterize the initial microbiome prior to experimental evolution or immediately

after the selection phase, which would have helped determine the magnitude of microbiome

evolution in both the control and selected lines before rearing on NutriFly. Additionally, this

experiment was not originally designed with microbiome evolution as a focal endpoint; thus, the

lack of sterility during culture, the possible introduction of environmental microbes, and the use

of food preservatives may have contributed to uncontrolled microbial variation. Nevertheless,

detectable and consistent patterns across selection regimes and time points long after cessation

of selection strengthen the inference that historical diet and subsequent inbreeding have left

persistent legacies on both survival and microbiome structure.

Future work should disentangle host and microbial contributions to longevity by directly

manipulating microbial communities in selection lines, ideally using gnotobiotic models.

Reciprocal microbiome transplants between lines could further test whether microbial legacies

from past selection can transfer phenotypes to naïve hosts. It would also be informative to re-

examine the lines on their original selection diets and to assess whether the microbiota

themselves evolved genetically or simply underwent species sorting. Finally, metagenomic

profiling would clarify whether observed shifts reflect changes in microbial function or taxonomic

identity.

**Conclusion**

This study demonstrates that historical dietary selection and inbreeding leave persistent effects

on the host-associated microbiome, which we consider the most robust outcome of this

experiment. Selection lines exhibited microbial restructuring, and potential microbiome

imbalance, while changes in survival and sex-specific life history strategies provide

complementary context for interpreting these microbial patterns. These findings reinforce the

idea that dietary history can imprint lasting host–microbiome legacies, influencing lifespan and

physiological trajectories through both host genetic and microbial pathways. Given the rising

relevance of diet-induced metabolic disorders in humans and pets, and changing climatic

conditions for natural species, understanding how nutrition shapes microbial dynamics offers a

promising path for translational insights.

**List of abbreviations**

C – control group, CH – constant high availability, DA – deteriorating availability, FA –

fluctuating availability, DSPR – Drosophila Synthetic Population Resource, SIL – selection

inbred line, ΔAIC – delta Akaike Index, ASV – amplicon sequence variant, ISA – indicator

species analysis.

**Data availability**

All 16S Illumina raw reeds are available in the NCBI Sequence Reads Archive (SRA) BioProject

ID PRJNA1233815. Computer code to reproduce all analyses is available as a GitHub

repository at https://github.com/nochet/selPhenotyping/ and analysis steps laid out in a

ProjSteps.md file.

**Declarations**

**Ethics approval and consent to participate**

Not applicable. Clinical trial number: not applicable.

**Consent for publication**

All authors give the publisher the right to publish this work.

**Competing interests**

The authors declare no competing interests.

**Acknowledgments**

Authors thank Dr. Elizabeth G. King for discussions and Arthur K. Miller for technical assistance

with experimental set up.

**Funding**

The study was supported by a University of Missouri start-up award and a University of

Washington Nathan Shock Center Pilot Project award to EN. Funding bodies were not involved

in the design, collection of data, analysis, interpretation, and writing of the manuscript.

**Authors' contributions**

PKW performed 16S sequencing and analysis and wrote the manuscript. GM performed all fly

experiments. GSK helped with data analysis and performed critical revision of the manuscript.

EN conceived the study, designed experiments, analyzed data, and wrote the manuscript.

## References

**References**

1. Ley RE, Hamady M, Lozupone C, Turnbaugh PJ, Ramey RR, Bircher JS, et al. Evolution of
mammals and their gut microbes. Science. 2008;320:1647–51.
https://doi.org/10.1126/science.1155725.

2. Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L, et al. Diet drives
convergence in gut microbiome functions across mammalian phylogeny and within humans.
Science. 2011;332:970–4. https://doi.org/10.1126/science.1198719.

3. Gould AL, Zhang V, Lamberti L, Jones EW, Obadia B, Korasidis N, et al. Microbiome
interactions shape host fitness. Proceedings of the National Academy of Sciences.
2018;115:E11951–60. https://doi.org/10.1073/pnas.1809349115.

4. Rudman SM, Greenblum S, Hughes RC, Rajpurohit S, Kiratli O, Lowder DB, et al.
Microbiome composition shapes rapid genomic adaptation of Drosophila melanogaster.
Proceedings of the National Academy of Sciences. 2019;116:20025–32.
https://doi.org/10.1073/pnas.1907787116.

5. Petersen C, Hamerich IK, Adair KL, Griem-Krey H, Torres Oliva M, Hoeppner MP, et al. Host
and microbiome jointly contribute to environmental adaptation. ISME J. 2023;17:1953–65.
https://doi.org/10.1038/s41396-023-01507-9.

6. Härer A, Thompson KA, Schluter D, Rennison DJ. Associations between gut microbiota
diversity and a host fitness proxy in a naturalistic experiment using threespine stickleback fish.
Molecular Ecology. 2024;33:e17571. https://doi.org/10.1111/mec.17571.

7. Storelli G, Defaye A, Erkosar B, Hols P, Royet J, Leulier F. Lactobacillus plantarum promotes
Drosophila systemic growth by modulating hormonal signals through TOR-dependent nutrient
sensing. Cell Metab. 2011;14:403–14. https://doi.org/10.1016/j.cmet.2011.07.012.

8. de Weerth C. Do bacteria shape our development? Crosstalk between intestinal microbiota
and HPA axis. Neuroscience & Biobehavioral Reviews. 2017;83:458–71.
https://doi.org/10.1016/j.neubiorev.2017.09.016.

9. Rowe M, Veerus L, Trosvik P, Buckling A, Pizzari T. The reproductive microbiome: an
emerging driver of sexual selection, sexual conflict, mating systems, and reproductive isolation.
Trends in Ecology & Evolution. 2020;35:220–34. https://doi.org/10.1016/j.tree.2019.11.004.

10. Suyama R, Cetraro N, Yew JY, Kai T. Microbes control Drosophila germline stem cell
increase and egg maturation through hormonal pathways. Commun Biol. 2023;6:1–18.
https://doi.org/10.1038/s42003-023-05660-x.

11. Erkosar B, Storelli G, Defaye A, Leulier F. Host-intestinal microbiota mutualism: "learning on
the fly." Cell Host Microbe. 2013;13:8–14. https://doi.org/10.1016/j.chom.2012.12.004.

12. Spragge F, Bakkeren E, Jahn MT, B. N. Araujo E, Pearson CF, Wang X, et al. Microbiome
diversity protects against pathogens by nutrient blocking. Science. 2023;382:eadj3502.
https://doi.org/10.1126/science.adj3502.

13. Christensen SM, Srinivas SN, McFrederick QS, Danforth BN, Buchmann SL, Vannette RL.
Symbiotic bacteria and fungi proliferate in diapause and may enhance overwintering survival in
a solitary bee. The ISME Journal. 2024;18:wrae089. https://doi.org/10.1093/ismejo/wrae089.

14. Clark RI, Salazar A, Yamada R, Fitz-Gibbon S, Morselli M, Alcaraz J, et al. Distinct shifts in
microbiota composition during Drosophila aging impair intestinal function and drive mortality.
Cell Reports. 2015;12:1656–67. https://doi.org/10.1016/j.celrep.2015.08.004.

15. Aleman FDD, Valenzano DR. Microbiome evolution during host aging. PLOS Pathogens.
2019;15:e1007727. https://doi.org/10.1371/journal.ppat.1007727.

16. Broderick NA, Buchon N, Lemaitre B. Microbiota-induced changes in Drosophila
melanogaster host gene expression and gut morphology. mBio. 2014;5:10.1128/mbio.01117-14.
https://doi.org/10.1128/mbio.01117-14.

17. Shapira M. Gut microbiotas and host evolution: scaling up symbiosis. Trends in Ecology &
Evolution. 2016;31:539–49. https://doi.org/10.1016/j.tree.2016.03.006.

18. Okin D, Medzhitov R. Evolution of inflammatory diseases. Current Biology. 2012;22:R733–
40. https://doi.org/10.1016/j.cub.2012.07.029.

19. Wong A, Chaston J, Douglas A. The inconstant gut microbiota of Drosophila species
revealed by 16S rRNA gene analysis. The ISME journal. 2013;7.
https://doi.org/10.1038/ismej.2013.86.

20. Martino ME, Joncour P, Leenay R, Gervais H, Shah M, Hughes S, et al. Bacterial adaptation
to the host's diet is a key evolutionary force shaping Drosophila-Lactobacillus symbiosis. Cell
Host Microbe. 2018;24:109-119.e6. https://doi.org/10.1016/j.chom.2018.06.001.

21. Wong AC-N, Dobson AJ, Douglas AE. Gut microbiota dictates the metabolic response of
Drosophila to diet. Journal of Experimental Biology. 2014;217:1894–901.
https://doi.org/10.1242/jeb.101725.

22. Lee KP, Simpson SJ, Clissold FJ, Brooks R, Ballard JWO, Taylor PW, et al. Lifespan and
reproduction in Drosophila: new insights from nutritional geometry. Proc Natl Acad Sci U S A.
2008;105:2498–503. https://doi.org/10.1073/pnas.0710787105.

23. Simpson SJ, Raubenheimer D. Macronutrient balance and lifespan. Aging (Albany NY).
2009;1:875–80. https://doi.org/10.18632/aging.100098.

24. Piper MDW, Partridge L, Raubenheimer D, Simpson SJ. Dietary restriction and aging: a
unifying perspective. Cell Metab. 2011;14:154–60. https://doi.org/10.1016/j.cmet.2011.06.013.

25. Skorupa DA, Dervisefendic A, Zwiener J, Pletcher SD. Dietary composition specifies
consumption, obesity, and lifespan in Drosophila melanogaster. Aging Cell. 2008;7:478–90.
https://doi.org/10.1111/j.1474-9726.2008.00400.x.

26. Grandison RC, Piper MDW, Partridge L. Amino-acid imbalance explains extension of
lifespan by dietary restriction in Drosophila. Nature. 2009;462:1061–4.
https://doi.org/10.1038/nature08619.

27. Sannino DR, Dobson AJ, Edwards K, Angert ER, Buchon N. The Drosophila melanogaster
gut microbiota provisions thiamine to its host. mBio. 2018;9:10.1128/mbio.00155-18.
https://doi.org/10.1128/mbio.00155-18.

28. Arias-Rojas A, Iatsenko I. The role of microbiota in Drosophila melanogaster aging.
Frontiers in Aging. 2022;3.

29. King EG, Macdonald SJ, Long AD. Properties and power of the Drosophila Synthetic
Population Resource for the routine dissection of complex traits. Genetics. 2012;191:935–49.
https://doi.org/10.1534/genetics.112.138537.

30. King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, et al. Genetic dissection
of a model complex trait using the Drosophila Synthetic Population Resource. Genome Res.
2012;22:1558–66. https://doi.org/10.1101/gr.134031.111.

31. Ng'oma E, Fidelis W, Middleton KM, King EG. The evolutionary potential of diet-dependent
effects on lifespan and fecundity in a multi-parental population of Drosophila melanogaster.
Heredity. 2019;122:582–94. https://doi.org/10.1038/s41437-018-0154-2.

32. Pool JE. The mosaic ancestry of the Drosophila Genetic Reference Panel and the D.
melanogaster Reference Genome reveals a network of epistatic fitness interactions. Molecular
Biology and Evolution. 2015;32:3236–51. https://doi.org/10.1093/molbev/msv194.

33. Bass TM, Grandison RC, Wong R, Martinez P, Partridge L, Piper MDW. Optimization of
dietary restriction protocols in Drosophila. J Gerontol A Biol Sci Med Sci. 2007;62:1071–81.
https://doi.org/10.1093/gerona/62.10.1071.

34. Chapman T, Liddle LF, Kalb JM, Wolfner MF, Partridge L. Cost of mating in Drosophila
melanogaster females is mediated by male accessory gland products. Nature. 1995;373:241–4.
https://doi.org/10.1038/373241a0.

35. Wigby S, Chapman T. Sex peptide causes mating costs in female Drosophila melanogaster.
Curr Biol. 2005;15:316–21. https://doi.org/10.1016/j.cub.2005.01.051.

36. Jones A, Donnell D, King EG, Ng'oma E. Larval protein restriction interacts with adult diet to
increase fitness in an outbred multiparent population. 2024;:2024.02.10.579751.
https://doi.org/10.1101/2024.02.10.579751.

37. Walters WA, Caporaso JG, Lauber CL, Berg-Lyons D, Fierer N, Knight R. PrimerProspector:
de novo design and taxonomic analysis of barcoded polymerase chain reaction primers.
Bioinformatics. 2011;27:1159–61. https://doi.org/10.1093/bioinformatics/btr087.

38. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al.
Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
Proceedings of the National Academy of Sciences. 2011;108 supplement_1:4516–22.
https://doi.org/10.1073/pnas.1000080107.

39. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13:581–3.
https://doi.org/10.1038/nmeth.3869.

40. Simhadri RK, Fast EM, Guo R, Schultz MJ, Vaisman N, Ortiz L, et al. The gut commensal
microbiome of Drosophila melanogaster is modified by the endosymbiont wolbachia. mSphere.
2017;2:e00287-17. https://doi.org/10.1128/mSphere.00287-17.

41. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. PLOS ONE. 2013;8:e61217.
https://doi.org/10.1371/journal.pone.0061217.

42. Greenacre M, Martínez-Álvaro M, Blasco A. Compositional data analysis of microbiome and
any-omics datasets: a validation of the additive logratio transformation. Front Microbiol.
2021;12. https://doi.org/10.3389/fmicb.2021.727398.

43. Karwowska Z, Aasmets O, Metspalu M, Metspalu A, Milani L, Esko T, et al. Effects of data
transformation and model selection on feature importance in microbiome classification data.
Microbiome. 2025;13:2. https://doi.org/10.1186/s40168-024-01996-6.

44. McKnight DT, Huerlimann R, Bower DS, Schwarzkopf L, Alford RA, Zenger KR. Methods for
normalizing microbiome data: An ecological perspective. Methods in Ecology and Evolution.
2019;10:389–400. https://doi.org/10.1111/2041-210X.13115.

45. Paradis E, Claude J, Strimmer K. APE: analyses of phylogenetics and evolution in R
language. Bioinformatics. 2004;20:289–90. https://doi.org/10.1093/bioinformatics/btg412.

46. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin P, O'Hara R, et al. Vegan:
community ecology package. R package vegan, vers. 2.2-1. 2015.

47. Anderson MJ. Distance-based tests for homogeneity of multivariate dispersions. Biometrics.
2006;62:245–53. https://doi.org/10.1111/j.1541-0420.2005.00440.x.

48. Cáceres MD, Legendre P. Associations between species and groups of sites: indices and
statistical inference. Ecology. 2009;90:3566–74. https://doi.org/10.1890/08-1823.1.

49. Sultanova Z, Dönertaş HM, Hita A, Aguilar P, Dag B, Lucas-Lledo JI, et al. Age-dependent
gut microbiota dynamics and their association with male fitness traits in Drosophila
melanogaster. 2025;:2025.05.22.655607. https://doi.org/10.1101/2025.05.22.655607.

50. Shen X, Miao J, Wan Q, Wang S, Li M, Pu F, et al. Possible correlation between gut
microbiota and immunity among healthy middle-aged and elderly people in southwest China.
Gut Pathogens. 2018;10:4. https://doi.org/10.1186/s13099-018-0231-3.

51. Gibbons J, Yoo JY, Mutka T, Groer M, Ho TTB. A pilot study to establish an in vitro model to
study premature intestinal epithelium and gut microbiota interactions. mSphere.
2021;6:10.1128/msphere.00806-21. https://doi.org/10.1128/msphere.00806-21.

52. Stanley PD, Ng'oma E, O'Day S, King EG. Genetic dissection of nutrition-induced plasticity
in insulin/insulin-like growth factor signaling and median life span in a *Drosophila* multiparent
population. Genetics. 2017;206:587–602. https://doi.org/10.1534/genetics.116.197780.

53. Ng'oma E, Williams-Simon PA, Rahman A, King EG. Diverse biological processes
coordinate the transcriptional response to nutritional changes in a Drosophila melanogaster

multiparent population. BMC Genomics. 2020;21:84. https://doi.org/10.1186/s12864-020-6467-
6.

54. Rose MR, Charlesworth B. Genetics of life history in Drosophila melanogaster. I. sib
analysis of adult females. Genetics. 1981;97:173–86.

55. Kawecki TJ, Ebert D. Conceptual issues in local adaptation. Ecology Letters. 2004;7:1225–
41. https://doi.org/10.1111/j.1461-0248.2004.00684.x.

56. Leitão-Gonçalves R, Carvalho-Santos Z, Francisco AP, Fioreze GT, Anjos M, Baltazar C, et
al. Commensal bacteria and essential amino acids control food choice behavior and
reproduction. PLOS Biology. 2017;15:e2000862. https://doi.org/10.1371/journal.pbio.2000862.

57. Staubach F, Baines JF, Künzel S, Bik EM, Petrov DA. Host species and environmental
effects on bacterial communities associated with Drosophila in the laboratory and in the natural
environment. PLOS ONE. 2013;8:e70749. https://doi.org/10.1371/journal.pone.0070749.

58. Lee H-Y, Lee S-H, Lee J-H, Lee W-J, Min K-J. The role of commensal microbes in the
lifespan of Drosophila melanogaster. Aging (Albany NY). 2019;11:4611–40.
https://doi.org/10.18632/aging.102073.

59. Blum JE, Fischer CN, Miles J, Handelsman J. Frequent replenishment sustains the
beneficial microbiome of Drosophila melanogaster. mBio. 2013;4:10.1128/mbio.00860-13.
https://doi.org/10.1128/mbio.00860-13.

60. Chandler JA, Lang JM, Bhatnagar S, Eisen JA, Kopp A. Bacterial communities of diverse
Drosophila species: Ecological context of a host–microbe model system. PLOS Genetics.
2011;7:e1002272. https://doi.org/10.1371/journal.pgen.1002272.

61. Brummel T, Ching A, Seroude L, Simon AF, Benzer S. Drosophila lifespan enhancement by
exogenous bacteria. Proceedings of the National Academy of Sciences. 2004;101:12974–9.
https://doi.org/10.1073/pnas.0405207101.

62. Shin SC, Kim S-H, You H, Kim B, Kim AC, Lee K-A, et al. Drosophila microbiome modulates
host developmental and metabolic homeostasis via insulin signaling. Science. 2011;334:670–4.
https://doi.org/10.1126/science.1212782.

63. Newell PD, Douglas AE. Interspecies interactions determine the impact of the gut microbiota
on nutrient allocation in Drosophila melanogaster. Appl Environ Microbiol. 2014;80:788–96.
https://doi.org/10.1128/AEM.02742-13.

64. Obata F, Fons CO, Gould AP. Early-life exposure to low-dose oxidants can increase
longevity via microbiome remodelling in Drosophila. Nat Commun. 2018;9:975.
https://doi.org/10.1038/s41467-018-03070-w.

65. Mukhopadhya I, Hansen R, El-Omar EM, Hold GL. IBD-what role do Proteobacteria play?
Nat Rev Gastroenterol Hepatol. 2012;9:219–30. https://doi.org/10.1038/nrgastro.2012.14.

66. Shin N-R, Whon TW, Bae J-W. Proteobacteria: microbial signature of dysbiosis in gut
microbiota. Trends Biotechnol. 2015;33:496–503. https://doi.org/10.1016/j.tibtech.2015.06.011.

67. Bradley PH, Pollard KS. Proteobacteria explain significant functional variability in the human
gut microbiome. Microbiome. 2017;5:36. https://doi.org/10.1186/s40168-017-0244-z.

68. Caselli E, Soffritti I, D'Accolti M, Piva I, Greco P, Bonaccorsi G. Atopobium vaginae and
Porphyromonas somerae induce proinflammatory cytokines expression in endometrial cells: a
possible implication for endometrial cancer? Cancer Manag Res. 2019;11:8571–5.
https://doi.org/10.2147/CMAR.S217362.

69. Moses RP, Y. Dawood H, Fashemi T, N. Fichorova R, Onderdonk AB, S. Yamamoto H.
Lactobacillus curbs inflammatory responses to atopobium vaginae. Obstetrics & Gynecology.
2015;125:3S. https://doi.org/10.1097/01.AOG.0000463525.50901.93.