1 **Title: Nutritional stress adaptation shapes host microbiome stability and life history**
2 **dynamics**
3
4 **Running title: Adaptive microbiota dynamics to dietary stress**
5

6 Peyton K. Warren, Garrison Miller[1], Gaurav S. Kandlikar[2], and Enoch Ng'oma*
7 Division of Biological Sciences, University of Missouri-Columbia, MO, USA
8
9 *Correspondence: ngomae@missouri.edu

10
11
12 **Author contributions:** PKW performed 16S sequencing and analysis and wrote the
13 manuscript. GM performed all fly experiments. GSK helped with data analysis and performed
14 critical revision of the manuscript. EN conceived the study, designed experiments, analyzed
15 data, and wrote the manuscript.
16

17 **Acknowledgments:** The study was supported by a University of Missouri start-up award and a
18 University of Washington Nathan Shock Center pilot project award to EN. Authors thank E.G.
19 King for discussions, A K. Miller for technical assistance with experimental set up.

20
21 *Conflict of interest*: The authors declare no conflicts of interest.
22
23 **Data Accessibility Statement**
24 All 16S Illumina raw reeds are available in the NCBI Sequence Reads Archive (SRA)
25 BioProjectID PRJNA1233815. Computer code to reproduce all analyses is available as a
26 GitHub repository at https://github.com/nochet/selPhenotyping/ and analysis steps laid out in a
27 ProjSteps.md file.
28
29
30
31

---

[1] PKW and GM should be considered joint first author

[2] Now, at Department of Biological Sciences, Louisiana State University, LA

Abstract

Dietary environments can influence host-microbe evolution by shaping adaptive fitness responses. However, host-microbe dynamics on evolutionary time scales remain under-characterized. We studied *Drosophila melanogaster* experimentally evolved under three nutrient regimes: constant high (CH), deteriorating (DA), and fluctuating (FA) availability, alongside unselected controls (C). DA and FA increased longevity relative to CH, with DA enhancing late-life survival, though all selected groups showed higher mortality than C, suggesting fitness costs of dietary adaptation. Sex-specific effects emerged, with FA favoring early-life female survival and DA supporting late-life female longevity. Microbial richness and composition shifted with selection and time. While Firmicutes and Proteobacteria dominated generally, selection regimes showed higher Firmicute and lower Proteobacteria abundances compared to C, correlating with dysbiotic microbiomes starting early in adulthood. DA flies maintained a relatively stable microbial diversity over time, potentially supporting longevity. In contrast, CH and FA both exhibited low microbial diversity that destabilized with age. Indicator species analysis revealed treatment- and age-specific genera, with CH and FA showing greater dysbiosis, marked by potentially pathogenic taxa such as *Leptotrichia* and *Mobiluncus*. Our findings highlight the importance of early life stages in maintaining microbiome stability and host fitness, suggesting dietary stress drives distinct microbiota patterns and survival strategies.

Keywords: experimental evolution, nutritional environment, longevity, microbiota dynamics, adaptive response

Introduction

Microorganisms are deeply integrated into host biology across taxonomic and functional scales, playing critical roles in organismal adaptation and fitness (Gould et al. 2018). Rapidly changing environmental conditions increase the pressure for hosts to adapt to new conditions faster than their genomes can evolve. Microbial genomes, which evolve more rapidly than their hosts', represent a potential adaptive variation, enabling hosts to respond more quickly to environmental changes. For example, microbiotas can contribute functional hologenomes capable of modulating host genomic responses to environmental change (Petersen et al., 2023). Indeed, variation in microbiome composition has been linked to allele frequency shifts in genes associated with local adaptation in *Drosophila* (Rudman et al. 2019). More broadly, microbiotas can influence host life-history traits by mediating development (de Weerth 2017), reproduction (Rowe et al. 2020; Suyama et al. 2023), survival (Aleman and Valenzano 2019; Christensen et al. 2024), and defense against pathogens (Spragge et al. 2023; Christensen et al. 2024).

Microbiotas facilitate various vital functions for their hosts, including nutritional intake (Leitão-Gonçalves et al. 2017), metabolic regulation (Wong et al. 2014), and maintaining fertility under poor diet (Gnainsky et al. 2021). For instance, in *Drosophila melanogaster*, environmental microbes enhance oogenesis by promoting germline stem cell proliferation and egg maturation through increased ovarian cell division and repression of apoptosis (Suyama et al. 2023). During diapause in a ground-nesting solitary bee species, immature stages maintain a core microbiome of Actinobacteria (*Streptomyces* and *Nocardiodes*) that inhibit pathogenic fungal growth while supporting fungal groups that increase cold tolerance and overwintering survival (Christensen et al. 2024). In some amphibians, environmental and host-derived microbiomes shape defense against pathogens, such as the skin surface bacterium *Janthinobacterium lividum*, which protects the frog *Rana muscosa* from a lethal skin fungus (Harris et al., 2009). These examples illustrate the adaptive roles of beneficial microbe-host interactions in diverse environmental contexts.

Nutritional environments play a dominant role in shaping the composition, richness, and abundance of beneficial microbiota, ultimately determining host fitness and life-history outcomes. For example, *Drosophila* harbor microbial communities shaped by diet rather than host genotype, as seen in the facultative symbiosis between the host and *Lactobacillus plantarum* in which microbial production of N-acetyl-amino acids from diet enhance host growth

(Martino et al. 2018). Similarly, in honeybees, gut microbiotas biosynthesize essential amino

acids critical for host nutrition (Parish et al. 2022). Environmental microbiomes can also shape

host immunity, as shown in amphibians, where climatic variation impacts microbial species

richness and functional traits (Kueneman et al. 2019). Thus, by directly modulating microbial

composition and function, environmental conditions can have profound indirect effects on host

adaptability and resilience to stressors.

Intrinsic host factors, including host age, physiology, and life-history strategies, also shape

microbiome composition and function. In humans, age-related changes in microbiota parallel

physiological shifts; younger individuals harbor microbiomes enriched with Bacteroidetes, while

older adults exhibit increased prevalence of *Streptococcus* (Zhang et al. 2021). Similarly, age-

related shifts in *D. melanogaster* behavior, locomotion, and cognition coincide with shifts in

microbiome dominance (Chiang et al. 2022), pointing to a potentially interactive role of host and

microbiota effects on key organismal traits. Microbiotas can also mediate life-history trade-offs.

For example, in seasonal killifish, gut microbiomes regulate longevity and aging rate, reflecting

trade-offs shaped by environmental conditions and microbial dynamics (Aleman and Valenzano

2019). These examples highlight the intricate interplay between microbiota and host biology,

with long-term implications for adaptation and fitness.

Here, we exposed a population derived from the *Drosophila Synthetic Population Resource*

(DSPR) to 65 generations of dietary selection followed by 38 generations of inbreeding in

common garden to investigate the effects of long-term dietary selection on microbiota richness

and host longevity under stressful nutritional conditions. The dietary regimes included

deteriorating nutrient availability (DA), fluctuating nutrient availability (FA), and constant high

nutrient availability (CH) (see Methods). We hypothesized that adaptation to these dietary

environments would result in distinct microbial communities and survival patterns shaped by the

interplay of diet, microbiota, and host physiology. Specifically, we predicted that: 1) the DA

regime would enrich for microbial taxa promoting somatic maintenance and longevity, supported

by lower pro-inflammatory dynamics; 2) the FA treatment, characterized by cyclical resource

availability, would prioritize reproduction over longevity, leading to divergent microbial

compositions and increased pro-inflammatory taxa compared to DA; and 3) the CH treatment

would exhibit stable microbiota compositions with reduced microbial diversity, reflecting

selection against traits favoring resource storage or stress resilience. Additionally, we

anticipated that age-related shifts in microbiota would interact with these dietary regimes,

revealing unique dynamics in survival across treatments. This experiment provides insight into

how environmental pressures, mediated by diet, shape the evolution of host-microbiota

interactions and their consequences for fitness under stress.

Materials and Methods

Study population

The study population was derived from a long-term, experimentally evolved population of

Drosophila melanogaster (Ng'oma et al. 2019). The base population was established by mass-

intercrossing 835 recombinant inbred lines from the B population (pB) of the Drosophila

Synthetic Population Resource (DSPR) for five generations in common garden (Figure 1A).

From this genetically diverse base, 36 populations were established and subjected to one of

three dietary selection regimes (described below) for 65 generations (Figure 1B). Following

selection, populations were inbred for 38 generations (~19 months) under common garden

conditions to generate Selection Inbred Lines (SILs).

Figure 1: Study design. **A.** A total of 835 recombinant inbred lines of the DSPR were interbred into a genetically heterogenous population which was subjected to experimental evolution for 65 generations. **B.** Three selection treatments involved diet alternations over a 21-day cycle per generation involving fluctuating (FA), deteriorating (DA) and constant high (CH) nutrient availability. An unselected group resulting from the base population was used as a control group (C). **C.** Multiple replicate selection lines maintained in common garden were assayed at three time points for survival and microbiome analysis at 15, 35, and 60 days post-eclosion.

For this study, six randomly selected SILs from each selection regime FA, DA, and CH (i.e.,

fluctuating, deteriorating, and constant high availability, respectively) were used alongside a

non-selected control (C) population (maintained at N > 4,000 flies). Each SIL was split into six

replicate vials (24 females and 6 males per vial): three vials were assigned to microbiome

assays, and three to lifespan estimation (Figure 1C). Microbiota sampling was performed on

females only to capture microbiomes of naturally mating individuals while avoiding effects of

male mating pressure. Whole flies were collected at three time points post-eclosion (15, 35, and

60 days) based on prior studies indicating key variability in lifespan and fecundity responses to

diet. At each time point, four females were flash-frozen and stored at -80°C for analysis. Surface

sterilization was omitted as environmental microbiomes in vials closely resemble gut

microbiomes in *Drosophila* (Douglas 2018).

Selection treatment regimes

Selection effects were analyzed using three diet regimes adapted from Bass et al. (2007) and

Skorupa et al. (2008), and previously used in our studies (Stanley et al. 2017; Ng'oma et al.

2019, 2020). Relative to the control diet (C, 200 g/L yeast, 50 g/L sucrose), the restricted (DR)

diet contained 0.5x less yeast and 1x sucrose. The high sugar (HS) diet included 1x yeast, but

7.2x sugar compared to the C diet. Thus, DR differed from C only in protein content, while HS

differed in sugar content. The C diet was considered optimal for growth and reproduction and

provided a reference for nutritionally stressful diets DR and HS due to restriction or oversupply

of key nutrients (previously characterized in Ng'oma et al. (2019). The FA treatment alternated

nutrient availability as follows: flies received C until day 14 post-oviposition (po), DR from days

14–18 po, and C again until day 21 po when eggs were collected for the next generation. The

DA treatment provided C until day 18 po, followed by DR until day 21 po. The CH treatment

provided HS until day 21 po. Diets were prepared with inactivated SAFPro Relax + YF 73050

yeast (Lesaffre Yeast Corp., Milwaukee, USA), stored at 4°C, and used within 14 days. Evolved

lines were held on a common Nutri-fly molasses formulation (Genesee Scientific) based on a

Bloomington Drosophila Stock Center Nutri-fly molasses protocol (12.4 g/L yeast, 61.3 g/L

yellow cornmeal, and 75.2 ml/L molasses). Across all assays (experimental evolution, survival

and microbiome assays), flies were maintained in a temperature-controlled chamber at 23°C

under continuous light. These conditions mirrored those under which the DSPR was originally

generated and maintained (King et al. 2012). Flies were supplied with fresh food three times per

week (Monday, Wednesday, Friday).

Phenotype measurement and 16S sequencing

We measured lifespan and microbiota composition (species diversity and richness) in flies

subjected to 65 generations of dietary selection followed by 38 generations of inbreeding in

common garden. For lifespan, three replicate vials per SIL, each with 24 females and 6 males,

were monitored thrice weekly (Monday, Wednesday, Friday) until all flies died. For microbiota

analysis, vials with the same initial composition were used for periodic female collections. DNA

was extracted from pools of 2–4 whole flies using the Qiagen DNeasy Blood & Tissue Kit (Cat.

No 69504) following the Quick-Start Protocol. Samples were manually homogenized with a

glass pestle in ATL buffer, incubated for 4 hours, and processed per protocol. DNA yield and

quality were assessed using a NanoDrop Spectrophotometer (ND-1000) and Qubit 4

Fluorometer with the Qubit 1X dsDNA HS Assay Kit (Cat. No Q33231). Samples were

normalized as needed with elution buffer and sequenced on a MiSeq instrument (PE250-v2

Nano flow cell, 25 cycles) targeting the V4 region of the bacterial 16S rRNA gene with primers

515F and 807R. Sequencing was performed at the University of Missouri Genomics Technology

Core.

Data Analysis

Lifespan

We analyzed lifespan data (N = 2,427 individuals; 2,242 deaths, 185 censored) using

survival models. Preliminary Kaplan-Meier analysis visualized global survival patterns

and sex-filtered survival at key time points (15, 35, 60 days). To assess evolved

differences in mortality risk after 65 generations of selection, we fitted a Cox

proportional hazards model: `~ regime + strata(sex)` and `~ regime*sex`. Stratification

allowed sex-specific baseline hazards while maintaining proportionality (Mills 2010;

Gordon 2021). To evaluate shifts in baseline hazards due to selection, we tested eight

parametric survival models (e.g., Gompertz, generalized gamma) using the **flexsurv**

and **muhaz** R libraries. The generalized gamma and Gompertz models provided the

best fit (Appendix Figure 1a). Our recent study testing effects of larval nutrition on adult

phenotypes in this population (Jones et al. 2024) showed that patterns of survival can

vary significantly across shorter periods of time or adult stages. To understand evolved

patterns of survival in each selection treatment, we examined percentile survival

differences in ten-day intervals in which each of our sampling time points formed the

median: 10-20, 30-40, and 55-65 days, post eclosion. We tested for significant

differences between treatments in each time interval I1, I2 and I3 using either analysis

of variance (Anova) or Kruskall-Wallis tests followed by an appropriate post-hoc test

(Turkey's HSD or Dunn's Test, respectively)

Microbiome analysis

We processed Illumina reads using the DADA2 pipeline (Callahan et al. 2016), leveraging its

amplicon sequence variant (ASV) approach to achieve higher resolution and reproducibility in

microbial community profiling. ASVs provide exact biological sequences, avoiding the limitations

of traditional OTU clustering. Reads were quality-filtered and trimmed at positions 225 (forward)

and 200 (reverse), and chimeras were removed. Taxonomy was assigned using the Silva

database (v138.1). Samples with fewer than 3,000 reads were excluded. To ensure a more

accurate representation of microbial communities, taxa identified as Wolbachia were removed

(Simhadri et al. 2017). Sequences were imported into **phyloseq** (McMurdie and Holmes 2013)

for analysis and visualization. Read counts were normalized using a log-transformation: `log(1`

`+ x)/sum(log(1 + x))`, to obtain relative abundances (i.e., the proportion of specific microbial

groups within samples). This approach combines two commonly used normalization equations,

log(1 + x) and (x)/sum(x), into one step and effectively scaling data to relative abundances while

revealing subtle differences (McKnight et al. 2019).

Indicator species analysis (ISA) was performed using **indicspecies** (Cáceres and Legendre

2009) to identify taxa enriched in specific treatments or time points. Alpha diversity (Shannon

and observed richness) was visualized with **phyloseq**, capturing richness and evenness. Beta

diversity was assessed using principal coordinate analysis (PCoA) based on UniFrac distances

obtained from phylogenetic trees generated by **ape** (Paradis et al. 2004). **vegan** (Oksanen et al.

2015) was used for PERMANOVA to test group differences in centroids and dispersion, and

PERMDISP to evaluate variability in dispersion. Visual inspection aided interpretation of

whether differences reflected means, dispersion, or both.

Results

Evolution of longevity following diet selection

Lifespan performance in the non-selected base population (C) served as a benchmark for

evaluating selection treatments (Figure 1). Summary survival patterns are presented in

Supplementary Table S1. Overall, DA- and FA-selected populations lived significantly longer

than CH-selected flies (Log-Rank tests: CH vs DA, p = 7.6e-45; CH vs FA, p = 3.3e-27; CH vs

C, p = 4.5e-35). DA flies also outlived FA flies (p = 4.1e-5), but the control group exhibited the

longest lifespan, significantly exceeding both DA (p = 0.025) and FA (p = 2.5e-07) regimes

(Figure 2a). Early adulthood (~38 days post-eclosion) showed higher male survival in controls

(Figure 2c&e), whereas greater female mortality under CH conditions likely reflected metabolic
stress from the high-sugar diet.

Figure 2: Survival performance in each treatment: **a** sexes together, **b** males, **c** females,
d in bins of 10 days in which the sampling time point is the median (only bins containing
assayed time points of 15, 35 and 60 days are shown (I1=10-20 days, I2=30-40 days,

245   To understand evolved patterns of survival in each selection treatment, we examined percentile
246   differences in survival in ten-day intervals in which each of our sampling time points formed the
247   median (I1: 10–20 days, I2: 30–40 days, I3: 55–65 days, Figure 2g). Two key patterns
248   emerged:

249   1. *Selection regime differences*: DA and FA regimes showed reduced early mortality in I1
250   and I2, favoring longevity, but diverged in I3, with FA experiencing higher late mortality
251   while DA maintained slower rates (Supplementary Table S2). Control flies exhibited the
252   lowest mortality across I1 and I2. In contrast, CH flies suffered heavy mortality across all
253   intervals.
254   2. *Sex-based variation*: CH and FA treatments favored male survival in I1 and I2, reversing
255   in I3 with higher female survival. Conversely, DA and control regimes showed greater
256   female survival later in life (I2, I3; Figure 2e&g, Supplementary Figure S1), suggesting
257   adaptation to dietary stress through reduced early mortality (DA, FA) or combined early
258   and late-life mortality reduction (DA, control).
These findings suggest that dietary stress adaptation involves distinct survival strategies, with
DA and FA regimes prioritizing early-life resilience and DA further extending late-life survival.

Sex significantly influenced mortality rates, with rank-order effects varying by selection regime

(Figure 2e-h, Supplementary Figure S2). To evaluate the impact of dietary selection on mortality

relative to the control (C) group, we computed hazard ratios using Cox proportional hazards

models accounting for sex. All selection regimes (CH, FA, DA) showed higher mortality risk

compared to controls (Table 1, Appendix Figure 1b, Supplementary Table S3), as indicated by

positive coefficients. The DA coefficient was marginally significant (p = 0.052), suggesting that

while DA approached control-like outcomes, it achieved them via potentially distinct adaptive

strategies. Notably, males in the DA and FA regimes exhibited nearly identical survival

trajectories (Appendix Figure 2c), suggesting that despite differing selection pressures, they

underwent similar adaptive responses to diet stress, resulting in comparable survival outcomes.

Sex-specific differences within regimes were evident and suggested interactions between sex

and selection treatment (Figure 2, Appendix Figure 2). Modeling survival as a function of

regime-by-sex interaction revealed significant effects in most categories, except for C males and

DA males (Table 1, Supplementary Table S3). CH flies displayed significantly high mortality risk

for both sexes (p < 0.0001). Overall, these results highlight significant adaptation in DA and FA

regimes but not in CH, with sex playing a critical role in shaping survival responses.

Table 1: Patterns of mortality risk in lifespan adaptation to diet selection.

Variable	Coeff.	HR[1]	95% CI[1]	p-value
Regime + strata(sex)				
CH	0.99	2.69	2.29, 3.17	<2e-16
DA	0.16	1.18	1.00, 1.39	0.0515
FA	0.40	1.48	1.26, 1.74	<9.16e-07
Sex				
M	0.23	1.26	1.15, 1.39	<1.68e-06
Regime * sex interaction				
C * F	-0.64	0.53	0.42, 0.66	3.33e-08
CH * F	0.44	1.56	1.31, 1.86	6.52e-07
DA * F	-0.50	0.61	0.51, 0.73	9.51e-08
FA * F	-0.18	0.84	0.70, 1.00	0.0457
C * M	-0.22	0.81	0.58, 1.12	0.1941
CH * M	0.49	1.64	1.33, 2.02	4.67e-06
DA * M	-0.02	0.98	0.79, 1.21	0.8501
FA * M	—	—	—	—

[1]HR = Hazard Ratio, CI = Confidence Interval

Microbial composition across selection regimes and time points

To assess microbial composition, we analyzed the average relative abundance of phyla across

time points and selection regimes (Figure 3). Proteobacteria and Firmicutes strongly dominated

microbial composition in all samples (Figs. 3, 4). Proteobacteria and Firmicutes were the most

abundant across all regimes but exhibited distinct temporal trends. In the control group (C), Fir-

micutes were initially low, increasing at the second time point before slightly decreasing at the

final time point (Figure 4a). In contrast, all selection treatments began with higher Firmicutes

levels and declined similar declines with age (Figure 4a). These results are also consistent with

the relative abundance profiles before being averaged (Supplementary Figure S3).

Figure 3: Phylum-level microbial relative abundance over time (t1-t3) for each of the four dietary regimes, including **a.** constant high sugar, **b.** decreasing nutrient availability, **c.** fluctuating nutrient availability, and **d.** control non-selected group.

Proteobacteria was most abundant in the control group, showing a decline at the second time

point followed by a partial recovery at the third (Figure 4b). In contrast, the FA regime had the

lowest initial Proteobacteria levels but exhibited a marked increase over time (Figure 4b). The

DA and CH regimes showed steady increases in Proteobacteria abundance as they aged (Fig-

ure 4a-b). Minor phyla, including Actinobacteria, Bacteroidota, and Fusobacteriota, were more

abundant in the selection treatments than in the control group but generally declined with age

(Figure 4c-d). These results highlight divergent microbial dynamics between control and selec-

tion regimes over time.

Based on prior studies linking microbial composition to immune function (Shen et al. 2018;

Gibbons et al. 2021), we hypothesized that Proteobacteria and Firmicutes could influence hosts

positively or negatively depending on their relative proportions. We calculated phyla ratios as

age-specific indicators of treatment effects (Figure 4f-g). In the selection regimes, the

Figure 4: Changes in the percentage of phyla over time (t1-t3) among dietary regimes (a-e): **a.** Firmicutes, **b.** Proteobacteria, **c.** Actinobacteriota, **d.** Bacteroidota, and **e.** Fusobacteriota. Phyla ratios over time between dietary treatments (f-g): **f.** Proteobacteria to Firmicutes, **g.** Bacteroidota to Firmicutes. Time points: t1, 15 days; t2, 35 days; and t3, 60 days.

Proteobacteria-to-Firmicutes ratio generally increased over time (Figure 4f), while the

Bacteroidota-to-Firmicutes ratio decreased between T1 and T2, except for DA increasing

between T2 to T3 (Figure 4g). The control group began with higher ratios than the treatments

and exhibited a general decline, suggesting an age-related increase in Firmicutes relative to

Proteobacteria and Bacteroidota. These patterns reveal contrasting dynamics between

Proteobacteria and Bacteroidota that are influenced by Firmicutes.

To explore functional significance of observed relative abundance dynamics, we performed an

indicator species analysis at the genus level. We identified 40 genera significantly different

across time points (Supplementary Table S4a) and 8 across selection regimes (Supplementary

Table S4b). Of the 40 time-related genera, 85% (34) were enriched at T1, one at T3, five at both

T1 and T3, and none at T2, indicating significant early-life microbial changes that may prime

later traits like lifespan. Among the 8 treatment-related genera, we found: *Fructilactobacillus* in

CH; *Mobiluncus* in FA; *Leptotrichia* in CH + FA; *Levilactobacillus*, *Acinetobacter*, and *Neisseria*

in CH + DA + FA; *Lactobacillus* in CH + FA + C; and *Halomonas* in DA + FA + C. This pattern

suggests a mix of treatment-specific and conserved functional effects across regimes.

Taxon diversity across time points and dietary treatments

To assess the effects of selection treatments on microbial diversity over time, we calculated

observed alpha diversity (ASV richness) and the Shannon-Wiener Index. ASV richness

significantly differed within DA samples across time points (Wilcoxon rank-sum: T1 vs. T2, *p* =

Figure 5: Alpha diversity between time points (t1-t3) across all dietary regimes. **a.** Observed diversity, **b.** Shannon-Weiner diversity. *P*-values between time points denote Wilcoxon rank-sum tests.

0.0001; T1 vs. T3, p = 0.0094; Figure 5a). The Shannon Index was significantly different between T1 and T2 in all regimes (CH: p = 0.0073; DA: p = 0.013; FA: p = 0.003; Figure 5b), but not between T2 and T3 or T1 and T3. These findings suggest selection-related diversity changes occur early in life, with CH and FA treatments lacking T3 data due to limited lifespan.

Figure 6: Beta diversity metric showing species diversity between time points (a-d) and across all dietary regimes (e-g). Note: For 6g, CH and FA groups did not reach t3, hence no comparisons.

We next examined between-group beta diversity using PCoA. Shifts in microbial composition were observed from T1 to T2 across treatments (Figure 6). At T3, the DA and C treatments diverged significantly. PERMANOVA confirmed significant differences by selection regime (r^2 =

0.12061, $p = 0.001$) and time point ($r^2 = 0.11005$, $p = 0.001$). these differences were due to

group means rather than dispersion (PERMDISP; CH vs. C: $p = 0.015$; DA vs. C: $p = 0.003$; FA

vs. C: $p = 0.001$). Significant differences were also observed between T1 and later time points

(T1 vs. T2: $p = 0.001$; T1 vs. T3: $p = 0.001$).

Discussion

We hypothesized that adaptation to stressful diet environments selects for distinct strategies in

survival, microbial community composition, and richness. To test this, we compared various se-

lection treatments against an unselected progenitor population (DSPR) from which the SILs

were derived. Although none of the selection treatments surpassed the control in longevity, the

DA treatment nearly matched it by generation 65. It is important to note that, after experimental

evolution, all selected lines were reared in a common nutritional environment - different from the

diets used in the selection regimes - which may influence the patterns we report. Nonetheless,

our previous studies have characterized key fitness traits (lifespan and fecundity) in component

diets across inbred (Stanley et al. 2017), outbred (Ng'oma et al. 2019, 2020), and evolved

(Ng'oma et al., unpublished) DSPR lines, and the patterns we observe here are consistent with

those findings. For instance, the DR diet prolonged lifespan by 37.5% in DSPR RILs (Stanley et

al. 2017), while the HS diet markedly reduced both lifespan and fecundity (Ng'oma et al. 2019).

These outcomes align with our observations: the DA selection treatment, which provided longer

exposure to the DR diet each generation, was associated with an extended lifespan, and similar

effects were seen in the CH treatment where a high-sugar environment was maintained. To-

gether, these findings support the expectation of persistent evolved patterns in response to

stressful dietary environments.

Effect of diet selection on longevity

Notwithstanding the context outlined above, the CH treatment was particularly detrimental,

especially for females, suggesting either incomplete adaptation or weak selection pressure, with

the latter likely contributing to slower evolution by depleting beneficial alleles. Where new

mutations are expected, stronger selection should lead to faster adaptation as it would favor

fitter mutation-genotypes to survive (Hartl and Clark 2007; Stephan 2021). In this study, we

expect negligible mutation-driven evolution. Rather, genetic variation should primarily be

maintained by recombination. In this case, stronger selection can lead to slower evolution, as

intense selection can rapidly deplete beneficial alleles leaving few options for further adaptation

(Ueda et al. 2017). This is especially possible in a complex trait like longevity where beneficial

combinations may be disrupted by recombination. Thus, reduced survival in the CH regime

could indicate greater levels of metabolic stress imposed by the high sugar diet such that

reproduction is severely depressed early, around the peak reproductive age (the first 3 weeks

post-eclosion) as genetic variation for longevity is removed with those that die early for multiple

other reasons.

The DA and FA treatments closely paralleled the control in longevity, with higher survival around

T1 (15 days pe, Figure 2a), reducing early mortality and maintaining lifespan variation. It should

be noted that males in the CH regime experienced significantly reduced mortality in the interval

around median lifespan (Figure 2b), but this advantage was probably late to significantly

improve median survival.

Sex-based effects of selection on longevity

Sex-specific diet responses, which are typically expected in *Drosophila*, revealed differing

evolved strategies. First, the FA selection shows a pattern of greater male survival earlier and

lower survival later which may support sustained reproduction in younger females and incurring

lower survival (Figure 2f). This pattern is less pronounced in the C regime (Figure 2h). Second,

males of the DA treatment survived poorly earlier in life compared to females which can lead to

observed reduced female reproduction, potentially necessitated by cues of a worsening

nutritional environment. These patterns suggest that DA and FA selection set different survival

patterns in early life. Longevity analysis suggests that adaptation to stressful diet alternations

may occur by reducing early mortality (DA, FA) or balancing higher early-life mortality with lower

mortality later in life.

Effect of diet selection on microbial community composition

Given that flies acquire microbiotas through mother-larvae transmission and from their diet

medium (Fischer et al. 2017; Guilhot et al. 2023), we anticipated selection-specific host-

microbiota interactions under a common nutritional environment. The DA regime was expected

to enrich longevity-promoting, low-inflammatory taxa, while the FA treatment, with its cyclical

resource availability, was predicted to shift the microbiota toward reproductive prioritization, with

increased pro-inflammatory taxa. In contrast, the CH treatment was projected to maintain a

stable but less diverse microbiome, indicating selection against resource storage or stress

resilience traits. Additionally, we expected age-related microbial shifts to interact with these

regimes, resulting in distinct survival strategies. These expectations were generally conformed

as microbial richness shifted significantly with selection and fly age, although selection regimes
differed from the control group.

The microbiome composition of the selection regimes differed markedly from the control group.
Survival patterns across dietary regimes were strongly aligned with distinct microbial richness
and diversity. The longest-living group, C, exhibited lower richness in Firmicutes and higher
richness in Proteobacteria compared to the selection regimes, particularly early in life post-
eclosion. This aligns with prior studies showing that fruit fly microbiomes are primarily
dominated by Firmicutes and Proteobacteria (Staubach et al. 2013; Adair et al. 2018; Lee et al.
2019). The dietary selection treatments likely contributed to the increased number of phyla
observed in our study.

One explanation for this increase is the colonization of pathogenic groups in the selection
regimes, leading to evolutionary stress and dysbiosis. Studies have shown that early increases
in pathogenic abundance can inhibit the development of other microbiota, disrupting the
protective roles of beneficial microbes (Spragge et al. 2023). For example, host microbiomes in
human and mouse guts can compete with pathogens for nutritional resources, with niche
exclusion being slower if pathogenic species establish dominance early (Spragge et al. 2023).
Additionally, compositional differences in the microbiome may be influenced by lab and strain
effects. Inconsistencies in fruit fly microbiomes are often attributed to dietary variation (Arias-
Rojas and Iatsenko 2022) and microbial load (Lee et al. 2019), both of which impact lifespan.
For instance, Lee et al. (2019) reported increased lifespan in lab strains with higher colonization
of commensal bacteria. Keebaugh et al. (2019) found that diet composition, such as the addition
of nutritional yeast, altered microbial load and lifespan outcomes by increasing environmental
microbe diversity. Conversely, other studies found no impact of microbial load on lifespan (Ren
et al. 2007). These findings highlight the complex interplay between diet, microbiome
composition, and host state, particularly the role of pathogenic colonization and dietary
interventions in shaping microbiota and their effects on lifespan.

Microbial selection effects depend on host stage
Adaptation strategies in response to dietary treatments appear to favor the optimization of
microbial community dynamics in younger host stages. For selection regimes that facilitated
more flies surviving to old age, such as DA and FA (to a lesser extent), maintaining microbial
composition in older flies seemed crucial for survival. In past research (Je et al. 2013; Na et al.

2014), the early period is important for establishment and proper succession of microbial

richness in adult flies. These patterns observed in microbiome analysis align with the observed

patterns in survival. For instance, selection treatments associated with longer lifespan (C, DA,

FA) all reduced early mortality compared with the least longevity treatment, CH (discussed

earlier). Thus, adaptation strategies to these treatments (especially longevity) likely optimizes

microbial community dynamics in younger host stages. For selection regimes that facilitated

more flies to survive to old age (i.e. DA, and FA to a lesser extent), it seems that the ability to

adapt depended on maintenance of microbial composition at older age.

Major microbial groups coordinated by diet selection

The dominant phyla identified in this study were Proteobacteria, Firmicutes, Actinobacteria,

Bacteroidota, and Fusobacteriota. Among these, Proteobacteria and Firmicutes were most

abundant across all regimes, though they exhibited distinct temporal trends. Previous research

has established correlations between certain phyla and immune responses. For instance, Shen

et al. (2018) reported a positive association between Bacteroidota (formerly Bacteroidetes) and

both immunoglobulin G and cytotoxic T cell production as well as a positive correlation of the

ratio of Bacteroidetes to Firmicutes and immunoglobulins G and M. Similarly, Gibbons et al.

(2021) observed increased cytokine and immunoglobulin production linked to the ratio of

Proteobacteria to Firmicutes.

In our study, the Bacteroidota-to-Firmicutes ratio generally decreased over consecutive time

points, suggesting that as fruit flies age, they may experience an increase in antibody

production but a reduction in cytotoxic T cell activity. Additionally, the Proteobacteria-to-

Firmicutes ratio declined in non-selected flies but increased in dietary regime flies, pointing to

potential differences in immune responses across treatments. However, none of the dietary

regimes exhibited levels of these ratios as high as the non-selected group, suggesting that the

dietary regimes negatively impacted immune response and survival in adult flies.

Notably, non-selected flies began with exceptionally high levels of Proteobacteria, a core

microbiome member in fruit flies alongside Firmicutes (Staubach et al. 2013; Wong et al. 2013;

Adair et al. 2018; Lee et al. 2019; Chiang et al. 2022). The dietary regime flies, on the other

hand, appeared to experience dysbiosis, potentially linked to their increased microbial diversity.

This imbalance may negatively affect immune function, gut health, and overall host health.

Specifically, higher levels of Firmicutes in the selection regime flies could contribute to reduced

immune responses and poorer survival outcomes compared to non-selected flies. These

findings highlight the complex interplay between microbiome composition, immune function, and

host survival, with Firmicutes emerging as a key player in shaping these dynamics.

Selection and dynamics of Proteobacteria and Bacteriodota

Our results reveal an inverse relationship between Proteobacteria and Bacteroidota in our

selection groups, a pattern that likely has important implications for host fitness. Overgrowth of

Proteobacteria has been associated with metabolic disturbances and inflammation

(Mukhopadhya et al. 2012; Shin et al. 2015), whereas Bacteroidota are generally linked to

healthier microbiomes (Zafar and Saier Jr 2021). The lower abundance of Proteobacteria in our

selected flies relative to C may indicate antagonistic interactions that contribute to their reduced

lifespan under nutritional stress, but not under normal diet. Similar inverse dynamics have been

reported in mice, where high-fat diets and inflammatory conditions favor Proteobacteria at the

expense of Bacteroidota (Bradley and Pollard 2017; Méndez-Salazar et al. 2018). Moreover, the

ability of certain probiotics to modulate this ratio (Gryaznova et al. 2022) supports a mechanistic

link between microbial balance, dietary factors, and host survival. Collectively, these

observations suggest that shifts in microbiome composition could play a critical role in mediating

the effects of our selection regimes.

Indicator species analysis

Our indicator species analysis revealed significant structural changes in the microbiome early in

adulthood across all treatments, with 34 of 40 microbial groups undergoing turnover at T1,

indicating early succession dynamics during microbiome establishment. This aligns with prior

research showing that *Drosophila melanogaster* larvae and pupae harbor rich environmental

microbial communities, but newly eclosed adults acquire minimal microbiota, establishing their

communities through food sources shortly after emergence (Blum et al. 2013; Broderick et al.

2014). In contrast, only eight microbial groups were enriched in at least one treatment,

suggesting relative stability in selection-specific microbiomes. These findings highlight the role

of stage-specific microbial shifts and conserved dynamics in long-term dietary adaptation, where

early-life succession and later stability influence host physiology and traits such as lifespan.

Similarly, prior studies emphasize early adulthood as a critical period for establishing microbial

diversity and longevity in flies (Brummel et al. 2004). Consistent with this, our diversity

measures showed increased microbiome diversity early in adulthood, particularly in the

selection regime treatments. However, members of some bacterial genera identified in our
analysis, such as *Porphyromonas* and *Atopobium*, are associated with inflammation and
disease in humans (Caselli et al. 2019; Salliss et al. 2021). An exception is *Lactobacillus*,
reported to inhibit these harmful groups (Moses et al. 2015; Mendi et al. 2016).
These results suggest that selection regime flies may have been exposed to harmful bacteria
throughout adulthood, contributing to inflammation and reduced survival. While the DA group
sustained more survivors, the other two dietary regimes appeared less effective at managing
infections. This underscores the complex interplay between microbiome dynamics, immune
function, and longevity, shaped by temporal and treatment-specific factors.
514
Although our design resulted in fewer flies in the CH and FA treatments reaching T3, our
findings reveal that adaptation to stressful dietary environments leads to distinct microbial
community dynamics, sex-specific survival patterns, and longevity outcomes. Selection regimes
such as DA and FA improved early survival and maintained greater lifespan variation compared
to the CH regime, which exhibited high early mortality, particularly in females. These patterns
underscore the complex interplay between dietary stress, genetic adaptation, and microbial
composition. Microbiome analysis highlights significant shifts during early adulthood, consistent
with the critical role of this period in microbiome establishment and its influence on longevity.
Selection regimes increased microbial diversity but also introduced pathogenic taxa,
contributing to dysbiosis and reduced survival compared to the control group. The inverse
relationship between Proteobacteria and Bacteroidota, particularly in selection regimes, may
reflect antagonistic microbial interactions linked to dietary stress during adaptation. These
findings underscore the importance of both stage-specific microbial succession and long-term
microbial community stability in shaping host fitness and survival.
529
530 Overall, our study provides new insights into the evolutionary and ecological dynamics of diet-
531 microbiome interactions, highlighting how dietary stressors can shape host-microbiome
532 relationships and influence complex traits like longevity. Future research should explore the
533 mechanistic links between microbial composition, immune function, and host physiology to
534 better understand the adaptive responses to dietary perturbations.
535
536
References

Adair, K. L., M. Wilson, A. Bost, and A. E. Douglas. 2018. Microbial community assembly in wild

populations of the fruit fly Drosophila melanogaster. ISME J 12:959–972. Nature

Publishing Group.

Aleman, F. D. D., and D. R. Valenzano. 2019. Microbiome evolution during host aging. PLOS

Pathogens 15:e1007727. Public Library of Science.

Arias-Rojas, A., and I. Iatsenko. 2022. The role of microbiota in Drosophila melanogaster aging.

Frontiers in Aging 3.

Bass, T. M., R. C. Grandison, R. Wong, P. Martinez, L. Partridge, and M. D. W. Piper. 2007.

Optimization of dietary restriction protocols in Drosophila. J Gerontol A Biol Sci Med Sci

62:1071–1081.

Blum, J. E., C. N. Fischer, J. Miles, and J. Handelsman. 2013. Frequent replenishment sustains

the beneficial microbiome of Drosophila melanogaster. mBio 4:10.1128/mbio.00860-13.

American Society for Microbiology.

Bradley, P. H., and K. S. Pollard. 2017. Proteobacteria explain significant functional variability in

the human gut microbiome. Microbiome 5:36.

Broderick, N. A., N. Buchon, and B. Lemaitre. 2014. Microbiota-induced changes in Drosophila

melanogaster host gene expression and gut morphology. mBio 5:10.1128/mbio.01117-

14. American Society for Microbiology.

Brummel, T., A. Ching, L. Seroude, A. F. Simon, and S. Benzer. 2004. Drosophila lifespan

enhancement by exogenous bacteria. Proceedings of the National Academy of Sciences

101:12974–12979. Proceedings of the National Academy of Sciences.

Cáceres, M. D., and P. Legendre. 2009. Associations between species and groups of sites:

indices and statistical inference. Ecology 90:3566–3574.

Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes.

2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nat

Methods 13:581–583. Nature Publishing Group.

Caselli, E., I. Soffritti, M. D'Accolti, I. Piva, P. Greco, and G. Bonaccorsi. 2019. Atopobium

vaginae And Porphyromonas somerae Induce Proinflammatory Cytokines Expression In

Endometrial Cells: A Possible Implication For Endometrial Cancer? Cancer Manag Res

11:8571–8575.

Chiang, M.-H., S.-M. Ho, H.-Y. Wu, Y.-C. Lin, W.-H. Tsai, T. Wu, C.-H. Lai, and C.-L. Wu. 2022.

Drosophila model for studying gut microbiota in behaviors and neurodegenerative

diseases. Biomedicines 10:596.

Christensen, S. M., S. N. Srinivas, Q. S. McFrederick, B. N. Danforth, S. L. Buchmann, and R.

L. Vannette. 2024. Symbiotic bacteria and fungi proliferate in diapause and may

enhance overwintering survival in a solitary bee. The ISME Journal 18:wrae089.

de Weerth, C. 2017. Do bacteria shape our development? Crosstalk between intestinal

microbiota and HPA axis. Neuroscience & Biobehavioral Reviews 83:458–471.

Douglas, A. E. 2018. The Drosophila model for microbiome research. Lab Anim (NY) 47:157–

164.

Fischer, C. N., E. P. Trautman, J. M. Crawford, E. V. Stabb, J. Handelsman, and N. A.

Broderick. 2017. Metabolite exchange between microbiome members produces

compounds that influence Drosophila behavior. eLife 6:e18855. eLife Sciences

Publications, Ltd.

Gibbons, J., J. Y. Yoo, T. Mutka, M. Groer, and T. T. B. Ho. 2021. A pilot study to establish an in

vitro model to study premature intestinal epithelium and gut microbiota interactions.

mSphere 6:10.1128/msphere.00806-21. American Society for Microbiology.

Gnainsky, Y., N. Zfanya, M. Elgart, E. Omri, A. Brandis, T. Mehlman, M. Itkin, S. Malitsky, J.

Adamski, and Y. Soen. 2021. Systemic Regulation of Host Energy and Oogenesis by

Microbiome-Derived Mitochondrial Coenzymes. Cell Reports 34:108583.

Gordon, M. 2021. How to use the timeSplitter. https://cran.r-

project.org/web/packages/Greg/vignettes/timeSplitter.html.

Gould, A. L., V. Zhang, L. Lamberti, E. W. Jones, B. Obadia, N. Korasidis, A. Gavryushkin, J. M.

Carlson, N. Beerenwinkel, and W. B. Ludington. 2018. Microbiome interactions shape

host fitness. Proceedings of the National Academy of Sciences 115:E11951–E11960.

Proceedings of the National Academy of Sciences.

Gryaznova, M., Y. Dvoretskaya, I. Burakova, M. Syromyatnikov, E. Popov, A. Kokina, E.

Mikhaylov, and V. Popov. 2022. Dynamics of changes in the gut microbiota of healthy

mice fed with lactic acid bacteria and bifidobacteria. Microorganisms 10:1020.

Guilhot, R., A. Xuéreb, A. Lagmairi, L. Olazcuaga, and S. Fellous. 2023. Microbiota acquisition

and transmission in *Drosophila* flies. iScience 26:107656.

Hartl, D. L., and A. G. Clark. 2007. Principles of Population Genetics. Fourth Edition. Oxford

University Press, Oxford, New York.

Je, B., F. Cn, M. J, and H. J. 2013. Frequent replenishment sustains the beneficial microbiome

of Drosophila melanogaster. mBio 4. mBio.

Jones, A., D. Donnell, E. G. King, and E. Ng'oma. 2024. Larval protein restriction interacts with

adult diet to increase fitness in an outbred multiparent population. bioRxiv.

Keebaugh, E. S., R. Yamada, and W. W. Ja. 2019. The nutritional environment influences the

impact of microbes on Drosophila melanogaster life span. mBio 10:10.1128/mbio.00885-

19. American Society for Microbiology.

King, E. G., S. J. Macdonald, and A. D. Long. 2012. Properties and power of the Drosophila

Synthetic Population Resource for the routine dissection of complex traits. Genetics

191:935–949.

Kueneman, J. G., M. C. Bletz, V. J. McKenzie, C. G. Becker, M. B. Joseph, J. G. Abarca, H.

Archer, A. L. Arellano, A. Bataille, M. Becker, L. K. Belden, A. Crottini, R. Geffers, C. F.

B. Haddad, R. N. Harris, W. M. Holden, M. Hughey, M. Jarek, P. J. Kearns, J. L. Kerby,

J. Kielgast, A. Kurabayashi, A. V. Longo, A. Loudon, D. Medina, J. J. Nuñez, R. G. B.

Perl, A. Pinto-Tomás, F. C. E. Rabemananjara, E. A. Rebollar, A. Rodríguez, L. Rollins-

Smith, R. Stevenson, C. C. Tebbe, G. Vargas Asensio, B. Waldman, J. B. Walke, S. M.

Whitfield, K. R. Zamudio, I. Zúñiga Chaves, D. C. Woodhams, and M. Vences. 2019.

Community richness of amphibian skin bacteria correlates with bioclimate at the global

scale. Nat Ecol Evol 3:381–389.

Lee, H.-Y., S.-H. Lee, J.-H. Lee, W.-J. Lee, and K.-J. Min. 2019. The role of commensal

microbes in the lifespan of Drosophila melanogaster. Aging (Albany NY) 11:4611–4640.

Leitão-Gonçalves, R., Z. Carvalho-Santos, A. P. Francisco, G. T. Fioreze, M. Anjos, C. Baltazar,

A. P. Elias, P. M. Itskov, M. D. W. Piper, and C. Ribeiro. 2017. Commensal bacteria and

essential amino acids control food choice behavior and reproduction. PLOS Biology

15:e2000862. Public Library of Science.

Martino, M. E., P. Joncour, R. Leenay, H. Gervais, M. Shah, S. Hughes, B. Gillet, C. Beisel, and

F. Leulier. 2018. Bacterial Adaptation to the Host's Diet Is a Key Evolutionary Force

Shaping Drosophila-Lactobacillus Symbiosis. Cell Host Microbe 24:109-119.e6.

McKnight, D. T., R. Huerlimann, D. S. Bower, L. Schwarzkopf, R. A. Alford, and K. R. Zenger.

2019. Methods for normalizing microbiome data: An ecological perspective. Methods in

Ecology and Evolution 10:389–400.

McMurdie, P. J., and S. Holmes. 2013. phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. PLOS ONE 8:e61217. Public Library

of Science.

Méndez-Salazar, E. O., M. G. Ortiz-López, M. de los Á. Granados-Silvestre, B. Palacios-

González, and M. Menjivar. 2018. Altered Gut Microbiota and Compositional Changes in

Firmicutes and Proteobacteria in Mexican Undernourished and Obese Children. Front

Microbiol 9:2494.

Mendi, A., S. Köse, D. Uçkan, G. Akca, D. Yilmaz, L. Aral, S. E. Gültekin, T. Eroğlu, E. Kiliç, and

S. Uçkan. 2016. Lactobacillus rhamnosus could inhibit Porphyromonas gingivalis derived

CXCL8 attenuation. J Appl Oral Sci 24:67–75.

Mills, M. 2010. Introducing survival and event history analysis. SAGE.

Moses, R. P., H. Y. Dawood, T. Fashemi, R. N. Fichorova, A. B. Onderdonk, and H. S.

Yamamoto. 2015. Lactobacillus curbs inflammatory responses to atopobium vaginae.

Obstetrics & Gynecology 125:3S.

Mukhopadhya, I., R. Hansen, E. M. El-Omar, and G. L. Hold. 2012. IBD-what role do

Proteobacteria play? Nat Rev Gastroenterol Hepatol 9:219–230.

Na, B., B. N, and L. B. 2014. Microbiota-induced changes in drosophila melanogaster host gene

expression and gut morphology. mBio 5. mBio.

Ng'oma, E., W. Fidelis, K. M. Middleton, and E. G. King. 2019. The evolutionary potential of

diet-dependent effects on lifespan and fecundity in a multi-parental population of

Drosophila melanogaster. Heredity 122:582–594. Nature Publishing Group.

Ng'oma, E., P. A. Williams-Simon, A. Rahman, and E. G. King. 2020. Diverse biological

processes coordinate the transcriptional response to nutritional changes in a Drosophila

melanogaster multiparent population. BMC Genomics 21:84.

Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. Minchin, R. O'Hara, G. Simpson, P.

Solymos, M. Stevens, and H. Wagner. 2015. Vegan: community ecology package. R

package vegan, vers. 2.2-1.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: Analyses of Phylogenetics and Evolution in

R language. Bioinformatics 20:289–290.

Parish, A. J., D. W. Rice, V. M. Tanquary, J. M. Tennessen, and I. L. G. Newton. 2022. Honey

bee symbiont buffers larvae against nutritional stress and supplements lysine. The ISME

Journal 16:2160–2168.

Petersen, C., I. K. Hamerich, K. L. Adair, H. Griem-Krey, M. Torres Oliva, M. P. Hoeppner, B. J.

M. Bohannan, and H. Schulenburg. 2023. Host and microbiome jointly contribute to

environmental adaptation. The ISME Journal 17:1953–65.

Ren, C., P. Webster, S. E. Finkel, and J. Tower. 2007. Increased internal and external bacterial

load during *Drosophila* aging without life-span trade-off. Cell Metabolism 6:144–152.

Rowe, M., L. Veerus, P. Trosvik, A. Buckling, and T. Pizzari. 2020. The reproductive

microbiome: an emerging driver of sexual selection, sexual conflict, mating systems, and

reproductive isolation. Trends in Ecology & Evolution 35:220–234. Elsevier.

Rudman, S. M., S. Greenblum, R. C. Hughes, S. Rajpurohit, O. Kiratli, D. B. Lowder, S. G.

Lemmon, D. A. Petrov, J. M. Chaston, and P. Schmidt. 2019. Microbiome composition

shapes rapid genomic adaptation of Drosophila melanogaster. Proceedings of the

National Academy of Sciences 116:20025–20032. Proceedings of the National Academy

of Sciences.

Salliss, M. E., J. D. Maarsingh, C. Garza, P. Łaniewski, and M. M. Herbst-Kralovetz. 2021.

Veillonellaceae family members uniquely alter the cervical metabolic microenvironment

in a human three-dimensional epithelial model. npj Biofilms Microbiomes 7:1–11. Nature

Publishing Group.

Shen, X., J. Miao, Q. Wan, S. Wang, M. Li, F. Pu, G. Wang, W. Qian, Q. Yu, F. Marotta, and F.

He. 2018. Possible correlation between gut microbiota and immunity among healthy

middle-aged and elderly people in southwest China. Gut Pathogens 10:4.

Shin, N.-R., T. W. Whon, and J.-W. Bae. 2015. Proteobacteria: microbial signature of dysbiosis

in gut microbiota. Trends Biotechnol 33:496–503.

Skorupa, D. A., A. Dervisefendic, J. Zwiener, and S. D. Pletcher. 2008. Dietary composition

specifies consumption, obesity, and lifespan in Drosophila melanogaster. Aging Cell

7:478–490.

Spragge, F., E. Bakkeren, M. T. Jahn, E. B. N. Araujo, C. F. Pearson, X. Wang, L. Pankhurst,

O. Cunrath, and K. R. Foster. 2023. Microbiome diversity protects against pathogens by

nutrient blocking. Science 382:eadj3502. American Association for the Advancement of

Science.

Stanley, P. D., E. Ng'oma, S. O'Day, and E. G. King. 2017. Genetic dissection of nutrition-
induced plasticity in insulin/insulin-like growth factor signaling and median life span in a
Drosophila multiparent population. Genetics 206:587–602.

Staubach, F., J. F. Baines, S. Künzel, E. M. Bik, and D. A. Petrov. 2013. Host species and
environmental effects on bacterial communities associated with Drosophila in the
laboratory and in the natural environment. PLOS ONE 8:e70749. Public Library of
Science.

Stephan, W. 2021. Rapid evolutionary adaptation in response to selection on quantitative traits.
Life 11:797.

Suyama, R., N. Cetraro, J. Y. Yew, and T. Kai. 2023. Microbes control Drosophila germline
stem cell increase and egg maturation through hormonal pathways. Commun Biol 6:1–
18. Nature Publishing Group.

Ueda, M., N. Takeuchi, and K. Kaneko. 2017. Stronger selection can slow down evolution
driven by recombination on a smooth fitness landscape. PLoS ONE 12:e0183120.

Wong, A., J. Chaston, and A. Douglas. 2013. The inconstant gut microbiota of Drosophila
species revealed by 16S rRNA gene analysis. The ISME journal 7. ISME J.

Wong, A. C.-N., A. J. Dobson, and A. E. Douglas. 2014. Gut microbiota dictates the metabolic
response of Drosophila to diet. Journal of Experimental Biology 217:1894–1901.

Zafar, H., and M. Saier Jr. 2021. Gut Bacteroides species in health and disease. Gut Microbes
13:1848158.

Zhang, F., J. L. Weckhorst, A. Assié, C. Hosea, C. A. Ayoub, A. S. Khodakova, M. L. Cabrera,
D. Vidal Vilchis, M.-A. Félix, and B. S. Samuel. 2021. Natural genetic variation drives
microbiome selection in the Caenorhabditis elegans gut. Current Biology, doi:
10.1016/j.cub.2021.04.046.

Appendix

Figure A1: Survival dynamics over time: **a.** Comparison of kernel density and parametric hazard function estimates. The black line represents the non-parametric kernel density estimate of the hazard function, while colored lines correspond to hazard functions estimated from various parametric survival models. The Generalized gamma and Gompertz fit the kernel estimate best suggesting a good fit to the data. **b.** Predicted hazard functions from a Gompertz survival model across different regimes. Each colored line represents the estimated hazard function for a specific regime, showing how the instantaneous risk of mortality changes over time. Diverging hazard trajectories suggest differences in survival patterns between regimes.

Figure A2: Hazard ratio estimates from Cox proportional hazards models evaluating the effect of selection regime and sex on survival. (a) Model stratified by sex, (b) Model with additive effects of regime and sex, (c) Model with interaction between regime and sex. Points represent hazard ratio (HR) estimates with 95% confidence intervals. HR > 1 indicates increased mortality risk relative to the reference category. Overall, these results suggest that dietary adaptation strongly influences survival, with `CH` and `FA` reducing lifespan relative to `C`, while `DA` showed a weaker effect. Males consistently exhibited higher mortality risk across regimes.