Antifungal biosynthesis by root-associated *Streptomyces* and *Pseudomonas* is elicited upon plant colonization

Wendalina Tigani[1], Chao Du[1], Somayah S. Elsayed[1], Paolo Innocenti[1], Victor Carrion-Bravo[1, 2], Iulian Rimboi[1], Nathaniel I. Martin[1], Jos M. Raaijmakers[3] and Gilles P. van Wezel[1,3, #]

[1] Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE, Leiden, The Netherlands.
[2] Department of Microbiology, University of Málaga, Málaga, Spain
[3] Department of Microbial Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, The Netherlands.

author for correspondence. Tel +31 715274310; g.wezel@biology.leidenuniv.nl

ABSTRACT

Plants are colonized by a diverse microbiome, with microorganisms residing inside and outside of plant tissues. Plants can harness the protective traits of their microbial inhabitants to ward off insect pests and fungal pathogens. However, current understanding of the role of commensal interactions on activating the desired microbial genomic traits remains limited. Here we show that biosynthesis of the antifungal (L)2-5 dihydro-phenylalanine (DHP) by the endophytic *Streptomyces* sp. PG2 is strongly induced upon colonization of *Arabidopsis thaliana*. This protects the plant from infection by the fungal root pathogen *Rhizoctonia solani*, both *in vitro* and *in vivo*. Interaction with sugar beets also enhanced DHP production in *Streptomyces* sp. PG2. We identified the DHP biosynthetic gene cluster (BGC) and showed that heterologous expression of the BGC in DHP non-producer *Streptomyces coelicolor* conferred plant-inducible DHP production. The BGC was also found in plant-associated Gram-negative bacteria, and in *Pseudomonas syringae* FF5 we again observed strongly enhanced DHP production upon plant colonization. Collectively, these data provide proof of concept that plant colonization can enhance the biosynthetic potential of root-associated microbe, thereby enhancing plant resilience. This may increase the competitiveness of the microbes within the rhizosphere.

INTRODUCTION

Plants are home to taxonomically and functionally diverse communities of bacteria, fungi, protists, nematodes and viruses, collectively referred to as the plant microbiome. These plant-associated microorganisms can provide various beneficial functions for their host by enhancing plant growth, facilitating nutrient acquisition, and bolstering resistance against pathogens[1]. Plants secrete compounds that actively shape the microbiome of the rhizosphere [2]. In return, plant growth-promoting bacteria (PGPB) suppress disease caused by microbial pathogens [3, 4] through several mechanisms, including competition for nutrients and iron, the production of lytic enzymes and the biosynthesis of bioactive molecules such as antibiotics or antifungals [5, 6]. Members of the genus *Streptomyces*, the largest genus within the Actinobacteria, are prolific producers of secondary metabolites, with bioactivities such as antibacterial, antifungal, antiviral, anticancer or nematicidal [7, 8]. Streptomycetes are prominent members of the plant microbiome, and play an important role in enhancing plant tolerance to biotic stresses[9-11]. However, most knowledge of their bioactivities has arisen from studies under controlled laboratory conditions, whereas genomic analyses point to a much larger biosynthetic potential[11-13]. Better understanding of the ecology of *Streptomyces*, their interactions with the plant host and the underlying chemistry is required to uncover and unleash their largely untapped functional potential[14].

Every step of the association between plants and microorganisms is tightly regulated by the exchange of specific signaling molecules. Plant-associated microorganisms perceive signals released by the plant in the root or shoot exudates and actively move toward the plant[15, 16]. Root exudate constituents are key for recruiting specific microorganisms from the diverse bulk soil microbial reservoir, resulting in distinct signatures of microbiome assembly for a given plant species or even cultivars within a plant species[17]. Plant signals can also trigger bacterial attachment and biofilm formation on the root surface[18]. Following colonization, plants may undergo metabolic changes, with the rhizosphere microbiome impacting root exudation profile[19], plant secondary metabolism[20] and

immunity (Induced Systemic Resistance, ISR) [6]. However, the reciprocal effects of the host plant on the chemistry and activities of the recruited microorganisms remain largely elusive.

According to the "cry for help" hypothesis [21], plants harness their microbiome to protect themselves against pathogens by eliciting the biosynthesis of bacterial secondary metabolites. Indeed, plant hormones have been shown to elicit the biosynthesis of bioactive compounds [11, 22]. We recently demonstrated *in vitro* studies that jasmonic acid (JA) and methyl jasmonate (MeJA) modify growth and trigger the production of antibiotics in streptomycetes [23], while the ubiquitous plant metabolite catechol also elicits natural product biosynthesis in streptomycetes [24]. However, how plants ensure that beneficial microbes in their microbiome produce disease-suppressive natural products in the rhizosphere and especially at the time of infection, remains a major question to be resolved.

We sought to investigate the ability of root-associated bacteria to interact with plants *in vivo* and protect against fungal infections. For this, we studied the ability of root-associated *Streptomyces* to protect *Arabidopsis thaliana* against infection with the fungal root pathogen *Rhizoctonia solani*. Using genetics, phylogenomics and metabolomics, we found that interaction with *A. thaliana* strongly enhances the biosynthesis of the antifungal (L)-2,5 dihydrophenylalanine (DHP) in the endophytic *Streptomyce*s PG2. DHP inhibited hyphal growth of *R. solani* and suppressed its infection of plants, both *in vitro* and *in vivo*. We observed remarkable conservation of DHP activation by plant colonization in among others *Pseudomonas*, with the inducible production of DHP conserved between different microbial taxa and plant species.

RESULTS

Colonization of *Arabidopsis thaliana* triggers antifungal activity by an endophytic streptomycete

To establish whether bioactive compounds are produced by bacteria upon root colonization, we screened a collection of streptomycetes isolated from the endophytic root compartment of *A. thaliana* for their ability to produce bioactive compounds inhibiting the growth of the fungal pathogen *Rhizoctonia solani*. Bioactivity was assessed in a dual culture assay by growing isolates of the collection on the edge of a petri dish on minimal medium. Five isolates were inoculated per petri dish, equidistant from one another, with *Streptomyces griseus* and *Streptomyces coelicolor* used as controls. After five days, *R. solani* was inoculated in the middle of the plates. Inhibition of fungal growth was measured after a further three days of incubation. Five out of 35 strains inhibited hyphal growth of *R. solani* **(Fig. S1)**. Following this, sterile seven-days-old *A. thaliana* seedlings were inoculated with *Streptomyces* spores and then moved to sterilized soil when colonization was visible **(Fig. 1a)**. After seven days, *R. solani* plugs were added to the pots just below the soil surface at of 0.5 cm from the stem base. *R. solani* infected sterile control plants, leading to plant death within 10 days. One of the isolates, *Streptomyces* sp. PG2, inhibited root infection by the fungus **(Fig. 1b).** The four other strains that showed bioactivity against *R. solani* during initial *in vitro* screening did not prevent plant infection **(Fig. S2)**, demonstrating the importance of *in vivo* validation. Based on this finding, *Streptomyces* sp. PG2 was selected for further investigation. To visualize the inhibition of fungal root infections in more detail, the experiment was repeated on agar plates. For this, seedlings inoculated with *Streptomyces sp.* PG2 spores, non-inoculated sterile *A. thaliana* seedlings, or *Streptomyces sp.* PG2 alone, were grown on half-strength MS agar plates, with an agar plug of *R. solani* inoculated in the centre of the plate. While plants or *Streptomyces* alone were readily overgrown by the fungal pathogen, a clear zone of hyphal growth inhibition of the fungus was observed around the plants inoculated with *Streptomyces sp.* PG2 **(Fig. 1c)**.

Plant colonization elicits the biosynthesis of DHP in *Streptomyces* sp. PG2

To characterize the antifungal activity, we investigated the metabolites released by *Streptomyces sp.* PG2 in liquid half-strength MS medium with or without *A. thaliana* seedlings. One-week old spent media were used to assay bioactivity on agar plates. Interestingly, while spent cell-free media from the bacterial cultures partially inhibited fungal growth, no inhibition was seen for the plant-derived media. However, antifungal activity was strongly enhanced in spent media obtained from plant-*Streptomyces* co-cultures **(Fig. 2a)**. These results suggested that the antifungal activity originated from *Streptomyces* sp. PG2, and that its biosynthesis was elicited only in the presence of *A. thaliana*. It is important to note that the plants did not show any signs of distress related to *Streptomyces* sp. PG2 colonization or antimicrobial production, remaining healthy throughout the duration of the experiment, when compared to the control group.

To elucidate the basis of the antifungal activity, the same extracts were analyzed by LC-MS. Comparison of the metabolomes of the different extracts revealed a significant correlation between a mass feature with an m/z 168.1019 and the level of bioactivity **(Fig. 2a-b)**. This mass feature corresponded to the bioactive compound (L)-2,5 dihydrophenylalanine (DHP)[25]. DHP was not present in extracts obtained from the plant alone and in only small quantities in the *Streptomyces* extracts, while it was strongly enriched in the extracts of the co-culture **(Fig. 2b)**. The antifungal activity of DHP likely arises from false feedback inhibition of the shikimate pathway as it serves as an analogue of the amino acid phenylalanine (Phe)[26]. DHP interferes with translation as it can be incorporated into proteins instead of Phe[27], which stops fungal growth via disruption of microtubule assembly[28].

To validate the activity of DHP towards *R. solani*, we used chemically synthesized DHP as a quantitative standard in LC-MS and showed that approximately 0.4 ng/ml (2.4 nM) of DHP was produced in the individual bacterial cultures and approximately 7 ng/ml (42 nM) in the co-culture of the bacteria and *A. thaliana*. Administration of the same concentrations of pure DHP to *R. solani* cultures reproduced the inhibition pattern described in **Fig. 2a**, confirming the sensitivity of the fungus to DHP at the concentrations measured in our samples **(Fig. 2c)**. To further validate that DHP was

indeed the sole antifungal agent, and that it acts by mimicking L-Phe, we supplemented the pure DHP samples with 2 mM L-Phe, which fully suppressed the antifungal activity in the extracts, strongly suggesting that L-Phe antagonized DHP and also that no other antifungals were produced, at least not at levels sufficient to inhibit growth of *R. solani* **(Fig. 2c)**. Similarly, L-Phe complementation of samples from pure bacterial/plant culture and bacterial-plant co-culture fully rescued the fungal growth **(Fig. 2d)**, suggesting that DHP was most likely solely responsible for the antifungal activity of the extracts.

Identification of the DHP gene cluster in *Streptomyces* sp. PG2

DHP production has been reported for *Streptomyces*[25, 29] and certain Gammaproteobacteria[30-32]. The DHP biosynthetic gene cluster (BGC) was identified for the Gammaproteobacterial *Erwinia amylovora* and *Photorhabdus luminescens*[30], but not in Actinobacteria. To identify the DHP BGC in *Streptomyces*, we obtained the whole genome sequence of *Streptomyces* sp. PG2, with 7 contigs and a genome size of 10.58 Mb. AntiSMASH[33] predicted 29 BGCs, but no BGC for DHP was detected **(Fig. S3)**. However, searching the PG2 genome with the DHP BGC from *E. amylovora* and *P. luminescens* as queries resulted in the identification of a putative BGC, of which five genes showed significant homology to genes of *E. amylovora* and *P. luminescens*. Of these, *plu3042* and *plu3043* encode a branched-chain amino acid aminotransferase IlvE and a prephenate decarboxylase, respectively, and were previously reported to be necessary for DHP production in *P. luminescens*[30]. The genes are preceded in the operon by the *plu3041* (*paaK*) gene, which encodes phenylacetate-coenzyme A ligase PaaK. The genomic environment of the loci and the conservation of the genes are shown in in **Fig. 3a.**

To ascertain that this gene cluster specifies the biosynthesis of DHP, a mutant was created whereby the orthologue of *plu3041* was replaced by double recombination with the apramycin resistance cassette *aac(C)3* using the unstable multi-copy plasmid pWHM3[34] (see Materials and Methods for details). This resulted in a mutant whereby nt positions -4 to +1326 bp relative to the start of gene TMLOC_04724 were replaced by the apramycin resistance cassette, which is expected to also transcriptionally block the downstream genes *plu3042-3043*. The resulting mutant was designated

PG2_Δ*plu3041*. The null mutant was then grown alone or in combination with *A. thaliana* seedlings and challenged with *R. solani*. Mutant PG2_Δ*plu3041* failed to inhibit the growth of the pathogen **(Fig. 3b)**. Metabolomic analysis of PG2_Δ*plu3041* confirmed that the mutant failed to produce any DHP, alone or in combination with *A. thaliana* **(Fig. S4a)**. Finally, to ensure that the phenotype owed solely to the mutation of gene TMLOC_04724 or the disruption of the expression of genes in this area, we genetically complemented the mutant, by introducing the whole gene cluster back into the knock-out mutant. For this we created a pSET152[35] derivative containing targeted region from TMLOC_04720 to TMLOC_04729, namely pE-DHP_Hyg. The plasmid was then introduced into PG2_Δ*plu3041* via conjugation. Genetic complementation restored DHP production in the mutant, enabling DHP production specifically in response to plant co-cultivation at levels equivalent to those of the wild-type strains **(Fig. S4a** and **S5b)** showing that indeed the BGC was solely responsible for DHP biosynthesis. As expected, the plant-dependent antifungal activity against *R. solani* was also restored by the genetic complementation of the BGC **(Fig. S6)**. Taken together, these data show that the BGC is required for production of DHP, and is required for inhibition of *R. solani* growth.

Exploring DHP elicitation by *Arabidopsis thaliana*

We next investigated the cues that could mediate DHP induction upon colonization. To investigate whether interaction with live plants is required, strain PG2 was grown in the presence of either spent media from axenic plants that contain metabolites released by the plant irrespective of colonization, or in the presence of dead plant material (following freezing). In both conditions, DHP production was not induced **(Fig. S5a)**. This excludes the possibility that DHP is elicited either by physical contact with inactive plants, or by plant metabolites produced in the absence of the bacterium. Since plant-microbe interactions often involve volatile compounds (VCs), we next conducted a split plate assay where *Streptomyces* sp. PG2 was separated from PG2-inoculated *A. thaliana* by a plastic barrier. In this way, the only way for the plant to activate DHP biosynthesis in the bacterium would be through the air, i.e. by producing eliciting VCs. However, *Streptomyces* sp. PG2 exposed to VCs released from PG2-

colonized *A. thaliana* did not produce more DHP than when the strain was exposed to the controls (sterile *A. thaliana*, *Streptomyces* sp. PG2 monoculture or MS medium without biomass, **Fig. S5a**). This demonstrates that VCs are not the (sole) causative agents of the DHP elicitation.

We then sought to investigate whether the elements required for recognition of the plant signals were conserved in the BGC itself. To test this, we introduced the plasmid pE-DHP_Hyg carrying the DHP BGC into *Streptomyces coelicolor* M145. This model streptomycete[36] does not have its own copy of the DHP gene cluster and does not produce DHP. Heterologous expression of the DHP BGC resulted in the production of DHP in *S. coelicolor*. This provided further evidence that the BGC is solely responsible for DHP production in *Streptomyces.* Excitingly, DHP levels were significantly enhanced when *S. coelicolor* was co-cultivated with *A. thaliana* **(Fig. S4b** and **S5b)**.

Finally, to explore whether plant colonization-dependent DHP elicitation was specific for *A. thaliana* or could also be mediated by colonization of other plant species, *Streptomyces* sp. PG2 was grown in association with sugar beet (*Beta vulgaris*). Again, DHP production was enhanced to a similar extent as during interaction with *A. thaliana* **(Fig. 4b)**. Collectively, these results strongly suggest that interaction with live plants is critical for DHP induction and that plant colonization-dependent elicitation of the BGC is conserved between streptomycetes, including in species that do not have the DHP BGC.

Colonization-dependent elicitation of DHP biosynthesis in other plant-associated bacteria

A search with three core genes of the DHP BGC against the RefSeq database[37] showed that around 1% (n=720) of all bacterial genomes (including WGS records) contain a likely DHP gene cluster. DHP gene clusters were found in the phyla Actinobacteria, Gammaproteobacteria and Betaproteobacteria. The major genera were *Erwinia* (230) and *Pseudomonas* (186) within the Gammaproteobacteria, *Streptomyces* (94) within the Actinobacteria and *Photorhabdus* (73) within the Betaproteobacteria. The diversity of the gene clusters was analyzed using BiG-SCAPE[38] **(Fig. 4a)**, with 0.3 clustering distance cut-off (default value). The DHP gene clusters found in *Streptomyces* genomes showed significant

diversity, separating into multiple network clusters, whereby a large number of clusters shared almost the same sequence (distance = 0). Many of the clusters were found in genomes obtained in a few "BioProjects" for comparative genome analyses, especially for the plant pathogen *Pseudomonas syringae*. Of note, most of these projects are collection of bacterial samples from plants **(Table S1)**, suggesting an association between the DHP BGC and plant-associated bacteria. However, further validation using an unbiased database is necessary to ascertain this connection.

DHP biosynthesis in conserved in plant-associated phyla such as *Pseudomonas*. We therefore wondered whether plant colonization-dependent production of DHP was also conserved in other phyla. Therefore, we analysed DHP production in the plant pathogen *Pseudomonas syringae* pv. *syringae* strain FF5. *P. syringae* FF5 harbors the DHP BGC, but so far has never been reported to produce DHP[30]. To see if expression of the DHP BGC of *P. syringae* FF5 could be elicited upon plant colonization, the strain was grown in MS medium in monoculture or in co-culture with *A. thaliana* seedlings. While DHP production by FF5 monocultures was extremely low, near background noise levels, the production of DHP was strongly enhanced in co-culture with the plant, showing up to 41-fold increase as compared to the monoculture **(Fig. 4b)**. Collectively, these findings reveal conservation of elicitation of the DHP gene cluster specifically upon plant colonization across bacteria belonging to different phyla.

DISCUSSION

Members of the genus *Streptomyces*, the largest genus of Actinobacteria, are well-known producers of a broad range of bioactive metabolites[11, 39]. Considering the significant potential of their antimicrobial arsenal in managing plant pathogens and their ability to colonize different plant species[9, 10, 40-42], understanding how to harness plant-microbe interactions for biocontrol of crops is crucial. Still, despite the exponential growth of microbial genome sequence information, our understanding of the

influence of host-microbe interactions on activating the desired genomic traits in the microbiome remains limited.

In this work, we show that plants stimulate the production of antimicrobials by members of their microbiome, resulting in protection against microbial pathogens **(Fig. 5)**. We identified the *Streptomyces* sp. PG2 as an efficient bioprotectant, producing the antifungal DHP specifically in response to plant colonization. Importantly, we also observed plant colonization-dependent induction of DHP production in *Pseudomonas*, showing that the concept is conserved between different bacterial phyla. Furthermore, introduction of the DHP BGC into *S. coelicolor* that lacks the BGC, led to plant-inducible DHP production, showing conservation of (i) perception of external cues; (ii) subsequent signal transduction; and (iii) regulation of DHP biosynthesis. Most of the experiments were done with *A. thaliana*. However, we also observed DHP stimulation upon colonization of other plant species, such as *Beta vulgaris*, possibly opening the door to broader application in crop protection.

Phylogenomic analysis revealed an intriguing correlation between the presence of the DHP BGC and a plant-associated lifestyle. We could only identify the BGC in the genomes of species of *Erwinia* (230), *Pseudomonas* (186), *Streptomyces* (94) and *Photorhabdus* (73), many of which are known to be plant-associated[11, 17, 43, 44]. Most of the DHP clusters from *Pseudomonas* where found in the genomes of bacteria obtained from plant samples, *Erwinia* is a genus of bacteria mostly composed of plant pathogenic species, for *Streptomyces* it is hard to draw a conclusion as for many strains the ecology is unknown, while *Photorhabdus* - best known as a symbiont of nematodes – was recently shown to respond to plant signals, contributing chitin degradation and inhibition of fungal growth[45]. *Photorhabdus* species indirectly interact with plant signals through their symbiotic relationship with obligate parasites of herbivorous insect larvae, and *Photorhabdus* and its symbiont *Heterorhabditis* are being studied as potential biocontrol agent[43, 46] for crop protection. We propose that both beneficial and pathogenic microbes may exploit DHP production to diversify their metabolic capabilities, thereby capitalizing on plant cues to enhance the resilience of their host or to improve their competitiveness within the rhizosphere, respectively.

The mechanism by which DHP biosynthesis is activated during plant colonization is yet unknown. Elicitation of DHP in *Streptomyces* or *Pseudomonas* requires interaction with living plants, as association with dead plant material did no elicit the production of DHP. Our work also shows that extracts from axenic plants fail to elicit DHP production, while at the same time elicitation could not be mediated by volatile compounds produced by the plant alone or by *Streptomyces*-colonized plants, as shown by split plate assays. This suggests that the activation of the DHP gene cluster requires more than the perception of a single signal, such as the provision or depletion by the plant for certain nutrients, or a joint action of microbe and plant. Heterologous expression of the gene cluster in the DHP nonproducer *S. coelicolor* showed that also in this case biosynthesis was activated in a plant-dependent manner, suggesting that the regulatory circuitry is located within the BGC itself. Regulatory networks coordinating critical cellular processes are typically preserved over evolutionary time scales and are often regulated by ancestral transcription factors[55]. The evolutionary conservation of the control of its biosynthesis indicates that DHP may play an important role in the lifestyle of plant-associated microbes. We recently showed that regulatory networks may be harnessed for the functional prediction of BGCs, using iron control as the basis for the discovery of novel BGCs involved in siderophore biosynthesis [47]. Therefore, elucidating how the plant-dependent activation of antifungal biosynthesis is governed will enable the search for BGCs associated with similar regulatory networks as candidates for roles in plant protection.

To the best of our knowledge, our work provides the first example of a natural product that is produced by microbes specifically in response to plant colonization, and whose stimulation is conserved both in beneficial and pathogenic plant-associated microbes. Deeper investigation into the chemical interplay regulating this phenomenon will be required to fully elucidate this inter-kingdom dialogue. Our study offers important new insights into the dynamics occurring amongst the several players populating the rhizosphere, and encourages further exploration of the biosynthesis of cryptic bioactive molecules specifically during plant-microbe interactions. This also offers great promise for novel approaches for the sustainable protection of crops against pathogens and pests.

MATERIALS & METHODS

Microbial strains and culturing conditions

The isolation of endophytic bacteria was employed as described [48]. Briefly, root tissue of *A. thaliana* ecotype mossel (Msl) obtained from a natural ecosystem was cleaned, sonicated and ground with mortar and pestle in 1 mL phosphate buffer (per liter: 6.33 g of $NaH_2PO_4·H2O$, 10.96 g of $Na_2HPO_4·2H_2O$ and 200 µL Silwet L-77). The plant material was spread onto the surface of a range of selective isolation media. Initial selection was done on media supplemented with the antifungal agent nystatin (50 µg/mL) and the antibacterial agent nalidixic acid (10 µg/mL). Plates were incubated at 30 °C for 4–25 days. Single strains were isolated based on morphology, and grown on soya flour mannitol (SFM) for seven days at 30 °C spores collected[49]. The 35 strains that were obtained were grouped under the name ATMOS collection. *Streptomyces coelicolor* A3(2) M145[36] was obtained from the John Innes centre strain collection and the host for heterologous expression experiments. The fungal plant pathogen *Rhizoctonia solani* was used in this work as an antifungal activity reporter. The fungus *R. solani* AG2-2IIIB was obtained from the collection of the Netherlands Institute of Ecology (NIOO-KNAW). Agar plugs with fungal hyphae used as inoculum in the bioassays were prepared by growing *R. solani* on 1/5th potato dextrose agar (PDA) medium for three days at 25°C. *Pseudomonas syringae pv syringae* strain FF5 was obtained from the IHSM-UMA-CSIC, Malaga, Spain.

Plant materials and growth conditions

Wild-type *Arabidopsis thaliana* Columbia-0 (Col-0) ecotype and *Beta vulgaris* (sugar beet) cultivar Camelia were employed in this study. *A. thaliana* seeds were obtained from Netherlands Institute of Ecology (NIOO-KNAW). Sugar beet seeds are from the Sugar beet Research Institute (IRS, The Netherlands), and were kindly provided by Victor Carrion Bravo group. The seeds were surface-sterilized by soaking for 10 min in 40 % bleach. After bleach removal they were rinsed five-seven times with sterile distilled water, and stored at 4 °C. Half-strength Murashige-Skoog (MS) medium was used

for plant growth. One liter of MS medium contains: 2.2 g MS powder (Duchefa), 10 g sucrose; the pH was adjusted to 5.7 and 10.8 g Micro Agar (Duchefa) was added for solid medium prior to autoclaving. Following the seed sterilization, all the plants in this work were germinated on half-strength MS plates for one week. After germination, seedlings for LC-MS analysis and bioassays were moved to liquid culture in Erlenmeyer flasks and incubated in shaking mode. For soil experiments, one-week-old seedlings were transferred from plates in liquid medium to 24 multi-well plates, incubated in shaking mode for one week and, eventually, moved on twice-autoclaved mixture of 9:1 substrate soil and sand (Holland Potgrond). *A. thaliana* and *J. vulgaris* on agar plates, liquid medium or soil, were cultivated at 21 °C, 16 h photoperiod, and 50% relative humidity; *B. vulgaris* was grown at 24 °C, a 16 h photoperiod, and 70% relative humidity.

Bioassays

All bioassays are presented in the **Supplemental Methods**. These include *in vitro* **bioassays**, namely: ATMOS collection screening; co-culture spent media; DHP bioactivity assays; and chemical complementation with L-Phenylalanine to chemically complement DHP and rescue the fungal growth; and *in vivo* **bioassays**, namely agar plate assays and in soil assays.

Analysis of secondary metabolites

Pure bacterial/plant cultures and plant-bacteria co-cultures were set up to analyze the differential metabolite production underlying the increased bioactivity of plant-bacterial co-culture spent media. Samples were prepared in triplicates. In brief, 15 one-week old seedlings were moved to flasks in 15 ml half-strength MS liquid medium and inoculated with 4.5 10^3 *Streptomyces* spores. For plant-*Pseudomonas* FF5 co-cultures, 50 µl of FF5 o/n culture adjusted to OD_{600} 0.5 in half-strength MS medium was added to the medium instead of *Streptomyces* spores. Pure bacterial cultures were prepared by adding the same amount of inoculum to sterile media. The flasks were incubated in shaking mode at 21 °C, 16 h photoperiod, and 50% relative humidity (*A. thaliana*), or at 24 °C, 16 h

photoperiod, and 70% relative humidity (*B. vulgaris*) for seven days. The experimental setup for the study of possible interaction via the production of **Volatile Compounds** (VCs) was as follows, using the same media. Spores from *Streptomyces* sp. PG2 were inoculated on one side of a petri dish and *A. thaliana* inoculated with PG2 was grown on the other side, with an impermeable sector dividing them in the middle. After 10 days of incubation, metabolites were extracted from the side with the *Streptomyces* sp. PG2 monoculture and checked for DHP.

For the study of soluble metabolites in liquid-grown cultures, supernatants were collected, 5% w/v Diaion® HP20 (Resindion) was added to the cultures and shaken for three h at room temperature. HP20 was filtered off the liquid media with glass wool, washed with distilled water, then soaked with 100% MeOH three times overnight. For extraction of either soluble and volatile metabolites from solid-grown cultures, the agar was chopped into very small pieces and submerged in 100% methanol three times overnight. MeOH extracts were dried and dissolved in 75% MeOH in H_2O to a final concentration of 1 mg/mL and filtrated (0.22 μm) for LC-MS analysis. Samples from sterile half-strength MS medium were similarly prepared and extracted to serve as negative controls.

LC-MS/MS acquisition was performed as described [24, 50]. A Shimadzu Nexera X2 UHPLC system was used, with attached PDA, coupled to a Shimadzu 9030 QTOF mass spectrometer, equipped with a standard ESI source unit, in which a calibrant delivery system (CDS) is installed. For details we refer to the Supplemental Methods.

Construction of the DHP knock-out mutant

The construct for gene disruption was obtained using a method published previously [51]. In brief, approximately 1.5 kb regions up- and downstream of the gene TMLOC_04724 from *S*. PG2 genomic DNA were amplified, using primer pairs 1+2 and 3+4, respectively (Table S2). The apramycin resistance cassette *aac(3)IV* flanked by *loxP* sites, was amplified with primer pair 5+6 from pWHM3-*oriT*[34]. The three DNA fragments were assembled into EcoRI-HindIII digested pWHM3 using NEBuilder® HiFi DNA Assembly. The obtained construct was verified by DNA sequencing, transformed in *E. coli*

ET12567/pUZ8002 [52, 53] and subsequently introduced into in *Streptomyces* PG2 by conjugation [49]. After double recombination, nt positions 5308194 to 5309523 of the PG2 chromosome were replaced by the apramycin resistance cassette, corresponding to nt positions -4 to +1326 bp relative to the start of gene TMLOC_04724. The correct mutant was selected by resistance to apramycin (50 µg/ml) and named PG2_Δ*plu3041*.

Expression of the DHP gene cluster in *Streptomyces*

For genetic complementation we cloned the genomic region between TMLOC_04720 - TMLOC_04729 into the conjugative vector pSET152[54], which harbours apramycin resistance cassette *aac(C)IV* as selection marker. For this, three contiguous fragments of this region were amplified with primer pairs 7+8, 9+10 and 11+12 (Table S2). The PCR products were assembled using NEBuilder® HiFi DNA Assembly into BamHI-EcoRI-digested pSET152. The plasmid was named pE-DHP and introduced via conjugation into *S. coelicolor* M145. Positive clones were selected on apramycin (50 µg/ml). The construct sequence was confirmed by restriction digestion and Sanger sequencing, and the obtained strain was named SCO-DHP. SCO-DHP was grown in liquid culture with *A. thaliana* seedlings as described for PG2 strain. Pure bacterial/plant cultures and plant-bacteria co-cultures were set up in triplicates as already described for the *in vitro* bioassays, followed by metabolite extraction and LC-MS run.

To ensure the specificity of the mutation in TMLOC_04724 in PG2_Δ*plu3041*, we reintroduced the entire BGC into the mutant. Since the mutant was already resistant to apramycin, a modified version of pE-DHP was created whereby the hygromycin resistance was inserted into the EcoRV site located inside the apramycin resistance gene *aac(C)IV*. The resulting plasmid pE-DHP_Hyg was then introduced via conjugation into PG2_Δ*plu3041*. The ability of the clone to restore DHP production to PG2_Δ*plu3041* was tested by growing the strain in liquid culture with *A. thaliana* seedlings as described for PG2 strain, followed by LC-MS analysis of the culture extracts.

Computational methods

Phylogenomic distribution of the DHP gene cluster

Genomes were analyzed using AntiSMASH (version 6.0.1)[33] to obtain BGC predictions. Program cblaster (version 1.3.18, 10.1093/bioadv/vbab016) was used for searching the DHP cluster against RefSeq database. Default settings of cblaster are used for BLAST hit. In general, for a protein to match query sequences, minimal identity is set to 30, minimal coverage is set to 50, e-value should not be higher than 0.01. To reduce the complexity of searches, query term "txid2[orgn]" is used to restrict the search within Bacteria superkindom. Predictions were used as input for BiG-SCAPE [55] (version 1.1.15, 10.1038/s41589-019-0400-9) with default clustering cutoff 0.3, for the creation of a sequence similarity network, with distance matrix cutoff set to 0.25. The resulting full network was visualized by Cytoscape (version 3.10.1) with default layout[56].

ACKNOWLEDGEMENTS

We are grateful to Victor Carrion Bravo for providing *Pseudomonas syringae* and sugar beet seedlings. The work was supported by grant OCENW.GROOT.2019.063 from the Netherlands Organization for Scientific Research (NWO) to JMR and GPvW.

REFERENCES

1. Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T.M. & Singh, B.K. Plant-microbiome interactions: from community assembly to plant health. *Nat Rev Microbiol* **18**, 607-621 (2020).
2. Hu, L. et al. Root exudate metabolites drive plant-soil feedbacks on growth and defense by shaping the rhizosphere microbiota. *Nat Commun* **9**, 2738 (2018).
3. Kloepper, J.W., Leong, J., Teintze, M. & Schroth, M.N. Enhanced plant growth by siderophores produced by plant growth-promoting rhizobacteria. *Nature* **286**, 885-886 (1980).
4. de Weger, L.A. et al. Siderophores and outer membrane proteins of antagonistic, plant-growth-stimulating, root-colonizing *Pseudomonas* spp. *J Bacteriol* **165**, 585-94 (1986).
5. Lugtenberg, B. & Kamilova, F. Plant-growth-promoting rhizobacteria. *Annu Rev Microbiol* **63**, 541-56 (2009).
6. Pieterse, C.M. et al. Induced systemic resistance by beneficial microbes. *Annu Rev Phytopathol* **52**, 347-75 (2014).

7.   Barka, E.A. et al. Taxonomy, physiology, and natural products of the Actinobacteria. *Microbiol Mol Biol Rev* **80**, 1-43 (2016).

8.   Bérdy, J. Bioactive microbial metabolites. *J Antibiot (Tokyo)* **58**, 1-26 (2005).

9.   Kinkel, L.L., Schlatter, D.C., Bakker, M.G. & Arenz, B.E. Streptomyces competition and co-evolution in relation to plant disease suppression. *Res Microbiol* **163**, 490-499 (2012).

10.   Newitt, J., Prudence, S., Hutchings, M. & Worsley, S. Biocontrol of Cereal Crop Diseases Using Streptomycetes. *Pathogens* **8**, 78 (2019).

11.   van der Meij, A., Worsley, S.F., Hutchings, M.I. & van Wezel, G.P. Chemical ecology of antibiotic production by actinomycetes. *FEMS Microbiol Rev* **41**, 392-416 (2017).

12.   Gavriilidou, A. et al. Compendium of specialized metabolite biosynthetic diversity encoded in bacterial genomes. *Nat Microbiol* **7**, 726-735 (2022).

13.   Nett, M., Ikeda, H. & Moore, B.S. Genomic basis for natural product biosynthetic diversity in the actinomycetes. *Nat Prod Rep* **26**, 1362-84 (2009).

14.   van Bergeijk, D.A., Terlouw, B.R., Medema, M.H. & van Wezel, G.P. Ecology and genomics of Actinobacteria: new concepts for natural product discovery. *Nat Rev Microbiol* **18**, 546-558 (2020).

15.   Tsai, A.Y.-L., Oota, M. & Sawa, S. Chemotactic Host-Finding Strategies of Plant Endoparasites and Endophytes. *Front Plant Sci* **11** (2020).

16.   Gupta Sood, S. Chemotactic response of plant-growth-promoting bacteria towards roots of vesicular-arbuscular mycorrhizal tomato plants. *FEMS Microbiol Ecol* **45**, 219-227 (2003).

17.   Berendsen, R.L., Pieterse, C.M.J. & Bakker, P.A.H.M. The rhizosphere microbiome and plant health. *Trends Plant Sci* **17**, 478-486 (2012).

18.   Kulkarni, O.S. et al. Volatile methyl jasmonate from roots triggers host-beneficial soil microbiome biofilms. *Nat Chem Biol* **20**, 473-483 (2024).

19.   Korenblum, E. et al. Rhizosphere microbiome mediates systemic root metabolite exudation by root-to-root signaling. *Proc Natl Acad Sci U S A* **117**, 3874-3883 (2020).

20.   Etalo, D.W., Jeon, J.S. & Raaijmakers, J.M. Modulation of plant chemistry by beneficial root microbiota. *Nat Prod Rep* **35**, 398-409 (2018).

21.   Rolfe, S.A., Griffiths, J. & Ton, J. Crying out for help with root exudates: adaptive mechanisms by which stressed plants assemble health-promoting soil microbiomes. *Curr Opin Microbiol* **49**, 73-82 (2019).

22.   Ramirez-Estrada, K. et al. Elicitation, an Effective Strategy for the Biotechnological Production of Bioactive High-Added Value Compounds in Plant Cell Factories. *Molecules* **21**, 182 (2016).

23.   van der Meij, A. et al. The plant stress hormone jasmonic acid evokes defensive responses in streptomycetes. *Appl Environ Microbiol* **89**, e0123923 (2023).

24.   Van Bergeijk, D.A. et al. The ubiquitous catechol moiety elicits siderophore and angucycline production in *Streptomyces*. *Comms Chem* **5**, 14 (2022).

25.   Scannell, J.P., Pruess, D.L., Demny, T.C., Williams, T. & Stempel, A. L-3-(2,5-dihydrophenyl)alanine, an antimetabolite of L-phenylalanine produced by a streptomycete. *J Antibiot (Tokyo)* **23**, 618-9 (1970).

26.   Genghof, D.S. 2,5-Dihydrophenylalanine as an Inhibitor of Microbial Growth. *Can J Microbiol* **16**, 545-& (1970).

27.   Pine, M.J. Incorporation of L-2,5-dihydrophenylalanine into cell proteins of *Escherichia coli* and sarcoma 180. *Antimicrob Agents Chemother* **7**, 601-605 (1975).

28.   Kiso, T., Fujita, K.I., Ping, X., Tanaka, T. & Taniguchi, M. Screening for microtubule-disrupting antifungal agents by using a mitotic-arrest mutant of *Aspergillus nidulans* and novel action of phenylalanine derivatives accompanying tubulin loss. *Antimicrob Agents Chemother* **48**, 1739-1748 (2004).

29.   Yamashita, T., Miyairi, N., Kunugita, K., Shimizu, K. & Sakai, H. L-1,4-cyclohexadiene-1-alanine, an antagonist of phenylalanine from *Streptomyces*. *J Antibiot (Tokyo)* **23**, 537-41 (1970).

30. Crawford, J.M., Mahlstedt, S.A., Malcolmson, S.J., Clardy, J. & Walsh, C.T. Dihydrophenylalanine: A Prephenate-Derived Photorhabdus luminescens Antibiotic and Intermediate in Dihydrostilbene Biosynthesis. *Chem Biol* **18**, 1102-1112 (2011).

31. Feistner, G.J. in Plant Pathogenic Bacteria: Proceedings of the Sixth International Conference on Plant Pathogenic Bacteria, Maryland, June 2–7, 1985 (eds. Civerolo, E.L., Collmer, A., Davis, R.E. & Gillaspie, A.G.) 509-519 (Springer Netherlands, Dordrecht, 1987).

32. Onishi, N., Watanabe, T., Izaki, K. & Takahashi, H. L-1,4-Cyclohexadiene-1-Alanine, an Antimetabolite of L-Phenylalanine Produced by a *Pseudomonas*. *J Antibiot (Tokyo)* **35**, 903-904 (1982).

33. Blin, K. et al. antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic Acids Res* **49**, W29-W35 (2021).

34. Vara, J., Lewandowska-Skarbek, M., Wang, Y.G., Donadio, S. & Hutchinson, C.R. Cloning of genes governing the deoxysugar portion of the erythromycin biosynthesis pathway in *Saccharopolyspora erythraea* (*Streptomyces erythreus*). *J Bacteriol* **171**, 5872-81 (1989).

35. Kim, M.K., Ha, H.S. & Choi, S.U. Conjugal transfer using the bacteriophage phiC31 att/int system and properties of the attB site in *Streptomyces ambofaciens*. *Biotechnol Lett* **30**, 695-9 (2008).

36. Hoskisson, P.A. & van Wezel, G.P. *Streptomyces coelicolor*. *Trends Microbiol* **27**, 468-469 (2019).

37. O'Leary, N.A. et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Research* **44**, D733-D745 (2016).

38. Navarro-Muñoz, J.C. et al. A computational framework to explore large-scale biosynthetic diversity. *Nat Chem Biol* **16**, 60-+ (2020).

39. Hutchings, M.I., Truman, A.W. & Wilkinson, B. Antibiotics: past, present and future. *Curr Opin Microbiol* **51**, 72-80 (2019).

40. Kim, D.-R. et al. A mutualistic interaction between *Streptomyces* bacteria, strawberry plants and pollinating bees. *Nat Comms* **10**, 4802 (2019).

41. Law, J.W.-F. et al. The Potential of *Streptomyces* as Biocontrol Agents against the Rice Blast Fungus, *Magnaporthe oryzae* (*Pyricularia oryzae*). *Front Microbiol* **8** (2017).

42. Schrey, S.D. & Tarkka, M.T. Friends and foes: streptomycetes as modulators of plant disease and symbiosis. *Antonie van Leeuwenhoek* **94**, 11-19 (2008).

43. Abd-Elgawad, M.M.M. Photorhabdus spp.: An Overview of the Beneficial Aspects of Mutualistic Bacteria of Insecticidal Nematodes. *Plants (Basel)* **10** (2021).

44. Kado, C.I. in The Prokaryotes: A Handbook on the Biology of Bacteria Volume 6: Proteobacteria: Gamma Subclass (eds. Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H. & Stackebrandt, E.) 443-450 (Springer New York, New York, NY, 2006).

45. Regaiolo, A., Dominelli, N., Andresen, K. & Heermann, R. The biocontrol agent and insect pathogen interacts with plant roots. *Appl Environ Microbiol* **86** (2020).

46. Askary, T.H. & Abd-Elgawad, M.M.M. Opportunities and challenges of entomopathogenic nematodes as biocontrol agents in their tripartite interactions. *Egypt J Biol Pest Co* **31** (2021).

47. Augustijn, H.E. et al. Prediction of gene cluster function based on transcriptional regulatory networks uncovers a novel locus required for desferrioxamine B biosynthesis. *BioRXiv*, 2024.06.10.598258 (2024).

48. van der Meij, A. et al. Inter- and intracellular colonization of *Arabidopsis* roots by endophytic actinobacteria and the impact of plant hormones on their antimicrobial activity. *Antonie Van Leeuwenhoek* **111**, 679-690 (2018).

49. Kieser, T., Bibb, M.J., Buttner, M.J., Chater, K.F. & Hopwood, D.A. Practical streptomyces genetics (John Innes Foundation Norwich, 2000).

50. Machushynets, N.V. et al. Discovery of actinomycin L, a new member of the actinomycin family of antibiotics. *Sci Rep* **12**, 2813 (2022).

51.   Swiatek, M.A., Tenconi, E., Rigali, S. & van Wezel, G.P. Functional analysis of the N-acetylglucosamine metabolic genes of *Streptomyces coelicolor* and role in the control of development and antibiotic production. *J Bacteriol* **194**, 1136-1144 (2012).

52.   Macneil, D.J. et al. Analysis of Streptomyces-Avermitilis Genes Required for Avermectin Biosynthesis Utilizing a Novel Integration Vector. *Gene* **111**, 61-68 (1992).

53.   Paget, M., Chamberlin, L.C.L., Atrih, A., Foster, S.J. & Buttner, M.J. Evidence that the Extracytoplasmic Function Sigma Factor SigE Is Required for Normal Cell Wall Structure in *Streptomyces coelicolor* A3(2). *J Bacteriol* **181**, 204 - 211 (1999).

54.   Bierman, M. et al. Plasmid cloning vectors for the conjugal transfer of DNA from *Escherichia coli* to *Streptomyces* spp. *Gene* **116**, 43-9. (1992).

55.   Navarro-Muñoz, J.C. et al. A computational framework to explore large-scale biosynthetic diversity. *Nat Chem Biol* **16**, 60-68 (2020).

56.   Shannon, P. et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* **13**, 2498-2504 (2003).

FIGURE LEGENDS

Figure 1. Plant-inducible suppression of *Rhizoctonia solani* infection by *Streptomyces* sp. PG2*. a) *A. thaliana* seedlings colonized by *Streptomyces* species. The colonization is visible as white spots on the root system. The magnification on the right shows the mycelium growing on the roots. **b)** *In vivo* bioassay in soil. Plants colonized by *Streptomyces* sp. PG2 (middle) are more resistant to infection by the fungal pathogen *R. solani* as compared to sterile plants (left) and those inoculated with a control endophytic *Streptomyces* that does not produce antifungals (right). **c)** Bioassay on agar plate. Top left, *A. thaliana* grown alone; top right, *Streptomyces* PG2 grown alone; bottom right, *A. thaliana* inoculated with *Streptomyces* PG2. *R. solani* was spotted in the middle and allowed to grow for 5 days. Note that while *R. solani* overgrows both the plant and the bacterium alone, it fails to grow near *A. thaliana* colonized by *Streptomyces* sp. PG2.

Figure 2. Identification of DHP as antifungal agent produced by *Streptomyces* sp. PG2. a) Bioassays against *R. solani* grown on media with extracts obtained from *A. thaliana* grown alone (left), *Streptomyces* PG2 grown alone (middle), or the plant-*Streptomyces* co-culture (right). Note that *R. solani* fails to grow on agar supplemented with extracts from the plant-*Streptomyces* coculture, while it grows well on extracts from either monoculture. The numbers correspond to the respective LC-MS profile in Panel b. **b)** Extracted ion chromatogram (EIC) of DHP (168.1019 m/z; chemical structure is shown) produced by *Streptomyces* PG2 grown as monoculture (trace 2) or in the presence of *A. thaliana* (trace 3); no peak was detected in samples from pure plant cultures (trace 1). **c)** Antifungal activity of chemically synthesized DHP on growth of *R. solani* and chemical complementation with L-Phe. Top, *R. solani* growth is impaired by increasing concentrations of chemically synthetized DHP, with low inhibition at 0.4 ng/ml and strong inhibition at 7 ng/ml DHP. The control (no DHP) is shown on the left. Bottom, chemical complementation with 2 mM L-Phe fully rescues fungal growth. DHP concentrations are the same as in the top row. **d)** Addition of L-Phe restores growth of *R. solani* in the

presence of extracts from liquid-grown cultures. Top row, same as in Panel a, with extracts from *A. thaliana* grown alone (left), *Streptomyces* grown alone (middle) or the co-culture (right). Bottom row, the same as in the top row, but with 2 mM L-Phe. Note that L-Phe again fully restores growth to *R. solani*. These experiments provide strong evidence that DHP is the antifungal agent that inhibits fungal growth.

Figure 3. DHP cluster conservation and mutational analysis. a) Alignment and conservation of DHP gene clusters identified in *Erwinia amylovora, Photorhabdus luminescens* and *Streptomyces* sp. PG2. The *plu3041-3043* operon is highlighted in blue. **b)** The DHP gene cluster is responsible of the antifungal activity. DHP Knock-out mutant PG2_Δ*plu3041* of *Streptomyces* sp. PG2 fails to inhibit growth of *R. solani*. Within 3 days of growth, the fungus overgrows the plant alone (top right), the mutant alone (top left) and the plant grown in co-culture with the mutant (bottom right). In line with the results shown in Figure 1b, the fungus cannot grow near the plant-*Streptomyces* co-culture due to production of antifungals (bottom left). Slight inhibition of the fungus is shown for *Streptomyces* sp. PG2 grown alone (left).

Figure 4. Phylogenomic analysis of DHP gene cluster distribution and plant-mediated elicitation of DHP biosynthesis. a) Network of DHP clusters from all bacterial genomes produced by BiG-SCAPE. DHP gene clusters were analyzed with a default cut-off (0.3) (BiG-SCAPE 10.1038/s41589-019-0400-9). Each node represents a single DHP gene cluster from one genome. The phyla are coded using node shapes, genera are coded using node colors. BiG-SCAPE calculated cluster distances are shown as different line darkness. PG2 genome was added to the collection before analysis. Singletons are not included in the graph. **b)** Fold-change in DHP production in several co-culture conditions compared to pure bacterial cultures. For *Streptomyces* PG2 grown in co-culture with *A. thaliana* (blue box), each data point represents the outcome of an experiment conducted in technical triplicates. For *Streptomyces* sp. PG2 with *Bacillus vulgaris* and *Pseudomonas* FF5 with *A. thaliana* (red and green boxes, respectively), the

experiment was run three times, whereby replicates belonging to the same experiments are represented with the same color. DHP quantification was performed via integration of the LC-MS peak.

Figure 5. Model for DHP gene cluster activation. (1) Root colonization; (2) the molecular cross-talk between the plant and its associated microbes results in the activation of DHP gene cluster; (3) DHP is released in the surroundings of the colonized host, and it prevents soil borne pathogens to reach the root system (4).

a

b

c

Sterile *A. thaliana*

Streptomyces PG2

R. solani

A. Thaliana + S. PG2

a

Plant culture	Bacterial culture	Co-culture
1	2	3

c

	No DHP	0.4 ng/ml	7 ng/ml
MS medium			
MS medium + L-Phe			

b

d

	Plant culture	Bacterial culture	Co-culture
MS medium			
MS medium + L-Phe			

a

b

a

PG2 DHP cluster

b