Title: Young Asian elephant calves show differentiated social relationships with conspecifics

Running title: Differentiated calf-conspecific social relationships in Asian elephants

Authors: T. Revathe[†] and T.N.C. Vidya

Affiliation: Evolutionary and Organismal Biology Unit, Jawaharlal Nehru Centre for
Advanced Scientific Research (JNCASR), Bengaluru, India.
Present address: [†]Development and Evolution of Cognition Group, Max Planck Institute of
Animal Behavior, Konstanz, Germany

Corresponding author
T.N.C. Vidya
Evolutionary and Organismal Biology Unit
Jawaharlal Nehru Centre for Advanced Scientific Research
Bengaluru, India
Email: tncvidya@jncasr.ac.in

ORCID TR, 0000-0003-2471-9194; TNCV, 0000-0002-7143-9008

Data availability statement
The raw data will be uploaded on Dryad or the publicly available JNCASR repository after
acceptance and are available immediately to the reviewers as uploaded files.

Funding
This work was supported by Council of Scientific and Industrial Research, Government of
India, under Grant No. 37(1613)/13/EMR-II, Science and Engineering Research Board
(SERB), Government of India, under Grant No. CRG/2022/005975, and JNCASR intramural
funds. TR was supported as a Ph.D. student by JNCASR. The funders had no role in study
design, data collection and analysis, preparation of the manuscript, or decision to publish.

Conflict of interest
The authors declare that the research was conducted in the absence of any commercial or
financial relationships that could be construed as a potential conflict of interest.

Ethical note

Field data collection was observational in nature. Therefore, no animal handling or manipulation was involved. Researchers maintained a minimum distance of 40 m to the elephants to avoid any disturbance. Fieldwork was carried out under permit number PCCF/WL/E2/CR-121/2014-15 dated 09/05/2014, issued by the office of the Principal Chief Conservator of Forests (Wildlife) and Chief Wildlife Warden, Karnataka.

Author contributions

Conceptualization: TR, TNCV; Data Curation: TR; Formal Analysis: TR; Funding Acquisition: TNCV; Investigation: TR; Methodology: TR, TNCV; Supervision: TNCV; Validation: TR, TNCV; Writing – Original Draft Preparation: TR; Writing – Review & Editing: TR, TNCV.

Acknowledgements

This work was funded primarily by the Council of Scientific and Industrial Research, Government of India, under Grant No. 37(1613)/13/EMR-II, and the Jawaharlal Nehru Centre for Advanced Scientific Research (JNCASR), and in part by the Science and Engineering Research Board (SERB), Government of India, under Grant No. CRG/2022/005975. TR was supported as a student by JNCASR. This work is part of TR's Ph.D. thesis. We thank the offices of the PCCF(WL) and APCCF(WL), Karnataka Forest Department, and of the Conservators of Forests of Nagarahole and Bandipur National Parks and Tiger Reserves for field permits. We also thank other officials and staff of Nagarahole and Bandipur National Parks for their support. We thank Ranga, Krishna, Shankar, Pramod, and others for field assistance. We thank Hansraj Gautam for recording some focal calf videos and members of the Animal Behaviour Lab, especially Ankana Sanyal, Anvitha S, Athira TK, Divya Choudhary, and Jabili Chowdari, for help with checking field data. We thank Amitabh Joshi for suggestions on statistical analyses.

Abstract

Social relationships may increase longevity and reproductive success of individuals, thereby
providing crucial benefits in social species. Adults of many social species have differentiated
social relationships; however, when and how these relationships form and are maintained are
often not understood. We studied spatial and behavioural interactions between young calves
(≤6 months old) and conspecific females (≥5 years old) to understand social ontogeny in a wild
Asian elephant (*Elephas maximus*) population in Nagarahole and Bandipur National Parks,
southern India. We found that elephant groups with calves comprised females – mother and
'escort', who provided active care, and females who did not provide overt, active care – 'other
females'. Escorts, apart from coordinating their movement with the calf, provided allomaternal
care by initiating positive and helpful interactions with the calf. Other females rarely initiated
interactions, which were usually negative, towards calves. Primarily calves, rather than
conspecific females, initiated and terminated proximity contacts and behavioural interactions.
Calves were near and interacted similarly with their mothers and escorts, under feeding, resting,
and social contexts, and almost never interacted with other females. Similarly, we found
remarkable similarities between mothers and escorts in their behaviours towards calves, with
the primary difference being that escorts did not provide milk. Our results suggest that calves
begin to form differentiated social relationships from a young age. The presence of calves and
their interactions with escorts might result from existing close social relationship between
mothers and escorts or may establish new relationships. Allomaternal care may also reinforce
relationship differentiation in such social species. As calf-escort interactions involved
behaviours that may potentially be helpful in protection and feeding skill acquisition, the
development and survival of calves may be affected by the presence of, and interactions, with
allomothers.

Keywords

Allomaternal care, calves, ontogeny, escort, social relationships, Asian elephant

Graphical Abstract

We studied spatial and behavioural interactions between young calves (≤6 months old) and

conspecific females to understand social ontogeny in a wild Asian elephant population. We

found differentiated relationships between calves and conspecifics even at a young age.

Escorts, who showed coordinated movement with calves, engaged in other positive interactions

with calves, with remarkable similarity to the mothers, whereas other females rarely initiated

such interactions. Escorts are thus allomothers in the Kabini elephant population. Calves, rather

than conspecifics, initiated and terminated most of the proximity and behavioural interactions

and were thus precocial in forming new relationships.

Introduction

The development of social relationships is a central component of the ontogeny of social mammals (Bekoff, 1972). Differentiated social relationships may be beneficial (Connor et al., 2001; Langergraber et al., 2013), and such relationships may exist from a young age itself or form gradually (Berman, 1982a, 1982b; Berman et al., 1997; de Waal, 1990; Roatti et al., 2023). Social relationships involving care are of particular importance to developing individuals. In mammals, young individuals form their closest relationship with their mother, who provides a majority of the care (van Noordwijk, 2012). In some species, parental care may be supplemented by alloparental care or allocare, which refers to care contributed by conspecifics apart from the biological parent(s) in raising offspring that are not one's own.

Alloparental care is documented in over 100 social mammals and include behaviours such as babysitting or accompanying the young one, carrying, active food provisioning, allosuckling, and defending against conspecifics or predators that result in direct interactions between allocarers and the young one (Brotherton et al., 2001; Clutton-Brock et al., 2000; Erb & Porter, 2020; Estes & Goddard, 1967; Gadgil & Nair, 1984; Gero et al., 2013; Hill & Campbell, 2014; Hodge, 2005; Hrdy, 1976; Lee, 1987; Pal et al., 2021; Pusey & Packer, 1994; Rood, 1978; Rowell et al., 1964; Whitehead, 1996; Xiang et al., 2019). Allocare may be provided by one or more or all the members of a social group, and allocarers may be generalised or may specialise in caretaking tasks (Erb & Porter, 2020; Garber, 1997; Gero et al., 2009). Young ones may solicit help from specific allocarers and/or allocarers may approach specific young ones to initiate affiliative interactions (Brotherton et al., 2001; Gilchrist, 2004). Ultimately, allocarers can interact differently than other conspecifics with the young ones and *vice versa*, resulting in the development of differentiated social relationships at a young age.

Among social mammals, primates are known for their highly differentiated social relationships (Berman, 1982a, 1982b; Roatti et al., 2023; Silk et al., 2009, 2010). Like many primates, adult female elephants show complex and differentiated social structure (Archie et al., 2011; de Silva et al. 2011; Nandini et al., 2017, 2018; Wittemyer et al., 2005), but when and how relationships form and become differentiated are not well studied. Elephants live in matrilineal societies with female philopatry and male dispersal (Archie et al., 2005; Vidya & Sukumar, 2005). Males associate only temporarily with female groups after dispersal (Keerthipriya et al., 2021) and do not provide parental care (McKay, 1973). Female elephants give birth to a single calf at a time;

twines are very rare. Calves are precocial at birth but are nutritionally and socially dependent on
their mothers (Gadgil & Nair, 1984; Lee & Moss, 2011; Nair, 1989; Revathe et al., 2020).
Young individuals (≤2.5 years of age) face high predation risk from lions in Africa (Loveridge
et al., 2006) and tigers in India (Williams, 1950). Young ones are weaned at only 4-5 years of
age or after the birth of a sibling (Mar et al., 2012) and constantly associate with the mother
until then. Thus, female elephants invest heavily in their young for an extended duration,
matched or exceeded only by certain great apes and cetaceans (Mann, 2019; van Noordwijk et
al., 2018).

Allomaternal care has been conjectured to be important in the evolution and cohesiveness of
female elephant societies (Dublin, 1983; Gadgil & Nair, 1984) and has been documented in
many African and Asian elephant populations (Dublin, 1983; Eisenberg, 1980; Gadgil & Nair,
1984; Jayantha et al., 2009; Lee, 1987; Lee & Moss, 2011; Rapaport & Haight, 1987; Vidya,
2014; Woodford & Trevor, 1970). However, there are only a few detailed studies of calf-
conspecific interactions in elephants (Lee, 1987; Lee & Moss, 2011 on African savannah
elephants (*Loxodonta africana*); Gadgil & Nair, 1984 on semi-captive Asian elephants
(*Elephas maximus*); Chelluri 2009 on wild African forest elephants (*Loxodonta cyclotis*);
Webber 2017 on wild Asian elephants in Uda Walawe, Sri Lanka, and captive Asian and
African savannah elephants) and hardly any studies quantifying allomaternal care and detailing
how such care results in calves experiencing differentiated relationships (but see Lee, 1987;
Lee & Moss, 2011).

We, therefore, investigated the ontogeny of social relationships with a focus on caretaking
behaviours in a long-term monitored wild Asian elephant population in southern India (Kabini
Elephant Project: Keerthipriya and Vidya, 2022; Vidya et al., 2014). We did so by comparing
behaviours of different categories of subadult and adult females towards calves up to 6 months
of age and *vice versa*. Specifically, we examined 1) whether proximity contacts and interactions
were initiated and terminated primarily by adult and subadult female conspecifics or calves (a,
d, and f in Results), 2) how care (and hence, calf-conspecific interactions) was distributed
among female conspecifics in the group and whether individuals that showed one form of care
(escorting, see below) also provided other forms of care (b, c, d, f in Results), and 3) whether
certain kinds of behavioural interactions were associated with specific categories of
conspecifics (e in Results).

In our study, we focused on calves <6 months old, as significant behavioural development
commences during the first six months of life (Revathe et al., 2020). This critical
developmental window is also marked by high mortality (Kabini Elephant Project, unpublished
data), necessitating heightened care. The latter might be expected to result in female
conspecifics initiating proximity contacts and interactions, whereas the precociality hypothesis
(Hill & Campbell, 2014) predicted that calves would initiate such contacts and interactions as
they are precocial and quick to explore their surroundings after birth (Nair, 1989; Revathe et
al., 2020). It was previously found in the Kabini population that certain non-mother females
showed coordinated movement with particular calves, beyond that resulting from coordinated
group movement (Revathe, 2022). We termed these females 'escorts'. In this population, at
least one escort was present in around three-fourths of the female group sightings that had a
calf (Revathe, 2022). Because of differentiated relationships among adult females (Nandini et
al., 2017, 2018) and the consistent presence of specific escorts for specific calves across
elephant groups (Revathe, 2022), we predicted that calves would have differentiated social
relationships with the conspecific females in their clan from a young age. We expected calves
to initiate more spatial and behavioural interactions with their mother – the reliable/familiar
social partner, source of nourishment for the calf, and female with the highest incentive to show
positive interactions to the calf – than with other conspecific categories of females in the group,
and *vice versa*. We did not have an *a priori* expectation about whether escorts would show
more affiliative behaviour towards calves, and *vice versa*, than would other females in the
group. As being near calves by coordinating movement might be thought of as a form of care
by itself, it was not necessary that escorts should show other forms of care, and other
individuals could potentially provide such care. We did not expect the trajectory of social
ontogeny to diverge between females and males within the first six months of life based on the
similar pattern of development across sexes in calves (up to 1 year old) (Revathe et al., 2020).
However, there could be an effect of age on calf relationship differentiation, as calves start to
explore their physical and social environment.
200
In essence, to gain a richer understanding of social ontogeny, we examined certain behaviours
(initiation and termination of proximity and interactions and responses to interactions) from
the perspective of both the calves and conspecific females, as it takes two to form a relationship.
Such an approach can help clarify the relative contributions of individuals (i.e., calf vs
conspecific female vs both; see Hinde & Spencer-Booth, 1967) to the development of
differentiated social relationships.

Methods

Field data collection

We collected field data from December 2015 to June 2018 in Nagarahole and Bandipur National Parks and Tiger Reserves, southern India, where the tiger density is high (Jhala et al., 2008). We carried out field sampling across these two parks following systematic, fixed sampling routes, from ~6:30 AM to ~5:45-6:45 PM (see Nandini et al., 2017 for further details). Female elephant groups were defined as one or more females, often accompanied by young ones, that showed coordinated movement or behaviour and were usually within 50-100 m of one another (Nandini et al., 2017, 2018; equivalent to 'party' in the primate literature). We aged, sexed, and identified the individuals seen based on natural physical characteristics (Vidya et al., 2014). The Kabini Elephant Project has recorded hundreds of individually identified elephants since its inception in 2009 (see Vidya et al., 2014). We age-classified females as subadults (5-<10 years) and adults (>=10 years). Since we were interested in the development of social relationships, we examined calves of different ages. We categorized calves according to differences in behavioural development based on a previous study on this population (Revathe et al., 2020). Therefore, in this study, we refer to those less than three months old as newborn calves and to those between 3 and <6 months of age as infant calves.

The subjects of the study were 20 calves ≤6 months old, belonging to 9 clans. Most of the behavioural data presented here were collected from around the Kabini backwaters (which separate Nagarahole and Bandipur) because of good visibility. Thus, we sampled clans that used this area. We used focal animal sampling (Altmann, 1974) to record interactions between calves and conspecifics within groups. We recorded observations using a video camera (SONY HDR-XR 100E or SONY HDR-PJ 540E).

We defined escorts based on coordinated movement between calf-conspecific pairs, beyond that required of group members (i.e., any distance less than 50-100 m that the group showed) (Revathe, 2022). Thus, one or more non-mother subadult or adult females who moved along with the focal calf (either accompanied by the calf's mother or not) was/were identified as escort(s) in each focal session (Figure 1a) as previously done in whales (e.g., Gero et al., 2009). We refer to females other than the focal calf's mother and escort(s) in a focal session as 'other females' (Figure 1a). Based on this qualitative difference observed in the field (Revathe, 2022),

we differentiated between escorts and other females because not acknowledging this distinction

and clubbing all conspecific females, other than the mother, into one category could mask the

presence of differentiated social relationships, if present, of calves with conspecific females.

Although an escort is a female that showed coordinated movement with the calf, she did not

have to necessarily initiate any, and specifically, allocare, interaction towards the calf. Mother-

escort comparisons in their behaviours towards calves and *vice versa* allowed us to understand

how comparable allomaternal care (if shown by escorts) was to maternal care, and if it was

primarily the escorts, and not other females, that showed allomaternal care (we defined this to

require more than coordinated movement with the calf). It should be noted that although, by

definition, other females did not show coordinated movement with the focal calf, they were

still part of the same group as the focal calf and, hence, could reach the calf to initiate (allocare)

interactions (Figure 1a). Similarly, it was possible for calves to easily go up to other females

to initiate interactions. Essentially, it was entirely feasible for conspecific females to show

escorting behaviour but not allomaternal care and *vice versa*. Data could not be recorded

blindly because our study involved focal animals in the field.

Fieldwork was carried out with relevant Forest Department permissions and in accordance with

India's wildlife protection laws.

Focal video scoring and data analysis

We used Windows Media Player 12.0 to score focal videos to quantify calf-conspecific

proximity and interactions (see below). All the videos were scored by TR. We used only those

focals in which all the group members were clearly visible, and in which all three conspecific

categories of females (Mother, Escort, Other female) were present for the focal calf because

we wanted to make mother-escort, mother-other female, and escort-other female comparisons.

Since a majority of the calves born during the study period had at least one escort in almost all

their sightings (Revathe, 2022) and as calves without an escort often did not survive past a few

months (Kabini Elephant Project, unpublished data), it was not feasible to obtain multiple

focals of calves without an escort. However, since such calves are only a minority in the study

population, we do not expect the lack of their focals in the analyses presented below to

substantially affect our results. Importantly, we use our results to only infer whether calves

with an escort develop differentiated relationships and whether escorting behaviour could be

used as a proxy to identify allomothers in the field. It is likely that calves without an escort also

eventually have differentiated social relationships, however that is beyond the scope of the
current study.

Calf-conspecific proximity contacts

We scored 3 focal videos per calf, recorded on three different days on a second-by-second basis
for 20 calves who were up to 6 months of age, and noted down who initiated movements
(approach or leave; *N*=299 movements) (Supplementary Material 1). The calf belonged to the
same age-class during all 3 focals, and each calf appeared in only one age-class. Each focal
lasted 20 minutes (72,000 seconds scored in all), during which we noted down the focal calf's
positions in terms of who was near the calf (see Supplementary Material 2). We considered a
focal calf to be near a conspecific if the distance between the calf and the conspecific was up
to one calf-body length (≤1 m). A calf could be near multiple individuals simultaneously.

We used the Hinde's proximity index (Hinde & Spencer-Booth, 1967; the percentage of
approaches made by the calf towards a conspecific minus the percentage of leavings made by
that calf away from that conspecific) to find out whether the calf more frequently approached
or left a particular conspecific (see Supplementary Material 3). Additionally, we calculated
Brown's proximity index (Brown, 2001) to measure the relative contributions of the calf and
conspecific to changes in proximity contacts between the pair (see Supplementary Material 3).

Since these indices could be calculated with all three conspecific categories in each focal only
for two calves (as the remaining did not show proximity contact changes with other females),
we could not use parametric tests. We instead tested for differences in Hinde's and Brown's
indices across conspecific categories (Mother, Escort, Other Female) using the non-parametric
Friedman's ANOVA, using the averaged value across focals for each calf and matched for calf
identity. We additionally performed ANOVAs on the logit Hinde's index and Brown's index
to examine possible differences between only mothers and escorts, for which the sample sizes
were larger. Here, conspecific category was a fixed factor with two levels – Mother and Escort.
Calf identity was included as a random factor because there were multiple calves, although
each calf was sampled only in two or three focals. However, we were not actually interested in
variation amongst calves, nor can such variation be tested without many focals per calf.
Therefore, in this and all the other following analyses whenever calf identity was used,
inferences were not made about calf identity or its interactions. Only calves that had Hinde's
and Brown's index values in at least two of their focals were used. Due to low sample size, the

effects of calf age and sex on proximity initiation could not be assessed.

Calf-conspecific proximity

During each 20-minute focal, we took a scan every 4 minutes to obtain six independent calf

positions in terms of the conspecific(s) who were near the calf (4 minutes was chosen based on

how quickly calves changed their positions: this corresponded to 95% of the durations to

changes in positions; see Supplementary Material 2). We thus used a total of 360 calf positions

(6 positions x 3 focals x 20 calves, of which 10 calves were newborns, less than 3 months old,

and 10 calves were infant calves, 3-<6 months old) for this analysis. In order to find out whether

calf proximity differed with respect to conspecific category, we carried out an ANOVA on the

logit proportion of calf-scans near a conspecific (dependent variable; proportions were logit

transformed as they were not normally distributed), with calf age-class (<3 and 3-<6 months)

and conspecific category (Mother, Escort, and Other Female) as fixed factors, and calf identity

nested within age-class as a random factor.

Calf-conspecific interactions

To examine calf-conspecific interactions, we scored two 30-minute focal videos per calf for

the same 20 calves as those used for proximity analyses, during which we noted down all the

interactions that occurred between a focal calf and females of the three conspecific categories,

the identities of the initiator and terminator of each interaction, and the duration of the

interaction. Behaviours were classified into three classes: feeding-related (sucking from a

conspecific, taking plucked grass from a conspecific, sniffing grass from the mouth of a

conspecific, etc.), resting-related (calf leaning against a conspecific, calf lying down near/under

a conspecific, and calf sliding off an individual to lie down), and social (for e.g., touching a

conspecific, rubbing against a conspecific, conspecific standing guard over a calf etc.; see

Supplementary Material 4). We also noted down whether conspecific-initiated interactions, as

well as the responses shown by conspecifics towards calf-initiated interactions, were positive

(for e.g., rushing towards a calf when the calf was in distress, stopping one's activity to allow

the calf to suck, keeping the calf within reachable distance, etc.), negative (for e.g., lashing out

at a calf, kicking a calf, pushing a calf, etc.), or neutral (see Supplementary Material 4).

Conspecifics did not always show a response, and 'no response' was also included under the

category 'neutral' during analysis.

<u>*Initiation of interactions*</u>

In order to find out whether calves initiated more interactions towards conspecifics or *vice*
versa and to find out whether there was a difference amongst mothers, escorts, and other
females in the numbers of interactions that calves initiated towards them and the numbers of
interactions that they initiated towards calves, we ran a nested ANOVA on the log-transformed
numbers of calf-conspecific interactions (as they were not normally distributed), with calf age-
class (<3, 3-<6 months), initiator category (Calf, Conspecific), and conspecific category
(Mother, Escort, and Other Female) as fixed factors, and calf identity nested within age-class
as a random factor. We also tested interaction effects.

Conspecific-initiated interactions: frequency in different contexts
We then classified conspecific-initiated interactions towards calves into four different
interaction types (Supplementary Material 4, Table 2) from the perspective of the calf, namely
positive (potentially benefitting the calf; such as checking the calf, chaperoning the calf),
neutral (no obvious benefit to the calf; such as sniffing the calf), and negative (not benefitting
the calf, or at least not in the immediate interest of the calf). Since not all the negative
interactions cause physical distress, we also split them into non-aggressive negative (not
causing physical distress, such as nudging the calf to move, mother nudging the calf towards
the escort) and aggressive negative (causing physical distress, such as kicking the calf, lashing
out at the calf) interactions. We found that most of the calf-conspecific interactions were
initiated by calves (see Results); hence, we could not use ANOVA to test the effect of
conspecific category on interaction type with calves. We, therefore, used an R x C test of
independence using *G*-test to test the null hypothesis that interaction type with calves was
independent of conspecific category by combining interactions across all focals (of all calves)
in each conspecific category and interaction type.

Calf-initiated interactions: frequency in different contexts, and terminations and types of
responses shown by conspecifics
To understand social preferences of calves, we examined whether the number of calf-initiated
interactions varied based on conspecific category for the different behavioural classes. To do
so, we carried out a nested ANOVA on the log-transformed numbers of calf-initiated
interactions, with calf age-class, conspecific category, and behavioural class of interaction
(Feeding, Resting, and Social) as fixed factors, and calf identity nested within age-class as a
random factor.

For each focal, we also calculated the proportion of calf-initiated interactions towards females

of a conspecific category terminated by females of that category (calves that did not initiate

any interaction towards females in a focal were not included in the analysis, N=2 calves in each

age-class). These proportions of calf-initiated interactions terminated by mothers and escorts

were compared. Terminations by other females were not used as only two calves initiated

interactions with all three conspecific categories of females in all their focals. We carried out

a nested ANOVA on the logit-transformed proportions of calf-initiated interactions terminated

by conspecifics, with calf age-class (<3 and 3-<6 months) and conspecific category (Mother,

Escort) as fixed factors and calf identity nested within age-class as a random factor.

We also calculated the proportions of calf-initiated interactions towards mothers and escorts

that elicited positive, neutral, or negative responses from that conspecific category (these

proportions would add up to 1 for each conspecific category; therefore, the proportion of

negative responses was not included in the analysis). Again, as above, responses by other

females could not be compared as calves did not initiate sufficient interactions towards them.

To see if mothers and escorts showed similar types of responses, we ran a nested ANOVA on

the logit proportions of calf-initiated interactions that elicited a positive or a neutral response,

with calf age-class, conspecific category, and type of response (Positive, Neutral) as fixed

factors, and calf identity nested within age-class as a random factor.

Thus, to check whether females of different conspecific categories behaved in a similar manner

towards calves, we compared conspecific-initiated interactions towards calves, proportion of

calf-initiated interactions that were terminated by conspecifics, and the proportion of calf-

initiated interactions that elicited different kinds of responses, across the conspecific categories.

To check whether calves behaved in a similar manner towards different categories of

conspecific females, we compared calf proximity initiation and leaving, calf proximity, and the

number of calf-initiated interactions – overall and under different behavioural classes, across

the conspecific categories.

Calf-conspecific non-suckling interactions

It was possible that nutritional dependence of calves on their mother might mask the presence

of otherwise comparable relationships of calves with escorts or other females. To investigate

this possibility, we modified the original dataset by removing all the suckling-related

interactions that the focal calves initiated towards the conspecifics and investigated whether

the pattern of social interactions of calves remained the same even when suckling was not
considered. The same analyses that were performed on calf-initiated interactions were also
performed on this modified dataset. We report these results in the supplementary materials and
present only graphs and *P*-values in the results in the main text.

The ANOVAs were carried out by obtaining sums of squares using Statistica (7.0, StatSoft,
Inc. 2004), and the *F* test calculations were carried out based on Neter *et al.* (1990, Chapter 27,
pgs. 1010-1029).

Results

a)   Initiatiation and termination of proximity contacts by calves and conspecifics, and
differences across conspecific categories (mothers, escorts, other females) in proximity contact
initiation

There was no significant difference in the Hinde's (Friedman ANOVA: χ^2= 2.000, *N*=7 calves,
df=2, *P*=0.368, Figure 1b) or Brown's (Friedman ANOVA: χ^2= 4.571, *N*=7 calves, *df*=2,
P=0.102, Figure 1c) proximity indices among mothers, escorts, and other females. ANOVAs
on the logit Hinde's index with mothers and escorts based on data from a larger number of
calves (17 calves, 44 focals) also showed no significant effect of conspecific category
($F_{1,16}$=3.849, *P*=0.067, Figure 1d). However, the average Hinde's index of the calves with their
mothers was not significantly different from zero, suggesting that calves approached and left
their mothers to similar extents, while the Hinde's index with the escorts was significantly
greater than zero, suggesting that calves were more likely to approach than to leave escorts
(confidence intervals in Figure 1d). The average Brown's proximity indices of calves with the
mother, escort, and other females were greater than 50 (Figure 1c,e), indicating that calves
contributed more than their conspecifics to changes in proximity. However, there was no
significant effect of conspecific category (mothers and escorts) on this index either (ANOVA
on logit Brown's index: $F_{1,16}$=0.888, *P*=0.360; 17 calves, 44 focals; Figure 1d).

a.

b.

c.

d.

e.

Figure 1. a) Photo of an elephant group showing the positions of group members. Left to Right:
other female, mother, calf, escort, other female. Photo: Divya Choudhary, © Kabini Elephant

Project; b) average Hinde's and c) average Brown's proximity indices of 7 calves for which

data were available with all three conspecific categories; d) Hinde's and e) Brown's indices of

17 calves with mothers and escorts. Error bars are 95% CI. Dashed lines indicate the expected

Hinde's index of 0 and expected Brown's index of 50 in the respective graphs.

b) Effects of calf age-class and conspecific category on calf-conspecific proximity

Calves were within easy reachable distance ($\leq$1 m) of at least one female conspecific almost

all the time (average proportion of scans near at least one female conspecific $\pm$ 95% CI:

newborn calves: 0.978 $\pm$ 0.022; infant calves: 0.930 $\pm$ 0.030). There was no significant effect

of calf age-class on the proportion of scans when calves were near conspecific females

(ANOVA: $F_{1,18}$=0.205, P=0.656), with newborn calves and infant calves spending similar

proportions of scans near conspecific females (average of the proportion of scans near mother,

escort, and other females $\pm$ 95% CI = 0.475 $\pm$ 0.076 for newborn calves and 0.439 $\pm$ 0.069 for

infant calves; these values were lower than the average proportion of scans near at least one

female conspecific because of averaging across conspecific categories, with other females not

spending much time near calves). However, there was a significant main effect of conspecific

category ($F_{2,36}$=103.315, P<0.001), and its interaction with calf age-class ($F_{2,36}$=8.139,

P=0.001; Figure 2).

The proportions of scans in which calves were near their mothers or escorts (which were not

significantly different from each other; P=0.529) were significantly higher than that near other

females (mother versus other females: P<0.001; escort versus other females: P<0.001).

Newborn calves spent a greater proportion of their time near mothers than escorts (P=0.043)

or other females (P<0.001), and a greater proportion of their time near escorts than other

females (P<0.001; Figure 2). Infant calves spent a similar proportion of their time near mothers

and escorts (P=0.642), but these were higher than that spent near other females (P<0.001 for

both the comparisons, Figure 2). Newborn calves spent a significantly greater proportion of

time near their mothers than infant calves did (P=0.015), but both spent similar proportions of

time near their escorts (P=0.871) and near other females (P=0.529; Figure 2). When tested for

infant calves separately, there was no effect of calf sex on the proportion of scans spent near

conspecifics (Supplementary Material 5).

Figure 2. Logit proportions of scans that a calf spent near the three conspecific categories of

females for 20 calves (<3 months: *N*=10 calves; 3-<6 months: *N*=10 calves). Error bars are

95% CI. In this and the following graphs, letters above the data points indicate pattern of

statistical significance based on Tukey's HSD tests (a<b<c), unless otherwise specified. Shared

letters above the bars indicate no significant difference between comparisons. The

untransformed value (average ± 95% CI) is shown below the graph for each category.

(Averaging was done over all the focals of all the calves in the category in this and the following

graphs.)

c)   Differences across conspecific categories in the number and nature of interactions (i.e.,

positive / neutral / negative) initiated towards calves

We observed 93 positive, 10 neutral, 35 non-aggressive negative, and 10 aggressive negative

conspecific-initiated interactions towards calves. The non-aggressive negative interactions

included only nudge (to make the calf move from a spot), whereas the aggressive negative

interactions included kick, push, pull trunk, beat with tail, and lash (Supplementary Material 4,

Table 1). We found that the type of conspecific-initiated interactions towards calves was

dependent on conspecific category ($G_{corrected}$ = 45.930, *P*<0.001). Escorts initiated a majority

(77.4%) of the 93 positive interactions towards calves, mothers initiated 20.4% of them, and

other females initiated only 2.2% of them. Escorts also initiated most of the 10 neutral

interactions (80%) towards calves, whereas mothers and other females initiated only one

neutral interaction each in total. Escorts also initiated a majority (60%) of the 35 negative non-

aggressive interactions towards calves, followed by mothers (31.4%), and other females

(8.6%). Aggressive negative interactions towards calves were rare (10 interactions), and mostly

(90%) initiated by other females; one interaction was initiated by a mother and none by an

escort.

d) Initiation of behavioural interactions by calves and conspecifics

Of the total of 1332 interactions between calves and their mothers, escorts, and other females

in their groups that we observed during the total focal duration of 1200 minutes, calves initiated

1184 interactions. Most of these interactions were of very short duration (average duration of

interaction ± 95% CI: calf-initiated: 20.4 ± 4.14 seconds; conspecific-initiated: 24.0 ± 14.5

seconds; Supplementary Material 6). We found significant main effects of initiator category,

calf age-class, and conspecific category on the log number of calf-conspecific interactions

(Table 1).

Calves initiated significantly more interactions towards conspecific females (average number

of interactions ± 95% CI per 30-min focal: 29.60 ± 4.931) than conspecific females initiated

towards calves (3.70 ± 1.634; Table 1, Figure 3a). Newborn calves were involved in a

significantly greater number of interactions (39.4 ± 8.87) with conspecific females (initiated

by either) than were infant calves (27.3 ± 6.64; Table 1). The numbers of calf-mother (14.88 ±

3.644) and calf-escort (16.95 ± 3.814) interactions were similar (95% CI around difference

between means for conspecific category: 0.312; Tukey's HSD: $P>0.05$), and both were

significantly higher ($P<0.05$ for both) than the number of calf-other female interactions (1.48

± 0.848).

There was also a significant interaction effect between initiator and conspecific categories

(Table 1). Calves initiated similar numbers of interactions towards mothers and escorts (95%

CI around difference between means for initiator category x conspecific category: 0.348;

$P>0.05$; Figure 3a), and a smaller number of interactions towards other females ($P<0.05$ for

both the comparisons; Figure 3a). On the other hand, escorts initiated a greater number of

interactions towards calves than mothers or other females did ($P<0.05$ for both the

comparisons; previous paragraph, Figure 3a), with the latter two not being significantly

different from each other ($P>0.05$; Figure 3a). When tested separately, there was no effect of

calf sex on the numbers of calf-conspecific interactions involving infant calves (Supplementary

Material 7).

Table 1. Results of the nested ANOVA on the log numbers of all calf-conspecific interactions.

Calf ID (random factor) was nested within age-class, and age-class, initiator category, and

conspecific category were fixed factors. Significant *P* values are marked in bold. The asterisks

in the *P* values column indicate significance in the ANOVA on the log numbers of non-suckling

calf-conspecific interactions (see Supplementary Material 8) for comparison. (Inferences

should not be drawn about Calf ID or its interactions in this and the following tables as each

calf was sampled in only two focals.)

Effect	Effect (F/R)	*SS*	*df*	*MS*	*F*	*P*
Calf ID(age-class)	Random	12.932	18	0.718	1.499	0.102*
Age-class	Fixed	4.573	1	4.573	6.366	**0.021***
Initiator category	Fixed	97.587	1	97.587	164.480	**<0.001***
Conspecific category	Fixed	79.952	2	39.976	61.366	**<0.001***
Calf ID(age-class) x Initiator category	Random	10.679	18	0.593	1.238	0.243
Calf ID(age-class) x Conspecific category	Random	23.452	36	0.651	1.359	0.112*
Age-class x Initiator category	Fixed	0.962	1	0.962	1.621	0.219
Age-class x Conspecific category	Fixed	2.345	2	1.172	1.800	0.180
Initiator category x Conspecific category	Fixed	36.471	2	18.236	68.065	**<0.001***
Calf ID(age-class) x Initiator category x Conspecific category	Random	9.645	36	0.268	0.559	0.977
Age-class x Initiator category x Conspecific category	Fixed	1.350	2	0.675	2.520	0.095
Error	_	57.516	120	0.479	_	_

Figure 3. Log numbers of a) all interactions and b) non-suckling interactions initiated by calves

towards the three conspecific categories of females, and the log numbers of interactions

initiated by these conspecific females towards calves per focal for 20 calves (<3 months: N=10

calves; 3-<6 months: N=10 calves). Error bars are 95% CI. The untransformed number of

interactions (average ± 95% CI) for each category is shown below the X-axis.

e) Calf initiation of interactions of certain behavioural classes (feeding / resting / social)

towards conspecific categories

Among the 1184 calf-initiated interactions were 539 feeding interactions, 89 resting

interactions, and 556 social interactions. The log number of calf-initiated interactions was

significantly affected by the behavioural class of interaction, conspecific category, their

interaction, and calf identity (Table 2). Calves initiated more interactions towards mothers and

escorts than towards other females as seen above (Initiator category x Conspecific category in

Table 1, Figure 3a). This pattern was also found separately in feeding (95% CI around

difference between means for behavioural class x conspecific category: 0.502; Mother versus

Other Females: P<0.05; Escort versus Other Females P<0.05, Figure 4a) and social interactions

(Mother versus Other Females: P<0.05; Escort versus Other Females P<0.05, Figure 4a), but

the number of resting-related interactions initiated by calves was higher only towards escorts

than towards other females (P<0.05), and was similar between mothers and other females

(P>0.05) and mothers and escorts (P>0.05, Figure 4a).

Newborn and infant calves initiated similar numbers of interactions towards conspecific

females (in keeping with the lack of an Age-class x Initiator category effect in Table 1). Calves

initiated similar numbers (average ± 95% CI per 30-minute focal) of feeding (13.48 ± 3.473)

and social interactions (13.90 ± 3.308; 95% CI around difference between means for

behavioural class: 0.310; Tukey's HSD: P>0.05) towards conspecific females, which were both

significantly higher than resting-related interactions (2.23 ± 1.208; P<0.05 for both the

comparisons); this pattern was not different across calf age-classes (Table 2). There was no

effect of sex on the number of calf-initiated interactions for infant calves examined separately

(see Supplementary Material 10).

Table 2. Results of the nested ANOVA on the log number of calf-initiated interactions towards

the three conspecific categories of females. Significant *P* values are marked in bold. The

asterisks in the *P* values column indicate significance in the ANOVA on the log numbers of

calf-initiated non-suckling interactions (see Supplementary Material 11) for comparison.

Effect	Effect (F/R)	*SS*	*df*	*MS*	*F*	*P*
Age-class	Fixed	1.151	1	1.151	1.153	0.297
Calf ID(Age-class)	Random	17.973	18	0.999	2.171	**0.005***
Behavioural-class	Fixed	52.362	2	26.181	27.138	**<0.001***
Conspecific category	Fixed	86.172	2	43.086	59.997	**<0.001***
Behavioural-class* Conspecific category	Fixed	22.078	4	5.519	11.206	**<0.001***
Age-class*Behavioural-class	Fixed	1.379	2	0.689	0.715	0.496
Age-class*Conspecific category	Fixed	0.539	2	0.269	0.375	0.690
Age-class*Behavioural-class*Conspecific category	Fixed	2.166	4	0.542	1.100	0.363
Calf ID(Age-class)*Behavioural-class	Random	34.730	36	0.965	2.098	**0.001***
Calf ID(Age-class)*Conspecific category	Random	25.853	36	0.718	1.561	**0.031***
Calf ID(Age-class)*Behavioural-class*Conspecific category	Random	35.464	72	0.493	1.071	0.353
Error	_	82.788	180	0.460	_	_

Figure 4. Log numbers of a) all calf-initiated interactions and b) calf-initiated non-suckling

interactions of three behavioural classes towards the three conspecific categories of females for

20 calves (<3 months: $N=10$ calves; 3-<6 months: $N=10$ calves). Error bars are 95% CI.

Untransformed numbers (average ± 95% CI) are shown below the graphs.

f) Termination of interactions by calves and conspecifics, and differences across conspecific

categories in the kinds of responses shown towards calf-initiated interactions

<u>Terminations</u>

Calves themselves terminated most ($N=868$) of the interactions that they initiated ($N=1184$),

while conspecifics terminated the remaining. We found that mothers and escorts terminated

similar proportions of calf-initiated interactions towards them ($P=0.271$; Figure 5,

Supplementary Material 12). The same pattern was seen in the termination of non-suckling

interactions (Supplementary Material 12). In both the cases, mothers and escorts terminated

similar proportions of interactions initiated by newborn and infant calves.

606
607

608
609

Figure 5. Logit proportions of calf-initiated interactions with their mothers and escorts that

were terminated by the calves and by the conspecific females for all calf-initiated interactions

for 16 calves (<3 months: $N=8$ calves; 3-<6 months: $N=8$ calves). Error bars are 95% CI. Letters

indicate pattern of statistical significance based on ANOVA. Proportions of calf-initiated

interactions terminated by the calf and mother, and by the calf and escort, respectively, (average

± 95% CI) are shown below the graph.

<u>Responses by conspecifics</u>

Calves received different types of responses, but most of their interactions did not elicit a

response from conspecific females (885 out of 1184 calf-initiated interactions received no

response). There were more negative (192) than positive (107) responses overall, but only 17

of the 192 negative responses were aggressive, such as kick, lash, push, pull trunk, and beat

with tail.

We found that the proportions of calf-initiated interactions towards mothers and escorts that

elicited a neutral response (about 80%) were significantly higher than the proportion that

elicited a positive response from them (about 10%; $P<0.001$; Figure 6, Supplementary Material

13). There was no significant effect of conspecific category or its interaction with response

type (Figure 6, Supplementary Material 13) on the proportion of calf-initiated interactions that

elicited a response.

When we pooled together negative responses to calf-initiated interactions and negative

conspecific female-initiated interactions towards calves across focals, we found that calves less

than 6 months of age received the lowest rate of aggression from their mothers (0.15/h),

followed by escorts (0.4/h). Calves received the highest rate of aggression from other females

(0.8/h), despite calves spending the least proportion of time near other females.

a)

Figure 6. Logit proportions of a) all calf-initiated interactions and b) non-suckling calf-initiated

interactions with their mothers and escorts that elicited a positive, neutral, and negative
response from them for 16 calves (<3 months: *N*=8 calves; 3-<6 months: *N*=8 calves). Error
bars are 95% CI. Letters above the data points indicate patterns of statistical significance (a<b).
Proportions of interactions that elicited a positive, neutral, and negative response from mothers
and escorts are written below the graph (average ± 95% CI).
646
647

Discussion

We examined the relationship between calves that were up to 6 months old and conspecific
females (that were at least 5 years old) to understand the ontogeny of social relationships in
wild Asian elephants. Through analysing calf-female spatial and behavioural interactions, we
found that calves were precocial in forming new relationships. Differentiated social
relationships emerged through escorting females also exhibiting other positive behavioural
interactions with the calf as much as or more than the calf's mother, through mothers and
escorts being similar in their behaviours towards calves and calves towards them, and through
the presence of other females who engaged in more negative than positive interactions with
calves. Thus, our study on social ontogeny shows that differentiated social relationships are
formed from a young age in Asian elephants.

Elephant calves are precocial in the formation of new social relationships
<u>Proximity</u>
We found that calves were responsible for initiating more changes in proximity with mothers,
escorts, and other females than *vice versa* from a young age, in line with the predictions of the
precociality hypothesis (Hill & Campbell, 2014). Offspring have been found to contribute more
to proximity changes than mothers in other precocial species such as belugas (*Delphinapterus*
leucas; Hill & Campbell, 2014), Saharan arrui (*Ammotragus leruia sahariensis*; Cassinello,
1997), and sorraia horses (*Equus caballus*; Heitor & Vicente, 2008). We further found that
calves approached their mothers, escorts, and other females more frequently than they left
them. Contrary to this, calves left their mothers more often than they approached them during
the first 6 months of life in the African savannah elephant (Douglas-Hamilton 1972). However,
the distance at which approaches and leavings by calves were defined in that study was 5 m,
whereas we used a distance of one calf body length, which is about 1 m. Results from the
African savannah elephants hint at the possibility that calves engage in different behaviours

when they are within 1 m than when within 5 m of the mother, as calves were more likely to

have an age mate as a neighbour in the latter scenario than in the former (Lee, 1987). This

suggests that calves might engage in play when they are at a 5-m distance, resulting in them

leaving their mothers more than they approach them. It is further possible that the group's

social environment – i.e., the number of individuals of different age-sex classes and the risk

posed by them to the calves – may dictate conspecific female-calf approach-leave behaviour

(e.g., Berman et al., 1997).

<u>Behavioural interactions</u>

Similar to proximity maintenance, we found that calves initiated and terminated most of the

behavioural interactions, showing their precocial nature in forging social relationships. Similar

results were reported by Gadgil and Nair (1984) in semi-captive Asian elephants (birth to 1.25

years) and by Lee (1986) in the Amboseli elephants (birth to 5 years). We found that the number

of interactions initiated by calves decreased within the first six months of life, which may be a

result of their increasing independence (e.g., Guarino et al., 2017; Lee, 1987). This would be

consistent with the calf exploring its physical rather than social environment, as the proportion

of time spent near the mother also decreased. As expected, males and females did not differ in

the number of interactions initiated, consistent with previous studies (Förster & Cords, 2005;

Lee, 1987), although differences between the sexes may develop with age, as found in play

behaviours in the African savannah elephant (Lee & Moss, 2014).

Calf interactions were differentially distributed among female conspecifics in the group

<u>Proximity</u>

Calves had non-random neighbours and were often positioned between their mothers and

escorts, as previously observed in semi-captive Asian elephants (Gadgil & Nair, 1984). As

young individuals face predation risk from tigers (Williams, 1950), being positioned near

conspecific females in addition to the mother might further reduce the risk of predation (see

Lee, 1987). Overall, calves showed similar proximity with mothers and escorts, but we do not

have similar comparisons from other elephant populations. Newborn calves spent a greater

proportion of time near their mothers than near escorts, which was also seen in semi-captive

elephants (Gadgil & Nair, 1984). As the calves grew (to 3-<6 months of age), they spent more

time away from their mothers; this period coincided with an increase in trunk motor skills and

adult-like feeding behavioural expression (Revathe et al., 2020). Increase in skill competence

has been found to be associated with an increase in mother-offspring distance in other species

also (e.g., *Pongo abelii*: van Noordwijk & van Schaik, 2005). Seeking out other social partners,

such as age-mates, for play, might also result in decreasing time spent near one's mother (e.g.,

Arroyo-Rodríguez et al., 2007). An effect of calf age on mother-calf proximity has also been

reported in wild African savannah elephants (Lee 1986) and wild and captive Asian elephants

in Sri Lanka and UK facilities, respectively (Webber 2017).

We found that the proportion of time spent near escorts did not decrease significantly at least

till 6 months of age. It is possible that escorts continue to actively keep track of and accompany

calves as calves age, perhaps to support maternal foraging freedom while ensuring calf

protection, as suggested in sperm whales (*Physeter macrocephalus*; Whitehead, 1996).

Frequent care from alloparents during later stages of infant development has also been seen in

other species (*Cebus olivaceus*: O'Brien & Robinson, 1991; *Canis familiaris*: Pal et al., 2021).

However, our results from the Hinde's index (albeit for calves of all ages up to 6 months) that

calves were more likely to approach than to leave escorts suggest that calves may be following

escorts rather than escorts moving along with calves. More data are required to differentiate

between these scenarios.

The proportion of time spent in proximity to other females did not increase with calf age.

Whether heterogeneity in calf-conspecific female (≥5 years) relationships is maintained as

calves grow up (e.g., Berman, 1982a; Wey & Blumstein, 2010) or whether it reduces with age

(Berman, 1982b; Roatti et al., 2023) remains to be seen. Overall, our findings are in line with

that found in the African savannah elephant, in which mother-calf distance increased but the

distance between calves and other conspecific females did not vary much with calf age

(Douglas-Hamilton 1972, Lee, 1987). As expected, and in line with a previous study (Webber

2017), we did not find sex-based differences in calf-female conspecific proximity at the young

ages we examined.

Escorting females, but not others, are allomothers

<u>Behavioural interactions</u>

Despite mothers and escorts spending similar proportions of time near calves, escorts initiated

more interactions (of all kinds) towards calves than did mothers. They also initiated the highest

proportion of positive interactions and initiated more positive than negative interactions.

Mothers also initiated more positive than negative interactions, but the difference was small.

As lactation is energetically expensive and requires increased time spent foraging (Wisniewska

et al., 2015) and as primarily mothers provide milk in elephants (including in the study

population), other caretaking behaviours might be shared with escorts, leading to more positive

interactions from escorts than mothers. A previous study in the Kabini elephant population

found that females in certain clans had their first- or second-order relatives as their top

associates (Nandini, 2016). If such females are escorts, they might accrue indirect fitness

benefits through positive interactions with calves.

Unlike African savannah elephant calves (≤12 months old) that had been found to interact with

their mothers more frequently than with any other age-sex class (Lee 1986, 1987), we found

that calves (<6 months) interacted with their mothers and escorts at similarly high rates.

Mothers and escorts showed similar extents of positive responses (that often required them to

stop feeding) to calf-initiated interactions, terminated comparable proportions of calf-initiated

interactions, and rarely directed aggression towards the calf. Mothers and escorts, but not other

females, also guarded calves during long periods of calf resting. As calves <6 months old are

likely in a crucial stage of learning and development, mothers and escorts may be tolerant of

calf interactions through which calves explore and understand their physical and social

environment and learn survival skills. Whether this relationship changes when calves attain

foraging and/or social independence, as that found in great apes (Mikeliban et al., 2021) – who

have learning-intensive, prolonged developmental period similar to elephants – remains to be

seen.

Other females initiated fewer interactions and more negative than positive or neutral

interactions than escorts did towards calves. Some of the differences in the responses of

conspecifics to calves could have arisen from the fact that calves initiated similar numbers of

interactions with their mothers and escorts but far fewer interactions (almost 14 times less

frequent than with escorts) with other females. However, this would not explain the higher

frequency of negative interactions by other females towards calves. The fact that not all the

females other than the calf's mother interacted positively – or interacted at all – with calves

suggests that either conspecific females and/or the calves may actively shape their relationship,

perhaps as a result of the differences in relatedness/familiarity between mothers and other

conspecific females (e.g., Berman, 1982b; Dunayer & Berman, 2017). Differences in female

reproductive status might also contribute to the differential nature of escort and other female

interactions with calves, as females who were escorts did not have a calf of their own during

the study period. However, not all the other females had a calf either during the study period.

Calves' role in the emergence of differentiated social relationships with female conspecifics

779 Calves interacted similarly with mothers and escorts in feeding, resting, and social contexts. Feeding interactions involved passive food sharing behaviours such as calves taking grass that had been scraped off the ground from a conspecific or feeding in the same patch as the conspecific. These behaviours required social tolerance and close contact between calves and conspecifics, which are prerequisites for social transmission of knowledge and skills (van Schaik, 2003), and were almost always initiated towards mothers and escorts, and almost never towards other females. Social interactions through which calves probably sought protection (from heat or predation) were also primarily with mothers and escorts, and such interactions interfered with the conspecific's foraging, as they partly restricted their leg movement (feeding requires kicking at and scraping short grass with the foot around the Kabini backwaters). Close proximity and interactions with escorts may contribute to not just protection of young ones but also social learning opportunities in this long-lived species with a prolonged developmental period.

793 We found that suckling interactions accounted for about half of the calf-initiated interactions with their mothers. Calves also sucked from escorts and rarely from other females, but these were non-lactating females; thus, calves received milk only from their mothers. Calves were quickly rejected by other females the few times calves tried to suck from them, as was seen in the African savannah elephant (Lee, 1987). This is expected as lactation is energetically costly. Calves initiated more non-suckling interactions towards their escorts than even their mothers, despite similar calf-mother and calf-escort proximity. Furthermore, almost all the play interactions with conspecific females (≤5 years) were also initiated towards escorts.

802 We do not yet know if escorts are more closely related or more familiar to calves than other females, and whether this drives the difference in interactions. In many species, interactions of developing individuals with familiar/related individuals are more frequent than that with unfamiliar/unrelated individuals (Bădescu et al., 2015; Berman, 1982a; Konrad et al., 2019; Lee, 1987). Allomaternal care may, therefore, reinforce existing relationship differentiation. It is also possible that escorts were younger than other females, and were, therefore, preferred by calves (see Lee, 1987). Furthermore, although other females do not show overt care, they may still participate in cooperative defense of calves against predators (see McComb et al., 2011). Since our study included 20 calves from 9 elephant clans, our results are generalizable.

However, we did not have enough sample size to test the effect of calf age and sex within the

same model or the potential confounding effects of clan, seasonality, and year. These effects

should be investigated with more data points.

In conclusion, we found that elephant groups with calves comprised females who provided

active care, them being mothers and escorts, and those who did not provide overt, active care,

them being other females. Females who were classified as escorts in a focal, based on

coordinated movement, always provided other forms of allomaternal care during that focal.

Overall, there were remarkable similarities between mothers and escorts in their behaviours

towards calves and in the behaviours of calves towards them, with the primary difference being

that escorts did not provide milk. These results together show that there are differentiated social

relationships – i.e., variation in the quantity and quality of interactions between calves and

conspecific females – from a young age. The presence of calves and their interactions with

escorts in the group might increase close social interactions between mothers and escorts,

which may establish new relationships or strengthen existing relationships (see also Gadgil &

Nair, 1984; Lee, 1987). As calf-escort interactions involved behaviours that may potentially be

helpful in feeding skill acquisition and protection, the development and survival of calves may

be affected by the presence of, and/or interactions, with allomothers (Hrdy 1976, Lee 1987,

Hodge 2005). The emergence of differentiated social relationships in animal societies has often

been studied from the perspective of rank, sex, and age differences among the social partners

of the maturing individuals, apart from differential maternal relations (including the effect of

relatedness) with conspecifics. Our study highlights an additional factor - allomaternal care -

which may potentially lead to the formation of new or reinforcement of existing relationship

structure in animal societies. Our results, therefore, bear significance for several other

mammalian study systems that show both alloparental care and differentiated adult social

relationships.

References

1. Altmann, J. (1974). Observational study of behavior: Sampling methods. *Behaviour*, *49*(3–

4), 227–266. https://doi.org/10.1163/156853974X00534

2. Archie, E. A., Moss, C. J., & Alberts, S. C. (2005). The ties that bind: Genetic relatedness

predicts the fission and fusion of social groups in wild African elephants. *Proceedings of*

the Royal Society B: Biological Sciences, *273*(1586), 513–522.
https://doi.org/10.1098/rspb.2005.3361

3. Archie, E. A., Moss, C. J., & Alberts, S. C. (2011). Friends and relations: Kinship and the
nature of female elephant social relationships. In C. J. Moss, H. Croze, and P. C. Lee (Eds.),
The Amboseli elephants: A long-term perspective on a long-lived mammal, (pp. 238–245).
University of Chicago Press. https://doi.org/10.7208/9780226542263-017

4. Arroyo-Rodríguez, V., Serio-Silva, Carlos., J, Álamo-García, J., & Ordano, M. (2007).
Exploring immature-to-mother social distances in Mexican mantled howler monkeys at Los
Tuxtlas, Mexico. *American Journal of Primatology*, *69*(2), 173–181.
https://doi.org/10.1002/ajp.20344

5. Bădescu, I., Sicotte, P., Ting, N., & Wikberg, E. C. (2015). Female parity, maternal kinship,
infant age and sex influence natal attraction and infant handling in a wild colobine (*Colobus*
vellerosus). *American Journal of Primatology*, *77*(4), 376–387.
https://doi.org/10.1002/ajp.22353

6. Bekoff, M. (1972). The development of social interaction, play, and metacommunication
in mammals: An ethological perspective. *The Quarterly Review of Biology*, *47*(4), 412–
434. https://doi.org/10.1086/407400

7. Berman, C. M. (1982a). The ontogeny of social relationships with group companions
among free-ranging infant rhesus monkeys I. Social networks and differentiation. *Animal*
Behaviour, *30*(1), 149–162. https://doi.org/10.1016/S0003-3472(82)80250-9

8. Berman, C. M. (1982b). The ontogeny of social relationships with group companions
among free-ranging infant rhesus monkeys II. Differentiation and attractiveness. *Animal*
Behaviour, *30*(1), 163–170. https://doi.org/10.1016/S0003-3472(82)80251-0

9. Berman, C. M., Rasmussen, K. L. R., & Suomi, S. J. (1997). Group size, infant
development and social networks in free-ranging rhesus monkeys. *Animal Behaviour*,
53(2), 405–421. https://doi.org/10.1006/anbe.1996.0321

10. Brotherton, P. N. M., Clutton-Brock, T. H., O'Riain, M. J., Gaynor, D., Sharpe, L., Kansky,
R., & McIlrath, G. M. (2001). Offspring food allocation by parents and helpers in a
cooperative mammal. *Behavioral Ecology*, *12*(5), 590–599.
https://doi.org/10.1093/beheco/12.5.590

11. Brown, G. R. (2001). Using proximity measures to describe mother-infant relationships.
Folia Primatologica, *72*(2), 80–84. https://doi.org/10.1159/000049926

12. Cassinello, J. (1997). Mother-offspring conflict in the Saharan Arrui, *Ammotragus lervia*
sahariensis: Relation to weaning and mother's sexual activity. *Ethology*, *103*(2), 127–137.
https://doi.org/10.1111/j.1439-0310.1997.tb00012.x

13. Chelluri, G. (2009). Ontogeny of the mother-calf relationship in African forest elephants,
Loxodonta africana cyclotis. Honours thesis, Cornell University, Cornell.

14. Clutton-Brock, T. H., Brotherton, P. N. M., O'Riain, M. J., Griffin, A. S., Gaynor, D.,
Sharpe, L., Kansky, R., Manser, M. B., & McIlrath, G. M. (2000). Individual contributions
to babysitting in a cooperative mongoose, *Suricata suricatta. Proceedings of the Royal*
Society of London. Series B: Biological Sciences, *267*(1440), 301–305.
https://doi.org/10.1098/rspb.2000.1000

15. Connor, R. C., Heithaus, M. R., & Barre, L. M. (2001). Complex social structure, alliance
stability and mating access in a bottlenose dolphin 'super-alliance.' *Proceedings of the*
Royal Society of London. Series B: Biological Sciences.
https://doi.org/10.1098/rspb.2000.1357

16. de Waal, F. B. M. (1990). Do rhesus mothers suggest friends to their offspring? *Primates*,
31(4), 597–600. https://doi.org/10.1007/BF02382543

17. Douglas-Hamilton, I. (1972). On the ecology and behaviour of the African elephant: The
elephants of Lake Manyara. Ph.D. thesis, University of Oxford, Oxford.

18. Dublin, H. T. (1983). Cooperation and reproductive competition among female African
elephants. In S.K. Wasser, *Social Behavior of Female Vertebrates*, (pp. 291–313).
Academic Press, New York. https://doi.org/10.1016/B978-0-12-735950-2.50016-5

19. Dunayer, E. S., & Berman, C. M. (2017). Infant handling enhances social bonds in free-
ranging rhesus macaques (Macaca mulatta). *Behaviour*, *154*(7–8), 875–907.
https://doi.org/10.1163/1568539X-00003448

20. Eisenberg, J. F. (1980). Ecology and Behavior of the Asian Elephant. *Elephant*, *1*(5), 36–
56. https://doi.org/10.22237/elephant/1521731835

21. Erb, W. M., & Porter, L. M. (2020). Variable infant care contributions in cooperatively
breeding groups of wild saddleback tamarins. *American Journal of Primatology*, *82*(12),
e23190. https://doi.org/10.1002/ajp.23190

22. Estes, R. D., & Goddard, J. (1967). Prey Selection and hunting behavior of the African wild
dog. *The Journal of Wildlife Management*, *31*(1), 52. https://doi.org/10.2307/3798360

23. Förster, S., & Cords, M. (2005). Socialization of infant blue monkeys (*Cercopithecus mitis*
stuhlmanni): Allomaternal interactions and sex differences. *Behaviour*, *142*(7), 869–896.
https://doi.org/10.1163/1568539055010138

24. Gadgil, M., Hegde, M., Joshi, N. V., & Gadgil, S. (1985). On the communication of well-
being.   *Proceedings: Animal   Sciences*,   *94*(6),   575–586.
https://doi.org/10.1007/BF03191862

25. Gadgil, M., & Nair, P. V. (1984). Observations on the social behaviour of free ranging
groups of tame Asiatic elephant (*Elephas maximus Linn*). *Proceedings: Animal Sciences*,
93(3), 225–233. https://doi.org/10.1007/BF03186282

26. Garber, P. A. (1997). One for all and breeding for one: Cooperation and competition as a
tamarin reproductive strategy. *Evolutionary Anthropology: Issues, News, and Reviews*,
5(6),   187–199.   https://doi.org/10.1002/(SICI)1520-6505(1997)5:6<187::AID-
EVAN1>3.0.CO;2-A

27. Gero, S., Engelhaupt, D., Rendell, L., & Whitehead, H. (2009). Who Cares? Between-
group variation in alloparental caregiving in sperm whales. *Behavioral Ecology*, *20*(4),
838–843. https://doi.org/10.1093/beheco/arp068

28. Gero, S., Gordon, J., & Whitehead, H. (2013). Calves as social hubs: Dynamics of the social
network within sperm whale units. *Proceedings of the Royal Society B: Biological*
Sciences, *280*(1763), 20131113. https://doi.org/10.1098/rspb.2013.1113

29. Gilchrist, J. S. (2004). Pup escorting in the communal breeding banded mongoose:
Behavior,   benefits,   and   maintenance.   *Behavioral   Ecology*,   *15*(6),   952–960.
https://doi.org/10.1093/beheco/arh071

30. Guarino, S., Hill, H. M., & Sigman, J. (2017). Development of sociality and emergence of
independence in a killer whale (*Orcinus orca*) calf from birth to 36 months: Killer whale
calf   sociality   and   independence.   *Zoo   Biology*,   *36*(1),   11–20.
https://doi.org/10.1002/zoo.21338

31. Heitor, F., & Vicente, L. (2008). Maternal care and foal social relationships in a herd of
Sorraia horses: Influence of maternal rank and experience. *Applied Animal Behaviour*
Science, *113*(1–3), 189–205. https://doi.org/10.1016/j.applanim.2007.11.005

32. Hill, H. M., & Campbell, C. A. (2014). The frequency and nature of allocare by a group of
Belugas (*Delphinapterus leucas*) in human care. *International Journal of Comparative*
Psychology, *27*(4). https://doi.org/10.46867/ijcp.2014.27.04.08

33. Hinde, R. A., & Spencer-Booth, Y. (1967). The behaviour of socially living rhesus
monkeys in their first two and a half years. *Animal Behaviour*, *15*(1), 169–196.
https://doi.org/10.1016/S0003-3472(67)80029-0

34. Hodge, S. J. (2005). Helpers benefit offspring in both the short and long-term in the
cooperatively breeding banded mongoose. *Proceedings of the Royal Society B: Biological*
Sciences, *272*(1580), 2479–2484. https://doi.org/10.1098/rspb.2005.3255

35. Hrdy, S. B. (1976). Care and exploitation of nonhuman primate infants by conspecifics
other than the mother. In J. S. Rosenblatt, R. A. Hinde, E. Shaw, & C. Beer (Eds.), *Advances*
in the Study of Behavior (Vol. 6, pp. 101–158). Academic Press.
https://doi.org/10.1016/S0065-3454(08)60083-2

36. Jayantha, D., Dayawansa, P. N., Padmalal, U. K. G. K., & Ratnasooriya, W. D. (2009).
Social relationships of wild juvenile Asian Elephants (*Elephas maximus*) in the Udawalawa
National Park, Sri Lanka. *Journal of Threatened Taxa*, *1*(4), 211–214.
https://doi.org/10.11609/JoTT.o1831.211-4

37. Jhala, Y.V., Gopal, R., & Qureshi, Q. (2008). Status of tigers, co-predators and prey in
India. Report. National Tiger Conservation Authority and Wildlife Institute of India.
TR08/001.

38. Keerthipriya, P., Nandini, S., & Vidya, T. N. C. (2021). Effects of male age and female
presence on male associations in a large, polygynous mammal in Southern India: The Asian
elephant. *Frontiers in Ecology and Evolution*, *9*, 616666.
https://doi.org/10.3389/fevo.2021.616666

39. Keerthipriya, P. & T.N.C. Vidya (2022). Kabini Elephant Project: A long-term programme
for understanding Asian elephant behaviour and ecology. Trumpet Project Elephant
Division Vol. III(1-2), 65–71.

40. Konrad, C. M., Frasier, T. R., Whitehead, H., & Gero, S. (2019). Kin selection and allocare
in sperm whales. *Behavioral Ecology*, *30*(1), 194–201.
https://doi.org/10.1093/beheco/ary143

41. Langergraber, K. E., Mitani, J. C., Watts, D. P., & Vigilant, L. (2013). Male–female socio-
spatial relationships and reproduction in wild chimpanzees. *Behavioral Ecology and*
Sociobiology, *67*(6), Article 6. https://doi.org/10.1007/s00265-013-1509-6

42. Lee, P.C. (1986). Early social development among African elephant calves. National
Geographic Research 2, 388-401.

43. Lee, P. C. (1987). Allomothering among African elephants. *Animal Behaviour, 35*(1), 278–
291. https://psycnet.apa.org/doi/10.1016/S0003-3472(87)80234-8

44. Lee, P. C., & Moss, C. J. (2014). African elephant play, competence and social complexity.
Animal Behavior and Cognition, 2(2), 144. https://doi.org/10.12966/abc.05.05.2014

45. Lee, P., & Moss, C. (2011). Calf development and maternal rearing strategies. In C. J.
Moss, H. Croze, and P. C. Lee (Eds.), *The Amboseli elephants: A long-term perspective on*
a   long-lived   mammal,   (pp.   224–237).   University   of   Chicago   Press.
https://doi.org/10.7208/chicago/9780226542263.003.0014

46. Loveridge, A. J., Hunt, J. E., Murindagomo, F., & Macdonald, D. W. (2006). Influence of
drought on predation of elephant (*Loxodonta africana*) calves by lions (*Panthera leo*) in an
African   wooded   savannah.   *Journal   of   Zoology*,   *270*(3),   523–530.
https://doi.org/10.1111/j.1469-7998.2006.00181.x

47. Mann, J. (2019). Maternal care and offspring development in Odontocetes. In B. Würsig
(Ed.), *Ethology and Behavioral Ecology of Odontocetes* (pp. 95–116). Springer
International Publishing. https://doi.org/10.1007/978-3-030-16663-2_5

48. Mar, K. U., Lahdenperä, M., & Lummaa, V. (2012). Causes and correlates of calf mortality
in   captive   Asian   elephants   (*Elephas   maximus*).   *PLoS   ONE*,   *7*(3),   e32335.
https://doi.org/10.1371/journal.pone.0032335

49. McComb, K., Shannon, G., Durant, S. M., Sayialel, K., Slotow, R., Poole, J., & Moss, C.
(2011). Leadership in elephants: The adaptive value of age. *Proceedings of the Royal*
Society   B: Biological   Sciences,   *278*(1722),   3270–3276.
https://doi.org/10.1098/rspb.2011.0168

50. McKay, G. M. (1973). Behavior and ecology of the Asiatic elephant in southeastern
Ceylon. Smithsonian Contributions to Zoology, 125, 1–113. 10.5479/si.00810282.125

51. Mikeliban, M., Kunz, B., Rahmaeti, T., Uomini, N., & Schuppli, C. (2021). Orangutan
mothers adjust their behaviour during food solicitations in a way that likely facilitates
feeding   skill   acquisition   in   their   offspring.   *Scientific   Reports*,   *11*(1),   23679.
https://doi.org/10.1038/s41598-021-02901-z

52. Nair, P. V. (1989). Development of nonsocial behaviour in the Asiatic elephant. *Ethology*,
82(1), 46–60. https://doi.org/10.1111/j.1439-0310.1989.tb00486.x

53. Nandini, S. (2016). Social structure, genetic relatedness, and dominance relationships in
female Asian elephants in Nagarahole and Bandipur National Parks, southern India. Ph.D.
thesis, Jawaharlal Nehru Centre for Advanced Scientific Research, Bengaluru.

54. Nandini, S., Keerthipriya, P., & Vidya, T. N. C. (2017). Seasonal variation in female Asian
elephant social structure in Nagarahole-Bandipur, southern India. *Animal Behaviour*, *134*,
135–145. https://doi.org/10.1016/j.anbehav.2017.10.012

55. Nandini, S., Keerthipriya, P., & Vidya, T. N. C. (2018). Group size differences may mask
underlying similarities in social structure: A comparison of female elephant societies.
Behavioral Ecology, *29*(1), 145–159. https://doi.org/10.1093/beheco/arx135

56. O'Brien, T. G., & J. G. Robinson (1991). Allomaternal care by female wedge-capped
capuchin monkeys: Effects of age, rank and relatedness. *Behaviour*, *119*(1/2), 30–50.
http://www.jstor.org/stable/4534974

57. Pal, S. K., Roy, S., & Ghosh, B. (2021). Pup rearing: The role of mothers and allomothers
in free-ranging domestic dogs. *Applied Animal Behaviour Science*, *234*, 105181.
https://doi.org/10.1016/j.applanim.2020.105181

58. Pusey, A. E., & Packer, C. (1994). Non-offspring nursing in social carnivores: Minimizing
the costs. *Behavioral Ecology*, *5*(4), 362–374. https://doi.org/10.1093/beheco/5.4.362

59. Rapaport, L., & Haight, J. (1987). Some observations regarding allomaternal caretaking
among captive Asian elephants (*Elephas maximus*). *Journal of Mammalogy*, *68*(2), 438–
442. https://doi.org/10.2307/1381494

60. Revathe, T. (2022). Calf development, calf-conspecific interactions, and the effect of calves
on female social structure in the Kabini Asian elephant population, southern India. Ph.D.
thesis, Jawaharlal Nehru Centre for Advanced Scientific Research, Bengaluru.
https://doi.org/10.13140/RG.2.2.15873.43366

61. Revathe, T., Anvitha, S., & Vidya, T. N. C. (2020). Development of motor control and
behaviour in Asian elephants in the Kabini elephant population, southern India. *The
International Journal of Developmental Biology*, *64*(4-5–6), 367–382.
https://doi.org/10.1387/ijdb.190274tv

62. Roatti, V., Cowlishaw, G., Huchard, E., & Carter, A. (2023). Social network inheritance
and differentiation in wild baboons. *Royal Society Open Science*, *10*(5), 230219.
https://doi.org/10.1098/rsos.230219

63. Rood, J. P. (1978). Dwarf mongoose helpers at the den. *Zeitschrift Für Tierpsychologie*,
48(3), 277–287. https://doi.org/10.1111/j.1439-0310.1978.tb00260.x

64. Rowell, T. E., Hinde, R. A., & Spencer-Booth, Y. (1964). "Aunt"-infant interaction in
captive rhesus monkeys. *Animal Behaviour*, *12*(2–3), 219–226.
https://doi.org/10.1016/0003-3472(64)90004-1

65. Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R.,
Wittig, R. M., Seyfarth, R. M., & Cheney, D. L. (2009). The benefits of social capital:
Close social bonds among female baboons enhance offspring survival. *Proceedings of the*

Royal Society B: Biological Sciences, *276*(1670), 3099–3104.
https://doi.org/10.1098/rspb.2009.0681

66. Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R.,
Wittig, R. M., Seyfarth, R. M., & Cheney, D. L. (2010). Strong and consistent social bonds
enhance the longevity of female baboons. *Current Biology*, *20*(15), 1359–1361.
https://doi.org/10.1016/j.cub.2010.05.067

67. Sukumar, R. (2003). Mothers, children, and aunts. In The Living Elephants (pp. 125–185).
Oxford University Press, New York.

68. van Noordwijk, M. A. (2012). From maternal investment to lifetime maternal care. n J. C.
Mitani, J. Call, P. M. Kappeler, R. A. Palombit, & J. B. Silk (Eds.), The evolution of primate
societies (pp. 321–42). University of Chicago Press. https://doi.org/10.5167/UZH-70319

69. van Noordwijk, M. A., Utami Atmoko, S. S., Knott, C. D., Kuze, N., Morrogh-Bernard, H.
C., Oram, F., Schuppli, C., van Schaik, C. P., & Willems, E. P. (2018). The slow ape: High
infant survival and long interbirth intervals in wild orangutans. *Journal of Human*
Evolution, *125*, 38–49. https://doi.org/10.1016/j.jhevol.2018.09.004

70. van Noordwijk, M. A., & van Schaik, C. P. (2005). Development of ecological competence
in Sumatran orangutans. *American Journal of Physical Anthropology*, *127*(1), 79–94.
https://doi.org/10.1002/ajpa.10426

71. van Schaik, C.P. (2003). Local traditions in orangutans and chimpanzees: Social learning
and social tolerance. In D.M., Fragaszy & S., Perry (Eds.), The Biology of Traditions:
Models and Evidence (pp. 297–328). Cambridge University Press, Cambridge.

72. Vidya, T. N. C. (2014). Novel behaviour shown by an Asian elephant in the context of
allomothering. *Acta Ethologica*, *17*, 123-127. https://doi.org/10.1007/s10211-013-0168-y

73. Vidya, T. N. C., Prasad, D., & Ghosh, A. (2014). Individual identification in Asian
elephants. *Gajah, 40*, 3-17

74. Vidya, T. N. C., & Sukumar, R. (2005). Social organization of the Asian elephant (*Elephas*
maximus) in southern India inferred from microsatellite DNA. *Journal of Ethology*, *23*(2),
Article 2. https://doi.org/10.1007/s10164-005-0144-8

75. Webber, C.E. (2017). A Comparison of behavioural development of elephant calves in
captivity and in the wild: Implications for welfare. Ph.D. thesis, University of Stirling,
Stirling.

76. Wey, T. W., & Blumstein, D. T. (2010). Social cohesion in yellow-bellied marmots is
established through age and kin structuring. *Animal Behaviour*, *79*(6), 1343–1352.
https://doi.org/10.1016/j.anbehav.2010.03.008

77. Whitehead, H. (1996). Babysitting, dive synchrony, and indications of alloparental care in
sperm whales. *Behavioral Ecology and Sociobiology*, *38*(4), 237–244.
https://doi.org/10.1007/s002650050238
78. Williams, J.H. (1950). Elephant Bill. Rupert Hart-Davis, London.
79. Wisniewska, M., Freeman, E. W., & Schulte, B. A. (2015). Behavioural patterns among
female African savannah elephants: The role of age, lactational status, and sex of the
nursing calf. *Behaviour*, *152*(12–13), 1719–1744. https://doi.org/10.1163/1568539X-
00003301
80. Wittemyer, G., Douglas-Hamilton, I., & Getz, W. M. (2005). The socioecology of
elephants: Analysis of the processes creating multitiered social structures. *Animal
Behaviour*, *69*(6), 1357–1371. https://doi.org/10.1016/j.anbehav.2004.08.018
81. Woodford, M. H., & Trevor, S. (1970). Fostering a baby elephant. *African Journal of
Ecology*, *8*(1), 204–205. https://doi.org/10.1111/j.1365-2028.1970.tb00846.x
82. Xiang, Z., Fan, P., Chen, H., Liu, R., Zhang, B., Yang, W., Yao, H., Grueter, C. C., Garber,
P. A., & Li, M. (2019). Routine allomaternal nursing in a free-ranging Old World monkey.
Science Advances, *5*(2), eaav0499. https://doi.org/10.1126/sciadv.aav0499
1090
1091

1092 **Supplementary Material**

1093

1094 Supplementary Material 1. Details of focal calves.

1095

1096 Supplementary Material 1, Table 1. Details of focal calves including their clan identities, age-

1097 class (at the time of sampling), and sex that were sampled to get durations of calf positions,

1098 calf proximity, and calf behavioural interactions data.

1099

S.No.	Calf identity	Clan identity	Calf sex	Calf age-class
1	Genette_2017_F	Patricia	F	<3 months
2	Georgina_2017_F	Patricia	F	<3 months
3	Ilaena_2016_F	Menaka	F	<3 months
4	Alena_2017_M	Alexandra	M	<3 months
5	Amarilla_2016_M	Osanna	M	<3 months
6	Gemini_2017_M	Patricia	M	<3 months
7	Iliora_2015_M	Menaka	M	<3 months
8	Ipomoea_2018_M	Victoria	M	<3 months
9	Kai_2016_M	Nakshatra	M	<3 months
10	Kasturi_2018_M	Kasturi	M	<3 months
11	Floppy_ears_2018_F	Victoria	F	3-<6 months
12	Marlene_2015_F	Tilottama	F	3-<6 months
13	Namrata_2017_F	Nakshatra	F	3-<6 months
14	Suhrita_2016_F	Victoria	F	3-<6 months
15	Zarin_2015_F	Victoria	F	3-<6 months
16	Camila_2017_M	Patricia	M	3-<6 months
17	Kausalya_2015_M	Kasturi	M	3-<6 months
18	Orlanda_2015_M	Lisa	M	3-<6 months
19	Valerie_2016_M	Victoria	M	3-<6 months
20	Vanessa_2015_M	Victoria	M	3-<6 months

1100

1101

1102 Supplementary Material 1, Table 2. Details of focal calves, including their clan identities and
1103 age-, sex-class that were sampled to get durations of calf positions to calculate the time to
1104 independence of calf positions.
1105

Calf identity	Clan identity	Calf sex	Calf age (in months)	Calf age-class	No. of different calf positions
Kausalya_2015_M	Kasturi	M	4.33	<6 months	25
Kausalya_2015_M	Kasturi	M	4.43	<6 months	12
Kausalya_2015_M	Kasturi	M	4.97	<6 months	18
Hannah_2016_F	Tilottama	F	3.80	<6 months	16
Hannah_2016_F	Tilottama	F	3.97	<6 months	20
Hannah_2016_F	Tilottama	F	4.63	<6 months	22
Vanessa_2015_M	Victoria	M	5.23	<6 months	6
Vanessa_2015_M	Victoria	M	5.73	<6 months	27
Vanessa_2015_M	Victoria	M	5.97	<6 months	45
Genette_2017_F	Patricia	F	0.10	<6 months	22
Genette_2017_F	Patricia	F	1.00	<6 months	8
Genette_2017_F	Patricia	F	1.17	<6 months	22
Leena_2017_F	Lisa	F	3.37	<6 months	15
Leena_2017_F	Lisa	F	3.60	<6 months	9
Leena_2017_F	Lisa	F	4.00	<6 months	32

1106
1107

1108   Supplementary Material 2. Calf position codes and duration to independence of positions.
1109

1110

Supplementary Material 2, Table 1. Focal calf's position codes and their descriptions. Mother

refers to the focal calf's mother; escort/s refers to the focal calf's escort/s; other female/s refers

to females (≥ 5 years) other the focal calf's mother or escorts in a group; conspecific male

refers to males (≥ 5 years) in a group; juvenile refers to a female or a male juvenile in the group;

and another calf refers to a female or a male calf other than the focal calf in the group. Near

refers to the focal calf standing/sitting/lying down within one calf body length of any of the

conspecifics. The order in which conspecific categories (Calf/Juvenile/Mother/Escort/Other

female/Male) appear in the position code do not signify anything.

1119

S.No.	Focal calf position code	Code description
1	ALO	Alone
2	SNC	Near another calf
3	SCE	Near another calf and an escort
4	CCJ	Near another calf and a juvenile
5	FCJ	Near another calf, a juvenile, and an other female
6	SCM	Near another calf and the mother
7	CCC	Near another calf and two other females
8	SNE	Near an escort and not near any other individual
9	ESC	Near an escort and another calf
10	SEE	Near two escorts
11	SEC	Near an escort and an other female
12	EFC	Near an escort, an other female, and another calf
13	EFJ	Near an escort, an other female, and a juvenile
14	ECC	Near an escort and two other females
15	SNJ	Near a juvenile
16	SJE	Near a juvenile and an escort
17	SJJ	Near two juveniles
18	SJM	Near a juvenile and the mother

S.No.	Focal calf position code	Code description
19	SML	Near a conspecific male
20	SME	Near a conspecific male and an escort
21	SMM	Near a conspecific male and the mother
22	SMF	Near a conspecific male and an other female
23	MFE	Near a conspecific male, an other female, and an escort
24	SNM	Near the mother and not near any other individual
25	SEM	Near an escort and the mother
26	MEC	Near the mother, an escort, and another calf
27	MEE	Near the mother and two escorts
28	MEJ	Near the mother, an escort, and a juvenile
29	MEM	Near the mother, an escort, and a conspecific male
30	MEF	Near the mother, an escort, and an other female
31	SMC	Near the mother and an other female
32	MFC	Near the mother, an other female, and another calf
33	MFJ	Near the mother, an other female, and a juvenile
34	MCC	Near the mother and two other females
35	MCO	Near the mother, two other females, and a calf
36	SCO	Near an other female and not near any other individual
37	SFC	Near an other female and another calf
38	FEE	Near an other female and two escorts
39	SFJ	Near an other female and a juvenile
40	SCC	Near two other females

1120

1121   While a focal calf could be near other calves or juveniles or subadult/adult males (as in the
1122   table above), as we were not interested in looking at these contacts, we did not include them in
1123   the analysis. Only changes in calf positions when alone or involving conspecific females were
1124   considered for the analysis.

1125

1126

1127

1128

Durations of calf positions

We constructed cumulative frequency distributions of durations of calf positions, calculated

from second-to-second scoring of calf position changes for calves <6 months of age. Position

changes were included irrespective of whether the calf or the conspecific moved to bring about

the change. We found that 95% of the calf positions (near conspecific females ≥5 years or

alone) changed within 200-210 seconds for calves <6 months (see figure below). Therefore,

we considered two subsequent calf positions to be independent if they were separated by 4

minutes.

1137

1138

1139

1140 Supplementary Material 2, Figure 1. Cumulative frequency distributions of the time durations

1141 of calf positions for calves that were <6 months of age.

1142

1143

1144

1145   Supplementary Material 3. Details of Hinde's and Brown's proximity indices.

1146

1147   We examined calf-conspecific proximity initiations using Hinde's proximity index and
1148   Brown's proximity index.

1149

Hinde's proximity index

Hinde's proximity index (Hinde and Spencer-Booth 1967) was originally developed as a
measure to understand the dynamics of mother-infant relationships in primates. Whenever the
calf's position changed, we recorded the time and whether the change was due to the calf's or
conspecific's movement. An approach (Ap) was recorded when the distance between a calf-
conspecific pair decreased from >1 to ≤1 calf-body length (either due to the calf's action – Ap_c
or conspecific's action – Ap_{cf}), while a leaving (L; L_c when a calf left, and L_{cf} when conspecific
left) was recorded when the calf-conspecific distance increased from ≤1 to >1 calf-body length.
Hinde's proximity index for each calf-conspecific pair was calculated as follows to see whether
calves showed more approaches or leavings towards each conspecific category:

1160
1161   Hinde's index = the percentage of approaches (% Ap_c) made by a young one (calf here) towards
1162   a conspecific – the percentage of leavings made by that young one away from that conspecific
1163   (% L_c).

1164
1165   % Approaches by a calf towards a conspecific (Ap_c)=

1166 $$\frac{\text{No. of approaches made by a calf towards a conspecific}}{\text{Total no. of approaches made by a calf-conspecific pair towards each other}} \times 100$$

1167

1168   % Leavings by that calf away from that conspecific (L_c)=

1169 $$\frac{\text{No. of leavings made by that calf away from that conspecific}}{\text{Total no. of leavings made by the calf-conspecific pair away from each other}} \times 100$$

1170

The index indicates whether the calf more frequently approached or left a particular
conspecific. Hinde's proximity index varies between -100 to +100 (Figure). A value of 0
indicates that the young one makes and breaks proximity contacts with a conspecific equally;
a negative value indicates that the calf more often leaves than approaches the conspecific, and
a positive value that the calf more often approaches than leaves the conspecific. While the total
number of approaches would be similar to the total number of leavings ($Ap = L$ (±1) in a focal)

for a calf-conspecific pair (Hinde and Atkinson 1970), the number of approaches by the calf

(Ap_c) could differ from the number of leavings by the calf (L_c) for the pair.

1179

Brown's proximity index

Hinde's index does not measure the relative contributions of the calf and conspecific to changes

in proximity contacts between the pair. Therefore, we also the calculated Brown's proximity

index (Brown, 2001) to measure the relative contributions of the calf and conspecific to

changes in proximity contacts between the pair. This calculates the total percentage of changes

in proximity contacts that were due to the movement of the calf.

Brown's proximity index $= \dfrac{Ap_c + L_c}{Ap_c + L_c + Ap_{cf} + L_{cf}} \times 100$

where,

Ap_c and L_c are as above,

Ap_{cf} is the number of approaches made by the conspecific towards the calf

L_{cf} is the number of leavings made by the conspecific away from the calf.

1191

1192
1193   Supplementary Material 3, Figure 1. A schematic of a) Hinde's and b) Brown's proximity

1194   indices.

1195

Brown's index ranges from 0 to 100, with 0 indicating that the calf was not responsible for any

of the changes in proximity (numerator will be zero), a value of 50 indicating that young one

and conspecific were equally responsible for changes in proximity contacts, any value between

0 and 50 indicating that the conspecific was more responsible for changes in proximity contacts

than the young one, a value of 100 indicating that the calf was responsible for all the changes

in proximity, and any value between 50 and 100 indicating that the calf was more responsible

for changes in proximity contacts than the conspecific. While Brown's proximity index

measures the relative contribution of calves and conspecifics to the changes in proximity

contacts, it does not differentiate between calves approaching or leaving a conspecific to a

greater extent. Hinde's and Brown's proximity indices complement each other by indicating

whether the calf was responsible for making or breaking contact with a particular conspecific

and whether the calf or conspecific contributed more to the changes in proximity between them.

The values of both the indices may change with the identity of the conspecific (Rowell *et al*.

1964) and the young one's age (Rowell *et al*. 1964, Hinde and Spencer-Booth 1967, Hinde and

Atkinson 1970, Douglas-Hamilton 1972), and the indices are not correlated (Brown 2001). The

relative contributions of the young one versus the conspecific in proximity contact changes

would also differ based on the distance at which approaches and leavings were measured

(Brown 2001).

1214

If a focal calf had more than one escort within a focal sample, the Hinde's index and Brown's

index were averaged across escorts. Similarly, averages were used when a focal calf had more

than one 'other female' with which it was involved in a proximity initiation/termination. As

mentioned in the main text, in certain focals, despite the presence of three conspecific

categories of females in a focal for a focal calf, there was no proximity initiation/termination

between the calf and one or more categories of females. Therefore, average Hinde's and

Brown's proximity indices could be calculated only for 7 calves (*N*: 3 females, 4 males; 3

newborn calves (<3 months), 4 infant calves (3-<6 months)) that each had proximity

initiation/termination with all three conspecific categories of females in at least two of their

focals. There was no significant difference in the Hinde's (Friedman ANOVA: χ^2= 2.000, *N*=7,

df=2, *P*=0.368) or Brown's (Friedman ANOVA: χ^2= 4.571, *N*=7, *df*=2, *P*=0.102) proximity

indices among mothers, escorts, and other females. ANOVAs on the logit Hinde's index with

mothers and escorts based on data from a larger number of calves (17 calves, 44 focals) also

showed no significant effect of conspecific category ($F_{1,16}$=3.849, *P*=0.067). There was no

effect of calf ID ($F_{16,54}$=0.992, *P*=0.479) or its interaction with conspecific category in the

ANOVA ($F_{16,54}$=0.724, *P*=0.757), but inferences cannot be drawn based on these as each calf

was sampled only twice. ANOVA on the logit Brown's index with mothers and escorts (17

calves, 44 focals) also showed no significant effect of conspecific category ($F_{1,16}$=0.888,

P=0.360), calf ID ($F_{16,54}$=1.484, *P*=0.140), or their interaction ($F_{16,54}$=0.365, *P*=0.985).

1234

1235 Supplementary Material 4. Details of behavioural interactions recorded.

1236

As mentioned in the main text, we used focal animal sampling (Altmann, 1974) to record

interactions between calves and conspecifics within groups. We recorded observations using a

video camera (SONY HDR-XR 100E or SONY HDR-PJ 540E). Focal videos were taken in

such a way that a focal calf was always in frame to capture all the calf-conspecific (i.e., subadult

or adult female) interactions that occurred. We also ensured that the focal calf's mother was

also in the frame, except in cases when the calf ventured far away from its mother, in which

case, the mother's behaviours were written down. Calf-initiated interactions, conspecific-

initiated interactions, and responses by conspecifics are detailed in the tables below.

1245

1246

1247 Supplementary Material 4, Table 1. Calf-initiated interactions towards mother, escorts, and

1248 other females, the frequencies with which they appear in the data, behavioural classes, and

1249 descriptions of the interactions.

1250

Code	Interaction	Frequency	Behavioural class	Description
INV	Investigate food	38	Feeding	Investigate another animal's food (in its mouth)/smell the grass that an individual is scraping off, but NOT taking someone else's food (which is TGC).
NDG	Nudge	1	Feeding	Nudge another animal especially when competing for food and feeding very close by and almost always supplanting the animal and feeding there (not dominance related in case of calves).
NPL	Pull at nipple	5	Feeding	Pull or hold nipple.
NPR	Locate nipple correctly	192	Feeding	Try to locate nipple in the correct direction. This includes sniffing in the direction of the nipples also.
NPW	Search for nipple in the	9	Feeding	Try to locate nipple between the hind legs (also includes trying to suck from there).

Code	Interaction	Frequency	Behavioural class	Description
	wrong direction			
SUA	Allosuck	45	Feeding	Suck from the escort/allomother.
SUF	Suck from female	2	Feeding	Suck from a female conspecific other than the mother or escort.
SUM	Suck from mother	137	Feeding	Suck from mother.
TGC	Take grass from a conspecific	89	Feeding	Take grass from a conspecific (and feed on it; PLO if they just throw it away PLO).
TGS	Take grass from a spot	21	Feeding	Feed from the exact spot where another is feeding from (mostly with mothers/escorts).
LEN	Lean	27	Resting	Stand or sit leaning on another animal.
LIE	Lie down	54	Resting	Lie down on land (as opposed to lying in the water, which is bathe BTH).
SLD	Slide	8	Resting	Slide off an individual's leg to lie down.
CHE	Bite	1	Social	Try to chew or bite any body part of another individual.
CHK2	Check	12	Social	Check a conspecific but NOT in dominance.
DST	Express distress	3	Social	Express distress through vocalisations.
EXT	Extend trunk	1	Social	Extend trunk towards a conspecific.
KIC	Kick conspecific	1	Social	Kick a conspecific, even if it misses.
PCL	Play climb	2	Social	Climb/roll on other calves (play).
PSH	Push	16	Social	Push with the head.
PTS	Pass	36	Social	Pass from one side of an older animal to the other through the space between that animal's trunk and forelegs or through the space under its belly.
RUB	Rub	151	Social	Rub against a conspecific (no dominance).

Code	Interaction	Frequency	Behavioural class	Description
SBE	Stand with body contact	46	Social	Stand between the trunk and leg of an escort.
SBM	Stand with body contact	11	Social	Stand between the trunk and leg of the mother.
SNI	Sniff air	62	Social	Sniff the air (could be in the direction of a conspecific or heterospecific).
STE	Stand with escort body contact	46	Social	Stand under the escort
STM	Stand with mother body contact	39	Social	Stand under the mother.
TOU	Touch	129	Social	Touch with trunk in areas apart from the mouth (in mouth would be TRM) and genitals (genitals would be CHK or CHQ), and not in dominance (dominance would be TCH).

1251

1252

1253

1254

 Supplementary Material 4, Table 2. Conspecific- (mother, escorts, and other females) initiated

 interactions towards calves, the frequencies with which they appear in the data, behavioural

 classes, type, and descriptions of the interactions.

| Code | Interaction | Frequency | | | Behav-ioural class | Type | Description |
		Mom	Esc-ort	Other F			
NDG	Nudge	9	21	3	Feeding	Negative non-aggressive	Nudge another animal, especially when competing for food and feeding very close by and almost always supplanting the animal and feeding there (not dominance-related in the case of calves).
KIC	Kick conspecific	0	0	5	Social	Negative aggressive	Kick a conspecific, even if it misses.
LSH	Lash	0	0	1	Social	Negative aggressive	Lash out at a conspecific with the trunk.
NGE	Nudge calf towards escort	2	0	0	Social	Negative non-aggressive	Mother nudges or pushes the calf towards an escort (different from chaperoning because the mother does not walk along with the calf).
PSH	Push	1	0	3	Social	Negative aggressive	Push with the head.
SNI	Sniff air	1	8	1	Social	Neutral	Sniff the air (could be in the direction of a conspecific or heterospecific).
CHK2	Check	0	5	0	Social	Positive	Check a conspecific but NOT in dominance.
CHP	Chaperone	1	8	0	Social	Positive	Follow the calf, or steer the calf while walking, or move in such a way as to place oneself between the calf and males or other group

members during dominance, or wait for and take the calf along while walking.

Code	Interaction	Frequency			Behav-ioural class	Type	Description
		Mom	Esc-ort	Other F			
CPT	Check for conspecific presence	0	1	0	Social	Positive	Check for a conspecific's presence using the tail.
ETM	Escort calf towards mother	0	8	0	Social	Positive	Escort takes the calf to its mother.
KRD	Keep calf within reachable distance	0	3	0	Social	Positive	Keep the calf within reachable distance by stopping it from moving away (when there is nobody else nearby).
RES	Rescue	1	3	0	Social	Positive	Walk/run towards the calf when distress is expressed and reach the calf.
SMA	Smell anus	1	6	2	Social	Positive	Check the calf by smelling its anus.
STG	Stand guard	3	7	0	Social	Positive	Stand guard while the calf sleeps. The calf should be next to at least one of the legs or the trunk. If the female stands over the calf with the calf in between the four legs, it is always counted as stand guard. If the female is standing next to the calf, it is counted as stand guard only if others are not nearby, for e.g., if others are moving away and the female continues to be near the calf. This

| | | Frequency | | | Behav- | | |
Code	Interaction	Mom	Esc-ort	Other F	ioural class	Type	Description
							is to eliminate the case when a calf may be lying down near someone without that individual actively doing anything.
TOU	Touch	5	19	0	Social	Positive	Touch with trunk in areas apart from mouth (in mouth would be TRM) and genitals (genitals would be CHK or CHQ), and not in dominance (dominance would be TCH).
TRM	Trunk in mouth	0	3	0	Social	Positive	Place trunk tip in the mouth.
TRN	Turn towards calf	2	5	0	Social	Positive	Turn towards the calf as the calf experiences distress. Might step forward too but not reach the calf (RES).
WTG	Wait	0	3	0	Social	Positive	Turn towards the direction of and wait for a conspecific to follow, or stop walking and wait for someone to join.
WUC	Wake up calf	6	1	0	Social	Positive	Wake up calf when about to move or when other group members approach.

It must be noted that as in the case of proximity, it was possible, although uncommon, for more than one conspecific to simultaneously interact with the same calf.

Supplementary Material 4, Table 3. Responses received by the calves to the interactions initiated towards mothers, escorts, and other females, the frequencies with which these responses appear in the data, and the type and the descriptions of the responses.

Code	Response	Frequency	Type	Description
AVO	Avoid	38	Negative	Turn away/walk away.
KIC	Kick	4	Negative	Kick a conspecific even if it misses.
LSH	Lash	4	Negative	Lash out at a conspecific with trunk.
MVL	Move leg	99	Negative	Move leg to stop calf from rubbing/leaning/sucking.
NDG	Nudge	13	Negative	Nudge another animal especially when competing for food and feeding very close by and almost always supplanting the animal and feeding there (not dominance related in the case of calves).
PSH	Push	5	Negative	Push with the head.
PTR	Pull trunk	1	Negative	Pull (hold) the trunk and stop the individual from feeding.
SFT	Shift	25	Negative	Shift body/trunk but not move away from the spot; adjust one's position in such a way that it stops the calf's body contact with the recipient.
TLB	Beat with tail	3	Negative	Beat a conspecific with one's tail.
NOR	No response	884	Neutral	No response.
SNI	Sniff air	1	Neutral	Sniff the air (could be in the direction of a conspecific or heterospecific).
ATN	Pay attention	3	Positive	Stand next to the calf and watch it (do something); the individual has to stop whatever it is doing and pay attention to the calf.
CCA	Cooperate with calf	14	Positive	Cooperate with calf action by stopping what one is doing.
CSU	Cooperate with suckling	62	Positive	Move foreleg forward or stop feeding for the calf to suck, or stand without moving once the calf starts sucking.

Code	Response	Frequency	Type	Description
RES	Rescue	1	Positive	Walk/run towards the calf when distress is expressed and reach the calf.
SMA	Smell anus	1	Positive	Check the calf by smelling the anus.
STG	Stand guard	16	Positive	Stand guard while the calf sleeps. The calf should be next to at least one of the legs or the trunk. If the female stands over the calf with the calf in between the four legs, it is always counted as stand guard. If the female is standing next to the calf, it is counted as stand guard only if others are not nearby, for e.g., if others are moving away and the female continues to be near the calf. This is to eliminate the case when a calf may be lying down near someone without that individual actively doing anything.
TOU	Touch	2	Positive	Touch with trunk in areas apart from mouth (in mouth would be TRM) and genitals (genitals would be CHK or CHQ), and not in dominance (dominance would be TCH).
TRM	Trunk in mouth	4	Positive	Place trunk tip in the mouth.
WUC	Wake up calf	4	Positive	Wake up calf when about to move or when other group members approach.

1267

1268

1269 Supplementary Material 5. Effect of calf sex on calf-conspecific proximity.

1270

1271 Supplementary Material 5, Table 1. Results of the nested ANOVA on the logit proportion of

1272 scans that calves spent near the three conspecific categories of females using 5 female and 5

1273 male infant calves (3-<6 months old). Significant *P* values are marked in bold.

1274

Effect	Effect (F/R)	*SS*	*df*	*MS*	*F*	*P*
Calf sex	Fixed	10.354	1	10.354	2.956	0.124
Conspecific category	Fixed	352.577	2	176.289	32.175	**<0.001**
Calf sex*Conspecific category	Fixed	2.358	2	1.179	0.215	0.809
Calf ID (calf sex)	Random	28.020	8	3.502	0.942	0.490
Calf ID (calf sex)*Conspecific category	Random	87.665	16	5.479	1.473	0.140
Error	_	223.134	60	3.719	_	_

1275

1276 Note: Inferences should not be drawn about Calf ID or interactions with this factor as each calf

1277 was sampled in only two focals.

1278

Note: As a calf could have more than one escort and/or other female in a focal, we calculated

the proportions of scans that the focal calf spent near each of the three conspecific categories

of females as the sum of the number of scans near each of the females in that conspecific

category / (total number of females in that conspecific category x 6; where 6 was the number

of total scans in each focal). Since calves could simultaneously be near more than one category

of conspecifics sometimes, the sum of the three proportions could exceed 1.

Supplementary Material 6. Durations of calf-conspecific interactions.

1286

1287

1288

1289

1290   Supplementary Material 6, Figure 1. Frequency distribution of the durations of calf-conspecific

1291   interactions (initiated by either the calf or the conspecific).

1292

1293

1294   Analyses carried out on the durations of interactions (variables: calf-conspecific interactions

1295   and calf-initiated interactions under different behavioural classes) gave similar results as those

1296   on the numbers of calf-conspecific interactions and numbers of calf-initiated interactions under

1297   different behavioural classes. Therefore, they are not reported here.

1298

1299

1300 Supplementary Material 7. Effect of calf sex on the numbers of calf-conspecific interactions.

1301

1302

We ran a nested ANOVA with the log-transformed numbers of calf-conspecific interactions as

the dependent variable, calf sex, initiator category (Calf, Conspecific), and conspecific

category (Mother, Escort, and Other F) as fixed factors, and calf ID nested within sex as a

random factor. We also checked the interaction effects. We could not test the effect of sex for

newborn calves, as there were only 3 female (and 7 male) calves. We ran a nested ANOVA

with only infant calves (3-<6 months), which contained 5 female and 5 male calves. The effect

of calf sex was not significant. However, even these are small sample sizes and it is desirable

to test the effects of calf sex on proximity and behavioural interactions with larger sample sizes

in the future.

1312

1313

1314 Supplementary Material 7, Table 1. Results of the nested ANOVA on the log no. of all calf-

1315 conspecific interactions using 5 female and 5 male infant calves (3-<6 months).

1316

Effect	Effect (F/R)	SS	df	MS	F	P
Calf ID(calf sex)	Random	7.209	8	0.901	2.195	**0.040**
Calf sex	Fixed	0.074	1	0.074	0.082	0.782
Initiator category	Fixed	58.964	1	58.964	133.247	**<0.001**
Conspecific category	Fixed	27.505	2	13.753	35.559	**<0.001**
Calf ID(calf sex)*Initiator category	Random	3.540	8	0.443	1.078	0.391
Calf ID(calf sex)*Conspecific category	Random	6.188	16	0.387	0.942	0.528
Calf sex*Initiator category	Fixed	0.333	1	0.333	0.753	0.411
Calf sex*Conspecific category	Fixed	0.313	2	0.156	0.404	0.674
Initiator category*Conspecific category	Fixed	20.936	2	10.468	37.391	**<0.001**
Calf ID(calf sex)*Initiator category*Conspecific category	Random	4.479	16	0.280	0.682	0.800
Calf sex*Initiator category* Conspecific category	Fixed	0.204	2	0.102	0.365	0.700
Error	_	24.627	60	0.410	_	_

1317

1318 Note: Inferences should not be drawn about Calf ID or interactions with this factor as each calf

1319 was sampled in only two focals.

1320

1321

1322   Supplementary Material 8. Non-suckling interactions.

1323

1324

Of the 1184 calf-initiated interactions, there was a total of 794 non-suckling interactions and

390 suckling interactions towards mothers, escort, and other females. The nested ANOVA on

the number of non-suckling interactions showed almost the same pattern of results as that on

all the interactions. There was a significant main effect of calf age-class, initiator category, and

conspecific category on the log number of non-suckling calf-conspecific interactions (Table 1

below). Newborn calves initiated a significantly higher number of non-suckling interactions

with conspecific females than infant calves (Figure 1 below). Even after removing suckling

interactions, calves continued to initiate a significantly higher number of interactions (average

$\pm$ 95% CI) towards conspecific females (19.85 $\pm$ 3.770) than the latter did towards calves (3.70

$\pm$ 1.634). There was again a significant interaction effect between initiator category and

conspecific category (Table 1 below). Calves initiated a significantly higher number of non-

suckling interactions towards their mothers and escorts than towards other females ($P<0.05$ for

both the comparisons), whereas escorts initiated a higher number of interactions towards calves

than mothers and other females did ($P<0.05$ for both the comparisons). Unlike the case of all

interactions, calves initiated more non-suckling interactions towards escorts than towards their

mothers ($P<0.05$). There was a significant effect of calf identity and also its interaction with

conspecific category, but the other random factors were not significant.

1342

1343

1344 Supplementary Material 8, Table 1. Results of the nested ANOVA on the log numbers of non-

1345 suckling calf-conspecific interactions. Significant *P* values are marked in bold.

1346

Effect	Effect (F/R)	*SS*	*df*	*MS*	*F*	*P*
Calf ID(age-class)	Random	14.521	18	0.807	2.082	**0.010**
Age-class	Fixed	4.596	1	4.596	5.697	**0.028**
Initiator category	Fixed	57.675	1	57.675	103.559	**<0.001**
Conspecific category	Fixed	62.637	2	31.319	44.727	**<0.001**
Calf ID(age-class)*Initiator category	Random	10.025	18	0.557	1.437	0.127
Calf ID(age-class)*Conspecific category	Random	25.208	36	0.700	1.807	**0.009**
Age-class*Initiator category	Fixed	0.952	1	0.952	1.709	0.208
Age-class*Conspecific category	Fixed	2.501	2	1.251	1.786	0.182
Initiator category*Conspecific category	Fixed	19.035	2	9.517	21.867	**<0.001**
Calf ID(age-class)*Initiator category*Conspecific category	Random	15.668	36	0.435	1.123	0.314
Age-class*Initiator category*Conspecific category	Fixed	1.755	2	0.877	2.016	0.148
Error	_	46.503	120	0.388	_	_

1347

1348 Note: Inferences should not be drawn about Calf ID or interactions with this factor as each calf

1349 was sampled in only two focals.

1350

27.9±6.46   19.2±6.39

1351

1352 Supplementary Material 8, Figure 1. Log numbers of non-suckling interactions between

1353 newborn and infant calves and conspecific females. Error bars are 95% CI. Letters above the

1354 data points indicate the pattern of statistical significance based on Tukey's HSD tests (a<b).

1355 The values of the number of interactions (average ± 95% CI) are shown below the graphs.

1356

1357 Supplementary Material 9. Details of focals that had calf-conspecific interactions.
1358

1359

Supplementary Material 9, Table 1. Numbers of focals (out of 40) in which calves and
conspecific females initiated interactions towards each other, and the numbers of all, non-
suckling, and suckling interactions initiated by calves towards the three conspecific categories
of females in these 40 focals. 74.5% of actual suckling was initiated by calves with the mother,
24.5% was initiated with the escorts, which is much higher than that observed in the Amboseli
elephants (3.7% of all suckling bouts with females other than the mother; Lee 1987), and only
1% was initiated with other females.

1367

Variable	Mother	Escort	Other F
No. of focals in which focal calves initiated interactions towards conspecifics	38	40	14
No. of focals in which the conspecifics initiated interactions towards calves	15	25	11
Number of calf-initiated interactions	563	577	44
Number of calf-initiated non-suckling interactions	296	460	38
Number of calf-initiated suckling interactions	267	117	6
Proportion of calf-initiated non-suckling interactions	0.53	0.80	0.86
Proportion of calf-initiated suckling interactions	0.47	0.20	0.14

1368

1369

1370

1371

1372 Supplementary Material 10. Effect of calf sex on the numbers of calf-initiated interactions
1373 under different behavioural classes.

1374

1375

1376 As mentioned in the main text (see Methods), we ran a nested ANOVA with the log-
1377 transformed numbers of interactions under different behavioural classes as the dependent
1378 variable to check the effect of calf sex. As before, we could only test the effect of sex on infant
1379 calves due to sample size constraints. We found that the effect of calf sex was not significant
1380 (Table 1 below).

1381

1382 Supplementary Material 10, Table 1. Results of the ANOVA on the log number of different
1383 behavioural classes of calf-initiated interactions towards the three conspecific categories of
1384 females using 5 males and 5 female infant calves.

1385

Effect	Effect (F/R)	SS	df	MS	F	P
Calf ID(calf sex)	Random	7.013	8	0.877	2.106	**0.043**
Calf sex	Fixed	0.458	1	0.458	0.522	0.490
Behavioural-class	Fixed	27.479	2	13.740	17.371	**<0.001**
Conspecific category	Fixed	39.014	2	19.507	33.393	**<0.001**
Calf ID(calf sex)*Behavioural-class	Random	12.655	16	0.791	1.900	**0.030**
Calf ID(calf sex)*Conspecific category	Random	9.347	16	0.584	1.403	0.158
Calf sex*Behavioural-class	Fixed	0.483	2	0.242	0.305	0.741
Calf sex*Conspecific category	Fixed	0.791	2	0.395	0.677	0.522
Behavioural-class*Conspecific category	Fixed	14.876	4	3.719	9.170	**<0.001**
Calf ID(calf sex)*Behavioural class*Conspecific category	Random	12.979	32	0.406	0.974	0.517
Calf sex*Behavioural-class*Conspecific category	Fixed	1.132	4	0.283	0.698	0.599
Error	_	37.472	90	0.416	_	_

1386

1387 Note: Inferences should not be drawn about Calf ID or interactions with this factor as each calf
1388 was sampled in only two focals.

1389

1390   Supplementary Material 11. Behavioural classes of calf-initiated non-suckling interactions.
1391

Of the calf-initiated interactions, there were a total of 539 feeding interactions (Mother: 328,

Escort: 198, Other F: 13) for a total of 92.83 minutes (Mother: 63.15 minutes, Escort: 27.88

minutes, Other F: 1.80 minutes), 89 resting interactions (Mother: 36, Escort: 50, Other F: 3)

for a total of 198.28 minutes (Mother: 80.48 minutes, Escort: 113.45 minutes, Other F: 4.35

minutes), and 556 social interactions (Mother: 199, Escort: 329, Other F: 28) for a total of

106.88 minutes (Mother: 34.52 minutes, Escort: 70.35 minutes, Other F: 2.02 minutes).

1398

Of the 539 feeding interactions, 390 were suckling interactions, and a majority of them was

with mothers (Supplementary Material 7). As mentioned in the main text (see Methods), we

performed a nested ANOVA with only calf-initiated non-suckling interactions of the three

behavioural classes with conspecific females and found that the pattern of results was the same

as that of all calf-initiated interactions (see Table 2 and Figure 6 in the main text and Table 1

and Figure 1 below). Here again, newborn calves initiated similar numbers of interactions as

infant calves towards conspecific females under the three behavioural classes (Table 1, Figure

1 below). Tukey's HSD tests involving feeding interactions showed a different pattern of

results from that seen in the analysis using all calf-initiated interactions. Calves initiated similar

numbers of feeding and social interactions (95% CI around difference between means for

behavioural class: 0.310; Tukey's HSD: $P>0.05$, Table 2) towards conspecific females when

all the calf-initiated interactions were considered. However, this comparison became

significant when only calf-initiated non-suckling interactions were considered (95% CI around

difference between means for behavioural class: 0.342; Tukey's HSD: $P<0.05$), as a

majority of the calf-initiated feeding interactions were suckling interactions (see

Supplementary Material 7). So, calves initiated a greater number of social interactions than

non-suckling feeding interactions towards conspecific females (Figure 6b). Similarly, calves

initiated a greater number of feeding interactions towards their mothers than towards other

females (95% CI around difference between means for behavioural class x conspecific

category: 0.502; $P<0.05$, Figure 6a). However, this comparison became non-significant when

only calf-initiated non-suckling feeding interactions were considered (95% CI around

difference between means for behavioural class: 0.452; $P>0.05$, Figure 6b), as a majority of

the suckling interactions of calves were with their mothers (see Supplementary Material 7).

1422

1423 Supplementary Material 11, Table 1. Results of the ANOVA on the log number of non-suckling

1424 calf-initiated interactions towards the three conspecific categories of females under different

1425 behavioural classes. Significant *P* values are marked in bold.

Effect	Effect (F/R)	*SS*	*df*	*MS*	*F*	*P*
Age-class	Fixed	0.283	1	0.283	0.434	0.518
Calf ID(Age-class)	Random	11.717	18	0.651	1.872	**0.021**
Behavioural-class	Fixed	53.409	2	26.704	22.801	**<0.001**
Conspecific category	Fixed	43.610	2	21.805	31.139	**<0.001**
Behavioural-class* Conspecific category	Fixed	11.796	4	2.949	7.385	**<0.001**
Age-class*Behavioural-class	Fixed	3.639	2	1.820	1.554	0.225
Age-class*Conspecific category	Fixed	0.390	2	0.195	0.278	0.759
Age-class*Behavioural-class*Conspecific category	Fixed	2.857	4	0.714	1.789	0.140
Calf ID(Age-class)*Behavioural-class	Random	42.162	36	1.171	3.368	**<0.001**
Calf ID(Age-class)*Conspecific category	Random	25.209	36	0.700	2.013	**0.002**
Calf ID(Age-class)*Behavioural-class*Conspecific category	Random	28.749	72	0.399	1.148	0.232
Error	_	62.601	180	0.348	_	_

1426

1427 Note: Inferences should not be drawn about Calf ID or interactions with this factor as each calf

1428 was sampled in only two focals.

1429

1430

Average ± 95% CI no. of interactions	Feeding		Resting		Social	
	2.2 ± 1.87	5.3 ± 4.84	2.8 ± 2.28	1.7 ± 1.22	17.1 ± 5.38	10.7 ± 4.19

1432
1433

1434   Supplementary Material 11, Figure 1. Log numbers of calf-initiated non-suckling interactions

1435   of the three behavioural classes by newborn and infant calves towards conspecific females.

1436   Error bars are 95% CI. NS indicates lack of statistical significance.

1437
1438

1439 Supplementary Material 12. Termination of calf-initiated interactions.

1440

All interactions

Supplementary Material 12, Table 1. Results of the nested ANOVA on the logit proportion of

calf-initiated interactions towards mothers and escorts that were terminated by these females.

The asterisks in the *P* values column indicate significance in the ANOVA on the logit

proportion of calf-initiated non-suckling interactions (see Supplementary Material 12, Table 2)

for comparison.

Effect	Effect (F/R)	*SS*	*df*	*MS*	*F*	*P*
Calf ID(Age-class)	Random	1.121	14	0.080	1.387	0.216*
Age-class	Fixed	0.236	1	0.236	2.953	0.108
Conspecific category	Fixed	0.036	1	0.036	1.314	0.271
Age-class*Conspecific category	Fixed	0.012	1	0.012	0.440	0.518
Calf ID(Age-class)*Conspecific category	Random	0.388	14	0.028	0.480	0.927
Error	_	1.848	32	0.058	_	_

1447

1448

Non-suckling interactions

Of the 794 calf-initiated non-suckling interactions with mothers, escorts, and other females,

calves terminated 563 interactions, and conspecifics terminated 231. There was no difference

between newborn and infant calves in the logit proportions of their non-suckling interactions

terminated by their mothers and escorts (Table 1 below). Mothers and escorts terminated

similar proportions of calf-initiated non-suckling interactions (Table 1, Figure 1 below). There

were insufficient data to examine the effect of calf sex on terminations.

1456

1457 Supplementary Material 12, Table 2. Results of the nested ANOVA on the logit proportion of

1458 calf-initiated non-suckling interactions towards mothers and escorts that were terminated by

1459 these conspecific females. Significant *P* values are marked in bold.

Effect	Effect (F/R)	SS	df	MS	F	P
Calf ID(Age-class)	Random	156.403	12	13.034	2.986	**0.008**
Age-class	Fixed	50.021	1	50.021	3.838	0.074
Conspecific category	Fixed	7.087	1	7.087	0.831	0.380
Age-class*Conspecific category	Fixed	4.301	1	4.301	0.504	0.491
Calf ID(Age-class)*Conspecific category	Random	102.316	12	8.526	1.954	0.071
Error	_	122.205	28	4.364	_	_

1460

1461 Note: Inferences should not be drawn about Calf ID or interactions with this factor as each calf

1462 was sampled in only two focals.

1463

1464

1465

1466 Supplementary Material 12, Figure 1. Logit proportions of calf-initiated non-suckling

1467 interactions towards their mothers and escorts that were terminated by the calves and by the

1468 conspecific females. Error bars are 95% CI. Letters above the data points indicate pattern of

1469 statistical significance. Proportions of calf-initiated non-suckling interactions terminated by

1470 calves and conspecific females are written below the graph (average ± 95% CI).

1471

1472

1473 Supplementary Material 13. Types of responses from conspecific females for calf-initiated

1474 interactions.

1475

All interactions

Supplementary Material 13, Table 1. Results of the ANOVA on the logit proportions of all

calf-initiated interactions towards mothers and escorts that elicited a positive or a neutral

response. Significant *P* values are marked in bold. The asterisks in the *P* values column indicate

significance in the ANOVA on the log numbers of calf-initiated non-suckling interactions (see

Supplementary Material 13, Table 2) for comparison.

1482

Effect	Effect (F/R)	*SS*	*df*	*MS*	*F*	*P*
Calf ID(Age-class)	Random	17.477	14	1.248	0.241	0.997
Age class	Fixed	1.363	1	1.363	1.092	0.314
Response type	Fixed	766.890	1	766.890	43.888	**<0.001***
Conspecific category	Fixed	0.002	1	0.002	0.002	0.966
Calf ID(Age-class)*Response type	Random	244.635	14	17.474	3.374	**<0.001***
Calf ID(Age-class)*Conspecific category	Random	13.358	14	0.954	0.184	0.999
Age class*Response type	Fixed	76.427	1	76.427	4.374	0.055
Age class*Conspecific category	Fixed	0.273	1	0.273	0.286	0.601
Response type*Conspecific category	Fixed	6.574	1	6.574	1.117	0.298
Age class*Response type*Conspecific category	Fixed	1.882	1	1.882	0.320	0.581
Calf ID(Age-class)*Response type*Conspecific category	Random	82.386	14	5.885	1.136	0.345
Error	_	331.429	64	5.179	_	_

1483

Non-suckling interactions

After removing the suckling interactions, of the 756 calf-initiated interactions towards mothers

and escorts (Supplementary Material 7), there was a positive or a negative response for 174 of

the interactions. Similar to the analysis using the full data, we found that there was a significant

effect of response type and the interaction between calf ID and response type (Table 2 below,

Figure 1 below). Mothers (0.36% of the 563 calf-initiated interactions towards mothers) and

escorts (1.39% of the 577 calf-initiated interactions towards escorts) showed a much smaller

proportion of aggressive responses than other females (15.91% of the 44 calf-initiated

interactions towards other females) towards the calf-initiated interactions.

None of the other effects was significant.

Supplementary Material 13, Table 2. Results of the ANOVA on the logit proportions of non-

suckling calf-initiated interactions towards mothers and escorts that elicited a positive or a

neutral response. Significant *P* values are marked in bold.

1497

Effect	Effect (F/R)	*SS*	*df*	*MS*	*F*	*P*
Calf ID(Age-class)	Random	41.365	12	3.447	0.600	0.833
Age class	Fixed	9.395	1	9.395	2.726	0.125
Response type	Fixed	784.004	1	784.004	40.673	**<0.001**
Conspecific category	Fixed	1.993	1	1.993	0.888	0.365
Calf ID(Age-class)*Response type	Random	231.309	12	19.276	3.355	**0.001**
Calf ID(Age-class)*Conspecific category	Random	26.932	12	2.244	0.391	0.961
Age class*Response type	Fixed	74.367	1	74.367	3.858	0.073
Age class*Conspecific category	Fixed	3.073	1	3.073	1.369	0.265
Response type*Conspecific category	Fixed	10.609	1	10.609	1.136	0.294
Age class*Response type*Conspecific category	Fixed	1.880	1	1.880	0.201	0.662
Calf ID(Age-class)*Response type*Conspecific category	Random	112.050	12	9.338	1.625	0.111
Error	_	321.774	56	5.746	_	_

1498

1499 Note: Inferences should not be drawn about Calf ID or interactions with this factor as each calf

1500 was sampled in only two focals.

1501

1502

1503 Supplementary Material 13, Figure 1. Logit proportions of calf-initiated non-suckling

1504 interactions towards mothers and escorts that elicited a positive, neutral, and negative responses

from the conspecific females. Error bars are 95% CI. Letters above the data points indicate patterns of statistical significance (a<b). Proportions of calf-initiated non-suckling interactions that elicited a positive, neutral, and negative responses from the conspecific females are written below the graph (average ± 95% CI).

References

1. Brown, G. R. (2001). Using proximity measures to describe mother-infant relationships. *Folia Primatologica*, *72*(2), 80–84. https://doi.org/10.1159/000049926

2. Douglas-Hamilton, I. (1972). *On the ecology and behaviour of the African elephant: The elephants of Lake Manyara*. Ph.D. thesis, University of Oxford, Oxford.

3. Hinde, R. A., & Atkinson, S. (1970). Assessing the roles of social partners in maintaining mutual proximity, as exemplified by mother-infant relations in rhesus monkeys. *Animal Behaviour*, *18*(1), 169-176. https://doi.org/10.1016/0003-3472(70)90087-4

4. Hinde, R. A., & Spencer-Booth, Y. (1967). The behaviour of socially living rhesus monkeys in their first two and a half years. *Animal Behaviour*, *15*(1), 169–196. https://doi.org/10.1016/S0003-3472(67)80029-0

5. Rowell, T. E., Hinde, R. A., & Spencer-Booth, Y. (1964). "Aunt"-infant interaction in captive rhesus monkeys. *Animal Behaviour*, 12(2–3), 219–226. https://doi.org/10.1016/0003-3472(64)90004-1