Pervanadate-induced oxidation relieves autoinhibition of SRC protein tyrosine kinase

Katie E. Mulholland[1], Maxime Bourguet[2], Nuo Cheng[3], Oisharja Rahman[1], Daria Ezeriņa[4,5,6], Leonard A. Daly[7], Tiffany Lai[1], Silvia Aldaz Casanova[1], Pau Creixell[3], Claire E. Eyers[7], Joris Messens[4,5,6], Patrick A. Eyers[8], Dominic P. Byrne[8*], Hayley J. Sharpe[1*].

Affiliations:
[1] Signalling Programme, Babraham Institute; Cambridge, UK
[2] MRC Laboratory of Molecular Biology; Cambridge, UK
[3] CRUK Cambridge Institute; Cambridge, UK
[4] VIB-VUB Center for Structural Biology, Vlaams Instituut Voor Biotechnologie; Brussels, Belgium
[5] Brussels Center for Redox Biology, Vrije Universiteit Brussel; Brussels, Belgium
[6] Structural Biology Brussels, Vrije Universiteit Brussel; Brussels, Belgium
[7] Centre for Proteome Research, Institute of Systems, Molecular and Integrative Biology, University of Liverpool; Liverpool, UK.
[8] Department of Biochemistry, Cell and Systems Biology, Institute of Systems, Molecular and Integrative Biology, University of Liverpool; Liverpool, UK.

*Corresponding authors. Email: hayley.sharpe@babraham.ac.uk and d.byrne@liverpool.ac.uk.

Highlights

- Pervanadate disrupts redox homeostasis promoting cysteine oxidation
- Cysteine 188 and 280 of SRC are sequentially oxidized by pervanadate
- Oxidation leads to conformational relief of SRC autoinhibition
- SRC cysteines are essential for overcoming contact inhibitory Hippo signaling

Summary
Dynamic regulation of protein tyrosine phosphorylation (pTyr) by phosphatases (PTPs) and kinases enables cells to sense and respond to environmental changes. The widely used chemical probe Pervanadate (PV) induces accumulation of high levels of pTyr in cells, an effect primarily attributed to its properties as a PTP inhibitor. This led to the assertion that PTPs are the master gatekeeper of intracellular pTyr homeostasis. Here, we use diverse approaches to reveal that PV potently and directly activates SRC family tyrosine kinases *via* oxidation of specific cysteine residues. Using mass spectrometry and biophysical approaches, we show that oxidation activates SRC by disrupting autoinhibition and altering phosphopeptide binding by its SH2 domain. We further establish that redox-sensitive cysteine residues are essential for SRC to promote cellular overgrowth. Our findings call for a re-evaluation of PV-based experiments and provide compelling evidence that oxidation is a crucial mechanism in controlling the oncogenic properties of SRC.

Key words:
Redox, kinase, phosphatase, pervanadate, SRC, phosphotyrosine, signaling, mass spectrometry

Introduction

Phosphorylation of protein tyrosine residues (pTyr) is a critical posttranslational modification that orchestrates diverse cellular processes[1] and is often dysregulated in disease[2,3]. Dynamic pTyr levels are controlled by the activities of protein tyrosine phosphatases (PTPs) and kinases (PTKs). Disruption of both PTPs and PTKs can lead to pathological conditions, highlighting the importance of understanding their regulation in homeostasis and disease. The development of specific PTP inhibitors has been challenging, with only a few compounds advancing to clinical trials[4,5]. As a result, generic PTP inhibitors remain invaluable in gaining insight into pTyr functions and regulation. Of these, orthovanadate (OV), commonly added to cell lysis buffers, and pervanadate (PV), have become central tools[6]. OV acts as a phosphate mimetic that competitively inhibits PTPs, while PV is produced by combining hydrogen peroxide (H_2O_2) with OV, followed by the addition of catalase to remove excess H_2O_2 (**Figure 1A**)[6,7].

PV exists in multiple oxidation states, containing up to three peroxogroups[8,9]. In cells, PV treatment rapidly elevates pTyr levels, enabling PTP substrates to be analyzed through 'trapping' and dephosphorylation approaches[10]. PV is established to inhibit PTPs by hyperoxidizing the low pKa catalytic cysteine to sulfonic acid[11]. Due to the structural similarity between phosphate and vanadate, it has been proposed that PV exhibits high affinity for the PTP active site[6], leading to the conclusion that its effects are predominantly driven through PTP inhibition[12]. Moreover, the capacity of PV to reproduce key cellular activation states has fostered the prevailing view that PTPs are constitutively active, constantly removing pTyr from proteins[13]. Furthermore, this concept is a key pillar in the kinetic segregation model, which proposes that steric exclusion of the highly abundant receptor PTP, CD45, is sufficient to trigger T cell activation[14], based on experiments showing that PV mimics T cell activation[15,16].

Cellular effects of PV are often indirectly linked to PTK activation, since PTPs are assumed to suppress activity[12]. However, PTP function is more complex, with several PTPs, including CD45, activating PTKs by dephosphorylation of inhibitory phosphosites[17,18]. Furthermore, the catalytic activities of certain PTKs, including EGFR and SRC, are enhanced following reversible oxidation of conserved regulatory cysteine residues[19-24]. Given that PV possesses peroxo groups with oxidizing potential, we hypothesized that its cellular effects on pTyr levels might extend beyond PTP inhibition to include direct oxidation of redox-sensitive cysteines in other proteins[25,26], including PTKs. Here, we show that the SFKs SRC and YES1 are directly activated by PV. Using protein dynamics analysis and biophysical approaches, we elucidated the mechanisms by which PV-induced oxidation relieves SRC autoinhibition. Additionally, we identified a key role for redox-active cysteines in SRC-regulated cell growth and control of Hippo signaling. Our study defines new redox-based mechanisms for SRC activation and highlights the importance of cysteine oxidation in oncogenic transformation. In addition, our data call for a mechanistic reinterpretation of the effects of pervanadate on phosphotyrosine and redox proteomes.

Figure 1

88
89
Figure 1. PV perturbs redox homeostasis and induces tyrosine phosphorylation in SRC
dependent manner.
(A) Schematic of mono-peroxo and di-peroxo-vanadium (PV compounds) generated using
ChemDraw. **(B)** HEK293T cells were treated as shown, lysed and subjected to immunoblot
analysis using the indicated antibodies. Image representative of N=3 experiments. (**C**)
Representative false colour images of F488/405 ratio (scaled 1-9) from **videos S1 and S2** of
U2OS cells stably expressing plasma membrane (PM) localised HyPer7 and treated with 100
µM PV or media. Scale bar = 20 µm. (**D**) Quantification of the ratio of the two excitation maxima

(F488/405), normalized against initial value, of PM-HyPer7 expressing U2OS cells over a time
course, imaged every 20 s. Treatments were added at 60 s (arrow). Mean of n=3 ± SD plotted.
** $p \leq 0.001$ Unpaired t-test at t= 400 s. (**E**) Ratio of the two excitation maxima (F500/400) of
recombinantly expressed HyPer7 (2.5 µM) following its treatment with the indicated reagents.
The bars represent the mean of n=3 ± SD. *** $p \leq 0.001$; ns – non-significant. The p-value is
based on Dunnett's multiple comparisons test of a one-way ANOVA. (**F**) Quantification of the
ratio of the two excitation maxima (F488/405), normalized against initial value, of PM-HyPer7
expressing U2OS cells over a time course, imaged every 20 s. Treatments were added at 60
s (arrow). Mean of n=3 ± SD plotted. (**G**) Cells were subjected to the indicated PV time course,
washed with PBS with catalase and NEM, prior to lysis without NEM. Samples were resolved
on non-reducing or reducing gels as indicated to distinguish PRDX1 monomers from disulfide-
linked dimers. Total and phospho PRDX1 blots are from separate gels but the same samples.
Representative of n=3 experiments. (**H**) Densitometric quantification of PRDX1 dimer and
monomer bands from G. Mean of n=3 ± SD plotted. (**I**) HEK293T cells were treated with the
indicated tyrosine kinase inhibitors at IC_{50} (**+**) and 2.5x IC_{50} (**‡**) concentrations for 2 h prior to
addition of 100 µM PV for 15 min before lysis and immunoblotting with annotated antibodies.
(**J**) Dasatinib targets based on[71] and mapped onto the tyrosine kinase arm of the Human
kinome using kinmap (http://kinhub.org/kinmap/). (**K**) WT and Src/Yes/Fyn (SYF) WT and KO
cells were stimulated with 100 µM PV for the indicated times prior to lysis and immunoblotting
with indicated antibodies. Red arrow indicates 130 kDa. Representative of 3 experiments.
118
119　See also Figures S1-S3.
120

Results

Pervanadate promotes cysteine thiol oxidation

We initially set out to characterize cellular effects of the PTP inhibitors PV and OV. As expected, PV induced a rapid pTyr accumulation in HEK293T cells with concomitant irreversible PTP oxidation (OxPTP) (**Figure 1B**). In contrast, the PTP inhibitor OV had minimal effect on global pTyr despite the induction of irreversible PTP oxidation at higher doses (**Figure 1B**). Longer treatment (4h) with high doses of OV, resulted in more pronounced upregulation of pTyr and PTP hyperoxidation, consistent with previous reports of oxidative damage by OV[27] (**Figure S1A**). We next asked whether our OV preparations inhibit PTPs. First, we confirmed that OV inhibits the phosphatase activity of recombinant purified PTPRK *in vitro*, albeit less potently than PV (**Figures S1B-S1C**), aligning with previous reports[28]. Furthermore, Flag-PTPRK immunoprecipitated from OV or PV treated cells was catalytically inactive (**Figures S1D-S1E**). Importantly, and in contrast to PV, OV did not induce extensive tyrosine phosphorylation at this concentration (**Figures S1F-S1G**). Collectively these data suggest that PTP inhibition alone is necessary, but not sufficient, for high-level global induction of pTyr. We next explored the possibility that PV could have functions beyond PTP inhibition that promote supraphysiological levels of pTyr accumulation.

Since the pTyr-inducing function of PV is usually attributed to PTP oxidation, we investigated whether it has broader oxidizing activities in cells. To this end, we used HyPer7, a ratiometric biosensor that is oxidized by peroxides and reduced by thioredoxins and reduced glutathione (GSH) within cells[29-31]. Treating U2OS cells expressing myristoylated, plasma membrane-anchored HyPer7 with 100 µM PV led to rapid biosensor oxidation compared to media-treated cells (**Figures 1C-1D and videos S1-S2**). In contrast, treatment of cells with 50-500 fold higher levels of OV slightly decreased the ratio of oxidized:reduced HyPer7 compared to media alone (**Figure S1H**). *In vitro*, PV directly oxidized purified recombinant HyPer7, although far less potently than H_2O_2 (**Figure 1E**). By comparison, both 20 µM PV and H_2O_2 rapidly oxidized HyPer7 in cells equivalently (**Figure 1F**). Thus, despite lower direct reactivity with HyPer7, PV is as effective as H_2O_2 (at 20 µM) at inducing HyPer7 oxidation in cells. This indicates that PV elicits its effects on HyPer7 in cells by perturbing antioxidant pathways, such as the thioredoxin pathway.

We next performed a time course of PV treatment and monitored the oxidation state of peroxiredoxin 1 (PRDX1), an endogenous substrate of the thioredoxin pathway. Upon oxidation by H_2O_2, PRDX forms disulfide-linked dimers, which are detectable as slower migrating species after non-reducing SDS-PAGE. To block post-lysis oxidation, samples were treated with catalase and N-ethyl maleimide (NEM). Within 30s, PV increased PRDX1 disulfide-linked dimers that were undetectable under reducing conditions (**Figures 1G, 1H and S1I**). We also detected a time dependent increase in PRDX1 phosphorylation (pY194), which is associated with impaired PRDX catalytic function[32]. Together these data suggest that PV fundamentally perturbs redox homeostasis. Strikingly, PTPs were irreversibly oxidized 30 s post PV treatment, several minutes before the peak in global tyrosine phosphorylation is evident, suggesting sustained or induced PTK activity. Since several kinases also possess redox-active cysteines[19-24], we next focused on the role of PTKs in driving the PV pTyr response.

PV-induced pTyr depends on specific PTKs, including SRC

To determine which PTKs might contribute to PV-induced increases in pTyr, we took an
unbiased approach pre-treating cells with a panel of pharmacological PTK inhibitors. PV was
equally effective at oxidizing PTPs in cells treated with the inhibitors but only dasatinib and
saracatanib profoundly limited PV-induced pTyr levels (**Figures 1I and S2**). Dasatinib and
saracatanib inhibit SFKs, ABL and EPH kinases (**Figure 1J**), suggesting they may contribute
to PV-induced pTyr signaling. SRC activation has previously been linked to PV-induced pTyr
signaling[12,33], so we employed genetic manipulation to validate the role of SFKs in this process.
We observed an attenuated pTyr response when treating Src/Yes/Fyn knockout (KO) MEFs
compared to WT cells when treated with 100 μM PV, despite equivalent levels of PTP
oxidation (**Figure 1K**). We also detected increased Abl1 expression in KO MEFs (**Figure
S3A**), which could explain the hyperphosphorylated protein band at ~130 kDa induced by PV
(**Figure 1K, red arrow**). However, knockdown of Abl1 in SYF KO MEFs did not further
attenuate the PV response or affect the intensity of the 130 kDa band (**Figure S3B**). Knocking
out SRC in HAP1 cells also decreased the levels of pTyr induced in response to PV (**Figure
S3C**). The knockdown of SRC and YES1, but not FGR, in HEK293T cells led to reduced PV
stimulated pTyr (**Figure S3D**). Thus, SFKs are key mediators of the cellular PV response.

Figure 2

184
Figure 2. SRC is oxidized on specific cysteines and activated by PV.
(**A**) 6xHis-SRC (3 ng) dependent phosphorylation of a specific fluorescent substrate (2 µM).
Data shown is relative SRC activity in the presence of the indicated concentrations of PV
compared to a buffer only control. SRC was preincubated for 0, 10, 60 or 120 mins prior to
initiation of the kinase assay. Means ± SD n=3 experiments are shown. (**B**) Dose response
curve for H_2O_2 pre-incubated with recombinant 6xHis-SRC (3 ng) for the indicated time
periods. Reactions were initiated with the addition of peptide substrate and ATP, and
phosphorylation was measured after 15- and 25-mins assay time. Data shown is relative SRC
activity in the presence of the indicated concentrations of H_2O_2 compared to a buffer only
control. Means of n=3 ± SD shown. (**C**) HEK293T cells treated with the indicated
concentrations of PV for 15 min, were lysed in catalase and NEM-containing buffer. Reversibly

oxidised thiols were reduced with TCEP, and newly regenerated thiol(ate)s were labelled with
Iodoacetyl-PEG2-Biotin. Lysates and biotinylated proteins were then affinity purified using
streptavidin beads and subjected to immunoblotting with indicated antibodies. Blot
representative of 3 experiments. (**D**) HEK293T cells transiently expressing StrepII-tag fused
SRC were treated as indicated, and SRC affinity purified by streptactin pull down. Purified
protein was then subjected to tryptic digest and analysed by LC/MS-MS for triply oxidised
cysteine (sulphonic acid). (**E**) SRC isolated from transfected HEK293T cells by streptactin
affinity purification was incubated with or without 5 mM PV, followed by tryptic digest and
analysis by LC MS/MS. Values are normalized against abundance of non-cysteine containing
SRC peptides. (**F**) Schematic of SRC kinase indicating the position of two key regulatory
phosphosites (Circle with P) as well as the two most readily oxidixed cysteine residues (red
stars). (**G**) Surface electrostatic potential[72] for regions surrounding Cys 280 (PDB:2SRC) and
Cys 188 and Cys241 (PDB:1Y57). Units = K_bT/e_c. (**H**) Phosphorylation of the fluorescent
peptide substrate by WT (black), C188A (blue) and C280A (red) SRC isolated from HEK293T
cells by streptactin affinity purification was measured after 30 min. Means of n=3 ± SD are
shown. (**I**) Dose-response curves for PV. WT (black), C188A (blue) and C280A (red) SRC
isolated from HEK-293T cells by streptactin affinity purification and incubated with the
indicated concentrations of PV for 10 mins. Phosphorylation of the fluorescent peptide
substrate was measured after 15 min. Data shown is % SRC activity compared to a buffer
control. Means of n=3 ±SD are shown. FL = full-length. (**J**) SYF KO MEFs were retrovirally-
transduced with specified SRC expression constructs and treated with PV as shown, before
lysis and immunoblotting with indicated antibodies. Red arrow indicates relevant band.
Representative of >3 experiments. (**K**) Densitometric quantification of western blot in Figure
2J. Ratios of phosphorylated cortactin/total tubulin levels are shown.
220
221   See also Figure S3.
222

Pervanadate activates SRC via cysteine oxidation *in vitro* and in cells

We hypothesized that PV induces potent pTyr accumulation in cells by directly activating SFKs. We focused our initial experiments on SRC since its redox sensing via specific cysteines is the best characterised amongst SFKs[20]. Using purified recombinant SRC, we determined that PV dose-dependently increased SRC kinase activity *in vitro* (**Figure 2A**). Introducing a pre-incubation period increased activation at lower PV concentrations but decreased catalytic activity at higher PV concentrations (**Figures 2A and S3E**). OV treatment had no effect on SRC activation (**Figure 2A**). Similar activity profiles have been reported for SRC and EGFR in cells treated with H_2O_2[21,22], where higher peroxide concentrations were inhibitory potentially due to oxidation of additional cysteines. In support of a redox mechanism, we also observed H_2O_2 stimulation of SRC activity, but only at higher concentrations than PV (**Figures 2B**). To test whether PV induces cellular SRC oxidation, we used differential alkylation to reveal TCEP-reversible thiol modifications by biotin labelling, or 'biotin switch'. At 0-250 µM PV, SRC and the PTP SHP2 were increasingly enriched by streptavidin, indicating increased reversible oxidation (**Figure 2C**). Higher PV concentrations decreased streptavidin-enrichment of both proteins, suggesting thiols had either become reduced, or irreversibly oxidized such that they are resistant to reduction and alkylation. These trends were recapitulated widely as indicated by the enrichment of numerous biotinylated, reversibly oxidized, proteins. Notably, we observed oxidation of both SRC and SHP2 in the absence of PV, indicating a basal level of oxidative signaling in HEK293T cells, consistent with the presence of PRDX1 dimers in untreated cells (**Figure 1G**). Together, these data demonstrate reversible oxidation of SRC by PV in cells.

To gain insight into the molecular mechanism(s) of SRC oxidation, we used a targeted approach to identify cysteines oxidized by PV. At high concentrations of PV, SRC cysteine thiols appeared to be hyperoxidized to sulfinic or sulfonic acid (**Figure 2C**), which are identifiable by mass spectrometry (MS) without the need for differential alkylation. We therefore stimulated cells overexpressing SRC with 5 mM PV for 30 min and analyzed affinity purified protein by MS (**Figure 2D and Table S1**). Peptides containing Cys 188 and Cys 403 were identified, however, only Cys 188 showed a large increase in sulfonic acid modification following PV treatment. We next analyzed purified, untagged SRC treated with PV, which identified sulfonic acid modification of cysteines 188, 280 and 241 (**Figure 2E and Table S2**). SRC Cys 241 was less sensitive to hyperoxidation by PV than Cys 188 and Cys 280 (**Figure 2E**), indicating distinct accessibility or reactivity of individual cysteines[34]. SRC comprises N-terminal, SRC homology (SH)3, SH2 and kinase domains (**Figure 2F**). Cys 188 is located within the SH2 domain pTyr binding pocket and is unique to SRC amongst SFKs, whilst Cys 280 resides in the kinase domain P-loop of SRC, YES1 and FGR (**Figures 2F and S3F**). Both Cys 280 and Cys 188 are found close to or within positively charged pockets on the surface of SRC (**Figure 2G**). In contrast, Cys 241, which was not oxidized by PV, is surface exposed in an electrostatically neutral region of the SH2 domain (**Figure 2G**). Together this reveals selective oxidation of Cys 188 and 280 by PV.

The conservation of Cys 280 in SFKs prompted us to also evaluate sensitivity of YES1 activity to oxidation, especially given its role in the PV response in cells (**Figure S3D**). Like SRC, PV dose dependently increased YES1 activity (**Figure S3G**). In contrast, H_2O_2 had a negligible impact on activity (**Figure S3H**), indicating differences in sensitivity to each oxidant. To directly assess the impact of Cys 188 and Cys 280 on SRC kinase activity *in vitro*, we affinity purified untagged WT (SRC^{WT}) and mutant (SRC^{C188A} and SRC^{C280A}) forms. All three proteins

displayed equivalent kinase activity (**Figure 2H**). However, when treated with PV, a dose-
dependent increase in SRC[WT] activity was evident, which was absent for SRC[C280A] and
impaired for SRC[C188A] (**Figure 2I**). At higher PV concentrations, all proteins were inhibited,
potentially indicating oxidation of additional cysteines. Thus, PV-induced SRC activation
requires Cys 188 and Cys 280. We next asked whether these cysteine residues are important
for the cellular pTyr response to PV. To this end, we introduced chicken Src and mutants
equivalent to C188A and C280A into SYF KO MEFs. Human SRC residue numbering is used
hereafter for clarity and consistency. In response to PV, reintroduction of SRC[WT] increased
global pTyr levels and phosphorylation of the SRC substrate cortactin, however, this was
impaired for SRC[C188A] and SRC[C280A] indicating a role for these cysteines in driving signaling in
response to PV (**Figures 2J and 2K**). Importantly, PTP catalytic cysteine hyperoxidation
detected by oxPTP was comparable in all cell lines. Together, our data suggest that oxidation
of Cys 188 and 280 by PV are important for activating SRC and driving cellular increases in
pTyr-containing substrates.
285

Figure 3

A 0.5 mM PV

Closed conformation (2SRC)   Open conformation (1Y57)

B 2.5 mM PV

Closed conformation (2SRC)   Open conformation (1Y57)

Figure 3. PV induces specific conformational changes in SRC.
(A, B) HDX-MS analysis of pervanadate treated SRC. Relative fractional uptake (RFU) differences between 5 µM SRC and SRC incubated with 0.5 mM (**A**) and 2.5 mM (**B**) PV were mapped on the closed (Left) and the open conformations (Right) of SRC (PDB: 2SRC and 1Y57, respectively). Regions showing significant protection and becoming less solvent exposed are colored in blue and regions showing significant deprotection and becoming more solvent exposed are presented in red. Regions where no significant differences were observed are shown in white, while grey denotes parts of SRC for which no peptides were observed. Key cysteines are highlighted in purple and key phosphorylation sites are highlighted in light blue. The overall coverages were 80% and 60% for the 0.5 mM and 2.5 mM PV HDX-MS experiments, respectively.

See also Figure S4.

PV triggers conformational changes that relieve SRC autoinhibition

We next performed a mechanistic investigation into SRC activation by oxidation. Previous molecular dynamics simulation suggested that cysteine oxidation plays a role in relieving autoinhibition[21], a key step in SRC activation. SRC autoinhibition is mediated by intramolecular contacts between SH2 and SH3 domains and the kinase domain[35,36], with extensive conformational changes occurring during activation. To evaluate whether oxidation promotes conformational changes in SRC, we initially used a thermal shift assay using affinity purified proteins. ATP and Dasatinib stabilized full-length recombinant SRC, whereas PV and H_2O_2 led to destabilization, indicating oxidation-induced conformational changes (**Figure S4A**). To obtain a detailed molecular understanding of conformational changes caused by PV, we used hydrogen-deuterium exchange MS (HDX-MS) on full-length human SRC. The relative deuterium uptake differences were mapped onto the 'closed' (2SRC[37]) and 'open' (1Y57[38]) SRC conformations (**Figures 3A-3B**). At 0.5 mM (100x molar ratio of PV to SRC), PV increased solvent exposure of the SH2-kinase domain linker, which spans the polyproline-II helix. This helix engages the SH3 domain in the autoinhibited conformation and is released upon activation (**Figures 3A *right,* and S4B**) [35,36]. In the SH2 domain, PV induced exposure of the αA helix while protecting the βB-βC loop, indicating redox-sensitive conformational changes near Cys 188. This suggests that 0.5 mM PV drives two key structural alterations involved in SRC activation.

At 2.5 mM (500x molar ratio of PV to SRC), PV induced extensive solvent exposure across SRC regions associated with conformational transition from inactive to active kinase (**Figure 3B**). The RT loop and the βE strand within the SH3 domain showed increased exchange, indicating release from the polyproline-II linker[39], a more pronounced effect than with 0.5 mM PV (**Figure 3A**). Changes in the SH2 domain were also more prominent at this higher PV concentration (**Figures 3B and S4C**). In the linker between the SH3 and kinase domains, a region spanning Trp 263 was exposed by PV. This key conserved residue is involved in forming a new β turn during the transition from closed to open conformations[40]. The activation loop, including Tyr 419, was also more exposed aligning with previous MD simulations predicting Cys 280 oxidation-induced unfolding of this loop[21]. The catalytic loop HRD motif and the αEF helix, which stabilize the activation segment in an active state, were also more exposed upon PV treatment[37,38,41] (**Figure 3B, *right***). The C-terminal tail, including Tyr 530, was more accessible along with the αD helix (**Figure 3B, *right***). Overall, PV induced conformational changes that increased flexibility and solvent accessibility, explaining the oxidation-induced destabilization observed by thermal shift (**Figure S4A**). These effects were increased at 2.5 mM PV, consistent with the dose-dependent effects observed in kinase assays (**Figure 2A**). Together, these results support a model in which PV drives the relief of SRC autoinhibition via specific conformational alterations across the polypeptide. Furthermore, aligning with the greater sensitivity of Cys 188 (**Figures 2E and 2F**), at lower levels of oxidation we observed conformational changes largely in the vicinity of Cys 188, with putative effects mediated by Cys 280 requiring increased amounts of PV. This supports a hierarchy of cysteine PV sensitivity within SRC.

Figure 4

345
Figure 4. **Cys 188 oxidation or mutation impacts SRC-SH2 phosphopeptide binding.**
(**A**) SRC-SH2 with highlighted regions protected (blue) and deprotected (red) from deuterium
exchange upon 0.5 mM PV treatment (PDB:2SRC) based on data from Fig. 3. (**B**) Alphafold3[42]
model of SRC SH2[WT] bound to the phosphorylated SRC C-terminal peptide:
FTSTEPQpYQPGENL. ipTM = 0.73. (**C**) Residues predicted to form hydrogen bonds (blue
dashed lines) based on AF3 models. Top: As in B. Middle: AF3 model with SRC s-nitroso
cysteine 188 and C terminal peptide. ipTM = 0.75. Bottom: AF3 model with C188A and C
terminal peptide. ipTM = 0.72. (**D**) Isothermal titration calorimetry (ITC) titration curves of the
interaction between SRC-SH2 and indicated phosphopeptides. The SH2 domain was
pretreated with 1 mM TCEP or 0.1 mM PV prior to the assay. Upper: baseline-corrected
differential power plotted over time. Lower: normalized binding curve showing the integrated
change in enthalpy against the molar ratio. Stoichiometry (N), mean enthalpy change (ΔH)
and mean K_D ±SD shown based on n =3 assays. (**E**) AF3 model of hydrogen bonding network
between residues within the JAK1 SH2 domain engaging glutamate on IFNR1. ipTM = 0.69.
360
361 See also Figure S5.
362

Cys 188 oxidation impacts SRC-SH2 affinity for phosphopeptides

We next focused on how the conformational changes within the SRC SH2 domain revealed by our HDX-MS data may contribute to relief of autoinhibition (**Figure 4A**). Cys 188 resides within the pTyr binding pocket and is ideally placed to mediate SRC interactions with phosphopeptides (**Figures 4A-4B**). We used AlphaFold3 to explore the binding of SRC-SH2 domain with the phosphorylated SRC C-terminal peptide, which interact in the autoinhibited state[42]. Cys188 is predicted to participate in a hydrogen bonding network that engages the phosphate group of the bound peptide (**Figures 4C, *Top* and S5A**). To mimic the effects of oxidation, we modelled Cys188 with an S-nitroso modification in AlphaFold3 (**Figures 4C, *middle* and S5B**). We infer that Cys oxidation causes extensive remodelling of the hydrogen-bonding network, reducing the overall number of contacts with the phosphate group. In contrast, modelling C188A predicts hydrogen bonding network adjustments that incorporate new interactions, including with Thr 182 (**Figures 4C, *bottom* and S5C**), which might impact phosphopeptide binding kinetics.

We next used isothermal titration calorimetry (ITC) to quantify effects of PV on phosphopeptide interaction dynamics using WT and C188A SRC-SH2 domains (**Figure 4D**). As predicted by our AlphaFold3 analysis (**Figure 4C, *bottom***), under reducing conditions (1 mM TCEP), SRC-SH2^{C188A} showed a 5-fold increase in affinity for the SRC C-terminal phosphopeptide (spanning pY530) compared to SH2^{WT} (**Figure 4D**). Strikingly, oxidation of SH2^{WT} with 0.1 mM PV abolished phosphopeptide binding, consistent with uncoupling of the inhibitory complex. In marked contrast, there was still detectable binding to SH2^{C188A}, albeit with ~3.5-fold lower affinity than under reducing conditions. Next, we examined interactions with an optimised SRC binding phosphopeptide (EPQ**pY**EEIP)[43]. We detected a ~10-fold increase in affinity with SH2^{C188A} compared to SH2^{WT}, consistent with a previous study[44] (**Figure 4D, *lower***). Strikingly, PV treatment eliminated phosphopeptide binding by SH2^{WT}, but not SH2^{C188A}, which showed a ~3-fold decrease in affinity compared to TCEP-treated samples (**Figure 4D**). Both Cys 241 and Cys 248 are present within the SH2 domain and may contribute to this residual PV sensing. Nevertheless, our results support Cys 188 in mediating SRC-SH2 phosphopeptide binding and sensitivity to PV. These findings provide a plausible mechanistic explanation for the impact of Cys 188 oxidation on the activity of SRC in cells.

Finally, we asked whether other human SH2 domains possess putative redox sensitive cysteines that could alter phosphopeptide interactions. Interestingly, SH2D1A, SH2D1B and SH2D5, all possess a cysteine analogous to SRC Cys188 (**Figure S5D**). Furthermore, a mutation affecting this cysteine in the SH2D1A domain is linked to lymphoproliferative disease[45]. Several signaling proteins also possess SH2 domains with cysteine residues within their pTyr binding pockets[46] (**Figure S5E**), including the PTKs Zap70 and Syk, which exhibit reduced phosphopeptide binding following oxidation by H_2O_2[47]. CLNK and JAK1 also have cysteine residues[46] in their pTyr binding pocket that could undergo oxidative modification to affect signaling (**Figures 4E and S5E**). Thus, redox-regulated phosphopeptide interactions with SH2 domains may be present across numerous proteins, beyond SRC.

Figure 5

406
Figure 5. SRC cysteines are required to overcome contact inhibition.
(**A**) SYF KO MEFs were retrovirally transduced with specified SRC expression constructs and
treated with 100 µM PV as shown, before lysis and immunoblotting with indicated antibodies.
(**B**) Cell numbers were determined over time in culture for SYF KO MEFs, retrovirally
transduced with specified SRC expression constructs (as in A). (**C**) Confocal microscopy
images of SYF KO MEFs complemented with the indicated constructs grown to post-
confluence, fixed, permeabilized and stained with the DAPI (nuclei; blue) and Phalloidin (actin;
red). (**D**) Quantification of nuclei taken from 5 or 6 fields of view per cell line. **** p.Adj <
0.001. Ordinary one-way ANOVA with Tukey's correction for multiple comparisons. (**E**) SYF
KO MEFs expressing the indicated constructs were grown to post-confluence, lysed and
immunoblotted with the indicated antibodies. Representative blot from n=3 experiments.
418
419 See also Figure S6.

Mutational relief of SRC autoinhibition bypasses the functions of Cys 188 and Cys 280 in cells.

Mutating SRC Tyr 530 (Y530F) favours its open, active conformation[48]. We hypothesized that this mutation would render SRC less sensitive to the effects of cysteine mutation and activation by oxidation. To test this, we introduced SRC^{Y530F} and dual mutant variants SRC$^{Y530F;C188A}$ and SRC$^{Y530F;C280A}$ into SYF KO MEFs. As expected, basal pY419 and global pTyr signal in cells expressing SRC^{Y530F} was increased compared to those expressing SRC^{WT} (**Figures 5A and S6A**). SRC^{C188A} and SRC^{C280A} showed a striking increase in pY530 and reciprocally diminished pY419, indicating increased autoinhibition. In contrast, introducing the Y530F mutation overcame the inhibitory effects of C188A and C280A in unstimulated cells, as judged by pY419 levels (**Figures 5A and S6A**). PV stimulated global pTyr equivalently elevated for all SRC^{Y530F} variants independently of Cys 188 or Cys 280 mutations. In contrast, PV-induced pTyr levels were lower for SRC^{WT} cysteine mutations. We also noted that SRC pY530, a substrate of numerous PTPs[49-51], was unchanged by PV treatment (**Figure 5A**). These results support that Cys 188 and Cys 280 regulate SRC activity at the level of autoinhibition in cells. Aligning with this, SRC Cys 188 and Cys 280 have been implicated in transducing signals downstream of ATP[21] and IL33[52] stimulations.

SRC Cys 188 and Cys 280 are required for overcoming cell contact inhibition

Reintroducing SRC^{WT} into SYF KO MEFs stimulated proliferation (**Figure 5B**). This effect was not recapitulated by expression of SRC^{C188A} or SRC^{C280A}, reflecting the reduced basal cellular activity of these mutants (**Figure 5B**). In contrast, constitutively active SRC^{Y530F} increased cell growth regardless of C188A and C280A mutations. Notably, SRC$^{Y530F; C188A}$ cells exhibited the highest growth rate (**Figure 5B**), potentially reflecting a negative impact of Cys 188 oxidation, which would impair pY530 binding but would subsequently also limit binding to and recruitment of other phosphoproteins (**Figure 4**). We observed that differences in cell number were largely driven by post-confluent growth. Indeed, SRC^{WT} and SRC^{Y530F} expressing cells did not undergo contact inhibition, whereas SYF KO, SRC^{C188A} and SRC^{C280A} cell lines did (**Figure 5B and videos S3-S6**). Following growth to post-confluence, SRC^{WT} expressing cultures had significantly more nuclei supporting increased cell proliferation compared to either the SYF KO or SRC^{C188A} and SRC^{C280A} (**Figures 5C-5D**). Together, these findings demonstrate that SRC Cys 188 and Cys 280 are required to drive cell overgrowth.

The Hippo pathway is a conserved regulator of contact inhibition of cell growth[53]. We next tested whether SRC-dependent cellular overgrowth phenotypes were associated with YAP activation[53]. Compared to WT, SYF KO MEFs displayed increased YAP phosphorylation at Ser 127 (**Figure 5E**), a site associated with its cytoplasmic sequestration and inactivation. This trend was recapitulated by phosphorylation of the upstream Hippo pathway kinase LATS1 (**Figures 5E and S6B**). Consistently, expression of SRC^{WT} was sufficient to suppress YAP phosphorylation, whereas SRC^{C188A} and SRC^{C280A} resembled SYF KO cells, with elevated levels of YAP phosphorylation, despite comparable SRC expression levels (**Figure 5E**). In conclusion, SRC redox-sensitive cysteines are required for YAP activation, correlating with the ability of cells to overcome contact inhibition of proliferation.

Discussion

PV is a commonly used PTP inhibitor that has shaped our understanding of pTyr-based regulation and the study of PTP-substrate interactions[54,55]. Our work demonstrates that PV can disrupt cellular redox homeostasis, leading to oxidation of HyPer7 and PRDX1, but also shows selectivity, targeting cysteines within positively charged protein surfaces, including SRC Cys 188 and Cys 280. Using a series of mutational approaches, enzyme assays and cellular studies, we find that the potent effects of PV on global pTyr are best explained by the combined inhibition of PTPs and the direct oxidation and activation of SFKs, and perhaps other PTKs, such as Eph receptors[56]. This finding should prompt a re-evaluation of the use and interpretation of experimental data generated using PV treatments.

Based on the oxidation of both HyPer7 and PRDX1, PV must interfere with antioxidant pathways, particularly thioredoxins that can reduce both proteins. One possibility is that it reacts with and depletes reduced glutathione (GSH) or NADPH. Another possibility is that PV evades H_2O_2-reactive PRDX, enabling it to target additional cysteine thiols similarly to what has been observed for *cis*Trp-OOH[57]. This would be consistent with the rapid PTP hyperoxidation. Over time, PV induces PRDX1 tyrosine phosphorylation, potentially enabling endogenous H_2O_2 to oxidise cysteine thiols[32]. *cis*Trp-OOH shows stereospecific oxidation of thiols. Interestingly, previous observations of sulfenic acid modification of SRC cysteine residues induced by H_2O_2 showed a distinct pattern of sensitivity than PV[21]. Therefore, we propose that PV preferentially targets PTP active sites, as well as other phosphate-binding pockets, including the SRC kinase active site (Cys 280) and SH2 domain (Cys 188), correlating with protein regions with high positive charge (**Figure 2G**). In contrast, mercuric chloride activates SRC independently of autoinhibition via Cys 486 and 501, which are surrounded by negative charge, particularly in the open conformation[58]. Therefore, SRC responds differentially to distinct oxidants.

Cys 188 is located at an equivalent position to Thr 42 in the N-terminal SHP2 SH2 domain, which is mutated in disease and alters phosphopeptide specificity[59]. Likewise, SRC-SH2[C188A] altered affinity for phosphopeptides, but also impacted redox sensitivity. Interestingly, Cys 188 reportedly forms a disulfide bridge with a cysteine residue of its substrate cortactin[60], which could explain the greater influence of mutating this cysteine in cells than *in vitro*. We therefore speculate that Cys 188 oxidation plays two roles: firstly, it relieves autoinhibition and secondly, it can mediate disulfide bridge formation with new interaction partners, an intriguing area for further investigation. Amongst SFK cysteine residues, Cys 188 exhibits the highest level of reversible oxidation in mouse tissues, supporting the physiological relevance of our findings[20,61]. We further identified other SH2 domain pTyr pocket cysteine residues. Zap70 possesses a redox-sensitive pTyr binding pocket in its N terminal SH2 domain, but has a distinct mode of autoinhibition compared to SRC[62]. Thus, oxidation of this cysteine may negatively impact ZAP70 recruitment to phosphorylated immunoreceptor tyrosine-based activation motifs (ITAMs)[47], but may also mediate intermolecular disulfide interactions. Interestingly, CSK, which promotes SRC Tyr 530 phosphorylation and subsequent inhibition, is regulated *in vitro* via an intramolecular disulfide bond[63] involving its SH2 domain and could also contribute to redox and pTyr signalling integration. Thus, redox regulation of SH2 domains is a wider area for future investigation.

SRC[C188A] and SRC[C280A] expressing cells showed a striking increase in pY530, which differs from previously reports[21]. One explanation for this difference is that we 'rescued' SRC-

deficient cells, whereas other studies were performed in the presence of endogenous SRC[21].
Furthermore, we have employed untagged SRC wherever possible, since both the N and C
termini of SRC contain important regulatory elements. For example, a C terminal affinity tag
could interfere with autoinhibition, whilst N-terminal tags would block N-myristoylation, which
is important for localisation[64]. This striking increase in pY530 (**Figure 5A**) may be consistent
with the preservation of an autoinhibited state and protection from dephosphorylation by a
stabilized SH2 domain interaction. Alternatively, it could represent an increase in CSK activity,
although this has not been mechanistically explored in this study.
520
In *Drosophila*, Src42A Cys 471, equivalent to human SRC Cys 490, is implicated in wound
healing responses[65], and may form an intramolecular disulphide with Cys 248[19]. However,
neither Cys 188 nor Cys 280 are conserved in *Drosophila* Src (42A or 64B), but they are found
in zebrafish and chicken, indicating multiple layers of SRC redox regulation acquired during
evolution. SRC can be S-glutathionylated at Cys 400 leading to inhibition of
autophosphorylation, linking its cellular regulation to this pathway[66]. How SRC cysteines
undergo oxidation and reduction *in vivo* remains an important question, particularly given the
relatively high oxidant concentrations required to elicit changes in activity *in vitro*.
529
SRC is a proto-oncogene, yet rarely mutated in human cancers[67]. Cys 188 and Cys 280
mutations prevent SRC mediated cell overgrowth, strongly suggesting that the oxidative
environment typical of some cancers[68] could drive relief of SRC autoinhibition, phenocopying
Y530F or C-terminal truncations[67]. Our findings with the conserved Cys 280 residue also have
implications for YES1, for which we also demonstrate oxidative activation by PV (**Figure S3G**).
Interestingly, the equivalent cysteine in YES1 (Cys 287) has previously been targeted with a
covalent inhibitor[69]. Our finding that non-catalytic redox-sensing Cys residues in PTKs is a
dominant signaling mechanism potentially increases their attractiveness as therapeutic
targets, especially if greater potential for selectivity than active site inhibitors can be
achieved[70].
540
541 **Resource availability:**

542 Mass spectrometry data have been deposited to the ProteomeXchange Consortium through
543 the PRIDE partner repository with the dataset identifiers PXD059641(Cysteine S-sulfonate
544 modifications) and PXD059840 (HDX-MS). All other materials are available upon request to
545 the corresponding authors.
546
Acknowledgements:
We thank the Babraham Institute imaging facility, J. Clark in the Babraham Institute biological
chemistry facility, C. Switzer for discussions about redox chemistry, T. Chessa for advice on
retroviral transduction, K. Wojdyla for advice on differential alkylation and phosphopeptide
enrichment, and E. Veal, P. Hawkins, A. Tuncay, R. Williams and members of the Sharpe lab
for critical feedback on the manuscript. This work was supported by a Sir Henry Dale
Fellowship jointly funded by the Wellcome Trust and the Royal Society [109407/A/15/A]
awarded to HJS, an ESPRC Frontier guarantee grant [EP/Z000114/1], a BBSRC institutional
programme grant [BBS/E/B/000C0433] and core capability grant [BB/CCG2210/1]. OR is
supported by a studentship funded by the Lister Institute of Preventive Medicine. MB in the
Roger Williams group is supported by the Medical Research Council (MC_U105184308 to
RLW) and Cancer Research UK (grant DRCPGM\100014 to RLW). PC and NC were
supported by Cancer Research UK grants C9685/A26398, C9685/A27415, and

C9545/A29580. JM and DE are being supported by a VIB grant. LAD/DPB/CEE and PAE
acknowledge UKRI equipment and project funding through BBSRC grants BB/S018514/1
BB/R000182/1 and BB/X002780/1. HJS is an EMBO Young Investigator and Lister Institute
Prize Fellow.
564
Author contributions:
Conceptualization: KEM, DPB, HJS
Methodology: KEM, MB, NC, OR, DE, LAD, SAC, DPB
Investigation: KEM, NC, OR, DE, DPB
Formal analysis: KEM, MB, NC, OR, DE, LAD, SAC, DPB
Visualization: KEM, MB, NC, OR, DE, TL, SAC, DPB, HJS
Validation: KEM, DPB
Data curation: HEJ, HJS
Funding acquisition: PC, CEE, JM, PAE, HJS
Project Administration: HJS
Supervision: PC, CEE, JM, PAE, HJS
Resources: PC, CEE, JM, PAE, HJS
writing – original draft: KEM, DPB, HJS
writing – review and editing: KEM, MB, NC, CEE, JM, PAE, DPB, HJS
579
580 **Declaration of interests:**
581 The authors declare that they have no competing interests.
582
Supplemental information
Document S1. Figures S1 to S6
Video S1. TIRF microscopy of U2OS cells expressing PM-HyPer7 treated with media. Related
to Figure 1.
Video S2. TIRF microscopy of U2OS cells expressing PM-HyPer7 treated with 100 µM PV.
Related to Figure 1.
Video S3. Brightfield timelapse of SYF KO MEF cells in culture. Related to Figure 5.
Video S4. Brightfield timelapse of SYF KO + SRC MEF cells in culture. Related to Figure 5.
Video S5. MEFs Brightfield timelapse of SYF KO + SRC[C188A] MEF cells in culture. Related to
Figure 5.
Video S6. MEFs Brightfield timelapse of SYF KO + SRC[C280A] MEF cells in culture. Related to
Figure 5.
Table S1. Mass spectrometry data for cysteine oxidation of affinity purified SRC.
596
STAR methods
Plasmids and constructs
Amino acid (aa) numbering is based on the following sequences: Human PTPRK; UniProt ID:
Q15262-3, Human SRC; Uniprot ID: P12931-1, Chicken Src Uniprot ID: P00523. Generation
of bacterially expressed (aa 864–1446) and mammalian overexpression PTPRK intracellular
domain (aa 776-1446) constructs were previously described[73,74]. Full length human SRC
containing a 3C-protease cleavable, N-terminal, tandem Strep-tag II was cloned into pcDNA3.
Human rhinovirus 14 3C protease plasmid (pET-NT*-HRV3CP) was a gift from Gottfried Otting
(Addgene # 162795).
606

The retroviral pLNCX chicken Src Y527F plasmid was a gift from Joan Brugge (Addgene
#13660). HyPer7 [30] (a gift from Vsevolod Belousov (Addgene #136464)) was fused in-frame
downstream of the SRC myristylation sequence to make the 'PM-HyPer7' biosensor and
subcloned into pLJM1 (gift from Joshua Mendell (Addgene #91980)). N-terminally His-tagged
HyPer7 was cloned into the pQE30 vector. The cDNA encoding the human SRC SH2 domain
(Gln147-Lys252) was subcloned into the pETM11 vector (a gift from Elisa Izaurralde (Addgene
#145950)). The QuikChange II mutagenesis kit (Agilent) was used to construct the mutant
SRC SH2 domains.
615
616 All other point mutations were introduced by polymerase chain reaction (PCR) using either Q5
617 High-Fidelity DNA (New England Biolabs, UK), CloneAmp (Takara) or Phusion Hot Start II
618 DNA (Thermo Fisher Scientific) polymerases as per manufacturer's protocol.
619
Cell culture and treatments
HEK293T, U2OS and SYF KO MEFs (CRL-2469) cells were purchased from ATCC. WT MEFs
were generated in-house, described below. SRC WT and KO HAP1 cells were purchased
from Horizon (HZGHC000348c023). Plat-E cells were donated by P. Hawkins and L. Stephens
(Babraham Institute) and were purchased from Cell Biolabs Inc (RV-101). Cells were
maintained in 75 cm^2 vented tissue culture flasks in a 37°C humidified 5% CO_2 ventilator and
passaged using trypsin-EDTA solution (Sigma-Aldrich), prior to reaching confluence every 2-
4 days. HEK293T and MEF cells were grown in DMEM containing 10% (v/v) fetal bovine serum
(FBS; Sigma-Aldrich), 2 mM L-glutamine (Sigma-Aldrich). U2OS cells were grown in McCoy's
5A medium modified with 10 % (v/v) FBS (Sigma-Aldrich). HAP1 cells were cultured in IMDM
containing 10% (v/v) fetal bovine serum (FBS; Sigma-Aldrich, with Plat-E cells cultured in
DMEM (high glucose with L-glutamine and pyruvate supplemented) containing 10 % heat-
inactivated FBS (Sigma-Aldrich) with additional blasticidin (10 mg/ml) and puromycin (1
mg/ml) when in culture for more than 2 passages. All cell lines are routinely mycoplasma
tested using MycoProbe Mycoplasma Detection Kit (R&D system) or MycoAlertTM (Lonza
Bioscience).
636
MEF line generation
WT (Bl6/N) MEFs were generated in-house. Following dissection, E13-14 embryos were
placed into PBS on ice. Placental and maternal tissues and head were removed prior to
tryspinization and dissociation with a razor blade. DNAseI was then added followed by
incubation at 37°C for 10 min. DMEM with 10 % FBS was added to quench trypsin. Cells were
pipetted through a 40 μm filter and pelleted at 140 x g for 5 min. Finally, cells were
resuspended in DMEM with 10% FBS, 2 mM L-glutamine and pen-strep then seeded for
culturing. Cells were cryopreserved in liquid N2 in 50% FBS:10% DMSO:40% DMEM for future
use.
646
Small molecules and growth factors
The following inhibitors were purchased from Sigma: Dasatinib (SML2589-50MG), focal
adhesion kinase inhibitor 1 (324877-5MG), Gefitinib (SML1657-10MG), Ibrutinib (PZ0242-
5MG), Imatinib (CDS022173-25MG), Lapatinib (SML2259-10MG), Nilotinib (SML3799-
10MG), PP2 inhibitor (529573-1MG), Saracatinib (SML3195-5MG) and SU6656 (S9692-5mg).
Inhibitors were re-suspended in DMSO. The following growth factors were purchased from
Qkine: PDGF (Qk044), IGF1 (Qk047), FGF2 (Qk025), TGFβ2 (Qk072). Insulin was purchased

from Scientific Laboratory supplies (I5500) and EGF was purchased from Peprotech (AF-100-
15).
656

Antibodies and dyes
Antibodies against the following target proteins or epitopes were used in the study:
Phosphotyrosine (Cell Signaling Technology (CST) #8954S), α Tubulin (Sigma #T6199), Flag
(CST #8146S), Pan-SFK (CST #2108), SRC family pY416 (#CST D49G4), SRC pY530 (CST
#2105), SRC-specific (CST #2123 & #2120), oxPTP (R&D Systems # MAB2844), His (CST
#12698S), ABL1 (c-Abl) (CST #2862), ABL1 (c-Abl) pY412 (CST #2865S), Cortactin (CST
#3503), Cortactin pY421 (CST #4569), LATS1 (CST #3477), LATS1 pY909 (CST#9157) , YAP
(CST #4912), YAP pS127 (CST #13008), AKT (CST #9272), AKT pS473 (CST #4060), ERK
pT202/pY204 (CST #4370), ERK (CST #4696), EGFR (CST #2232S), EGFR pY1068 (CST
#3777S), SHP2 (CST #3397), PRDX1 pY194 (CST #14041), PRDX1 (CST #8499). HRP-
conjugated secondary antibodies were purchased from Jackson Immunoresearch(#715-035-
150 & #711-035-152). Hoechst and BODIPY Phalloidin 558/568 (B3475) was purchased from
Thermo Fisher Scientific.
670

Pervanadate stock generation
Pervanadate was generated as previously described[6]. To avoid decomposition, it was
prepared immediately prior to use. In a 1.5 ml tube, 100 µl of activated pH 10 100 mM sodium
orthovanadate (Alfa Aesar 15403009) was combined with 103 µl 0.49 M H_2O_2 in 20 mM
HEPES, pH 7.3, mixed by gentle inversion and incubated at RT for 5 min. Excess H_2O_2 was
quenched by addition of 23 µl of 0.5 mg/ml catalase in 50 mM potassium phosphate, followed
by mixing by gentle inversion, yielding a 44.2 mM pervanadate stock solution for immediate
use.
679

Recombinant proteins and synthesized peptides
Peptides were obtained at 95% purity from Biosynth. The SRC peptide substrate 5-FAM-
EGIYGVLFKKK-CONH2 was synthesized by Genscript. Recombinant N-terminal His6-tagged
SRC (1-536) kinase fusion protein (DU19041) and N-terminal His6-tagged YES1 (1-543)
(DU5884) was purified from Sf21 cells was purchased from Medical Research Council Protein
Phosphorylation and Ubiquitylation Unit (PPUU) reagents and services (University of
Dundee). N-terminally His-tagged Hyper7[30], PTPRK ICD[73], SRC SH2 domain and His6-tagged
3C protease[75] were purified using standard affinity and size exclusion chromatography
methods and stored −80°C as follows*:
689

M15 Escherichia coli cells were transformed with N-terminally His-tagged Hyper7 construct
and cultured in LB Broth at 37°C to OD600nm ~0.5-0.8 and induced with 400 µM isopropyl-
thio-β-D-galactopyranoside (IPTG Thermo Fisher Scientific, dioxane-free) for 14 h at 16°C.
Escherichia coli BL21(DE3) Rosetta cells transformed with SRC SH2 construct or the other
aforementioned constructs were cultured at 37°C/220 rpm in LB broth containing 34 µg/ml
chloramphenicol until the OD600 nm reached 0.5–0.8 or cultured at 30°C/220 rpm in 1 L
2XTY medium containing 50 µg/ml carbenicillin and 34 µg/ml chloramphenicol until the
OD600 reached 0.6–0.7 respectively. Cultures for SRC SH2 were transferred to 16°C/220
rpm and induced with 100 µM IPTG (Thermo Fisher Scientific, dioxane-free) for 14 h. All
other constructs were transferred to 20°C/220 rpm and allowed to equilibrate, prior to the
addition of 1 mM IPTG (Generon) for 20h. PTPRK cultures were also supplemented with 200

µM of D-biotin (Sigma-Aldrich) upon IPTG addition. Cells were harvested by centrifugation at
4000 x g for 20-30 mins and bacterial pellets were stored at – 20°C until required.
703
For Hyper7, pellets were resuspended in lysis buffer (40 mM HEPES, pH 7.4, 400 mM NaCl,
10% (v/v) glycerol, 1 mM benzamidine, 1 mg/mL Leupeptin, 0.1 mg/mL 4-(2-aminoethyl)
benzenesulfonyl fluoride hydrochloride (AESBF), 100 mg/mL DNaseI, 20 mM $MgCl_2$) or
oxidizing or reducing lysis buffer for SRC SH2 domains (20 mM HEPES (pH 7.5), 100 mM
NaCl, and 0.1 mM pervanadate; reducing SH2 buffer: 20 mM HEPES (pH 7.5), 100 mM
NaCl, and 1 mM TCEP). Once re-suspended in lysis buffer, samples were lysed by
sonication and clarified by centrifugation at 39,000 x g for 30 min at 4°C. For PTPRK and 3C
protease proteins, cells were lysed in purification buffer (50 mM HEPES pH 7.5, 500 mM
NaCl, 5% (v/v) glycerol and 0.5 mM tris(2-carboxyethyl)phosphine (TCEP)), containing
EDTA-free protease inhibitor tablets (Roche) followed by cell disruption using a Constant
Systems cell disruptor and then lysate clarification via centrifugation at 40000 x g for 30 min
at 4°C. His-HyPer7 were purified by affinity chromatography (Ni^{2+}- Sepharose (GE
Healthcare) and imidazole-based elution) and size exclusion chromatography using a
HiLoad 16/600 Superdex 200 column (GE Healthcare) equilibrated in 50 mM Tris-HCl (pH
7.4), 100 mM NaCl, and 10% (v/v) glycerol. SH2 domains were purified with pre-equilibrated
nickel column (His-Trap HP, GE Healthcare) and the 6x His-tag was removed by digestion
with home-made tobacco etch virus protease at 4°C for 12 h accompanied with oxidizing
buffer and reducing buffer.
722
SH2 domains were further purified on a Superdex 75 10/300 GL size exclusion column (GE
Healthcare) with oxidizing buffer and reducing buffer. For PTPRK and 3C protease, the
supernatant was removed and incubated with 0.5 ml of Ni-NTA agarose (Qiagen) for 1 h at
4°C. Ni-NTA Agarose was then pelleted via centrifugation at 500 x g for 5 min at 4°C and
packed into a gravity flow column. Ni-NTA agarose was then washed with 10 volumes of
purification buffer containing 5 mM imidazole, followed by 20 volumes of purification buffer
containing 20 mM imidazole; prior to elution in purification buffer containing 250 mM
imidazole. The eluted protein was then subjected to size exclusion chromatography (SEC)
using a Superdex 200 16/600 column (GE Healthcare Life Sciences, Thermo Fisher
Scientific). Columns were equilibrated in SEC buffer (50 mM HEPES pH 7.5 (50 mM Tris pH
7.4 for SH2 domains), 150 mM NaCl, 5% (v/v) glycerol, 5 mM DTT). Protein was
concentrated to 2–10 mg/ml using an Ultracel-10K regenerated cellulose centrifugal filter
(Merck Millipore). Proteins were either used fresh or snap-frozen for storage at -80°C until
required.
737

Generation of stable cell lines

SRC-rescued SYF KO MEF cell lines were generated using the retroviral expression system.
Retrovirus was generated through the reverse transfection of Plat-E cells (12 x 10^6) in 15 cm^2
dishes with 16.8 µg pLNCX construct DNA into 2 ml OPTIMEM (Gibco) with 50.4 µl
Lipofectamine 2000 transfection reagent (Invitrogen) following the manufacturer's instructions.
Media was replaced with 9 ml complete MEF media per 15 cm^2 dish 20-24 hours post-
transfection. 48-72 h post-transfection, retrovirus was harvested from Plat-E cell media by
centrifugation for 5 min at 800 x g (RT) and filtered through a 0.2 µm syringe filter (Sartorius)
before either freezing at -80°C or adding directly onto MEF SYF KO cells after dilution 1:1 into
MEF culture media, in 6-well dishes. Media was replaced 24 h after adding viral supernatant,
followed by at least two passages before experimental analysis.

For lentiviral generation, 1.5×10^7 HEK293T cells were seeded in 12 ml of complete growth
medium per 15 cm^2 dish (two dishes per lentivirus) and incubated for 24 h at 37°C with 5%
CO$_2$. Each 15 cm^2 dish was then transfected with 6 µg of pLJM.PM-HyPer7, 12 µg psPAX2
packing plasmid (Addgene #12260, deposited by Didier Trono) and 3 µg pMD2.G envelope
plasmid (Addgene #12259, deposited by Didier Trono) using the GeneJuice transfection
reagent (Merck Millipore, UK) as per manufacturer's instructions. After 24 h, the medium was
then replaced with 16 ml complete growth medium. At 48–72 h post-transfection, culture
medium was collected and filtered through a 0.2 µm syringe filter (Sartorius). Viral particles
were concentrated using Lenti-X concentrator as per manufacturer's instructions (Takara Bio).
Lentivirus was centrifuged at 1500 x g for 45 min at 4°C, resuspended with complete DMEM
media and aliquoted and stored at −80°C until required. 2×10^5 U2OS cells were seeded per
well of a six-well plate in 900 µl of growth medium, prior to the drop-wise addition of 100 µl
concentrated lentivirus. After 30 min at room temperature, the cells were returned to the
incubator. 72 h later, cells were reseeded in 1 µg/ml puromycin (Thermo Fisher Scientific)
selection medium. Cells were subsequently maintained in media supplemented with 1 µg/ml
puromycin.
766

Lipid-based transfection of siRNA pools
MEF SYF KO cells were reverse transfected with siRNAs (ON-TARGETplus SMARTpool,
Dharmacon Horizon Discovery) using lipofectamine RNAiMAX as per the manufacturer's
instructions (Thermo Fisher Scientific 13778030). Briefly, for each well of a 6-well plate, 6 µl
RNAiMAX was used to transfect 10 nM siRNA. After 24 hr, media was replaced with complete
growth medium and left to recover for another 24 hr prior to cell treatments and processing for
analysis.
774

pNPP phosphatase activity assay
Recombinant PTP domains were made up to 500 µl in assay buffer (50 mM Tris–HCl, pH 7.4,
150 mM NaCl, 5% [v/v] glycerol, 5 mM DTT) at 1 µM. PTPRK intracellular domain and 20 mM
pNPP substrate (NEB P0757S) (in assay buffer) were equilibrated to 30°C for 15 min in an
orbital shaking heat block at 500 RPM. To initiate reactions, 500 µl of 20 mM pNPP substrate
was added to PTP containing tubes (0.5 µM PTP and 10 mM pNPP final concentrations).
Reactions were carried out for 120 min at 30°C in an orbital shaking heat block at 500 RPM.
At each time point (0, 2, 5, 10, 15, 30, 60, 90 and 120 min), 100 µl of the total reaction was
transferred to a 96-well microplate well containing 50 µl 0.58 M NaOH, terminating the
reaction. After the final time point, absorbance of each sample was measured at 405 nm in a
Spectramax M5 plate reader (Molecular Devices). Product formation was calculated by
interpolation of absorbance values using a 4-nitrophenol standard curve of known
concentration.
788

SDS-PAGE and immunoblotting
Protein or cell lysate samples were resuspended in an appropriate volume of 5X SDS-PAGE
sample buffer (0.25 M Tris-HCl pH 6.8, 10% (w/v) SDS, 20% (v/v) glycerol, 0.1% (w/v)
bromophenol blue, 10% (v/v) β-mercaptoethanol) and incubated at 92°C for 10 min. Samples
were run on a 10% (v/v) SDS-polyacrylamide resolving gel with a 5% (v/v) SDS-PAGE
stacking gel and subjected to electrophoresis at 125 V for 1.5 h in 25 mM Tris, 190 mM glycine,
0.1% (w/v) SDS. Proteins were transferred onto 0.2 µm reinforced nitrocellulose membranes
(GE Healthcare) at 300 mA for 3 h at 4°C in 25 mM Tris, 190 mM glycine, 20% (v/v) methanol.

Membranes were briefly rinsed in TBS-T (20 mM Tris pH 7.6, 137 mM NaCl, 0.1% (v/v) Tween-
20) prior to incubation for 30 min in 5% (w/v) skimmed milk/TBS-T to block non-specific
antibody binding. The blocking solution was removed and membranes rinsed in TBS-T prior
to primary antibody incubation (overnight at 4°C). Membranes were then subjected to 3 × 10
min washes in TBS-T, prior to incubation with HRP-conjugated species-specific anti-IgG
antibodies (1 h at RT). Membranes were then subjected to 3 × 10 min washes in TBS-T, prior
to being incubated with combined EZ-ECL solution (Cytiva) and imaged using a G:BOX Chemi
XX6 (Syngene).
805
Peroxiredoxin disulphide shift
HEK293T cells were subjected to a pervanadate time course followed by incubation with
PBS with or without 10 µg/ml catalase (SLS #C3155) and 100 mM NEM for 15 mins at room
temperature. Catalase was added to the PBS 30 min prior to use with NEM added
immediately prior to use. Samples were lysed in 2 % SDS lysis buffer with and without 10
mM NEM. Samples underwent immunoblotting on 12 % gels with either non-reducing or
reducing conditions and probed with PRDX1.

813
PTPRK Flag overexpression, immunoprecipitation and pNPP assay
Flag IP-phosphatase assays were performed as previously described[74]. HEK293T (6 x10^6)
cells were plated in 10 cm^2 dishes, 24 h prior to transfection. 8 µg expression constructs for
3x FLAG tagged ICDs were transfected using 24 µl GeneJuice (Sigma) in 1 ml OPTIMEM
(Gibco) serum free media. Media was replaced 4 h post-transfection. For immunoprecipitation
(IP), cells were transferred to ice 48 h post transfection and washed twice with ice-cold 1X
phosphate buffered saline (PBS). Cells were lysed in ice-cold lysis buffer (50 mM Tris-HCl pH
7.4, 150 mM NaCl, 10 % [v/v] glycerol, 1 % [v/v] Triton X-100, 1 mM EDTA, 10 mM NaF, 1
mM PMSF, 1X-EDTA-free protease inhibitor) on ice for 30 mins with periodic agitation. Lysates
were collected and cleared by centrifugation at 14000 x g for 15 min at 4 °C and supernatants
were transferred to chilled tubes. Total protein concentration was quantified by bicinchoninic
acid (BCA) assay. Lysates were adjusted to a final concentration of 5 mM DTT to prevent air
oxidation. Equal amounts of total cell lysate for each sample (~2 mg) were combined with 50
µl (100 µl of 50 % slurry) of washed FLAG-M2 magnetic beads (Sigma Aldrich) in a total
volume of 1 ml made up in lysis buffer and incubated for 30 mins at 4 °C with rotation. Beads
were then collected on a magnetic stand and supernatants removed. Beads were then
resuspended and washed once with 1 ml lysis buffer and then four times with lysis buffer
containing 500 mM NaCl. For pNPP assays, IPs were then washed in 1 ml assay buffer (50
mM Tris-HCl pH 7.4, 150 mM NaCl, 5% [v/v] glycerol, 5 mM DTT] and then resuspended in
500 µl assay buffer. Samples were then subject to pNPP assays as described above. After
the assay beads were collected by magnet, washed in 1 ml TBS and resuspended in 1X SDS
PAGE sample buffer and incubated at 95°C for 5 mins. Beads were collected by magnet and
supernatants used for SDS-PAGE and immunoblot analysis.
837
HyPer7 imaging and analysis
U2OS cells expressing PM-HyPer7 were seeded on 35 mm glass bottom IBIDI dishes (81158)
prior to imaging and maintained at 37°C and 5% CO$_2$. Fluorescence excitation was measured
at 488 nm and 405 nm, and λem = 525 nm as described previously[30]. Treatments were
prepared at 3x and added gently to the side of the culture dish after 1 min imaging. Images
were acquired using Total internal reflection fluorescence (TIRF) microscopy (Nikon Eclipse
STORM-TIRF microscope), using a 40x oil immersion objective. All image analysis was

conducted using ImageJ/FIJI. HyPer7 ratiometric images (λex=488/ λex=405) were false
coloured using mpl-Magma look-up table (purple to yellow). GraphPad Prism (Ver. 10) was
used for statistical analysis.
848
HyPer7 and pervanadate reaction *in vitro*
Purified recombinant HyPer7 was buffer-exchanged into Ar-flushed 100 mM NaPOi, pH 7.4,
150 mM NaCl, 0.1 mM diethylene triamine penta-acetic acid (DTPA) using a 5 ml HiTrap
desalting column (GE Healthcare). The protein concentration was determined with the
Bradford reagent (Bio-Rad), using bovine serum albumin as a standard. The protein was
diluted to 2.8 µM and incubated with 0.1 mM DTT for 30 min at RT. Pervanadate, H_2O_2 and
catalase controls (subsequently referred to as "reactants") were prepared at a 10x
concentration. HyPer7 (final concentration 2.5 µM) and the reactants were mixed in a quartz
cuvette, incubated for 1 min at RT and the fluorescence excitation spectrum was recorded
from λex = 350 nm to 505 nm with λem = 512 nm using a Cary Eclipse Fluorescence
Spectrophotometer (Agilent). The ratio of the two excitation maxima at 400 nm and 498.5 nm
was the final readout. GraphPad Prism (Ver. 10) was used for statistical analysis.
861
Protein kinase assays
Kinase assays were performed using nonradioactive real-time mobility shift-based microfluidic
assays, as described previously[75], in the presence of 2 mM of the appropriate fluorescent-
tagged peptide substrate (5-FAM-EGIYGVLFKKK-$CONH_2$) and 1 mM ATP (unless specified
otherwise). Pressure and voltage settings were adjusted manually to afford optimal separation
of phosphorylated and non-phosphorylated peptides. All assays were performed in 50 mM
HEPES (pH 7.4), 0.015% (v/v) Brij-35, and 5 mM $MgCl_2$, using 3 ng of SRC, and real-time or
end point peptide phosphorylation was determined using the ratio of the
phosphopeptide:peptide. Kinase activity in the presence of different redox reagents or inhibitor
compounds was quantified by monitoring the generation of phosphopeptide during the assay,
relative to controls, with phosphate incorporation into the peptide generally limited to <20% to
prevent ATP depletion and maintain assay linearity. The effects of variable exposure times on
kinase activity were monitored by preincubating SRC and redox reagent for 10, 60 or 120 mins
at room temperature (RT), and then initiating substrate phosphorylation with the addition of
ATP and peptide substrate. Relative changes of SRC activity in the presence of increasing
concentrations of pervanadate were monitored in real time over 4 h using 20 µM peptide
substrate. The rate of substrate phosphorylation at each time point was calculated as pmol
phosphate incorporation per min and plotted as a function of total assay time. Counter screen
assays using 50 ng Aurora A kinase were performed as for SRC.
881
Differential alkylation in cell lysates
Differential alkylation methodology was performed as previously described[76]. HEK293T cells
were plated at 2 x10^6 per 10 cm^2 dish until full confluency followed by PV treatments. Cells
were washed twice in ice-cold PBS followed by harvesting into pH 7.0 lysis buffer [30 mM Tris-
HCL (pH 7), 120 mM NaCl, 2 mM EDTA, 2 mM KCl, 10 % (v/v) glycerol, 1 % (v/v) triton, 1 %
SDS (v/v), 5 mM NaF, 1X EDTA-free protease inhibitor tablet cocktail (Roche; 1836170001)
and PhosSTOP tablets (Roche; 4906845001)] containing NEM (Sigma; E3876) 5 mM for
alkylation. Lysates were then sonicated and incubated on ice for 20 min in the dark to enable
NEM alkylation, followed by centrifugation at 13000 x g to pre-clear the lysate. During this
incubation, Zeba™ spin desalting columns (Thermo Fisher Scientific; 89882) were
equilibrated with lysis buffer pH 8.0 following the manufacturer's instructions. 0.5 mg protein

per sample was buffer exchanged by filtering through three Zeba™ spin columns per sample to remove excess NEM. 50 mM TCEP (Merck; 646547) was added to each sample followed by incubation at room temp for 30 mins to enable sample reduction. Samples were then filtered through three pH 8.0 lysis buffer equilibrated Zeba™ columns per sample to remove excess TCEP. Finally, samples were incubated with 10 mM EZ-Link™ Iodoacetyl-PEG2-Biotin (Thermo Fisher Scientific; 21334) for 90 mins at RT in the dark, followed by buffer exchange through three equilibrated Zeba™ spin columns to remove excess alkylating agent. Lysates were then subjected to streptavidin enrichment using the following method. Protein concentration was measured again for each sample using BCA assay and 400 µg each sample was subject to pull-down. 10 % lysate was removed and added to sample buffer for input sample. Remaining lysate was incubated with 50 µl neutravidin beads (Pierce; 29202) rotating for 2 h at 4°C. Equivalent of 10 % starting lysate concentration was removed for supernatant sample and added to sample buffer. The remaining beads were washed 4x with RIPA buffer [50 mM Tris-HCl, pH 7.4, 1% (v/v) Triton, 1% Na deoxycholate, 0.1% SDS, 150 mM NaCl, 1 mM EDTA] followed by elution in 5x sample buffer with 25 mM biotin (Merck; B4639) with incubation at 95°C for 10 mins. Input, supernatant and pull-down samples then underwent analysis by immunoblotting.

SRC affinity precipitation from HEK293T cells

Affinity precipitation experiments used HEK293T cells transfected using a 3:1 polyethyleneimine [average Mw (weight-average molecular weight), ~25,000 Da; Sigma-Aldrich] to DNA ratio (30:10 mg, for a single 10 cm^2 culture dish). Proteins were harvested 48 h post transfection using a lysis buffer containing 50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 0.1% (v/v) Triton X-100, and 5% (v/v) glycerol and supplemented with a protease inhibitor cocktail tablet and a phosphatase inhibitor tablet (Roche). Lysates were briefly sonicated on ice and clarified by centrifuged at 20,817g for 20 min at 4°C, and the resulting supernatants were incubated with Strep-TactinXT 4Flow affinity resin (IBA Lifesciences) for 1 h with gentle end over end mixing at 4°C. Beads containing bound protein were collected and washed three times in 50 mM tris-HCl (pH 7.4) and 500 mM NaCl and then equilibrated in storage buffer [50 mM Tris-HCl (pH 7.4), 100 mM NaCl, and 5% (v/v) glycerol]. The purified kinases were then proteolytically eluted from the beads over a 1 h period using 3C protease (0.5 mg) at 4°C with gentle agitation and then assayed using real-time microfluidic peptide assay as described above. For MS-dependent detection of cysteine modifications, HEK293T cells were treated with pervanadate or H$_2$O$_2$ for 30 mins, and protein harvested in 50 mM Tris-HCl (pH 6.5), 150 mM NaCl, 0.1% (v/v) Triton X-100, and 5% (v/v) glycerol supplemented with a protease and phosphatase inhibitor tablets (Roche), and 100 mM iodoacetamide to alkylate thiols. After brief sonication, lysates were incubated in the dark for 1 h at 4 °C, clarified by centrifugation and affinity purified as above.

Identification of affinity enriched SRC cysteine oxidation state using mass spectrometry

Affinity purified SRC preparations were diluted to 180 µl in 100 mM ammonium bicarbonate pH 8.0 and reduced and alkylated with dithiothreitol and iodoacetamide, as previously described by[77]. Samples were subject to SP3-bead digestion as described by [78] with the adaptions of: using 100 mM ammonium bicarbonate pH 8.0, 0.5 µg Trypsin gold (Promega) and post-digest bead washing in 1% (w/v) Rapigest (Waters). Once supernatants were combined, samples were acidified by addition of a final concentration of 0.5% TFA and incubated for 30 min each at 37 °C with 600 rpm shaking and on ice. Samples were centrifuged

at 13000 x g for 10 min at 4 °C and cleared supernatant collected into fresh tubes prior to
vacuum centrifugation. Dried peptides were solubilized in 20 μl of 3% (v/v) acetonitrile and
0.1% (v/v) TFA in water, sonicated for 10 minutes, and centrifuged at 13000 x g for 15 min at
4 °C prior to reversed-phase HPLC separation using an Ultimate3000 nano system (Dionex)
over a 60 min gradient, as described by[79]. All data acquisition was performed using a Thermo
QExactive HF mass spectrometer (Thermo Fisher Scientific), with higher-energy C-trap
dissociation (HCD) fragmentation set at 29% normalized collision energy for 2+ to 5+ charge
states. MS1 spectra were acquired in the Orbitrap (60K resolution at 200 m/z) over a m/z
range of 350 to 2000, AGC target = 3e6, maximum injection time = 100 ms, with an intensity
threshold for fragmentation of 1e3. MS2 spectra were acquired in the Orbitrap (30K resolution
at 200 m/z), maximum injection time = 45 ms, AGC target = 1e5 with a 20 s dynamic exclusion
window applied with a 10-ppm tolerance. Data was analyzed using Proteome Discoverer 2.4
(Thermo Fisher Scientific) in conjunction with MASCOT[80]; searching the UniProt Human
Reviewed database (updated weekly, accessed March 2022) with constant modifications =
carbamidomethyl (C), variable modifications = oxidation (M), phosphorylation (ST), trioxidation
(C), instrument type = electrospray ionization–Fourier-transform ion cyclotron resonance (ESI-
FTICR), MS1 mass tolerance = 10 ppm, MS2 mass tolerance = 0.01 Da with the ptmRS node
active. Label free quantification was performed using the Minora feature detector node,
calculating the area under the curve for m/z values, total protein abundance was determined
using the HI3 method[81]. For cysteine trioxidation-containing peptides identified in at least 3
replicates, peptide abundance was normalized against the calculated total protein abundance
to account for potential protein load variability during analysis prior to normalization against
untreated control. For the best MASCOT scoring PSM of a C188-trioxidation containing
peptide, the spectrum was extracted in *.mgf format from the raw file and imported into a
custom R script for re-drawing and manual annotation.
966

Differential scanning fluorimetry
Thermal shift assays were performed using a StepOnePlus real-time polymerase chain
reaction (PCR) machine (Life Technologies) using SYPRO Orange dye (Invitrogen) and
thermal ramping (0.3°C in step intervals between 25°C and 94°C). Recombinant SRC proteins
were assayed at a final concentration of 2.5 mM in 50 mM tris-HCl (pH 7.4) and 100 mM NaCl
in the presence or absence of the indicated concentrations of ATP, H_2O_2, DTT, orthovanadate,
pervanadate or Dasatinib [final dimethyl sulfoxide (DMSO) concentration no higher than 4%
(v/v)] as described previously[75]. Normalized data were processed using the Boltzmann
equation to generate sigmoidal denaturation curves, and average Tm/ΔTm values were
calculated as previously described[82] using GraphPad Prism software.
977

Hydrogen-Deuterium exchange mass spectrometry
A preincubation was performed by adding to a concentrated SRC sample (57 μM SRC in 50
mM Tris pH 7.5, 150 mM NaCl) a 4.5 mM PV solution to yield a final solution of 5 μM SRC (20
mM Hepes pH 7, 150 mM NaCl) in presence of 0.5 mM or 2.5 mM of PV. Samples were
incubated for 15 minutes at room temperature and then exposed for different deuteration times
(3 seconds on ice, 3, 30, 300 and 3000 seconds at room temperature) to a deuterated buffer:
50 mM Tris pH 7.5, 150 mM NaCl, with D_2O at a final concentration of 90% (Thermo Fisher
Scientific, 166300100) to yield a final solution that was 0.5 μM of SRC in absence/presence
of 0.05 mM or 0.25 mM of PV. Triplicate samples were aliquoted for each deuteration
timepoint. Exchange reactions were quenched in 2 M Guanidine-Hydrochloride, 100 mM
Glycine, formic acid (Fisher Chemical, 10596814), LC-MS grade water (Romil, H949), pH 2.3

(final pH was 2.5). Samples were then flash-frozen in liquid nitrogen immediately and stored
at -80 °C until analysis. Prior to LC-MS analysis, samples were quickly thawed and injected
on a HDX sample manager coupled to an Acquity ultra performance liquid chromatography
(UPLC) M-Class system (Waters) set at 0.1 °C. Samples were then digested at 15 °C using a
20 mm x 2.0 mm home-made pepsin column (Pepsin from Thermo Scientific, 20343 and
hardware from Upchurch Scientific, C130-B) and loaded on a UPLC pre-column (Acquity
UPLC BEH C18 VanGuard pre-column, 2.1 mm I.D. × 5 mm, 1.7 µm particle diameter, Waters,
186003975) for 2 min at 100 µL/min. Digested peptides were then eluted from the pre-column
onto an Acquity UPLC BEH C18 column (1.0 mm I.D. × 100 mm, 1.7 µm particle diameter,
Waters, 186002346) using a 5-43% gradient of Acetonitrile (Romil, H050), 0.1% formic acid
over 12 min. MS data acquisition was done on a Synapt G2Si-HDMS (Waters) with positive
electrospray ionization, using HDMS^E acquisition mode over a *m*/*z* range of 50–2000 and a
100 fmol.µL^{-1} Glu-FibrinoPeptide solution was used for lock-mass correction and calibration.
The following parameters were used during the acquisition: capillary voltage, 3 kV; sampling
cone voltage, 40 V; source temperature, 90°C; desolvation gas, 150°C and 650 L.h^{-1}; scan
time, 0.3 s; trap collision energy ramp, 20–45 eV. Peptide identification was performed using
ProteinLynx Global Server 3.0.1 (Waters) with a home-made protein sequence library
containing SRC and pepsin sequences, with peptide and fragment tolerances set
automatically by PLGS. Deuterium uptakes for all identified peptides were then filtered and
validated manually using DynamX 3.0 (Waters) as follows: only peptides identified with a
minimum fragment per amino acid of 0.2, a minimum intensity of 10^3, a length between 5 and
25 residues, a minimum PLGS score of 6.62 and a MH$^+$ tolerance of 5 ppm were kept. An
initial automated spectral processing step was conducted by DynamX followed by a manual
inspection of individual peptides for sufficient quality where only one charge state per peptide
was kept. Deuterium uptake is not corrected for back-exchange and is reported as relative
values. HDX-MS results were statistically validated using an in-house program
(Archaeopteryx), where the t-statistic threshold was set to 0.05. Only statistically significant
peptides showing a difference greater than 0.25 Da and 2% for at least two timepoints were
kept. HDX-MS results were illustrated on SRC open and closed conformations (PDB: 2SRC
and 1Y57 respectively) using PyMOL 2.5.4 (www.pymol.org).
1019
AlphaFold3 modeling
SRC SH2 and C-terminal pTyr complexes were modelled from sequence using AlphaFold3
through its public web server (https://www.alphafoldserver.com). Standard AlphaFold
procedures were followed, and the outputs were manually inspected to confirm the correct
orientation of the SH2 domain bound to peptide. Post-translational modifications were added
using the PTMs function on the selected residues. For each complex, five models were
generated, and their confidence metrics were evaluated using the AlphaFold3 ranking scores.
The highest-ranking model, determined by its pLDDT score, was selected for further analysis.
Structural representations were prepared using UCSF Chimera, and alignment was performed
using MatchMaker in Chimera X.
1030
Isothermal Titration Calorimetry
ITC experiment was measured by MicroCal PEAQ-ITC Automated at 25℃. After treatment,
SRC SH2 domains and peptides were buffer exchanged into fresh experimental buffer using
ultrafiltration prior to titrations. ITC experiments were performed by stepwise titration of the
peptides (400 µM) into an adiabatic cell containing SRC SH2 (40 µM), and the heat energy
change accompanying the reaction was detected upon each injection by comparison with a

reference cell. The stoichiometry of binding (n) was between 0.84 and 1.1 for all experiments.
Protein solution was placed in the 380 µl calorimeter cell and stirred at 750 rpm to ensure
rapid mixing, and 2 µl aliquots of the peptides were injected with a 120 s interval between each
injection until saturation. The heat changes were integrated after subtracting values obtained
when peptides were titrated into buffer, data were analyzed with the MicroCal PEAQ-ITC
software.
1043
Cell counting and Incucyte® imaging
MEFs were plated on fibronectin (Roche #10838039001) 5 µg/cm^2 coated 6-well plates at 0.2
x10^6 and grown over 6 days with a media change on day 3. On daily intervals, one well was
trypsinized, resuspended in normal growth medium and counted with trypan blue using the
countess cell counter (Invitrogen) for the manual growth curves. For live imaging, MEFs were
plated on fibronectin (Roche; 10838039001) 5 µg/cm^2 coated CytoOne 96-well plates at 2.5
x10^5 cells per well. The following day, the plate was transferred to an Incucyte® S3 live cell
imaging and analysis system (Sartorious; IC70682) and 4 regions were imaged per well every
2 h for 6 days. Media was refreshed on day 3.
1053
Immunofluorescence microscopy
Cells were plated on fibronectin (Roche #10838039001) 5 µg/cm^2 coated coverslips at 0.2
x10^6 and let to grow for 6 days with a media change on day 3. Then fixed using 4%
paraformaldehyde in PBS for 10 min. The cells were then permeabilized in 0.5% Triton X-100
and 3% BSA in PBS for 2 min and blocked in 0.2% Triton X-100 and 3% BSA in PBS (blocking
buffer) for 1 h. They were stained using Alexa Fluor™ Plus 647 Phalloidin (1:200) and Hoechst
(1:2000) for 1 h then washed 3 times in blocking buffer for 1 h. Images were taken using and
Leica Stellaris 8 with a 40X objective and 1.5 zoom. The nucleus of 6 different fields per
condition were counted.
1063
1064 **Ethical Statement**
1065 All animal experiments at The Babraham Institute were reviewed and approved by The Animal
1066 Welfare and Ethics Review Body and performed under Home Office Project license
1067 PP4624082.
1068

References

1069

1. Hunter, T. (2009). Tyrosine phosphorylation: thirty years and counting. Curr Opin Cell
Biol *21*, 140-146. 10.1016/j.ceb.2009.01.028.
2. Baselga, J. (2006). Targeting Tyrosine Kinases in Cancer: The Second Wave.
Science *312*, 1175-1178. 10.1126/science.1125951.
3. Östman, A., Hellberg, C., and Böhmer, F.D. (2006). Protein-tyrosine phosphatases
and cancer. Nature Reviews Cancer *6*, 307-320. 10.1038/nrc1837.
4. Baumgartner, C.K., Ebrahimi-Nik, H., Iracheta-Vellve, A., Hamel, K.M., Olander, K.E.,
Davis, T.G.R., McGuire, K.A., Halvorsen, G.T., Avila, O.I., Patel, C.H., et al. (2023).
The PTPN2/PTPN1 inhibitor ABBV-CLS-484 unleashes potent anti-tumour immunity.
Nature *622*, 850-862. 10.1038/s41586-023-06575-7.
5. Chen, Y.N., LaMarche, M.J., Chan, H.M., Fekkes, P., Garcia-Fortanet, J., Acker,
M.G., Antonakos, B., Chen, C.H., Chen, Z., Cooke, V.G., et al. (2016). Allosteric
inhibition of SHP2 phosphatase inhibits cancers driven by receptor tyrosine kinases.
Nature *535*, 148-152. 10.1038/nature18621.
6. Huyer, G., Liu, S., Kelly, J., Moffat, J., Payette, P., Kennedy, B., Tsaprailis, G.,
Gresser, M.J., and Ramachandran, C. (1997). Mechanism of Inhibition of Protein-
tyrosine Phosphatases by Vanadate and Pervanadate. Journal of Biological
Chemistry *272*, 843-851. 10.1074/jbc.272.2.843.
7. Fantus, I.G., Kadota, S., Deragon, G., Foster, B., and Posner, B.I. (1989).
Pervanadate [peroxide(s) of vanadate] mimics insulin action in rat adipocytes via
activation of the insulin receptor tyrosine kinase. Biochemistry *28*, 8864-8871.
10.1021/bi00448a027.
8. Irving, E., and Stoker, A.W. (2017). Vanadium Compounds as PTP Inhibitors.
Molecules *22*. 10.3390/molecules22122269.
9. Posner, B.I., Faure, R., Burgess, J.W., Bevan, A.P., Lachance, D., Zhang-Sun, G.,
Fantus, I.G., Ng, J.B., Hall, D.A., Lum, B.S., and et al. (1994). Peroxovanadium
compounds. A new class of potent phosphotyrosine phosphatase inhibitors which are
insulin mimetics. J Biol Chem *269*, 4596-4604.
10. Blanchetot, C., Chagnon, M., Dubé, N., Hallé, M., and Tremblay, M.L. (2005).
Substrate-trapping techniques in the identification of cellular PTP targets. Methods
35, 44-53. 10.1016/j.ymeth.2004.07.007.
11. Karisch, R., Fernandez, M., Taylor, P., Virtanen, C., St-Germain, J.R., Jin, L.L.,
Harris, I.S., Mori, J., Mak, T.W., Senis, Y.A., et al. (2011). Global proteomic
assessment of the classical protein-tyrosine phosphatome and "Redoxome". Cell
146, 826-840. 10.1016/j.cell.2011.07.020.
12. Boulven, I., Robin, P., Desmyter, C., Harbon, S., and Leiber, D. (2002). Differential
involvement of Src family kinases in pervanadate-mediated responses in rat
myometrial cells. Cellular Signalling *14*, 341-349. https://doi.org/10.1016/S0898-
6568(01)00269-8.
13. Shisheva, A., and Shechter, Y. (1993). Mechanism of pervanadate stimulation and
potentiation of insulin-activated glucose transport in rat adipocytes: dissociation from
vanadate effect. Endocrinology *133*, 1562-1568. 10.1210/endo.133.4.8404595.
14. Davis, S.J., and van der Merwe, P.A. (2006). The kinetic-segregation model: TCR
triggering and beyond. Nature Immunology *7*, 803. 10.1038/ni1369
https://www.nature.com/articles/ni1369#supplementary-information.
15. Imbert, V., Peyron, J.F., Farahi Far, D., Mari, B., Auberger, P., and Rossi, B. (1994).
Induction of tyrosine phosphorylation and T-cell activation by vanadate peroxide, an
inhibitor of protein tyrosine phosphatases. Biochem J *297 (Pt 1)*, 163-173.
10.1042/bj2970163.
16. Secrist, J.P., Burns, L.A., Karnitz, L., Koretzky, G.A., and Abraham, R.T. (1993).
Stimulatory effects of the protein tyrosine phosphatase inhibitor, pervanadate, on T-
cell activation events. J Biol Chem *268*, 5886-5893.

17. Courtney, A.H., Amacher, J.F., Kadlecek, T.A., Mollenauer, M.N., Au-Yeung, B.B.,
Kuriyan, J., and Weiss, A. (2017). A Phosphosite within the SH2 Domain of Lck
Regulates Its Activation by CD45. Mol Cell *67*, 498-511.e496.
10.1016/j.molcel.2017.06.024.
18. Courtney, A.H., Shvets, A.A., Lu, W., Griffante, G., Mollenauer, M., Horkova, V., Lo,
W.-L., Yu, S., Stepanek, O., Chakraborty, A.K., and Weiss, A. (2019). CD45
functions as a signaling gatekeeper in T cells. Science Signaling *12*, eaaw8151.
10.1126/scisignal.aaw8151.
19. Giannoni, E., Buricchi, F., Raugei, G., Ramponi, G., and Chiarugi, P. (2005).
Intracellular Reactive Oxygen Species Activate Src Tyrosine Kinase during Cell
Adhesion and Anchorage-Dependent Cell Growth. Molecular and Cellular Biology *25*,
6391-6403. 10.1128/MCB.25.15.6391-6403.2005.
20. Heppner, D.E. (2021). Structural insights into redox-active cysteine residues of the
Src family kinases. Redox Biol *41*, 101934. 10.1016/j.redox.2021.101934.
21. Heppner, D.E., Dustin, C.M., Liao, C., Hristova, M., Veith, C., Little, A.C., Ahlers,
B.A., White, S.L., Deng, B., Lam, Y.W., et al. (2018). Direct cysteine sulfenylation
drives activation of the Src kinase. Nat Commun *9*, 4522. 10.1038/s41467-018-
06790-1.
22. Truong, T.H., Ung, P.M., Palde, P.B., Paulsen, C.E., Schlessinger, A., and Carroll,
K.S. (2016). Molecular Basis for Redox Activation of Epidermal Growth Factor
Receptor Kinase. Cell Chem Biol *23*, 837-848. 10.1016/j.chembiol.2016.05.017.
23. Yoo, S.K., Starnes, T.W., Deng, Q., and Huttenlocher, A. (2011). Lyn is a redox
sensor that mediates leukocyte wound attraction in vivo. Nature *480*, 109-112.
10.1038/nature10632.
24. Zhang, H., Davies, K.J.A., and Forman, H.J. (2015). TGFβ1 rapidly activates Src
through a non-canonical redox signaling mechanism. Archives of Biochemistry and
Biophysics *568*, 1-7. https://doi.org/10.1016/j.abb.2015.01.001.
25. Mikalsen, S.-O., and Kaalhus, O. (1998). Properties of Pervanadate and
Permolybdate: CONNEXIN43, PHOSPHATASE INHIBITION, AND THIOL
REACTIVITY AS MODEL SYSTEMS *. Journal of Biological Chemistry *273*, 10036-
10045. 10.1074/jbc.273.16.10036.
26. Tao, G.-Z., Zhou, Q., Strnad, P., Salemi, M.R., Lee, Y.M., and Omary, M.B. (2005).
Human Ran Cysteine 112 Oxidation by Pervanadate Regulates Its Binding to
Keratins. Journal of Biological Chemistry *280*, 12162-12167.
27. Alcántara-Mejía, V.A., Rodríguez-Mercado, J.J., Mateos-Nava, R.A., Álvarez-
Barrera, L., Santiago-Osorio, E., Bonilla-González, E., and Altamirano-Lozano, M.A.
(2024). Oxidative damage and cell cycle delay induced by vanadium(III) in human
peripheral blood cells. Toxicology Reports *13*, 101695.
https://doi.org/10.1016/j.toxrep.2024.101695.
28. Faure, R., Baquiran, G., Bergeron, J.J., and Posner, B.I. (1992). The
dephosphorylation of insulin and epidermal growth factor receptors. Role of
endosome-associated phosphotyrosine phosphatase(s). Journal of Biological
Chemistry *267*, 11215-11221. https://doi.org/10.1016/S0021-9258(19)49898-0.
29. Kritsiligkou, P., Shen, T.K., and Dick, T.P. (2021). A comparison of Prx- and OxyR-
based H_2O_2 probes expressed in
S. cerevisiae. Journal of Biological Chemistry *297*.
10.1016/j.jbc.2021.100866.
30. Pak, V.V., Ezeriņa, D., Lyublinskaya, O.G., Pedre, B., Tyurin-Kuzmin, P.A., Mishina,
N.M., Thauvin, M., Young, D., Wahni, K., Martínez Gache, S.A., et al. (2020).
Ultrasensitive Genetically Encoded Indicator for Hydrogen Peroxide Identifies Roles
for the Oxidant in Cell Migration and Mitochondrial Function. Cell Metabolism *31*,
642-653.e646. 10.1016/j.cmet.2020.02.003.
31. Zhuravlev, A., Ezeriņa, D., Ivanova, J., Guriev, N., Pugovkina, N., Shatrova, A.,
Aksenov, N., Messens, J., and Lyublinskaya, O. (2024). HyPer as a tool to determine

1176		the reductive activity in cellular compartments. Redox Biology *70*, 103058.
1177		https://doi.org/10.1016/j.redox.2024.103058.
1178	32.	Woo, H.A., Yim, S.H., Shin, D.H., Kang, D., Yu, D.Y., and Rhee, S.G. (2010).
1179		Inactivation of peroxiredoxin I by phosphorylation allows localized H(2)O(2)
1180		accumulation for cell signaling. Cell *140*, 517-528. 10.1016/j.cell.2010.01.009.
1181	33.	Fan, C., Li, Q., Ross, D., and Engelhardt, J.F. (2003). Tyrosine Phosphorylation of
1182		IκBα Activates NFκB through a Redox-regulated and c-Src-dependent Mechanism
1183		Following Hypoxia/Reoxygenation*. Journal of Biological Chemistry *278*, 2072-2080.
1184		https://doi.org/10.1074/jbc.M206718200.
1185	34.	Bertoldo, J.B., Rodrigues, T., Dunsmore, L., Aprile, F.A., Marques, M.C., Rosado,
1186		L.A., Boutureira, O., Steinbrecher, T.B., Sherman, W., Corzana, F., et al. (2017). A
1187		Water-Bridged Cysteine-Cysteine Redox Regulation Mechanism in Bacterial Protein
1188		Tyrosine Phosphatases. Chem *3*, 665-677. 10.1016/j.chempr.2017.07.009.
1189	35.	Hubbard, S.R. (1999). Src autoinhibition: let us count the ways. Nat Struct Biol *6*,
1190		711-714. 10.1038/11468.
1191	36.	Moarefi, I., LaFevre-Bernt, M., Sicheri, F., Huse, M., Lee, C.H., Kuriyan, J., and
1192		Miller, W.T. (1997). Activation of the Src-family tyrosine kinase Hck by SH3 domain
1193		displacement. Nature *385*, 650-653. 10.1038/385650a0.
1194	37.	Xu, W., Doshi, A., Lei, M., Eck, M.J., and Harrison, S.C. (1999). Crystal Structures of
1195		c-Src Reveal Features of Its Autoinhibitory Mechanism. Molecular Cell *3*, 629-638.
1196		https://doi.org/10.1016/S1097-2765(00)80356-1.
1197	38.	Cowan-Jacob, S.W., Fendrich, G., Manley, P.W., Jahnke, W., Fabbro, D., Liebetanz,
1198		J., and Meyer, T. (2005). The crystal structure of a c-Src complex in an active
1199		conformation suggests possible steps in c-Src activation. Structure *13*, 861-871.
1200		10.1016/j.str.2005.03.012.
1201	39.	Boggon, T.J., and Eck, M.J. (2004). Structure and regulation of Src family kinases.
1202		Oncogene *23*, 7918-7927. 10.1038/sj.onc.1208081.
1203	40.	Meng, Y., and Roux, B. (2016). Computational study of the W260A activating mutant
1204		of Src tyrosine kinase. Protein Sci *25*, 219-230. 10.1002/pro.2731.
1205	41.	Meng, Y., and Roux, B. (2014). Locking the active conformation of c-Src kinase
1206		through the phosphorylation of the activation loop. J Mol Biol *426*, 423-435.
1207		10.1016/j.jmb.2013.10.001.
1208	42.	Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger,
1209		O., Willmore, L., Ballard, A.J., Bambrick, J., et al. (2024). Accurate structure
1210		prediction of biomolecular interactions with AlphaFold 3. Nature *630*, 493-500.
1211		10.1038/s41586-024-07487-w.
1212	43.	Payne, G., Stolz, L.A., Pei, D., Band, H., Shoelson, S.E., and Walsh, C.T. (1994).
1213		The phosphopeptide-binding specificity of Src family SH2 domains. Chem Biol *1*, 99-
1214		105. 10.1016/1074-5521(94)90047-7.
1215	44.	Bradshaw, J.M., Mitaxov, V., and Waksman, G. (1999). Investigation of
1216		phosphotyrosine recognition by the SH2 domain of the Src kinase11Edited by P. E.
1217		Wright. Journal of Molecular Biology *293*, 971-985.
1218		https://doi.org/10.1006/jmbi.1999.3190.
1219	45.	Morra, M., Simarro-Grande, M., Martin, M., Chen, A.S.-I., Lanyi, A., Silander, O.,
1220		Calpe, S., Davis, J., Pawson, T., Eck, M.J., et al. (2001). Characterization of SH2D1A
1221		Missense Mutations Identified in X-linked Lymphoproliferative Disease Patients*.
1222		Journal of Biological Chemistry *276*, 36809-36816.
1223		https://doi.org/10.1074/jbc.M101305200.
1224	46.	Bajusz, D., Pándy-Szekeres, G., Takács, Á., de Araujo, E.D., and Keserű, G.M.
1225		(2023). SH2db, an information system for the SH2 domain. Nucleic Acids Res *51*,
1226		W542-w552. 10.1093/nar/gkad420.
1227	47.	Visperas, P.R., Winger, J.A., Horton, T.M., Shah, N.H., Aum, D.J., Tao, A., Barros,
1228		T., Yan, Q., Wilson, C.G., Arkin, M.R., et al. (2015). Modification by covalent reaction
1229		or oxidation of cysteine residues in the tandem-SH2 domains of ZAP-70 and Syk can
1230		block phosphopeptide binding. Biochem J *465*, 149-161. 10.1042/bj20140793.

1231   48. Koudelková, L., Pataki, A.C., Tolde, O., Pavlik, V., Nobis, M., Gemperle, J.,
1232   Anderson, K., Brábek, J., and Rosel, D. (2019). Novel FRET-Based Src Biosensor
1233   Reveals Mechanisms of Src Activation and Its Dynamics in Focal Adhesions. Cell
1234   Chem Biol *26*, 255-268.e254. 10.1016/j.chembiol.2018.10.024.
1235   49. Arias-Romero, L.E., Saha, S., Villamar-Cruz, O., Yip, S.-C., Ethier, S.P., Zhang, Z.-
1236   Y., and Chernoff, J. (2009). Activation of Src by Protein Tyrosine Phosphatase 1B Is
1237   Required for ErbB2 Transformation of Human Breast Epithelial Cells. Cancer
1238   Research *69*, 4582-4588. 10.1158/0008-5472.CAN-08-4001.
1239   50. Playford, M.P., and Schaller, M.D. (2004). The interplay between Src and integrins in
1240   normal and tumor biology. Oncogene *23*, 7928-7946. 10.1038/sj.onc.1208080.
1241   51. Zheng, X.M., Wang, Y., and Fallen, C.J. (1992). Cell transformation and activation of
1242   pp60c-src by overexpression of a protein tyrosine phosphatase. Nature *359*, 336-
1243   339. 10.1038/359336a0.
1244   52. Dustin, C.M., Habibovic, A., Hristova, M., Schiffers, C., Morris, C.R., Lin, M.J., Bauer,
1245   R.A., Heppner, D.E., Daphtary, N., Aliyeva, M., and van der Vliet, A. (2021).
1246   Oxidation-Dependent Activation of Src Kinase Mediates Epithelial IL-33 Production
1247   and Signaling during Acute Airway Allergen Challenge. J Immunol *206*, 2989-2999.
1248   10.4049/jimmunol.2000995.
1249   53. Gumbiner, B.M., and Kim, N.-G. (2014). The Hippo-YAP signaling pathway and
1250   contact inhibition of growth. Journal of Cell Science *127*, 709-717.
1251   10.1242/jcs.140103.
1252   54. Blanchetot, C., Chagnon, M.I.J., Dubé, N., Hallé, M., and Tremblay, M.L. (2005).
1253   Substrate-trapping techniques in the identification of cellular PTP targets. Methods
1254   *35 1*, 44-53.
1255   55. Mercan, F., and Bennett, A.M. (2010). Analysis of protein tyrosine phosphatases and
1256   substrates. Curr Protoc Mol Biol *Chapter 18*, Unit 18.16.
1257   10.1002/0471142727.mb1816s91.
1258   56. Han, Y., Gao, Q., Xu, Y., Chen, K., Li, R., Guo, W., and Wang, S. (2024). Cysteine
1259   sulfenylation contributes to liver fibrosis via the regulation of EphB2-mediated
1260   signaling. Cell Death & Disease *15*, 602. 10.1038/s41419-024-06997-9.
1261   57. Queiroz, R.F., Stanley, C.P., Wolhuter, K., Kong, S.M.Y., Rajivan, R., McKinnon, N.,
1262   Nguyen, G.T.H., Roveri, A., Guttzeit, S., Eaton, P., et al. (2021). Hydrogen peroxide
1263   signaling via its transformation to a stereospecific alkyl hydroperoxide that escapes
1264   reductive inactivation. Nature Communications *12*, 6626. 10.1038/s41467-021-
1265   26991-5.
1266   58. Senga, T., Hasegawa, H., Tanaka, M., Rahman, M.A., Ito, S., and Hamaguchi, M.
1267   (2008). The cysteine-cluster motif of c-Src: Its role for the heavy metal-mediated
1268   activation of kinase. Cancer Science *99*, 571-575. https://doi.org/10.1111/j.1349-
1269   7006.2007.00714.x.
1270   59. van Vlimmeren, A.E., Voleti, R., Chartier, C.A., Jiang, Z., Karandur, D., Humphries,
1271   P.A., Lo, W.L., and Shah, N.H. (2024). The pathogenic T42A mutation in SHP2
1272   rewires the interaction specificity of its N-terminal regulatory domain. bioRxiv.
1273   10.1101/2023.07.10.548257.
1274   60. Evans, J.V., Ammer, A.G., Jett, J.E., Bolcato, C.A., Breaux, J.C., Martin, K.H., Culp,
1275   M.V., Gannett, P.M., and Weed, S.A. (2012). Src binds cortactin through an SH2
1276   domain cystine-mediated linkage. Journal of cell science *125*, 6185-6197.
1277   10.1242/jcs.121046.
1278   61. Xiao, H., Jedrychowski, M.P., Schweppe, D.K., Huttlin, E.L., Yu, Q., Heppner, D.E.,
1279   Li, J., Long, J., Mills, E.L., Szpyt, J., et al. (2020). A Quantitative Tissue-Specific
1280   Landscape of Protein Redox Regulation during Aging. Cell *180*, 968-983.e924.
1281   10.1016/j.cell.2020.02.012.
1282   62. Yan, Q., Tiago, B., R., V.P., Sebastian, D., A., K.T., Arthur, W., and and Kuriyan, J.
1283   (2013). Structural Basis for Activation of ZAP-70 by Phosphorylation of the SH2-
1284   Kinase Linker. Molecular and Cellular Biology *33*, 2188-2201. 10.1128/MCB.01637-
1285   12.

63.   (!!! INVALID CITATION !!! 44,45).

64.   Kaplan, K.B., Bibbins, K.B., Swedlow, J.R., Arnaud, M., Morgan, D.O., and Varmus,
H.E. (1994). Association of the amino-terminal half of c-Src with focal adhesions
alters their properties and is regulated by phosphorylation of tyrosine 527. Embo j *13*,
4745-4756. 10.1002/j.1460-2075.1994.tb06800.x.

65.   Hunter, M.V., Willoughby, P.M., Bruce, A.E.E., and Fernandez-Gonzalez, R. (2018).
Oxidative Stress Orchestrates Cell Polarity to Promote Embryonic Wound Healing.
Developmental Cell *47*, 377-387.e374. 10.1016/j.devcel.2018.10.013.

66.   Yang, Y., Dong, X., Zheng, S., Sun, J., Ye, J., Chen, J., Fang, Y., Zhao, B., Yin, Z.,
Cao, P., and Luo, L. (2020). GSTpi regulates VE-cadherin stabilization through
promoting S-glutathionylation of Src. Redox Biology *30*, 101416.
https://doi.org/10.1016/j.redox.2019.101416.

67.   Wheeler, D.L., Iida, M., and Dunn, E.F. (2009). The role of Src in solid tumors.
Oncologist *14*, 667-678. 10.1634/theoncologist.2009-0009.

68.   Glorieux, C., Liu, S., Trachootham, D., and Huang, P. (2024). Targeting ROS in
cancer: rationale and strategies. Nature Reviews Drug Discovery *23*, 583-606.
10.1038/s41573-024-00979-4.

69.   Rao, S., Gurbani, D., Du, G., Everley, R.A., Browne, C.M., Chaikuad, A., Tan, L.,
Schröder, M., Gondi, S., Ficarro, S.B., et al. (2019). Leveraging Compound
Promiscuity to Identify Targetable Cysteines within the Kinome. Cell Chemical
Biology *26*, 818-829.e819. https://doi.org/10.1016/j.chembiol.2019.02.021.

70.   Boike, L., Henning, N.J., and Nomura, D.K. (2022). Advances in covalent drug
discovery. Nature Reviews Drug Discovery *21*, 881-898. 10.1038/s41573-022-00542-
z.

71.   Li, J., Rix, U., Fang, B., Bai, Y., Edwards, A., Colinge, J., Bennett, K.L., Gao, J.,
Song, L., Eschrich, S., et al. (2010). A chemical and phosphoproteomic
characterization of dasatinib action in lung cancer. Nature Chemical Biology *6*, 291-
299. 10.1038/nchembio.332.

72.   Jurrus, E., Engel, D., Star, K., Monson, K., Brandi, J., Felberg, L.E., Brookes, D.H.,
Wilson, L., Chen, J., Liles, K., et al. (2018). Improvements to the APBS biomolecular
solvation software suite. Protein Sci *27*, 112-128. 10.1002/pro.3280.

73.   Fearnley, G.W., Young, K.A., Edgar, J.R., Antrobus, R., Hay, I.M., Liang, W.C.,
Martinez-Martin, N., Lin, W., Deane, J.E., and Sharpe, H.J. (2019). The homophilic
receptor PTPRK selectively dephosphorylates multiple junctional regulators to
promote cell-cell adhesion. Elife *8*. 10.7554/eLife.44597.

74.   Hay, I.M., Fearnley, G.W., Rios, P., Köhn, M., Sharpe, H.J., and Deane, J.E. (2020).
The receptor PTPRU is a redox sensitive pseudophosphatase. Nature
Communications *11*, 3219. 10.1038/s41467-020-17076-w.

75.   Byrne, D.P., Shrestha, S., Galler, M., Cao, M., Daly, L.A., Campbell, A.E., Eyers,
C.E., Veal, E.A., Kannan, N., and Eyers, P.A. (2020). Aurora A regulation by
reversible cysteine oxidation reveals evolutionarily conserved redox control of
Ser/Thr protein kinase activity. Sci Signal *13*. 10.1126/scisignal.aax2713.

76.   Wojdyla, K., and Rogowska-Wrzesinska, A. (2015). Differential alkylation-based
redox proteomics – Lessons learnt. Redox Biology *6*, 240-252.
https://doi.org/10.1016/j.redox.2015.08.005.

77.   Ferries, S., Perkins, S., Brownridge, P.J., Campbell, A.E., Eyers, P.A., Jones, A.R.,
and Eyers, C.E. (2017). Evaluation of Parameters for Confident Phosphorylation Site
Localization Using an Orbitrap Fusion Tribrid Mass Spectrometer. Journal of
proteome research *16 9*, 3448-3459.

78.   Daly, L.A., Byrne, D.P., Perkins, S., Brownridge, P.J., McDonnell, E., Jones, A.R.,
Eyers, P.A., and Eyers, C.E. (2023). Custom Workflow for the Confident Identification
of Sulfotyrosine-Containing Peptides and Their Discrimination from
Phosphopeptides. J Proteome Res *22*, 3754-3772. 10.1021/acs.jproteome.3c00425.

79.   Ferries, S., Perkins, S., Brownridge, P.J., Campbell, A., Eyers, P.A., Jones, A.R., and
Eyers, C.E. (2017). Evaluation of Parameters for Confident Phosphorylation Site

Localization Using an Orbitrap Fusion Tribrid Mass Spectrometer. J Proteome Res
16, 3448-3459. 10.1021/acs.jproteome.7b00337.
80.   Perkins, D.N., Pappin, D.J., Creasy, D.M., and Cottrell, J.S. (1999). Probability-based
protein identification by searching sequence databases using mass spectrometry
data. Electrophoresis *20*, 3551-3567. 10.1002/(sici)1522-
2683(19991201)20:18<3551::Aid-elps3551>3.0.Co;2-2.
81.   Silva, J.C., Gorenstein, M.V., Li, G.Z., Vissers, J.P., and Geromanos, S.J. (2006).
Absolute quantification of proteins by LCMSE: a virtue of parallel MS acquisition. Mol
Cell Proteomics *5*, 144-156. 10.1074/mcp.M500230-MCP200.
82.   Murphy, James M., Zhang, Q., Young, Samuel N., Reese, Michael L., Bailey,
Fiona P., Eyers, Patrick A., Ungureanu, D., Hammaren, H., Silvennoinen, O.,
Varghese, Leila N., et al. (2013). A robust methodology to subclassify pseudokinases
based on their nucleotide-binding properties. Biochemical Journal *457*, 323-334.
10.1042/BJ20131174.