Comparative Virome Profiling and Discovery of Novel Viruses in

Managed and Wild Bees using RNA-seq

Zixiao Zhao[1]; M. Catherine Farrell[2]; Rameshwor Pudasaini[1]; Jayla Marvin[2] ; Sandra M.

Rehan[3]; Hongmei Li-Byarlay[1,2 *]

[1]Agricultural Research Development Program, Central State University, Wilberforce,

OH, USA

[2]Department of Agricultural and Life Sciences, Central State University, Wilberforce,

OH, USA

[3]Department of Biology, York University, Toronto, ON, Canada

Author contact information

ZZ: zxzhao5@gmail.com

https://orcid.org/0000-0002-3627-5503

MCF: mfarrell@centralstate.edu

https://orcid.org/0000-0002-0081-3285

RP: rpudasaini@centralstate.edu

https://orcid.org/0000-0001-9957-1185

JM: JaylaMarvin832@gmail.com

SMR: sandra.rehan@gmail.com

https://orcid.org/0000-0002-6441-5155

HL-B: Hli-byarlay@centralstate.edu

https://orcid.org/0000-0001-6579-5172

*Corresponding Author

Hli-byarlay@centralstate.edu

Abstract

Insect pollinators face growing challenges from viral pathogens that may contribute to population declines, yet comparative data on virome composition in managed versus wild bees remain limited. Using RNA-Seq-based metatranscriptomic analysis, we compared the virome of *Apis mellifera* L. and *Ceratina calcarata* Robertson collected from agricultural and non-agricultural landscapes. We identified multiple known bee pathogens including Black Queen Cell Virus and Deformed Wing Virus in *A. mellifera*, and discovered three novel insect viruses—two iflaviruses and one entomopoxvirus—associated with *C. calcarata*. Notably, one novel iflavirus was identified in *A. mellifera*. Our findings reveal species- and landscape-specific viral prevalence, highlight viral diversity beyond canonical bee pathogens, and provide genomic resources for future functional studies. These data offer new insights into virus-host dynamics and underscore the utility of virome profiling in understanding pollinator health at the molecular level.

Keywords: Honey bees, agricultural landscape, transcriptomics, bee virus, iflavirus, entomopoxvirus

Introduction

Animal pollinators have critical ecological functions in maintaining the diversity of global ecosystems [1, 2]. About 87.5% of flowering plants rely on animal pollinators for successful reproduction [3]. In agriculture, approximately 75% of crop species require pollinators, contributing about 30-35% of total crop production [4, 5]. Bees (Hymenoptera: Apidae) are considered the most abundant and important pollinators globally. Honey bee species (*Apis* spp.), with a global distribution due to domestication, contribute about 80% of total pollination and create about $700 million value annually in the US [6]. However, the ecological significance of non-*Apis* wild bee species shouldn't be neglected.   Wild bees provide pollination services to many crops, even when managed honey bees are not available [7, 8]. Among the over 17,000 wild bee species worldwide, with a wide range of social structures from eusocial to solitary, eusocial species may be more susceptible to viral infections due to their high colony densities, frequent social interactions and genetic factors [19]. For last several decades, both wild and managed honey bees are suffering from colony declines [9-12], which not only causes economical losses to apiary industry, but more importantly, reduces pollination service and imposes potential risks for global food security.

The causes of   bee populations decline are complex. Factors responsible for bee populations decline include habitat loss and land fragmentation, increased use of chemical pesticides and other agrochemicals, introduction of pests and pathogens, and climate change [11, 13]. Among these stressors, pathogenic viruses are considered the direct and one of the most significant hazards to both managed and wild bee populations (Ullah et al. 2020). Currently identified bee pathogenic viruses belong to positive-sense single-stranded RNA (+ss RNA) virus category. The majority of the pathogenic bee viruses are from two families under the order Picornavirales:

Iflaviridae and Dicistroviridae. Iflaviridae has only one genus: *Iflavirus*, which includes

Sacbrood virus (SBV), Deformed wing virus (DWV), Varroa destructor virus-1 (VDV-1),

and Slow bee paralysis virus (SBPV). While the family Dicistroviridae includes   Black

queen cell virus (BQCV), Kashmir bee virus (KBV), Acute bee paralysis virus (ABPV),

and Israeli acute paralysis virus (IAPV) [14]. Two additional   pathogenic viruses of

bees are Chronic bee paralysis virus (CBPV), an unclassified virus with two genomic

RNA segments [15], and Lake Sinai viruses (LSVs), members of the order

Nodamuvirales [14], with eight identified lineages [16]. The symptoms caused by

pathogenic viruses vary, including body deformities, color changes, uncapped cells

with larvae or pupae, paralysis of adults, which eventually increase mortality and

lead to colony degradation. IAPV infections have been discovered association with

colony collapse disorder (CCD), which responsible for large-scale honey bee colony

loss in the US [17].

Despite managed honey bees, the pathogenic viruses can also infect wild bee

species [18-20]. For example, DWV has been reported to infect a wide range of

hymenopterans, including some social and solitary bees as well as several wasp

species [18, 20, 21]. Dolezal et al. [19] reported DWV and   SBV were widely

detected in wild bees collected in Iowa, while IAPV, LSV and BQCV were also detected

in wild be species. Furthermore, DWV, BQCV and SBV can be deposited to pollen and

remain infective, which can contribute to inter-taxa transmission among bees, and

wasps [20]. These studies suggested viruses can be a potential factor responsible for

colony declines in wild bee species.

Although agriculture provides important foraging resources for bees, it is also

considered one of the major reasons for colony declines. It has converted natural

ecosystems to farmland primary for food and fiber productions. This process is

accompanied by loss of biodiversity and use of agrochemicals causing habitat loss,

fragmentation and degradation. Agriculture can also alter host-pathogen interactions

at multiple levels. A study   demonstrated that the relevance of plant viruses is

negatively correlated with the biodiversity in agricultural landscape [22]. However,

whether agricultural activities affect viral prevalence in insects, especially in honey

bees and other bee species, is still unknown.

The small carpenter bee, *Ceratina calcarata* Robertson (Hymenoptera: Apidae),

is a wild bee species native to the eastern part of North America. It is considered as

an indicator species of healthy ecosystems and important pollinators for several

native and agricultural plants [23]. Due to the feature of subsociality, *C. calcarata* is

used for studying pollinator ecology, behavior, evolution, genomics and nutrition

[24-27]. However, little is known about the virology of this species. The impact of

viruses on *C. calcarata* populations and whether the viral prevalence is related to

agricultural activities are not clear.

Here we presented our metagenomic studies on viruses associated with western

honey bee (*A. mellifera* L.) and *C. calcarata* in response to different agricultural

intensifications. Bees were collected from three landscape types representing levels

of agricultural intensification: conventional farming, organic farming and roadside

with no agricultural activities. The viral RNA was extracted from collected bee

samples and subjected to high-throughput sequencing. We analyzed known and

novel viruses associated with both bee species, which were highly specific to location

and bee species. Our study provides a new methodology and unique knowledge for

bee-virus interactions.

Results

Reconstruct virome from viral RNA-Seq reads

The RNA-Seq of extracted viral RNA generates approximately 17 to 36 million

raw read pairs for each *A. mellifera* and *C. calcarata* sample. After quality trimming

and partially removing reads from respective host genome (Amel HAv3.1

PRJNA471592 and Ccalc.v3 PRJNA791561), 5 to 25 million pairs were kept for each

sample (**Supplementary file S1**). Trinity *de nove* assembled 366,289 contigs for *A.*

mellifera and 203,317 contigs for *C. calcarata* virome, which consists of 188,433 and

91,314 non-redundant clusters. The assembled contigs were subject to BLASTN

searches against non-redundant databases (nr), which claimed 541 clusters and 1187

contigs from *A. mellifera* assembly potentially belong to viruses. In contrast, 62

clusters and 84 contigs from *C. calcarata* assembly were potentially from viruses. A

large portion of assembled contigs were from either host genes or the plants they

have visited.

Viruses infecting bees and other insects detected in A. mellifera and C. calcarata

virome assemblies

BLASTN search identified four known bee pathogenic viruses from *A. mellifera*

virome: BQCV, DWV, SBV, and LSV3 with fully assembled genome sequences. Two

other members of the LSV group: LSV1 and LSV2 were detected. But their sequences

were only partially assembled. Positive and negative strands were detected in all of

the above viruses except LSV2, from which only positive strand was detected. We

also detected *Apis mellifera* filamentous virus (AmFV) [28] and Hubei partiti-like virus

34 [29] in *A. mellifera*. The AmFV were partially assembled and the Hubei partiti-like

virus 34 demonstrated a fully assembled genome (**Table 1A**). In *C. calcarata*, BQCV

was the only detected bee pathogenic virus and appeared as partial-assembled as

negative-strand contig (**Table 1B**).

Plant viruses discovered in A. mellifera and C. calcarata virome assemblies

A range of plant viruses were detected in virome assemblies of both bee species.

In *A. mellifera,* white clover cryptic virus 1 (WCCV-1) and white clover cryptic virus 2

(WCCV-2) from genus *Alphapartitivirus* (family: Partitiviridae, +ssRNA), as well as

white clover mosaic virus (WCMV) from genus *Potexvirus* (family: Alphaflexiviridae,

+ssRNA) were fully assembled. Blackberry chlorotic ringspot virus (BCRV) from genus

Ilarvirus (family: Bromoviridae, +ssRNA) and *Raphanus sativus* cryptic virus 1 (RsPV1)

from Partitiviridae (+ssRNA) have one genomic segment fully assembled, while other

segments were either partially assembled or completely missing. Partially assembled

plant viruses include 7 +ssRNA viruses: peanut stunt virus (PSV) and tobacco ringspot

virus (TRSV) from Bromoviridae, clover yellow mosaic virus (ClYMV) from

Alphaflexiviridae, red clover nepovirus A (RCNVA) , tobacco streak virus (TSV), and

tomato ringspot virus (ToRSV) from genus *Nepovirus* (family: Secoviridae), red clover

vein mosaic virus (RCVMV) from Betaflexiviridae, and one -ssRNA virus: *Pinus flexilis*

virus 1 (PiFleV1) from Rhabdoviridae (**Table 1A**). However, the plant viruses in *C.*

calcarata virome were considerably less diverse compared to *A. mellifera*. WCCV-2

was the only fully assembled plant virus, while WCCV-1, BCRV, ToRSV, and WClMV

were partially assembled in *C. calcarata* (**Table 1B**).

Novel viruses in A. mellifera and C. calcarata virome assemblies

The genus *Iflavirus* (family: Iflaviridae) consists of +ssRNA viruses infecting

arthropods, including three bee pathogenic viruses SBV, DWV, and SBPV. Two novel

viruses under genus *Iflavirus*, referred to as unknown iflavirus 1 and unknown

iflavirus 2, were identified in *C. calcarata* virome assembly as positive strands. The

phylogenetic analysis using translated polyprotein amino acid sequence suggests

both viruses are in the same group as SBV. The unknown iflavirus-1 has the highest

identity with Darwin bee virus-2, an unclassified picornavirus infecting honey bee [30]

(**Figure 1A**). This virus was also detected in *A. mellifera* virome assembly as

a negative strand, suggesting that it can infect and replicate in honey bees.

In *C. calcarata*, we detected a novel poxvirus (a double-strand DNA virus). Its

DNA-dependent RNA polymerase RPO132 is partially assembled. Phylogenetic

analysis showed this virus is closely related to the Yalta virus, an uncategorized

poxvirus from *Drosophila melanogaster* and *Diachasmimorpha* entomopoxvirus

(DIEPV), from parasitoid wasp *Diachasmimorpha longicaudata*. The three viruses

form a sister cluster to other entomopoxviruses (EPVs) (genus: *Alphaentomopoxvirus*,

Betaentomopoxvirus, and *Deltaentomopoxvirus*), while all the vertebrate poxviruses

(genus: *Avipoxvirus*, *Capripoxvirus*, and *Parapoxvirus*) are clustered into a separate

group. The result suggests this novel virus belongs to the subfamily

Entomopoxvirinae (EPV) that primarily infect insects (**Figure 1E**).

Three additional novel plant viruses were detected in the *A. mellifera* virome,

including one from the genus *Nepovirus* (unknown nepovirus 1, family: Secoviridae),

one from family Partitiviridae (unknown parititivirus 1, family: Partitiviridae), and one

of genus *Amalgavirus* (unknown Amalgavirus 1, family: Amalgaviridae). Nepovirus

has two positive-strand genomic RNAs [31]. For unknown nepovirus 1, the open

reading frame (ORF) of polyprotein RNA2 was fully assembled. However, the

correspondent RNA1 encoding viral coat protein was not detected. The phylogeny of

the polyprotein sequence shows that it is most closely related to Blueberry leaf

mottle virus and Grapevine Bulgarian latent virus, forming a separate cluster (**Figure**

1B). Similarly, only RNA1 encoding RNA-dependent RNA polymerase (RdRp) was fully

assembled for another potential plant virus: unknown parititivirus 1. The

phylogenetic analysis of RdRp showed that it is clustered with several uncategorized

partitivirus viruses associated with plant: *Polygonatum kingianum* cryptic virus 1,

Panax cryptic virus 3, and *Dichroa partitivirus* 1. This cluster is a sister group to

plant-infecting genus *Alphapartitivirus* (**Figure 1D**). The genome of unknown

amalgavirus 1 was fully assembled with two complete ORF: the ORF1+2 fusion

protein responsible for genome replication, and a coat protein. The phylogeny of

fusion proteins showed that it is the most closed to Vicia cryptic virus M (VCV-M) and

Medicago sativa amalgavirus 1 (MsAV1) and have a clear separation with

Zygosaccharomyces bailii virus Z (ZbV-Z), a yeast-infecting virus belongs to the genus

Zybavirus (**Figure 1C**).

In addition, other viruses, including phages and a human virus: Merkel cell

polyomavirus (Family: Polyomaviridae) were identified in both virome assemblies.

The viral genome consists of circular double-stranded RNA (dsRNA), which resulted in

the assembly of two identical full-length viral sequences into one contig in *C.*

calcarata virome. In *A. mellifera*, only partial sequence was assembled. How the

Merkel cell polyomavirus was introduced into virome and its effects in both species,

are still unknown.

Abundance of viruses in response to different agricultural landscapes

The distributions of the above detected viruses were highly site-specific. In *A.*

mellifera, among all identified insect viruses, five known bee viruses: BQCV, DWV,

LSVs, SBV, and AmFV were detected at all of the experimental sites regardless

of agricultural practices, with AmFV only marginally detected in conventional farm C1.

In contrast, the novel unknown ilfavirus 1 was only detected in samples from organic

farm O2 (**Figure 2A**). For five known bee viruses, the edgeR did not show any

significant differences at virus level. However, a separate 2834-bp LSV3 segment,

containing ORFs of two viral proteins, was significantly increased in bees collected

from organic sites compared to bees collected from roadside site (FDR: 0.04). The

Hubei partiti-like virus was also identified at all sites except conventional farm C1

(**Figure 2A**). Its full-length sequence was significantly higher in bee samples collected

from organic farms compared to bee samples collected from both conventional and

roadside sites (FDR: 0.006 to conventional, 0.003 to roadside).

The presence and levels of plant viruses were highly dependent on the sampling

locations. For instance, TSV and PiFleV1 were detected only at one separate roadside

site each, while RCNVA and TRSV were found only at one conventional and one

organic site, respectively. RCVMV was identified across three conventional sites, but

at relatively low levels. WCIMV, WCCV-2, and ToRSV were expressed at all

experimental sites (**Figure 2B**). WCCV-1 was the only significant plant virus detected

by edgeR. The levels of both genome segments decreased significantly in organic

sites, compared to both conventional (FDR, RNA1: 0.011, RNA2: 0.006) and roadside

sites (FDR, RNA1: 0.006, RNA2: 0.003). The viral levels of BRCV at roadside sites were

either very low or undetected, but the overall level is too low to perform statistical

analysis in edgeR.

C. calcarata hosted relatively fewer viruses compared to *A. mellifera*. The

presence of insect viruses was also highly site-specific. BQCV and unknown

entomopoxvirus 1 were only found at one conventional site, whereas two novel

ilfavirus were detected in roadside sites and conventional farm C1. None of the

viruses were detected from bee samples collected from organic sites, conventional

farm C3, and roadside R2 (**Figure 2C**). Regarding the plant viruses, WCCV-1 and

WCCV-2 had a similar pattern, being detected only at conventional farms, while

WCIMV and ToRSV were detected in few bee samples collected from roadside sites as

well (**Figure 2D**). However, their levels were not statistically significant. Similarly,

none of the plant viruses were detected in bee samples collected from organic sites.

Discussion

The metagenomic approach based on viral RNA-Seq was adopted for identifying and quantifying known viruses, as well as discovering novel viruses. The present study surveyed the abundance of viruses associated with two bee species. Findings show that the virome compositions were site-specific. However, the *A. mellifera* carried a wider range of both insect and plant viruses than *C. calcarata*. The results also indicate that the *A. mellifera* may transmit plant diseases more effectively than *C. calcarata*.

The Honey bee viruses in A. mellifera and spillover *to C. calcarata*

The levels of BQCV, DWV and SBV were in high levels in *A. mellifera* virome at all experimental sites, suggesting the high prevalence of these viruses in *A. mellifera*. In contrast, none of the acute bee paralysis virus complex (ABPV, KBV and IAPV) were detected. This result is consistent with other studies in the Northeast of the US [20, 32]. LSV is a highly variable, multi-strain bee virus that does not have any significant symptoms but is likely to weaken colonies. The LSV contigs assembled in this study are most closely related to LSV1, LSV2 and LSV3. The LSV3 was detected from all experimental sites and the only LSV with full-length genome sequence assembled. The LSV3 was originally identified in Slovenia [33] and has been detected in the Western US [34]. The study confirmed LSV3 has been spread to the New England area, in the US.

Two known viruses associated with *A. mellifera* were also detected: AmFV and Hubei partiti-like virus 34. The AmFV is a weak pathogen in *A. mellifera* and rarely causes significant symptoms. Its infection has been reported related to BQCV and DWV [28, 35]. We detected AmFV from all sites, while its viral loads were from extremely low at conventional farm C1 to medium-high at conventional farm C3. The

Hubei partiti-like virus 34 was first discovered as an arthropod RNA virus [36] and

was reported associated with hymenopteran species, including *A. mellifera* and Asian

hornet (*Vespa velutina*) in East Asia and South Europe [29, 37]. The current study

reported the low to moderate infection of Hubei partiti-like virus 34 from all sites

except conventional farm C3 site. The potential impact of this virus on honey bee

health is still unknown.

The spillover of managed honey bee pathogen to wild bees has been considered

as a threat to native bee populations [38, 39]. In genus *Ceratina*, DWV was found

spillover to two species *C. dupla* [20] and *C. smaragdula* [40]. The current study

showed that, given the high prevalence of both BQCV and DWV in *A. mellifera*, BQCV

may have spilled over to *C. calcarata*, while no evidence of DWV spillover was

observed. BQCV was only detected from conventional farm C2 in relatively low

prevalence and viral load, indicating that this spillover to *C. calcarata* is not a

common scenario.

Potential viruses infecting A. mellifera and C. calcarata

The primary hosts for iflaviruses are arthropods. We discovered two closely

related novel ifla-like viruses. Unknown iflavirus 1 was detected in both *A. mellifera*

and *C. calcarata*, while the unknown iflavirus 2 was detected only in *C. calcarata*. The

prevalence of both viruses was low, however, the viral load of unknown ilfavirus 2

was slightly higher in *C. calcarata* samples collected from conventional farm C1 and

roadside site 1, compared to the unknown ilfavirus 1 in the same samples.

The EPVs are double-stranded DNA (dsDNA) viruses that infect insects and

predominantly insect pathogens. The unknown entomopoxivirus 1 detected in *C.*

calcarata are more closely related to Yalta viruses and DIEPV than other

entomopathogenic viruses. Although the role of Yalta virus is still unknown, the

DIEPV has been found replicated in both parasitoid wasp *D. longicaudata* and its host

fruit fly (*Anastrepha suspensa*). The replication of virus is highly virulent to the host

fruit fly and no negative impact to wasp host, suggesting a mutualistic symbiotic

relationship of DIEPV to parasitoid wasp [41]. Although both *C. calcarata* and *D.*

longicaudata are from order Hymenoptera, no such parasitoid relationship was

observed in *C. calcarata.* Whether this novel entomopoxivirus primary infects *C.*

calcarata or just an overspill event to *C. calcarata* yet to be studied.

Although our findings reveal the presence of viral sequences in bee samples, it does

not confirm active infection or pathogenicity in these bees. Some of the viral

sequences may originate from environmental exposure, gut contents, or external

body surfaces, rather than systemic infection. Therefore, the present study does not

infer a direct relationship between viral presence and bee health or stress levels

(Olgun et al., 2020).

Bees carrying plant viruses

Many plant viruses can travel within or attach to pollen grains [42, 43], including

BRCV, TSV, PSV, WCCV-1, WCCV-2, TRSV, ToRSV, and WClMV as observed in the

present study   However, only TSV, TRSV, and ToRSV have been proven vertically and

horizontally transmitted by pollen. The rest were only detected by RNA-Seq [43].

These findings show that pollinators such as *A. mellifera* posing potential risks due to

these viruses. However, none of the detected viruses were on the list of known

honey bee-transmitted viruses, according to Fetters et al [43]. TRSV was reported

infect honey bee [44], however whether cross-kingdom infection assist transmitting

diseases is unknown. On the other hand, bees are a powerful tool for monitoring

plant diseases [45]. The plant viruses detected in *A. mellifera* resembles to the results

of other metagenomic studies of bees and pollen at family and order levels [45, 46],

while the species composition is determined by plant and prevalence of plant viruses

at the location.

Compared to *A. mellifera*, plant viruses in *C. calcarata* are much less diverse.

Previous studies have shown that pollen from clover and rose is a major food

resource for bees [47], and white clover pollen significantly increases the fitness of

C. calcarata [23]. Consistent with that, among four plant viruses detected in *C.*

calcarata virome, WCCV-1, WCCV-2, WClMV primarily infect white clover (*Trifolium*

repens), and the host range of ToRSV include rose and red clover (*T. pratense*). None

of the detected viruses originated beyond *C. calcarata's* foraging range, suggesting

that *C. calcarata* may be an ideal bee species for monitoring white clover-associated

pathogens within localized areas.

Material and method

Sample collection

Both honey bee (*A. mellifera*) and small carpenter bee (*C. calcarata*) samples

were collected from sites representing three different types of land use: conventional

farm, organic farm and roadside. Honey bees and the small carpenter bees were

collected by sweep nets from May to July 2018 in Strafford County, New Hampshire.

We collected both species within three conventional farms and three organic farms.

The roadside samples were collected in the urbanized area within the University of

New Hampshire campus. The detailed about locations and GPS coordinates are

described in previous publications [48, 49]. At least ???? bees were collected at each

sampling site, and to minimize age-related variation, all bee samples were collected

from forager bees. Bee samples were flash-frozen in liquid nitrogen and temporarily

stored at -80 °C.

RNA extraction, sequencing and data cleaning

The viral RNA was extracted from *A. mellifera* and *C. calcarata* individually by

Quick-RNA viral kit (ZYMO RESEARCH, Irvine, CA). DNase I (ZYMO RESEARCH)

in-column digestions were applied to remove genomic and environmental DNA. For

mRNA library generation, 200ng of total RNA was treated with NEBNext Poly mRNA

Magnetic Isolation Module E7490L (New England Biolabs, USA) following the

manufacturer protocol. Subsequently, the isolated mRNA was fragmented for 10

minutes. cDNA was synthesized and amplified for 12 PCR cycles using NEBNext Ultra

II Directional (stranded) RNA Library Prep Kit E7760L for Illumina (New England

Biolabs, USA) with NEBNext Multilex oligos indexes kit E6440S following the

manufacturer directions. Distributions of the template length and adapter-dimer

contamination were assessed using an Agilent 2100 Bioanalyzer and High Sensitivity

DNA kit (Agilent Technologies, Inc). The concentration of cDNA libraries was

determined using Invitrogen Qubit dsDNA HS reagents and read on a Qubit

Fluorometer (Thermo Fisher), and the cDNA libraries were paired-end 150bp format

sequenced on a Novaseq SP system (Illumina, San Diego, CA). In total, 17 honey bee

samples and 16 *C. calcarata* samples were sequenced (**Supplementary File S1**). Raw

reads were subject to adapter and quality trimming by Trimmomatic [50]. Sample

metadata from collection was used to assign each sample to either *A. mellifera* or *C.*

calcarata. To remove endogenous transcripts from bee hosts, the reads were

respectively aligned to honey bee and small carpenter bee [25] genomes (including

mitochondrial sequences) using Bowtie2 [51]. Read pairs that were concordantly

aligned to bee genomes were considered as endogenous transcripts and removed

from further analysis.

Virome assembly and viral identification

378 To reconstruct the virome of *A. mellifera* or *C. calcarata,* we first merged all the

379 pre-cleaned reads from the same organism to increase the sequence coverage. Then

380 two *de novo* assembler: Trinity [52] and metaSPAdes [53] were used with default

381 setting. The results were comparable; however, the total running time for Trinity was

382 lower. Therefore, we chose Trinity for the subsequent analyses.

383 Despite the pre-cleaning process, the assembled virome still contains host

384 sequences. To identify virus sequences, we used BLASTN to search assembled contigs

385 to default nr/nt database with E-value cutoff of 1e-7. Contigs with first three top hits

386 to viral sequence from database was retained as viral contigs. For further virus

387 identification, we examined the blast results of contigs >500 nt. A contig was

388 assigned as a known virus if it aligned stringently (E-value = 0) to existing viruses. If

389 the alignment is less stringent (0<E-value<1e-7), it was assigned as a novel virus. The

390 "full-length" or "partial" labels were based on the completeness and intactness of

391 viral protein of each viral contig. For potential novel viral contigs, ORF finder and

392 BLASTP were used to translate and predict viral protein. MEGA 11 [54] was used to

393 construct Maximum-likelihood phylogenetic tree based on the sequence of

394 homologous proteins of known and novel viruses.

396 *Abundance of viruses and differential analysis between landscapes*

397 To quantify the abundance of each contigs, the clean reads were mapped to

398 virome assembly of respective species using Bowtie 2 [51]. Multiple alignment was

399 allowed to obtain maximum sensitivity. The read counts were obtained by

400 FeatureCounts [55] and were normalized and converted to count per million reads

401 (CPM) by R edgeR package [56]. We also used edgeR to conduct differential analysis

402 between different landscape types. Prior to that, each contig, including endogenous

403 and viral, was filtered based on its expression level [57]. Only contigs with sufficiently

high expression levels were retained for statistical tests. The contigs with FDR less

than 0.05 were considered significantly difference between landscapes. The analysis

then focused on viral contigs as defined by the viral identification step.

Acknowledgements:

We are grateful for the help received from Stephanie Gardner, Katherine Odanaka,

and Wyatt Shell with collecting bee samples. We thank the Genomic Shared Resource

Shared Resource at The Ohio State University Comprehensive Cancer Center,

Columbus, OH for generating the virus RNA-seq data reported in this

publication. Research reported in this publication was supported by USDA Evans

Allen Fund NI231445XXXXG004, NI241445XXXXG004 for HL-B and ZZ, USDA CBG

GRANT13202344 for HL-B, USDA award 2020-67014-31557 for MCF, and the NSF

Louis Stokes Alliances for Minority Participation (LSAMP) program Ohio Alliance

scholarship for JM, the Ohio State University Comprehensive Cancer Center and the

National Institutes of Health under grant number P30 CA016058.

Author contributions:

HL-B and SMR designed the experiment. SMR conducted the field studies and

provided the samples. MCF and JM prepared samples for RNA-seq. ZZ conducted

data analysis and wrote the first draft of the manuscript. All authors reviewed the

final manuscript.

 Figures and Tables:

 Figure 1. Phylogenetic trees of novel viruses discovered from *A. mellifera* and *C. calcarata* virome

 assemblies by Maximum-likelihood analysis of sequences of viral proteins: **(A)** polyprotein (iflavirus),

 (B) polyprotein from RNA2 (neprovirus), **(C)** RNA-dependent RNA polymerase (amalgavirus), **(D)**

 RNA-dependent RNA polymerase from RNA1 (partitivirus), and **(E)** RNA-dependent RNA polymerase

 submit RPO132 (poxvirus). The names in bold correspond to honeybee-associated viruses.

Abbreviations and protein accessions of known viruses are included in **Supplementary File S2**.

Figure 2. Heatmaps demonstrating the relative abundance of **(A)** insect viruses in *A. mellifera*, **(B)**

plant viruses in *A. mellifera*, **(C)** insect viruses in C. *calcarata*, and **(D)** plant viruses in *C. calcarata* at

each sampling site. Average normalized count-per-million reads (cpm) from RNA-Seq was calculated

for each virus of each site and used in the heatmap. White color indicates the virus is undetected on

that site. Potential insect and plant viruses were included.

Table 1. List of identified insect and plant viruses in **(A)** *A. mellifera* and **(B)** *C. calcarata* virome

assemblies, assembled from RNA-Seq data

A

Virus name	Abbr.	Type	Family	Genus	Host	Assemble status	Strand
Black queen cell virus	BQCV	(+)ssRNA	Dicistroviridae	*Triatovirus*	Bees	Full	+/-
Deformed wing virus	DWV	(+)ssRNA	Iflaviridae	*Iflavirus*	Bees	Full	+/-
Sacbrood virus	SBV	(+)ssRNA	Iflaviridae	*Iflavirus*	Bees	Full	+/-
Lake Sinai virus 3	LSV3	(+)ssRNA	Sinhaliviridae	*Sinaivirus*	Bees	Full	+/-
Lake Sinai virus 1	LSV1	(+)ssRNA	Sinhaliviridae	*Sinaivirus*	Bees	Partial	+/-
Lake Sinai virus 2	LSV2	(+)ssRNA	Sinhaliviridae	*Sinaivirus*	Bees	Partial	+
Apis mellifera filamentous virus	AmFV	dsDNA	*unclassified*	*unclassified*	Bees	Partial	+/-
Hubei partiti-like virus 34	-	RNA	*unclassified*	*unclassified*	Bees	Full	+
Blackberry chlorotic ringspot virus	BCRV	(+)ssRNA	Bromoviridae	*Ilarvirus*	Plants	RNA1: Full	+
						RNA2: Partial	-
						RNA3: Partial	+
Tobacco streak virus	TSV	(+)ssRNA	Bromoviridae	*Ilarvirus*	Plants	RNA1: Partial	+/-
						RNA2: Partial	+/-
						RNA3: Partial	+
Peanut stunt virus	PSV	(+)ssRNA	Bromoviridae	*Cucumovirus*	Plants	RNA1: Partial	+
						RNA2: Partial	+
						RNA3: Partial	-
Raphanus sativus cryptic virus 1	RasCV1	dsRNA	Partitiviridae	*unclassified*	Plants	RNA1: Full	-
						RNA2: Missing	
White clover cryptic virus 1	WCCV-1	dsRNA	Partitiviridae	*Alphapartitivirus*	Plants	RNA1: Full	-
						RNA2: Full	-

Virus name	Abbr.	Type	Family	Genus	Host	Assembly status	Strand
White clover cryptic virus 2	WCCV-2	dsRNA	Partitiviridae	*Alphapartitivirus*	Plants	RNA1: Full	-
						RNA2: Full	-
Red clover nepovirus A	RCNVA	(+)ssRNA	Secoviridae	*Nepovirus*	Plants	RNA1: Partial	+
						RNA2: Partial	+/-
Tobacco ringspot virus	TRSV	(+)ssRNA	Secoviridae	*Nepovirus*	Plants	RNA1: Partial	+/-
						RNA2: Partial	-
Tomato ringspot virus	ToRSV	(+)ssRNA	Secoviridae	*Nepovirus*	Plants	RNA1: Partial	+/-
						RNA2: Partial	-
White clover mosaic virus	WClMV	(+)ssRNA	Alphaflexiviridae	*Potexvirus*	Plants	Full	+/-
Clover yellow mosaic virus	ClYMV	(+)ssRNA	Alphaflexiviridae	*Potexvirus*	Plants	Partial	+/-
Red clover vein mosaic virus	RCVMV	(+)ssRNA	Betaflexiviridae	*Carlavirus*	Plants	Partial	+/-
Pinus flexilis virus 1	PiFleV1	(-)ssRNA	Rhabdoviridae	*unclassified*	Plants	Partial	+/-

445   **B**

Virus name	Abbr.	Type	Family	Genus	Host	Assembly status	Strand
Black queen cell virus	BQCV	(+)ssRNA	Dicistroviridae	*Triatovirus*	Bees	Partial	-
Tomato ringspot virus	ToRSV	(+)ssRNA	Secoviridae	*Nepovirus*	Plants	RNA1: Partial	+/-
						RNA2: Partial	+/-
White clover cryptic virus 1	WCCV-1	dsRNA	Partitiviridae	*Alphapartitivirus*	Plants	RNA1: Full	-
						RNA2: Missing	
White clover cryptic virus 2	WCCV-2	dsRNA	Partitiviridae	*Alphapartitivirus*	Plants	RNA1: Full	+
						RNA2: Full	-
White clover mosaic virus	WClMV	(+)ssRNA	Alphaflexiviridae	*Potexvirus*	Plants	Partial	+/-

448   **Table 2.** List of novel viruses detected in *A. mellifera* and *C. calcarata* virome assemblies assembled

449   from RNA-Seq data

Virus	Potential host	Length (bp)	Viral protein ORF status	Detected in
Merkel cell polyomavirus	Vertebrate	3330	Partial	*A. mellifera*
		10809	Full	*C. calcarata*
Unknown iflavirus 1	Arthropods	1574	Partial	*A. mellifera*
		616	Partial	*C. calcarata*
Unknown iflavirus 2	Arthropods	7343	Full	*C. calcarata*
Unknown nepovirus 1	Plants	6501	Full	*A. mellifera*
Unknow partitivirus 1	Plants	1895	Full	*A. mellifera*
Unknown amalgavirus 1	Plants	3425	Full	*A. mellifera*
Unknown entomopoxvirus 1	Arthropods	2543	Partial	*C. calcarata*

Reference:

1. Ollerton J. Pollinator diversity: distribution, ecological function, and conservation. Annual Review of Ecology, Evolution, and Systematics. 2017;48:353-76. doi: https://doi.org/10.1146/annurev-ecolsys-110316-022919.

2. Katumo DM, Liang H, Ochola AC, Lv M, Wang Q-F, Yang C-F. Pollinator diversity benefits natural and agricultural ecosystems, environmental health, and human welfare. Plant Diversity. 2022;44(5):429-35. doi: https://doi.org/10.1016/j.pld.2022.01.005.

3. Ollerton J, Winfree R, Tarrant S. How many flowering plants are pollinated by animals? Oikos. 2011;120(3):321-6. doi: https://doi.org/10.1111/j.1600-0706.2010.18644.x.

4. Khalifa SAM, Elshafiey EH, Shetaia AA, El-Wahed AAA, Algethami AF, Musharraf SG, et al. Overview of bee pollination and its economic value for crop production. Insects. 2021;12(8):688. PubMed PMID: doi:10.3390/insects12080688.

5. Klein A-M, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, et al. Importance of pollinators in changing landscapes for world crops. Proceedings of the Royal Society B: Biological Sciences. 2007;274(1608):303-13. doi: doi:10.1098/rspb.2006.3721.

6. Breeze TD, Bailey AP, Balcombe KG, Potts SG. Pollination services in the UK: How important are honeybees? Agriculture, Ecosystems & Environment. 2011;142(3):137-43. doi: https://doi.org/10.1016/j.agee.2011.03.020.

7. Winfree R, Williams NM, Dushoff J, Kremen C. Native bees provide insurance against ongoing honey bee losses. Ecology Letters. 2007;10(11):1105-13. doi: https://doi.org/10.1111/j.1461-0248.2007.01110.x.

8. Brittain C, Kremen C, Klein A-M. Biodiversity buffers pollination from changes in environmental conditions. Global Change Biology. 2013;19(2):540-7. doi: https://doi.org/10.1111/gcb.12043.

9. Goulson D, Nicholls E, Botías C, Rotheray EL. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. Science. 2015;347(6229):1255957. doi: 10.1126/science.1255957.

10. Roulston TaH, Goodell K. The role of resources and risks in regulating wild bee populations. Annual Review of Entomology. 2010;56(1):293-312. doi: 10.1146/annurev-ento-120709-144802.

11. Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. Global pollinator declines: trends, impacts and drivers. Trends in Ecology & Evolution. 2010;25(6):345-53. doi: 10.1016/j.tree.2010.01.007.

12. Drossart M, Gérard M. Beyond the decline of wild bees: optimizing conservation measures and bringing together the actors. Insects. 2020;11(9):649. PubMed PMID:

doi:10.3390/insects11090649.

13.   Brown MJF, Dicks LV, Paxton RJ, Baldock KCR, Barron AB, Chauzat M-P, et al. A
horizon scan of future threats and opportunities for pollinators and pollination. PeerJ.
2016;4:e2249. doi: 10.7717/peerj.2249.

14.   Walker PJ, Siddell SG, Lefkowitz EJ, Mushegian AR, Adriaenssens EM,
Alfenas-Zerbini P, et al. Recent changes to virus taxonomy ratified by the
International Committee on Taxonomy of Viruses (2022). Archives of Virology.
2022;167(11):2429-40. doi: 10.1007/s00705-022-05516-5.

15.   Seitz K, Buczolich K, Dikunová A, Plevka P, Power K, Rümenapf T, et al. A
molecular clone of Chronic Bee Paralysis Virus (CBPV) causes mortality in honey bee
pupae (*Apis mellifera*). Scientific Reports. 2019;9(1):16274. doi:
10.1038/s41598-019-52822-1.

16.   Chen N-C, Wang C-H, Yoshimura M, Yeh Y-Q, Guan H-H, Chuankhayan P, et al.
Structures of honeybee-infecting Lake Sinai virus reveal domain functions and capsid
assembly with dynamic motions. Nature Communications. 2023;14(1):545. doi:
10.1038/s41467-023-36235-3.

17.   Cox-Foster DL, Conlan S, Holmes EC, Palacios G, Evans JD, Moran NA, et al. A
metagenomic survey of microbes in honey bee colony collapse disorder. Science.
2007;318(5848):283-7. doi: doi:10.1126/science.1146498.

18.   Tehel A, Brown MJF, Paxton RJ. Impact of managed honey bee viruses on wild
bees. Current Opinion in Virology. 2016;19:16-22. doi:
https://doi.org/10.1016/j.coviro.2016.06.006.

19. Dolezal AG, Hendrix SD, Scavo NA, Carrillo-Tripp J, Harris MA, Wheelock MJ, et al.
Honey bee viruses in wild bees: viral prevalence, loads, and experimental Inoculation.
PLOS ONE. 2016;11(11):e0166190. doi: 10.1371/journal.pone.0166190.

20.   Singh R, Levitt AL, Rajotte EG, Holmes EC, Ostiguy N, vanEngelsdorp D, et al. RNA
viruses in hymenopteran pollinators: evidence of inter-taxa virus transmission via
pollen and potential impact on non-apis hymenopteran species. PLOS ONE.
2010;5(12):e14357. doi: 10.1371/journal.pone.0014357.

21.   Zhang X, He SY, Evans JD, Pettis JS, Yin GF, Chen YP. New evidence that deformed
wing virus and black queen cell virus are multi-host pathogens. Journal of
Invertebrate Pathology. 2012;109(1):156-9. doi:
https://doi.org/10.1016/j.jip.2011.09.010.

22.   Susi H, Laine AL. Agricultural land use disrupts biodiversity mediation of virus
infections in wild plant populations. New Phytologist. 2021;230(6):2447-58.

23.   Lawson SP, Kennedy KB, Rehan SM. Pollen composition significantly impacts the
development and survival of the native small carpenter bee, *Ceratina calcarata*.
Ecological Entomology. 2021;46(2):232-9. doi: https://doi.org/10.1111/een.12955.

24.   Rehan SM, Richards MH. Nesting biology and subsociality in *Ceratina calcarata*

(Hymenoptera: Apidae). The Canadian Entomologist. 2010;142(1):65-74. Epub

2012/04/02. doi: 10.4039/n09-056.

25.   Rehan SM, Glastad KM, Lawson SP, Hunt BG. The genome and methylome of a

subsocial small carpenter bee, *Ceratina calcarata*. Genome Biology and Evolution.

2016;8(5):1401-10. doi: 10.1093/gbe/evw079 %J Genome Biology and Evolution.

26.   Lawson SP, Helmreich SL, Rehan SM. Effects of nutritional deprivation on

development and behavior in the subsocial bee *Ceratina calcarata* (Hymenoptera:

Xylocopinae). Journal of Experimental Biology. 2017;220(23):4456-62. doi:

10.1242/jeb.160531.

27.   Shell WA, Rehan SM. The price of insurance: costs and benefits of worker

production in a facultatively social bee. Behavioral Ecology. 2017;29(1):204-11. doi:

10.1093/beheco/arx146.

28.   Hartmann U, Forsgren E, Charrière J-D, Neumann P, Gauthier L. Dynamics of *Apis*

mellifera filamentous virus (AmFV) infections in honey bees and relationships with

other parasites. Viruses [Internet]. 2015; 7(5):[2654-67 pp.].

29.   Marzoli F, Forzan M, Bortolotti L, Pacini MI, Rodríguez-Flores MS, Felicioli A, et al.

Next generation sequencing study on RNA viruses of *Vespa velutina* and *Apis*

mellifera sharing the same foraging area. Transboundary and Emerging Diseases.

2021;68(4):2261-73. doi: https://doi.org/10.1111/tbed.13878.

30.   Roberts JMK, Anderson DL, Durr PA. Metagenomic analysis of Varroa-free

Australian honey bees (*Apis mellifera*) shows a diverse Picornavirales virome. Journal

of General Virology. 2018;99(6):818-26. doi: https://doi.org/10.1099/jgv.0.001073.

31.   Rowhani A, Daubert SD, Uyemoto JK, Al Rwahnih M, Fuchs M. American

Nepoviruses. In: Meng B, Martelli GP, Golino DA, Fuchs M, editors. Grapevine Viruses:

Molecular Biology, Diagnostics and Management. Cham: Springer International

Publishing; 2017. p. 109-26.

32.   Drummond FA, Lund J, Eitzer B. Honey bee health in Maine wild blueberry

production. Insects. 2021;12(6). doi: 10.3390/insects12060523.

33.   Šimenc L, Kuhar U, Jamnikar-Ciglenečki U, Toplak I. First complete genome of

Lake Sinai virus lineage 3 and genetic diversity of Lake Sinai virus strains from honey

bees and bumble bees. Journal of Economic Entomology. 2020;113(3):1055-61. doi:

10.1093/jee/toaa049.

34.   Daughenbaugh KF, Martin M, Brutscher LM, Cavigli I, Garcia E, Lavin M, et al.

Honey bee infecting Lake Sinai viruses. Viruses. 2015;7(6):3285-309. doi:

10.3390/v7062772.

35.   Abou Kubaa R, Molinatto G, Solaiman Khaled B, Daher-Hjaij N, Heinoun K,

Saponari M. First detection of black queen cell virus, *Varroa destructor* macula-like

virus, *Apis mellifera* filamentous virus and *Nosema ceranae* in Syrian honey bees *Apis mellifera syriaca*. Bulletin of Insectology. 2018;71(2):217–24.

36. Shi M, Lin X-D, Tian J-H, Chen L-J, Chen X, Li C-X, et al. Redefining the invertebrate RNA virosphere. Nature. 2016;540(7634):539-43. doi: 10.1038/nature20167.

37. Kwon M, Jung C, Kil E-J. Metagenomic analysis of viromes in honey bee colonies (*Apis mellifera*; Hymenoptera: Apidae) after mass disappearance in Korea. Frontiers in Cellular and Infection Microbiology. 2023;13. doi: 10.3389/fcimb.2023.1124596.

38. Nanetti A, Bortolotti L, Cilia G. Pathogens spillover from honey bees to other arthropods. Pathogens [Internet]. 2021; 10(8).

39. Fleites-Ayil FA, Medina-Medina LA, Quezada Euán JJG, Stolle E, Theodorou P, Tragust S, et al. Trouble in the tropics: Pathogen spillover is a threat for native stingless bees. Biological Conservation. 2023;284:110150. doi: https://doi.org/10.1016/j.biocon.2023.110150.

40. Santamaria J, Villalobos EM, Brettell LE, Nikaido S, Graham JR, Martin S. Evidence of Varroa-mediated deformed wing virus spillover in Hawaii. Journal for Parasitology: Parasites and Wildlife. 2018;151:126-30.

41. Coffman KA, Burke GR. Genomic analysis reveals an exogenous viral symbiont with dual functionality in parasitoid wasps and their hosts. PLOS Pathogens. 2020;16(11):e1009069. doi: 10.1371/journal.ppat.1009069.

42. Card SD, Pearson MN, Clover GRG. Plant pathogens transmitted by pollen. Australasian Plant Pathology. 2007;36(5):455-61. doi: 10.1071/AP07050.

43. Fetters AM, Ashman T-L. The pollen virome: A review of pollen-associated viruses and consequences for plants and their interactions with pollinators. American Journal of Botany. 2023;110(6):e16144. doi: https://doi.org/10.1002/ajb2.16144.

44. Li JL, Cornman RS, Evans Jay D, Pettis Jeffery S, Zhao Y, Murphy C, et al. Systemic spread and propagation of a plant-pathogenic virus in European honeybees, *Apis mellifera*. mBio. 2014;5(1):10.1128/mbio.00898-13. doi: 10.1128/mbio.00898-13.

45. Smadi M, Lee E, Phelan J, Wang A, Bilodeau GJ, Pernal SF, et al. Plant virus diversity in bee and pollen samples from apple (*Malus domestica*) and sweet cherry (*Prunus avium*) agroecosystems. Frontiers in Plant Science. 2024;15:1335281.

46. Fetters AM, Cantalupo PG, Wei N, Robles MTS, Stanley A, Stephens JD, et al. The pollen virome of wild plants and its association with variation in floral traits and land use. Nature Communications. 2022;13(1):523. doi: 10.1038/s41467-022-28143-9.

47. Lawson SP, Ciaccio KN, Rehan SM. Maternal manipulation of pollen provisions affects worker production in a small carpenter bee. Behavioral Ecology and Sociobiology. 2016;70(11):1891-900. doi: 10.1007/s00265-016-2194-z.

48. Tucker EM, Rehan SM. Wild Bee Community Assemblages Across Agricultural

Landscapes1. Journal of Agricultural and Urban Entomology. 2017;33(1):77-104.

49.   Tucker EM, Rehan SM. Farming for bees: annual variation in pollinator

populations across agricultural landscapes. Agricultural and Forest Entomology.

2018;20(4):541-8.

50.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014. doi: 10.1093/bioinformatics/btu170.

51.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature

Methods. 2012;9(4):357-9. doi: 10.1038/nmeth.1923

http://www.nature.com/nmeth/journal/v9/n4/abs/nmeth.1923.html#supplementar

y-information.

52.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al.

Full-length transcriptome assembly from RNA-Seq data without a reference genome.

Nature Biotechnology. 2011;29(7):644-52.

53.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile

metagenomic assembler. Genome Research. 2017;27(5):824-34. doi:

10.1101/gr.213959.116.

54.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular evolutionary genetics

analysis version 11. Molecular Biology and Evolution. 2021;38(7):3022-7. doi:

10.1093/molbev/msab120.

55.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30.

doi: 10.1093/bioinformatics/btt656.

56.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics.

2010;26(1):139-40. doi: 10.1093/bioinformatics/btp616.

57.   Chen Y, Lun AT, Smyth GK. From reads to genes to pathways: differential

expression analysis of RNA-Seq experiments using Rsubread and the edgeR

quasi-likelihood pipeline. F1000Research. 2016;5:1438. Epub 2016/08/11. doi:

10.12688/f1000research.8987.2. PubMed PMID: 27508061; PubMed Central PMCID:

PMCPMC4934518.

Khabir M, Hussain R, Khan HU, Hameed M, Anjum SI. Viral impacts on honey bee

populations: A review. Saudi J Biol Sci. 2021 Jan;28(1):523-530. doi:

10.1016/j.sjbs.2020.10.037.

Olgun T, Everhart SE, Anderson T, Wu-Smart J. Comparative analysis of viruses in four

bee species collected from agricultural, urban, and natural landscapes. PLoS One.

2020 Jun 12;15(6):e0234431. doi: 10.1371/journal.pone.0234431. PMID: 32530936;

PMCID: PMC7292363.

643