**Comparing virome composition between honey bees (*Apis mellifera* L.)**

**and small carpenter bees (*Ceratina calcarata* Robertson) in response to**

**various agricultural landscapes**

Zixiao Zhao[1]; M. Catherine Farrell[2]; Jayla Marvin[2] ; Sandra M. Rehan[3]; Hongmei

Li-Byarlay[1,2] [*]

[1]Agricultural Research Development Program, Central State University, Wilberforce,

OH, USA

[2]Department of Agricultural and Life Sciences, Central State Univesity, Wilberforce,

OH, USA

[3]Department of Biology, York University, Toronto, ON, Canada

**Author contact information**

ZZ: zxzhao5@gmail.com

https://orcid.org/0000-0002-3627-5503

MCF: mfarrell@centralstate.edu

https://orcid.org/0000-0002-0081-3285

JM: JaylaMarvin832@gmail.com

SMR: sandra.rehan@gmail.com

https://orcid.org/0000-0002-6441-5155

HL-B: Hli-byarlay@centralstate.edu

https://orcid.org/0000-0001-6579-5172

**\*Corresponding Author**

Hongmei Li-Byarlay

Hli-byarlay@centralstate.edu

**Abstract:**

Managed pollinators, such as honeybees (*Apis mellifera* L.) and wild pollinators, such as small carpenter bees (*Ceratina calcarata* Robertson), are important for food and agriculture in North America. Viral prevalence as a pathogenic stress and agricultural landscapes such as organic and conventional farms as foraging resources for pollinators are important for their health. However, it is unclear if the virus is distributed differently between managed and wild pollinators when they share similar foraging area resources. We investigated the virome composition of *Apis* and *Ceratina* collected from conventional, organic farms, or roadside (as a control without agricultural activity) using RNA-Seq analysis. The result demonstrates that honeybees carry a wide range of insect and plant viruses. The black queen cell virus is the only pathogenic virus discovered from both species in one conventional farm. Three novel insect viruses (1 entomopoxvirus and 2 iflaviruses) were detected in small carpenter bees, 1 novel iflavirus of which was identified in honey bee samples. The biological and ecological significance of novel viruses are yet to be studied. The results of our project shed light on the status of the biotic stress of viruses on pollinators.

# Keywords:

Honey bees, agricultural landscape, transcriptomics, bee virus, iflavirus, entomopoxvirus

## Introduction:

Animal pollinators have critical ecological functions in maintaining the diversity of global ecosystems [1, 2]. About 87.5% of flowering plants rely on animal pollinators to support reproduction [3]. In agriculture, approximately 75% of crop species require pollinators, contributing about 30-35% of total crop production yield [4, 5]. Bees (Hymenoptera: Apidae) are considered the most abundant pollinators. Honey bee species (*Apis* spp.), with a global distribution due to domestication, contribute about 80% of total pollination and create about $700 million value annually in the US [6]. However, the ecological significance of non-*Apis* wild bee species shouldn't be neglected. It is estimated over 17,000 wild bee species worldwide, with diverse social structures from eusocial to solitary. Wild bees provide pollination services to many crops, especially when honey bees are not available [7, 8]. For last several decades, both wild and honey bees are suffering from colony declines [9-12], which not only causes economical losses to apiary industry, but more importantly, reduces pollination service and imposes potential risks for global food security.

The causes for pollinator colony declines are complicated. Factors include habitat loss and fragmentation, increasing use of chemical pesticides and other agrochemcials, introductions of pests and pathogens, and climate changes [11, 13]. Among those stressors, pathogenic viruses are considered the direct and the most significant harrzard to both wild and honey bee populations. Currently identified bee pathogenic viruses belong to positive-sense single-stranded RNA (+ss RNA) virus category. The majority of the pathogenic bee viruses are from two families under the order Picornavirales: Iflaviridae and Dicistroviridae. Iflaviridae has only one genus: *Iflavirus*, which contains Sacbrood virus (SBV), Deformed wing virus (DWV), Varroa destructor virus-1 (VDV-1), and Slow bee paralysis virus (SBPV). While the family

Dicistroviridae contains Black queen cell virus (BQCV), Kashmir bee virus (KBV), Acute

bee paralysis virus (ABPV), and Israeli acute paralysis virus (IAPV) [14]. Two extra

pathogenic viruses of bees: Chronic bee paralysis virus (CBPV) is an unclassified virus

with two genomic RNA segments [15], and Lake Sinai viruses (LSVs) are from a

separate order Nodamuvirales [14], with 8 identified linages [16]. The symptoms

caused by pathogenic viruses vary, including body deformities, color changed,

uncapped cell with larvae or pupae, paralysis adults, which eventually increase

mortality and the degradation of the colony. IAPV infections have been discovered

association with colony collapse disorder (CCD), which responsible for large-scale

honey bee colony loss in the US [17].

Despite the focus on honey bees, the pathogenic viruses can also infect wild bee

species [18-20]. For example, DWV have been reported infecting a wide range of

hymenopteran insects, including some social and solitary bees and several non-bee

wasp species [18, 20, 21]. Dolezal et al. [19] reported DWV as well as SBV were

widely detected in wild bees collected in Iowa, while IAPV, LSV and BQCV were also

detected in wild be species. Furthermore, DWV, BQCV and SBV can be deposited to

pollen and remain infective, which can contribute to inter-taxa transmission among

bees, wild bees, and wasps [20]. These studies suggested virus can be a potential

factor responsible for colony declines in wild bee species.

Agriculture are considered one of the major reason for colony declines. It have

converted natural ecosystems to farmland primary for food and fiber productions.

This process is accompanied by loss of biodiversity and use of agrochemicals causing

habitat loss, fragmentation and degradation. Agriculture can also alter host-pathogen

interactions at multiple levels. Study in the plants demonstrated the revalence of

plant viruses is negatively correlated with the biodiversity in agricultural landscape

[22]. However, whether agricultural activities affect viral prevalence in insects,

especially in wild and honey bees, is unknown.

The small carpenter bee, *Ceratina calcarata* Robertson, is a wild bee species

native to eastern part of North America. It is considered as indicator species and

pollinators for several native and agricultural plants [23]. Due to the feature of

subsociality, *C. calcarata* is used for studying pollinator ecology, behavior, evolution

and nutrition [24-27]. Little is known about the virology of this species. The impact of

viruses to *C. calcarata* populations and whether the viral prevalence is related to

agricultural activities are not clear.

Here we presented our metagenomic studies on viruses associated with western

honey bee (*A. mellifera* L.) and *C. calcarata* in response to different agricultural

intensification. Bees were collected from three landscape types representing levels of

agricultural intensification: conventional farming, organic farming and roadside with

no agricultural activities. The viral RNA was extracted from collected samples and

subjected to high-throughput sequencing. We analyzed known and novel viruses

associated with both pollinators, which are highly specific to location and pollinator

species. Our study provided new methodology and unique knowledge for

pollinator-virus interactions.

## Results:

*Reconstruct virome from viral RNA-Seq reads*

The RNA-Seq of extracted viral RNA generates approximately 17 to 36 million

raw read pairs for each *A. mellifera* and *C. calcarata* sample. After quality trimming

and partially removing reads from respective host genome (Amel HAv3.1

PRJNA471592 and Ccalc.v3 PRJNA791561), 5 to 25 million pairs were kept for each

sample (**Supplementary file S1**). Trinity *de nove* assembled 366,289 cotings for *A.*

*mellifera* and 203,317 contigs for *C. calcarata* virome, which consists of 188,433 and

91,314 non-redundant clusters. The assembled contigs were subject to BLASTN

searches against non-redundant database (nr), which claimed 541 clusters and 1187

contigs from *A. mellifera* assembly poteintially belong to viruses. In contrast, 62

clusters and 84 contigs from *C. calcarata* assembly are potentially from viruses. A

large portion of assembled contigs of are from either host genes or the plants they

have visited.

*Viruses infecting bees and other insects detected in A. mellifera and C. calcarata*

*virome assemblies*

BLASTN search identified four known bee pathogenic viruses from *A. mellifera*

virome: BQCV, DWV, SBV, and LSV3 with fully assembled genome sequences. Two

other members of the LSV group: LSV1 and LSV2 were detected. But their sequences

were only partially assembled. Positive and negative strands were detected in all of

the above viruses except LSV2, from which only positive strand is detected. We also

detected *Apis mellifera* filamentous virus (AmFV) [28] and Hubei partiti-like virus 34

[29] in *A. mellifera*. The AmFV were partially assembled and the Hubei partiti-like

virus 34 demonstrated a fully assembled genome (**Table 1A**). In *C. calcarata*, BQCV

was the only detected bee pathogenic virus and appeared as partial-assembled as

negative-strand contig (**Table 1B**).

*Plant viruses discovered in A. mellifera and C. calcarata virome assemblies*

A variety of plant viruses were detected in virome assemblies of both species. In

*A. mellifera,* white clover cryptic virus 1 (WCCV1) and white clover cryptic virus 2

(WCCV2) from genus *Alphapartitivirus* (family: Partitiviridae, +ssRNA), as well as

white clover mosaic virus (WCMV) genus *Potexvirus* (family: Alphaflexiviridae,

+ssRNA) were fully assembled. Blackberry chlorotic ringspot virus (BCRV) from genus

*Ilavirus* (family: Bromoviridae, +ssRNA) and *Raphanus sativus* cryptic virus 1 (RsPV1)

from Partitiviridae (+ssRNA) have one genomic segment fully assembled, while other

segments were either partially assembled or completely missing. Partially assembled

plant viruses include 7 +ssRNA viruses: peanut stunt virus (PSV) and tobacco ringspot

virus (TRSV) from Bromoviridae, clover yellow mosaic virus (ClYMV) from

Alphaflexiviridae, red clover nepovirus A (RCNVA) , tobacco streak virus (TSV), and

tomato ringspot virus (ToRSV) from genus *Nepovirus* (family: Secoviridae), red clover

vein mosaic virus (RCVMV) from Betaflexiviridae, and one -ssRNA virus: *pinus flexilis*

virus 1 (PiFleV1) from Rhabdoviridae (**Table 1A**). The plant viruses in *C. calcarata*

virome are much less diversed. WCCV-2 is the only fully-assembled plant viruses,

while WCCV-1, BCRV, ToRSV, and WClMV were partially assembled (**Table 1B**).

*Novel viruses in A. mellifera and C. calcarata virome assemblies*

The genus *Iflavirus* (family: Iflaviridae) consists of +ssRNA viruses infecting

arthropods, including three bee pathogenic viruses SBV, DWV, and SBPV. Two novel

Iflavirus, referred to as unknown iflavirus 1 and unknown inflavirus 2, were identified

in *C. calcarata* virome assembly as positive strands. The phylogenetic analysis using

translated polyprotein amino acid sequence suggests both viruses are in the same

group with SBV. The unknown iflavirus-1 has the highest identity with Darwin bee

virus-2, an unclassified picornavirus infecting honey bee [30] (**Figure 1A**). This virus is

also detected in *A. mellifera* virome assembly as a negative strand, suggesting that it

can infect and replicate in honey bees.

In *C. calcarata*, we detected a novel poxvirus (double-strand DNA virus). Its

DNA-dependent RNA polymerase RPO132 is partially assembled. Phylogenetic

analysis showed this virus is closely related to the Yalta virus, an uncategorized

poxvirus from *Drosophila melanogaster* and *Diachasmimorpha* entomopoxvirus

(DIEPV), from parasitoid wasp *Diachasmimorpha longicaudata*. The three viruses

form a sister cluster to other entomopoxvirus (genus: *Alphaentomopoxvirus*,

*Betaentomopoxvirus*, and *Deltaentomopoxvirus*), while all the vertebrate poxvirus

(genus: *Avipoxvirus*, *Capripoxvirus*, and *Parapoxvirus*) are clustered into a separate

group. The result suggests this novel virus belongs to the subfamily

Entomopoxvirinae (EPV) that primarily infect insects (**Figure 1E**).

Three additional novel plant viruses were detected in the *A. mellifera* virome,

including one from the genus *Nepovirus* (unknown nepovirus 1, family: Secoviridae),

one from family Partitiviridae (unknown parititivirus 1, family: Partitiviridae), and one

of genus *Amalgavirus* (unknown Amalgavirus 1, family: Amalgaviridae). Nepovirus

has two positive-strand genomic RNAs [31]. For unknown nepovirus 1, the ORF of

polyprotein RNA2 was fully assembled. However, the correspondent RNA1 encoding

viral coat protein was not detected. The phylogeny of polyprotein sequence shows it

is most close related to Blueberry leaf mottle virus and Grapevine Bulgarian latent

virus, forming a separate cluster (**Figure 1B**). Similarly, only RNA1 encoding

RNA-dependent RNA polymerase (RDRP) were fully assembled for another potential

plant virus: unknown parititivirus 1. The phylogenetic analysis of RDRP showed it is

clustered with several uncategorized partitivirus viruses associated with plant:

*Polygonatum kingianum* cryptic virus 1, Panax cryptic virus 3, and Dichroa partitivirus

1. This cluster is sister group to plant-infecting genus *Alphapartitivirus* (**Figure 1D**).

The genome of unknown amalgavirus 1 was fully assembled with two complete ORF:

the ORF1+2 fusion protein responsible for genome replication, and a coat protein.

The phylogeny of fusion proteins showed the it is the most closed to Vicia cryptic

virus M (VCV-M) and Medicago sativa amalgavirus 1 (MsAV1) and have a clear

separation with *Zygosaccharomuces bailii* virus Z (ZbV-Z), a yeast-infecting virus

belongs to the genus *Zybavirus* (**Figure 1C**).

In addition, other viruses, including phages and a human virus: Merkel cell

polymavirus (Family: Polyomaviridae) were identified in both virome assemblies. The

viral genome consists of circular dsRNA, which resulted in two identical full-length

viral sequences were assembled into one contig in *C. calcarata* virome. In *A. mellifera*,

only partial sequence was assembled. How the Merkel cell polymavirus introduced

into virome and their effects in both species are unknown.

*Quantification of viruses in response to different agricultural landscape*

The distributions of the above detected viruses are highly site-specific. In *A.*

*mellifera*, among all identified insect viruses, five known bee viruses: BQCV, DWV,

LSVs, SBV, and AmFV were detected from all of the experimental sites regardless

of the type of agricultural use, with AmFV only marginally detected in conventional

farm C1. In contrast, the novel unknown ilfavirus 1 was only detected in samples

from organic farm O2 (**Figure 2A**). For five known bee viruses, the edgeR did not

show any significantly differences at virus level. While, a separate 2834-bp LSV3

segment, containing ORFs of two viral proteins, were significantly increased in

organic samples comparing to roadside samples (FDR: 0.04). The Hubei partiti-like

virus was also identified on all sites except conventional farm C1 (**Figure 2A**). Its

full-length sequence significantly increased in organic farms comparing to both

conventional and roadside samples (FDR: 0.006 to conventional, 0.003 to roadside).

The presence and viral levels of plant viruses are highly depended to the

location. For instance, TSV and PiFleV1 were only detected respectively on one

roadside site, RCNVA and TRSV were only found respectively on one conventional

and one organic site. RCVMV were identified across on three conventional sites, but

in relatively low level. WCIMV, WCCV2, and ToRSV were expressed on all

experimental sites (**Figure 2B**). The WCCV-1 is the only significant plant virus

detected by edgeR. The levels of both genome segments decreased significantly in

organic sites, comparing to both conventional (FDR, RNA1: 0.011, RNA2: 0.006) and

roadside (FDR, RNA1: 0.006, RNA2: 0.003). The viral levels of BRCV at road sites were

either very low or undetected, but the overall level is too low to carry out statistical

analysis in edgeR.

*C. calcarata* hosts much less viruses. The presence of insect viruses was high

site-specific. BQCV and unknown entomopoxvirus 1 were only found respectively in

one conventional site, where two novel ilfavirus were detected in roadside sites and

conventional farm C1. None of the viruses were detected from samples collected

from both organic sites, conventional farm C3, and roadside R2 (**Figure 2C**).

Regarding to the plant viruses, WCCV1 and WCCV2 have a similar pattern, being

detected only on conventional farms, while WCIMV and ToRSV were detected in

some roadside samples as well (**Figure 2D**). However, their levels were not claimed

significant. Similarly, none of the plant viruses were detected from samples collected

organic sites.

**Discussion**

The metagenomic approach based on viral RNA-Seq was adopted for identifying

and quantifying known viruses, as well as discovering novel viruses. The presented

study surveyed the abundance of viruses associated with two pollinator species. The

virome compositions are site-specific. However, the *A. mellifera* carries a wider range

of both insect and plant viruses than *C. calcarata*. The results also indicate that the *A.*

*mellifera* may transmit plant diseases more effectively than *C. calcarata*.

*The Honey bee viruses in A. mellifera and spillover to C. calcarata*

The levels of BQCV, DWV and SBV are high in *A. mellifera* virome at all

experimental sites, suggested the high prevalence of above viruses. In contrast, none

of the acute bee paralysis virus complex (ABPV, KBV and IAPV) were detected. This

result is consistent with other studies in the Northeast of the US [20, 32]. LSV is a

highly-variable multi-strain bee virus that do not have any significant symptom, but

likely cause week colonies. The LSV contigs assembled in this study are mostly closely

related to LSV1, LSV2 and LSV3. The LSV3 were detected from all experimental sites

and the only LSV with full-length genome sequence assembled. The LSV3 were

originally identified Slovenia [33] and has been detected in the Western US [34]. The

study confirmed LSV3 has been spread to the New England area.

Two known viruses associated with *A. mellifera* were also detected: AmFV and

Hubei partiti-like virus 34. The AmFV is a weak pathogen in *A. mellifera* and rarely

causes significant symptoms. Its infection has been reported related to BQCV and

DWV [28, 35]. We detected AmFV from all sites, while its viral loads were from

extremely low at conventional farm C1 to medium-high at conventional farm C3. The

Hubei partiti-like virus 34 was first discovered as an arthropod RNA virus [36] and

was reported associated with hymenopteran species, including *A. mellifera* and Asian

hornet *Vespa velutina* in East Asia and South Europe [29, 37]. The current study

reported the low to moderate infection of Hubei partiti-like virus 34 from all except

conventional farm C3 site. The potential impact of this virus to honey bee health is

still unknown.

The spillover of honey bee pathogen to wild bees has been considered as a

threaten to native pollinator populations [38, 39]. In genus *Ceratina*, DWV was found

spillover to two species *C. dupla* [20] and *C. smaragdula* [40]. The current study

showed given the high prevalence of both BQCV and DWV in *A. mellifera*, BQCV can

spilled over to *C. calcarata*, while no DWV spill over was observed. BQCV is only

detected from conventional farm C2 in relatively low prevalence and viral load,

indicating that this spillover to *C. calcarata* is not a common scenario.

*Potential viruses infecting A. mellifera and C. calcarata*

The primary hosts for iflaviruses were arthropods. We discovered two closely

related novel ifla-like viruses. Unknown iflavirus 1 were detected in both *A. mellifera*

and *C. calcarata*. The unknown iflavirus 2 were only detected in *C. calcarata*. The

prevalence of both viruses was low, however, the viral load of unknown ilfavirus 2

were slightly higher of *C. calcarata* samples collected from conventional farm C1 and

roadside site 1, comparing to unknown ilfavirus 1 of the samples.

The EPVs are dsDNA viruses infecting insects and predominantly insect

pathogens. The unknown entomopoxivirus 1 detected in *C. calcarata* are more

closely related to Yalta viruses and DIEPV than other entomopathogenic viruses.

Although the role of Yalta virus is still unknown, the DIEPV has been found replicate

in both parasitoid wasp *D. longicaudata* and its host fruit fly *Anastrepha suspensa*.

The replication of virus is highly virulent to the host fruit fly and no negative impact

to wasp host, suggesting a mutualistic symbiotic relationship of DIEPV to parastoid

wasp [41]. Although both *C. calcarata* and *D. longicaudata* are from order

Hymenoptera, no such parasitoid relationship was observed in *C. calcarata*. Whether

this novel entomopoxivirus primary infects *C. calcarata* or just an overspill event to *C.*

*calcarata* yet to be studied.

*Pollinators carrying plant viruses*

Many plant viruses can travel within or attach to pollen grains [42, 43], including

BRCV, TSV, PSV, WCCV1, WCCV2, TRSV, ToRSV, and WClMV from presented. However,

only TSV, TRSV, and ToRSV have been proven vertically and horizontally transmitted by the pollens. The rest were only detected by RNA-Seq [43]. Pollinator such as *A. mellifera* posing potential risks. However none of the detected viruses were on the list of pollinator-transmitting virus, according to Fetters et al [43]. TRSV was reported infect honey bee [44], however whether cross-kingdom infection assist transmitting diseases is unknown. On the other hand, pollinators are a powerful tool for monitoring plant diseases [45]. The plant viruses detected in *A. mellifera* resembles to the results of other metagenomic studies of bees and pollen at family and order levels [45, 46], while the species composition is determined by plant and prevalence of plant viruses at the location.

Comparing to *A. mellifera*, plant viruses in *C. calcarata* are much less diverse. Previous studies have shown pollens from clover and rose are major food recourses [47], and white clover pollen significantly increases the fitness of *C. calcarata* [23]. In consist with that, among four plant viruses detected in *C. calcarata* virome, WCCV1, WCCV2, WClMV primarily infect white clover (*Trifolium repens*), and the host range of ToRSV include rose and red clover (*T. pratense*). None of the viruses beyond *C. calcarata's* food range were detected. This suggested *C. calcarata* is better for monitoring white clover pathogens in a smaller area.

## Material and method:

*Sample collection*

Both honey bee and small carpenter bee samples were collected from sites representing three different types of land use: conventional farm, organic farm and roadside. Honey bees and the small carpenter bees were collected by sweep nets from May to July, 2018 in Strafford County, New Hampshire. We collected both

species within three conventional farms and three organic farms. The roadside

samples were collected in the urbanized area within the University of New

Hampshire campus. The detailed location and GPS Coordinates are described in

previous publication [48, 49]. Bee samples were flash-frozen in liquid nitrogen and

temporarily stored at -80 ℃.

*RNA extraction, sequencing and data cleaning*

The viral RNA was extracted from *A. mellifera* and *C. calcarata* individually by

Quick-RNA viral kit (ZYMO RESEARCH, Irvine, CA). DNase I (ZYMO RESEARCH)

in-column digestions were applied to remove genomic and environmental DNA. For

mRNA library generation, 200ng of total RNA was treated with NEBNext Poly mRNA

Magnetic Isolation Module E7490L (New England Biolabs, USA) following the

manufacturer protocol. Subsequently, the isolated mRNA was fragmented for 10

minutes. cDNA was synthesized and amplified for 12 PCR cycles using NEBNext Ultra

II Directional (stranded) RNA Library Prep Kit E7760L for Illumina (New England

Biolabs, USA) with NEBNext Multilex oligos indexes kit E6440S following the

manufacturer directions. Distributions of the template length and adapter-dimer

contamination were assessed using an Agilent 2100 Bioanalyzer and High Sensitivity

DNA kit (Agilent Technologies, Inc). The concentration of cDNA libraries was

determined using Invitrogen Qubit dsDNA HS reagents and read on a Qubit

Fluorometer (Thermo Fisher), and the cDNA libraries were paired-end 150bp format

sequenced on a Novaseq SP system (Illumina, San Diego, CA). In total, 17 honey bee

samples and 16 *C. calcarata* samples were sequenced (**Supplementary File S1**). Raw

reads were subject to adapter and quality trimming by Trimmomatic [50]. To remove

endogenous transcripts from bee host, the reads were respectively aligned to honey

bee and small carpenter bee [25] genomes (including mitochondrial sequences)

respectively using Bowtie2 [51]. Read pairs concordantly aligned to bee genomes

were considered as endogenous transcripts and subject to removal.

*Virome assembly and viral identification*

To reconstruct the virome of *A. mellifera* or *C. calcarata,* we first merged all the

pre-cleaned reads from the same organism to increase the sequence coverage. Then

two *de novo* assembler: Trinity [52] and metaSpade [53] were used with default

setting. The results are comparable but the total runtime for Trinity is lower. We

decided to adopt Trinity for following proceeding analysis.

With the pre-cleaning process the assembled virome still contains host

sequences. To identify virus sequences, we use BLASTN to search assembled contigs

to default nr/nt database with E-value cutoff of 1e-7. Contigs with first three top hits

to viral sequence from database will be retained as viral contigs. To further virus

identification, we exam the blast results of contigs >500 nt. A contig will be assigned

as known virus if it aligned stringently (E-value = 0) to existing virus. If the alignment

is less stringent (0<E-value<1e-7), it will be assigned as novel virus. The "full-length"

or "partial" labels are based on the completeness and intactness of viral protein of

each viral contig. For potential novel viral contigs, ORF finder and BLASTP were used

to translate and predict viral protein. MEGA 11 [54] was used to construct

Maximum-likelihood phylogenetic tree based on the sequence of homologous

proteins of known and novel viruses.

*Quantification of virus and differential analysis between landscapes*

To quantify the abundance of each contigs, the clean reads were mapped to

virome assembly of respective species using Bowtie 2 [51]. Multiple alignment is

allowed to obtain maximum sensitivity. The read counts were obtained by

featureCount [55] and were normalized and converted to count per million reads

(CPM) by R edgeR package [56]. We also use edgeR to conduct differential analysis

between different landscape type. Prior to that, each contig, including endogenous

and viral, was subject to the filter based on its expression level [57]. Only contigs

with enough high levels were retained in statistical test. The contigs with FDR less

than 0.05 were claim significantly difference between landscapes. The viral contigs,

defined by viral identification step, were focused.

## Acknowledgements:

We are grateful for the help received from Stephanie Gardner, Katherine Odanaka,

and Wyatt Shell with collecting bee samples. We thank the Genomic Shared Resource

Shared Resource at The Ohio State University Comprehensive Cancer Center,

Columbus, OH for generating the virus RNA-seq data reported in this

publication. Research reported in this publication was supported by USDA Evans

Allen Fund NI231445XXXXG004, NI241445XXXXG004 for HL-B and ZZ, USDA CBG

GRANT13202344 for HL-B, USDA award 2020-67014-31557 for MCF, and the NSF

Louis Stokes Alliances for Minority Participation (LSAMP) program Ohio Alliance

scholarship for JM, the Ohio State University Comprehensive Cancer Center and the

National Institutes of Health under grant number P30 CA016058.

## Author contributions:

HL-B and SMR designed the experiment. SMR conducted the field studies and

provided the samples. MCF and JM prepared samples for RNA-seq. ZZ conducted

data analysis and wrote the first draft of the manuscript. All authors reviewed the

final manuscript.

**Figures and Tables:**

**Figure 1.** Phylogenetic trees of novel viruses discovered from A. mellifera and C. *calcarata* virome

assemblies by Maximum-likelihood analysis of sequences of viral proteins: **(A)** polyprotein (iflavirus),

**(B)** polyprotein from RNA2 (neprovirus), **(C)** RNA-dependent RNA polymerase (amalgavirus), **(D)**

RNA-dependent RNA polymerase from RNA1 (partitivirus), and **(E)** RNA-dependent RNA polymerase

submit RPO132 (poxvirus). The names in bold correspond to honeybee-associated viruses.

Abbreviations and protein accessions of known viruses are included in **Supplementary File S2**.

**Figure 2.** Heatmaps demonstrating the relative abundance of **(A)** insect viruses in *A. mellifera*, **(B)**

plant viruses in *A. mellifera*, **(C)** insect viruses in *C. calcarata*, and **(D)** plant viruses in *C. calcarata* at

each sampling site. Average normalized count-per-million reads (cpm) from RNA-Seq was calculated

for each virus of each site and used in the heatmap. White color indicates the virus is undetected on

that site. Potential insect and plant viruses were included.

**Table 1**. List of identified insect and plant viruses in **(A)** *A. mellifera* and **(B)** *C. calcarata* virome

assemblies, assembled from RNA-Seq data

**A**

| Virus name | Abbr. | Type | Family | Genus | Host | Assemble status | Strand |
|---|---|---|---|---|---|---|---|
| Black queen cell virus | BQCV | (+)ssRNA | Dicistroviridae | *Triatovirus* | Bees | Full | +/- |
| Deformed wing virus | DWV | (+)ssRNA | Iflaviridae | *Iflavirus* | Bees | Full | +/- |
| Sacbrood virus | SBV | (+)ssRNA | Iflaviridae | *Iflavirus* | Bees | Full | +/- |
| Lake Sinai virus 3 | LSV3 | (+)ssRNA | Sinhaliviridae | *Sinaivirus* | Bees | Full | +/- |
| Lake Sinai virus 1 | LSV1 | (+)ssRNA | Sinhaliviridae | *Sinaivirus* | Bees | Partial | +/- |
| Lake Sinai virus 2 | LSV2 | (+)ssRNA | Sinhaliviridae | *Sinaivirus* | Bees | Partial | + |
| Apis mellifera filamentous virus | AmFV | dsDNA | *unclassified* | *unclassified* | Bees | Partial | +/- |
| Hubei partiti-like virus 34 | - | RNA | *unclassified* | *unclassified* | Bees | Full | + |
| Blackberry chlorotic ringspot virus | BCRV | (+)ssRNA | Bromoviridae | *Ilarvirus* | Plants | RNA1: Full | + |
| | | | | | | RNA2: Partial | - |
| | | | | | | RNA3: Partial | + |
| Tobacco streak virus | TSV | (+)ssRNA | Bromoviridae | *Ilarvirus* | Plants | RNA1: Partial | +/- |
| | | | | | | RNA2: Partial | +/- |
| | | | | | | RNA3: Partial | + |
| Peanut stunt virus | PSV | (+)ssRNA | Bromoviridae | *Cucumovirus* | Plants | RNA1: Partial | + |
| | | | | | | RNA2: Partial | + |
| | | | | | | RNA3: Partial | - |
| Raphanus sativus cryptic virus 1 | RasCV1 | dsRNA | Partitiviridae | *unclassified* | Plants | RNA1: Full | - |
| | | | | | | *RNA2: Missing* | |
| White clover cryptic virus 1 | WCCV1 | dsRNA | Partitiviridae | *Alphapartitivirus* | Plants | RNA1: Full | - |
| | | | | | | RNA2: Full | - |

| Virus name | Abbr. | Type | Family | Genus | Host | Assembly status | Strand |
|---|---|---|---|---|---|---|---|
| White clover cryptic virus 2 | WCCV2 | dsRNA | Partitiviridae | *Alphapartitivirus* | Plants | RNA1: Full | - |
| | | | | | | RNA2: Full | - |
| Red clover nepovirus A | RCNVA | (+)ssRNA | Secoviridae | *Nepovirus* | Plants | RNA1: Partial | + |
| | | | | | | RNA2: Partial | +/- |
| Tobacco ringspot virus | TRSV | (+)ssRNA | Secoviridae | *Nepovirus* | Plants | RNA1: Partial | +/- |
| | | | | | | RNA2: Partial | - |
| Tomato ringspot virus | ToRSV | (+)ssRNA | Secoviridae | *Nepovirus* | Plants | RNA1: Partial | +/- |
| | | | | | | RNA2: Partial | - |
| White clover mosaic virus | WClMV | (+)ssRNA | Alphaflexiviridae | *Potexvirus* | Plants | Full | +/- |
| Clover yellow mosaic virus | ClYMV | (+)ssRNA | Alphaflexiviridae | *Potexvirus* | Plants | Partial | +/- |
| Red clover vein mosaic virus | RCVMV | (+)ssRNA | Betaflexiviridae | *Carlavirus* | Plants | Partial | +/- |
| Pinus flexilis virus 1 | PiFleV1 | (-)ssRNA | Rhabdoviridae | *unclassified* | Plants | Partial | +/- |

428    **B**

| Virus name | Abbr. | Type | Family | Genus | Host | Assembly status | Strand |
|---|---|---|---|---|---|---|---|
| Black queen cell virus | BQCV | (+)ssRNA | Dicistroviridae | *Triatovirus* | Bees | Partial | - |
| Tomato ringspot virus | ToRSV | (+)ssRNA | Secoviridae | *Nepovirus* | Plants | RNA1: Partial | +/- |
| | | | | | | RNA2: Partial | +/- |
| White clover cryptic virus 1 | WCCV-1 | dsRNA | Partitiviridae | *Alphapartitivirus* | Plants | RNA1: Full | - |
| | | | | | | *RNA2: Missing* | |
| White clover cryptic virus 2 | WCCV-2 | dsRNA | Partitiviridae | *Alphapartitivirus* | Plants | RNA1: Full | + |
| | | | | | | RNA2: Full | - |
| White clover mosaic virus | WClMV | (+)ssRNA | Alphaflexiviridae | *Potexvirus* | Plants | Partial | +/- |

431    **Table 2.** List of novel viruses detected in *A. mellifera* and *C. calcarata* virome assemblies assembled

432    from RNA-Seq data

| Virus | Potential host | Length (bp) | Viral protein ORF status | Detected in |
|---|---|---|---|---|
| Merkel cell polyomavirus | Vertebrate | 3330 | Partial | *A. mellifera* |
| | | 10809 | Full | *C. calcarata* |
| Unknown iflavirus 1 | Arthropods | 1574 | Partial | *A. mellifera* |
| | | 616 | Partial | *C. calcarata* |
| Unknown iflavirus 2 | Arthropods | 7343 | Full | *C. calcarata* |
| Unknown nepovirus 1 | Plants | 6501 | Full | *A. mellifera* |
| Unknow partitivirus 1 | Plants | 1895 | Full | *A. mellifera* |
| Unknown amalgavirus 1 | Plants | 3425 | Full | *A. mellifera* |
| Unknown entomopoxvirus 1 | Arthropods | 2543 | Partial | *C. calcarata* |

**Reference:**

1.    Ollerton J. Pollinator diversity: distribution, ecological function, and
conservation. Annual Review of Ecology, Evolution, and Systematics. 2017;48:353-76.
doi: https://doi.org/10.1146/annurev-ecolsys-110316-022919.
2.    Katumo DM, Liang H, Ochola AC, Lv M, Wang Q-F, Yang C-F. Pollinator diversity
benefits natural and agricultural ecosystems, environmental health, and human
welfare. Plant Diversity. 2022;44(5):429-35. doi:
https://doi.org/10.1016/j.pld.2022.01.005.
3.    Ollerton J, Winfree R, Tarrant S. How many flowering plants are pollinated by
animals? Oikos. 2011;120(3):321-6. doi:
https://doi.org/10.1111/j.1600-0706.2010.18644.x.
4.    Khalifa SAM, Elshafiey EH, Shetaia AA, El-Wahed AAA, Algethami AF, Musharraf
SG, et al. Overview of bee pollination and its economic value for crop production.
Insects. 2021;12(8):688. PubMed PMID: doi:10.3390/insects12080688.
5.    Klein A-M, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C,
et al. Importance of pollinators in changing landscapes for world crops. Proceedings
of the Royal Society B: Biological Sciences. 2007;274(1608):303-13. doi:
doi:10.1098/rspb.2006.3721.
6.    Breeze TD, Bailey AP, Balcombe KG, Potts SG. Pollination services in the UK: How
important are honeybees? Agriculture, Ecosystems & Environment.
2011;142(3):137-43. doi: https://doi.org/10.1016/j.agee.2011.03.020.
7.    Winfree R, Williams NM, Dushoff J, Kremen C. Native bees provide insurance
against ongoing honey bee losses. Ecology Letters. 2007;10(11):1105-13. doi:
https://doi.org/10.1111/j.1461-0248.2007.01110.x.
8.    Brittain C, Kremen C, Klein A-M. Biodiversity buffers pollination from changes in
environmental conditions. Global Change Biology. 2013;19(2):540-7. doi:
https://doi.org/10.1111/gcb.12043.
9.    Goulson D, Nicholls E, Botías C, Rotheray EL. Bee declines driven by combined
stress from parasites, pesticides, and lack of flowers. Science.
2015;347(6229):1255957. doi: 10.1126/science.1255957.
10.    Roulston TaH, Goodell K. The role of resources and risks in regulating wild bee
populations. Annual Review of Entomology. 2010;56(1):293-312. doi:
10.1146/annurev-ento-120709-144802.
11.    Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. Global
pollinator declines: trends, impacts and drivers. Trends in Ecology & Evolution.
2010;25(6):345-53. doi: 10.1016/j.tree.2010.01.007.
12.    Drossart M, Gérard M. Beyond the decline of wild bees: optimizing conservation
measures and bringing together the actors. Insects. 2020;11(9):649. PubMed PMID:

doi:10.3390/insects11090649.

13.   Brown MJF, Dicks LV, Paxton RJ, Baldock KCR, Barron AB, Chauzat M-P, et al. A

horizon scan of future threats and opportunities for pollinators and pollination. PeerJ.

2016;4:e2249. doi: 10.7717/peerj.2249.

14.   Walker PJ, Siddell SG, Lefkowitz EJ, Mushegian AR, Adriaenssens EM,

Alfenas-Zerbini P, et al. Recent changes to virus taxonomy ratified by the

International Committee on Taxonomy of Viruses (2022). Archives of Virology.

2022;167(11):2429-40. doi: 10.1007/s00705-022-05516-5.

15.   Seitz K, Buczolich K, Dikunová A, Plevka P, Power K, Rümenapf T, et al. A

molecular clone of Chronic Bee Paralysis Virus (CBPV) causes mortality in honey bee

pupae (*Apis mellifera*). Scientific Reports. 2019;9(1):16274. doi:

10.1038/s41598-019-52822-1.

16.   Chen N-C, Wang C-H, Yoshimura M, Yeh Y-Q, Guan H-H, Chuankhayan P, et al.

Structures of honeybee-infecting Lake Sinai virus reveal domain functions and capsid

assembly with dynamic motions. Nature Communications. 2023;14(1):545. doi:

10.1038/s41467-023-36235-3.

17.   Cox-Foster DL, Conlan S, Holmes EC, Palacios G, Evans JD, Moran NA, et al. A

metagenomic survey of microbes in honey bee colony collapse disorder. Science.

2007;318(5848):283-7. doi: doi:10.1126/science.1146498.

18.   Tehel A, Brown MJF, Paxton RJ. Impact of managed honey bee viruses on wild

bees. Current Opinion in Virology. 2016;19:16-22. doi:

https://doi.org/10.1016/j.coviro.2016.06.006.

19.   Dolezal AG, Hendrix SD, Scavo NA, Carrillo-Tripp J, Harris MA, Wheelock MJ, et al.

Honey bee viruses in wild bees: viral prevalence, loads, and experimental Inoculation.

PLOS ONE. 2016;11(11):e0166190. doi: 10.1371/journal.pone.0166190.

20.   Singh R, Levitt AL, Rajotte EG, Holmes EC, Ostiguy N, vanEngelsdorp D, et al. RNA

viruses in hymenopteran pollinators: evidence of inter-taxa virus transmission via

pollen and potential impact on non-apis hymenopteran species. PLOS ONE.

2010;5(12):e14357. doi: 10.1371/journal.pone.0014357.

21.   Zhang X, He SY, Evans JD, Pettis JS, Yin GF, Chen YP. New evidence that deformed

wing virus and black queen cell virus are multi-host pathogens. Journal of

Invertebrate Pathology. 2012;109(1):156-9. doi:

https://doi.org/10.1016/j.jip.2011.09.010.

22.   Susi H, Laine AL. Agricultural land use disrupts biodiversity mediation of virus

infections in wild plant populations. New Phytologist. 2021;230(6):2447-58.

23.   Lawson SP, Kennedy KB, Rehan SM. Pollen composition significantly impacts the

development and survival of the native small carpenter bee, *Ceratina calcarata*.

Ecological Entomology. 2021;46(2):232-9. doi: https://doi.org/10.1111/een.12955.

24.  Rehan SM, Richards MH. Nesting biology and subsociality in *Ceratina calcarata*
(Hymenoptera: Apidae). The Canadian Entomologist. 2010;142(1):65-74. Epub
2012/04/02. doi: 10.4039/n09-056.

25.  Rehan SM, Glastad KM, Lawson SP, Hunt BG. The genome and methylome of a
subsocial small carpenter bee, *Ceratina calcarata*. Genome Biology and Evolution.
2016;8(5):1401-10. doi: 10.1093/gbe/evw079 %J Genome Biology and Evolution.

26.  Lawson SP, Helmreich SL, Rehan SM. Effects of nutritional deprivation on
development and behavior in the subsocial bee *Ceratina calcarata* (Hymenoptera:
Xylocopinae). Journal of Experimental Biology. 2017;220(23):4456-62. doi:
10.1242/jeb.160531.

27.  Shell WA, Rehan SM. The price of insurance: costs and benefits of worker
production in a facultatively social bee. Behavioral Ecology. 2017;29(1):204-11. doi:
10.1093/beheco/arx146.

28.  Hartmann U, Forsgren E, Charrière J-D, Neumann P, Gauthier L. Dynamics of *Apis*
*mellifera* filamentous virus (AmFV) infections in honey bees and relationships with
other parasites. Viruses [Internet]. 2015; 7(5):[2654-67 pp.].

29.  Marzoli F, Forzan M, Bortolotti L, Pacini MI, Rodríguez-Flores MS, Felicioli A, et al.
Next generation sequencing study on RNA viruses of *Vespa velutina* and *Apis*
*mellifera* sharing the same foraging area. Transboundary and Emerging Diseases.
2021;68(4):2261-73. doi: https://doi.org/10.1111/tbed.13878.

30.  Roberts JMK, Anderson DL, Durr PA. Metagenomic analysis of Varroa-free
Australian honey bees (*Apis mellifera*) shows a diverse Picornavirales virome. Journal
of General Virology. 2018;99(6):818-26. doi: https://doi.org/10.1099/jgv.0.001073.

31.  Rowhani A, Daubert SD, Uyemoto JK, Al Rwahnih M, Fuchs M. American
Nepoviruses. In: Meng B, Martelli GP, Golino DA, Fuchs M, editors. Grapevine Viruses:
Molecular Biology, Diagnostics and Management. Cham: Springer International
Publishing; 2017. p. 109-26.

32.  Drummond FA, Lund J, Eitzer B. Honey bee health in Maine wild blueberry
production. Insects. 2021;12(6). doi: 10.3390/insects12060523.

33.  Šimenc L, Kuhar U, Jamnikar-Ciglenečki U, Toplak I. First complete genome of
Lake Sinai virus lineage 3 and genetic diversity of Lake Sinai virus strains from honey
bees and bumble bees. Journal of Economic Entomology. 2020;113(3):1055-61. doi:
10.1093/jee/toaa049.

34.  Daughenbaugh KF, Martin M, Brutscher LM, Cavigli I, Garcia E, Lavin M, et al.
Honey bee infecting Lake Sinai viruses. Viruses. 2015;7(6):3285-309. doi:
10.3390/v7062772.

35.  Abou Kubaa R, Molinatto G, Solaiman Khaled B, Daher-Hjaij N, Heinoun K,
Saponari M. First detection of black queen cell virus, *Varroa destructor* macula-like

virus, *Apis mellifera* filamentous virus and *Nosema ceranae* in Syrian honey bees *Apis mellifera syriaca*. Bulletin of Insectology. 2018;71(2):217–24.

36. Shi M, Lin X-D, Tian J-H, Chen L-J, Chen X, Li C-X, et al. Redefining the invertebrate RNA virosphere. Nature. 2016;540(7634):539-43. doi: 10.1038/nature20167.

37. Kwon M, Jung C, Kil E-J. Metagenomic analysis of viromes in honey bee colonies (*Apis mellifera*; Hymenoptera: Apidae) after mass disappearance in Korea. Frontiers in Cellular and Infection Microbiology. 2023;13. doi: 10.3389/fcimb.2023.1124596.

38. Nanetti A, Bortolotti L, Cilia G. Pathogens spillover from honey bees to other arthropods. Pathogens [Internet]. 2021; 10(8).

39. Fleites-Ayil FA, Medina-Medina LA, Quezada Euán JJG, Stolle E, Theodorou P, Tragust S, et al. Trouble in the tropics: Pathogen spillover is a threat for native stingless bees. Biological Conservation. 2023;284:110150. doi: https://doi.org/10.1016/j.biocon.2023.110150.

40. Santamaria J, Villalobos EM, Brettell LE, Nikaido S, Graham JR, Martin S. Evidence of Varroa-mediated deformed wing virus spillover in Hawaii. Journal for Parasitology: Parasites and Wildlife. 2018;151:126-30.

41. Coffman KA, Burke GR. Genomic analysis reveals an exogenous viral symbiont with dual functionality in parasitoid wasps and their hosts. PLOS Pathogens. 2020;16(11):e1009069. doi: 10.1371/journal.ppat.1009069.

42. Card SD, Pearson MN, Clover GRG. Plant pathogens transmitted by pollen. Australasian Plant Pathology. 2007;36(5):455-61. doi: 10.1071/AP07050.

43. Fetters AM, Ashman T-L. The pollen virome: A review of pollen-associated viruses and consequences for plants and their interactions with pollinators. American Journal of Botany. 2023;110(6):e16144. doi: https://doi.org/10.1002/ajb2.16144.

44. Li JL, Cornman RS, Evans Jay D, Pettis Jeffery S, Zhao Y, Murphy C, et al. Systemic spread and propagation of a plant-pathogenic virus in European honeybees, *Apis mellifera*. mBio. 2014;5(1):10.1128/mbio.00898-13. doi: 10.1128/mbio.00898-13.

45. Smadi M, Lee E, Phelan J, Wang A, Bilodeau GJ, Pernal SF, et al. Plant virus diversity in bee and pollen samples from apple (*Malus domestica*) and sweet cherry (*Prunus avium*) agroecosystems. Frontiers in Plant Science. 2024;15:1335281.

46. Fetters AM, Cantalupo PG, Wei N, Robles MTS, Stanley A, Stephens JD, et al. The pollen virome of wild plants and its association with variation in floral traits and land use. Nature Communications. 2022;13(1):523. doi: 10.1038/s41467-022-28143-9.

47. Lawson SP, Ciaccio KN, Rehan SM. Maternal manipulation of pollen provisions affects worker production in a small carpenter bee. Behavioral Ecology and Sociobiology. 2016;70(11):1891-900. doi: 10.1007/s00265-016-2194-z.

48. Tucker EM, Rehan SM. Wild Bee Community Assemblages Across Agricultural

Landscapes1. Journal of Agricultural and Urban Entomology. 2017;33(1):77-104.

49.   Tucker EM, Rehan SM. Farming for bees: annual variation in pollinator

populations across agricultural landscapes. Agricultural and Forest Entomology.

2018;20(4):541-8.

50.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014. doi: 10.1093/bioinformatics/btu170.

51.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature

Methods. 2012;9(4):357-9. doi: 10.1038/nmeth.1923

http://www.nature.com/nmeth/journal/v9/n4/abs/nmeth.1923.html#supplementar

y-information.

52.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al.

Full-length transcriptome assembly from RNA-Seq data without a reference genome.

Nature Biotechnology. 2011;29(7):644-52.

53.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile

metagenomic assembler. Genome Research. 2017;27(5):824-34. doi:

10.1101/gr.213959.116.

54.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular evolutionary genetics

analysis version 11. Molecular Biology and Evolution. 2021;38(7):3022-7. doi:

10.1093/molbev/msab120.

55.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30.

doi: 10.1093/bioinformatics/btt656.

56.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics.

2010;26(1):139-40. doi: 10.1093/bioinformatics/btp616.

57.   Chen Y, Lun AT, Smyth GK. From reads to genes to pathways: differential

expression analysis of RNA-Seq experiments using Rsubread and the edgeR

quasi-likelihood pipeline. F1000Research. 2016;5:1438. Epub 2016/08/11. doi:

10.12688/f1000research.8987.2. PubMed PMID: 27508061; PubMed Central PMCID:

PMCPMC4934518.

618