A molecular circuit regulates fate plasticity in emerging and adult AT2 cells

Amitoj S Sawhney[1#], Brian J Deskin[2#], Junming Cai[1], Daniel Gibbard[1], Gibran Ali[1], Annika

Utoft[1], Aaron Olson[1], Hannah Hausman[1], Liberty Sabol[1], Shannon Holmberg[1], Ria Shah[1],

Rachel Warren[1], Stijn De Langhe[1], Rui Benedito[5], Zintis Inde[2,3], Kristopher A Sarosiek[2,3], Evan

Lemire[2], Adam L Haber[2], Liu Wang[4], Zong Wei[4], and Douglas G Brownfield[1,2+]*

[1]Division of Pulmonary and Critical Care Medicine, Department of Medicine, Mayo Clinic

College of Medicine and Science, Rochester, MN 55905, USA

[2] Molecular and Integrative Physiological Sciences Program, Harvard T.H. Chan School of

Public Health, Boston, MA, USA

[3]John B. Little Center for Radiation Sciences, Harvard T.H. Chan School of Public Health,

Boston, MA, USA

[4]Department of Physiology and Biomedical Engineering, Mayo Clinic College of Medicine and

Science, Scottsdale, AZ 85259, USA.

[5]Molecular Genetics of Angiogenesis Group, Centro Nacional de Investigaciones

Cardiovasculares (CNIC), Madrid E28029, Spain

21
22 # These authors contributed equally to this work.
23 [+] Current address: Division of Pulmonary and Critical Care Medicine, Departments of
24 Physiology and Biomedical Engineering and of Biochemistry and Molecular Biology, Mayo
25 Clinic College of Medicine and Science, Rochester, MN 55905
26
* Corresponding author. Email: brownfield.douglas@mayo.edu

Abstract

Alveolar AT1 and AT2 cells are vital for lung gas exchange and become compromised in several

diseases. While key differentiation signals are known, their emergence and fate plasticity are

unclear. Here we show in the embryonic lung that single AT2s emerge at intermediate zones,

extrude, and connect with nearby epithelium via interlumenal junctioning. We observe that AT2s

retain fate plasticity until the bZIP transcription factor C/EBPα suppresses Notch signaling at a

novel *Dlk1* enhancer. Both *Dlk1* and *Cebpa* are regulated by the polycomb repressive complex

(PRC2), which together form a "pulse generator" circuit that times *Dlk1* expression and thus

Notch activation, resulting in a "salt and pepper" pattern of AT1 and AT2 fate. In injured adult

lungs, C/EBPα downregulation is required to re-access AT2 fate plasticity and is mediated by the

dominant negative C/EBP family member CHOP. Finally, *Cebpa* loss also activates a "defender"

AT2 state, distinct from its reparative state, and we propose AT2s toggle between either state

following infection to protect and repair alveoli.

Introduction

Alveoli are the sites of gas exchange in the lung that are lined by two distinct epithelial cell types. The flat surface of alveolar type 1 (AT1) cells facilitates gas exchange while cuboidal alveolar type 2 (AT2) cells secrete surfactant to prevent alveolar collapse. Defective alveoli underly several maladies—bronchopulmonary dysplasia (BPD) in infants, chronic obstructive pulmonary disease (COPD)/emphysema[1], pulmonary fibrosis[2], lung adenocarcinoma[3, 4], and respiratory infections like Severe Acute Respiratory Syndrome (SARS)[5, 6] and COVID-19[7]. Deciphering how these crucial alveolar cell types emerge as well as the mechanisms that regulate their fate is imperative for developing therapies that will not only slow or stop disease progression but may even promote lung repair.

As alveologenesis begins, AT1 and AT2 cells arise from a common pool of distal progenitors (DP)[4, 8, 9, 10, 11], with differentiation detected as early as embryonic day 16.5 (e16.5) in mice, save for one study reporting an earlier timepoint[12]. Upon differentiation, AT1 and AT2 cells are arranged in a salt and pepper pattern[13] along a lumenal surface which, as alveoli mature, interconnect with lumens of anatomically distinct branches to facilitate gas conduction between alveoli[14]. After development, mature AT2 cells can act as antigen presenting cells in an immune response[15, 16] (partly directed interferon signaling[17, 18]), as well as facultative stem cells following injury[4, 11, 19, 20 21] (directed by Wnt[22] and Notch[23] signaling). While both AT2 states are critical after infection for alveolar recovery, a recent study found interferon stimulation suppresses repair of the lung epithelium following infection[24, 25], suggesting that signals driving these states may oppose one another at the cellular level.

Here we show nascent AT2 cells are first detected molecularly at e15.5 as singletons at stereotyped intermediate regions of distal branches. Following basal extrusion, we observe

nascent AT2s can form *de novo* attachments to nearby but anatomically separate epithelia, a step

that is likely critical in stitching together alveolar lumen and that we term "interlumenal

junctioning", followed by sporadically patterned AT2 maturation, distinct from the "proximal-to-

distal wave" pattern known to initiate differentiation of AT1s and AT2s earlier[13].

We determine that nascent AT2s retain fate plasticity into the first perinatal week,

revising the lineage model to include a "window of fate plasticity" spanning from distal

progenitors to the AT2 lineage. From screening transcriptional regulators around this window,

we identify a critical negative regulator of AT2 fate plasticity, the bZIP transcription factor

C/EBPα, that when lost results in AT2-to-AT1 conversion. Previously C/EBPα has been reported

to play a role in epithelial homeostasis[26] and is implicated in fate regulation in multiple tissues,

including the blood and liver[27]. Further, we found that the observed AT2-to-AT1 differentiation

was mediated in a non-cell autonomous manner by the Notch regulator DLK1, a cell-surface

transmembrane protein[28] that has been associated with brain and muscle development, stem cell

maintenance, lung cancer, and abnormal tissue repair[23, 29]. Observing a pulsed expression of *Dlk1*

during alveologenesis, we find *Dlk1* and *Cebpa* are both regulated by the polycomb repressive

complex (PRC2), and determine these components together constitute an incoherent feed

forward loop. Upon abrogation of this "pulsed generator" circuit, we observe a disruption of

AT1/AT2 differentiation. PRC2 is a well-established transcriptional repressor that has been

shown to regulate DLK1 expression in neuronal fate selection[30] and recent evidence suggests it

also modulates C/EBPα expression in the context of adipocyte differentiation[31].

Following injury in the adult lung C/EBPα must be downregulated in AT2s for

conversion to AT1. We identify that the dominant negative member of the C/EBP family,

CHOP, is induced in AT2s following injury and its upregulation promotes AT2-to-AT1

differentiation. Finally, we observe *Cebpa* loss also activates a "defensive" AT2 state that is

enriched in antipathogen response genes, likely regulated by interferon signaling, and is mutually

exclusive from the AT1-associated DATP state[32].

Results

Timing and pattern of AT2 cell emergence

To delineate newly forming AT2 cells from distal progenitors and their mature counterparts, we

analyzed scRNAseq data of the distal epithelial lineage from a prior study sampling the

embryonic, perinatal, and adult timepoints of mouse lung development[33] (Fig. 1a). Clustering

analysis of the distal epithelium identified both DP and AT1 populations, as well as stratified

AT2s into two groups (Fig. 1b) that are also largely separate in time (Fig. 1c). While these AT2

groups positively correlate in gene expression (Pearson coefficient = 0.6, Suppl. Fig. 1a), their

distinction as a separate cluster was confirmed by co-assignment probability (Suppl. Fig. 1b).

Taken together with the temporal pattern, we refer to the earlier cluster as the nascent AT2

(nAT2) state, and the later as the mature AT2 (mAT2) state. From this distinction, we were able

to identify hundreds of genes unique to either DP, nAT2, or mAT2 cluster (Fig. 1d, Suppl. Fig.

1c, and Suppl. Table 2), and initially focused on two for protein-level validation: *Retnla*

expressed by nAT2s and *Cd74* by mAT2s (Fig. 1e, Suppl. Fig. 1d). *Retnla* encodes a regulator of

inflammation, resistin-like molecule (RELM) α, while CD74 is critical in MHC-II antigen

processing. In contrast to commonly used markers that label both DP and AT2 lineage starting

around e17.5 (such as SFTPC)[4, 10], RELMα uniquely labels nAT2s at this stage and can be used

to distinguish them from both DPs and AT1s, as can be seen at an e17.5 transition zone where

the mAT2 marker CD74 is not highly expressed (Fig. 1f, Suppl. Fig. 1e). Further, flow

cytometry confirmed at e18.5 that a substantial proportion of marker-expressing AT2s are

RELMα[+] CD74[−] (Suppl. Fig. 1g), which shifts to RELMα[−] CD74[+] in adult AT2s (Suppl. Fig. 1h).

115   Given the distinct labeling of nAT2s by RELMα in the embryonic lung, we used it to

identify when and where AT2 cells first emerge during alveologenesis. Careful immunostaining

revealed nAT2s emerge as early as e15.5 (Fig. 1g), which steadily increases over time (Fig. 1h).

At first emergence, SOX9[low]/RELMα[+] and nAT2s are singletons (Suppl. Fig. 1e, f) with retained

columnar morphology (Fig. 1i) positioned at anatomical intermediate zones, located between

airway smooth muscle (ASM)-covered distal branch stalks and terminal end buds (Fig. 1g,

asterisks). After their initial emergence, nAT2s next arise (~e17.5) within distal stalks following

ASM to myofibroblast remodeling as previously reported[34], then finally at terminal end buds

soon after birth[4]. These observations support a sequential, three zone model of AT2

differentiation (Fig. 1j, left, Suppl. Fig. 1d, e) that integrates previously conflicting findings that

reported AT2 differentiation first occurring either at terminal end buds[12] or in a proximal-to-

distal pattern[4, 9].

Nascent AT2s junction with nearby but anatomically distinct epithelium

While imaging RELMα at these timepoints, we also observed nAT2s forming connections to

lumens across anatomically separate branches (Fig. 2a), a behavior which increased in frequency

over time (Fig. 2b). While it is known that during alveolar maturation AT2 cells are capable of

simultaneously integrating into lumens of multiple alveoli[14, 34], the process by which it occurs

has yet to be described. Careful immunostaining of junctioning nAT2s revealed their negativity

for AT1 markers (PDPN, RAGE, and HOPX) and positivity for AT2 markers (SFTPC, LAMP1,

and MUC1) that is often higher than non-junctioned AT2s, suggesting they may be further along

in differentiation (Suppl. Fig. 2a, b). Using high resolution imaging and 3D reconstruction, we

observed that junctions occurred either directly with the second lumen, or indirectly by

attachment to another extruded nAT2 (Fig. 2c, Suppl. Fig. 2c, Suppl. Video 1-4).

139   To further investigate the dynamics of nAT2 junctioning, we performed timelapse

confocal microscopy on Precision Cut Lung Slices (PCLS) isolated from e17.5 murine lungs

wherein the distal lung epithelium was labeled with a membrane tethered GFP ($Tg^{Nkx2.1-Cre}$

$Rosa26^{mTmG}$) wherein nAT2s could be readily visualized to extrude and junction (Fig. 2d). We

observed both types of nAT2 junctioning - direct when the interstitial distance was short (Fig.

2e) and indirect when greater than a cell length away (Fig. 2f, Suppl. Fig. 2d, Suppl. Video 5).

Finally, we observed for most, if not all junctioning nAT2s, a significant reduction in length of

MUC1 domains indicative of apical constriction (Fig. 2g, h). Taken together, we believe our

observations are of an initial "interlumenal junctioning" event critical in the process of

integrating alveolar lumenal surfaces (Fig. 2i, j) that relies upon basal extrusion of nAT2s to

junction. Further, we hypothesize the manner of junctioning (direct or indirect) is dependent on

interstitial distance, and that defective nAT2 junctioning could result in alveolar simplification

similar to what is observed in BPD and COPD.

Nascent and mature AT2 state differ in fate plasticity and BH3 regulation.

Following their initial specification, nAT2s undergo maturation in the first week or so after birth,

which can be observed by the switch from RELMα to CD74 expression (Fig. 3a, b). As RELMα

expression is lost in mAT2s, we observe its concomitant upregulation in the airway epithelium

(Suppl. Fig. 3a). In contrast to the initial proximal-to-distal wave pattern that AT1/AT2 fate

selection is reported to occur[4, 13], we observe AT2 maturation does not follow this pattern but

rather occurs in a sporadic salt-and-pepper pattern (Fig. 3c, d, Suppl. Fig. 3b), suggesting

maturation may be governed by a distinct mechanism.

Given the molecular distinction between nAT2 and mAT2 states, we next sought to

determine whether functional differences also exist. One manner in which time-dependent cell

states can vary is in fate plasticity, as has been observed in neutrophils[35] in vivo and lung

epithelial iPSCs[36] in culture. To determine whether and how nAT2s and mAT2s differ in fate

plasticity, we analyzed two previously generated scRNAseq datasets that sample timepoints

either during[12] (Fig. 3e - g, Suppl. Fig. 3c - g) or far beyond[37] (Fig. 3h - j) alveolar epithelial

differentiation. Analyzing first the distal epithelium prior to (e15.5) and during (e17.5) alveolar

epithelial differentiation, we performed standard clustering as well as UMAP visualization and

found that distal epithelial cells isolated from these timepoints only rarely co-clustered (Suppl.

Fig. 3e), indicating no e15.5 progenitor population had significantly adopted either AT1 or AT2

transcriptional program, a central tenet of the early lineage specification model proposed in a

prior study[12] . A weak correlation between one of the e15.5 progenitor clusters (DE3) and the

e17.5 progenitor cluster (0.42) was observed (Suppl. Fig. 3e, yellow box), suggesting a shared

transcriptional program.

Single-cell transcriptional dynamics analysis was next analyzed to determine when and

how the AT1 and nAT2s clusters segregate. Performing pseudotime (Suppl. Fig. 3f), velocity

(Suppl. Fig. 3g), as well as integrated PAGA-Velocity analysis (Fig. 3e) all identified e15.5

clusters as the starting timepoint, as well as identified the AT1 cluster as the predicted endpoint

(Suppl. Fig. 3f, g). Further, velocity analysis found a substantial proportion of nAT2s to be on a

transcriptional trajectory towards the AT1 cluster, suggesting fate plasticity was retained (Fig.

3e, red arrow). To determine whether a subset of nAT2s displayed a biased trajectory towards

AT1 differentiation, we established an AT1 and AT2 score based on well-established marker

genes[9] and observed a clear stratification within the nAT2 cluster (Fig. 3f). Separating the nAT2

cluster by this ratio into two subgroups (High AT1 or Low AT1 gene expression, > 0.5 = High

AT1/AT2 score), we compared velocity trajectories and observed a clear separation (Fig. 3g),

indicating nAT2 subset with a high AT1/AT2 ratio are leaving the nAT2 state and transitioning

to AT1.

After observing that embryonic nAT2s possibly retain fate plasticity, we sought to

determine how long AT2s retain fate plasticity after birth. Using a previously reported dataset

that samples across the lifetime of the distal epithelium of the mouse lung[37], we performed

similar processing as before to remove duplicates and non-distal epithelial cells before

performing UMAP visualization and clustering analysis (Fig. 3h, i). Analyzing more widely over

time, we again observed earlier progenitor populations, mature AT1 and AT2 clusters, as well as

a cluster transitioning between them as previously reported[37]. Velocity analysis of this

intermediate cluster found the cellular trajectories are oriented away from AT2 and towards AT1

(Fig. 3i, red box) like our prior observation. Finally, analysis of the timepoints of cells within the

AT2→AT1 transitioning clusters revealed most were from the early perinatal period, suggesting

fate plasticity was lost as AT2s mature within the first week after birth (Fig. 3j). To confirm this,

we performed alveolosphere differentiation assays following our previously described protocol[10]

and found that both nAT2s as well as early mAT2s (PN3) could give rise to AT1 cells (Fig. 3k,

l), confirming fate plasticity was lost primarily in a time-dependent manner, independent of

nAT2 or mAT2 state. Taken together, our analysis indicates that fate plasticity, a property of

DPs, is retained in newly formed AT2 cells for a period of 1-2 weeks after birth, revising the

model of the alveolar epithelial fate selection model to include a window of plasticity (Fig. 3m).

Another property known to vary from early life to maturity is mitochondrial sensitivity to

apoptotic stimuli (apoptotic priming)[38], which is reduced during maturation in several organs but

can be retained in certain stem cell populations[39, 40]. To determine whether apoptotic sensitivity

is retained in nAT2s, BH3 profiling was performed on isolated DPs, nAT2s, as well as adult

mAT2s. As expected, DPs and nAT2s exhibited heightened sensitivity to pro-apoptotic BH3

peptides as indicated by higher rates of cytochrome c release, while mAT2s were largely

unresponsive (Suppl. Fig. 3h). Taken together, our findings indicate the nAT2 state retains both

fate plasticity as well as sensitivity to BH3 regulated apoptosis despite being functionally

differentiated.

C/EBPα is required to maintain but not select AT2 fate

Given the loss of fate plasticity as nAT2s mature into mAT2s, we next sought to determine how

this paradigm is regulated. Referencing the scRNAseq data we previously used to distinguish

nAT2 and mAT2 clusters (Fig. 1a, b) as well as AT2s sequenced more deeply by the Tabula

Muris Consortium[41], we screened putative transcriptional and epigenetic regulators from a

curated list (1,678 genes) and identified 67 genes with ≥ 2-fold enrichment along the AT2

(Suppl. Table 3) lineage (DP→nAT2→mAT2) compared to AT1 (Fig. 4a). Of these, we then

considered only the 32 genes with known embryonic lethal or lung defects when knocked out in

mice (Fig. 4a, Suppl. Fig. 4a). These genes can be stratified into four categories based on

expression pattern over time: constitutive, DP-restrictive, DP/nAT2-restrictive, and

nAT2/mAT2-restrictive (Fig. 4a, lower and Suppl. Fig. 4a). Of these, we focused on genes

expressed in the observed window of fate plasticity (Suppl. Fig. 4b), which narrowed the list to

four genes, *Etv5* and *Elf5* (constitutive) as well as, *Cebpa* and *Nupr1* (nAT2/mAT2-Restrictive)

(Fig. 4b). While both constitutively expressed genes (*Etv5* and *Elf5*) have been investigated in

the lung, findings suggest regulatory roles other than fate plasticity, as ETV5 is required

primarily to maintain AT2 survival[42] while ELF5 appears to broadly regulate aspects of lung

epithelial differentiation[43]. Further, we excluded study of *Nupr1* given its primary role is in death

regulation, not differentiation[44, 45]. This left *Cebpa* as the putative regulator, for which we

validated the timing of expression via immunostaining in the embryonic lung (Fig. 4c).

Previous *Cebpa* research in the developing mouse lung performed knockouts via a Tet-

On approach and concluded it was required for alveolar epithelial differentiation[46, 47]. These

findings are conflicting with the expression pattern of *Cebpa*, which is first expressed after AT2

differentiation, not before (Fig. 4b, c, Suppl. Fig. 6b). Given this discrepancy, as well as a

subsequent study that questioned the viability of the prior Tet-On approach in the lung[48], we first

sought to confirm whether *Cebpa* was required for alveolar epithelial differentiation via a more

refined method of gene deletion. Using a lineage labeling with efficient recombination approach

(*Nkx2.1-Cre*; *Rosa26*$^{mTmG/mTmG}$; *Cebpa*$^{fl/fl}$) wherein distal bipotent progenitors are both

fluorescently labeled and floxed gene recombined[10], we tested whether *Cebpa* loss indeed

resulted in suppressed differentiation. Surprisingly, we observed that both AT2 and AT1 fate

selection occurred despite the absence of *Cebpa* (Fig. 4d, Suppl. Fig. 4c), both at the molecular

and morphological level.

As we expected *Cebpa* to play a role later in development consistent to when it is

expressed, we next performed sparsely labeling with efficient deletion experiment wherein

Cebpa was deleted in a subset of AT2 cells starting in the early perinatal period (~PN5, *LyzM*^{Cre};

Rosa26$^{mTmG/mTmG}$; *Cebpa*$^{fl/fl}$). While lineage-traced control AT2 cells maintain their fate into

adulthood, a significant portion of AT2 cells lacking *Cebpa* converted into AT1 cells (Fig. 4e, f),

suggesting an increase in AT2 fate plasticity. Finally, we confirmed the result using an AT2-

restrictive tamoxifen-inducible Cre recombinase line ($Sftpc^{CreER}$) which again demonstrated an

increase in conversion to AT1 fate both morphologically and molecularly (Suppl. Fig. 4d). Taken

together, we determined that C/EBPα, a bZIP transcription factor expressed following AT2 cell

specification is required during the perinatal period to maintain, not select, AT2 fate, suggesting

it plays a role in suppressing fate plasticity during the transition from nascent to mature AT2

state.

C/EBPα represses a non-cell autonomous AT1 differentiation program driven by DLK1

To determine more precisely how C/EBPα regulates AT2 fate plasticity, we conducted

scRNAseq of the alveolar epithelium from both control ($Cebpa^{wt/fl}$) and AT2-floxed ($Cebpa^{fl/fl}$)

lungs, wherein Cre-mediated deletion was induced in the perinatal period using $Sftpc^{CreER}$ in

combination with the Tg(iSuRe-Cre)[49] allele. Retrospective identification of cell types via

Louvain clustering using Seurat was able to distinguish control (AT2 – $Cebpa^{wt/fl}$) and $Cebpa$

floxed (AT2 – $Cebpa^{fl/fl}$) AT2s as well as AT1s (Fig. 5a-i, ii), a subset of which we confirmed

were derived from AT2 cells following Cre induction by expression of the Cre-inducible reporter

Tg(iSuRe-Cre) (Suppl. Fig. 5a), confirming our prior lineage tracing experiments. We observed a

substantial down- and upregulation of 117 and 116 genes respectively upon loss of $Cebpa$

(Suppl. Fig. 5b). To determine how AT2 or AT1 gene expression overall during development

(Suppl. Fig. 5c) and was impacted following $Cebpa$ loss, we first used gene scores to broadly

assess changes in either AT2 or AT1 related genes (Suppl. Fig. 5e). While no significant

upregulation of AT1 genes was observed (suggesting C/EBPα likely does not repress fate

plasticity by directly repressing the AT1 gene program), a reduction in overall AT2 genes was

observed in AT2s lacking *Cebpa* (Fig. 5a-ii). To further determine how AT2-associated genes

were altered, we separately assessed genes restrictively enriched in either nAT2 or mAT2 state

(Suppl. Fig. 5d). Upon *Cebpa* loss, AT2 cells reduce expression of mAT2 genes while

concomitantly upregulating nAT2 gene expression (Fig. 5a-ii), suggesting re-accessing of a

plastic, nascent-like AT2 state.

To better understand the behavior of these plastic AT2s as well as focus our investigation

of their molecular regulators, we next sought to determine whether the AT2-to-AT1

differentiation observed upon *Cebpa* loss was mediated by a cell autonomous or non-

autonomous mechanism. While a common experimental approach in *Drosophila* research[50, 51] to

delineate between mechanisms targeting internal signaling or proliferation (cell autonomous)

versus expression of an extracellular signal (non-cell autonomous), such experiments are less

common in the mouse and have been difficult using inducible Cre lines until the development

transgenic alleles such as Tg(iSuRe-Cre) [49]. Usage of a CreER line alone under sparse labeling

conditions can result in incomplete or mosaic recombination of alleles and label "escaper

cells"[52], confounding interpretations of the gene's necessity. To overcome this, We next

performed sparse labeling with efficient *Cebpa* deletion using the Tg(iSuRe-Cre) allele

(*Sftpc^{CreER}; Tg(iSuRe-Cre); Cebpa^{fl/fl}*) at a tamoxifen dose we determine labels ~20% of AT2s

validated by immunostaining (Fig. 5b, c, Suppl. Fig. 5f, g). This method of sparse labeling with

efficient recombination can be used to delineate whether a phenotype is regulated in a cell

autonomous (i.e. via a cell intrinsic mechanism, in which case the phenotype would occur in both

sparse and non-sparse labeling experiments) and non-cell autonomous programs (i.e. via a cell

extrinsic mechanism, in which case the phenotype would occur significantly less if at all in the

sparse labeling experiment). In contrast to when *Cebpa* is removed from most AT2s (~99%), we

observed that removal of *Cebpa* from sparse numbers of AT2s (~20%) resulted in a substantial

reduction of AT2-to-AT1 differentiation as assessed by percentage of lineage labeled AT1s and

proliferation (Fig. 5b, Suppl. Fig. 5h). This finding strongly indicates that AT2 fate plasticity acts

via a cell non-autonomous mechanism, likely an extracellular signal, to promote AT2-to-AT1

differentiation.

Given this, we focused our screen on genes upregulated in *Cebpa* deleted AT2s that

encode proteins with an extracellular localization (GO:0005615), which returned 22 genes, 6 of

which were also expressed in the nAT2s from the prior scRNAseq timecourse, and of these 4 are

localized on the cell membrane or secreted (*Dlk1*, *Col23a1*, *Cd14*, and *Apoc1*) (Fig. 5d, Suppl.

Fig. 5i, j). The most highly upregulated of these genes is *Dlk1*, which encodes a single pass type

I membrane protein localized to the outer cell surface that is known to regulate Notch signaling[53]

and AT2-to-AT1 differentiation[23] (Fig. 5d, red).

Finally, we confirmed DLK1 is also upregulated at the protein level following *Cebpa* loss

in culture (Suppl. Fig. 5k), and further observe its expression is maintained under sparse labeling

with efficient *Cebpa* deletion in vivo, suggesting C/EBPα directly regulates DLK1 in a cell

autonomous manner (Fig. 5e, f). Taken together, we hypothesize that C/EBPα begins to block

fate plasticity of specified AT2s by direct suppression of *Dlk1* and its subsequent activation of

Notch signaling.

After establishing C/EBPα likely directly represses *Dlk1* expression, we sought to

determine where within the genome it is mediated. We performed bulk ATAC-seq on both

control and *Cebpa* floxed AT2s in the same experiment timeline as for scRNAseq (Suppl. Fig.

5l), but AT2-lineage cells were further purified by FACS using the genetically encoded

fluorophore tdTomato. After confirming correlation between ATACseq replicates (Suppl. Fig.

5m), we identified 23,764 sites within the genome that significantly changed in accessibility

following *Cebpa* loss (Suppl. Fig. 5n), with decreased accessibility associated with 3,986 genes,

consistent with C/EBPα's well established role as a positive regulator of gene expression[54]. We

observed 369 DEGs shared between our ATACseq and scRNAseq datasets (Suppl. Fig. 5o) and

mapped the proportions of genomic features that are up- and downregulated (Suppl. Fig. 5p). We

also observed a smaller but substantial number of genes (2,384) with associated upregulation in

chromatin accessibility following *Cebpa* loss, indicating repressive regulation, including the

transcriptional start site (TSS) of *Dlk1* (Fig. 5g). Further, the change in enrichment of

transcription factor binding sites following *Cebpa* loss is consistent with prior work on Nkx2.1

and AP-1 acting as a bivalent co-factor to reinforce fate of both AT1 (via Yap/Taz interactions)

and AT2 (via C/EBPα) cells[55] (Suppl. Fig. 5q, r, s).

In investigating putative regulatory sites near *Dlk1*, we initially considered the nearby

intergenic differentially methylated region (IG-DMR) previously demonstrated to strongly

regulate *Dlk1* expression[56]. While no change in accessibility was detected in the IG-DMR locus,

a region nearby was found to decrease following *Cebpa* loss (Fig. 5g). Further, this region

included a predicted regulatory site containing a C/EBPα binding motif (Fig. 5g, dashed red)

where it could possibly act to repress *Dlk1* and that we validated C/EBPα binds to in AT2 cells

in a previously published ChIPseq dataset[52].

PRC2 and C/EBPα act as a *Dlk1* "pulse generator" critical for patterning alveolar

epithelial fate

Given the perinatal repression of *Dlk1* by C/EBPα, we were curious of its expression earlier in

lung development. As previously reported[57, 58], we observed that *Dlk1* is lowly expressed at

stages prior to alveolar epithelial differentiation (e12.5-e15.5) both at the protein (Fig. 6a) and

RNA level (Fig. 6b, dashed grey box). Further, *Dlk1* expression appeared to peak in the period

from e17.5 to PN1 (Fig. 6a, b, Suppl. Fig. 6a). As *Cebpa* is not detected at earlier stages ($\leq$

e15.5) (Suppl. Fig. 6b), we were curious how *Dlk1* downregulation is mediated. Suppression of

Dlk1 has also been found to be mediated by the polycomb repressive complex (PRC2), including

the H3-K-27 methyltransferase EZH2[59]. We observed that genes encoding PRC2 members were

enriched at earlier stages (Fig. 6b, red text), and further that EZH2 protein is substantially

downregulated after e16.5 (Fig. 6c).

As EZH2 and the PRC2 complex are enriched earlier when *Dlk1* is downregulated, we

next tested whether EZH2 activity is required to suppress DLK1 in culture. Isolated e15.5 DPs

were cultured as organoids and treated either with FGF7 alone or with the EZH2 inhibitor

GSK126 (10 µM) for 4 days (Fig. 6d). While control organoids had detectable DLK1 protein

($\sim$12% DLK1$^+$ cells), levels significantly increased when treated with the EZH2 inhibitor ($\sim$22%,

Fig. 6e), indicating EZH2 methyl transferase activity is required for DLK1 suppression.

When characterizing the expression of *Ezh2* alongside *Cebpa*, we noticed a consistent

mutually exclusive pattern (Fig. 6b, Suppl. Fig. 6c), suggesting EZH2/PRC2 might also suppress

Cebpa. We analyzed a published ENCODE epigenetic dataset that mapped the PRC2

modification H3K27me3 across development[60] and observed its time-dependent reduction at the

Cebpa locus concomitant to when *Cebpa* expression is upregulated (Suppl. Fig. 6d). Further,

analysis of ChIPseq database of mouse tissues and cells[61] confirmed significant enrichment in

binding of EZH2 as well as other PRC2 components near the *Cebpa* locus (Fig. 6f). To

determine whether EZH2/PRC2 is also necessary for repressing C/EBPα, we again conducted

organotypic culture with e15.5 DPs during which differentiation was promoted by addition of

FGF7 as previously described[10]. Blocking of EZH2 activity by GSK126 significantly increased

the percentage of C/EBPα$^+$ cells compared to control (55% versus 34% respectively, Fig. 6g, h).

Finally, we confirmed whether *Ezh2* is required in vivo to suppress C/EBPα by using a sparse

deletion approach wherein a Cre line we previously found to sparsely label DPs[10] (*Tg$^{Nkx2.1-Cre}$*)

was bred to a transgenic line to delete *Ezh2* and lineage label by GFP (*Tg$^{Nkx2.1-Cre}$*

Rosa26$^{mTmG/mTmG}$; Ezh2$^{fl/fl}$). Immunofluorescence conducted at e15.5, a timepoint where no

C/EBPα is observed (Suppl. Fig. 6b), found significant upregulation of C/EBPα in *Ezh2* deleted

(GFP$^+$) versus control (RFP$^+$) DPs residing within the same end buds (Fig. 6i, j).

The interactions between EZH2/PRC2, C/EBPα, and DLK1 we observed appeared to fit

the definition of a type 2 incoherent feed-forward loop[62] (I2-FFL, Fig. 6k). An incoherent

feedforward loop (I-FFL) is a type of network motif found in biological systems, such as gene

regulatory networks. It contains three components, a regulator, an intermediate , and a target –

we propose these components in our model to be PRC2, C/EBPα, and *Dlk1*, respectively (Fig.

6k). The term incoherent means that the regulator (PRC2) acts in a conflicting manner to the

target — by both inhibiting (directly on *Dlk1*) and activating it (indirectly through repression of

the repressive intermediate C/EBPα. One emergent property of an I-FFL is the capacity to

generate a pulse of the target gene's expression (*Dlk1*) following downregulation of the upstream

regulator (EZH2/PRC2). To more precisely assess whether the observed temporal pattern of

Dlk1 expression matched the pulse predicted by and I2-FFL, we perform modeling using the

biocircuits package (https://biocircuits.github.io/). After establishing reasonable values for

parameters describing the relationship of factors in the I2-FFL (see Methods), the model was

found to fit well to the observed *Dlk1* expression pattern as assessed by root mean squared error

(RMSE = 0.12, Fig. 6l, Suppl. Fig. 6e). Next, we wanted to assess the importance of this

molecular circuit in patterning alveolar epithelial differentiation. Immunostaining of e15.5 DPs

in organotypic culture for markers of AT2 and AT1 fate (SFTPC and RAGE, respectively)

revealed that blocking EZH2 activity, with or without FGF7 treatment, significantly increased

the proportion of molecularly specified AT2s while concomitantly reducing AT1s compared to

FGF7 alone (Fig. 6m, n). Finally, we wanted to assess whether lateral inhibition of Notch

signaling occurs in AT1 cells, as it has been well demonstrated to promote a spatial "salt and

pepper" pattern of cell fates[63]. Analysis of scRNAseq timecourse datasets found that nascent

AT1 cells upregulate the Notch target gene *Hes1* suggesting signaling upregulation as expected

from prior studies[64] as well as downregulation of Notch ligands *Jag1* and *Dll4* that is consistent

with lateral inhibition of Notch signaling (Suppl. Fig. 6f, g). Further, we also observe in nAT1s

changes in Notch regulators that is also consistent with lateral inhibition, specifically

downregulation of *Dlk1* and upregulation of *Dlk2* (Suppl. Fig. 6f, g) - the latter has been found to

act as a cis-inhibitor of DLK1[65]. Therefore, our findings suggest that the potential model of

Cebpa-mediated fate regulation is as follows (Fig. 6o): In the *Cebpa*$^{fl/fl}$ background, the AT2 fate

plasticity window is reopened, and the pulsing of DLK1 upon the removal of repressors converts

a subset of AT2s into Notch signaling sending cells. Through further lateral inhibition, the

molecular circuit depicted in Fig. 6k likely contributes to AT1/AT2 patterning during lung

development. Taken together, our findings support the model that EZH2/PRC and C/EBPα act as

a molecular circuit (specifically an I2-FFL) to generate a DLK1 pulse during late-stage

embryogenesis that is critical to patterning the proportion of AT2 and AT1 cells.

C/EBPα suppression is required for accessing both reparative and defensive AT2 states

Given the role of C/EBPα in suppressing plasticity in AT2 cells, we sought to determine if and

how it is overridden following injury in the adult lung, where AT2s are well established to act as

facultative progenitors to regenerate lost AT1 cells[4, 22]. To first determine if C/EBPα

downregulation is required following injury in adulthood for AT2-to-AT1 conversion, we

infected adult mouse lungs harboring a Cre-dependent fluorescent reporter ($Rosa26^{mTmG/mTmG}$)

with AAVs targeting AT2 cell expression via a minimal *Sftpc* promoter as previously described[10]

to express either Cre alone or Cre and C/EBPα. After five days, we injured lungs via

intratracheal instillation of the irreversible FGFR inhibition Fiin1 (14.75 mg/kg, as previously

characterized by our group[10]) or intraperitoneal injection of BHT (225 mg/kg) and performed

immunostaining 14 days afterwards (Fig. 7a upper, Suppl. Fig. 7a, b). Intratracheal instillation of

AAVs efficiently labeled AT2s with high specificity (Suppl. Fig. 7c). While injured lungs

showed AT2-to-AT1 conversion when treated with the control AAV, it was significantly reduced

in those given C/EBPα overexpressing AAVs (Fig. 7a, b).

Next, we examined whether C/EBPα expression is downregulated following injury using

the 1X Bleomycin model, which results in patchy regions of injury and that resolve within 4-6

weeks. Bleomycin treated lungs were imaged 16 days after administration and injured areas

detected by alveolar simplification and reduction in E-cadherin as previously observed[66].

Immunostaining for C/EBPα revealed a significant downregulation in injured versus uninjured

regions (Fig. 7c, d). This finding is in alignment with analysis of a recent scRNAseq timecourse

study[20] following single-dose bleomycin, during which *Cebpa* expression is downregulated in

AT2 cells as early as 2 days post bleomycin (Fig. 7e).

After establishing the presence and necessity of C/EBPα downregulation following

injury, bioinformatic screening of the bleomycin injury timecourse for transcriptional regulators

known to suppress C/EBPα that are also upregulated in the previously identified AT2-to-AT1

transitionary[20] state (Krt8[+] ADI, also known as DATP[32] or PATS[67]) identified three genes, *Atf3,*

Atf4, and *Ddit3* (Fig. 7e, Suppl. Fig. 7e). Both activating transcription factors 3 and 4 are known

to bind the *Ddit3* promoter to drive expression[68, 69], and DNA damage-inducible transcript 3

protein, also known as C/EBP homologous protein (or CHOP), has been well established to

directly bind and act as a dominant negative inhibitor of C/EBP transcription factors[70]. Given its

direct interaction and inhibition of C/EBPα, we next wanted to determine whether induction of

CHOP is alone sufficient to drive AT2-AT1 conversion. A day after lineage labeling AT2 cells

in lungs of perinatal mice (PN11, *Sftpc[CreER]; Rosa26[mTmG]*), intraperitoneal injections of either

vehicle alone or a small molecule inducer of CHOP via PERK activation CCT020312 (2 mg/kg)

were administered every four days and lungs isolated at PN25 (Fig. 7f). While little to no lineage

labeled AT1s were observed in vehicle treated lungs, there was a significant increase in those

treated with CCT020312 (Fig. 7f, g), suggesting PERK/CHOP induction can promote AT2-to-

AT1 conversion.

To precisely map AT2 transcriptional trajectories and programs following *Cebpa* loss, we

conducted Velocity analysis as well as Leiden clustering on our scRNAseq dataset and

confirmed a cellular trajectory from AT2 to AT1 cluster as expected (Fig. 7h). Unexpectedly, we

observed another distinct trajectory terminating on a distinct AT2 cell cluster that was enriched

in genes associated with pathogen defense response and interferon signaling by GO analysis

(Fig. 7i, Suppl. Table 2), leading us to name the "defender" state AT2[def] (Fig. 7h, red).

Visualizing the expression of AT2[def] enriched genes versus those associated with the damage-

associated transitionary progenitor (DATP) state[32], we observed these programs were largely

mutually exclusive, apparent by both UMAP (Fig. 7j) and SPRING plot (Fig. 7k, l). Given the

distinct trajectories, as well as prior research indicating interferon signaling negatively impacts

lung epithelial repair[24], we were interested whether stimulation of the AT2^{def} state by interferon

signaling would block AT1 conversion in culture. Given the expression pattern of interferon

receptors as well as transcriptional targets in the AT2^{def} cluster (Suppl. Fig. 7d, e), we treated

cultures with interferon-γ (Fig. 7m). While perinatal AT2s could give rise to AT1s following

FGF7 treatment in organotypic culture, addition of exogenous interferon-γ resulted in a

significant reduction AT1 conversion, consistent with the model that AT2^{def} and AT2-to-AT1

molecular program are antagonistic to each other at the cellular level (Fig. 7m, n). To assess the

degree to which these programs are present in the same or separate AT2 cells following injury,

we analyzed scRNAseq datasets wherein lungs were injured in distinct ways, including

diphtheria toxin ablation of AT1 cells[71], influenza[72], LPS[73], and Bleomycin[20]. In support of our

findings following *Cebpa* deletion, DATP and Defender states were found to primarily exist in

separate AT2 cells during the repair process across all forms of injury (Suppl. Fig. 7f).

Discussion

These results show that AT2 fate plasticity exists in a time-window from late embryonic to early

postnatal life consistent in time with nascency. The timing and length of this window is dictated

by a molecular circuit driven by PRC2 and C/EBPα that together generate a pulse of *Dlk1,* a

known regulator of adult AT2 fate plasticity. Downregulation of C/EBPα occurs following injury

in the adult lung, is required to regenerate AT1 cells, and is mediated in part by the dominant

negative C/EBP family member CHOP. C/EBPα also suppresses an AT2 defensive state that is

mutually exclusive from fate plasticity that appears to be driven by interferon signaling.

From characterizing its nascent state, we determined AT2 cells first emerge around e15.5

at intermediate regions between the tips and stalks of the distal branches. Our findings

consolidate differing models on AT2 emergence, with our observed timing supporting the

bipotent model[4, 9, 10, 33, 37] and our observed pattern suggesting an intermediate starting point,

distinct from that proposed in both the bipotent (more proximal) and earlier differentiation[12]

(distal tip) models. After initial emergence (e15.5-16.5), subsequent AT2s next arise at proximal

regions coinciding with ASM-to-myofibroblast differentiation as previously reported[34]. Taken

together, these findings suggest more than one cue may be present to suppress alveolar epithelial

differentiation, one distal and another proximal in position. While it has been established that

distal tips are enriched in Wnt activity and thus retain stemness[74], it is unclear how more

proximally positioned DPs are suppressed. One compelling possibility is that ASM directly

suppresses differentiation of these proximally localized distal progenitors through direct

interaction on the basal epithelial surface, although it is unclear how said suppression is

molecularly mediated.

Our findings on AT2 emergence have also provided insight into the sequence of their

morphological changes. Given that basal extrusion occurs after AT2 molecular specification, it is

likely downstream of fate selection rather than an upstream regulator as previously proposed[75].

This observation is supported by prior work wherein AT2 fate selection was found to occur

under culture conditions that block basal extrusion[10]. If a byproduct of fate selection, what

purpose does basal extrusion serve? Following extrusion, we observe that some extruded AT2

cells engage with nearby epithelium from anatomically distinct branches, forming what we name

interlumenal junctions, or ILJs. We hypothesize basal extrusions are required to form these

junctions, that they are mediated in part by contraction of the AT2 lumenal surface, and likely

play a critical early role in postnatal alveologenesis, the stitching together of alveolar lumens[76].

While this process has been observed at the tissue level[14], as has the multilumenal nature of

mature AT2s[34], many questions remain as to the underlying cellular and molecular mechanisms

governing it. Further work on interlumenal junctions will be critical to understanding this stage

of alveolar maturation and assessing whether its disruption could underly diseases such as BPD

and COPD.

Following their emergence, nascent AT2 cells undergo transition towards their mature

AT2 transcriptional state over the course of ~7 days. This process occurs in a sporadic, salt and

pepper pattern distinct from the reported proximal-to-distal wave of alveolar epithelial fate

selection[13, 77] suggesting the cues driving maturation might arise from a different source, which is

likely to be immune given the genes enriched in mAT2s as well as prior studies demonstrating

interactions underlying one another's maturation[33] and function[15]. While AT2s have been shown

to impact macrophage maturation[33], an immune cell type has not been implicated in driving AT2

maturation during postnatal development. More work is needed to better understand which

immune cell types interact with AT2 cells during this process and determine how they impact

differentiation. While AT2 cells from both nascent and mature states retained plasticity, mAT2s

quickly lose this capability within the first week or so after birth, suggesting fate plasticity is

somewhat negatively correlated to AT2 maturation. Given these findings, we revised our model

of alveolar epithelial differentiation to include a fate plasticity window extending from e15.5

distal progenitors into AT2 nascency, and that tapers down over 1-2 weeks after birth. It is likely

this window is critical for postnatal alveologenesis and might be involved in the observed age-

dependent severity of COVID19[78].

While our bioinformatic screening for transcriptional regulators of the plasticity window

identified C/EBPα as a possible repressor, previous genetic deletions studies concluded that

Cebpa is required earlier for both AT1 and AT2 differentiation[46, 47]. As these findings conflict

with the observed expression pattern of C/EBPα, which is upregulated following AT2 fate

selection, we conducted sparse *Cebpa* deletion experiments in both DP and AT2 populations.

From these studies we found that *Cebpa* is dispensable for the selection but required for the

maintenance of AT2 fate, with the later finding recently confirmed in a manuscript by another

group[52]. The discrepancy is likely the result of the field's prior use of genetic tools now found to

cause toxicity in the developing mouse lung[48]. While our subsequent scRNAseq and ATACseq

analysis reveals repressive targets of C/EBPα, its positive transcriptional regulation of gene

expression has been demonstrated including recent publications [52]. While *Cebpa* loss does result

in a reduced expression of several AT2 mature genes, identified in both studies, it is important to

note that these AT2 cells do not concomitantly increase the expression of AT1 genes nor directly

convert into AT1 cells. Rather the AT2 program reverts to the nascent AT2 state, again

suggesting a repressive mechanism is at play that targets fate plasticity. While Sox9 was reported

in the other study to be upregulated following *Cebpa* loss, we only detected a modest increase at

the RNA level and screening of scRNAseq data referencing a DP gene list revealed no strong

upregulation, supporting our model that loss resulted in reversion to a more nAT2 state rather

than full dedifferentiation back to DP. Recent work has found that AT1s in the perinatal period

also retain fate plasticity and when stretch and YAP/TAZ signaling is lost they are capable to

AT1-to-AT2 differentiation[79]. While more work is needed to integrate those findings with ours

and others[52], one compelling model is that both perinatal AT2 and AT1 cells retain fate plasticity

for a period of time but are each suppressed by distinct repressors – C/EBPα for AT2s and

YAP/TAZ for AT1s.

Our investigation into the key gene(s) repressed by C/EBPα that could regulate fate

plasticity identified *Dlk1*, which encodes a cis-regulator of Notch signaling previously implicated

in AT2-to-AT1 conversion[23]. Notch is distinct from other signaling pathways in its use of cis-

regulators (like DLK1) for activation and has been shown to be critical in generating its well

characterized "salt and pepper" pattern[80]. While regulation of the *Dlk1* locus is well documented

to be mediated by an intergenic differentially methylation region (IG-DMR) downstream of the

gene[56, 81], ATAC-seq revealed no change in its accessibility nor any changes immediately

upstream of the gene's transcriptional start site upon loss of *Cebpa*. Rather, a dramatic change in

accessibility was observed in a previously uncharacterized region roughly 2 kilobases

downstream of the IG-DMR that the UCSC genome browser predicts is an enhancer. More work

is needed to determine how this element is regulated during development as well as repair

following injury in the adult lung.

Our investigation into developmental regulation of *Dlk1* expression determined that

PRC2 and C/EBPα act as a type 2 incoherent feedback loop to generate of pulse of DLK1

following downregulation of the PRC2 complex and likely its repressive epigenetic mark,

H3K27me3. While several feedback loops have been postulated and constructed in the field of

systems biology[62], most of the loops observed in eukaryotic cells are either type 1 coherent or

incoherent in configuration[62]. While the identified loop type (I2-FFL) is observed less frequently,

we believe the PRC2-C/EBPα regulation plays a critical role in timing and pattern of alveolar

epithelial differentiation, and future studies are needed to test whether reactivation of said loop

could be used to re-access AT2 fate plasticity in diseased lungs. We believe this feedback loop

constitutes a novel molecular circuit – a "pulse generator" of Notch activation, resulting in

spatial patterning of AT1 and AT2 differentiation through lateral inhibition. It is also conceivable

that disruption of this molecular circuit could result in an aberrant AT2 state that is disease

associated, such as those observed in models of IPF[82, 83].

Finally, our investigation into the adult lung demonstrates that C/EBPα downregulation

following injury is required for repair and is partly mediated by a C/EBP family member with

dominant negative activity, CHOP[70]. As CHOP is upregulated from various types of injury that

cause ER stress[84], it is still unclear how C/EBPα downregulation is stimulated following injury

of alveolar epithelial cell death, especially as we observe little to no C/EBPα downregulation

upon sparse AT2 ablation (Suppl. Fig. 7e). One compelling model requiring further inquiry is

that AT1 death sends a specific cue that downregulates C/EBPα to promote fate plasticity in

nearby AT2 cells and thus alveolar repair. Interestingly we found that *Cebpa* loss also allowed

AT2 cells to access what appears to be a pathogen defensive state driven by interferon signaling.

While presence of such a state is expected, as AT2 cells have been implicated in playing a role in

immune response in the lung[15], said state was not expected to be suppressed by C/EBPα nor to

be a mutually exclusive transcriptional state from the well-established regenerative DATP state[67,

85].

Taken together, these observations suggest that C/EBPα might suppress a larger program

that alternates between two states – one regenerative and the other defensive. If true, said

program would represent a more robust response to pathogen-mediated injury then previously

appreciated, as alternating between regenerative and defense states could ensure enough AT2s

are available at a given time to both promote clearance of the pathogen as well as repair damaged

alveoli (Fig. 7o).

We will briefly cover the limitations of this study to motivate and direct future work.

First, we dissociate murine lungs to form organoids which, while a common approach in the

field[10, 22, 34], might alter cellular phenotypes and promote fate plasticity. While the observed

behaviors of cultured DPs and AT2s largely match in vivo experiments[10, 22, 34] and velocity

analysis (Figure 3e-j), future work investigating fate plasticity more precisely in vivo would be

informative. Differences observed by normalized gene expression alone would benefit from

validation via ISH or immunostaining. As our study does not investigate whether AT1s derived

from DPs are distinct from those from AT2s, more work is needed to make said delineation. As

conclusions of scRNAseq cluster similarity can be impacted by cut-off points, more work

delineating differences in e15.5 DPs in an unbiased manner would be informative. While we

focused on the role of C/EBPα, future work on other genes from our screen would likely identify

other critical regulators of AT1/2 fate plasticity (see Suppl. Table 3). As BH3 death regulation is

studied only to determine its similarity between DPs, nAT2s, and mAT2s, more research is

needed to determine its role during development and repair. As CHOP expression was induced

using a smaller activator of PERK (CCT020312) that may stimulate off target molecules

conflating our experiments, further work is needed to precisely investigate the function of CHOP

in vivo. Finally, as our experiment revealing C/EBPα's regulation of a defensive AT2 state did

not directly introduce a pathogen, future work is needed to confirm said state is upregulated

following infection.

Methods

Ethics

All mouse experiments followed applicable regulations and guidelines and were approved by the

Institutional Animal Care and Use Committee at the Mayo Clinic (Protocol# A00006319-21).

Mice were house under a 12 h light/12 h dark cycle and lights were not used during the dark

cycle. Mouse housing temperature was kept at 65-75 °F (~18-23 °C) and humidity at 40–60%.

Mice were housed in filtered cages and all experiments were performed in accordance with

approved Institutional Animal Care and Use Committee protocols and ethical considerations at

the Mayo Clinic.

Mouse strains

Timed-pregnant C57BL/6 J females (abbreviated B6; Jackson Laboratories) were used for all

embryonic time points, with gestational age verified by crown–rump length. For studies of adult

wild-type lungs, B6 males and females were used. Sparse labeling and recombination studies

were conducted by Cre recombinase expression using gene-targeted alleles BAC-*Nkx2.1*^{Cre},

LyzM (also called Lyz2)-Cre, and *Sftpc*$^{Cre-ERT2-rtTA}$[58]. Cre-dependent target genes were the

conditional "floxed" *Cebpa*[86] and *Ezh2*[87] for removal of *Cebpa* and *Ezh2* respectively. Two Cre

reporters were used: *Rosa26*^{mTmG} [88] (JAX stock #037456) and Tg(iSuRe-Cre) (MGI:6361145) [49].

Upon recombination, the Tg(iSuRe-Cre) allele labels the cell with membrane and cytoplasmic

tdTomato (tdT) and constitutively expresses Cre recombinase in the targeted cell, thus ensuring

that all Cre-conditional alleles are recombined, even under sparse or clonal conditions. For

initiating recombination with *Sftpc*$^{Cre-ERT2}$ (*Sftpc*^{CreER}), intraperitoneal injection of tamoxifen was

administered either at 20 mg/kg for complete (high dose) recombination in adult mice (> PN60),

or at 30 ng/kg for sparse recombination in perinatal mice (P0-P21), except where noted. EdU

(Cayman Chemicals 20518) was administered via drinking water at 0.1 mg/mL for the indicated

period of time. Genomic DNA was extracted from ear punches followed by Proteinase K (KAPA

Biosystems) digestion, and genotyping was performed by polymerase chain reaction (PCR) using

published primer sets.

Single-cell RNA-seq bioinformatic analysis

Cell Isolation for library preparation

643   Cells from PN25 *Sftpc^{CreER}; Tg(iSuRe-Cre);Cebpa^{wt/fl}* and *Sftpc^{CreER}; Tg(iSuRe-Cre);Cebpa^{fl/fl}*

644   mice (2 weeks after TMX) were isolated and enriched (EpCAM$^+$) as per protocol described

645   below.

scRNAseq library preparation and processing

Single cell RNA sequencing was performed using the Chromium Single Cell 3' system (10x

Genomics) at the single cell sequencing core at Mayo Clinic according to manufacturer's

instruction (10x Genomics). Sequencing depth is an average of 75k reads per cell to detect lowly

expressed genes. Fastq files were generated using Cellranger v 7.2. The sequenced files were

mapped to the GRCm38/mm10, supplemented with tdTomato, PhiMut, and WPRE transcripts as

per the Tg(iSuRe-Cre) construct[49].

scRNAseq data analysis

We used Scanpy[89] and Seurat v.4.2.3[90] for processing and downstream analysis. In the Scanpy

pipeline, we identified highly variable genes using the scanpy function

scanpy.pp.highly_variable_genes with arguments min_mean = 0.0125, max_mean = 3, and

min_disp = 0.5. We restricted to this set of genes for downstream analysis. We further used

scanpy.pp.regress_out to regress out unwanted variation in the count of Unique Molecular

Identifiers (UMIs) and the percentage of mitochondrial reads. We ran scanpy's built in Principle

Component Analysis (PCA) algorithm with sc.tl.pca and default parameters. We then created a

nearest neighbor graph using sc.pp.neighbors with 200 neighbors and 50 Principle Components.

We then produced a UMAP dimensionality reduction with scanpy.tl.umap using min_dist =

0.001. We identified cell clusters through unsupervised clustering using the Leiden clustering

algorithm through with resolution = 0.8. We assigned cell labels to the clusters based on marker

gene expression, and removed several contaminating mesenchymal clusters. These clusters were

then visualized on the UMAP.

In the Seurat pipeline, doublets were estimated and removed using the DoubletFinder

Package[91]. Dead cells or outliers were excluded by removing cells with less than 500 features

and with more than 10% mitochondria gene content. Post alignment, the Seurat objects were

merged, normalized with LogNormalize method, and scaled. Top 3000 variable genes were used

to anchor. After PCA dimensional reduction with 30 principal components, we mapped our data

on UMAP projection with Louvain clustering algorithm. Subsequently canonical marker genes

for AT1 and AT2 cells[92] and sample conditions were used to assign cellular groups.

For RNA velocity, velocyto was used with default run command and run10x sub-

command to obtain velocity read counts[93]. For velocity analysis, we used Scanpy and scVelo

packages. The datasets were filtered by expression of < 200 genes or > 3200 genes, and genes

were expressed in < 3 cells. The dataset was subsequently normalized and the top 3000 highly

variable genes with a minimum 20 expressed count threshold were used in preprocessing. A

dynamical model was used to recover velocity estimate following Scvelo default parameters

(pp.moments(n_neighbors = 30), tl.recover_dynamics(fit_scaling = False), tl.velocity(mode =

'dynamical')).

684   We then produced velocity stream overlay on UMAP projection and latent time. We used

685   kmeans clustering from the SKlearn package to identify the two predominant directions of

686   velocity vectors within the nAT2 cell cluster. We then calculated an AT1/AT2 score in these

687   clusters based on expression of cell markers. We further performed partition-based graph

688   abstraction (PAGA)[94] based on the velocity vectors of each cell type to determine broader cell

689   type linkages[95].

690   For STITCH analysis (to generate SPRING plots), we ported in formatted datasets and

691   followed steps of logNormalization, and determined highly variable genes as instructed in the

692   Wagner 2018[96] example in MATLAB 2023a. The visualization was done using ForcedAtlas

693   layout in Gephi v7.2.1.

694   For analysis of lung development through embryonic, perinatal, and adult timepoints in

published scRNAseq datasets, the processed mRNA counts for each cell were used from

developing mouse lung datasets, (GSE119228)[33], (GSE113320)[12], and (GSE149563)[37]. Analysis

of bleomycin-induced lung injury on AT2 cells was performed on a published scRNAseq dataset

(GSE141259)[20].

Lung isolation and processing

For prenatal time points, individual embryos were staged by fetal crown–rump length before

sacrifice and lungs removed *en bloc*. For postnatal time points, mice were euthanized by either

carbon dioxide inhalation or intraperitoneal injection of Pentobarbital (Fatal-Plus Solution) and

dissected to both expose the lungs and allow for exsanguination via the abdominal aorta. 5-10

mL of phosphate buffered saline (PBS; Ca^{2+} and Mg^{2+} free, pH 7.4) was gently perfused into the

right ventricle of the heart by syringe with a 25-gauge needle until the lungs appeared white. For

postnatal time points ($\geq$ 8 days), the trachea was then cannulated with either a blunt 22- or 25-

guage catheter, and the lungs were gently inflated to full capacity with molten low melting point

agarose (Sigma, 2% in PBS). For all time points, lungs were then separated by lobe and fixed in

4% PFA at 4 °C for either 1 h (e15.5 – e16.5), 1.5 h (e17.5), 2 h (e18.5 – PN10), or 4 h (>

PN10). For sectioning, a vibrating microtome (Leica) was used to generate embryonic (150 µm)

or adult (250 µm) tissue sections of uniform thickness.

Immunostaining

Immunohistochemistry was performed as previously described in Treutlein et. al.[9] using primary

antibodies against the following epitopes (used at 1:500 dilution unless otherwise noted): proSPC

(rabbit, Sigma AB3786), RAGE (rat, R&D MAB1179), E-cadherin (rat, Life Technologies

ECCD-2), PDPN (hamster, DSHB 8.1.1), MUC1 ((hamster, Thermo Scientific HM1630 and

rabbit (GeneTex GTX100459)), Alpha-SMA (1:100, mouse IgG2a, Invitrogen 50-9760-82),

EZH2 (ENX-1; mouse IgG2b, Santa Cruz Biotechnology sc-166609), KI67 (rat, Thermo

Scientific 14-5698-82), RELMα (rabbit, PeproTech 500-P214), CD74 (rat, BD Biosciences

555317), DLK1 (rabbit, R&D Systems MAB8634), C/EBPα (rabbit, Cell Signaling 8178),

LAMP1 (rat, DSHB 1D4B-s), and HOPX (rabbit, Abcam ab307671). Briefly, fixed lung slices

were permeabilized and blocked overnight in blocking buffer (5% goat serum/PBS/0.5% Triton

X-100) at 4 °C. Primary antibody incubation steps were conducted in blocking buffer over 3

nights and washing steps were performed in five rounds (30, 30 60, 90, and 90 min respectively)

using the blocking buffer. Primary antibodies were subsequently detected using Alexa Fluor-

conjugated secondary antibodies used at 1:1000 dilution unless otherwise noted, including Goat

anti-rabbit IgG Alexa 555 conjugated (Invitrogen, A21428), Goat anti-rabbit IgG Alexa 647

conjugated (Invitrogen, A32733), Goat anti-rat IgG Alexa 488 conjugated (Invitrogen, A11006),

Goat anti-rat IgG Alexa 647 conjugated (Invitrogen, A21247), Goat anti-hamster Alexa 647

(Invitrogen, A21451), and Goat anti-hamster Alexa 594 (Invitrogen, A21113) unless noted

otherwise. Secondary antibody incubation steps were conducted in blocking buffer over 2 nights

and washing steps were performed in five rounds (30, 30 60, 90, and 90 min respectively) using

PBT (0.1% Tween-20/PBS). For co-staining using antibodies from the same rabbit origins, we

performed sequential staining with 1-day anti-rabbit blocking between processes. In brief, we

stain with primary and secondary antibodies of the 1st marker as described, and then we block the

stained tissue with 40 μg/mL Fab Fragment Goat Anti-Rabbit IgG (Jackson ImmunoResearch,

AffiniPure™, 111-007-008) prior to the 2nd marker staining. EdU was detected using the standard

Click-iT reaction (Thermo Fisher). Lung slices were then mounted in Vectashield Mounting

Medium (Vector labs, H1000) and sealed with a coverslip. Images were acquired using a laser-

scanning confocal microscope (Zeiss LSM 780 and Leica Stellaris 8) and subsequently processed

using ImageJ.

Timelapse Imaging

Fluorescently lineage labeled mouse lung were sectioned at 200 μm thick using compresstome.

When imaging, we image sectioned slices in serum-free DMEM media on top of Matrigel for

stability, using Timelapse setup in LAS-X software for 24 h consecutively.

Confocal Images 3D reconstruction

We reconstructed 3D representation of cell imaging using IMARIS 10.2.1. The z-stack images

are imported as .tif images and reconstructed into 3D representation of tissue images. Marker

intensities are individually normalized thresholds for intensity and contrast, smoothing cellular

structures for interpretation accuracy, and removal background noise.

Cell isolation and culture

For time points e15.5, e18.5, and PN3, lungs were isolated *en bloc* without perfusion and pooled

by litter (6-8 lungs) for further processing. The lungs were then dissociated in 50 U/mL Dispase

(Corning 354235) and triturated with flame sterilized glass Pasteur pipettes of decreasing

diameters (1,000 µm, 800 µm, and 600 µm) to achieve a single-cell suspension. For PN25 and

adult time points, the lung was perfused with DPBS (Ca^{2+} and Mg^{2+} free, pH 7.4) as described

above. The trachea was punctured and then cannulated with a blunt 22-guage catheter, and the

lungs were inflated to full capacity with Dispase (~1 mL), and the trachea was sealed using a

non-absorbable silk suture. The inflated lobes were incubated in Dispase for 45 min at 25 °C on

a shaker.

For all time points, digested lungs were minced with a razor blade, suspended in 7 mL of

digestion buffer (RPMI-1640 containing FBS (10%, Corning), penicillin-streptomycin (1 U/mL,

Thermo Scientific), and DNase I (0.33 U/mL, Roche)), incubated at 4 °C for 10 min on a flat

shaker and triturated briefly with a 5-mL pipette.

To deplete red blood cells, an equal volume of AT2 isolation media (RPMI-1640

supplemented with penicillin-streptomycin (1 U/mL, Thermo Scientific), and DNase I (0.33

U/mL, Roche)) was added to the lung single-cell suspension. The suspension was passed through

a 100 µm mesh filter to remove residual tissue fragments and centrifuged at 400×g, 4 °C for 10

min. Pelleted cells were resuspended in 5 mL of red blood lysis buffer (BD Biosciences),

incubated for 2 min at room temperature, and neutralized with an equal volume of AT2 isolation

media. The suspension was passed through a 40 µm mesh filter (Fisher), centrifuged at 400×g, 4

°C for 10 min, and then resuspended in MACS buffer (2 mM EDTA, 0.5% BSA in PBS, filtered

and degassed) for purification.

The resultant single-cell suspension was incubated with Mouse Fc Blocking Reagent

(Miltenyi Biotec, 130-092-575) for 5 min at 4 °C. Other cell types were first depleted with

antibodies against CD45 (eBioscience, 13-0451-85), CD31 (BD Biosciences, 557355), Ter119

(eBioscience, 13-5921-85), CD140a (Miltenyi Biotec, 130-101-905), and CD104 (Miltenyi

Biotec, 130-106-922), followed by anti-biotin (Miltenyi Biotec, 130-090-485) magnetic

MicroBeads. Cell suspensions were passed through a 35 µm cell strainer (BD Biosciences) prior

to being loaded on LD columns (Miltenyi Biotec) as per the vendor's protocol. The flow through

was centrifuged at 300×g, 4 °C for 5 min and the pelleted cells were resuspended in MACS

buffer.

For positive selection of cell types, the following antibodies were used: anti-EpCAM

(clone G8.8, eBioscience, 13-5791-82) for e15.5 distal bipotent progenitors, anti-RELMα

(PeproTech, 500-P214BT) for e18.5 nascent AT2 cells, and either anti-CD74 or anti-MHC-II

(clone M5/114.15.2, eBioscience, 13-5321-82) for mature AT2 cells. Following antibody

incubation, the cells were then incubated with either anti-biotin or anti-rat IgG MicroBeads and

loaded on MS columns (Miltenyi Biotec) as per the vendor's protocol. The flow-through was

discarded and the column was taken off the magnet. The cells remaining in the column were

eluted with MACS buffer, resulting in a highly enriched preparation of either bipotent

progenitors (EpCAM$^+$), nascent AT2s (RELMα$^+$), or mature AT2s (MHC-II$^+$), unless noted

otherwise.

For culturing bipotent progenitors, nAT2s, or mAT2s, cell density and viability were

calculated using a Vi-CELL XR Cell Viability Analyzer (Beckman Coulter). Cells were cultured

in eight-well #1.5 coverglass chambers (Cellvis, C8-1.5H-N), pre-coated with growth factor

reduced Matrigel (85 µL, Corning 354230) for 30 min at 37 °C. The cells were plated at varying

densities: 30,000 cells/well (for e15.5 and e18.5), 60,000 cells/well for PN3), and 80,000

cells/well (for PN25 and adult). Cells were supplemented with FGF7 (50 ng/mL, R&D Systems

5028-KG-025), GSK126 (10 µM, ApexBio A3446), KL001 (12 µM, MedChemExpress HY-

108468), and interferon-γ (20 ng/mL, PeproTech 315-05) to the indicated concentrations in 400

µL of DMEM/F12 supplemented with penicillin-streptomycin (1 U/mL, Thermo Scientific).

Excluding FGF7, all other small molecules listed above were added after a 12h incubation period

to allow for cells to seed and adhere to the layer of Matrigel. Cells were maintained in an

incubator at 37 °C with 5% CO2/air, with media changes every other day typically for 4 days,

except where indicated otherwise.

Flow cytometry

We perform flow cytometry of AT2 cells using Miltenyl's Tyto flow sorter. Filtered and

RBC depleted lung cell isolates are to be incubated with 20 µL of each of the following 1:50

diluted primary antibodies: anti-RELMα, anti-CD74, and anti-MHC-II for 30 min at 4 °C. Cells

are washed with sorting buffer (Miltenyi, 130-107-207) and subsequently incubated with 20 µL

of each of the following secondary antibodies: (Anti-rabbit Alexa Fluor 647, anti-rat Alexa Fluor

405) for 30 min at 4 °C. Subsequently the sample are washed, filtered, and resuspended in 1 mL

sorting buffer and transfer to the sorting compartment of the cartridge (Miltenyi, 130-104-791).

Gates for markers positivity are drawn using unstained cells as negative controls.

BH3 analysis

Lung samples from mice of different ages (e15.5, e18.5, and ≥ PN60) were dissociated into a

single-cell suspension and enriched using relevant markers as described above. Cells were

stained on ice for 25 min away from light, then centrifuged at 200×g for 5 min, and subjected to

flow cytometry–based BH3 profiling as previously described by the Sarosiek lab[78]. Briefly, cells

were treated with activator or sensitizer BH3 peptides (New England Peptide) for 60 min at 28

°C in mannitol experimental buffer (MEB) [10 mM Hepes (pH 7.5), 150 mM mannitol, 50 mM

KCl, 0.02 mM EGTA, 0.02 mM EDTA, 0.1% BSA, and 5 mM succinate] with 0.001% digitonin.

Peptide sequences are as follows: BIM (Ac-MRPEIWIAQELRRIGDEFNA-NH2), BID (Ac-

EDIIRNIARHLAQVGDSMDRY-NH2), PUMA (p53 upregulated modulator of apoptosis) (Ac-

EQWAREIGAQLRRMADDLNA-NH2), BAD (Ac-LWAAQRYGRELRRMSDEFEGSFKGL-

NH2), Hrk (Harakiri) (Ac-WSSAAQLTAARLKALGDELHQ-NH2), and MS1 (MCL-1

specific) (Ac-RPEIWMTQGLRRLGDEINAYYAR-NH2). After peptide exposure, cells were

fixed in 2% PFA for 15 min which was then neutralized by addition of N2 buffer [1.7 M tris base

and 1.25 M glycine (pH 9.1)]. Cells were stained overnight with DAPI (1:1000, Abcam) and

anti–cytochrome c–Alexa Fluor 647 (1:2000, clone 6H2.B4, BioLegend) in a saponin-based

buffer (final concentration, 0.1% saponin; 1% BSA) and then analyzed by flow cytometry.

Cytochrome c release in response to BIM treatment was measured on an Attune NxT flow

cytometer (Thermo Fisher Scientific).

Conditional deletion of *Cebpa* and *Ezh2 in vivo*

For conditional and efficient recombination of floxed alleles (*Cebpa^{flox}* or *Ezh2^{flox}*) at different

time points and cell types, we bred the floxed allele into one of four mouse lines, each with a

distinct combination of Cre and reporter alleles. For in vivo sparse cell labeling with efficient

recombination of either *Cebpa^{flox}* or *Ezh2^{flo}* alleles in distal progenitors, the *Tg$^{Nkx2.1-Cre}$*;

Rosa26$^{mTmG/mTmG}$ mouse line was used as we have previously found it sparsely labels DPs with

efficient flox allele recombination[10]. For sparse labeling of nAT2s with efficient recombination

of *Cebpa^{flox}*, the *LyzM^{Cre}*; *Rosa26$^{mTmG/mTmG}$* mouse line was used as previously described[10]. For

broad labeling of AT2 cells with efficient *Cebpa^{flox}* recombination following a high dose of

tamoxifen (20 mg/kg), the *Sftpc$^{Cre-ERT2}$*; *Rosa26$^{mTmG/mTmG}$* mouse line was used. Finally, to

sparsely label AT2s and retain efficient *Cebpa^{flox}* recombination following a low dose of

tamoxifen (30 ng/kg), the *Sftpc$^{Cre-ERT}$*; *Tg(iSuRe-Cre)* mouse line was used. Upon its

recombination, the *Tg(iSuRe-Cre)* allele constitutively expresses both the fluorescent lineage

label tdTomato as well as Cre recombinase, an approach that has been well established to

efficiently recombine multiple floxed under lower tamoxifen doses for targeting sparse cells[49].

To determine proliferation of AT2 cells, EdU (1 mg/mL) was administered in the drinking water

for the indicated period of time.

ATAC-seq and CHIP-seq analysis

Cells from PN25 *Sftpc^{CreER}; Tg(iSuRe-Cre); Cebpa$^{wt/fl}$* and *Sftpc^{CreER}; Tg(iSuRe-Cre); Cebpa$^{fl/fl}$*

mice (2 weeks after TMX) were isolated and enriched (EpCAM^{+}) as per protocol described

above. Raw sequencing reads were trimmed of sequencing adapters and bad quality bases with

cutadapt version 2.8 in Trim Galore version 0.6.6 using default parameters. Bowtie2 version

2.5.0 (parameters: --end-to-end --very-sensitive --no-mixed --no-discordant --phred33 -I 10 -X

700) was used to map trimmed reads to the reference genome Mus_musculus.GRCm38. Picard

MarkDuplicates (v.2.26.10; http://broadinstitute.github.io/picard) was used for duplicate

removal. Reads were shifted by + 4 bp for those mapping to the positive strand and – 5 bp for

those mapping to the negative strand. ATAC-seq analysis was performed with HOMER(v 4.11).

Tag directories were made for each ATAC-seq run using HOMER[97] "makeTagDirectory". Peaks

for individual replicates were identified using the "findPeaks" command. To generate high-

confidence peak lists across both replicate samples for downstream analysis, ATAC-seq peaks

were identified using the HOMER "getDifferentialPeaksReplicates.pl" command. This command

uses DESeq2[98, 99] and identifies peaks that pass 2-fold enrichment and false discovery rate (FDR)

< 0.05 cutoffs. For these analyses, peak lists combining all peaks from the comparison conditions

were assembled. One condition was used as the target, and the other was used as background. To

characterize enriched motifs near differentially enriched peaks, we used HOMER's

"findMotifsGenome.pl" command to scan the entire peak by using "-size given" with standard

background. ATAC-seq peak annotation was performed using the homer annotatePeaks.pl

function to assign the nearest gene name to the peaks. To estimate the enrichment of

transcription factors and epigenetic modifications on the murine genome, we used the Cistrome

DB Toolkit analysis browser[61, 100] using default settings.

Incoherent feed forward loop type 2 model fitting

To assess whether the temporal expression pattern of *Dlk1* fit that predicted by a feed forward

loop we used the publicly available biocircuits package (https://biocircuits.github.io/) to

numerically solve the differential equations for a type 2, incoherent feed forward loop following

a step reduction of PRC2. After solving for a function over time for *Dlk1*, an R script was

generated to align its peak value to that observed in the scRNAseq timecourse and calculate a

goodness of fit by root mean squared error (RMSE). Parameters of the I2-FFL model were

independently varied until a good fit was reached (beta = 1.9, gamma = 5.5, and kappa = 3).

Mouse lung injury experiments and viral transfections

Lung injury was induced in adult mice by single-dose administration of bleomycin sulfate

(0.25mg/kg; VWR J60727-MA), which was dissolved in sterile saline and administered via

intratracheal (i.t.) administration. Lungs were isolated 16 days post injury and processed for

immunostaining.

To test whether C/EBPα is required for proper injury repair, adult $Rosa26^{mTmG/mTmG}$ mice were

administered either an AAV-*Sftpc-Cre* (control), or an AAV-*Sftpc-Cebpa-T2A-Cre* (prepared by

the Boston Children's Hospital Viral Vector Core (1.5×10^{14} GC (genome copies)/mL, serotype

2/9) and intratracheally instilled (5 µL of viral solution diluted in 45 µL PBS)) to overdrive

Cebpa expression and induce GFP labeling of AT2 lineage. A minimal 320 bp *Sftpc* promoter

element was used for both AAV preps. 5 days later, all mice were administered Fiin1

hydrochloride (14.75 mg/kg; Tocris 4002) via intratracheal instillation as described in out prior

work[10] or BHT (225 mg/kg; Sigma B1378) via intraperitoneal injection[71] to induce lung injury.

Lungs were isolated 14 days post injury and processed for immunostaining. For all intratracheal

instillation experiments, mice were anesthetized with isoflurane and kept in recovery under a

heat lamp.

Small molecule induction of CHOP

To induce CHOP expression in AT2 cells, *Sftpc*$^{CreERT2-rtTA}$; *Rosa26*$^{mTmG/mTmG}$ mice were first

administered tamoxifen (20 mg/kg) at PN11 for labeling of AT2 cells with GFP. 24 h later, mice

were either given corn oil (vehicle) or CCT020312 (MedChem Express HY-119240), a PERK

activator found to upregulate CHOP expression[101], at 2 mg/kg dissolved in corn oil via

intraperitoneal injection every 4 days (4 doses total). Lungs were isolated 2 weeks later and

processed for immunostaining.

Statistical analysis

Data analysis and all statistical tests were performed with GraphPad Prism software or R.

Replicate experiments were biological replicated using different animals, and the quantitative

values are presented as mean ± SD unless indicated otherwise. In graphs displayed as box plots,

the horizontal lines represent the median and error bars depict the $75^{th} - 25^{th}$ interquartile range

(IQR). Two-sided Student's *t*-tests (for normally distributed data), Mann-Whitney U-tests (for

data not normally distributed), and Brown-Forsythe and Welch ANOVA (for comparisons

between ≥ 3 groups) with Dunnett's multiple comparisons test were used to determine P-values.

No statistical method was used to predetermine sample size, and data distribution was tested for

normality prior to statistical analysis and plotting.

Figure Legends

Figure 1: Emergence and patterning of nascent AT2 cells.

a scRNAseq data (GSE119228)[33] depicting embryonic, perinatal, and adult timepoints of

alveolar epithelial development. **b** Leiden Clustering analysis identifies four populations: mature

AT2s (mAT2), nascent AT2s (nAT2), distal progenitors (DP), and AT1s. **c** Stratification of the

AT2 lineage clusters across development depicting time separation of the DP, nAT2, and mAT2

cell populations. **d** Venn diagram showing number of genes detected for all clusters. **e** Dot plot

showing stage-restricted expression of selected surface markers for nAT2 and mAT2 cells,

Retnla and *Cd74* respectively. **f** e17.5 lung immunostained for nascent AT2 marker RELMα

(green), mature AT2 marker CD74 (red), and DP/AT1-restrictive luminal marker PDPN (white).

Bar, 25 μm. **g** e15.5 (left) and e16.5 (right) lungs immunostained for ASM cell marker smooth

muscle actin (SMA, red), RELMα (green), epithelial marker E-cadherin (ECAD, blue), and

PDPN (white). Close-up images for e15.5 lungs depict rare cases of single nAT2 cells (asterisk)

emerging in intermediate regions, but not at the terminal end buds (arrowheads). Bars, 100 μm

(left and right panels), 20 μm (close-ups). Mesothelium marked with white dash. **h**

Quantification of RELMα$^+$ nAT2 emergence in distal epithelial branches across development

from e15.5 to e17.5 determined by immunostaining (*n* represents branches scored for each

timepoint). ****$p \leq 0.0001$ (Brown-Forsythe and Welch ANOVA test). **i** Quantification of

RELMα$^+$ nAT2 anatomical positioning (columnar versus basally extruded) across development.

Extruded values presented as mean ± SD (*n* represents cells scored at each timepoint). **j**

Schematic showing region of AT2 Emergent Zone (green) within Sox9$^+$ distal epithelium at the

terminal stalk (blue) instead of Sox2$^+$ airway epithelium region. Around 17.5, nAT2 cell

(RELMα$^+$) basally extrude and then junction in distal epithelial branches and propagate emergent

zone proximally further. Near birth (PN0), as proximal regions further developed, emergent

zones shift to TEB.

Figure 2:

a e17.5 lung immunostained for RELMα (green), PDPN (white), and DAPI (blue). **a-i** depicts a

RELMα$^+$ nAT2 cell traversing across the interstitium from lumen A to B. Close-up panels (right)

show a nAT2 cell embedded in lumen A, at z-position +18 μm (top right), as well as in lumen B,

at z-position +24 μm (bottom right). Bar, 50 μm (left), 10 μm (close-ups; right). **b** Quantification

of nAT2 interlumenal junctioning (ILJ) events detected during development by immunostaining

(*n* represents nAT2 cells scored at each timepoint). ****$p = 4.8 \times 10^{-11}$ (one-way ANOVA, data

as mean ± SD). **c** 3D rendering immunostaining of single (upper) and double (lower)

interlumenal junctioning nAT2s for E-cadherin (green), RELMα (red), PDPN (white), and DAPI.

Lumen boundaries are marked in yellow dashes. **d** experimental schematics of timeplase

confocal microscopy on *Nkx2.1^{Cre}; Rosa26^{mTmG}* lineage labeled precision cut lung slices at distal

lung. **e** Single nAT2 junctions between two lumens in 12-hour snapshots. **f** Two nAT2s

established contact at 14.5 hr and junction between two lumens in 15-hour snapshots, contact

marked with white arrows. **g** Examples of luminal Muc1 domain length in non-junctioning (left)

and junctioning nAT2s **h** Quantification of (g) shows condensed MUC1 domain in ILJ nAT2s**.** (*n*

represents number of nAT2 cells scored at each timepoint). ****$p = 4.8 \times 10^{-11}$ (Student's two-

sided *t*-test, data as mean ± SD). **i** schematics of stages of interluminal junctioning, including

nAT2 basal extrusion and apical domain constriction **j** schematics of nAT2 extrusion and

junctioning outcomes at various interstitial positions.

Figure 3: Nascent AT2s retain fate plasticity.

a PN1 and adult ($\geq$ PN60) lungs immunostained for AT2 marker MUC1 (white), RELMα

(green), CD74 (red), and DAPI. Bar, 10 μm. **b** Timecourse quantification of (**a**) showing AT2

transitioning from nascent (RELMα$^+$) to mature (CD74$^+$). (*n* represents total AT2 cells scored at

each timepoint in experimental triplicate, values as mean $\pm$ SD). ****$p \leq 0.0001$ (one-way

ANOVA, data as mean $\pm$ SD). **c** PN1 lung immunostained for RELMα (green), CD74 (red), and

DAPI. Close-up showing both CD74$^+$ mAT2s and RELMα$^+$ nAT2s. Bar, 50 μm. **d** Alveolar fate

selection during late embryogenesis, contrasted with AT2 maturation during the early post-natal

period. **e** PAGA velocity of scRNAseq data of e15.5 and e17.5 epithelial cells. Arrow thickness

represents the relative fraction of cells with velocity from the original cluster to target cluster. A

transcriptional trajectory is observed starting from nAT2s to nAT1s (red arrow). **f** Velocity

magnitude and AT1/AT2 gene score for the nAT2 fraction from (**e**). **g** Velocity analysis of the

nAT2 fraction from (**e**) depicting AT1/AT2 score (cutoff 0.5). **h** scRNAseq data (GSE149563)[37]

of embryonic, perinatal, and adult time points of the distal epithelium. **i** Leiden clustering

including a distinct transitional AT2 cluster (tAT2 > 1; brown). Close-up depicts velocity

analysis (red dotted box), showing transition towards AT1 cluster. **j** Timepoint distribution of

cells observed within tAT2 > 1 cluster in (**i**). **k** Immunostained spheroids of e15.5 distal

progenitors (EpCAM$^+$), e18.5 nAT2s (RELMα$^+$), PN3 mAT2s (MHC-II$^+$), and adult mAT2s

(MHC-II$^+$). Spheroids were cultured on Matrigel for four days with media supplemented every

two days with FGF7 (50 ng/mL), followed by fixation and stained for PDPN (white), RAGE

(red), DAPI (blue), and AT2 marker SFTPC (green). Cells that remained distal progenitors

retained both AT1 markers (RAGE, PDPN) and the AT2 marker (SFTPC). Bars, 20 μm. **l**

Quantification of (**k**) showing percentage of cells per spheroid that differentiated into AT1 cells

determined by immunostaining (n = spheroids sampled for each condition in experimental

triplicate). ****$p \leq 0.0001$ (one-way ANOVA). **m** A revised model of the Fate Plasticity

Window depicting retained fate plasticity in nAT2 cells with the ability to differentiate into AT1

cells. All experiments were repeated at least three times.

1003

Figure 4: C/EBPα a is required to maintain but not select AT2 fate.

a A bioinformatic screen of 1,678 transcription factors or epigenetic regulators, out of which 67

are enriched along the AT2 lineage. A final list of 32 genes were selected that have known

embryonic lethal or lung defects upon knockout in mice. These 32 genes stratify into four time-

dependent categories: DP restrictive, DP-nAT2 restrictive, nAT2-mAT2 restrictive, and

constitutive. The Plasticity Window (red dotted box) spans from ~e16.5 to ~PN7. **b** Dot plot

showing expression of *Sox9* (DP restrictive; yellow), *Cebpa* and *Nupr1* (nAT2-mAT2 restrictive;

pink), *Etv5* and *Elf5* (constitutive; grey) over developmental and adult timepoints. The plasticity

window is depicted with a red dotted box. **c** e16.5 and e17.5lungs immunostained for PDPN

(white), C/EBPα (green), E-cadherin (ECAD, red), and DAPI (blue). C/EBPα$^+$ cells are marked

(arrowheads) within the proximal (P) and distal (D) regions of the epithelial branches. Bars, 100

μm. **d** Alveolar regions of PN0 control *Nkx2.1^{Cre}; Rosa26^{mTmG}; Cebpa$^{wt/fl}$* (left) and *Nkx2.1^{Cre};*

Rosa26^{mTmG}; Cebpa$^{fl/fl}$ (right) lungs immunostained for Cre reporter (mTmG) and MUC1

(white). GFP$^+$ (DP lineage) cells form both AT1 cells (flattening; "1") and AT2 cells (MUC1$^+$)

even upon *Cebpa* deletion. Bars, 50 μm. **e** Alveolar regions of adult control *LyzM^{Cre};*

Rosa26^{mTmG}; Cebpa$^{wt/fl}$ (left) and *LyzM^{Cre}; Rosa26^{mTmG}; Cebpa$^{fl/fl}$* (right) lungs immunostained

for MUC1 (white) and CD74 (blue). AT2-lineage derived GFP$^+$ cells are seen to differentiate

into flat AT1 cells (asterisk) upon *Cebpa* deletion. Bars, 100 μm. **f** Quantification of (**e**) showing

percent GFP⁺ cells that differentiate into AT1s upon *Cebpa* deletion (*n* represents number of

GFP⁺ cells sampled for the *Cebpa^{wt/fl}* and *Cebpa^{fl/fl}* conditions in experimental triplicate). *****p*

$= 1.4 \times 10^{-14}$ (Student's two-sided *t*-test, data as mean $\pm$ SD). All experiments were repeated at

least three times.

1026

Figure 5: C/EBPα blocks AT2 fate plasticity by repressing Dlk1.

a Experimental setup and timeline: Lungs were isolated 14 days post TMX (PN25) and alveolar

epithelial cells (EpCAM⁺) from control (*Sftpc^{CreER}; Tg(iSuRe-Cre); Cebpa^{wt/fl}*) and *Cebpa*

deleted (*Sftpc^{CreER}; Tg(iSuRe-Cre); Cebpa^{fl/fl}*) lungs were processed for scRNAseq. **a-i** depicts

the clusters (Louvain). **a-ii** shows gene scores across the 3 clusters in (**ai**). Upon *Cebpa* deletion,

AT2 and mAT2 scores decreases, while nAT2 score increases compared to control AT2s without

activation of AT1 score of the *Cebpa^{fl/fl}* cluster (red dashed box). **b** Experimental timeline and

TMX dosage. EdU water administered continuously for 14 days. Lungs immunostained for tdT

(red) and proliferation (detected by ClickIT in green). *Cebpa* deleted lungs with either high

(~99% AT2s) or sparse dose (~20% AT2s) TMX, depicting proliferating AT2-lineage cells

(asterisk) and flattening AT1 cells (arrowhead). Bars, 50 μm. **c** Quantification of (**b**) showing

percent tdT⁺ AT1 cells upon *Cebpa* deletion in either High Dose (fl/fl – H.D.) or Sparse (fl/fl –

Sp) deletions compared to control (wt/fl) AT2s (*n* represents tdT⁺ cells sampled for each

condition in experimental triplicate). ****$p \leq 0.0001$ (One-way ANOVA with Tukey's multiple

comparisons testing; data as mean $\pm$ SD). **d** Average gene expression values (log$_2$ fold change) of

the 22 (out of 107 total) genes that encode for extracellular proteins enriched in the *Cebpa^{fl/fl}*

AT2 cluster. Six genes (bold) are found to be upregulated in nAT2 cells and four of which

encodes secreted (green) or membrane bound (red) protein, with *Dlk1* (red, bold) having the

highest fold expression (log$_2$ fold change). **e** Alveolar regions of lungs from control *Sftpc^{CreER};*

Tg(iSuRe-Cre); Cebpa$^{wt/fl}$ (left) or *Sftpc^{CreER}; Tg(iSuRe-Cre); Cebpa$^{fl/fl}$* (right) mice, isolated 14

days post high-dose TMX (at PN25). Lungs were immunostained for DLK1 (green) and tdT

(red). Bars, 20 μm. **f** Quantification of (**e**) for percent TdT$^+$ AT2 cells that express DLK1 upon

loss of *Cebpa* in both high dose and sparse deletion conditions. Note similar percentage of tdT$^+$

AT2s express DLK1 in both high dose and sparse deletion conditions, 47.2% and 43.9%

respectively (*n* represents tdT$^+$ cells sampled for control, high dose and sparse deletion in

experimental triplicate). ****$p \leq 0.0001$, *$p = 0.015$ (one-way ANOVA, data as mean $\pm$ SD). All

experiments were repeated at least three times. **g** Bulk ATACseq tracks showing chromatin

accessibility of *Cebpa$^{wt/fl}$* (blue) and *Cebpa$^{fl/fl}$* (red) AT2 cells (*n* = 3 replicates for each

condition), showing the *Dlk1* transcriptional start site (TSS) and the intergenic differentially

methylated region (IG-DMR). The closest C/EBPα binding site to the *Dlk1* TSS is observed at a

distal trans-regulatory element (TRE, yellow), downstream of the IG-DMR, with reduced

chromatic accessibility upon loss of *Cebpa*. Further reanalysis of ChIP-seq dataset of the TRE

(red dashed box) shows elevated activity at binding sites, indicating their involvement in *Dlk1*

regulation.

1061

Figure 6: PRC2 and C/EBPα act as an incoherent FFL to generate a DLK1 pulse late in

development.

a Distal alveolar buds of e15.5 (top) and e17.5 (bottom) lungs immunostained for DLK1 (green)

and E-cadherin (ECAD, red). Close-up images show e15.5 tips have low DLK1 intensity (fire

LUT), and e17.5 tips have higher levels of DLK1 protein. Bars, 10 μm. **b** Dot plot showing RNA

expression levels of *Cebpa*, *Dlk1*, and PRC2 members genes (red) during developmental time-

course. Minimal expression of *Cebpa* and *Dlk1* observed (grey dashed box), in contrast to the

high PRC2 members genes before e16.5. **c** Epithelial branches of e16.5 and e17.5 lungs

immunostained for PDPN (white), EZH2 (green), and E-cadherin (ECAD, red). EZH2 presence

in the proximal (P) and distal (D) regions of the epithelial branches at e16.5 but absence by

e17.5. Bars, 50 μm. **d** Immunostained spheroids of e15.5 DPs (EpCAM$^+$) cultured on Matrigel

for four days with media supplemented every two days with FGF7 alone (Control, top) or FGF7

with GSK126 (10 μM), an EZH2 inhibitor (bottom). Note an increase in DLK1 (red) in cultures

with GSK126. Bars, 20 μm. **e** Quantification of (**d**) showing percent DLK1$^+$ cells per spheroid

for each condition in the culture at day 4 determined by immunostaining (*n* represents number of

spheroids for each condition in experimental triplicate). ****$p = 8.24 \times 10^{-15}$ (Student's two-

sided *t*-test, data as mean ± SD). **f** Cistrome DB Toolkit analysis at the *Cebpa* locus identifies

EZH2 (red) and the other PRC2 members (grey) to have enriched binding at this site. **g**

Immunostained spheroids of e15.5 DPs cultured as in (**d**). Note an increase in C/EBPα (red) in

GSK126 group. Bars, 20 μm. **h** Quantification of (**g**) for percent C/EBPα$^+$ cells per spheroid at

day 4 determined by immunostaining (*n* represents number of spheroids for each condition in

experimental triplicate). ****$p = 3.9 \times 10^{-36}$ (Student's two-sided *t*-test, data as mean ± SD). **i**

Distal epithelial branch of an e15.5 *Nkx2.1^{Cre}; Rosa26^{mTmG}; Ezh2$^{fl/fl}$* mouse lung, sparsely

labeling and deleting *Ezh2* in (GFP$^+$) alveolar epithelial cells. Lungs were immunostained for

C/EBPα (white) and the Cre reporter (mTmG). Bar, 10 μm. **j** Quantification of (**i**) showing

percent of DP cells in the distal alveolar buds expressing C/EBPα in lineage negative (RFP$^+$) or

lineage positive (GFP$^+$) cells (*n* represents number of distal alveolar buds sampled for both

groups). ****$p \leq 0.0001$ (Mann-Whitney U-test). **k** Interactions between EZH2/PRC2, C/EBPα,

and DLK1 that fits the model of an incoherent, type 2 Feed Forward Loop (I2-FFL). **l** Model

fitting of *Dlk1* expression from scRNAseq data of the developing lung[37], following the DP >

nAT2 > mAT2 lineage fitted onto an I2-FFL Model (red line). RMSE = 0.12 (High goodness of

fit if RMSE ≤ 0.75). **m** Spheroids of e15.5 DPs cultured as in (**d**), immunostained for PDPN

(white), SFTPC (green), RAGE (red), and DAPI. Note that treatment with GSK126 increases the

proportion of AT2 differentiation detected by an increase in SFTPC$^+$ (green) cells. Bars, 20 μm.

n Quantification of (**m**) showing a stacked plot of the proportion of DP (white), AT1 (grey) and

AT2 (black) cells determined by immunostaining. Compared to control, Both GSK126 alone and

FGF7 + GSK126 groups resulted in significantly more AT2 differentiation (*n* represents number

of spheroids for each condition in experimental triplicate). ****$p = 3.3 \times 10^{-43}$ for Control vs

GSK126, $p = 2.0 \times 10^{-36}$ for Control vs FGF7 + GSK126 (one-way ANOVA, data as

mean ± SD). **o** Schematic of DLK1 pulse upon removal of repressors reopens AT2 fate plasticity

window and convert a subset of AT2s into Notch signaling senders. In conjunction of further

lateral inhibition, the molecular circuit in (**k**) contributes to AT1/AT2 patterning in lung

development. All experiments were repeated at least three times.

1105

Figure 7: C/EBPα downregulation is required to re-access mutually exclusive Defender and

Regenerative states in response to injury.

a Experimental design (top). 5 days after AAV-SPC-Cre (Control; left) or AAV-SPC-Cebpa-

T2A-Cre (right) instillation into lungs of adult *Rosa26*^{mTmG} mice to induce constitutive GFP-Cre

expression and overdrive expression of *Cebpa* in AT2 cells, lungs were injured by Fiin1 (14.75

mg/kg) intratracheal instillation (i.t.). Lungs were isolated and immunostained for Cre reporter

(GFP) and AT1 markers 14 days post injury (dpi). Immunostained images (bottom) show GFP$^+$

AT1 cells (asterisk) in the control, but little to none when C/EBPα is overexpressed. Bars, 100

μm. **b** Quantification of (**a**) for percent GFP⁺ cells that differentiated into AT1 cells (*n* represents

GFP⁺ cells sampled for each condition in experimental triplicate). ****$p = 6.5 \times 10^{-10}$ (Student's

two-sided *t*-test, data as mean ± SD). **c** Alveolar regions of an adult mouse lung 16 days post 1X

Bleomycin (0.25 mg/kg, i.t.) injury, immunostained for E-cadherin (ECAD, green), C/EBPα

(red), and DAPI (blue). Injured (right) sites show simplified alveolar structures with reduced E-

cadherin expression compared to uninjured sites (left). C/EBPα breakout (fire LUT) below

depicts reduced C/EBPα expression in injured versus uninjured sites. Bars, 100 μm. **d**

Quantification of (**c**) for mean C/EBPα intensity in AT2 cells within uninjured or injured regions

(*n* represents cells scored for each condition in experimental triplicate). ****$p \leq 0.0001$ (Mann-

Whitney U-test). **e** Dot plot from a scRNAseq dataset (GSE141259)[20] of 1X bleomycin-induced

injury in AT2 cells, showing relative gene expression in vehicle (control) and various timepoints

post 1X Bleomycin injury. Spikes in expression of DATP state associated genes (*Ddit3*, *Atf3*,

Atf4, and *Ezh2*) are marked (dashed red box), whereas a temporary reduction of *Cebpa* is

observed upon injury. The DATP gene score (red, see Suppl. Table 1) is also plotted on this

timecourse, showing an activation of a DATP associated state upon injury, which resolves over

time. **f** Experimental design for lineage labeling AT2 cells at PN11 in *Sftpc^CreER; Rosa26^mTmG*

mice, followed by administration of corn oil (vehicle) or CCT020312 (2 mg/kg, intraperitoneal),

a Ddit3 agonist, 1 day post TMX, and every 4 days (4 doses total). Lungs were isolated 14 days

post TMX (PN25) and immunostained for Cre reported (mTmG) and MUC1 (white). Upon

treatment with CCT020312, an increase in GFP⁺ AT1 cells (asterisk) is observed. Bars, 50 μm. **g**

Quantification of (**f**) showing percent GFP⁺ AT1 cells for either vehicle or CCT020312

conditions (*n* represents GFP⁺ cells sampled for each condition). ****$p = 2.1 \times 10^{-11}$ (Student's

two-sided *t*-test, data as mean ± SD). **h** Leiden clustering of our scRNAseq dataset reveals two

distinct *Cebpa*$^{fl/fl}$ AT2 clusters: AT2^{DLK1+} (green) and AT2^{Def} (red). Velocity analysis (arrows)

show AT2^{DLK1+} cells going in two separate trajectories, towards the AT1 and AT2^{Def} clusters. **i**

Gene Ontology (GO) analysis of the AT2^{Def} cluster shows enriched pathogen defense response

(red) and interferon signaling (black) gene programs. **j** DATP- and Defender-associated gene

score co-expression shows a mutually exclusive pattern in *Cebpa*$^{fl/fl}$ cells. **k** SPRING plot of the

four clusters observed in (**h**), showing AT2^{Def} state (red) in between the control AT2 (orange)

and AT2^{DLK1+} (green) clusters. **l** SPRING plot of the DATP-associated gene score showing a

high DATP score between the control AT2 (middle) and AT1 (left) clusters. **m** Immunostained

spheroids of PN3 mAT2s cultured on Matrigel for four days with media supplemented every two

days with FGF7 alone (Control, top) or with interferon-γ (20 ng/mL) (bottom). Note a decrease

in AT1 cells (asterisk) in cultures treated with interferon-γ. Bars, 20 μm. **n** Quantification of (**m**)

showing percent of cells per spheroid that differentiated into AT1s for each condition (*n*

represents spheroids sampled per condition). ****$p = 1.2 \times 10^{-8}$ (Student's two-sided *t*-test, data

as mean ± SD). **o** Schematic showing AT2 oscillation between AT2^{Def} (purple) and AT2^{DATP}

(yellow) states, negatively regulated by C/EBPα. Upon AT2 loss (grey AT2s) due to infection,

remaining AT2s can oscillate (left) between the two states resulting in proper infection response

and regeneration. The inability to access either the AT2^{Def} or AT2^{DATP} states would result in

impaired pathogen response or impaired repair and regeneration respectively. All experiments

were repeated at least three times.

1156

Supplementary Figure 1: Dynamics of nascent AT2 state and emergence

a Cluster correlation analyzing the strength of correlation between each of the four clusters

detected: AT1, DP, nAT2, and mAT2. Pearson Correlation > 0.5 indicates a strong, positive

correlation. **b** Coassignment probability for the four clusters shown in (**a**), measuring the

probability of merging two clusters. Note that the nAT2 and mAT2 clusters do not have a

significant coassignment probability (asterisk), indicating they are distinct, stable cell

populations. Low coassignment probability (< 0.5) represents a stable cluster with respect to one

another. **c** scRNAseq expression for genes enriched for the clusters identified within the AT2

lineage. *Cd74* and *Retnla* are highlighted (bold) as being surface markers chosen for mature and

nascent AT2s respectively. **d** feature plot of *Retnla* and *Cd74* in mouse development timecourse

dataset from Cohen et al., showing high and exclusive expression patterns of the nascent and

mature AT2 cell markers. **e** e17.5 lung immunostained for PDPN (white), distal progenitor

marker SOX9 (green), nAT2 marker RELMα (red), and DAPI (blue). Nascent AT2 cells are

marked (arrowhead) to show them as RELMα^{+} and SOX9^{low}. **f** Quantification of nAT2 clustering

observed as either singlets, doublets, or in clusters of ≥ 3 across embryonic timepoints from

e15.5 to PN0 ($n = 30$ nAT2 clusters scored at e15.5, 155 at e16.5, 144 at e17.5, 228 at e18.5, 140

at e19.5, and 96 at P0). **g** flow cytometry of EpCAM^{+} nascent AT2s from *WT embryonic* mouse

lung, show moderate RELMα positivity, and low labeling efficiency by CD74, suggesting that

nascent AT2s can be labeled and sort by RELMα. **h** flow cytometry of tdT^{+} mature AT2s from

*Sftpc^{CreER}; Tg(iSuRe-Cre)*mouse lung, show little RELMα positivity, moderate labeling

efficiency by CD74, and high labeling efficiency by MHC-II, suggesting that MHC-II is more

effective as cell isolating marker comparing to CD74.

1179

Supplementary Figure 2: Spatial and Temporal Characterization of junctioning nAT2s

a Cross-section of e17.5, e18.5, or PN0 distal lungs slices immunostained for RELMα (red),

DAPI (blue), and markers of interests (green), including C/EBPα, EZH2, DLK1, SFTPC,

LAMP1, MUC1, PDPN, RAGE, and HOP. **b** Quantification of portion of nAT2 showing their

high positivity in AT2 character and fate plasticity and their absence of AT1 character. **c**

Additional 3D renderings on immunostaining of double interlumenal junctioning nAT2s for E-

cadherin (ECAD) or RAGE (green), RELMα (red), PDPN or MUC1 (white), and DAPI (blue).

Lumen boundaries are marked in yellow dashes. **d** Additional snapshots timeplase confocal

microscopy on $Nkx2.1^{Cre}$; $Rosa26^{mTmG}$ lineage labeled precision cut lung slices at distal lung

showing dynamics interlumenal junctioning.

1190

Supplementary Figure 3: Details on state dependent plasticity and death sensitivity

a Cross-section of an airway (dashed yellow) of e17.5 (left) and PN1 (right) lungs

immunostained for RELMα (green), CD74 (red) and DAPI (blue). Airway region at PN1 shows

presence of RELMα in the airway epithelium, but the absence at e17.5 confirms its specificity to

nAT2s during the AT2 fate selection window. Bars, 25 μm. **b** Proximal and distal epithelium of a

PN1 lung immunostained for RELMα (green), CD74 (red) and DAPI (blue), showing salt-and-

pepper pattern of AT2 maturation. Bar, 50 μm. **c** scRNAseq UMAP of distal lung epithelium at

e15.5 (blue) and e17.5 (orange) timepoints. **d** Merged clusters of scRNAseq dataset in (**c**),

showing three e15.5 distal epithelium (DE) clusters, and three e17.5 clusters (DP, nAT2, and

nAT1). **e** Cluster correlation of the six clusters observed in (**d**). No significant correlation

observed between either e15.5 DE cluster with the nAT2 or nAT1 clusters at e17.5. A weak

correlation (0.42, dashed box) observed between the e15.5 DE3 and e17.5 DP clusters. **f**

Pseudotime trajectory analysis showing e15.5 cells are the primary root cells of origin (left) and

e17.5 AT1 cells are endpoint cells (right). **g** Velocity trajectories showing transition from e15.5

DE clusters to e17.5 DP cells. Subsequent trajectories lead either directly to AT1 cells or

indirectly, transitioning through the nAT2 cluster. **h** BH3 death sensitivity assay of e15.5 DP,

e18.5 nAT2, and $\geq$ PN60 mAT2 cells, showing percent cytochrome c release upon treatment

with activator or sensitizer BH3 peptides. A higher cytochrome c release percentage indicates

increased sensitivity to BH3-mediated apoptosis. Alam and DFNA5 are positive controls; DMSO

is the vehicle control and PUMA 2A is an inactive peptide negative control. Cytochrome c

release (%) values are presented as mean $\pm$ SD. ****$p \leq 0.0001$, ***$p \leq 0.001$ (one-way

ANOVA, comparing DP and mAT2 cells).

1213

Supplementary Figure 4: Bioinformatic screen for regulators of AT2 plasticity and

perinatal *Cebpa* deletion.

a Heat map showing expression of all 32 genes from Figure 4a. Cells (columns) are arranged in

developmental pseudotime (DP, nAT2, mAT2; left to right) along the AT2 lineage. Note that the

"mAT2 (TM)" is from the Tabula Muris scRNAseq dataset (GSE109774)[41], which sequences

AT2 cells more deeply. Genes form four patterns: constitutive (grey), DP restrictive (blue), DP-

nAT2 restrictive (orange), and nAT2-mAT2 restrictive (pink). **b** Dot plot of the transcription

factors from (**a**) that are expressed within the plasticity window (red dotted box), showing

expression at different developmental and adult timepoints. **c** Alveolar regions of PN0 control

Nkx2.1^{Cre}; Rosa26^{mTmG}; Cebpa$^{wt/fl}$ (left) and *Nkx2.1^{Cre}; Rosa26^{mTmG}; Cebpa$^{fl/fl}$* (right) lungs

immunostained for MUC1 (white) and RAGE (red). GFP$^+$ (DP lineage) cells form both AT1

cells (flattening; asterisk "*") and AT2 cells (MUC1$^+$) even upon *Cebpa* deletion. Bars, 25 μm. **d**

Experimental design (top) and immunostains (bottom) for simultaneous *Cebpa* deletion and

lineage labeling AT2 cells at PN11 via intraperitoneal (i.p.) tamoxifen (TMX) induction. Lungs

were isolated 14 days post TMX (PN25) and immunostained for MUC1 (white) and RAGE

(red). Alveolar regions of control *Sftpc*^{*CreER*}*; Rosa26*^{*mTmG*}*; Cebpa*^{*wt/fl*} (left) and *Sftpc*^{*CreER*}*;*

Rosa26^{*mTmG*}*; Cebpa*^{*fl/fl*} (right) lungs immunostained as indicated, showing GFP$^+$/RAGE$^+$ AT1

cells (asterisk) upon *Cebpa* deletion. Close up (bottom left) panels for each condition show a

GFP$^+$/MUC1$^+$ AT2 (control; left) or GFP$^+$/RAGE$^+$ AT1 cell (*Cebpa* deletion; right). Bars, 100

μm. **e** Quantification of (**d**) showing percent GFP$^+$ cells that differentiated into AT1s upon

Cebpa deletion (n = 2856 GFP$^+$ cells sampled for control and 2105 for *Cebpa* deletion in

experimental triplicate). ****p = 6.4×10^{-16} (Student's two-sided *t*-test, data as mean $\pm$ SD). All

experiments were repeated at least three times. **f** Alveolar region of *Sftpc*^{*CreER*}*; Tg(iSuRe-Cre);*

Cebpa^{*fl/fl*} mouse lung immunostained for RAGE (white), RELMα (green) and Cre reporter (tdT).

Lineage labeled RAGE$^+$ (arrowheads) and RELMα$^+$ (asterisk) cells are mutually exclusive from

one another, showing the two distinct states upon loss of *Cebpa*.

1240

Supplementary Figure 5: *Cebpa* deletion and gene score validation.

a scRNAseq UMAP close-up of AT1 cluster (Figure 5a-i) depicting which cells are lineage

positive (black) that converted to AT1s upon deletion of *Cebpa*. **b** Volcano plot showing

downregulated (left) and upregulated (right) genes in AT2 cells upon *Cebpa* deletion. Genes are

considered to have significant down/upregulation if $-\text{Log}_{10} P > 100$, and Log_2 fold change > 1 or

< -1 (red). **c** AT1 and AT2 gene score co-expression on scRNAseq UMAP of the developing

lung[33] to assess changes in either AT2 or AT1 related genes. **d** nAT2 and mAT2 gene score co-

expression on scRNAseq UMAP of the developing lung[33] to confirm restrictive expression of

genes in either AT2 state. **e** AT1 and AT2 gene score (see Suppl. Table 1) co-expression on

scRNAseq UMAP, showing retained AT2 gene score, and lack of an AT1 score upon loss of

Cebpa in AT2 cells. **f** Alveolar regions of a PN25 *Sftpc*^{*CreER*}*; Tg(iSuRe-Cre)*mouse lung, 14 days

after Cre induction with a sparse dose of TMX. Lungs were immunostained for MUC1 (green),

tdT (red), and DAPI (blue). Bar, 25 μm. **g** Quantification of (**f**) for percent tdT$^+$ labeling of AT2

cells in either sparse or high doses of TMX (n = 1258 AT2 cells sampled for high dose TMX and

1670 for the sparse dose TMX). ****p = 1.73 × 10^{-9} (Student's two-sided t-test, data as

mean ± SD). **h** Quantification of Figure 5b, showing percent tdT$^+$/EdU$^+$ proliferating cells upon

Cebpa deletion in either High Dose or Sparse conditions. Note a significant reduction in

proliferation upon sparse *Cebpa* deletion (n = 2358 tdT$^+$ cells sampled for the high dose deletion

and 1115 for the sparse deletion in experimental triplicate). ****p = 2.47 × 10^{-9} (Student's two-

sided t-test, data as mean ± SD). **i** Dot plot showing expression of the six genes (encoding for

extracellular proteins) upregulated in the *Cebpa$^{fl/fl}$* cluster, also enriched in nAT2 cells (Figure

5d). **j** *Dlk1* gene expression on a scRNAseq developmental dataset confirming DLk1 expression

in mAT2 cells. **k** Quantification of Lineage labeled organoids generated from AT2s isolated

from *Sftpc^{CreER}; Tg(iSuRe-Cre); Cebpa$^{wt/fl}$* and *Sftpc^{CreER}; Tg(iSuRe-Cre); Cebpa$^{fl/fl}$* murine lung,

showing higher percent DLK1$^+$ AT2s cells upon *Cebpa* deletion (n = 435 tdT$^+$ cells sampled for

the *Cebpa* deletion and 435 for control in experimental triplicate). ****p = 0.0388 (Student's

two-sided t-test). **l** Experimental design for labeling and deleting *Cebpa* in AT2 cells of *Sftpc-*

CreER;Tg(iSuRe-Cre);Cebpa$^{wt/fl}$ and *Sftpc-CreER;Tg(iSuRe-Cre);Cebpa$^{fl/fl}$* mice at PN11,

isolating cells 14 days post TMX (at PN25) for scRNAseq and bulk ATACseq analyses. **m**

Hierarchical clustering of sample binding sites shows high correlation between conditions in the

ATAC-seq data. **n** Venn diagram of binding site overlaps in the ATAC-seq data between two

programs for differential accessibility analysis – edgeR and DESeq2. **o** Differentially expressed

gene (DEG) overlaps between RNA-seq and ATAC-seq data for the *Cebpa$^{fl/fl}$* vs *Cebpa$^{wt/fl}$* AT2

cells. **p** Feature distribution from the ATAC-seq data detailing the genomic features associated

with down- and upregulated binding sites upon loss of *Cebpa*. **q** HOMER analysis of the ATAC-

seq data shows motifs enriched in the *Cebpa*$^{wt/fl}$ (red) and *Cebpa*$^{fl/fl}$ (blue). AP-1 and Yap/Taz

interacting factors are highlighted via shape. **r** HOMER motifs and associated *p*-values highlight

a role of AP1, TEAD, and NKX2.1 signaling in both the WT and KO conditions. **s** Schematic

detailing the potential role of C/EBPα interacting with AP1 and NKX2.1 in AT2 (left) and AT1

(right) cells.

1281

Supplementary Figure 6: Embryonic expression of C/EBPα and DLK1.

a Distal epithelial branch of an e17.5 lung immunostained for DLK1 (green), PDPN (red), and

DAPI (blue). Bar, 10 μm. **b** Distal epithelial branch of an e15.5 lung immunostained for C/EBPα

(green), PDPN (red), E-cadherin (ECAD, white), and DAPI (blue). No C/EBPα$^+$ cells are

detected in the distal epithelial branches at this point in development. Bar, 10 μm. **c** *Cebpa*

(green) and *Ezh2* (red) co-expression on scRNAseq UMAP of the developing lung[33] shows

mutually exclusive expression. **d** Analysis of ENCODE epigenetic dataset mapping of the PRC2

modification H3K27me3 across developmental timepoints (e14.5 – P0) at the *Cebpa* locus. **e**

Max normalized gene expression for *Cebpa* (red), *Dlk1* (green) and *Ezh2* (blue) during

development from scRNAseq data of the developing lung[37], showing the relation of the

components forming the I2-FFL that generates a pulse of *Dlk1*.**f** Dot plot showing expression of

the Notch signaling pathway genes (ligands and its downstream activation) have distinct pattern

as nascent cell fates are specified. **g** schematics of our working model how a Dlk1 pulse

generator regulate AT2's accessibility to fate plasticity via Notch signaling.

1296

Supplementary Figure 7: AT2 defensive state and interferon signaling.

a Experimental design (top). 5 days after AAV-SPC-Cre (Control; left) or AAV-SPC-Cebpa-

T2A-Cre (right) instillation into lungs of adult $Rosa26^{mTmG}$ mice to induce constitutive GFP-Cre

expression and overdrive expression of *Cebpa* in AT2 cells, lungs were injured by BHT (225

mg/kg) intraperitoneal injection (i.p.). Lungs were isolated and immunostained for Cre reporter

(GFP) and AT1 markers 14 days post injury (dpi). Immunostained images (bottom) show GFP$^+$

AT1 cells (asterisk) in the control, but little to none when C/EBPα is overexpressed. Bars, 50

μm. **b** Quantification of (**a**) for percent GFP$^+$ cells that differentiated into AT1 cells ($n = 557$

GFP$^+$ cells sampled for the control condition and 631 for the *Cebpa* overdrive in experimental

triplicate). ****$p = 6.5 \times 10^{-10}$ (Student's two-sided *t*-test, data as mean ± SD). **c** Immunostained

adult lung showing GFP-labeling efficiency of AT2 cells (MUC1; white) by AAV-SPC-Cre

(Figure 7a). Bar, 100 μm. **d** Dot plot showing interferon receptors (left) and interferon-γ target

genes (right) for the AT2 clusters from the scRNAseq analysis. The AT2^{Def} cluster is highly

enriched in both receptors and target genes. Interferon-γ specific receptor genes (dashed red box)

are upregulated in the AT2^{Def} state upon loss of *Cebpa*. **e** Dot plot showing expression of DATP

and Defender gene scores (top), along with expression of *Cebpa*, *Ezh2*, and DATP-associated

genes (*Atf3*, *Atf4*, *Ddit3*) (bottom) in the clusters identified from the bleomycin injury timecourse

scRNAseq dataset[20]. **f** DATP- and Defender-associated gene score co-expression shows a

mutually exclusive pattern in four other lung injury model besides Cebpa deletion model (Fig 7j).

1316

Data Availability

The authors confirm that all relevant data are available in the paper or its Supplementary

Information files. The scRNAseq datasets are available in the GEO repository under accession

numbers "GSE119228", "GSE113320", "GSE109774", and "GSE149563" for developing and

adult mouse lung cells, "GSE295269" for CEBP/α knockout mouse lung epithelial cells,

"GSE141259" for adult bleomycin injury, "GSE218666" for AT1 injury, "GSE184384" for

influenza injury, and "GSE113049" for LPS injury models. Source Data are available for this

paper.

1325

1326 **Code Availability**

1327 Code to reproduce the analyses described in this manuscript can be accessed at

1328 https://github.com/JimmyMayoGit/Sawhney_2025

1329

Acknowledgments

The authors thank members of the Brownfield laboratory for helpful discussions and critical

reading of the manuscript. We thank D. Tenen (*Cebpa*$^{fl/fl}$) for the transgenic mouse strain;

Hannah Golding for performing organotypic culture experiments; and Gamaliel Taengwa for

data quantification. This work was supported by R00HL127267, R01HL171056, and a Pilot

Award from Mayo Clinic's Center for Biomedical Discovery (D.G.B).

1336

Author Contributions

ASS, BJD, and DGB conceived the study and designed the experiments. ASS, BJD, JC, DG,

GA, AO, HH, LS, XQ, SRH, ASU performed the experiments. ASS, BJD, and AO analyzed and

interpreted the data. JC, AH, EL, and DGB performed the scRNAseq analysis. DG and ZW

performed the ATACseq analysis. RW and SD performed bleomycin injury experiments, and

KAS and ZI performed BH3 profiling experiments. DGB, ASS, and BJD wrote the paper.

1343

1344   **Ethics Declarations**

1345   **Competing interests**

1346   The authors have no competing interests to declare.

1347

1348   **References:**

1349   1.   Barnes PJ*, et al.* Chronic obstructive pulmonary disease. *Nat Rev Dis Primers* **1**, 15076
1350   (2015).

1351
1352   2.   Martinez FJ*, et al.* Idiopathic pulmonary fibrosis. *Nature Rev Dis Primers* **3**, 17074 (2017).

1353
1354   3.   Siegel R, Naishadham D, Jemal A. Cancer statistics, 2013. *CA Cancer J Clin* **63**, 11-30
1355   (2013).

1356
1357   4.   Desai TJ, Brownfield DG, Krasnow MA. Alveolar progenitor and stem cells in lung
1358   development, renewal and cancer. *Nature* **507**, 190-194 (2014).

1359
1360   5.   Franks TJ*, et al.* Lung pathology of severe acute respiratory syndrome (SARS): a study of
1361   8 autopsy cases from Singapore. *Hum Pathol* **34**, 743-748 (2003).

1362
1363   6.   Hwang DM, Chamberlain DW, Poutanen SM, Low DE, Asa SL, Butany J. Pulmonary
1364   pathology of severe acute respiratory syndrome in Toronto. *Mod Pathol* **18**, 1-10 (2005).

1365
1366   7.   Tian S, Hu W, Niu L, Liu H, Xu H, Xiao SY. Pulmonary pathology of early-phase 2019 novel
1367   coronavirus (COVID-19) pneumonia in two patients with lung cancer. *J Thorac Oncol* **15**,
1368   700-704 (2020).

1369
1370   8.   Rawlins EL, Clark CP, Xue Y, Hogan BL. The Id2+ distal tip lung epithelium contains
1371   individual multipotent embryonic progenitor cells. *Development* **136**, 3741-3745 (2009).

1372
1373   9.   Treutlein B*, et al.* Reconstructing lineage hierarchies of the distal lung epithelium using
1374   single-cell RNA-seq. *Nature* **509**, 371-375 (2014).

1375
1376   10.   Brownfield DG, de Arce AD, Ghelfi E, Gillich A, Desai TJ, Krasnow MA. Alveolar cell fate
1377   selection and lifelong maintenance of AT2 cells by FGF signaling. *Nat Commun* **13**, 7137
1378   (2022).

1379
1380   11.   Zacharias WJ*, et al.* Regeneration of the lung alveolus by an evolutionarily conserved
1381   epithelial progenitor. *Nature* **555**, 251-255 (2018).

1382

1383 12. Frank DB*, et al.* Early lineage specification defines alveolar epithelial ontogeny in the
1384   murine lung. *Proc Natl Acad Sci U S A* **116**, 4362-4371 (2019).

1385

1386 13. Vila Ellis L, Chen J. A cell-centric view of lung alveologenesis. *Dev Dyn* **250**, 482-496
1387   (2021).

1388

1389 14. Weibel ER. On the tricks alveolar epithelial cells play to make a good lung. *Am J Respir*
1390   *Crit Care Med* **191**, 504-513 (2015).

1391

1392 15. Shenoy AT*, et al.* Antigen presentation by lung epithelial cells directs CD4+ TRM cell
1393   function and regulates barrier immunity. *Nature Communications* **12**, 5834 (2021).

1394

1395 16. Choi J*, et al.* Inflammatory Signals Induce AT2 Cell-Derived Damage-Associated Transient
1396   Progenitors that Mediate Alveolar Regeneration. *Cell Stem Cell* **27**, 366-382 e367 (2020).

1397

1398 17. Malik A*, et al.* Epithelial IFNγ signalling and compartmentalized antigen presentation
1399   orchestrate gut immunity. *Nature* **623**, 1044-1052 (2023).

1400

1401 18. Gessani S, Conti L, Del Cornò M, Belardelli F. Type I interferons as regulators of human
1402   antigen presenting cell functions. *Toxins (Basel)* **6**, 1696-1723 (2014).

1403

1404 19. Rock JR*, et al.* Multiple stromal populations contribute to pulmonary fibrosis without
1405   evidence for epithelial to mesenchymal transition. *Proc Natl Acad Sci U S A* **108**, E1475-
1406   1483 (2011).

1407

1408 20. Strunz M*, et al.* Alveolar regeneration through a Krt8+ transitional stem cell state that
1409   persists in human lung fibrosis. *Nat Commun* **11**, 3559 (2020).

1410

1411 21. Barkauskas CE*, et al.* Type 2 alveolar cells are stem cells in adult lung. *The Journal of*
1412   *clinical investigation* **123**, 3025-3036 (2013).

1413

1414 22. Nabhan AN, Brownfield DG, Harbury PB, Krasnow MA, Desai TJ. Single-cell Wnt signaling
1415   niches maintain stemness of alveolar type 2 cells. *Science* **359**, 1118-1123 (2018).

1416

1417   23.   Finn J*, et al.* Dlk1-Mediated Temporal Regulation of Notch Signaling Is Required for
1418   Differentiation of Alveolar Type II to Type I Cells during Repair. *Cell Rep* **26**, 2942-2954
1419   e2945 (2019).

1420
1421   24.   Major J*, et al.* Type I and III interferons disrupt lung epithelial repair during recovery
1422   from viral infection. *Science* **369**, 712-717 (2020).

1423
1424   25.   Jiang P*, et al.* Ineffectual Type 2-to-Type 1 Alveolar Epithelial Cell Differentiation in
1425   Idiopathic Pulmonary Fibrosis: Persistence of the KRT8(hi) Transitional State. *Am J Respir*
1426   *Crit Care Med* **201**, 1443-1447 (2020).

1427
1428   26.   Lourenco AR*, et al.* C/EBPa is crucial determinant of epithelial maintenance by
1429   preventing epithelial-to-mesenchymal transition. *Nat Commun* **11**, 785 (2020).

1430
1431   27.   Suh HC, Gooya J, Renn K, Friedman AD, Johnson PF, Keller JR. C/EBPalpha determines
1432   hematopoietic cell fate in multipotential progenitor cells by inhibiting erythroid
1433   differentiation and inducing myeloid differentiation. *Blood* **107**, 4308-4316 (2006).

1434
1435   28.   Baladron V*, et al.* dlk acts as a negative regulator of Notch1 activation through
1436   interactions with specific EGF-like repeats. *Exp Cell Res* **303**, 343-359 (2005).

1437
1438   29.   Grassi ES, Pietras A. Emerging Roles of DLK1 in the Stem Cell Niche and Cancer
1439   Stemness. *J Histochem Cytochem* **70**, 17-28 (2022).

1440
1441   30.   Yen YP*, et al.* Dlk1-Dio3 locus-derived lncRNAs perpetuate postmitotic motor neuron cell
1442   fate and subtype identity. *Elife* **7**, (2018).

1443
1444   31.   Zhao J*, et al.* PPARgamma and C/EBPalpha enable adipocyte differentiation upon
1445   inhibition of histone methyltransferase PRC2 in malignant tumors. *J Biol Chem* **300**,
1446   107765 (2024).

1447
1448   32.   Auyeung VC*, et al.* IRE1alpha drives lung epithelial progenitor dysfunction to establish a
1449   niche for pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol* **322**, L564-L580 (2022).

1450
1451   33.   Cohen M*, et al.* Lung single-cell signaling interaction map reveals basophil role in
1452   macrophage imprinting. *Cell* **175**, 1031-1044 e1018 (2018).

1453

1454   34.   Gillich A, St. Julien KR, Brownfield DG, Travaglini KJ, Metzger RJ, Krasnow MA. Alveoli
1455   form directly by budding led by a single epithelial cell. *bioRxiv*, 2021.2012.2025.474174
1456   (2021).

1457

1458   35.   Blanco MA*, et al.* Chromatin-state barriers enforce an irreversible mammalian cell fate
1459   decision. *Cell Reports* **37**, 109967 (2021).

1460

1461   36.   Hurley K*, et al.* Reconstructed Single-Cell Fate Trajectories Define Lineage Plasticity
1462   Windows during Differentiation of Human PSC-Derived Distal Lung Progenitors. *Cell*
1463   *Stem Cell* **26**, 593-608.e598 (2020).

1464

1465   37.   Zepp JA*, et al.* Genomic, epigenomic, and biophysical cues controlling the emergence of
1466   the lung alveolus. *Science* **371**, (2021).

1467

1468   38.   Sarosiek KA*, et al.* Developmental Regulation of Mitochondrial Apoptosis by c-Myc
1469   Governs Age- and Tissue-Specific Sensitivity to Cancer Therapeutics. *Cancer Cell* **31**, 142-
1470   156 (2017).

1471

1472   39.   Singh R*, et al.* Radiotherapy-Induced Neurocognitive Impairment Is Driven by
1473   Heightened Apoptotic Priming in Early Life and Prevented by Blocking BAX. *Cancer Res*
1474   **83**, 3442-3461 (2023).

1475

1476   40.   Basundra R*, et al.* Constitutive High Expression of NOXA Sensitizes Human Embryonic
1477   Stem Cells for Rapid Cell Death. *Stem Cells* **40**, 49-58 (2022).

1478

1479   41.   A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. *Nature* **583**,
1480   590-595 (2020).

1481

1482   42.   Zhang Z*, et al.* Transcription factor Etv5 is essential for the maintenance of alveolar type
1483   II cells. *Proc Natl Acad Sci U S A* **114**, 3903-3908 (2017).

1484

1485   43.   Metzger DE, Stahlman MT, Shannon JM. Misexpression of ELF5 disrupts lung branching
1486   and inhibits epithelial differentiation. *Dev Biol* **320**, 149-160 (2008).

1487

1488   44.   Santofimia-Castaño P*, et al.* Inactivation of NUPR1 promotes cell death by coupling ER-
1489   stress responses with necrosis. *Sci Rep* **8**, 16999 (2018).

45. Liu J*, et al.* NUPR1 is a critical repressor of ferroptosis. *Nature Communications* **12**, 647 (2021).

46. Basseres DS*, et al.* Respiratory failure due to differentiation arrest and expansion of alveolar cells following lung-specific loss of the transcription factor C/EBPalpha in mice. *Mol Cell Biol* **26**, 1109-1123 (2006).

47. Martis PC, Whitsett JA, Xu Y, Perl AK, Wan H, Ikegami M. C/EBPalpha is required for lung maturation at birth. *Development* **133**, 1155-1164 (2006).

48. Morimoto M, Kopan R. rtTA toxicity limits the usefulness of the SP-C-rtTA transgenic mouse. *Dev Biol* **325**, 171-178 (2009).

49. Fernández-Chacón M*, et al.* iSuRe-Cre is a genetic tool to reliably induce and report Cre-dependent genetic modifications. *Nat Commun* **10**, 2262 (2019).

50. Germani F, Bergantinos C, Johnston LA. Mosaic Analysis in Drosophila. *Genetics* **208**, 473-490 (2018).

51. Burke R, Basler K. Dpp receptors are autonomously required for cell proliferation in the entire developing Drosophila wing. *Development* **122**, 2261-2269 (1996).

52. Hassan D, Chen J. CEBPA restricts alveolar type 2 cell plasticity during development and injury-repair. *Nat Commun* **15**, 4148 (2024).

53. Nueda ML, Baladrón V, Sánchez-Solana B, Ballesteros MA, Laborda J. The EGF-like protein dlk1 inhibits notch signaling and potentiates adipogenesis of mesenchymal cells. *J Mol Biol* **367**, 1281-1293 (2007).

54. Ramji DP, Foka P. CCAAT/enhancer-binding proteins: structure, function and regulation. *Biochem J* **365**, 561-575 (2002).

55. Little DR, Lynch AM, Yan Y, Akiyama H, Kimura S, Chen J. Differential chromatin binding of the lung lineage transcription factor NKX2-1 resolves opposing murine alveolar cell fates in vivo. *Nat Commun* **12**, 2509 (2021).

1526

1527   56.   Aronson BE*, et al.* A bipartite element with allele-specific functions safeguards DNA
1528   methylation imprints at the Dlk1-Dio3 locus. *Dev Cell* **56**, 3052-3065.e3055 (2021).

1529

1530   57.   Hagan JP, O'Neill BL, Stewart CL, Kozlov SV, Croce CM. At least ten genes define the
1531   imprinted Dlk1-Dio3 cluster on mouse chromosome 12qF1. *PLoS One* **4**, e4352 (2009).

1532

1533   58.   Visel A, Thaller C, Eichele G. GenePaint.org: an atlas of gene expression patterns in the
1534   mouse embryo. *Nucleic Acids Res* **32**, D552-556 (2004).

1535

1536   59.   Sanli I*, et al.* Meg3 Non-coding RNA Expression Controls Imprinting by Preventing
1537   Transcriptional Upregulation in cis. *Cell Reports* **23**, 337-348 (2018).

1538

1539   60.   He Y*, et al.* Spatiotemporal DNA methylome dynamics of the developing mouse fetus.
1540   *Nature* **583**, 752-759 (2020).

1541

1542   61.   Zheng R*, et al.* Cistrome Data Browser: expanded datasets and new tools for gene
1543   regulatory analysis. *Nucleic Acids Research* **47**, D729-D735 (2018).

1544

1545   62.   Alon U. *An introduction to systems biology : design principles of biological circuits*,
1546   Second edition. edn. CRC Press, Taylor & Francis Group (2020).

1547

1548   63.   Chitnis AB. The role of Notch in lateral inhibition and cell fate specification. *Mol Cell*
1549   *Neurosci* **6**, 311-321 (1995).

1550

1551   64.   Ochi S, Imaizumi Y, Shimojo H, Miyachi H, Kageyama R. Oscillatory expression of Hes1
1552   regulates cell proliferation and neuronal differentiation in the embryonic brain.
1553   *Development* **147**, (2020).

1554

1555   65.   Nueda ML, Naranjo AI, Baladron V, Laborda J. The proteins DLK1 and DLK2 modulate
1556   NOTCH1-dependent proliferation and oncogenic potential of human SK-MEL-2
1557   melanoma cells. *Biochim Biophys Acta* **1843**, 2674-2684 (2014).

1558

1559   66.   Milara J*, et al.* Sphingosine-1-phosphate is increased in patients with idiopathic
1560   pulmonary fibrosis and mediates epithelial to mesenchymal transition. *Thorax* **67**, 147-
1561   156 (2012).

67. Kobayashi Y, *et al.* Persistence of a regeneration-associated, transitional alveolar epithelial cell state in pulmonary fibrosis. *Nat Cell Biol* **22**, 934-946 (2020).

68. Jiang HY, *et al.* Activating transcription factor 3 is integral to the eukaryotic initiation factor 2 kinase stress response. *Mol Cell Biol* **24**, 1365-1377 (2004).

69. Ma Y, Brewer JW, Diehl JA, Hendershot LM. Two distinct stress signaling pathways converge upon the CHOP promoter during the mammalian unfolded protein response. *J Mol Biol* **318**, 1351-1365 (2002).

70. Ron D, Habener JF. CHOP, a novel developmentally regulated nuclear protein that dimerizes with transcription factors C/EBP and LAP and functions as a dominant-negative inhibitor of gene transcription. *Genes Dev* **6**, 439-453 (1992).

71. Konkimalla A, *et al.* Transitional cell states sculpt tissue topology during lung regeneration. *Cell Stem Cell* **30**, 1486-1502 e1489 (2023).

72. Beppu AK, *et al.* Epithelial plasticity and innate immune activation promote lung tissue remodeling following respiratory viral infection. *Nat Commun* **14**, 5814 (2023).

73. Riemondy KA, *et al.* Single cell RNA sequencing identifies TGFbeta as a key regenerative cue following LPS-induced lung injury. *JCI Insight* **5**, (2019).

74. Goss AM, *et al.* Wnt2/2b and beta-catenin signaling are necessary and sufficient to specify lung progenitors in the foregut. *Dev Cell* **17**, 290-298 (2009).

75. Li J, Wang Z, Chu Q, Jiang K, Li J, Tang N. The strength of mechanical forces determines the differentiation of alveolar epithelial cells. *Dev Cell* **44**, 297-312 e295 (2018).

76. Konkimalla A, *et al.* Multi-apical polarity of alveolar stem cells and their dynamics during lung development and regeneration. *iScience* **25**, 105114 (2022).

77. Alanis DM, Chang DR, Akiyama H, Krasnow MA, Chen J. Two nested developmental waves demarcate a compartment boundary in the mouse lung. *Nat Commun* **5**, 3923 (2014).

78. Inde Z*, et al.* Age-dependent regulation of SARS-CoV-2 cell entry genes and cell death programs correlates with COVID-19 severity. *Sci Adv* **7**, (2021).

79. Penkala IJ*, et al.* Age-dependent alveolar epithelial plasticity orchestrates lung homeostasis and regeneration. *Cell Stem Cell* **28**, 1775-1789 e1775 (2021).

80. Sprinzak D*, et al.* Cis-interactions between Notch and Delta generate mutually exclusive signalling states. *Nature* **465**, 86-90 (2010).

81. Kagami M*, et al.* The IG-DMR and the MEG3-DMR at human chromosome 14q32.2: hierarchical interaction and distinct functional properties as imprinting control centers. *PLoS Genet* **6**, e1000992 (2010).

82. Yao C*, et al.* Senescence of Alveolar Type 2 Cells Drives Progressive Pulmonary Fibrosis. *Am J Respir Crit Care Med* **203**, 707-717 (2021).

83. Wang F*, et al.* Regulation of epithelial transitional states in murine and human pulmonary fibrosis. *The Journal of clinical investigation* **133**, (2023).

84. Kaufman RJ. Stress signaling from the lumen of the endoplasmic reticulum: coordination of gene transcriptional and translational controls. *Genes Dev* **13**, 1211-1233 (1999).

85. Toth A*, et al.* Alveolar epithelial progenitor cells require Nkx2-1 to maintain progenitor-specific epigenomic state during lung homeostasis and regeneration. *Nature Communications* **14**, 8452 (2023).

86. Zhang P*, et al.* Enhancement of hematopoietic stem cell repopulating capacity and self-renewal in the absence of the transcription factor C/EBP alpha. *Immunity* **21**, 853-863 (2004).

87. Shen X*, et al.* EZH1 mediates methylation on histone H3 lysine 27 and complements EZH2 in maintaining stem cell identity and executing pluripotency. *Mol Cell* **32**, 491-502 (2008).

88. Muzumdar MD, Tasic B, Miyamichi K, Li L, Luo L. A global double-fluorescent Cre reporter mouse. *Genesis* **45**, 593-605 (2007).

1635

1636   89.   Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data
1637   analysis. *Genome Biol* **19**, 15 (2018).

1638

1639   90.   Hao Y*, et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
1640   3587.e3529 (2021).

1641

1642   91.   McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell
1643   RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337.e324
1644   (2019).

1645

1646   92.   Haber AL*, et al.* A single-cell survey of the small intestinal epithelium. *Nature* **551**, 333-
1647   339 (2017).

1648

1649   93.   La Manno G*, et al.* RNA velocity of single cells. *Nature* **560**, 494-498 (2018).

1650

1651   94.   Wolf FA*, et al.* PAGA: graph abstraction reconciles clustering with trajectory inference
1652   through a topology preserving map of single cells. *Genome Biology* **20**, 59 (2019).

1653

1654   95.   Bergen V, Lange M, Peidli S, Wolf FA, Theis FJ. Generalizing RNA velocity to transient cell
1655   states through dynamical modeling. *Nat Biotechnol* **38**, 1408-1414 (2020).

1656

1657   96.   Wagner DE, Weinreb C, Collins ZM, Briggs JA, Megason SG, Klein AM. Single-cell
1658   mapping of gene expression landscapes and lineage in the zebrafish embryo. *Science*
1659   **360**, 981-987 (2018).

1660

1661   97.   Heinz S*, et al.* Simple combinations of lineage-determining transcription factors prime
1662   cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-
1663   589 (2010).

1664

1665   98.   Gontarz P*, et al.* Comparison of differential accessibility analysis strategies for ATAC-seq
1666   data. *Sci Rep* **10**, 10150 (2020).

1667

1668   99.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
1669   RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1670

1671   100.   Mei S*, et al.* Cistrome Data Browser: a data portal for ChIP-Seq and chromatin
1672   accessibility data in human and mouse. *Nucleic Acids Research* **45**, D658-D662 (2016).

1673

1674   101.   Lei Y, He L, Yan C, Wang Y, Lv G. PERK activation by CCT020312 chemosensitizes
1675   colorectal cancer through inducing apoptosis regulated by ER stress. *Biochem Biophys*
1676   *Res Commun* **557**, 316-322 (2021).

1677

1678