A Y-linked duplication of anti-Mullerian hormone is the sex determination gene in threespine

stickleback

Matthew J Treaster, Jenny McCann, Kyra S Solovei, Ryan J Palmieri, Michael A White*

Department of Genetics, University of Georgia, Athens, GA, 30602

Corresponding Author:

Michael A. White

Department of Genetics, University of Georgia

120 Green St,

Athens, GA 30602

whitem@uga.edu

Abstract

Many taxa have independently evolved genetic sex determination where a single gene located on a sex chromosome controls gonadal differentiation. The gene anti-Mullerian hormone (*amh*) has convergently evolved as a sex determination gene in numerous vertebrate species, but how this gene has repeatedly evolved this novel function is not well understood. In the threespine stickleback (*Gasterosteus aculeatus*), *amh* was duplicated onto the Y chromosome (*amhy*) ~22 million years ago. To determine whether *amhy* is the primary sex determination gene, we used CRISPR/Cas9 and transgenesis to show that *amhy* is necessary and sufficient for male sex determination, consistent with the function of a primary sex determination gene. Despite being indispensable for sex determination, we detected low levels of *amhy* expression throughout early development. This indicates the mechanism of sex determination is likely unrelated to overall dosage of *amhy* and its autosomal paralog, *amha*. Threespine stickleback have striking differences in behavior and morphology between sexes. The creation of sex reversed lines allow us to investigate the genetic basis of secondary sex characteristics. Here we show one of the classic traits important for reproductive success, male nuptial coloration, is controlled by both Y-linked genetic factors as well as hormonal factors independent of sex chromosome genotype. This research establishes stickleback as a model to investigate how *amh* regulates gonadal development and how this gene repeatedly evolves novel function in sex determination. Analogous to the four core genotypes model in house mice, sex-reversed threespine stickleback offer a new vertebrate model for investigating the separate contributions of gonadal sex and sex chromosomes to sexual dimorphism.

Introduction

For species that have two discrete sexes, sex determination is the process that designates whether an individual will become male or female [1,2]. While this process is essential to the fitness and survival of a species, it is regulated by a surprising variety of initial

cues and can incorporate a range of extrinsic signals as well as intrinsic genetic factors often

located on sex chromosomes [1–3]. Numerous eukaryotic taxa have convergently evolved sex

chromosomes, such as the XX-XY system found in mammals, to regulate this critical

developmental process [1,2]. Sex chromosomes are derived from a pair of autosomes after one

of these autosomes acquires a sex determining (SD) gene [4]. Suppression of recombination

around this gene leads to sequence divergence of the two autosomes into a pair of divergent

sex chromosomes [4]. Why this recombination suppression occurs is not clear; however, the

prevailing model is that suppressed recombination will be selected for to link an SD gene with

nearby sexually antagonistic mutations that are beneficial to one sex but detrimental to the other

[4–6]. One remarkable observation that has emerged from comparative studies is the SD gene

responsible for initiating development of testes or ovaries in their respective sex is incredibly

variable [1,7]. How the otherwise well conserved sex differentiation developmental network can

be regulated by such a diverse range of SD genes remains poorly understood.

While a wide variety of genes have evolved novel function as sex determination genes,

some genes appear predisposed to acquiring this role and have convergently evolved as sex

determination genes in distantly related taxa [7]. *Sox3*, the ancestor of the mammalian SD gene

Sry, also acts as the SD gene in the Indian rice fish (*Oryzias dancena*) [8]. SD genes derived

from *dmrt1* are found in birds [9], the African clawed frog (*Xenopus laevis*) [10], and medaka

(*Oryzias latipes*) [11,12]. The most prevalent SD gene among vertebrates by far is *anti-Müllerian*

hormone (*amh*) [7]. Independently acquired Y-linked copies of *amh* or its dedicated receptor,

amhrII, have been identified in at least twelve different clades of teleost fish [13–23] as well as

in monotremes [24]. These Y-linked copies of *amh* and *amhrII* are presumed to be SD genes in

all of these taxa. Although *amh* accounts for over one quarter of identified SD genes in

vertebrates, how *amh* controls this critical developmental process and evolves novel function in

sex determination is not clear [7].

Amh is a member of the transforming growth factor beta (TGF-β) family of proteins that

are shared across vertebrates (Reviewed in [25,26]). *Amh* is unique among TGF-β hormones as

it has only one identified type II receptor, *amhrII*, of which *amh* is the only known ligand [27–29].

Amh and its receptor were first identified for their role in reproductive tract development in

mammals, where secretion of *amh* by the developing testes causes regression of the primordial

female reproductive tract, the Müllerian ducts [30]. Disruption of *amh* signaling in teleosts, which

lack Müllerian ducts, causes over proliferation of germ cells in the gonad and can interfere with

sex determination, even in species were *amh* is not the SD gene [31–33]. This suggests the

ancestral function of *amh* is likely in germ cell proliferation and maintenance which, if disrupted,

can disrupt sex determination, but how this disruption occurs is not clear. It also is not known

whether Y-linked copies of *amh* that have convergently evolved function in sex determination

control the downstream network in the same way or if *amh* can regulate sex differentiation

through a variety of distinct mechanisms.

Species of stickleback fish (family Gasterosteidae) have multiple, independently evolved

sex chromosome systems, making them an ideal model for investigating the origins of sex

chromosomes and SD genes [20,34–37]. Two separate duplications of *amh* that are associated

with male sex have been identified within stickleback fish. In the genus *Gasterosteus*, *amh* was

duplicated from autosome 8 onto autosome 19 approximately 22 million years ago (Mya),

leading to the evolution of a Y chromosome [20,38]. An independent duplication of *amh* onto

chromosome 20 was identified in the brook stickleback, *Culea inconstans*. This duplication

differs from the ancestral copy on autosome 8 by only a handful of single nucleotide

polymorphisms (SNPs), suggesting this duplication occurred much more recently than in the

genus *Gasterosteus* [22,39]. Consistent with its recent acquisition, the duplicate copy of *amh* is

not a fully penetrant SD gene, and its association with male sex varies across populations [39].

This provides a unique opportunity to examine the convergence evolution of *amh* as an SD

gene at two dramatically different stages of sex chromosome evolution.

The threespine stickleback (*Gasterosteus aculeatus*) is a powerful emerging model for

evolutionary and developmental biology with a robust set of genomic and functional genetic

tools [40,41]. Their small size, large clutch sizes, and relatively short generation time make them

a highly tractable model for studying the developmental mechanisms of *amh* as an SD gene.

Threespine stickleback also have many well-studied sexually dimorphic traits and mating

behaviors that have provided substantial insight into the evolution of sexual dimorphism and its

effect on fitness [42–63]. Like other members of its genus, threespine stickleback have retained

a functional copy of autosomal *amh* on chromosome 8 (hereafter called *amha*) as well as the Y-

linked *amhy*. The AMH prodomain and TGF-β signaling domain within *amhy* are conserved (Fig

S1), indicating the Y-linked duplicate is likely functional [20]. In addition, *amhy* is expressed in

larvae around the time of sex determination, consistent with a role in initiating male

development [20]. Beyond this, it is unknown whether *amhy* has a functional role in male sex

determination. In this study, we show that *amhy* is both necessary and sufficient for male sex

determination in the threespine stickleback, demonstrating that it is the SD gene in this species.

We also perform RNA-seq of stickleback embryos and larvae to characterize expression of

amhy and early transcriptional differentiation of males and females. Our findings establish

stickleback fish as a premier model to study both the molecular mechanisms of *amh* based sex

determination as well as the convergent evolution of *amh* as a sex determination gene in

vertebrates.

Results

CRISPR/Cas9 knockout of *amhy* caused male-to-female sex reversal in XY fish.

If *amhy* is the SD gene in threespine stickleback, it should be both necessary and

sufficient for male sex determination. To test this, we first knocked out function of *amhy* in XY

fish using CRISPR/Cas9. We designed two sgRNAs targeting exon 1 and exon 3 of *amhy* (Fig

1). We injected stickleback embryos from lab-reared fish derived from two independent marine

populations (Port Gardner Bay and Japan Pacific Ocean) with CRISPR ribonucleoprotein (RNP)

made with one of the two sgRNAs. 8/17 (47%) of F0 XY fish injected with RNP exhibited male-

to-female sex reversal. For all eight XY sex reversed fish, we observed ovaries with all cell

types seen in wildtype ovaries (Fig S2).

F0 crispants are often mosaic for mutations induced by CRISPR/Cas9 gene editing,

which can cause variation in observed phenotypes. To generate non-mosaic F1 gene-edited

fish, we used two F0 XY males that had heritable germline mutations in *amhy* but were not sex

reversed, indicating they likely did not have *amhy* mutations in cell-lineages important for sex

determination. Sperm from these fish were used to fertilize eggs from wildtype XX females. F1

offspring from these crosses inherited frameshift mutations in *amhy*, identified by Sanger

sequencing (Fig S3). We analyzed five F1 males from one F0 crispant with mutations in exon 1

and 21 F1 males from one F0 crispant with mutations in exon 3. All 26 F1 fish exhibited

complete male-to-female sex reversal and developed ovaries morphologically and histologically

indistinguishable from wildtype XX females (Fig 1C). No F1 XY fish showed development of

testis or ambiguous gonad tissue with any visible spermatocyte production. We identified three

loss of function alleles in exon 1 of *amhy* (Fig 1A) and one in exon 3 (Fig 1B). In ten F1 fish, the

region of *amhy* in exon 3 targeted by CRISPR/Cas9 could not be amplified. PCR of exons 1 and

7 showed the presence of these distal portions *amhy*, but PCRs spanning a large putative

deletion all failed to amplify, suggesting a chromosomal rearrangement disrupted *amhy* in these

fish. As a specific mutation could not be positively identified, these samples were excluded from

further analysis and final counts.

We analyzed the expression of canonical male differentiation genes *amha* and *dmrt1*

and female differentiation genes *foxl2* and *cyp19a1a* using *in situ* hybridization on wildtype and

F1 XY *amhy*-KO ovaries (Fig 2, Fig S4). We also analyzed *gsdf* as a positive control, as this

gene is expected to be expressed in somatic gonadal cells of both sexes [64]. Both wildtype and

sex reversed ovaries showed expression of *foxl2* and *cyp19a1a* in somatic cells. Expression of

amha and *dmrt1* was minimal and restricted to oocytes when present as has been observed in

some species of fish [21,65–68]

Knockout of *amh* or *amhrII* in teleost species without amh-based sex determination has

caused male-to-female sex reversal, including in lab strains of zebrafish which lack sex

chromosomes (reviewed in [69]) and medaka which have an XX-XY system with a *dmrt1*

derived sex determination gene [31]. Thus, it is possible that off-target mutations of the

homologous *amha* could contribute to the sex reversal observed in F1 *amhy* mutants. While

both sgRNAs have multiple mismatches to their homologous region in *amha*, we verified that all

F1 XY *amhy*-KO females did not have mutations within *amha* by sanger sequencing (Fig S5)*.*

This demonstrates that the sex reversal phenotype was not caused by mutations of *amha*. We

also investigated whether the sex reversal phenotype was caused by a mutation at an unknown

autosomal locus. This autosomal mutation must be dominant since we observed male-to-female

sex reversal in heterozygous XY F1 fish. F1 XX siblings of sex reversed XY fish would also be

heterozygous for this mutation, and half of their F2 XY progeny would be sex reversed. We

crossed two F1 XX PG females with wildtype XY males and phenotyped F2 offspring for sex

reversal. All 16 16 F2 XY fish were phenotypically male, indicating that sex reversal in F1 XY

amhy-KO fish was not caused by an off-target autosomal mutation.

Introduction of an *amhy* transgene caused female-to-male sex reversal in XX fish.

To determine if *amhy* is sufficient for male sex determination, we created a construct

(pT2-5amhy2.5,Xla.Ef1a:EGFP) containing an 11 kb region of the Y chromosome which

includes *amhy*, 5 kb of upstream sequence, and 2.5 kb of downstream sequence (Fig S6F). By

including the surrounding non-coding sequences, we aimed to recapitulate the endogenous

expression patterns of *amhy.* An Xla.Ef1a:EGFP cassette downstream of *amhy* was included to

facilitate the screening of injected embryos for transgene integration. These components were

flanked by *Tol2* transposon sequences to facilitate random integration into the genome by the

Tol2 transposase [70,71]. 16 F0 XX fish showed positive GFP expression and successful

transgene integration, detected through PCR genotyping (Fig 3D). At reproductive age, all 16 F0

XX *Tg(amhy,EGFP)* were female-to-male sex reversed, developing testes rather than ovaries

(Fig 3A). All 10 control F0 XX fish injected with the control Xla.ef1a:EGFP transgene showed

positive presence of the transgene, but did not show any signs of sex reversal and developed

cytologically normal ovaries (Fig 3B). We also analyzed the expression of *amha, dmrt1, foxl2,*

cyp19a1a, and *gsdf* using *in situ* hybridization on wildtype and F0 XX *Tg(amhy,EGFP)* testes

(Fig 2, Fig S7). Both wildtype and sex reversed testes showed strong expression of *amha* and

dmrt1 in somatic cells and no discernable expression of *foxl2* or *cyp19a1a.*

We attempted to constitutively express *amhy* using a synthetic exon-only coding

sequence with an N-terminus FLAG tag joined to a 3' EGFP by a P2A self-cleaving peptide

[72,73]. Plasmids were made using the *D. rerio* ubiquitin promoter [74] or the *X. laevis* ef1a

promoter [75] to drive expression of the amhy,EGFP construct (Fig S6). Transgenes with the

respective promoter driving expression of only EGFP were used as controls. While somatic

EGFP expression was observed with both control transgenes indicating these promoters can

drive expression in stickleback, neither the Dre.Ubi:amhy,EGFP nor the Xla.Ef1a:amhy,EGFP

transgene produced any larvae with visible somatic EGFP, indicating that the *amhy* transgenes

were failing to integrate or be actively expressed in any embryos. We hypothesize that the

failure to constitutively express *amhy* may be due to cytotoxicity of extremely high and

widespread expression of *amh* using these non-native, constitutive promoters.

Sex reversed *amhy* mutants produce viable gametes but are subfertile.

Sex-reversed XY *amhy*-KO females produced mature oocytes and became gravid, but

they were subfertile compared to wildtype females. To assess fertility, we crossed a three F1 XY

amhy-KO females to wildtype XY males. These clutches all showed extremely low viability rates,

between 2% and 6% (Table S1), measured by presence of a gastrula and active epiboly 12 to

24 hours after fertilization (haf). One of the three F1 fish was crossed four additional times and

showed improvement in viability with the subsequent crosses. The viability rates of the fourth

and fifth cross were 78% and 79% respectively, producing a total of 148 viable F2 eggs (Table

S1). Three days after fertilization, 50 of the 148 eggs had arrested development. We were able

to genotype 48 of the failed embryos. Three-quarters of these (36) carried no X chromosome

and both the paternally transmitted Y chromosome (Y^p) and the maternally transmitted *amhy-*

KO Y chromosome (Y^m) (Fig 4). The remaining embryos were reared to adulthood and 92 were

genotyped for their sex chromosome complement. No adult YY fish were observed, indicating

that the YY genotype is embryonic lethal due to the missing X chromosome. XY^m females were

found to transmit the X and Y^m chromosome with equal frequency, indicating there was not any

biased transmission of the X or Y chromosome during female meiosis (36 Y^pY^m embryos out of

148 total does not deviate from a 1:4 Mendelian ratio; chi-squared = 0.0360; p = 0.8494; of the

92 adult F2s genotyped 36 XX, 27 XY^p , and 29 XY^m does not deviate from the expected 1:1:1

ratio; chi-squared = 1.457; p = 0.4827). Subfertility also extended to second generation sex-

reversed XY females. We crossed 11 F2 females with wildtype XY males. The clutch size did

not differ significantly between XY^m and XX females (Fig 4C; Table S2). However, the viability

rate was substantially lower in the F2 XY^m females, similar to the F1 generation (Fig 4D, Table

S2). Two XY^m F2 females were also eggbound and were unable to be stripped of eggs.

Female-to-male sex reversed F0 XX fish had varied fertility. We initially dissected eight

F0 XX *Tg(amhy,EGFP)* males at 6 months post fertilization. Seven of these XX males had

small, underdeveloped testes, whereas one had testes morphologically resembling those of a

fertile, wildtype male. Despite its mature morphology, the testis was composed only of

premeiotic spermatogonia, immature round spermatids, and no mature sperm (Fig 3A). Crosses

with this male only had a 3% viability rate, suggesting the XX male was not reproductively

mature. Of six XY fish injected with the control Ef1a:EGFP transgene that were dissected at six

months, only two had testes that appeared reproductively mature, and histology of these testes

also lacked mature sperm. XX females injected with the control transgene were also not fully

mature (Fig 3B).

We then assessed fertility of transgenic fish at 12 months of age to ensure the males

had adequate time to fully mature. At this age, XY *Tg(EGFP)* control fish showed nuptial

coloration, and both XX and XY *Tg(amhy,EGFP)* fish were reproductively mature. We analyzed

eight additional F0 XX *Tg(amhy,EGFP)* males at 12 months post fertilization. Three of these

males had morphologically and histologically normal testes with fully mature sperm (Fig 3A).

These three males showed normal viability rates when crossed with wildtype XX females (78

viable of 97 total eggs, 84 viable of 89 total eggs, and 231 viable of 235 total eggs). One male

also produced offspring with somatic GFP expression at a very low frequency (16/231)

indicating a heritable germline integration of the transgene had occurred. The remaining five

males had hypotrophic testes, so no crosses were performed with these males. Histology of the

hypotrophic testes was abnormal and did not show any clearly identifiable germ cells (Fig 3A).

This phenotypic variation could be a result of variable *amhy* expression influenced by transgene

copy number and insertion location.

Male nuptial coloration is present in both sex reversed genotypes

Male nuptial coloration is one of the most prominent sexually dimorphic characteristics in

threespine stickleback. During their breeding season, males from many stickleback populations

develop throats and dark blue eyes and bodies [76]. To investigate the separate contributions of

gonadal and genotypic sex on this trait, we compared coloration in mature sex reversed XY

amhy-KO females and XX *Tg(amhy,EGFP)* males to wildtype XX and XY fish (Fig 5). Nuptial

coloration is prominent in XY males by one year post hatching. While neither *amhy* mutant

genotype fully recapitulates wildtype XY male coloration, both XX males and XY females show

elements of male nuptial coloration and are distinct from wildtype XX females. In XX males,

nuptial coloration was present in all three fertile males, but not in any males with hypotrophic

testes. Together, these findings show that nuptial coloration is a product of both sex

chromosome genotype and gonad identity.

amhy does not have distinct expression timing relative to amha

Since Amh shows high specificity for its dedicated receptor, AmhrII, [27–29] we predicted

that *amhy* would be functionally constrained regarding coding sequence evolution and would

instead evolve novel function through regulatory changes. We investigated the degree of

sequence conservation within putative regulatory regions (2 kb upstream and downstream of

the coding sequence), of *amhy* relative to *amha*. We also compared sequence similarity

between *amha* and the autosomal *amh* in the ninespine stickleback (*Pungitius pungitius*). The

divergence time between threespine and ninespine stickleback is approximately 26 Mya [77],

similar to the 22 Mya divergence time between the threespine *amha* and *amhy* [20]. There was

less sequence identity between the coding sequences of threespine *amha* and *amhy* than

between threespine *amha* and ninespine *amh* (Fig 6, Table 1). This is consistent with weaker

purifying selection acting on the Y chromosome due to a lack of recombination [78]. This was

also supported by a higher dN/dS value between threespine *amha* and *amhy* than between

threespine *amha* and ninespine *amh*. The sequence identity of upstream and downstream

regions was much lower between *amha* and *amhy* than between *amha* and ninespine *amh*,

indicating *amhy* has undergone significant regulatory evolution since its duplication onto the Y

chromosome, which may cause differences between *amha* and *amhy* expression.

To determine whether expression of these genes differ throughout development, we

sequenced the total transcriptomes from prehatching embryonic stages 17 (early

somitogenesis, ~43 haf), 20 (onset of pigmentation, ~73 haf), and 23 (completion of yolk sac

circulation, ~122 haf) as well as larvae 0, 4, and 8 days after hatching (dah), where hatching

occured around 200 haf [79]. The earliest identified morphological difference between male and

female threespine stickleback is a difference in primordial germ cell count at 3 to 4 dah [80]. Sex

determination should begin sometime prior to this stage. We predicted that the unique

regulatory architecture of *amhy* resulted in earlier expression timing relative to *amha.* Thus,

amhy would be expressed at higher levels relative to *amha* early in development in order to

initiate sex determination. As the gonad differentiates, *amha* expression should increase relative

to *amhy* due to the canonical role of *amh* in male differentiation. This pattern has been observed

in Japanese flounder [15] and Patagonian pejerrey [13] which also have *amhy* SD genes.

Contrary to our prediction, we found that *amhy* expression was substantially lower than *amha* at

all observed timepoints pre- and post-hatch (Fig 7B). This difference was still present after

scaling read counts for differences in transcript length which is not accounted for by normalized

read counts (Table S3). We found *amhy* expression trended towards higher expression at later

developmental stages, consistent with a role in sex determination. However, this difference was

not significant (anova p = 0.0769). We then hypothesized that male sex determination could be

initiated by a higher overall dose of *amhy* when expressed simultaneously with *amha.* We

compared *amha* and transcript length adjusted *amhy* normalized read counts at each stage and

found no significant difference in total *amh* dosage between males and females at any timepoint

(Table S3), suggesting dosage alone is not a mechanism for initiating male differentiation.

Combined, our results indicate the unique regulatory elements that evolved for *amhy* do not

increase early expression of *amhy*. Instead, they may alter expression level of *amhy* at other

developmental stages and/or change spatial expression patterns, contributing to its new

function in sex determination.

Sex differentiation genes are not differentially expressed between males and females

early in development

Following sex determination, we expected to see divergence in the expression of overall

male and female transcriptomes. However, principal component analysis of autosomal gene

expression indicated that male and female larvae had similar expression profiles. Autosomal

genes were clustered by developmental time rather than separating by sex at any stage of

development (Fig 7A). Only sex chromosomes showed clear separation by both sex and

developmental time (Fig S8), since the X chromosome is expressed two-fold higher in females,

relative to males, and the Y chromosome is only expressed in males. The overall lack of sex-

specific expression on the autosomes suggests that many aspects of sex differentiation occur

well after the earliest noted phenotypic difference between males and females: the proliferation

of germ cells in the primordial gonad at 3 to 4 dah [80].

We examined whether any developmental stages had genes that were significantly

differentially expressed between males and females. Many male and female differentiation

genes are highly conserved across vertebrates [2] and should also be expressed during

development of the threespine stickleback. For instance, we expected differential expression of

key transcription factors canonically involved in differentiation, including the male biasing factors

dmrt1 and *sox9b* or the female biasing factors *foxl2a*, followed by divergence of downstream

genes such as the aromatase enzyme *cyp19a1a* and the TGF-b hormones *amha* and *gsdf*. In

total, we found 464 differentially expressed genes (DEGs) between males and females on the

autosomes (File S1). Key regulators of differentiation should be differentially expressed

throughout development and into adulthood in order to actively maintain gonadal fate [2].

However, 438 of the DEGs were only differentially expressed in a single stage (File S1),

suggesting these genes do not play a substantial role in sex differentiation. Additionally, while

we expected to see an increase in the number DEGs over time due to activation of male and

female specific gene networks, the number of DEGs was highest at stage 17 (152 DEGs),

decreasing in later stages until a small increase from 4 dah (40 DEGs) to 8 dah (72 DEGs) (Fig

S9). None of the canonical differentiation genes of interest were differentially expressed

between males and females at any timepoint examined in this study (Fig 7). Male differentiation

genes *dmrt1* and *amha* and female differentiation genes *foxl2* and *cyp19a1a* were differentially

expressed in adult gonads of both wildtype and sex reversed fish, indicating their function in

differentiation are conserved (Fig 2, Fig S4, Fig S7). Of the identified DEGs, only a small

handful have known associations with gonadal function, development, or differentiation. Two

genes associated with germ cell specification and maintenance [81], *ddx4* and *piwil1*, showed

female biased expression at 8 dah (Fig 2C). Interestingly, *amhrII,* which shows female biased

expression in embryonic mouse gonads [82], was also expressed more highly in females at 8

dah. While *cyp19a1a* was not differentially expressed, the steroidogenic enzyme *hsd17b1*,

which plays a role in both estrogen activation and androgen inactivation [83] and acts as the SD

gene in *Seriola* fishes [84], showed female biased expression at 8 dah. A lack of differential

expression in many of these key genes indicates that gonad differentiation occurs after 8 dah.

We identified some other DEGs that have known reproductive functions in other

vertebrates but do not have clear connections to sex differentiation and are only differentially

expressed at a single timepoint. Gonadotropin-releasing hormone II receptor (LOC120811940)

has known roles in the stimulating steroidogenesis in both testes and ovaries across mammals

[85] and had male biased expression at stage 17 (logFC = 2.74, adjusted P = 1.4E-3). TGF-beta

3 hormone is expressed more highly in male mice during gonadogenesis [86], and a TGF-beta 3

proprotein like gene (LOC120807995) was expressed slightly higher in females at stage 17

(logFC = -0.63, adjusted P = 0.044). *Piwil2*, another germ cell associated gene [81], was

expressed higher in males at stage 20 (logFC = 1.72, adjusted P = 3.2E-3). *Cdc25b* is

expressed at low levels in mouse ovaries and testes [87] and was male biased at stage 23

(logFC = 0.52, adjusted P = 0.012). Separin/separase (*espl1*) has known roles in mitotic and

meiotic chromosome segregation [88] and was expressed at higher levels in 8 dah females

(logFC = -0.56, adjusted P = 1.2E-4). Additional work will be necessary to understand what

function, if any, these genes have in threespine stickleback gonad development.

Discussion

***amhy* is the sex determination gene in threespine stickleback fish**

Our findings demonstrate that *amhy* is the SD gene in threespine stickleback, as it is

both necessary and sufficient for sex determination (Table 3). Independently derived *amhy*

genes have been functionally demonstrated to have a role in sex determination in three other

clades of teleost; however, threespine stickleback offers a novel perspective as their *amhy* is

more diverged, and genetic sex determination is more canalized in this species. Nile tilapia and

Japanese flounder have only one and nine SNPs, respectively, that differentiate their X- and Y-

linked copies of *amh* [15,16], and *amhy* in Patagonian pejerry shares 94% nucleotide identity

with its autosomal paralog [13]. In all three of these species, genetic sex determination can be

overridden to varying degrees by temperature [89–91]. In contrast, the threespine stickleback,

amha and *amhy* only share 78% nucleotide identity (Table 1). There is no evidence that

temperature can disrupt sex determination in threespine stickleback, and occurrences of sex

reversed or intersex stickleback in wild populations is rare [92,93]. This indicates that genetic

sex determination is highly canalized in threespine stickleback and less susceptible to

environmental signals that many other teleost species, making stickleback a robust model to

investigate how *amh* regulates gonadal development. Comparative studies across these taxa

will be invaluable for understanding how *amh* repeatedly evolves this novel function, and the

presence of an independently derived male associated *amh* duplication in the closely related

brook stickleback [22,39] makes the stickleback family an especially powerful tool for studying

the convergent evolution of *amh* as an SD gene.

Potential mechanisms of sex determination in stickleback

Germ cell density has been shown to affect sex determination in several teleost species,

with the male fate associated with lower germ cell count and the female fate with a higher count.

Common lab zebrafish strains are undifferentiated gonochorists, where all individuals begin

ovary development, but the gonad transitions early in males to a testis [94]. Complete ablation

of germ cells causes zebrafish to undergo male development [95], and a threshold of germ cells

at the time of gonadal differentiation is necessary to sustain ovary development [96]. *O. latipes*,

like sticklebacks, are differentiated gonochorists where either testis or ovary development

begins from a common undifferentiated primordium [97]. Ablation of germ cells in XX *O. latips*

causes female-to-male sex reversal [98], and overproliferation of germ cells in XY fish causes

male-to-female sex reversal [99]. The association of germ cells with sex has also been

observed outside of teleosts in the red eared slider turtle (*Trachemys scripta elegans*) which has

temperature dependent sex determination. Embryos incubated at female promoting

temperatures have higher germ cell counts than those incubated at male promoting

temperatures, and depleting germ cells at intermediate temperatures biased the sex ratio

towards males [100]. This suggests the presence of a widely conserved, ancestral mechanism

in vertebrates by which germ cell density governs gonad differentiation. A new duplication of

amh could increase overall *amh* dosage, reducing germ cell proliferation to co-opt this

underlying mechanism and become a male determining gene.

Consistent with this mechanism, we found evidence that *amhy* is expressed in

threespine stickleback when germ cell proliferation diverges in males and females. While *amhy*

levels were low at all stages observed, *amhy* expression was highest in post hatching stages

where differences in germ cell proliferation between males and females are first observable in

the developing gonad [80]. We also found increased expression of germ cell specific genes (e.g.

ddx4 and *piwil1*) in females at 8 dah, corroborating previous histological observations of

increased germ cell proliferation in female stickleback gonads beginning at 3 dah [80].

Overexpression of *amh* in mice [101] and chicken [102] causes reduced germ cell count and

gonad size. Studies misexpressing *amh* in telesots have primarily focused on sex determination;

however, gonadal hypertrophy observed in *amh* knockouts in other teleosts [31,33] alongside

gonadal hypotrophy with a lack of mature germ cells observed in some XX *Tg(amhy,EGFP)*

males in this study demonstrate the conserved function of *amh* suppressing germ cell

proliferation in stickleback and other teleosts.

Somatic cells may also be affected by *amhy* in other ways. In mice, low concentrations

of Amh promote proliferation and growth factor production in somatic cells, while increased

concentrations induce apoptosis [103]. Altering somatic cell proliferation could lead to

differences in the levels of hormones and growth factors that they produce, shaping gonadal

development. While Amh is not necessary for initial gonadal differentiation, it is important to

maintaining Sertoli cell fate in fetal mice [104]. Estrogen production in somatic cells also plays a

major role in sex differentiation and ovary development in teleosts (reviewed in [105,106]). In

tilapia, Amh signaling directly inhibits *cyp19a1a* production via Smad transcription factors [107].

While we did not detect a difference in *cyp19a1a* expression between males and females in

embryos or larvae, it was expressed more highly in adult ovaries than in testes, so *amhy* could

play a reinforcing role later in development to suppress estrogen synthesis in males. As we

were unable to isolate embryonic gonadal tissue for expression analysis, our results are

currently limited to characterizing total expression in the whole embryo. *Amh* can be expressed

in other developing tissues, such as the brain and heart [108,109]. If this is also the case in

threespine stickleback, this would obscure the true *amhy* to *amha* expression ratio in the gonad.

Further research characterizing the expression of *amha* and *amhy* in stickleback at spatial and

single cell levels will be essential for understanding how *amhy* functions in the developing

stickleback gonad.

The evolutionary changes that have given rise to novel function of *amhy* in stickleback

sex determination remain unclear. It is clear that substantial regulatory evolution has taken

place following the duplication of *amh* onto the Y chromosome; however, this regulatory

evolution has not lead to earlier or higher expression of *amhy* relative to *amha* as we initially

predicted. While we suspect a more detailed study of *amha* and *amhy* expression at tissue and

cell specific levels will lend insight into the role of this regulatory evolution, we also should not

rule out the significance of coding sequence evolution for *amhy*. In Nile tilapia, *amhy* expression

is lower than the X-linked *amh* throughout development, and misexpression of *amhy* but not

amh induces male sex determination [16]. This implies that the single missense SNP between

amh and *amhy* in tilapia is important to the function of *amhy* in sex determination. In Spotted

knifejaw (*Oplegnathus punctatus*), a Y linked copy of *amhrII* shows higher affinity for binding

amh than its X linked paralog [23]. In zebrafish which lack *amhrII*, *amh* may instead use the type

II TGF-β receptor *bmpr2a* [110,111]. Mutations that increase the stability of Amhy, increase its

affinity for the receptor, AmhrII, or promote interaction with a non-canonical receptor could also

affect gonadal development.

Sex differentiation occurs later than anticipated

We observed no differential expression in canonical vertebrate sex differentiation genes,

including *dmrt1, sox9b, foxl2a,* and *cyp19a1a.* While the order and timing of expression varies

across taxa, these core genes are reliably differentially expressed between males and females

before the onset of morphological gonadal differentiation [112–114]. This indicates that there is

some delay between the initial establishment of germ cell differences between males and

females and the later differentiation of the primordial gonad into either a testis or ovary. We

propose that sex determination in threespine stickleback occurs in two phases. In the early

critical period of sex determination, germ cell proliferation in the undifferentiated gonad diverges

between males and females due to the presence of *amhy* in males. Then, during the sensitive

period of gonadal differentiation, the difference in germ cell count between individuals dictates

gonadal fate, marking the end of the sex determination and the beginning of differentiation.

Unlike *Sry* in mammals which is expressed in a brief window and directly initiates gonadal

differentiation [115], *amhy* in stickleback appears to be expressed throughout this early period of

germ cell proliferation. These two phases of germ cell proliferation followed by gonad

differentiation are also present in other species with SD genes including Nile tilapia (reviewed in

[116]), and medaka (reviewed in [97]) as well as species without SD genes such as red eared

slider [100] where germ cell proliferation is affected by temperature. This suggests that this two-

phase system of differential germ cell proliferation followed by gonadal differentiation is broadly

conserved across vertebrates, but what combination of environmental and/or genetic factors

influences germ cell proliferation has changed between taxa. The ability of *amh* to immediately

integrate with this preexisting sex determination network through its existing function in

regulating germ cell proliferation along with possible reinforcing roles in regulating somatic cell

identity and estrogen synthesis would explain the prevalence of *amh* as an SD gene.

Fertility of sex reversed threespine stickleback

Both XY *amhy*-KO females and XX *Tg(amhy,EGFP)* male sex reversed fish were fertile

in our study. Fertility of sex reversed mutants is not uncommon in other teleost species;

however, it is often observed in species with younger or less diverged sex chromosomes. In

these instances, the X and Y chromosomes contain a similar complement of genes. In medaka,

the Y specific region is 280 kb and the only functional gene it contains is the sex determining

dmrt1by [12]. In Nile Tilapia, the sex determining region on linkage group 23 is 1.5 Mb and

contains 51 annotated genes, including the sex determining *amhy* [117]. In both species, sex

reversed fish are fertile and YY offspring are viable [11,118]. In contrast, the threespine

stickleback Y chromosome has a ~15 Mb non-recombining region [20]. Within this region, the Y

chromosome has lost half of the almost 1200 ancestral gametologs and has accumulated many

Y-specific genes not present on the X chromosome [20]. Sex chromosome-specific gene

content evolution could lead to improper expression in the opposite sex, affecting

gametogenesis. Given this divergence, the ability of both XY *amhy*-KO females and XX

Tg(amhy,EGFP) males to reproduce is surprising. In mouse, reduced fertility of XY females is a

result of meiotic errors between the X and Y chromosomes as well as the effect of aberrant Y-

chromosome genes in the ooplasm on early development (Reviewed in [119]). As clutch size in

XY females did not differ from XX siblings and some crosses from XY females have normal

viability, it seems unlikely that widespread meiotic failure occurs in XY female stickleback.

Furthermore, it is clear that Y chromosome transcripts or dosage differences from X

chromosome genes do not prevent embryo development. Threespine stickleback are batch

spawners that produce multiple clutches of eggs over a single breeding season [76]. If ovulated

eggs are not passed, they can become overripe and inviable [120,121]. Overripe eggs can also

harden and obstruct the fish, causing them to become eggbound. Many of the clutches from XY

amhy-KO females had this hardened, overripe appearance that is uncommon in our lab-reared,

wildtype fish. In addition, some XY females we attempted to cross had no cloaca dilation and

were eggbound. We suspect that subfertility of XY females is not due to an inability to produce

viable eggs, but due to an inability to pass these eggs once mature. While wildtype fish can

pass eggs on their own and begin another cycle of ovulation, XY females retain these eggs until

they are manually removed. Repeated and frequent crossing would prevent overripe eggs from

building up and allow the harvest of freshly ovulated, viable eggs. Hormones produced by the

gonad and pituitary gland appears to play a role in egg maturation and overripening for

stickleback [121,122], so while XY females are histologically similar to XX females, variation in

hormone production could inhibit proper production and release of eggs.

The mammalian Y chromosome has several genes essential for spermatogenesis and

sperm development [123–127]. Single cell sequencing has shown that many Y-specific genes

are expressed throughout spermatogenesis in stickleback [128], but these genes are apparently

not essential for this process. Several XX *Tg(amhy,EGFP)* males from this study had testes with

many fully mature spermatozoa that were able to fertilize eggs and generate viable progeny at

similar rates to wildtype sperm. This raises a question of the importance of the Y chromosome

to fitness and reproduction. Testes from fertile XX males did not show an obvious reduction in

sperm number indicating that widespread meiotic arrest does not occur in XX males.

Stickleback are external fertilizers in which sperm competition frequently occurs [129]. Variation

in sperm morphology [130] and motility [131] can contribute to reproductive success, so there

may be differences in sperm function that our *ex vivo* fertilization assays do not capture. Some

XX *Tg(amhy,EGFP)* males did have hypotrophic testes with a reduction in identifiable germ

cells, but since this phenotype was not universal, we suspect that variation in transgene copy

number and insertion location led to overexpression of *amhy* in these fish. The germline

Tg(amhy,EGFP) offspring we produced will allow us to determine if this phenotypic variation

exists in a consistent genetic background.

A four core genotypes stickleback model

Sex chromosomes impact a wide range of traits outside of fertility that are still critical to

fitness. By comparing XY males and XX females to sex reversed XY females and XX males, the

four core genotypes mouse model has allowed researchers to dissect the role of genotypic and

gonadal sex to a number of behavioral and physiological sex differences in mammals (reviewed

in [132,133]) including mating behavior [134] and parental care [135]. The creation of

reproductively mature XX males and XY females in this study offers a new vertebrate model to

study the separate contributions of gonadal sex and sex chromosomes to sexual dimorphism.

Many traits are sexually dimorphic in the threespine stickleback, including shape and

size of the head and body [42–45], bony armor features [42,43], and brain size [46,47] as well

as feeding behaviors [48,49] and boldness [50]. Male stickleback also show complex courtship

and mating behavior [51,52] and male-only paternal care [53], while females show a variety of

mate choice preferences [54–59]. Male nuptial coloration is a particularly interesting trait, as

strong coloration is beneficial for courtship [59,60], but has a cost of increased risk of predation

[61,62]. Therefore, sexually antagonistic selection would select for stronger, more conspicuous

coloration in males and inconspicuous coloration in females. Androgens, secreted from the

gonads, are shown influence nuptial coloration in stickleback [63]. These observations are

supported by the presence of nuptial coloration in XX males in our study, which lack all Y-linked

genes, but have functioning testes. In contrast, male nuptial coloration in XY females that lack

testes shows that there are also Y-linked factors that contribute to nuptial coloration independent

of gonadal sex. Future research with the four core genotypes stickleback will provide many

insights into the genetic basis of sexual dimorphism and how sexually antagonistic selection

shapes the evolution of sex chromosomes.

Methods

All experiments used lab-derived progeny from wild-caught threespine stickleback fish

from Port Gardner Bay (Washington, USA) and Japan Pacific Ocean (Akkeshi, Japan). All

procedures were approved by the University of Georgia Animal Care and Use Committee

(protocols A2021 07-031-Y3 and A2024 08-009-Y1). We collected fish from Washington State

over multiple years using Scientific Collection Permit numbers 21-122, 22-150, 23-147, and 24-

143.

CRISPR/Cas9 knockout of *amhy*

We microinjected stickleback embryos following standard procedures [40], with the

addition of KCl to the injection mix at a final concentration of 0.2 M [136]. Cas9 protein was

purchased from QB3 MacroLab (University of California, Berkeley). Synthetic sgRNAs were

designed using CHOPCHOP [137–139] and purchased from Synthego: amhy_sgRNA_77r (5'-

GATCAGCTGCTGTCCATGCA-3') and amhy_sgRNA_600r (5'- GGTCCTCTCTCGTCTTCGTT -

3'). We combined Cas9 protein and sgRNA at a final concentration of 10 μM each and

incubated for at least ten minutes at room temperature to allow the ribonucleoprotein (RNP) to

form. The injection mix was made to a final concentration of 4 μM RNP.

Synthesis of constructs and transgenesis

Plasmids pBT2 [140], pUbi:Switch [74], and pCS-zT2TP [8] were kindly shared by Mary

Goll. Plasmid pT2AL200R150G [70] (referred to as pT2Xla.Ef1a:EFGP) was kindly shared by

Tyler Square and Craig Miller. We purchased a synthetic amhy,EGFP construct from TWIST

Bioscience (San Francisco, CA, USA). This construct consisted of a Kozak consensus

sequence (5'-GCCGCCACCATGG-3'), N-terminus flag tag, the threespine stickleback *amhy*

coding sequence, a P2A self cleaving peptide [72,73], EGFP, and an SV40 poly(A) sequence,

all flanked by 5' BamHI and 3' NotI restriction sites to facilitate cloning. We performed all

restriction digests were following the manufacturer's guidelines (New England Biolabs). We

isolated all of the plasmids using a Monarch Plasmid Miniprep Kit (NEB #T1010) and gel

purified all DNA fragments using the Monarch Gel Extraction kit (NEB #T1020). DNA fragments

were ligated using T4 DNA Ligase following manufacturer recommended protcols (NEB

#M0202). We transformed all plasmids into DH10B competent cells (Thermo Scientific EC0113).

All plasmids were sequenced by Azenta Life Sciences (South Plainfield, NJ, USA) (File S2).

We assembled each construct as follows:

1. pT2Dre.Ubi:amhy,EGFP: The *D. rerio* ubiquitin promoter was excised from

pUbi:Switch and cloned into pBT2 using XhoI and BamHI. The synthetic amhy,EGFP

construct was excised from pTwist_amhy-P2A-EGFP and cloned into the

intermediate pT2Dre.Ubi backbone using BamHI and NotI.

2. pT2Dre.Ubi:EGFP: the *D. rerio* ubiquitin promoter was excised from pUbi:Switch and

cloned into pT2Xla.Ef1a:EGFP using XhoI and BamHI.

3. pT2Xla.Ef1a:amhy,EGFP: the synthetic amhy,EGFP construct was excised from

pT2Dre.Ubi:amhy,EGFP and cloned into pT2Xla.Ef1a:EGFP using BamHI and BglII.

4. pT2-5amhy2.5,Xla.Ef1a:EGFP: an 11.5 kb region of the Y chromosome containing

amhy along with 5 kb of upstream and 2.5 kb downstream DNA was excised from

bacterial artificial chromosome STB26-N21 from the CHORI-215 library [141] and

cloned into pBT2 using BamHI and XhoI. The Xla.Ef1a:EGFP cassette was excised

from pT2Xla.Ef1a:EGFP by digestion with XhoI and HpaI and ligated into the

intermediate pT2-5amhy2.5 plasmid following digestion with XhoI and SwaI.

We synthesized Tol2 mRNA using the mMessage mMachine SP6 Transcription kit

(Invitrogen AM1340) and purified the product with the MEGAclear Transcription Clean-Up Kit

(Invitrogen AM1908). We injected stickleback embryos were as in [40] with the addition of KCl to

a final concentration of 0.2 M [136]. All embryos were screened for EGFP expression five to

seven days after injections before hatching had occurred. We removed all embryos without

discernable somatic EGFP fluorescence.

Genotyping and sanger sequencing

We extracted DNA from fin clips or whole embryos using a HotSHOT isolation protocol

[142]. We performed all PCR reactions on an Analytik Jena Biometra Tone in a total volume of

20 µl using 0.8 µL of genomic DNA, 0.2 µM each primer, 0.2 mM each dNTPs, 0.5 units

DreamTaq DNA polymerase (Thermo Scientific EP), and 2 µl 10x DreamTaq Green Buffer with

20 mM $MgCl_2$. We genotyped fish for sex using the sex-linked marker *idh* [38] with the following

cycling conditions: 1 cycle of 95°C for 2 min; 30 cycles of 95°C for 30 s, 57.3°C for 30 s, and

72°C for 30 s; 1 cycle of 72°C for 5 minutes. All other reactions used: 1 cycle of 95°C for 2 min;

35 cycles of 95°C for 30 s, 60°C for 30 s, and 72°C for 30 s; 1 cycle of 72°C for 5 minutes.

Primer sequences for all reactions are listed in Table S4. We purified the PCR product using a

Monarch PCR & DNA cleanup kit (NEB #T1030). All samples were Sanger sequenced through

Azenta Life Sciences (South Plainfield, NJ, USA).

Phenotyping and crossing

To phenotype gonads, we euthanized adult stickleback fish at six- or twelve-months post

hatching in 0.05% MS-222 buffered to neutral. We dissected out the gonads and fixed one or

both gonads for 24 to 48 hours in 10% neutral buffered formalin. The remainder of the fish was

preserved in 75% ethanol. The gonads were processed, embedded in paraffin, sectioned at 5

µm, and stained with hematoxylin and eosin by the University of Georgia College of Veterinary

Medicine Histology Lab. We used a Zeiss Axio Scope A1 microscope and Zeiss Axiocam 305

Color camera to capture all images at the University of Georgia Biomedical Microscopy Core.

To make crosses, we euthanized mutant or wildtype male stickleback and gently

macerated testes in 500 µl of Hanks' Balanced Salt Solution. We stripped eggs from gravid

females and applied approximately 50 µl of sperm solution with a disposable transfer pipette.

We photographed adult stickleback with a Samsung Galaxy S22 under standardized lighting.

Riboprobe synthesis

We generated riboprobe templates for *gsdf, dmrt1, foxl2,* and *cyp10a1a* from cDNA from

adult gonads. We extracted total RNA from mature testis and ovary tissue using TRIzol Reagent

(Invitrogen 15596026) according to manufacturer's instructions. We synthesized cDNA using the

iScript cDNA Synthesis Kit (Bio-Rad 1708891) and amplified the targeted region for each gene

through PCR as described above (Table S4). We cloned the PCR product into plasmid pJC53.2

(kindly shared by Kendall Clay and Rachel Roberts-Galbraith) after digestion with Eam1105I

(Thermo Scientific FD0244) as in Collins *et al*. 2010 [143]. We verified each riboprobe template

after cloning with sanger sequencing by Azenta Life Sciences (South Plainfield, NJ, USA). For

amha, we ordered a synthetic gene fragment from TWIST Bioscience (San Francisco, CA, USA)

containing a 307 bp region of the *amha* cds flanked by a 5' T3 promoter and 3' T7 promoter to

facilitate riboprobe synthesis (File S2).

To synthesize riboprobes, we PCR amplified linear template from the appropriate

plasmid or used the synthetic gene fragment directly. We performed *in vitro* transcription from

these templates using T7 (New England Biolabs M0251S), T3 (New England Biolabs M0378S),

or SP6 (New England Biolabs M0207S) RNA polymerase according to manufacturer's

instructions, substituting 0.2 mM of UTP for DIG-labeled UTP (Roche 11209256910). We

purified riboprobes with a DNase treatment (New England Biolabs M0303S) and an ammonium

acetate precipitation.

In situ hybridization

We adapted a colorimetric *in situ* hybridization (ISH) protocol from Square, *et al.* 2021

[144]. We fixed the dissected gonads for 24-48 hours in 10% neutral buffered formalin. The

gonads were processed, embedded in paraffin, and sectioned at 5 μm by the University of

Georgia College of Veterinary Medicine Histology Lab. Unless otherwise stated, we performed

all steps in Lock-Mailer microscope slide jars in a volume of 9-11mL and at room temperature.

To deparaffinize, we warmed the slides in a 65°C incubator for five minutes, allowed them to

cool to room temperature, then washed the slides in Hemo-De once for 5 minutes then again for

10 minutes. We performed prehybridization washes indicated with an asterisk in a Tissue-Tek II

Slide Staining Station rather than Lock-Mailer jars: 100% EtOH for 5 minutes*, 80% EtOH for 10

minutes*, Milli-Q water for 10 minutes*, PBST for 5 minutes*, 15 ug/mL proteinase K in PBST

for 5 minutes, PBST rinse*, 4% PFA for 20 minutes, PBST twice for 10 minutes each*, 67°C pre-

heated hybridization buffer (50% formamide, 5X SSC, 0.1% Tween, 5mg/mL CHAPS, 0.1

mg/mL yeast RNA, 0.1mg/mL heparin, pH 6.0 with citric acid) twice for 5 minutes each, and pre-

heated hybridization buffer for 1 to 4 hours on a rotator at 67°C. We performed hybridization

with 100-500 ng/mL of riboprobe in hybridization buffer incubated overnight rotating at 67°C. We

stored hybridization solution at 20°C and reused up to 3 times.

Post-hybridization, we washed the slides 6 times with hybridization wash (50%

formamide, 5X SSC, 0.1% Tween) rotating at 67°C for 20-90 minutes each, totaling 5 to 6

hours. We rinsed the slides in MABT (maleic acid buffer with Tween) and washed them in MABT

for twice for 20 minutes each, first at 67°C, then at room temperature. We added blocking

solution (2% Blocking Reagent (Roche 11096176001) in MAB) to each slide, covered the slides

with parafilm, and incubated for 1 to 3 hours in a humidor. We poured off the blocking solution

and added 1:2000 anti-digoxigenin alkaline phosphate antibody (Roche 11093274910) in

blocking solution, covered the slides with parafilm, and left them in a humidor overnight at 4°C.

We performed five 20-50 minute post-antibody washes hybridization wash solution (50%

formamide, 5X SSC, 0.1% Tween, citric acid to pH 6.0) on a rotator for a total of three to four

hours, followed by an overnight MABT wash at 4°C.

Prior to coloration, we performed three 5-to-10-minute washes in NTMT (0.1 M NaCl, 0.1

M Tris pH 9.5, 0.05 M MgCl2, 0.1% Tween). We moved the slides into coloration solution

(blocking solution plus 25 ug/mL NBT and 175 ug/mL BCIP) and incubated 2 to 30 hours

covered to block light. We removed the slides from coloration solution once the color was stable

in the positive control (*gsdf*), but before any background color was seen in the negative control

(sense *gsdf*). Once coloration was complete, we rinsed the slides with PBST and washed for 10

minutes in PBST. We fixed the slides in 4% PFA in PBS at 4°C for one to five days.

To prepare slides for mounting, we performed three 5-minute washes in PBST and three

5-minute washes in deionized water. We mounted and coverslipped the slides using Vectashield

antifade mounting medium with DAPI. We used a Zeiss Axio Scope A1 microscope and Zeiss

Axiocam 305 Color camera to capture all images at the University of Georgia Biomedical

Microscopy Core.

Sequence Alignments

We retrieved sequences for *amha* (Gene ID: 120823390) and *amhy* (Gene ID:

120812167) from threespine assembly GAculeatus_UGA_version5 (GCF_016920845.1) and

amh (Gene ID: 119216504) from ninespine assembly fPunPun2.1 (GCA_949316345.1). We

performed all alignments and calculated sequence identity between *amh* homologues with

Muscle 5.1 [145] within Geneious Prime (v2024.0.5) using default parameters. We defined

intron and exon positions based upon the threespine *amha* sequence in each alignment. We

calculated dN/dS with the R package ape v5.8-1 [146] using the translation aligned coding

sequences of *amha* (XM_040183306.1)*, amhy* (XM_040167960.1), and ninespine *amh*

(XM_037469420.2). One additional base was included at the 5' end of exon 7 of the ninespine

amh cds in order to correct the predicted annotation which contains multiple premature stop

codons in this final exon. Alignment files are presented in File S2. We generated Vistaplots

[147,148] using LAGAN [149] alignments and a 100 bp window size with a 70% sequence

identity threshold.

RNAseq

We staged embryos according to established stickleback developmental stages [79] and

preserved the whole embryos in RNAlater Stabilization Solution (Invitrogen AM7021) following

manufacturer guidelines and stored at -80 °C until analysis. We extracted RNA using TRIzol

Reagent (Invitrogen 15596026) and the Direct-zol RNA Microprep Kit (Zymo R2062). We

created sequencing libraries using the Illumina TruSeq Stranded mRNA Library Prep kit

(20020594). We pooled the libraries from all samples and 150 bp paired end reads were

sequenced on two lanes using a NovaSeq X Plus by Azenta Life Sciences (South Plainfield, NJ,

USA). All of the samples were sequenced to an average of 34.7 M read depth (min 11.6 M, max

71.3 M, SD = 10.4 M) (File S1). We trimmed raw reads for quality and sequencing adapters with

Trimmomatic (v. 0.39) [150] using leading and trailing quality threshold of 3, a sliding window of

4 with quality threshold of 15, and minimum length cutoff of 36. We aligned paired and orphan

reads to the threespine stickleback reference genome (GAculeatus_UGA_version5, NCBI

RefSeq assembly GCF_016920845.1) using HISAT-3N [151] with default parameters. All

samples had an alignment rate between 73% and 92% (File S1). We generated Indexed BAM

files of the alignments with Samtools (v. 1.17). We generated transcript counts against the NCBI

annotation GTF file (release 100) for GCF_016920845.1 with HTseq (v. 2.0.2) [152] using

HTSeq-count with default parameters except --stranded=no. We performed all remaining

analyses in R (v. 4.4.1) and RStudio (v. 2024.04.2+764). We combined the counts from paired

and orphaned reads for each sample and used DESeq2 (v. 1.44.0) [153] to generate normalized

read counts and perform differential expression analysis. We identified DEGs using the standard

workflow with stage and sex combined as a single factor. We used the contrast function to

calculate log2 fold changes (logFC) and adjusted p values for differential expression between

males and females at each timepoint. We identified DEGs of interest using GO terms from the

NIH gene2go database (https://ftp.ncbi.nlm.nih.gov/gene/DATA/).

Material Availability

RNAseq data is available in the short reads archive under BioProject PRJNA1248589. All

scripts are available at https://github.com/MTreasterUGA/Amhy-sex-determination. All plasmids

are available upon request.

Acknowledgements

This work was funded by the National Science Foundation MCB 1943283 (M.A.W), the National

Institutes of Health 1T32GM142623 (M.J.T.), The University of Georgia Research Foundation

(M.J.T.), the Society for the Study of Evolution R.C. Lewontin Early Award (M.J.T.), and ARCS

foundation (M.J.T.).

Research reported in this publication was supported by the National Institute of General Medical

Sciences of the National Institutes of Health under Award Number R01GM147312 (M.A.W). The

content is solely the responsibility of the authors and does not necessarily represent the official

views of the National Institutes of Health.

This study was supported in part by resources and technical expertise from the Georgia

Advanced Computing Resource Center, a partnership between the University of Georgia's

Office of the Vice President for Research and Office of the Vice President for Information

Technology.

The imaging data used in this publication was produced in collaboration with the Biomedical

Microscopy Core at the University of Georgia.

We thank Koichi Kawakami (National Institute of Genetics, Japan) for use of the Tol2 elements

in this study.

743   Declaration of Interests

The authors declare no conflicts of interests

Figures and tables

Fig 1. CRISPR/Cas9 knockout of *amhy* caused male to female sex reversal in F1 XY fish.

We used CRISPR/Cas9 to target exon 1 (A) or exon 2 (B) of *amhy*. CRISPR target site and

PAM sequence are indicated by orange lines above wildtype reference sequence. Mutations

relative to wildtype are bolded. We identified four loss of function *amhy* alleles in F1 progeny.

One allele (-2282,+8) deleted the first six exons of *amhy*. Other alleles featured small frameshift

indels. C) We identified male to female sex reversal using gonad morphology and histology.

Ovaries in XX wildtype and XY *amhy*-KO fish feature oocytes throughout multiple stages of

development. P – primary oocyte, C – cortical alveolus stage V – vitellogenic stage, M – Mature

oocyte, G – Granulosa cells, T – Theca cells. Scale bars = 200 µm.

Fig 2. Gene expression in sex reversed gonads resembles expression in wild type gonads.

In situ hybridization of gonadal genes in wildtype and sex reversed stickleback gonads. A

positive control gene *gsdf* was expressed in somatic cells of both testes and ovaries. Wildtype

XY and sex reversed XX testes show strong expression of male differentiation genes *amha* and

dmrt1 but not female differentiaiton genes *foxl2* and *cyp19a1a*. Wildtype XX and sex reversed

XY ovaries strongly express female differentiation genes but not male differentiation genes.

Scale bars = 200 µm.

Fig 3. XX fish with an *amhy* transgene exhibit female to male sex reversal.

A) XX *Tg(-5amhy3.5,Xla.Ef1a:EGFP)* fish developed testes with varying cell composition. At 12 months post hatching, fish had either normal testes with mature sperm (middle) or hypotrophic testes without discernable germ cells (right). G – spermatogonia, C – spermatocyte, R – round spermatid, Z – Spermatozoa, L – Leydig cells, S – Sertoli cells Scale bars = 200 µm. B) Control XX *Tg(Xla.Ef1a:EGFP)* fish developed ovaries identical to wildtype XX counterparts. Scale bars = 200 µm. C) Embryos injected with pT2-5amhy3.5,Xla.Ef1a:EGFP seven days after fertilization under bright field (top) and EGFP excitation (bottom). Embryos positive for transgene integration show somatic EGFP expression. We removed embryos with no EGFP expression or EGFP

expression only in the yolk, which indicates the yolk was injected rather than the blastomere

[40]. D) We used PCR of the sex-linked marker *idh* to genotype XX and XY fish. PCR of EGFP

and *amhy* show the presence of the associated transgene. 1 – wildtype XX, 2 – wildtype XY, 3 –

XX *Tg(EGFP)*, 4 – XX *Tg(amhy,EGFP)*.

Fig 4. XY *amhy*-KO females can reproduce and transmit an X or Y chromosome.

YY embryos (A) at 48 hours post fertilization show arrested development compared to wildtype

embryos (B). Arrowheads denote anterior end of body axis. C) F2 XX and XY *amhy*-KO female

siblings do not show a significant difference in clutch size (wilcoxon test, p < 0.05). Clutches

from XY *amhy*-KO females have a significantly lower viability rate than those from XX female

siblings (z test, p <0.05). D) PCR of the sex linked marker IDH shows the absence of the X

chromosome in YY embryos. Both the paternal (wildtype) and maternal (+9,+5) loss of function

alleles can be detected in XY offspring. YY embryos show the presence of both amhy alleles as

well as a third band, likely a heterodimer of the maternal and paternal amplicons.

XX WT

XY WT

XX
Tg(amhy,EGFP)

XY
amhy-KO

Fig 5. Both male and female sex reversed stickleback show partial male nuptial coloration.

Wildtype and mutant threespine stickleback at one year post hatching. Wildtype XX females are

silver and may darken slightly when gravid. Wildtype XY males have dark blue eyes and bodies

and red throats. Sex reversed XX *Tg(amhy,EGFP)* males are less dark than wildtype males but

show blue coloration on the body and eyes. Sex reversed XY *amhy*-KO females show slight or

moderate darkening of the body with distinct blue eyes not typical of wildtype XX females.

Fig 6. *amha* and *amhy* show no sequence conservation outside of the coding sequence.

Vistaplots showing conserved sequence identity of threespine *amha* to threespine *amhy* (top)

and to ninespine *amh* (bottom) with sequence identity on the y axis and nucleotide position of

threespine *amha* on the x axis. Exon positions are indicated along the x axis in blue. Conserved

exon and noncoding sequences share 70% sequence identity in a 100 bp window.

Fig 7. Expression of gonadal growth and differentiation genes in embryonic and larval

stickleback.

(A) PCA plots of normalized read counts for all genes cluster by stage rather than sex. (B) and

(C) Normalized read counts of genes in embryonic stages 17, 20, 23, and larvae 0, 4, and 8

days after hatching. Stages with significant differences between males and females are marked

with an asterisk and bolded (Benjamini Hochberg adjusted p value < 0.05; Table 2). The orange

arrow (B) indicates stages where differences in germ counts have been observed between

males and females.

Table 1. Sequence conservation of the threespine *amha* to threespine *amhy* and ninespine *amh.*

Threespine *amha* and threespine *amhy*											
	Exon 1	Exon 2	Exon 3	Exon 4	Exon 5	Exon 6	Exon 7	CDS	Introns	2kb Upstream	2kb Downstream
Sequence identity	0.760	0.682	0.812	0.866	0.790	0.846	0.742	0.780	0.286	0.365	0.376
dN/dS	0.583	0.362	0.512	0.803	0.651	0.373	0.723	0.534	-	-	-
Threespine *amha* and ninespine *amh*											
	Exon 1	Exon 2	Exon 3	Exon 4	Exon 5	Exon 6	Exon 7	CDS	Introns	2kb Upstream	2kb Downstream
Sequence identity	0.791	0.769	0.890	0.920	0.870	0.905	0.936	0.879	0.668	0.595	0.660
dN/dS	0.597	0.425	1.003	0.441	0.266	0.565	0.278	0.467	-	-	-

821   Table 2. Benjamini Hochberg adjusted P values for differential expression between males and

822   females.

gene	gene id	stage 17	stage 20	stage 23	0 dah	4 dah	8 dah
amha	amh	1	1	0.679	1	1	1
amhrII	LOC120835457	0.952	1	0.322	1	0.509	**0.0280**
ddx4	ddx4	1	1	1	1	1	**0.00193**
piwil1	piwil1	1	1	1	1	1	**0.00403**
hsd17b1	hsd17b1	1	1	1	0.8681	1	**0.0238**
foxl2a	foxl2a	0.788	1	1	1	1	1
cyp19a1a	LOC120811580	0.662	1	1	1	1	1
dmrt1	dmrt1	1	1	1	1	1	1
sox9b	LOC120827768	0.930	1	1	1	1	1
gsdf	gsdf	1	1	0.467	1	1	1
nr5a1	nr5a1a	1	1	1	1	1	0.983

825   Table 3. Phenotypes of mutant stickleback from this study

Genotype	N	Male (%)	Female (%)
F0 XY *amhy* crispants	17	9 (53%)	8 (47%)
F1 XY *amhy*-KO	16	0 (0%)	16 (100%)
F0 XX *Tg(-5amhy2.5,Xla.Ef1a:EGFP)*	16	16 (100%)	0 (0%)
F0 XX *Tg(Xla.Ef1a:EGFP)*	10	0 (0%)	10 (100%)

References

1. Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman T-L, et al. Sex
Determination: Why So Many Ways of Doing It? PLoS Biol. 2014;12: e1001899.
doi:10.1371/journal.pbio.1001899

2. Capel B. Vertebrate sex determination: evolutionary plasticity of a fundamental switch.
Nat Rev Genet. 2017;18: 675–689. doi:10.1038/nrg.2017.60

3. Weber C, Capel B. Sex determination without sex chromosomes. Philosophical
Transactions of the Royal Society B: Biological Sciences. 2021;376: 20200109.
doi:10.1098/rstb.2020.0109

4. Charlesworth D, Charlesworth B, Marais G. Steps in the evolution of heteromorphic sex
chromosomes. Heredity. 2005;95: 118–128. doi:10.1038/sj.hdy.6800697

5. Charlesworth B, Charlesworth D. A Model for the Evolution of Dioecy and Gynodioecy.
The American Naturalist. 1978;112: 975–997. doi:10.1086/283342

6. Ponnikas S, Sigeman H, Abbott JK, Hansson B. Why Do Sex Chromosomes Stop
Recombining? Trends in Genetics. 2018;34: 492–503. doi:10.1016/j.tig.2018.04.001

7. Pan Q, Kay T, Depincé A, Adolfi M, Schartl M, Guiguen Y, et al. Evolution of master sex
determiners: TGF-β signalling pathways at regulatory crossroads. Philosophical
Transactions of the Royal Society B: Biological Sciences. 2021;376: 20200091.
doi:10.1098/rstb.2020.0091

8. Takehana Y, Matsuda M, Myosho T, Suster ML, Kawakami K, Shin-I T, et al. Co-option of
Sox3 as the male-determining factor on the Y chromosome in the fish Oryzias dancena.
Nat Commun. 2014;5: 4157. doi:10.1038/ncomms5157

9. Shan Z, Nanda I, Wang Y, Schmid M, Vortkamp A, Haaf T. Sex-specific expression of an
evolutionarily conserved male regulatory gene, DMRT1, in birds. Cytogenetics and Cell
Genetics. 2000;89: 252–257. doi:10.1159/000015626

10. Yoshimoto S, Okada E, Umemoto H, Tamura K, Uno Y, Nishida-Umehara C, et al. A W-
linked DM-domain gene, DM-W, participates in primary ovary development in Xenopus
laevis. Proceedings of the National Academy of Sciences. 2008;105: 2469–2474.
doi:10.1073/pnas.0712244105

11. Matsuda M, Nagahama Y, Shinomiya A, Sato T, Matsuda C, Kobayashi T, et al. DMY is a
Y-specific DM-domain gene required for male development in the medaka fish. Nature.
2002;417: 559–563. doi:10.1038/nature751

12. Nanda I, Kondo M, Hornung U, Asakawa S, Winkler C, Shimizu A, et al. A duplicated copy
of DMRT1 in the sex-determining region of the Y chromosome of the medaka, Oryzias

latipes. Proceedings of the National Academy of Sciences. 2002;99: 11778–11783.
doi:10.1073/pnas.182314699

13.   Hattori RS, Murai Y, Oura M, Masuda S, Majhi SK, Sakamoto T, et al. A Y-linked anti-
Mullerian hormone duplication takes over a critical role in sex determination. Proceedings
of the National Academy of Sciences. 2012;109: 2955–2959.
doi:10.1073/pnas.1018392109

14.   Hattori RS, Somoza GM, Fernandino JI, Colautti DC, Miyoshi K, Gong Z, et al. The
Duplicated Y-specific amhy Gene Is Conserved and Linked to Maleness in Silversides of
the Genus Odontesthes. Genes. 2019;10: 679. doi:10.3390/genes10090679

15.   Hattori RS, Kumazawa K, Nakamoto M, Nakano Y, Yamaguchi T, Kitano T, et al. Y-specific
amh allele, amhy, is the master sex-determining gene in Japanese flounder Paralichthys
olivaceus. Frontiers in Genetics. 2022;13. Available:
https://www.frontiersin.org/articles/10.3389/fgene.2022.1007548

16.   Li M, Sun Y, Zhao J, Shi H, Zeng S, Ye K, et al. A Tandem Duplicate of Anti-Müllerian
Hormone with a Missense SNP on the Y Chromosome Is Essential for Male Sex
Determination in Nile Tilapia, Oreochromis niloticus. Naruse K, editor. PLoS Genet.
2015;11: e1005678. doi:10.1371/journal.pgen.1005678

17.   Rondeau EB, Laurie CV, Johnson SC, Koop BF. A PCR assay detects a male-specific
duplicated copy of Anti-Müllerian hormone (amh) in the lingcod (Ophiodon elongatus).
BMC Research Notes. 2016;9: 230. doi:10.1186/s13104-016-2030-6

18.   Bej DK, Miyoshi K, Hattori RS, Strüssmann CA, Yamamoto Y. A Duplicated, Truncated
amh Gene Is Involved in Male Sex Determination in an Old World Silverside. G3
Genes|Genomes|Genetics. 2017;7: 2489–2495. doi:10.1534/g3.117.042697

19.   Pan Q, Feron R, Yano A, Guyomard R, Jouanno E, Vigouroux E, et al. Identification of the
master sex determining gene in Northern pike (Esox lucius) reveals restricted sex
chromosome differentiation. PLOS Genetics. 2019;15: e1008013.
doi:10.1371/journal.pgen.1008013

20.   Peichel CL, McCann SR, Ross JA, Naftaly AFS, Urton JR, Cech JN, et al. Assembly of
the threespine stickleback Y chromosome reveals convergent signatures of sex
chromosome evolution. Genome Biol. 2020;21: 177. doi:10.1186/s13059-020-02097-x

21.   Song W, Xie Y, Sun M, Li X, Fitzpatrick CK, Vaux F, et al. A duplicated amh is the master
sex-determining gene for Sebastes rockfish in the Northwest Pacific. Open Biology.
2021;11: 210063. doi:10.1098/rsob.210063

22.   Jeffries DL, Mee JA, Peichel CL. Identification of a candidate sex determination gene in
Culaea inconstans suggests convergent recruitment of an Amh duplicate in two lineages
of stickleback. Journal of Evolutionary Biology. 2022;n/a. doi:10.1111/jeb.14034

23.   Zhao H, Li J, Xiao Z, Xiao Y. Dose-dependent role of AMH and AMHR2 signaling in male
differentiation and regulation of sex determination in Spotted knifejaw (Oplegnathus
punctatus) with X1X1X2X2/X1X2Y chromosome system. Cell Communication and
Signaling. 2025;23: 59. doi:10.1186/s12964-025-02038-w

24. Cortez D, Marin R, Toledo-Flores D, Froidevaux L, Liechti A, Waters PD, et al. Origins
and functional evolution of Y chromosomes across mammals. Nature. 2014;508: 488–
493. doi:10.1038/nature13151

25. Mullen RD, Ontiveros AE, Moses MM, Behringer RR. AMH and AMHR2 mutations: A
spectrum of reproductive phenotypes across vertebrate species. Developmental Biology.
2019;455: 1–9. doi:10.1016/j.ydbio.2019.07.006

26. Howard JA, Hart KN, Thompson TB. Molecular Mechanisms of AMH Signaling. Frontiers
in Endocrinology. 2022;13. Available:
https://www.frontiersin.org/articles/10.3389/fendo.2022.927824

27. Baarends WM, Helmond MJL van, Post M, Schoot PJCM van der, Hoogerbrugge JW,
Winter JP de, et al. A novel member of the transmembrane serine/threonine kinase
receptor family is specifically expressed in the gonads and in mesenchymal cells adjacent
to the müllerian duct. Development. 1994;120: 189–197. doi:10.1242/dev.120.1.189

28. Josso N, di Clemente N, Gouédard L. Anti-Müllerian hormone and its receptors.
Molecular and Cellular Endocrinology. 2001;179: 25–32. doi:10.1016/S0303-
7207(01)00467-1

29. Mishina Y, Whitworth DJ, Racine C, Behringer RR. High Specificity of Müllerian-Inhibiting
Substance Signaling in Vivo*. Endocrinology. 1999;140: 2084–2088.
doi:10.1210/endo.140.5.6705

30. Jost A. The Age Factor in the Castration of Male Rabbit Fetuses. Proceedings of the
Society for Experimental Biology and Medicine. 1947;66: 302–303.
doi:10.3181/00379727-66-16071

31. Morinaga C, Saito D, Nakamura S, Sasaki T, Asakawa S, Shimizu N, et al. The hotei
mutation of medaka in the anti-Müllerian hormone receptor causes the dysregulation of
germ cell and sexual development. PNAS. 2007;104: 9691–9696.
doi:10.1073/pnas.0611379104

32. Pfennig F, Standke A, Gutzeit HO. The role of Amh signaling in teleost fish – Multiple
functions not restricted to the gonads. General and Comparative Endocrinology.
2015;223: 87–107. doi:10.1016/j.ygcen.2015.09.025

33. Yan Y-L, Batzel P, Titus T, Sydes J, Desvignes T, BreMiller R, et al. A Hormone That Lost
Its Receptor: Anti-Müllerian Hormone (AMH) in Zebrafish Gonad Development and Sex
Determination. Genetics. 2019;213: 529–553. doi:10.1534/genetics.119.302365

34. Ross JA, Urton JR, Boland J, Shapiro MD, Peichel CL. Turnover of Sex Chromosomes in
the Stickleback Fishes (Gasterosteidae). PLOS Genetics. 2009;5: e1000391.
doi:10.1371/journal.pgen.1000391

35. Dixon G, Kitano J, Kirkpatrick M. The Origin of a New Sex Chromosome by Introgression
between Two Stickleback Fishes. Molecular Biology and Evolution. 2019;36: 28–38.
doi:10.1093/molbev/msy181

36.   Natri HM, Merilä J, Shikano T. The evolution of sex determination associated with a
941   chromosomal inversion. Nat Commun. 2019;10: 145. doi:10.1038/s41467-018-08014-y

942   37.   Sardell JM, Josephson MP, Dalziel AC, Peichel CL, Kirkpatrick M. Heterogeneous
943   Histories of Recombination Suppression on Stickleback Sex Chromosomes. Molecular
944   Biology and Evolution. 2021;38: 4403–4418. doi:10.1093/molbev/msab179

945   38.   Peichel CL, Ross JA, Matson CK, Dickson M, Grimwood J, Schmutz J, et al. The Master
946   Sex-Determination Locus in Threespine Sticklebacks Is on a Nascent Y Chromosome.
947   Current Biology. 2004;14: 1416–1424. doi:10.1016/j.cub.2004.08.030

948   39.   Pigott GC, Akel MA, Rogers MGQ, Flanagan ME, Marlette EG, Treaster MJ, et al.
949   Evidence for Variation in the Genetic Basis of Sex Determination in Brook Stickleback
950   (Culaea inconstans). Ecology and Evolution. 2025;15: e70955. doi:10.1002/ece3.70955

951   40.   Erickson PA, Ellis NA, Miller CT. Microinjection for Transgenesis and Genome Editing in
952   Threespine Sticklebacks. J Vis Exp. 2016 [cited 30 Jan 2021]. doi:10.3791/54055

953   41.   Nath S, Shaw DE, White MA. Improved contiguity of the threespine stickleback genome
954   using long-read sequencing. G3 Genes|Genomes|Genetics. 2021;11.
955   doi:10.1093/g3journal/jkab007

956   42.   Aguirre WE, Ellis KE, Kusenda M, Bell MA. Phenotypic variation and sexual dimorphism
957   in anadromous threespine stickleback: implications for postglacial adaptive radiation.
958   Biological Journal of the Linnean Society. 2008;95: 465–478. doi:10.1111/j.1095-
959   8312.2008.01075.x

960   43.   Leinonen T, Cano JM, Merilä J. Genetic basis of sexual dimorphism in the threespine
961   stickleback Gasterosteus aculeatus. Heredity. 2011;106: 218–227.
962   doi:10.1038/hdy.2010.104

963   44.   Kitano J, Mori S, Peichel CL. Sexual Dimorphism in the External Morphology of the
964   Threespine Stickleback (Gasterosteus Aculeatus). Copeia. 2007;2007: 336–349.
965   doi:10.1643/0045-8511(2007)7[336:SDITEM]2.0.CO;2

966   45.   Aguirre WE, Akinpelu O. Sexual dimorphism of head morphology in three-spined
967   stickleback Gasterosteus aculeatus. Journal of Fish Biology. 2010;77: 802–821.
968   doi:10.1111/j.1095-8649.2010.02705.x

969   46.   Toli EA, Noreikiene K, DeFaveri J, Merilä J. Environmental enrichment, sexual
970   dimorphism, and brain size in sticklebacks. Ecology and Evolution. 2017;7: 1691–1698.
971   doi:10.1002/ece3.2717

972   47.   Kotrschal A, Räsänen K, Kristjánsson BK, Senn M, Kolm N. Extreme Sexual Brain Size
973   Dimorphism in Sticklebacks: A Consequence of the Cognitive Challenges of Sex and
974   Parenting? PLOS ONE. 2012;7: e30055. doi:10.1371/journal.pone.0030055

975   48.   McGee MD, Wainwright PC. Sexual dimorphism in the feeding mechanism of threespine
976   stickleback. Journal of Experimental Biology. 2013;216: 835–840. doi:10.1242/jeb.074948

977   49.   Whoriskey FG, FitzGerald GJ. Sex, cannibalism and sticklebacks. Behav Ecol Sociobiol.
978   1985;18: 15–18. doi:10.1007/BF00299233

979   50.   King AJ, Fürtbauer I, Mamuneas D, James C, Manica A. Sex-Differences and Temporal
980   Consistency in Stickleback Fish Boldness. PLOS ONE. 2013;8: e81116.
981   doi:10.1371/journal.pone.0081116

982   51.   FitzGerald GJ. The Reproductive Behavior of the Stickleback. Scientific American.
983   1993;268: 80–85.

984   52.   Bell MA, Foster SA. The Evolutionary Biology of the Threespine Stickleback. The Journal
985   of Animal Ecology. 1994;64: 418. doi:10.2307/5902

986   53.   Pressley PH. Parental Effort and the Evolution of Nest-Guarding Tactics in the Threespine
987   Stickleback, Gasterosteus aculeatus L. Evolution. 1981;35: 282–295.
988   doi:10.2307/2407838

989   54.   Bakker TCM, Künzler R, Mazzi D. Condition-related mate choice in sticklebacks. Nature.
990   1999;401: 234–234. doi:10.1038/45727

991   55.   Milinski M, Bakker TCM. Female sticklebacks use male coloration in mate choice and
992   hence avoid parasitized males. Nature. 1990;344: 330–333. doi:10.1038/344330a0

993   56.   Conte GL, Schluter D. Experimental Confirmation That Body Size Determines Mate
994   Preference via Phenotype Matching in a Stickleback Species Pair. Evolution. 2013;67:
995   1477–1484. doi:10.1111/evo.12041

996   57.   Andreou D, Eizaguirre C, Boehm T, Milinski M. Mate choice in sticklebacks reveals that
997   immunogenes can drive ecological speciation. Behavioral Ecology. 2017;28: 953–961.
998   doi:10.1093/beheco/arx074

999   58.   Milinski M. The function of mate choice in sticklebacks: optimizing Mhc genetics. Journal
1000   of Fish Biology. 2003;63: 1–16. doi:10.1111/j.1095-8649.2003.00215.x

1001   59.   Flamarique IN, Bergstrom C, Cheng CL, Reimchen TE. Role of the iridescent eye in
1002   stickleback female mate choice. Journal of Experimental Biology. 2013;216: 2806–2812.
1003   doi:10.1242/jeb.084889

1004   60.   Barker TCM, Milinski M. The advantages of being red: Sexual selection in the stickleback.
1005   Marine & Freshwater Behaviour & Phy. 1993 [cited 21 Apr 2025].
1006   doi:10.1080/10236249309378870

1007   61.   Candolin U. Reproduction under predation risk and the trade–off between current and
1008   future reproduction in the threespine stickleback. Proceedings of the Royal Society of
1009   London Series B: Biological Sciences. 1998;265: 1171–1175.
1010   doi:10.1098/rspb.1998.0415

1011   62.   Johnson S, Candolin U. Predation cost of a sexual signal in the threespine stickleback.
1012   Behavioral Ecology. 2017;28: 1160–1165. doi:10.1093/beheco/arx080

63. Borg B, Mayer I, Hoffmann E, Österman A. 11-ketotestosterone is not responsible for the
entire testicular effect on male reproductive behaviour in the threespine stickleback.
Behav. 2008;145: 509–525. doi:10.1163/156853908792451548

64. Shibata Y, Paul-Prasanth B, Suzuki A, Usami T, Nakamoto M, Matsuda M, et al.
Expression of gonadal soma derived factor (GSDF) is spatially and temporally correlated
with early testicular differentiation in medaka. Gene Expression Patterns. 2010;10: 283–
289. doi:10.1016/j.gep.2010.06.005

65. Guo Y, Cheng H, Huang X, Gao S, Yu H, Zhou R. Gene structure, multiple alternative
splicing, and expression in gonads of zebrafish *Dmrt1*. Biochemical and Biophysical
Research Communications. 2005;330: 950–957. doi:10.1016/j.bbrc.2005.03.066

66. von Hofsten J, Larsson A, Olsson P e. Novel steroidogenic factor-1 homolog (ff1d) is
coexpressed with anti-Mullerian hormone (AMH) in zebrafish. Developmental Dynamics.
2005;233: 595–604. doi:10.1002/dvdy.20335

67. Johnsen H, Seppola M, Torgersen JS, Delghandi M, Andersen Ø. Sexually dimorphic
expression of *dmrt1* in immature and mature Atlantic cod (*Gadus morhua* L.).
Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology.
2010;156: 197–205. doi:10.1016/j.cbpb.2010.03.009

68. von Schalburg KR, Gowen BE, Rondeau EB, Johnson NW, Minkley DR, Leong JS, et al.
Sex-specific expression, synthesis and localization of aromatase regulators in one-year-
old Atlantic salmon ovaries and testes. Comparative Biochemistry and Physiology Part B:
Biochemistry and Molecular Biology. 2013;164: 236–246. doi:10.1016/j.cbpb.2013.01.004

69. Kossack ME, Draper BW. Genetic regulation of sex determination and maintenance in
zebrafish (Danio rerio). Curr Top Dev Biol. 2019;134: 119–149.
doi:10.1016/bs.ctdb.2019.02.004

70. Urasaki A, Morvan G, Kawakami K. Functional Dissection of the Tol2 Transposable
Element Identified the Minimal cis-Sequence and a Highly Repetitive Sequence in the
Subterminal Region Essential for Transposition. Genetics. 2006;174: 639–649.
doi:10.1534/genetics.106.060244

71. Kawakami K, Takeda H, Kawakami N, Kobayashi M, Matsuda N, Mishina M. A
Transposon-Mediated Gene Trap Approach Identifies Developmentally Regulated Genes
in Zebrafish. Developmental Cell. 2004;7: 133–144. doi:10.1016/j.devcel.2004.06.005

72. Kim JH, Lee S-R, Li L-H, Park H-J, Park J-H, Lee KY, et al. High Cleavage Efficiency of a
2A Peptide Derived from Porcine Teschovirus-1 in Human Cell Lines, Zebrafish and Mice.
PLOS ONE. 2011;6: e18556. doi:10.1371/journal.pone.0018556

73. Liu Z, Chen O, Wall JBJ, Zheng M, Zhou Y, Wang L, et al. Systematic comparison of 2A
peptides for cloning multi-genes in a polycistronic vector. Scientific Reports. 2017;7:
2193. doi:10.1038/s41598-017-02460-2

74. Mosimann C, Kaufman CK, Li P, Pugach EK, Tamplin OJ, Zon LI. Ubiquitous transgene
expression and Cre-based recombination driven by the ubiquitin promoter in zebrafish.
Development. 2011;138: 169–177. doi:10.1242/dev.059345

75.   Ueda Y, Mizuno N, Araki M. Transgenic Xenopus laevis with the ef1-α promoter as an
1054   experimental tool for amphibian retinal regeneration study. genesis. 2012;50: 642–650.
1055   doi:10.1002/dvg.22024

1056   76.   Wootton RJ. A Functional Biology of Sticklebacks. 1st ed. University of California Press;
1057   1984. doi:10.2307/jj.8306147

1058   77.   Varadharajan S, Rastas P, Löytynoja A, Matschiner M, Calboli FCF, Guo B, et al. A High-
1059   Quality Assembly of the Nine-Spined Stickleback (Pungitius pungitius) Genome. Genome
1060   Biology and Evolution. 2019;11: 3291–3308. doi:10.1093/gbe/evz240

1061   78.   Bachtrog D. Y-chromosome evolution: emerging insights into processes of Y-chromosome
1062   degeneration. Nat Rev Genet. 2013;14: 113–124. doi:10.1038/nrg3366

1063   79.   Swarup H. Stages in the Development of the Stickleback Gasterosteus aculeatus (L.).
1064   Development. 1958;6: 373–383.

1065   80.   Lewis ZR, McClellan MC, Postlethwait JH, Cresko WA, Kaplan RH. Female-specific
1066   increase in primordial germ cells marks sex differentiation in threespine stickleback (
1067   *Gasterosteus aculeatus*). J Morphol. 2008;269: 909–921. doi:10.1002/jmor.10608

1068   81.   Shi D-L. Interplay of RNA-binding proteins controls germ cell development in zebrafish.
1069   Journal of Genetics and Genomics. 2024;51: 889–899. doi:10.1016/j.jgg.2024.06.020

1070   82.   Planells B, Gómez-Redondo I, Pericuesta E, Lonergan P, Gutiérrez-Adán A. Differential
1071   isoform expression and alternative splicing in sex determination in mice. BMC Genomics.
1072   2019;20: 202. doi:10.1186/s12864-019-5572-x

1073   83.   He W, Gauri M, Li T, Wang R, Lin S-X. Current knowledge of the multifunctional 17β-
1074   hydroxysteroid dehydrogenase type 1 (HSD17B1). Gene. 2016;588: 54–61.
1075   doi:10.1016/j.gene.2016.04.031

1076   84.   Koyama T, Nakamoto M, Morishima K, Yamashita R, Yamashita T, Sasaki K, et al. A SNP
1077   in a Steroidogenic Enzyme Is Associated with Phenotypic Sex in Seriola Fishes. Current
1078   Biology. 2019;29: 1901-1909.e8. doi:10.1016/j.cub.2019.04.069

1079   85.   Desaulniers AT, Cederberg RA, Lents CA, White BR. Expression and Role of
1080   Gonadotropin-Releasing Hormone 2 and Its Receptor in Mammals. Front Endocrinol.
1081   2017;8. doi:10.3389/fendo.2017.00269

1082   86.   Sarraj MA, Sarraj ,Mai A., Chua ,Hui Kheng, Sarraj ,Mai A., Chua ,Hui Kheng, Umbers
1083   ,Alexandra, et al. Differential expression of TGFBR3 (betaglycan) in mouse ovary and
1084   testis during gonadogenesis. Growth Factors. 2007;25: 334–345.
1085   doi:10.1080/08977190701833619

1086   87.   Wickramasinghe D, Becker S, Ernst MK, Resnick JL, Centanni JM, Tessarollo L, et al.
1087   Two CDC25 homologues are differentially expressed during mouse development.
1088   Development. 1995;121: 2047–2056. doi:10.1242/dev.121.7.2047

1089   88.   Namgoong S, Kim N-H. Meiotic spindle formation in mammalian oocytes: implications for
1090   human infertility†. Biology of Reproduction. 2018;98: 153–161. doi:10.1093/biolre/iox145

1091   89.   Strüssmann CA, Saito T, Usui M, Yamada H, Takashima F. Thermal thresholds and critical
1092   period of thermolabile sex determination in two atherinid fishes, Odontesthes bonariensis
1093   and Patagonina hatcheri. Journal of Experimental Zoology. 1997;278: 167–177.
1094   doi:10.1002/(SICI)1097-010X(19970615)278:3<167::AID-JEZ6>3.0.CO;2-M

1095   90.   Baroiller JF, D'Cotta H, Bezault E, Wessels S, Hoerstgen-Schwark G. Tilapia sex
1096   determination: Where temperature and genetics meet. Comparative Biochemistry and
1097   Physiology Part A: Molecular & Integrative Physiology. 2009;153: 30–38.
1098   doi:10.1016/j.cbpa.2008.11.018

1099   91.   Sun P, You F, Ma D, Li J, Zhang P. Sex steroid changes during temperature-induced
1100   gonadal differentiation in (Temminck & Schegel, 1846). Journal of Applied Ichthyology.
1101   2013;29: 886–890. doi:10.1111/jai.12128

1102   92.   Borg B, van den Hurk R. Oocytes in the Testes of the Three-Spined Stickleback,
1103   Gasterosteus aculeatus. Copeia. 1983;1983: 259–261. doi:10.2307/1444727

1104   93.   Gercken J, Sordyl H. Intersex in feral marine and freshwater fish from northeastern
1105   Germany. Marine Environmental Research. 2002;54: 651–655. doi:10.1016/S0141-
1106   1136(02)00156-3

1107   94.   Aharon D, Marlow FL. Sexual determination in zebrafish. Cell Mol Life Sci. 2021;79: 8.
1108   doi:10.1007/s00018-021-04066-4

1109   95.   Slanchev K, Stebler J, de la Cueva-Méndez G, Raz E. Development without germ cells:
1110   The role of the germ line in zebrafish sex differentiation. Proceedings of the National
1111   Academy of Sciences. 2005;102: 4074–4079. doi:10.1073/pnas.0407475102

1112   96.   Tzung K-W, Goto R, Saju JM, Sreenivasan R, Saito T, Arai K, et al. Early Depletion of
1113   Primordial Germ Cells in Zebrafish Promotes Testis Formation. Stem Cell Reports.
1114   2015;4: 61–73. doi:10.1016/j.stemcr.2014.10.011

1115   97.   Schartl M. A comparative view on sex determination in medaka. Mechanisms of
1116   Development. 2004;121: 639–645. doi:10.1016/j.mod.2004.03.001

1117   98.   Kurokawa H, Saito D, Nakamura S, Katoh-Fukui Y, Ohta K, Baba T, et al. Germ cells are
1118   essential for sexual dimorphism in the medaka gonad. Proceedings of the National
1119   Academy of Sciences. 2007;104: 16958–16963. doi:10.1073/pnas.0609932104

1120   99.   Nakamura S, Watakabe I, Nishimura T, Picard J-Y, Toyoda A, Taniguchi Y, et al.
1121   Hyperproliferation of mitotically active germ cells due to defective anti-Müllerian hormone
1122   signaling mediates sex reversal in medaka. Development. 2012;139: 2283–2287.
1123   doi:10.1242/dev.076307

1124   100.   Tezak B, Straková B, Fullard DJ, Dupont S, McKey J, Weber C, et al. Higher
1125   temperatures directly increase germ cell number, promoting feminization of red-eared
1126   slider turtles. Current Biology. 2023;0. doi:10.1016/j.cub.2023.06.008

1127   101.   Lyet L, Louis F, Forest MG, Josso N, Behringer RR, Vigier B. Ontogeny of Reproductive
1128   Abnormalities Induced by Deregulation of Anti-Müllerian Hormone Expression in

Transgenic Mice. Biology of Reproduction. 1995;52: 444–454.
doi:10.1095/biolreprod52.2.444

102. Han Y, Zhao M, Wang L, Yu Z, Wang J, Yu Q, et al. Overexpression of Anti-müllerian
Hormone Gene in vivo Affects Gonad Sex Differentiation in Undifferentiated Orange-
Spotted Groupers (Epinephelus coioides). Front Endocrinol. 2019;0.
doi:10.3389/fendo.2019.00210

103. Rehman Z ur, Worku T, Davis JS, Talpur HS, Bhattarai D, Kadariya I, et al. Role and
mechanism of AMH in the regulation of Sertoli cells in mice. The Journal of Steroid
Biochemistry and Molecular Biology. 2017;174: 133–140.
doi:10.1016/j.jsbmb.2017.08.011

104. Rodriguez KF, Brown PR, Amato CM, Nicol B, Liu C-F, Xu X, et al. Somatic cell fate
maintenance in mouse fetal testes via autocrine/paracrine action of AMH and activin B.
Nat Commun. 2022;13: 4130. doi:10.1038/s41467-022-31486-y

105. Guiguen Y, Fostier A, Piferrer F, Chang C-F. Ovarian aromatase and estrogens: A pivotal
role for gonadal sex differentiation and sex change in fish. General and Comparative
Endocrinology. 2010;165: 352–366. doi:10.1016/j.ygcen.2009.03.002

106. Li M, Sun L, Wang D. Roles of estrogens in fish sexual plasticity and sex differentiation.
General and Comparative Endocrinology. 2019;277: 9–16.
doi:10.1016/j.ygcen.2018.11.015

107. Liu X, Dai S, Wu J, Wei X, Zhou X, Chen M, et al. Roles of anti-Müllerian hormone and its
duplicates in sex determination and germ cell proliferation of Nile tilapia. Genetics. 2021;
iyab237. doi:10.1093/genetics/iyab237

108. Halm S, Rocha A, Miura T, Prat F, Zanuy S. Anti-Müllerian hormone (*AMH*/AMH) in the
European sea bass: Its gene structure, regulatory elements, and the expression of
alternatively-spliced isoforms. Gene. 2007;388: 148–158. doi:10.1016/j.gene.2006.10.018

109. Liu X, Xiao H, Jie M, Dai S, Wu X, Li M, et al. Amh regulate female folliculogenesis and
fertility in a dose-dependent manner through Amhr2 in Nile tilapia. Molecular and Cellular
Endocrinology. 2020;499: 110593. doi:10.1016/j.mce.2019.110593

110. Wu K, Yue Y, Zhou L, Zhang Z, Shan H, He H, et al. Disrupting Amh and androgen
signaling reveals their distinct roles in zebrafish gonadal differentiation and
gametogenesis. Commun Biol. 2025;8: 1–15. doi:10.1038/s42003-025-07719-3

111. Zhang Z, Wu K, Ren Z, Ge W. Genetic evidence for Amh modulation of gonadotropin
actions to control gonadal homeostasis and gametogenesis in zebrafish and its
noncanonical signaling through Bmpr2a receptor. Development. 2020;147: dev189811.
doi:10.1242/dev.189811

112. Ijiri S, Kaneko H, Kobayashi T, Wang D-S, Sakai F, Paul-Prasanth B, et al. Sexual
Dimorphic Expression of Genes in Gonads During Early Differentiation of a Teleost Fish,
the Nile Tilapia Oreochromis niloticus1. Biology of Reproduction. 2008;78: 333–341.
doi:10.1095/biolreprod.107.064246

113. Czerwinski M, Natarajan A, Barske L, Looger LL, Capel B. A timecourse analysis of
systemic and gonadal effects of temperature on sexual development of the red-eared
slider turtle Trachemys scripta elegans. Developmental Biology. 2016;420: 166–177.
doi:10.1016/j.ydbio.2016.09.018

114. Sánchez-Baizán N, Jarne-Sanz I, Roco ÁS, Schartl M, Piferrer F. Extraordinary variability
in gene activation and repression programs during gonadal sex differentiation across
vertebrates. Front Cell Dev Biol. 2024;12. doi:10.3389/fcell.2024.1328365

115. Hacker A, Capel B, Goodfellow P, Lovell-Badge R. Expression of Sry, the mouse sex
determining gene. Development. 1995;121: 1603–1614. doi:10.1242/dev.121.6.1603

116. Kobayashi T, Nagahama Y. Molecular Aspects of Gonadal Differentiation in a Teleost Fish,
the Nile Tilapia. Sexual Development. 2009;3: 108–117. doi:10.1159/000223076

117. Eshel O, Shirak A, Weller JI, Hulata G, Ron M. Linkage and Physical Mapping of Sex
Region on LG23 of Nile Tilapia (Oreochromis niloticus). G3 Genes|Genomes|Genetics.
2012;2: 35–42. doi:10.1534/g3.111.001545

118. Liu X, Zhou L, Li W, Zhang L, Li M, Wang D. Production of fertile YY pseudofemales in
Nile tilapia with LG23 as the sex chromosome. Aquaculture. 2024;587: 740870.
doi:10.1016/j.aquaculture.2024.740870

119. Yamazaki W, Tan SL, Taketo T. Role of the X and Y Chromosomes in the Female Germ
Cell Line Development in the Mouse (Mus musculus). Sexual Development. 2023;16:
355–364. doi:10.1159/000521151

120. Lam TJ, Nagahama Y, Chan K, Hoar WS. Overripe eggs and postovulatory corpora lutea
in the threespine stickleback, Gasterosteus aculeatus L., form trachurus. Can J Zool.
1978;56: 2029–2036. doi:10.1139/z78-273

121. Roufidou C, Sebire M, Katsiadaki I, Mustafa A, Schmitz M, Mayer I, et al. Overripening of
eggs and changes in reproductive hormones in the threespine stickleback, Gasterosteus
aculeatus. Evol Ecol Res. 2016;17: 583–601.

122. Roufidou C, Borg B. Ovarian fluid in the three-spined stickleback : effects of egg
overripening and sex steroid treatment. Journal of Fish Biology. 2019;94: 446–457.
doi:10.1111/jfb.13915

123. Levy ER, Burgoyne PS. The fate of XO germ cells in the testes of XO/XY and
XO/XY/XYY mouse mosaics: evidence for a spermatogenesis gene on the mouse Y
chromosome. Cytogenet Cell Genet. 1986;42: 208–213. doi:10.1159/000132280

124. Mazeyrat S, Saut N, Grigoriev V, Mahadevaiah SK, Ojarikre OA, Rattigan Á, et al. A Y-
encoded subunit of the translation initiation factor Eif2 is essential for mouse
spermatogenesis. Nat Genet. 2001;29: 49–53. doi:10.1038/ng717

125. Touré A, Szot M, Mahadevaiah SK, Rattigan A, Ojarikre OA, Burgoyne PS. A new deletion
of the mouse Y chromosome long arm associated with the loss of Ssty expression,
abnormal sperm development and sterility. Genetics. 2004;166: 901–912.

126. Vernet N, Mahadevaiah SK, Yamauchi Y, Decarpentrie F, Mitchell MJ, Ward MA, et al.
Mouse Y-Linked Zfy1 and Zfy2 Are Expressed during the Male-Specific Interphase
between Meiosis I and Meiosis II and Promote the 2nd Meiotic Division. PLOS Genetics.
2014;10: e1004444. doi:10.1371/journal.pgen.1004444

127. Yamauchi Y, Riel JM, Stoytcheva Z, Ward MA. Two Y Genes Can Replace the Entire Y
Chromosome for Assisted Reproduction in the Mouse. Science. 2014;343: 69–72.
doi:10.1126/science.1242544

128. Shaw DE, Ross WD, Lambert AV, White MA. Single cell RNA-sequencing reveals no
evidence for meiotic sex chromosome inactivation in the threespine stickleback fish.
bioRxiv; 2024. p. 2024.11.26.625488. doi:10.1101/2024.11.26.625488

129. Largiadèr CR, Fries V, Bakker TCM. Genetic analysis of sneaking and egg-thievery in a
natural population of the three-spined stickleback (Gasterosteus aculeatus L.). Heredity.
2001;86: 459–468. doi:10.1046/j.1365-2540.2001.00850.x

130. Mehlis M, Rahn AK, Bakker TCM. Sperm quality but not relatedness predicts sperm
competition success in threespine sticklebacks (Gasterosteus aculeatus). BMC Evol Biol.
2015;15: 74. doi:10.1186/s12862-015-0353-x

131. Mehlis M, Hilke LK, Bakker TCM. Attractive males have faster sperm in three-spined
sticklebacks Gasterosteus aculeatus. Current Zoology. 2013;59: 761–768.
doi:10.1093/czoolo/59.6.761

132. Arnold AP, Chen X. What does the "four core genotypes" mouse model tell us about sex
differences in the brain and other tissues? Frontiers in Neuroendocrinology. 2009;30: 1–9.
doi:10.1016/j.yfrne.2008.11.001

133. Arnold AP. Four Core Genotypes and XY* mouse models: Update on impact on SABV
research. Neuroscience & Biobehavioral Reviews. 2020;119: 1–8.
doi:10.1016/j.neubiorev.2020.09.021

134. Vries GJD, Rissman EF, Simerly RB, Yang L-Y, Scordalakes EM, Auger CJ, et al. A Model
System for Study of Sex Chromosome Effects on Sexually Dimorphic Neural and
Behavioral Traits. J Neurosci. 2002;22: 9005–9014. doi:10.1523/JNEUROSCI.22-20-
09005.2002

135. Gatewood JD, Wills A, Shetty S, Xu J, Arnold AP, Burgoyne PS, et al. Sex Chromosome
Complement and Gonadal Sex Influence Aggressive and Parental Behaviors in Mice. J
Neurosci. 2006;26: 2335–2342. doi:10.1523/JNEUROSCI.3743-05.2006

136. Suster ML, Abe G, Schouw A, Kawakami K. Transposon-mediated BAC transgenesis in
zebrafish. Nat Protoc. 2011;6: 1998–2021. doi:10.1038/nprot.2011.416

137. Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. CHOPCHOP: a CRISPR/Cas9
and TALEN web tool for genome editing. Nucleic Acids Research. 2014;42: W401–W407.
doi:10.1093/nar/gku410

138.   Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. CHOPCHOP v2: a web tool for
1244   the next generation of CRISPR genome engineering. Nucleic Acids Research. 2016;44:
1245   W272–W276. doi:10.1093/nar/gkw398

1246   139.   Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E. CHOPCHOP
1247   v3: expanding the CRISPR web toolbox beyond genome editing. Nucleic Acids Research.
1248   2019;47: W171–W174. doi:10.1093/nar/gkz365

1249   140.   Subedi A, Macurak M, Gee ST, Monge E, Goll MG, Potter CJ, et al. Adoption of the Q
1250   transcriptional regulatory system for zebrafish transgenesis. Methods. 2014;66: 433–440.
1251   doi:10.1016/j.ymeth.2013.06.012

1252   141.   Kingsley DM, Zhu B, Osoegawa K, De Jong PJ, Schein J, Marra M, et al. New Genomic
1253   Tools for Molecular Studies of Evolutionary Change in Threespine Sticklebacks.
1254   Behaviour. 2004;141: 1331–1344.

1255   142.   Archambeault SL, Bärtschi LR, Merminod AD, Peichel CL. Adaptation via pleiotropy and
1256   linkage: Association mapping reveals a complex genetic architecture within the
1257   stickleback Eda locus. Evolution Letters. 2020;4: 282–301. doi:10.1002/evl3.175

1258   143.   Collins III JJ, Hou X, Romanova EV, Lambrus BG, Miller CM, Saberi A, et al. Genome-
1259   Wide Analyses Reveal a Role for Peptide Hormones in Planarian Germline Development.
1260   PLOS Biology. 2010;8: e1000509. doi:10.1371/journal.pbio.1000509

1261   144.   Square TA, Sundaram S, Mackey EJ, Miller CT. Distinct tooth regeneration systems
1262   deploy a conserved battery of genes. EvoDevo. 2021;12: 4. doi:10.1186/s13227-021-
1263   00172-3

1264   145.   Edgar RC. Muscle5: High-accuracy alignment ensembles enable unbiased assessments
1265   of sequence homology and phylogeny. Nat Commun. 2022;13: 6968.
1266   doi:10.1038/s41467-022-34630-w

1267   146.   Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary
1268   analyses in R. Bioinformatics. 2019;35: 526–528. doi:10.1093/bioinformatics/bty633

1269   147.   Mayor C, Brudno M, Schwartz JR, Poliakov A, Rubin EM, Frazer KA, et al. VISTA :
1270   visualizing global DNA sequence alignments of arbitrary length. Bioinformatics. 2000;16:
1271   1046–1047. doi:10.1093/bioinformatics/16.11.1046

1272   148.   Frazer KA, Pachter L, Poliakov A, Rubin EM, Dubchak I. VISTA: computational tools for
1273   comparative genomics. Nucleic Acids Research. 2004;32: W273–W279.
1274   doi:10.1093/nar/gkh458

1275   149.   Brudno M, Do CB, Cooper GM, Kim MF, Davydov E, Program NCS, et al. LAGAN and
1276   Multi-LAGAN: Efficient Tools for Large-Scale Multiple Alignment of Genomic DNA.
1277   Genome Res. 2003;13: 721–731. doi:10.1101/gr.926603

1278   150.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
1279   data. Bioinformatics. 2014;30: 2114–2120. doi:10.1093/bioinformatics/btu170

151. Zhang Y, Park C, Bennett C, Thornton M, Kim D. Rapid and accurate alignment of
nucleotide conversion sequencing reads with HISAT-3N. Genome Res. 2021;31: 1290–
1295. doi:10.1101/gr.275193.120

152. Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F. Analysing high-throughput sequencing
data in Python with HTSeq 2.0. Bioinformatics. 2022;38: 2943–2945.
doi:10.1093/bioinformatics/btac166

153. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology. 2014;15: 550. doi:10.1186/s13059-014-
0550-8

1289

1290 **Supplemental figures and tables**
1291

1292
1293 Fig S1. Sequence conservation of the *amhy* transgene region with autosomal *amha*.
1294 Vistaplot showing conserved sequence identity of *amhy* to *amha* with sequence identity on the Y
1295 axis and base pair position of *amhy* on the x axis. Exon positions are indicated in dark blue, and
1296 the amh domain and TGF-β domain are indicated in light blue above the graph. Conserved
1297 sequences share 70% sequence identity in a 100 bp window.
1298

A)

B)

C)

1299
Fig S2. F0 *amhy* crispants show genetic mosaicism and variable sex reversal
A) XY crispants developed as females with ovaries or males with testes. Sanger sequencing of
exon 1 (B) and exon 3 (C) show various mutations relative to wildtype. Discordant
chromatograms in crispants are indicative of mosaic mutations. Mutations are bolded in orange.
CRISPR target site and PAM sequence are indicated by orange lines above wildtype reference
sequence.
1306

-2282,+8
GCTCAGCGCC**AGGTCGGG**TTCATTCAAACCCACAGGCCAGT

-4
CGGGGCTCTGTGTGGCCCTATTGACCTTGGACAGCAGCTGATCCC

-4,+5
CGGGGCTCTGTGTGGCCCTATTGACCTTG**ACCTT**GACAGCAGCTGATCCC

+9,+5
CCAGC**AGTGCTCAT**AGTGCTCATACCAAAC**ATACC**GAAGACGAGAGAGGACGC

1307
1308   Fig S3. Sanger sequencing of four F1 *amhy* mutant alleles.
1309   Inserted base pairs are bolded. Locations of deletions are indicated with red arrowheads.
1310

1311
1312 Fig S4. Expression of gonadal genes in wildtype and sex reversed ovaries
1313 *In situ* hybridization of gonadal genes in three biological replicates for wildtype XX ovaries and
1314 XY *amhy*-KO ovaries. Scale bars = 200 μm.
1315

1316

1317
1318 Fig S5. CRISPR/Cas9 sgRNAs for *amhy* have multiple mismatches to their homologous site in
1319 *amha*.
1320 PAM and sgRNA sequences are shown above the reference sequence. Mismatched nucleotides
1321 are bolded in orange text. Sanger sequencing of sex reversed XY F1 *amhy*-KO fish showed no
1322 off-target mutations in *amha* in exon 1 (A) or exon 3 (B).
1323

1324
Fig S6. Plasmid maps of transgenes constructed in this study.
Transgenes used the *X. laevis* Ef1a promoter (A and D), *D. rerio* Ubi promoter (B and E), or
endogenous *amhy* promoter (C and F). Control transgenes express only EGFP (A and B). Other
transgenes contain the *amhy* coding sequence (D and E) or the full *amhy* sequence including
introns (C and F). To screen for transgene integration, *amhy* transgenes include an EGFP joined
to *amhy* with a P2A self-cleaving peptide (D and E) or separate EGFP cassette (F). All
transgenes are flanked by Tol2 transposon arms to facilitate transgene integration.
1332

1333
1334 Fig S7. Expression of gonadal genes in wildtype and sex reversed testes.
1335 *In situ* hybridization of gonadal genes in three biological replicates for wildtype XY testes and
1336 XX *Tg(amhy,EGFP)* testes. For one wildtype replicate, the tissue was lost during staining for
1337 *cyp19a1a*. Scale bars = 200 μm.
1338

1339
Fig S8. PCA plots of normalized gene expression counts.
Principal components 1 and 2 with the Proportion of Variance Explained on the X and Y axes,
respectively. When analyzing all genes or autosomal genes only, samples group by
developmental stage and not sex. X chromosome genes show separation by both stage and
sex. Y chromosome genes group primarily by sex with male samples separate by
developmental stage.
1346

1347

1348
1349 Fig S9. Volcano plots of female to male differential expression by stage.
1350 Positive log2 Fold Change values indicate higher expression in females, and negative values
1351 indicate higher expression in males. Significant genes (adjusted p < 0.05) are shaded. The
1352 number of male biased, female biased, and total differentially expressed genes are shown
1353 above each plot.
1354

1355 Table S1. Viability and Hatching rates of crosses from F1 XY *amhy*-KO females with wildtype XY
1356 males

Fish	Cross number	Clutch Size	Viable	Hatched	Viable:total	Hatched:total	Hatched:viable
A	1	43	1	1	0.0233	0.0233	1.0000
B	1	51	2	0	0.0392	0.0000	0.0000
C	1	63	4	2	0.0635	0.0317	0.5000
C	2	99	24	21	0.2424	0.2121	0.8750
C	3	81	11	6	0.1358	0.0741	0.5455
C	4	100	78	54	0.7800	0.5400	0.6923
C	5	89	70	46	0.7865	0.5169	0.6571

1357
1358
1359 Table S2. Viability and Hatching rates of crosses from F2 XX or XY *amhy*-KO females a wildtype
1360 XY male

Genotype	Clutch Size	Viable	Hatched	Viable:total	Hatched:total	Hatched:viable
XX	57	57	57	1.0000	1.0000	1.0000
XX	112	105	34	0.9375	0.3036	0.3238
XX	56	55	55	0.9821	0.9821	1.0000
XX	136	90	47	0.6618	0.3456	0.5222
XX	94	94	79	1.0000	0.8404	0.8404
XY	84	6	2	0.0714	0.0238	0.3333
XY	65	0	0	0.0000	0.0000	NA
XY	98	0	0	0.0000	0.0000	NA
XY	22	0	0	0.0000	0.0000	NA
XY	72	6	4	0.0833	0.0556	0.6667
XY	92	0	0	0.0000	0.0000	NA
Total XX	455	401	272	0.8813	0.5978	0.6783
Total XY	433	12	6	0.0277	0.0139	0.5000

1361
1362

1363 Table S3. Combined *amha* and *amhy* expression show no significant difference between males and females

stage	stage 17		stage 20		stage 23		0 dah		4 dah		8 dah	
sex	female	male	female	male	female	male	female	male	female	male	female	male
amha normalized counts	24.2	31.0	32.5	62.5	142.8	98.1	250.9	248.8	314.6	408.4	348.0	350.2
	35.3	13.2	48.5	37.4	125.2	57.8	288.3	286.2	395.5	286.7	407.7	493.6
	27.5	28.4	54.0	33.1	98.0	109.8	283.8	331.6	318.2	364.8	525.1	431.9
Mean	29.0	24.2	45.0	44.3	122.0	88.6	274.3	288.9	342.8	353.3	426.9	425.2
SD	5.7	9.6	11.2	15.9	22.6	27.3	20.4	41.5	45.7	61.7	90.1	71.9
amhy normalized counts	0	9.4	1.0	3.8	0	14.5	0.9	22.5	0	40.8	0	13.5
	0	10.2	0	4.7	0	13.8	0	20.9	0	20.5	0	32.9
	0	5.7	0	5.2	0	13.0	2.9	8.7	0	16.3	0	9.4
Mean	0	8.4	0.3	4.5	0	13.8	1.3	17.4	0	25.9	0	18.6
sd	0	2.4	0.6	0.7	0	0.8	1.5	7.5	0	13.1	0	12.6
length adjusted *amhy* counts	0	9.38	1.01	3.79	0	14.50	0.85	22.50	0	40.84	0	13.54
	0	10.18	0	4.68	0	13.79	0	20.94	0	20.48	0	32.90
	0	5.68	0	5.17	0	12.96	2.93	8.73	0	16.27	0	9.39
Mean	0	8.42	0.34	4.55	0	13.75	1.26	17.39	0	25.86	0	18.61
SD	0	2.40	0.59	0.70	0	0.77	1.50	7.54	0	13.14	0	12.55
amha + adj. *amhy* normalized counts	24.21	45.59	34.06	68.43	142.76	120.73	252.26	283.91	314.60	472.16	348.00	371.36
	35.34	29.13	48.55	44.72	125.18	79.29	288.27	318.92	395.50	318.66	407.67	544.90
	27.52	37.28	54.01	41.15	97.97	130.05	288.35	345.21	318.15	390.16	525.11	446.56
Mean	29.02	37.33	45.54	51.44	121.97	110.02	276.29	316.01	342.75	393.66	426.93	454.27
SD	5.71	8.23	10.31	14.82	22.57	27.02	20.81	30.75	45.72	76.81	90.11	87.03
P value	0.4078		0.6054		0.5892		0.1474		0.3913		0.7246	

1364 Length adjusted *amhy* counts were calculated by multiplying *amhy* normalized read counts by transcript length of *amha* (3895 bp)
1365 divided by *amhy* (2496 bp). P values show two tailed t test for difference in total *amh* expression between males and females at each
1366 stage (P <0.05)
1367

1368   Table S4 – PCR primers used in this study

Assay	Forward primer	Reverse primer
XY *idh* genotyping	5'-GGGACGAGCAAGATTTATT-3'	5'-TTATCGTTAGCCAGGAGATGG-3'
Amhy exon 1 genotyping, amhy transgene detection	5'-TTGGTGGGTTGATGCTCTGC-3'	5'-CTATGTGAGAGACTGTGGCGG-3'
amhy exon 3 genotyping	5'-TCAGTTTGGAATCTGCACTGTT-3'	5'-GGATGCAGTGATTTACTGGTGA-3'
amhy 2.3 kb deletion genotyping	5'-CACAGTCCTTCGACTCCTTCG-3'	5'-GAGACCATGGTGACTGGCTC-3'
amha exon 1 genotyping	5'-GGCAGTGAGAAGGGAAGGTG-3'	5'-GGCCTCTGGTGATGGGAATG-3'
amha exon 3 genotyping	5'-CAGCCGGAGTCAGTTTGGAA-3'	5'-TCATGCATCGTCGACTGGAG-3'
EGFP transgene detection	5'-ATCATGGCCGACAAGCAGAA-3'	5'-AACTCCAGCAGGACCATGTG-3'
Dmrt1 riboprobe cloning	5'-AGCAGCACCGAGAAGCAG-3'	5'-CTCGTTGACGTTGCCCATGA-3'
Gsdf riboprobe cloning	5'-TTTTGTCACGGCGATGCTTC-3'	5'-AGGAAATGACGAGGGTGCTG-3'
Foxl2 riboprobe cloning	5'-GAGAAGGTCCCGGAGAAAGC-3'	5'-GCAGTGCATCATCGAGAGCT-3'
Cyp19a1a riboprobe cloning	5'-CCTTGTTTCTGTCTGGGCCT-3'	5'-TTGAGGAGCAGCAGCATGA-3'
Riboprobe template synthesis	5'-TAATACGACTCACTATAGGG-3'	5'-TAATACGACTCACTATAGGG-3'

1369
1370