1 **Low anabolic independence emerges when cultivating more than three bacterial species**
2 **together.**
3

4 Armando Pacheco-Valenciana[1*], Anna Tausch[2], Iva Veseli[3,4], Jennah E. Dharamshi[1], Fabian Bergland[1], Luis
5 Fernando Delgado-Zambrano[4], Alejandro Rodríguez-Gijón[1], Anders F. Andersson[4], Sarahi L. Garcia[1,2,3*].
6

[1] Department of Ecology, Environment, and Plant Sciences, Science for Life Laboratory, Stockholm University,
Stockholm, Sweden.
[2] Institute for Chemistry and Biology of the Marine Environment (ICBM), School of Mathematics and Science,
Carl von Ossietzky Universität Oldenburg, 26129, Oldenburg, Germany
[3] Helmholtz Institute for Functional Marine Biodiversity at the University of Oldenburg (HIFMB), Oldenburg,
Germany
[4] Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, 27570 Bremerhaven, Germany
[5] Department of Gene Technology, Science for Life Laboratory, KTH Royal Institute of Technology, Stockholm,
Sweden.
16
17 * Correspondence: armando.pacheco@su.se, sarahi.garcia@su.se

18 KEYWORDS: microbial model communities, metagenomics, low anabolic independence, microbial
19 interdependencies.
20

21 **ABSTRACT**
22

Microorganisms thrive in complex communities shaped by intricate interactions, yet the
extent and ecological implications of biosynthetic dependencies in natural communities
remain underexplored. Here, we used a dilution approach to cultivate 204 microbial model
communities from the Baltic Sea and recovered 527 metagenome-assembled genomes
(MAGs) that dereplicated into 72 species-clusters (>95% average nucleotide identity - ANI).
Of these species, at least 70% represent previously uncultivated lineages. Combined with
1073 MAGs from Baltic Sea metagenomes, we generated a genomic catalog of 701 species-
clusters. Our results show that cultures with more than 3 species included microorganisms
with smaller genome sizes, lower biosynthetic potential for amino acids and B vitamins, and
higher prevalence and abundance in the environment. Moreover, biosynthetic genes were
distributed across genomes within the same model community, indicating metabolic
interdependencies. Our results demonstrate that cultivating bacteria in dilution model
communities facilitates access to previously uncultivated but abundant species that likely
depend on metabolic partners for survival. Together, our findings highlight the value of
community-based cultivation for unraveling ecological strategies. Finally, we confirm that

metabolic interdependencies and genome streamlining are widespread features of successful

environmental microorganisms.

INTRODUCTION

Microbial communities in diverse environments operate as complex systems driven by multi-

species interactions[1]. Understanding such complex interactions is essential because

microorganisms play key roles in the biogeochemical cycles on Earth[2]. To unravel microbial

interactions, we need to investigate microbial communities at various levels of biological

organization[3,4], ranging from one-to-one species interactions to simplified multi-species

systems (e.g., model communities[5], synthetic communities[6], or microcosms[7]), and ultimately

to naturally occurring communities.

At the community level, metagenomics has become a powerful tool for uncovering the

genetic potential of microbial communities via shotgun sequencing[8-10]. Analyzing

metagenomic data reveals not only the vast diversity of microbial species[11] but also their

metabolic potential and co-occurrence networks, which are important for ecosystem

functioning[12]. To bridge the gap between broad metagenomic insights and detailed ecological

understanding, a few studies have explored genome-specific traits and potential interactions

by inferring auxotrophies. While auxotrophs have historically been experimentally identified

via cultures that require the addition of specific nutrients to grow[13], recent work based on

genomes and metagenomes has determined auxotrophies based on pathway completeness and

found smaller genomes to be more auxotrophic[6,14-19]. In all these studies, auxotrophy has

been treated as a binary trait, however, microbial biosynthetic capabilities in nature likely

span a spectrum. For example, many microorganisms can complete the biosynthesis of an

essential metabolite starting from a precursor or intermediate without needing the essential

metabolite itself[20,21]. Nevertheless, modeling work has shown that microbial communities

enriched in the so-called auxotrophs can exhibit greater robustness under ecological

disturbances, suggesting that these metabolic interdependencies may contribute to overall

community stability[22]. While metagenomics offers a broad understanding of microbial

communities, interactions cannot easily be inferred from co-occurrences within natural

complex ecosystems.

Experimental systems are needed to observe microbial interaction dynamics under controlled

conditions. Studies have increasingly turned to simplified systems[23-25], such as co-cultures[26-

[28] and mixed cultures[18,29,30], in order to identify specific types of interactions. To further

contextualize these findings, our literature review in **Table S1** provides a comprehensive

overview of publications where microbial interaction patterns were observed in different

experimental settings. Across these studies, cross-feeding mechanisms and mutualistic

interactions are mostly studied in cultures with two different populations or species[27,31,32]. To

a lesser extent, more complex metabolic interactions have also been studied by mixing

different isolated species or co-cultivating them in model ecosystems, with the goal of

increasing complexity to more closely resemble natural environments[29,33,34]. However, many

of these methods focus on cultured isolates, and the vast majority of microorganisms remain

uncultivated[35]. An alternative, yet underutilized method of establishing model ecosystems

composed of previously uncultivated microorganisms is through dilution cultivation. Dilution

to mixed-cultures of naturally co-occurring microorganisms has the potential to cultivate

previously uncultivated microorganisms as well as to allow observation of natural microbial

interactions[36]. Such cultures, also known as microbial model communities, represent a small

subset of the many interactions likely occurring in the natural system[33]. By studying a larger

number of microbial model communities, we can gain a more comprehensive understanding

of microbial interactions occurring in a natural environment.

In our study, we have focused on studying potential interdependencies at two levels of

biological organization by using high-throughput dilution cultivation of model communities

together with genome-resolved metagenomics to unravel ecological strategies of

microorganisms in the Baltic Sea. Moreover, we examine the biosynthesis of essential

metabolites or anabolic independence as a continuous spectrum rather than through

conventional binary classifications. For this, we use pathway completeness metrics rather

than assigning genomes as strictly prototrophic or auxotrophic. Finally, we identified

correlations between genome size, potential biosynthesis of essential metabolites, relative

abundance, and prevalence using genomes obtained from both microbial model communities

and metagenomic data from Baltic Sea pelagic samples. Our findings demonstrate that

microbial model communities are an effective cultivation technique to cultivate previously

uncultivated taxa and for identifying putative microbial interactions, including metabolic

interdependencies between biosynthetically dependent members.

RESULTS

A Baltic Sea MAG catalog

To generate the microbial model communities, we used the dilution-to-extinction cultivation technique in two formats. The first type, low inoculum size, involved inoculating between approximately 2 and 100 cells per well in 1 ml 96-well plates used for each of the inoculum sizes. The second type, high inoculum size, ranged from approximately 200 to 1×10^6 cells inoculated per microbial model community in 100 ml volumes. After a 4-week incubation period, we sent an aliquot of all 801 cultures for lysis and DNA amplification using multiple displacement amplification (MDA). Based on amplification success, 315 cultures passed the negative control threshold and were sent for sequencing. In total, only 204 microbial model communities together yielded 527 MAGs. Moreover, from the original sample used to establish the microbial model communities, we generated two metagenomes (each from a distinct DNA extraction method) that yielded 305 MAGs. To create a comprehensive genomic catalog, we also added 771 MAGs from 110 publicly available Baltic Sea metagenomes (**Figure 1A**, **Table S2**)[37–39].

Examining this comprehensive MAG catalog allowed us to assess microbial genomic characteristics of all microorganisms found both in the environmental samples and our 204 microbial model communities (**Table S3**). Altogether, the 1603 MAGs were dereplicated into 701 species-clusters (ANI >95%), which form the Baltic Sea genomic catalog (BalticMAG catalog)[40] used in this study (**Table S4**). The average completeness of the 701 species-cluster representative MAGs is 88%, and they were all used to analyze taxonomy, abundance, prevalence, and estimated genome size (**Figure S1**). The varying completeness of the MAGs has a very minor effect on estimated genome size and relative abundance, just as observed in other studies[17,41]. However, only 450 species-cluster representative MAGs are of high-quality (completeness >90% and contamination <5%) and were used to investigate anabolic potential (**Figure S1D**).

Examining the source of genomes in the BalticMAG catalog, we found that 33 of the species-clusters included MAGs exclusively from the microbial model communities, 629 included MAGs exclusively from the environmental metagenomes, and 39 (54% of all cultured species) included MAGs from both sources (**Figure 1C, Table S4**, **Figure S2).**

To investigate the relative abundance of the BalticMAG catalog, we mapped all the environmental metagenomics reads against the genome catalog. We observed that salinity

significantly co-varied with the proportion of metagenomic reads that mapped to all species

representative genomes (**Figure 1B**). While on average, 38.63% of the metagenomic reads

per sample mapped to the BalticMAG catalog, the highest mapping percentage (63.99% and

62.89‰) was observed at salinity concentrations 11.28‰ and 7.65‰, respectively. Notably,

the two metagenomes from this study (sample with salinity concentration 7.12‰) displayed

some of the highest mapping rates at 51.77% and 60.16%, reflecting that the BalticMAG

catalog is most complete for salinities between 6 and 11‰.

Despite salinity differences (**Figure 1D**), the average relative abundance of the 39 species-

clusters that included MAGs from both microbial model communities and environmental

metagenomes was significantly higher in the whole dataset (**Figure 1E),** but also when

comparing only the metagenomes from the location and salinity from which we sampled

(**Figure S3**). This group of species-clusters (from both sources) shows that our cultivation

method can capture some of the most abundant taxa from the environment. Altogether, the

diverse taxa cultivated in model communities accounted for ~20% of the total relative

abundance in the original environmental sample (**Figure S4**). Moreover, the 33 species-

clusters with MAGs sourced exclusively from microbial model communities were detected

across environmental metagenomes, albeit at significantly lower abundances. This indicates

that our cultivation approach also enables the recovery of species that are missed by assembly

and binning in metagenomic surveys.

Higher inoculum size increases community richness and uncovers a genome size

plateau.

In the low inoculum size microbial model communities, the more cells we inoculated, the

higher the number of cultures that yielded MAGs (**Figure 2A**). In total, 94 low inoculum size

model communities yielded only one MAG each, while 110 model communities resulted in

two or more MAGs. In model communities with more than one MAG, each MAG belonged

to a different species-cluster in our analysis. Therefore, for clarity, we refer to different

MAGs within a model community as different species. Starting from an inoculum size of

approximately 30 cells, microbial model communities with more than two species appear

more often (**Figure 2B** and **C**). Nearly 82% (n = 433) of the microbial model community

MAGs were obtained from multi-species cultures, and the highest number of co-occurring

species were found in the high inoculum size microbial model communities inoculated with

5000 cells (**Figure 2C**). Despite using a complex inoculum from the Baltic Sea, observing a

maximum of 13 co-occurring species suggests that our cultivation conditions may impose a

threshold on the complexity of model communities. Alternatively, since sequencing followed

MDA, there is also the possibility that some model communities included more species that

were not amplified or assembled and binned. Nevertheless, the increased growth success with

increasing inoculum size likely reflects a greater probability of including cells that can grow

in isolation or in the presence of a specific required community partner.

We found that as the number of retrieved MAGs per microbial model community increased,

more microorganisms with smaller genomes emerged, as indicated by a slight but significant

correlation between estimated genome size and the number of MAGs per microbial model

community (**Figure 2D**). In general, the estimated genome sizes ranged from 0.88 to 6.27

Mbp for MAGS from cultures and, from 0.61 to 12.32 Mbp for MAGs from the environment

(**Figure 2D** and **F**). While there was no significant difference between the average estimated

genome sizes of species-clusters from microbial model communities and Baltic Sea

metagenomes (around 3.00 Mbp), we found that species-clusters exclusively composed of

genomes from microbial model communities had significantly larger genome sizes than

species-clusters including genomes from both sources, and no statistically different genome

completeness or contamination (**Figure S5**). Further analysis reinforced the observation that

genomes from high inoculum size microbial model communities and from model

communities with more than 3 species had on average, smaller genome sizes (2.89 and 2.92

Mbp, respectively - **Figure S6** and **S7**). Genomes from high inoculum size microbial model

communities were also on average more prevalent and had higher average relative abundance

in the investigated Baltic Sea metagenomes. Finally, the average genome size per microbial

model community stabilized at around 3 Mbp in cultures containing more than three species

(**Figure 2E**).

Model communities with more than three species reveal distinct microbial diversity.

The 72 species from microbial model communities spanned five of the most abundant phyla

(*Pseudomonadota*,   *Bacteroidota*,   *Campylobacterota*,   *Cyanobacteriota*,   and

Verrucomicrobiota) present in the Baltic Sea environmental metagenomic sample[42,43]. These

species varied substantially in their observed growth strategies. Four of them were only found

in single-species cultures, forty-eight exclusively in multi-species cultures, and twenty

appeared growing both alone and in groups (**Figure 3**). Of the species growing consistently in groups, nine grew across different levels of community complexity (e.g., 2, 3, or more than three species per culture), and thirty-nine species were restricted to a single type of community complexity. Specifically, thirty-one species were recovered from microbial model communities with more than three species. Notably, 70% of these cultured species lacked a species-level assignment in the GTDB taxonomy, suggesting they represent previously uncultivated lineages with no characterized MAGs either.

To evaluate whether culturing bacteria in groups increases the cultivability and recovery of microbial diversity, we computed rarefaction curves for species accumulation across increasing numbers of cultures (**Figure 4**). After 50 cultures of the single-species type, we had in average recovered seventeen species, and every 12 new cultures would yield only three more species in average. As we cultivated more species together, the initial steep slope also increased. In fact, in 25 cultures with more than three species, forty-nine species were recovered. This demonstrates that culturing in groups is a powerful strategy for the recovering of microbial taxa that would normally not grow in axenic cultures.

Species strictly found growing in groups are more abundant and prevalent and have lower biosynthetic capacity.

The total number of genomes recovered per species-cluster varied considerably (**Figure 5A**), ranging from those detected only once (e.g., sc_576 and sc_639; taxa of species-clusters can be found in **Table S4** and **Figure 3**) to those found in over 40 model communities (e.g., sc_011 and sc_014). Fifteen species (21%) were present in more than 10 model communities, collectively accounting for approximately 63% of the total MAGs (330 out of 527). We found that most of the frequently retrieved MAGs showed the flexibility of growing independently or in groups (**Figure 5A** and **B**). However, our experiment did not systematically test if species only found on their own could also grow in groups, or if species found only in groups could also grow alone. Nevertheless, when we examined the relative abundance of all cultivated species across their source environmental sample, we observed a positive correlation to the total number of genomes recovered for species found in groups (**Figure 5C**). On the other hand, the two most abundant species (Pelagibacterales sc_139 and Methylacidiphilales sc_121) were found exclusively in groups but were cultivated very few times. This might reflect that while many abundant microorganisms are easier to cultivate

242 under our model community conditions, finding the right partners or getting the right
243 conditions might be more challenging for others.
244
245 When examining species prevalence across all environmental samples, a similar significant
246 positive correlation emerged with the number of cultured genomes per species (**Figure 5D**),
247 indicating that more prevalent taxa are more frequently retrieved through our cultivation
248 conditions.
249
Next, we compared estimated genome sizes and biosynthetic potential for species capable of
growing alone (singleton-capable) and those that strictly grow in groups. While there was no
significant difference in estimated genome sizes between these two categories (**Figure 5E**),
we observed a significant difference in average pathway completeness of custom biosynthesis
modules for both amino acids and vitamins, with species found strictly in groups exhibiting a
lower average pathway completeness (**Figure 5F**). For this analysis, we excluded three amino
acids (alanine, asparagine, and aspartate) since their biosynthesis is commonly mediated by
several generic transaminases[44] (see Methods for details). These findings suggest that the
ability to grow alone is most likely associated with greater anabolic independence, whereas
species that require growing in groups may depend biosynthetically on other members of the
community.
261
262 **Lowest anabolic independence and higher interdependencies in species growing in**
263 **groups**
264
We next examined the biosynthetic potential of the 305 high-quality genomes (completeness
>90%, contamination <5%) recovered from our microbial model communities, with a focus
on how this potential relates to community complexity. We found that genomes from single-
species and two-species cultures showed consistently higher anabolic independence seen in
amino acid biosynthesis and vitamin modules (**Figure 6** and **Figure S8**). In contrast,
genomes from three-species cultures showed lower biosynthesis potential, and cultures with
more than three species have the lowest anabolic independence. This reduction was
particularly pronounced in amino acid biosynthesis pathways when compared to their highest
average completeness value observed in cultures with one or two species. The nine
aminoacids with lowest pathway completeness were arginine (~25% lower), proline (~22%),
phenylalanine (~21%), tyrosine (~20%), threonine (~18%), leucine (~17%), tryptophan

(~16%), serine (~14%), and isoleucine (~12%) (**Figure S9**). Additionally, although vitamin

B12 showed a relative decrease, both single-species and more than three-species model

communities had low average pathway completeness (~22% down to ~12%), suggesting

generally limited biosynthetic capacity for B12, regardless of community complexity.

While it is assumed that metagenome assembly might work better with lower diversity

inputs, genomes from our more complex cultures (> 3 species) showed only slight differences

in completeness, contamination, and N50 values (**Figure S10**). To test whether this small

difference in genome quality could explain the reduced biosynthetic potential in complex

communities, we examined the relationship between genome completeness and the

completeness of each biosynthetic module individually (**Figure S11**). Across all 27 modules,

correlations with all genomes from model communities were weak (Spearman R typically

<0.3). Modules with similar completeness were often observed across genomes with varying

genome completeness, indicating that minor differences in MAG quality do not drive the

observed metabolic trends. We also assessed correlations for each biosynthetic module

separately across the four community complexity categories (1, 2, 3, and >3 species per

culture; **Figure S12**) and found only a few modules exhibited moderate correlations (R ≈

0.3–0.5), with no systematic bias toward the more complex communities (>3 species). These

results suggest that the lower biosynthetic capacity in multi-species cultures is a biological

signal driven by community composition rather than a technical artifact of genome quality.

To evaluate if microbial model communities with more than three species collectively encode

complete biosynthetic pathways, we evaluated gene content (based on individual module

steps) of all 262 genomes recovered from 37 microbial model communities regardless of

completeness. Even partial genomes were included because they can provide valuable

evidence for individual metabolic steps. When examining the data at the community-level,

we observed that in these cultures, all species collectively encoded the biosynthetic pathways

through a mosaic of partial contributions from different species (**Figure 7**). This community-

level stepwise completion indicates that biosynthetic capacity emerges collectively rather

than within individual genomes, suggesting that anabolic interdependencies support the idea

of facilitated community growth.

Ubiquitous species tend to have smaller estimated genome sizes and reduced

biosynthetic capacity.

For the species from cultures (57 high-quality), we found a significant negative correlation

between relative abundance and average pathway completeness for both amino acids and

vitamins in species that exclusively grew in groups and those that exclusively grew alone

(**Figure 8A**). This trend persisted for vitamin biosynthesis (**Figure S13C**), while amino acids

alone showed a weaker association (**Figure S13B**). Moreover, we observed a significant

positive correlation between estimated genome size and average pathway completeness for

species that exclusively grew in groups and alone, showing that smaller genomes encode

fewer biosynthetic pathways (**Figure 8B**). Notably, the correlation between vitamin

biosynthesis and estimated genome size was particularly strong in group-only species

(**Figure S13F**).

Scaling the observations from model communities to all species from the BalticMAG catalog

(n = 450 high-quality), we found similar trends, particularly among species detected only in

environmental metagenomes. In this group, we found that higher average relative abundances

(**Figure 8C**) and smaller estimated genome sizes (**Figure 8D**) were linked to lower average

completeness of biosynthetic pathways for amino acids and vitamins.

Finally, when we analyzed all 701 species-clusters, we observed a strong negative correlation

between estimated genome size and both relative abundance (**Figure 8E**) and prevalence

across samples (**Figure 8F**). These results align with prior observations, where the most

successful and widespread taxa have a streamlined genome with low biosynthetic potential[17].

Collectively, these findings suggest that biosynthetic dependencies may be a common

ecological strategy in the Baltic Sea.

Anabolic dependencies in the Baltic Sea: different paths to microbial success

Given that anabolic dependencies seem to be common in the Baltic Sea, an important

question arises: Are these dependencies uniform across different microorganisms, or do

different microorganisms adopt distinct metabolic strategies to achieve ecological success?

To explore variation in biosynthetic potential across the BalticMAG catalog, we analyzed the

450 high-quality species genomes. A principal component analysis (PCA) on the biosynthetic

completeness matrix revealed a clear separation of genomes into three distinct biosynthetic

completeness groups: Low (0–30%), Medium (30–62.5%), and High (>62.5%) (**Figure**

9A&D). Estimated genome size followed the trend, increasing from low to high across the

biosynthetic completeness groups (**Figure 9E**). In general, among the most frequently

incomplete or partially incomplete amino acid pathways were histidine, phenylalanine, and

tyrosine as well as vitamin B1 and B12 (**Figure S14**). The microorganisms in the different

biosynthetic groups showed distinct and clear taxonomic signatures (**Figure 9C**). The low

biosynthesis completeness group consisted exclusively of *Patescibacteria* and a single

Firmicutes genome, both lineages associated with symbiotic or highly host-dependent

lifestyles. The medium biosynthesis completeness group was dominated by *Bacteroidota* and

Planctomycetota, while the high biosynthesis group encompassed a broader diversity of

phyla, including *Proteobacteria* and *Actinobacteria* as major contributors. Interestingly,

microorganisms with low and high biosynthetic completeness showed no significantly

different relative abundance, possibly because of the small number of taxa in the former

group (**Figure 9F**).

To better visualize biosynthetic strategies, we projected the data using Uniform Manifold

Approximation and Projection (**Figure 9B**), which further confirmed the biosynthetic

groupings, with strong clustering patterns aligned with biosynthetic capacity. Interestingly,

the high biosynthetic group is further separated into two subclusters. One of them, composed

of 51 genomes, was taxonomically diverse yet tightly grouped. We refer to this subcluster as

Cluster A, and the remaining genomes from the High biosynthetic group as Cluster B. Cluster

A genomes had significantly larger genome sizes than those in Cluster B (**Figure 9H**) and

appeared to be the main producers of vitamin B12 (**Figure 9G** and **Figure S15**). Cluster B

genomes showed higher completeness for vitamin B3.

Finally, from the 57 high-quality genomes from cultures, we observed that we cultivated no

microorganisms from the low biosynthetic group. This might suggest that cultivation

techniques to grow them together with their possible host might be needed. While, most of

the microorganisms we cultivated belonged to the high biosynthetic group (**Figure 9J**), the

six microorganisms we cultivated from the medium biosynthetic group grew exclusively in

groups. Further, a PCA based on only the genomes from the microorganisms we cultivated

revealed a new set of clusters with biosynthesis values ranging from 50% to 90% (**Figure 9K**

and **9L**). Cluster 1, which had the lowest biosynthesis potential, was again mostly composed

of species that were capable of growing exclusively in groups. Cultivating a microorganism

with lower biosynthetic potential in pure culture might indicate residual amino acids and

vitamins in the filtered sterilized ocean water used as media. Nevertheless, our cultivation

strategy still showed that microorganisms with lower anabolic independence prefer to grow in

groups.

DISCUSSION

The field of microbiology has traditionally relied on the isolation of microorganisms from

nature in pure culture[46–51]. The practice has yielded foundational insights into microbial

physiology[52–54], metabolism[55–58], and ecological interactions such as mutualism[28,59,60], and

competition[61–63]. However, this reductionistic approach has limitations, particularly a strong

cultivation bias favoring microorganisms with larger genome sizes[41,64], broader biosynthetic

potential[65], and greater independence. As a result, a substantial portion of the most abundant

microbes in natural environments remains uncultivated[66].

Here, we demonstrate the utility of microbial model communities established through

dilution cultivation from a Baltic Sea pelagic sample for obtaining a wide range of previously

uncultivated but abundant and biosynthetically limited microorganisms. Despite using only

one type of ocean water as media (which did not allow us to control the presence of amino

acids or vitamins), our method allowed for the cultivation of groups of microorganisms that

showcased important ecological principles that govern microbial life in nature. Our results

suggest that by increasingly using high-throughput dilution cultivation of microbial model

communities, more diverse microorganisms with medium to high biosynthesis potential could

be cultivated. Moreover, by varying cultivation physicochemical parameters, such as adding

catalase[67], changing light regimes[68], or temperature[69], and perhaps by leveraging a better

understanding of bacterial host dynamics[70,71], we will be able to cultivate a greater proportion

of the abundant microorganisms found in nature.

Additionally, we found that species growing in microbial model communities composed of

more than three species exhibited lower biosynthetic potential for both amino acids and B

vitamins compared to species found growing in smaller groups or independently. The reduced

per-genome biosynthetic capacity in more complex communities suggests that

microorganisms with low anabolic independence are forming metabolic networks to support

their nutritional requirements[72]. Interestingly, a previous study has also found a threshold of

microbial diversity at which competition and complementation saturate[30]. Specifically, they

observed that beyond approximately 26 taxa, further increases in diversity had little detectable impact on an overall community function (respiration), suggesting a saturation of functional capacity. Although our work examines only amino acid and vitamin biosynthesis, both studies observe a diversity threshold. These different thresholds observed emphasize that microbial communities likely achieve a balanced state of interaction complexity beyond certain diversity levels, thereby optimizing ecological efficiency.

Our findings also align with the Black Queen Hypothesis, which posits that certain functions, particularly costly biosynthetic pathways, can be lost by some community members as long as the production of the metabolite at the community level is sufficient to sustain the individual[73]. In this context, the reduced genome sizes observed in the abundant species from the model communities and the environment likely reflect a genome streamlining process[74]. The findings further align with the hypothesis that obligate co-existing microbes have evolved to rely on their community for essential nutrients[75], potentially creating social networks that might bring microbial community stability[14,76–78]. To support this, our stepwise module analysis revealed that microbial model communities with more than three species collectively maintain biosynthetic potential for amino acids and vitamins. These observations also align with recent findings[79] from a study which examined the potential for metabolic complementarity among auxotrophic soil bacteria. The study analyzed 746 auxotrophic strains from 27 soil-derived communities that were grown in groups of 2, 3, 4, and 5 strains, and described a clear trend: larger groups of bacteria were more capable of collectively producing all necessary amino acids due to metabolic complementarity.

Importantly, our findings also complement large-scale studies investigating the ecological distribution of auxotrophies. A recent analysis of over 26,000 representative bacterial genomes across diverse environments found that auxotrophy is more common in host-associated environments while relatively rare in aquatic and soil ecosystems. However, this analysis draws a robust but strict boundary on how to categorize microorganisms in a binary model of prototrophs and auxotrophs (more than 40% genes missing per pathway to classify as auxotroph)[16]. While this categorization has been widely used, it might also be hiding potential interdependencies. In our study, we draw no boundary for defining auxotrophy. Instead, we studied the pathways by observing their completeness and comparing them across the different species in the dataset. We believe this has the potential to reveal a wide spectrum of possibilities where microorganisms might just need a precursor to complete biosynthesis[21].

Moreover, we found further support for the idea that metabolic interdependencies go beyond
the simple exchange of end-products in a recent study[20]. The authors used 25 engineered
strains of *E. coli* that were auxotrophic for specific amino acids (arginine, histidine,
isoleucine, proline, and tryptophan) and performed pairwise cocultures of strains auxotrophic
for the same amino acid. Strikingly, they found that growth complementation was often
achieved by sharing the intermediates within the biosynthetic pathway. For this reason, we
believe that moving away from binary categorization of auxotrophies might bring more
nuance to the study of anabolic dependencies.

In summary, our study highlights the larger potential of microbial model communities in
bridging the gap between laboratory cultivation and the environment. By cultivating naturally
assembled groups of microorganisms, we recovered ecologically dominant taxa with limited
biosynthetic capacity that are often overlooked by traditional isolation techniques. These
findings demonstrate that increasing community complexity is associated with different
forms of reduced metabolic autonomy. Moreover, we observed that biosynthetic
interdependencies are widespread in nature. In combination with recent experimental
evidence showing the exchange of biosynthetic intermediates among bacteria, our results
reinforce the idea that anabolic dependencies, rather than complete autonomy, are a
successful ecological strategy. Cultivating groups rather than individuals can offer a more
ecologically relevant understanding of how microbes survive, interact, and evolve in nature.

MATERIALS AND METHODS

Sampling and sample processing
We collected an environmental sample from the surface layer of the Baltic Sea close to Askö
in the Trosa archipelago (Lat 58°48.20'N, Lon 17°37.42'E) in October 2021 (**Figure 1A**).
For details on the chemical composition of the water at sampling time, please visit
https://shark.smhi.se/hamta-data/. The sample for our experiments was subsequently
processed in the laboratory for DNA extraction and metagenomic sequencing. Briefly, the
water sample was filtered through a 0.1 µm membrane and used to extract environmental
DNA using either the FastDNA® SPIN Kit for Soil (MP Bio) and the Dneasy PowerWater kit
(Qiagen). Additionally, we filtered water through a 0.1 µm hollow fiber cartridge (Cytiva)
and used the filtrate as media to establish cultures with different starting inoculum sizes.

Flow cytometry.

We used a CytoFLEX instrument manufactured by Beckman Coulter to process the

environmental sample and to calculate the cell count as the number of events/mL. Briefly, we

stained 50 µl of the samples with Syto13 at a final concentration of 0.025 mM. During the

flow cytometer acquisition process, we set the following parameters: FSC 2500, SSC 2500,

and FITC 800 with a flow rate of 60 µl/min.

Establishment of microbial model communities.

We used the dilution-to-extinction technique considering the cell count information from the

sample and diluting the cells in the filtered sterilized water, to achieve approximately the

desired starting number of cells in our microbial model communities. The model

communities were designed in two ways: low inoculum size and high inoculum size. The low

inoculum size model communities included hundreds of individual cultures in 96-well plates

with a starting number of cells of 0 (control), 2, 6, 10, 15, 20, 30, 50, and 100 cells/well. The

high inoculum size model communities were prepared in bottles with starting numbers of

cells of 0 (control), 200, 600, 1000, 1500, 2000, 3000, 5000, 10000, 100000, and 1000000

cells/bottle. Filtered water from the original environmental sample, without any additional

nutrients, was used as media, resulting in an undefined medium closely reflecting in situ

conditions. All cultures were incubated with light/dark cycles (light 6:42, dark 18:32 each

day) and 12.2 °C at light and 11.8 °C at dark for four weeks before further processing. This

regime was designed to emulate as best as possible the natural fluctuations in time,

temperature, and light that cells experience in their environment.

MDA, library preparation, and sequencing.

Before sending our samples for metagenomic sequencing, we performed multiple

displacement amplification (MDA) on all cultures to increase the concentration of DNA. The

MDA reaction consisted of 0.6 µL of culture and 4.4 µL of reaction mix using the Repli-g

Single Cell kit (Qiagen). After DNA amplification, we found that 315 of the cultures passed

the amplification threshold of the negative controls set in the MDA reaction. These cultures

were deemed positive and were sent for sequencing. We extracted DNA from the selected

cultures and our environmental sample with two different DNA extraction methods and

subjected them to library preparation using the TruSeq PCRfree DNA library preparation kit

(Illumina Inc.) followed by metagenomic Illumina sequencing at the SNP&SEQ Platform at

Uppsala University. This sequencing technology utilized cluster generation and 150 cycles of

paired-end sequencing on an SP flow cell, employing the NovaSeq 6000 system with v1.5

sequencing chemistry (Illumina Inc).

Genome resolved metagenomics pipeline.

We removed low-quality reads from the raw sequences using the software Trimmomatic

(v0.36)[80]   with   the   following   options: ILLUMINACLIP:TruSeq3-PE-

2.fa:2:30:10:2:keepBothReads   LEADING:3   TRAILING:3   SLIDINGWINDOW:4:15

MINLEN:50. We employed the MetaWRAP pipeline (v1.3.2)[81] to process our clean

metagenomic reads. First, clean reads were assembled in a single-sample assembly style

using the "metaWRAP_assembly" module with MegaHit (v1.1.3) for the environmental

samples[82] and with SPAdes (v3.15.3)[83] for the culture samples. The quality of the assemblies

was assessed with QUAST (v.5.0.2)[84]. Since the culture DNA was amplified using MDA, the

read information for coverage could not be used. For this reason, we included background

metagenomic data from previous projects in the Baltic Sea (**Figure 1A** and **Table S2**) for

binning. Subsequently, we used multiple-sample coverage binning to decrease the

contamination and increase the completeness of bins[85]. The reads were mapped against all

assemblies using the Input_POGENOM pipeline[86], which uses Bowtie2[87] with default

parameters. After mapping and obtaining the BAM files, the min coverage was calculated

using samtools (v1.9)[88]. Only coverage values for each assembly in each metagenomic

sample with mean coverage ≥20× and mean breadth ≥40% were retained, following

Input_POGENOM recommendations. These coverage values for each sample were combined

and processed with the "metaWRAP_binning" module, which uses three metagenomic

binning tools: metaBAT2[89], maxBIN2[90], and CONCOCT[91]. We consolidated all the bins

generated by these different tools using the "metaWRAP_bin_refinement" module. We

classified the resulting bins taxonomically with GTDB-tk v2.1.1[92]. Finally, we assessed the

quality of the bins using CheckM (v1.1.3)[93]. We considered bins as MAGs when they had a

completeness of >45% and a contamination of <10%, and these MAGs were included for

further analysis.

Complementing with previously published MAGs.

The de-replicated MAGs obtained here were supplemented with 771 MAGs from an earlier

study[37] that were based on metagenomics data from three studies[37–39]. We de-replicated the

collection of MAGs to obtain species-cluster representatives using ANI > 95% with

mOTUpan (v0.3.2)[94], and selected the genome with the highest quality as the species-cluster

representative genome.

Relative abundance analysis.

To calculate the relative abundance of our 701 species-clusters in the BalticMAG catalog, we

employed the mapping tool Strobealign (v0.14.0)[95] , which aligned the short metagenomic

reads to our species-clusters collection using a high-speed indexing method (**Table S5**).

Briefly, we created three different indexes with different lengths (100, 125, and 150) for our

701 species-clusters. We filtered out low-quality reads from our 112 environmental samples

with Trimmonatic (v0.36)[80] with the following options: ILLUMINACLIP:TruSeq3-PE-

2.fa:2:30:10:2:keepBothReads   LEADING:3   TRAILING:3   SLIDINGWINDOW:4:15

MINLEN:50. After this, we did a competitive mapping of all reads against our created index

to obtain the corresponding BAM file for each sample. We sorted the BAM files with the

Anvi'o platform (v7.1)[96] using the `anvi-init-bam` program, and we calculated the coverage

of each genome per sample with the program `anvi-profile-blitz`. We calculated each species-

cluster's relative abundance with the previously obtained output by dividing each genome's

mean coverage inner quartiles (i.e., q2q3_cov) by the overall sample mean coverage. We also

computed the "prevalence", defined as the frequency of each species-cluster across samples.

Specifically, prevalence represents the proportion of samples in which a species-cluster was

detected with a relative abundance > 0.

Custom functional annotation of KEGG Orthologs (KOs) and biosynthetic modules.

We used the Anvi'o platform (v7.1)[97] to perform functional annotation of KEGG Orthologs

(KOs) and to estimate metabolic potential. Initially, for each genome we used `anvi-gen-

contigs-database` to create a contigs database, which served as the basis for the subsequent

functional annotation steps. To annotate each genome with KOs from the KEGG KOfam

database[98], we ran the `anvi-run-kegg-kofams` program. We then predicted the metabolic

capabilities of each genome by running the `anvi-estimate-metabolism` program [99] (**Table**

S6).

In addition to default KEGG modules[100,101], we developed and implemented a custom set of

30 curated modules targeting the biosynthesis of amino acids and B vitamins (**Table S7**, **S8**,

and **S9**). These custom modules consolidate multiple KEGG modules and fill gaps for

reactions   without KEGG definitions. Custom module definitions and implementation files

are   publicly   available   in   the   accompanying   GitHub   repository:
.

Custom modules were integrated using the `anvi-setup-user-modules` command.
Completeness scores for each genome and custom module were calculated using `anvi-
estimate-metabolism` with the "--only-user-modules" flag. Although we initially created
custom modules for all 20 proteinogenic amino acids (plus the important precursor
chorismate), we subsequently excluded alanine, asparagine, and aspartate modules from
downstream analysis. These three amino acids are commonly produced by generic
transamination reactions with central metabolic intermediates (e.g., pyruvate or oxaloacetate),
and their biosynthesis often involves multiple redundant enzymes that are still challenging to
annotate accurately[44]. Because our custom definitions included only a very limited subset of
these enzymes, we observed artificially low completeness scores for these three modules.
Thus, for accuracy and consistency, we retained only 18 amino acid modules (plus the 9 B
vitamin modules) for downstream statistical comparisons (**Table S10**).

Statistical analyses.

All statistical analyses from this study were performed with R version (v4.4.0)[102] and
RStudio[103]. We used the Shapiro test to assess the normality of our data to be compared; if p
< 0.05, we interpret this as not normally distributed[104]. Since our data was not normally
distributed, we employed a non-parametric test, such as the Wilcoxon test, to find differences
between pairs of groups (e.g., culture vs environment)[105]. To assess statistical differences
among more than two groups, we first applied a Kruskal–Wallis rank-sum test to determine if
any group differed significantly[106]. When the Kruskal–Wallis test was significant ($p < 0.05$),
we performed post-hoc pairwise comparisons using Dunn's test with Bonferroni correction
for multiple testing[107]. To visualize pairwise group differences, we applied a compact letter
display (CLD), where groups that do not differ significantly share the same letter, and groups
with different letters are significantly different from each other.

We also performed a Kruskal-Wallis test to compare different groups, and the p-values were
corrected using the Benjamin-Hochberg procedure. Kruskal-Wallis tests were followed by
Dunn's post hoc test for pairwise comparisons, and significant differences are indicated by
the different letters when $p < 0.05$. All comparisons were reported to be significant if the p-
values were < 0.05.

Data availability. The paired-end sequences of both environmental (n =2) and culture (n =
204) metagenomic samples from the Baltic Sea samples from this study, along with the
corresponding 827 MAGs (>45% completeness and <10% contamination), have been
deposited in NCBI under the BioProject ID PRJNA1134408. All other metagenomes from the
Baltic Sea (n = 110) were downloaded from public repositories, and the metadata is included
in **Table S2**, including their publication reference.
620
621   **ACKNOWLEDGEMENTS**
622

This work was funded by SciLifeLab and by the Swedish Research Council VR (grant 2022-
03077). We thank Jakob Walve from the Marine Laboratory for sampling the Baltic Sea on
October 12, 2021 and providing the sample to us. The authors would like to acknowledge
support from the Genomics infrastructure services at Science for Life Laboratory in Uppsala.
The MDA was done by Claudia Bergin at the Microbial Single Cell facility, and sequencing
was performed by the SNP&SEQ Technology Platform in Uppsala. The facilities are part of
the National Genomics Infrastructure (NGI) Sweden and Science for Life Laboratory. The
SNP&SEQ Platform is also supported by the Swedish Research Council and the Knut and
Alice Wallenberg Foundation. The authors acknowledge support from SNIC/Uppsala
Multidisciplinary Center for Advanced Computational Science for access to the UPPMAX
computational infrastructure as well as the National Academic Infrastructure for
Supercomputing in Sweden (NAISS). Computational work and data handling were enabled
by resources in the projects SNIC 2022/5-392, 2023/5-126, and naiss2023-5-379 provided by
the Swedish National Infrastructure for Computing (SNIC) at UPPMAX, partially funded by
the Swedish Research Council through a grant agreement no. 2018-05973 and 2022-06725.

638
SLG conceptualized, designed, and supervised the research. FB and SLG performed the
cultivation work. AFA and LFD-Z provided the Baltic Sea MAGs that contributed to the
genomic catalog. AP-V, JED, LFD-Z, and SLG conducted bioinformatics analysis. AR-G
supported bioinformatic analysis. IV implemented the custom modules with input from AP-V.
AP-V, AT, and SLG performed the data interpretation and visualization. AP-V led the writing
of the manuscript with input from SLG. All co-authors contributed to the literature searches,
participated in editing and reviewing the manuscript, and approved the final version.
646
CONFLICTS OF INTERESTS

The authors declare no competing interests.

REFERENCES

1.   Konopka, A. What is microbial community ecology? *The ISME Journal* **3**, 1223–1230
(2009).

2.   Falkowski, P. G., Fenchel, T. & Delong, E. F. The Microbial Engines That Drive Earth's
Biogeochemical Cycles. *Science* **320**, 1034–1039 (2008).

3.   Gralka, M. Searching for Principles of Microbial Ecology Across Levels of Biological
Organization. *Integrative And Comparative Biology* **63**, 1520–1531 (2023).

4.   Herman, M. A. *et al.* A Unifying Framework for Understanding Biological Structures
and Functions Across Levels of Biological Organization. *Integrative and Comparative*
Biology **61**, 2038–2047 (2022).

5.   O'Toole, G. A. We have a community problem. *J Bacteriol* e00073-24 (2024)
doi:10.1128/jb.00073-24.

6.   Mee, M. T., Collins, J. J., Church, G. M. & Wang, H. H. Syntrophic exchange in
synthetic microbial communities. *Proc. Natl. Acad. Sci. U.S.A.* **111**, (2014).

7.   Mao, Z. *et al.* The selection of copiotrophs may complicate biodiversity-ecosystem
functioning relationships in microbial dilution-to-extinction experiments. *Environmental*
Microbiome **18**, 19 (2023).

8.   Benoit, G. *et al.* High-quality metagenome assembly from long accurate reads with
metaMDBG. *Nat Biotechnol* **42**, 1378–1383 (2024).

9.   Eren, A. M. & Banfield, J. F. Modern microbiology: Embracing complexity through
integration across scales. *Cell* **187**, 5151–5170 (2024).

10. Méheust, R., Castelle, C. J., Jaffe, A. L. & Banfield, J. F. Conserved and lineage-specific
hypothetical proteins may have played a central role in the rise and diversification of
major archaeal groups. *BMC Biol* **20**, 154 (2022).

11. Hug, L. A. *et al.* A new view of the tree of life. *Nat Microbiol* **1**, 16048 (2016).

12. Anantharaman, K. *et al.* Thousands of microbial genomes shed light on interconnected
biogeochemical processes in an aquifer system. *Nat Commun* **7**, 13219 (2016).

13. Davis, B. D. & Mingioli, E. S. Mutants of Escherichia coli requiring methionine or
vitamin B12. *J Bacteriol* **60**, 17–28 (1950).

14.   Giordano, N. *et al.* Genome-scale community modelling reveals conserved metabolic
cross-feedings in epipelagic bacterioplankton communities. *Nat Commun* **15**, 2721
(2024).

15.   Hessler, T. *et al.* Vitamin interdependencies predicted by metagenomics-informed
network analyses and validated in microbial community microcosms. *Nat Commun* **14**,
(2023).

16.   Ramoneda, J., Jensen, T. B. N., Price, M. N., Casamayor, E. O. & Fierer, N. Taxonomic
and environmental distribution of bacterial amino acid auxotrophies. *Nat Commun* **14**,
(2023).

17.   Rodríguez-Gijón, A. *et al.* The ecological success of freshwater microorganisms is
mediated by streamlining and biotic interactions. Preprint at
https://doi.org/10.1101/2025.03.24.644981 (2025).

18.   Garcia, S. L. *et al.* Auxotrophy and intrapopulation complementary in the 'interactome'
of a cultivated freshwater model community. *Mol Ecol* **24**, 4449–4459 (2015).

19.   Gómez☐Consarnau, L. *et al.* Mosaic patterns of B☐vitamin synthesis and utilization in
a natural marine microbial community. *Environmental Microbiology* **20**, 2809–2823
(2018).

20.   Hong, Y.-J., Cai, Y. & Antoniewicz, M. R. Cross-feeding of amino acid pathway
intermediates is common in co-cultures of auxotrophic Escherichia coli. *Metabolic*
Engineering **88**, 172–179 (2025).

21.   Wienhausen, G. *et al.* Ligand cross-feeding resolves bacterial vitamin B12 auxotrophies.
Nature (2024) doi:10.1038/s41586-024-07396-y.

22.   Oña, L. & Kost, C. Cooperation increases robustness to ecological disturbance in
microbial cross☐feeding networks. *Ecology Letters* **25**, 1410–1420 (2022).

23.   Aziz, F. A. A. *et al.* Interspecies interactions are an integral determinant of microbial
community dynamics. *Front. Microbiol.* **6**, (2015).

24.   Goldford, J. E. *et al.* Emergent simplicity in microbial community assembly. (2018).

25.   Pacheco, A. R., Osborne, M. L. & Segrè, D. Non-additive microbial community
responses to environmental complexity. *Nat Commun* **12**, 2365 (2021).

26.   Bayer, B. *et al.* Metabolite release by nitrifiers facilitates metabolic interactions in the
ocean. *The ISME Journal* wrae172 (2024) doi:10.1093/ismejo/wrae172.

27.   Grant, M. A. A., Kazamia, E., Cicuta, P. & Smith, A. G. Direct exchange of vitamin B12
is demonstrated by modelling the growth dynamics of algal–bacterial cocultures. *The*
ISME Journal **8**, 1418–1427 (2014).

28.   Hillesland, K. L. & Stahl, D. A. Rapid evolution of stability and productivity at the
716   origin of a microbial mutualism. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 2124–2129 (2010).

717   29.   Chang, C.-Y., Bajić, D., Vila, J. C. C., Estrela, S. & Sanchez, A. Emergent coexistence
718   in multispecies microbial communities. *Science* **381**, 343–348 (2023).

719   30.   Yu, X., Polz, M. F. & Alm, E. J. Interactions in self-assembled microbial communities
720   saturate with diversity. *The ISME Journal* **13**, 1602–1617 (2019).

721   31.   Hammarlund, S. P., Gedeon, T., Carlson, R. P. & Harcombe, W. R. Limitation by a
722   shared mutualist promotes coexistence of multiple competing partners. *Nat Commun* **12**,
723   619 (2021).

724   32.   Pande, S. *et al.* Fitness and stability of obligate cross-feeding interactions that emerge
725   upon gene loss in bacteria. *ISME J* **8**, 953–962 (2014).

726   33.   Garcia, S. L. *et al.* Model Communities Hint at Promiscuous Metabolic Linkages
727   between Ubiquitous Free-Living Freshwater Bacteria. *mSphere* **3**, e00202-18 (2018).

728   34.   Ratzke, C., Barrere, J. & Gore, J. Strength of species interactions determines
729   biodiversity and stability in microbial communities. *Nat Ecol Evol* **4**, 376–383 (2020).

730   35.   Steen, A. D. *et al.* High proportions of bacteria and archaea across most biomes remain
731   uncultured. *ISME J* **13**, 3126–3130 (2019).

732   36.   Garcia, S. Mixed cultures as model communities: hunting for ubiquitous
733   microorganisms, their partners, and interactions. *Aquat. Microb. Ecol.* **77**, 79–85 (2016).

734   37.   Alneberg, J. *et al.* Ecosystem-wide metagenomic binning enables prediction of
735   ecological niches from genomes. *Commun Biol* **3**, (2020).

736   38.   Alneberg, J. *et al.* BARM and BalticMicrobeDB, a reference metagenome and interface
737   to meta-omic data for the Baltic Sea. *Sci Data* **5**, (2018).

738   39.   Larsson, J. *et al.* Picocyanobacteria containing a novel pigment gene cluster dominate
739   the brackish water Baltic Sea. *The ISME Journal* **8**, 1892–1903 (2014).

740   40.   Pacheco-Valenciana, A., Garcia, S. L., Dharamshi, J. E., Delgado-Zambrano, L. F. &
741   Andersson, A. F. The BalticMAG catalog.
742   https://doi.org/10.17044/scilifelab.28746086.v1 (2025).

743   41.   Rodríguez-Gijón, A. *et al.* A Genomic Perspective Across Earth's Microbiomes Reveals
744   That Genome Size in Archaea and Bacteria Is Linked to Ecosystem Type and Trophic
745   Strategy. *Front. Microbiol.* **12**, 761869 (2022).

746   42.   Herlemann, D. P. *et al.* Transitions in bacterial communities along the 2000 km salinity
747   gradient of the Baltic Sea. *ISME J* **5**, 1571–1579 (2011).

43. Rodríguez-Gijón, A. *et al.* Linking prokaryotic genome size variation to metabolic
potential and environment. *ISME COMMUN.* **3**, 25 (2023).

44. Price, M. N., Deutschbauer, A. M. & Arkin, A. P. GapMind: Automated Annotation of
Amino Acid Biosynthesis. *mSystems* **5**, 10.1128/msystems.00291-20 (2020).

45. Somerville, V. *et al.* Long-read based de novo assembly of low-complexity metagenome
samples results in finished genomes and reveals insights into strain diversity and an
active phage system. *BMC Microbiol* **19**, 143 (2019).

46. Gich, F., Schubert, K., Bruns, A., Hoffelner, H. & Overmann, J. Specific Detection,
Isolation, and Characterization of Selected, Previously Uncultured Members of the
Freshwater Bacterioplankton Community. *Appl Environ Microbiol* **71**, 5908–5919
(2005).

47. Janssen, P. H., Yates, P. S., Grinton, B. E., Taylor, P. M. & Sait, M. Improved
Culturability of Soil Bacteria and Isolation in Pure Culture of Novel Members of the
Divisions *Acidobacteria* , *Actinobacteria* , *Proteobacteria* , and *Verrucomicrobia*. *Appl*
Environ Microbiol **68**, 2391–2396 (2002).

48. Kaeberlein, T., Lewis, K. & Epstein, S. S. Isolating 'Uncultivable' Microorganisms in
Pure Culture in a Simulated Natural Environment. *Science* **296**, 1127–1129 (2002).

49. Könneke, M. *et al.* Isolation of an autotrophic ammonia-oxidizing marine archaeon.
Nature **437**, 543–546 (2005).

50. Rappé, M. S., Connon, S. A., Vergin, K. L. & Giovannoni, S. J. Cultivation of the
ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**, 630–633 (2002).

51. Schut, F. *et al.* Isolation of Typical Marine Bacteria by Dilution Culture: Growth,
Maintenance, and Characteristics of Isolates under Laboratory Conditions. *Appl Environ*
Microbiol **59**, 2150–2160 (1993).

52. Mella-Flores, D. *et al.* Prochlorococcus and Synechococcus have Evolved Different
Adaptive Mechanisms to Cope with Light and UV Stress. *Front. Microbio.* **3**, (2012).

53. Orr, P. T. & Jones, G. J. Relationship between microcystin production and cell division
rates in nitrogen☐limited Microcystis aeruginosa cultures. *Limnology & Oceanography*
43, 1604–1614 (1998).

54. Sohm, J. A., Edwards, B. R., Wilson, B. G. & Webb, E. A. Constitutive Extracellular
Polysaccharide (EPS) Production by Specific Isolates of Crocosphaera watsonii. *Front.*
Microbio. **2**, (2011).

55. Fischer, F., Zillig, W., Stetter, K. O. & Schreiber, G. Chemolithoautotrophic metabolism
of anaerobic extremely thermophilic archaebacteria. *Nature* **301**, 511–513 (1983).

56.   Leschine, S. B., Holwell, K. & Canale-Parola, E. Nitrogen Fixation by Anaerobic
Cellulolytic Bacteria. *Science* **242**, 1157–1159 (1988).

57.   Zehnder, A. J. & Brock, T. D. Methane formation and methane oxidation by
methanogenic bacteria. *J Bacteriol* **137**, 420–432 (1979).

58.   Zhang, T., Shi, X.-C., Ding, R., Xu, K. & Tremblay, P.-L. The hidden
chemolithoautotrophic metabolism of *Geobacter sulfurreducens* uncovered by
adaptation to formate. *The ISME Journal* **14**, 2078–2089 (2020).

59.   Belzer, C. *et al.* Microbial Metabolic Networks at the Mucus Layer Lead to Diet-
Independent Butyrate and Vitamin B_{12} Production by Intestinal Symbionts. *mBio* **8**,
e00770-17 (2017).

60.   Kazamia, E. *et al.* Mutualistic interactions between vitamin B_{12} ⬚dependent algae and
heterotrophic bacteria exhibit regulation. *Environmental Microbiology* **14**, 1466–1476
(2012).

61.   Chodkowski, J. L. & Shade, A. Bioactive exometabolites drive maintenance competition
in simple bacterial communities. *mSystems* e00064-24 (2024)
doi:10.1128/msystems.00064-24.

62.   Harrison, F., Paul, J., Massey, R. C. & Buckling, A. Interspecific competition and
siderophore-mediated cooperation in *Pseudomonas aeruginosa*. *The ISME Journal* **2**,
49–55 (2008).

63.   Rao, D., Webb, J. S. & Kjelleberg, S. Competitive Interactions in Mixed-Species
Biofilms Containing the Marine Bacterium *Pseudoalteromonas tunicata*. *Appl Environ*
Microbiol **71**, 1729–1736 (2005).

64.   Han, K. *et al.* Extraordinary expansion of a Sorangium cellulosum genome from an
alkaline milieu. *Sci Rep* **3**, 2101 (2013).

65.   Bentkowski, P., Van Oosterhout, C. & Mock, T. A Model of Genome Size Evolution for
Prokaryotes in Stable and Fluctuating Environments. *Genome Biol Evol* **7**, 2344–2351
(2015).

66.   Lloyd, K. G., Steen, A. D., Ladau, J., Yin, J. & Crosby, L. Phylogenetically Novel
Uncultured Microbial Cells Dominate Earth Microbiomes. *mSystems* **3**, e00055-18
(2018).

67.   Kim, S., Kang, I., Seo, J.-H. & Cho, J.-C. Culturing the ubiquitous freshwater
actinobacterial acI lineage by supplying a biochemical 'helper' catalase. *ISME J* **13**,
2252–2263 (2019).

68. Bialevich, V., Zachleder, V. & Bišová, K. The Effect of Variable Light Source and Light
Intensity on the Growth of Three Algal Species. *Cells* **11**, 1293 (2022).

69. Jiang, L. & Morin, P. J. Temperature fluctuation facilitates coexistence of competing
species in experimental microbial communities. *Journal of Animal Ecology* **76**, 660–668
(2007).

70. Kuroda, K. *et al.* Microscopic and metatranscriptomic analyses revealed unique cross-
domain parasitism between phylum *Candidatus* Patescibacteria/candidate phyla
radiation and methanogenic archaea in anaerobic ecosystems. *mBio* **15**, e03102-23
(2024).

71. Man, D. K. W. *et al.* Enrichment of different taxa of the enigmatic candidate phyla
radiation bacteria using a novel picolitre droplet technique. *ISME Communications* **4**,
ycae080 (2024).

72. Johnson, W. M. *et al.* Auxotrophic interactions: a stabilizing attribute of aquatic
microbial communities? *FEMS Microbiology Ecology* **96**, fiaa115 (2020).

73. Morris, J. J., Lenski, R. E. & Zinser, E. R. The Black Queen Hypothesis: Evolution of
Dependencies through Adaptive Gene Loss. *mBio* **3**, e00036-12 (2012).

74. Giovannoni, S. J., Cameron Thrash, J. & Temperton, B. Implications of streamlining
theory for microbial ecology. *ISME J* **8**, 1553–1565 (2014).

75. Sokolovskaya, O. M., Shelton, A. N. & Taga, M. E. Sharing vitamins: Cobamides unveil
microbial interactions. *Science* **369**, (2020).

76. Kost, C., Patil, K. R., Friedman, J., Garcia, S. L. & Ralser, M. Metabolic exchanges are
ubiquitous in natural microbial communities. *Nat Microbiol* **8**, 2244–2252 (2023).

77. Zelezniak, A. *et al.* Metabolic dependencies drive species co-occurrence in diverse
microbial communities. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 6449–6454 (2015).

78. Zengler, K. & Zaramela, L. S. The social network of microorganisms — how
auxotrophies shape complex communities. *Nat Rev Microbiol* **16**, 383–390 (2018).

79. Yousif, G. *et al.* Obligate cross-feeding of metabolites is common in soil microbial
communities. Preprint at https://doi.org/10.1101/2025.01.29.635426 (2025).

80. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

81. Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for
genome-resolved metagenomic data analysis. *Microbiome* **6**, 158 (2018).

82. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-
node solution for large and complex metagenomics assembly via succinct *de Bruijn*
graph. *Bioinformatics* **31**, 1674–1676 (2015).

83. Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications
to Single-Cell Sequencing. *Journal of Computational Biology* **19**, 455–477 (2012).

84. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

85. Mattock, J. & Watson, M. A comparison of single-coverage and multi-coverage
metagenomic binning reveals extensive hidden contamination. *Nat Methods* **20**, 1170–
1173 (2023).

86. Sjöqvist, C., Delgado, L. F., Alneberg, J. & Andersson, A. F. Ecologically coherent
population structure of uncultivated bacterioplankton. *The ISME Journal* **15**, 3034–3049
(2021).

87. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**, 357–359 (2012).

88. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

89. Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient
genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

90. Wu, Y.-W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning
algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**,
605–607 (2016).

91. Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat*
Methods **11**, 1144–1146 (2014).

92. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927
(2020).

93. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

94. Buck, M., Mehrshad, M. & Bertilsson, S. mOTUpan: a robust Bayesian approach to
leverage metagenome-assembled genomes for core-genome estimation. *NAR Genomics*
and Bioinformatics **4**, lqac060 (2022).

95. Sahlin, K. Strobealign: flexible seed size enables ultra-fast and accurate read alignment.
Genome Biol **23**, 260 (2022).

96. Eren, A. M. *et al.* Anvi'o: an advanced analysis and visualization platform for 'omics
data. *PeerJ* **3**, e1319 (2015).

97. Eren, A. M. *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat*
Microbiol **6**, 3–6 (2020).

98. Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM
and adaptive score threshold. *Bioinformatics* **36**, 2251–2252 (2020).

99. Veseli, I. *et al.* Microbes with higher metabolic independence are enriched in human gut
microbiomes under stress. *eLife* **12**, RP89862 (2025).

100. Kanehisa, M. *et al.* Data, information, knowledge and principle: back to metabolism in
KEGG. *Nucl. Acids Res.* **42**, D199–D205 (2014).

101. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new
perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45**, D353–
D361 (2017).

102. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing (2024).

103. RStudio Team. RStudio: Integrated Development for R. (2024).

104. Royston, J. P. An Extension of Shapiro and Wilk's W Test for Normality to Large
Samples. *Applied Statistics* **31**, 115 (1982).

105. Bauer, D. F. Constructing Confidence Sets Using Rank Statistics. *Journal of the*
American Statistical Association **67**, 687–690 (1972).

106. Kruskal, W. H. & Wallis, W. A. Use of Ranks in One-Criterion Variance Analysis.
Journal of the American Statistical Association **47**, 583–621 (1952).

107. Dunn, O. J. Multiple Comparisons Using Rank Sums. *Technometrics* **6**, 241–252 (1964).

Figure Legends

Figure 1. The BalticMAG catalogue. (A) Map of the Baltic Sea showing the geographic location of
all the metagenomes analyzed in this study, including our sampling site (n =112). Sampling sites are
color-coded according to their salinity gradient (measured in ‰, equivalent to PSU or parts per
thousand). The shapes correspond to the different reference sources. (B) Boxplot showing the
percentage of mapped reads from all metagenomes to our species-cluster collection, categorized by
salinity concentration (Kruskal-Wallis test). (C) Venn diagram showing the overlap of species-clusters
presence among cultures (red), the environment (blue), and found in both (purple). (D) Relative
abundance of species-clusters across the collection of metagenomes, ordered by salinity gradient from
left to right (corresponding to the order in panel B). (E) Boxplot comparing the relative abundance of
species-clusters categorized by their presence in cultures, the environment, or both (Kruskal-Wallis
test). Kruskal-Wallis tests were followed by Dunn's post hoc test for pairwise comparisons. Groups

sharing at least one letter (e.g., a and ab) are not significantly different from each other, while groups
with different letters (e.g., a vs. b) are significantly different (p < 0.05).
920
Figure 2. Impact of inoculum size on microbial richness and genome size. (A) Relationship between
the number of inoculated cells (low inoculum size = 0 to 100 cells/well) and the total number of
cultures from which MAGs were assembled and binned. (B) Number of MAGs per culture for the low
inoculum size (yellow) and (C) the high inoculum size (pink) microbial model communities. (D)
Estimated genome size (circles) of MAGs from cultures (n = 527) in relation to the number of species
growing in the same culture, and (E) average estimated genome size (triangles) of MAGs growing in
the same culture. (F) Boxplot comparing the average estimated genome size of species-clusters from
cultures and those from the environment (Wilcoxon rank-sum test).
929
Figure 3. Microbial model communities of increased complexity host distinct sets of cultured species.
Heatmap showing the distribution of the 72 species-clusters across cultures grouped by community
complexity: 1, 2, 3, or more than 3 species per culture. Taxonomic affiliations (class, genus, and
species) based on GTDB-tk are displayed alongside their unique species-cluster IDs. The white-to-
black gradient indicates the number of genomes recovered per species-cluster in each culture
category. Species that exclusively grow alone are highlighted in a light orange box, while those only
found to grow in groups (≥2 genomes) are highlighted in light green.
937
Figure 4. Culturing in groups enables access to a greater diversity of microbial species. Rarefaction
curves show the cumulative number of unique species recovered from microbial model communities
grouped by community complexity (1, 2, 3, or more than 3 species), as cultured samples are
progressively added. For each category, species accumulation was calculated across 100 random
permutations (bootstrapped iterations), and shaded ribbons represent ±1 standard deviation from the
mean. Communities with a single species (1 genome) are shown in light orange, while increasingly
complex communities (≥2 genomes) are shaded in progressively darker greens.
945
Figure 5. Growth strategy, environmental distribution, and genomic features of cultured microbial
species. (A) Barplot showing the total number of cultured genomes recovered per species-cluster (n =
72). Bars are colored by the type of culture from which each genome was obtained: light orange for
single-genome cultures and light green for multi-genome cultures. Species-clusters were further
categorized based on their observed growth behavior: "Single" (light orange) for species that grew
exclusively alone, "Group" (light green) for those that grew exclusively in groups, and "Both"
(orange) for species that could grow alone and in groups. (B) Boxplot comparing the number of
cultured genomes per species-cluster across the three growth categories (Kruskal-Wallis test, followed
by Dunn's post hoc test with Bonferroni correction; p < 0.05). Letters indicate results of Dunn's post-
hoc test (Bonferroni-corrected); groups with different letters are significantly different. (C) The dot
plots display the relationship between the total number of cultured genomes per species-cluster and
their relative abundance in the sample of origin (D), as well as the prevalence of these species-clusters
in all environmental samples. Each data point represents one species-cluster. A non-linear regression
models (Spearman) were fitted to the data. (E) Boxplot comparing the estimated genome size of
species-cluster representative genomes (n = 72) growing exclusively in groups ("Group" = light
green) and capable of growing alone ("Singleton-capable" = light orange). (F) Boxplot comparing the
average pathway completeness of custom amino acid and vitamin biosynthesis modules between
high-quality species-cluster representatives (n = 57; completeness >90%, contamination <5%)
growing exclusively in groups and those capable of growing alone. Statistical significance was
assessed using the Wilcoxon rank-sum test.
966
Figure 6. Amino acid and vitamin biosynthetic potential across species from varying community
complexity. (A) Heatmap showing pathway completeness scores for 18 custom amino acid
biosynthesis modules (three-letter abbreviations) across high-quality genomes (n = 305; >90%
completeness, <5% contamination) grouped by community complexity: 1, 2, 3, or more than 3
genomes per culture. Each column represents a genome, and each row a biosynthetic module.
Completeness values range from white (0) to dark orange (1) for genomes from single-genome

cultures, and white to dark green for genomes from multi-genome cultures (≥2 genomes). (B) Boxplot summarizing the average amino acid pathway completeness per genome across the same four community complexity groups (Kruskal-Wallis test, followed by Dunn's post hoc test with Bonferroni correction; $p < 0.05$). Groups with different letters are significantly different. (C) Heatmap as in panel A, but for the 9 custom vitamin biosynthesis modules. Color gradients and grouping are defined identically. (D) Boxplot summarizing the average vitamin pathway completeness per genome across complexity groups. Statistical analysis performed as in panel B.

Figure 7. Stepwise module framework analysis of amino acid and B vitamin biosynthetic pathways across microbial model communities.
Heatmap shows the average presence of biosynthetic module steps across the 204 microbial model communities, including all genomes, regardless of genome completeness. Each column represents a specific KEGG module step (e.g., M000020_01) grouped by compound. Each row corresponds to an individual model community, grouped by community complexity (1, 2, 3, or more than 3 species per culture). Values represent the proportion of genomes in each community that encode the respective module step, ranging from 0 to 1. Single-genome communities are displayed using an orange gradient (white to dark orange), while multi-genome communities (≥2 genomes) use a green gradient (white to dark green). This stepwise framework uses canonical KEGG modules (non-customized), capturing the presence of standard and alternative steps within pathways.

Figure 8.
Link between pathway completeness, relative abundance, and estimated genome size of cultured species and the BalticMAG species catalog. (A) Correlation between average biosynthesis pathway completeness of custom amino acids and vitamin modules and their relative abundance of cultured species (n = 57, completeness >90%, contamination <5%). (B) Correlation between the average biosynthesis completeness for custom vitamin modules and the estimated genome size of cultured species-clusters. Each data point represents one species cluster (n = 57, completeness >90%, contamination <5%), which either grew exclusively on their own (light orange), exclusively in groups (light green), or both on their own as well as in groups (orange). Panels (C) and (D) replicate the analysis from panels (A) and (B) but include all species-clusters (n = 450, completeness >90%, contamination <5%) from both cultures and environmental samples. (E) Correlation between estimated genome size and average relative abundance and (F) correlation between estimated genome size and prevalence of all species-clusters (n = 701). Data points are color-coded to represent species clusters from cultures only (red), environments only (blue), and found in both (purple).

Figure 9. Anabolic dependencies and biosynthetic clustering of the BalticMAG species catalog inlcuding cultivated species. (A) Principal component analysis (PCA) of the biosynthetic module completeness, with genomes color-coded by biosynthetic completeness group: Low = light blue, Medium = light purple, High = pink. (B) Uniform Manifold Approximation and Projection (UMAP) of the same data, confirming the biosynthetic groupings. (C) Stacked bar plot showing the proportion and taxonomic composition of the three biosynthetic completeness groups, color-coded by phylum. (D–F) Boxplots comparing (D) average pathway completeness, (E) estimated genome size (Mbp), and (F) relative abundance of species-clusters grouped by biosynthetic completeness. Statistical significance was assessed using a Kruskal–Wallis test ($p < 0.05$) followed by Dunn's post hoc test with Bonferroni correction; groups with different letters are significantly different ($p < 0.05$). (G) Median completeness differences for KEGG custom biosynthesis modules between Cluster A and Cluster B genomes of the High biosynthetic group. Positive values indicate modules are more complete in Cluster A, while negative values indicate modules are more complete in Cluster B. (H–I) Boxplots comparing (H) estimated genome size (Mbp) and (I) relative abundance of species-clusters from Cluster A (dark pink) and Cluster B (light pink). Statistical significance was assessed using the Wilcoxon rank-sum test ($p < 0.05$). (J) PCA of biosynthetic completeness color-coded by species-cluster growth category: Single = light orange, Group = light green, Both = orange; environmental-only clusters are shown in grey. (K) PCA of the 57 high-quality genomes from microbial model communities, also color-coded by growth category. (L) Boxplot showing the average pathway completeness of the three biosynthetic clusters derived from cultivated species. Statistical significance

1028   was tested using Kruskal–Wallis with Dunn's post hoc test; groups with different letters are
1029   significantly different ($p < 0.05$).