Microbial model communities reveal widespread auxotrophies in abundant bacteria

Armando Pacheco-Valenciana[1*], Anna Tausch[2], Jennah E. Dharamshi[1], Fabian Bergland[1], Luis Fernando

Delgado-Zambrano[3], Alejandro Rodríguez-Gijón[1], Anders F. Andersson[3], Sarahi L. Garcia[1,2,4*].

[1] Department of Ecology, Environment, and Plant Sciences, Science for Life Laboratory, Stockholm University,

Stockholm, Sweden.

[2] Institute for Chemistry and Biology of the Marine Environment (ICBM), School of Mathematics and Science,

Carl von Ossietzky Universität Oldenburg, 26129, Oldenburg, Germany

[3] Department of Gene Technology, Science for Life Laboratory, KTH Royal Institute of Technology, Stockholm,

Sweden.

[4] Helmholtz Institute for Functional Marine Biodiversity at the University of Oldenburg (HIFMB), Oldenburg,

Germany

Correspondence: armando.pacheco@su.se, sarahi.garcia@su.se

KEYWORDS: microbial model communities, metagenomics, auxotrophs, microbial interdependencies.

ABSTRACT

Microorganisms exist in complex communities shaped by intricate interactions. Here, we

examined auxotrophies and metabolic interdependencies in microbial model communities

established from Baltic Sea pelagic sample dilutions. We recovered 527 metagenome-

assembled genomes (MAGs) from 204 microbial model communities, including some of the

most abundant taxa such as Pelagibacterales. Combined with 1073 environmental MAGs, we

created a genomic catalog of 701 species-clusters (95% average nucleotide identity - ANI). Our

results show that more complex communities included microorganisms with smaller genome

sizes and communities with more than three species exhibited lower average biosynthetic

potential for amino acids and B vitamins. Furthermore, microorganisms with lower

biosynthetic potential for those compounds are more abundant in the environment and have

smaller estimated genome sizes. Overall, our results highlight the value of cultivating

microorganisms in groups of naturally co-occurring microorganisms for unraveling ecological

strategies such as auxotrophies and genomic streamlining which are widespread features of

successful microorganisms.

INTRODUCTION

Microbial communities in diverse environments operate as complex systems driven by multi-species interactions[1]. Understanding such complex interactions is essential because microorganisms play key roles in the biogeochemical cycles on Earth[2]. To unravel microbial interactions, we need to investigate microbial communities at various levels of biological organization[3,4], ranging from one-to-one species interactions, to model communities[5], to naturally occurring communities.

At the community level, metagenomics has become a powerful tool for uncovering the genetic potential of microbial communities without the need for isolating individual species[6–8]. Analyzing metagenomic data reveals not only the vast diversity of microbial species[9] but also the metabolic potential and co-occurrence networks, which are important for ecosystem functioning[10]. To bridge the gap between broad metagenomic insights and detailed ecological understanding, a few studies have explored genome-specific traits and their ecological implications. Some of these studies have found a link between small genome sizes and a higher number of auxotrophies for both amino acids and vitamins[11–14]. Specifically, auxotrophs are often deficient in biosynthetic pathways required for producing energetically costly but essential compounds such as amino acids[15–17] and certain vitamins (e.g., B_1, B_7, B_{12})[18–20]. Auxotrophic organisms can acquire the essential metabolites from other community members to meet their nutritional requirements[21–23]. Beyond individual interactions, modeling work has shown that microbial communities enriched in auxotrophies can exhibit greater robustness under ecological disturbances, suggesting that these metabolic interdependencies may contribute to overall community stability[24]. While metagenomics offers a broad understanding of microbial communities, interactions cannot easily be derived from co-occurrences within natural complex ecosystems.

Experimental systems are needed to observe microbial interaction dynamics under controlled conditions. Studies have increasingly turned to simplified systems[25–27], such as co-cultures[28–30] and mixed cultures[16,31,32], in order to identify specific types of interactions. To further contextualize these findings, our literature review in **Table S1** provides a comprehensive overview of publications where microbial interaction patterns were observed in different experimental settings, including isolation attempts, co-cultures, and different versions of microbial model communities. Across these studies, cross-feeding mechanisms and mutualistic

interactions are mostly studied in cultures with two different populations or species[29,33,34]. To a lesser extent, more complex metabolic interactions have also been studied by mixing different isolated species together in model ecosystems, with the goal of increasing complexity to more closely resemble natural environments[31,35,36]. However, these methods focus on cultured isolates, and the vast majority of microorganisms remain uncultivated[37]. An alternative, yet underutilized, method of establishing model ecosystems composed of previously uncultivated microorganisms is through dilution cultivation. Dilution to mixed-cultures of naturally co-occurring microorganisms has the potential to allow observation of natural microbial interactions[37]. Such cultures, also known as microbial model communities, represent a small subset of the many interactions likely occurring in the natural system[35]. By studying a larger number of microbial model communities, we can gain a more comprehensive understanding of microbial interactions occurring in a natural environment.

In our study, we have focused on studying auxotrophies and potential interdependencies at two levels of biological organization by using high-throughput dilution cultivation of model communities together with genome-resolved metagenomics to unravel ecological strategies of microorganisms in the Baltic Sea. The 204 obtained microbial model communities revealed insights into community complexity. Moreover, we identified correlations between genome size, potential biosynthesis of essential metabolites, relative abundance, and prevalence using genomes obtained from both microbial model communities and metagenomic data from Baltic Sea pelagic samples. Our findings demonstrate that microbial model communities are an effective cultivation technique for identifying putative microbial interactions, including auxotrophies that lead to metabolic interdependencies and thereby promote stable microbial community structure and function.

RESULTS

A Baltic Sea MAG catalog

To generate the microbial model communities, we used the dilution-to-extinction cultivation technique in two formats. The first type, low inoculum size (LS), involved inoculating between approximately 2 and 100 cells per well in 1 ml 96-well plates used for each of the inoculum sizes. The second type, high inoculum size (HS), ranged from approximately 200 to 1×10^6 cells inoculated per microbial model community in 100 ml volumes. After a 4-week incubation period, we sent an aliquot of all cultures for lysis and DNA amplification. Out of 801 cultures,

315 passed the amplification threshold and were sent to sequencing. In total, 204 microbial

model communities together yielded 527 MAGs. Moreover, from the original sample used to

establish the microbial model communities, we generated two metagenomes (each from a

distinct DNA extraction method) that yielded 305 MAGs. To create a comprehensive genomic

catalog, we added 771 MAGs deriving from 110 Baltic Sea metagenomes (**Figure 1A**, **Table**

S2)[39–41].

Examining this comprehensive MAG catalog allowed us to assess microbial genomic

characteristics of all microorganisms found both in the environmental samples and our 204

microbial model communities (**Table S3**). Altogether, the 1603 MAGs were dereplicated into

701 species-clusters (ANI > 95%), which form the Baltic Sea genomic catalog (BalticMAG

catalog)[42] used in this study (**Table S4**). The average completeness of the 701 species-cluster

representative MAGs is 88%, and they were all used to analyze taxonomy, abundance,

prevalence, and estimated genome size (**Figure S1**). The varying completeness of the MAGs

has a very minor effect on estimated genome size and relative abundance, just as observed in

other studies[14,43]. However, only 450 species-cluster representative MAGs are of high-quality

(completeness >90% and contamination <5%) and were used to reliably investigate metabolic

potential (**Figure S1D**).

Examining the source of genomes in the BalticMAG catalog, we found that 33 of the species-

clusters included MAGs exclusively from the microbial model communities, 629 included

MAGs exclusively from the environmental metagenomes, and 39 (54% of all cultured species)

included MAGs from both sources (**Figure 1C, Table S4**, **Figure S2).**

To investigate the relative abundance of the BalticMAG catalog, we mapped all the

environmental metagenomics reads against the genome catalog. We observed that salinity

significantly co-varied with the proportion of metagenomic reads that mapped to all species

representative genomes (**Figure 1B**). While on average, 38.63% of the metagenomic reads per

sample mapped to the BalticMAG catalog, the highest mapping percentage (63.99% and

62.89%) was observed at salinity concentrations 11.28‰ and 7.65‰, respectively. Notably,

the two metagenomes from this study (sample with salinity concentration 7.12‰) displayed

some of the highest mapping rates at 51.77% and 60.16%, reflecting that the BalticMAG

catalog is most complete for salinities between 6 and 11‰.

Figure 1. The BalticMAG catalogue. (A) Map of the Baltic Sea showing the geographic location of all
the metagenomes analyzed in this study, including our sampling site (n =112). Sampling sites are color-
coded according to their salinity gradient (measured in ‰, equivalent to PSU or parts per thousand).
The shapes correspond to the different reference sources. (B) Boxplot showing the percentage of
mapped reads from all metagenomes to our species-cluster collection, categorized by salinity
concentration (Kruskal-Wallis test). (C) Venn diagram showing the overlap of species-clusters presence
among cultures (red), the environment (blue), and found in both (purple). (D) Relative abundance of
species-clusters across the collection of metagenomes, ordered by salinity gradient from left to right
(corresponding to the order in panel B). (E) Boxplot comparing the relative abundance of species-
clusters categorized by their presence in cultures, the environment, or both (Kruskal-Wallis test).

Despite salinity differences (**Figure 1D**), the average relative abundance of the 39 species-

clusters that included MAGs from both microbial model communities and environmental

metagenomes was significantly higher in the whole dataset (**Figure 1E),** but also when

comparing only the metagenomes from the location and salinity from which we sampled

(**Figure S3**). Moreover, the 33 species-clusters with MAGs sourced exclusively from microbial

model communities were detected across environmental metagenomes, albeit at significantly

lower abundances. Interestingly, 26 of these 33 species-clusters grew in model communities

with more than two members and were less abundant in the environment (**Figure 1E**),

indicating that the cultivation approach not only captures abundant taxa but also enables the

recovery of species that may be missed in metagenomic surveys.

Model communities with more than three species harbor the most diverse and prevalent

taxa

In the LS microbial model communities, the more cells we inoculated, the higher the number

of cultures that yielded MAGs (**Figure 2A**). In total, 94 LS model communities yielded only

one MAG each, while 110 model communities resulted in two or more MAGs. In model

communities with more than one MAG, each MAG belonged to a different species-cluster in

our analysis. Therefore, for clarity, we refer to different MAGs within a model community as

different species. On the one hand, microbial model communities with an inoculum size of less

than approximately 30 cells primarily resulted in single-species cultures (**Figure 2B**). On the

other hand, starting from an inoculum size of approximately 30 cells, microbial model

communities with more than two species appear more often (**Figure 2B** and **C**). Nearly 82%

(n = 433) of the microbial model community MAGs were obtained from multi-species cultures,

and the highest number of co-occurring species were found in the HS microbial model

communities inoculated with 5000 cells (**Figure 2C**). Despite using a complex inoculum from

the Baltic Sea, observing a maximum of 13 co-occurring species suggests that our cultivation

conditions may impose a threshold on model community complexity. Alternatively, since

sequencing followed MDA, there is also the possibility that some model communities included

more species that were not amplified or assembled and binned. Nevertheless, the increased

growth success at higher inoculum sizes may be attributed to a greater chance of inoculating

cells that will grow either alone or together.

Figure 2. Impact of inoculum size on microbial richness, genome size, and taxonomic diversity of cultures. (A) Relationship between the number of inoculated cells (low inoculum size = 0 to 100 cells/well) and the total number of cultures from which MAGs were assembled and binned. (B) Number of MAGs per culture for the low inoculum size (yellow) and (C) high inoculum size (pink) cultures. (D) Estimated genome size (circles) of MAGs from cultures (n = 527) in relation to the number of

species growing in the same culture, and (E) average estimated genome size (triangles) of MAGs growing in the same culture. (F) Boxplot comparing the average estimated genome size of species-clusters from cultures and those from the environment (Wilcoxon rank-sum test). (G) Barplot showing the total genome count in relation to the number of genomes recovered per culture. Genomes are color-coded by genera, and the legend shows abbreviated prefixed indicating the class of each genus (e.g., Alph_ for Alphaproteobacteria, Bact_ for Bacteroidia, Camp_ for Campylobacteria, Cyan_ for Cyanobacteria, Gamm_ for Gammaproteobacteria, and Verr_ for Verrucomicrobiae. (H) Rarefaction curves of cultured microbial genera by species-per-culture category. Each curve represents the mean cumulative number of unique genera recovered as cultured samples are progressively added. Cultures were categorized into four groups based on the number of species they contained: 1, 2, 3, or >3 species per culture. For each category, genus accumulation was calculated across 100 random permutations (bootstrapped iterations), and shaded ribbons represent ±1 standard deviation from the mean.

Among the 527 MAGs obtained from the microbial model communities, the estimated genome sizes ranged from 0.88 to 6.27 Mbp, with an average of 3.05 Mbp (**Figure 2D** and **F**). For the MAGs derived from Baltic Sea metagenomes, estimated genome sizes ranged from 0.61 to 12.32 Mbp, with an average of 3.00 Mbp. While there was no significant difference between the average estimated genome sizes of species-clusters from microbial model communities and Baltic Sea metagenomes, we found that species-clusters exclusively composed of genomes from microbial model communities had significantly larger genome sizes than species-clusters including genomes from both sources (**Figure S4**). Moreover, we found that as the number of retrieved MAGs per microbial model community increased, more microorganisms with smaller genomes emerged, as indicated by a slight but significant correlation between estimated genome size and the number of MAGs per microbial model community (**Figure 2D**). Further analysis reinforced the observation that genomes from HS microbial model communities and from model communities with more than 3 species had on average, smaller genome sizes (2.89 and 2.92 Mbp, respectively - **Figure S5** and **S6**). Genomes from HS microbial model communities were also on average more prevalent and had higher average relative abundance in the investigated Baltic Sea metagenomes. Additionally, the average genome size per microbial model community stabilized at around 3 Mbp in cultures containing more than three species (**Figure 2E**).

Furthermore, the taxonomic diversity in our microbial model communities spanned 51 genera across 5 of the most abundant phyla (*Pseudomonadota*, *Bacteroidota*, *Campylobacterota*, *Cyanobacteriota*, and *Verrucomicrobiota*) present in the Baltic Sea environmental metagenomic sample[44,45], with microbial model communities with more than three species exhibiting a higher cumulative and rarefied diversity at the genus level (**Figure 2G** and **H**).

The analysis illustrates how more diversity growing together allows for the cultivability of different taxa that would normally not grow in axenic cultures under the conditions tested.

The diverse taxa cultivated in model communities together accounted for a relative abundance of approximately 20% in the original environmental sample (**Figure S7**). Notably, one of our HS model communities yielded a MAG that was found to be very abundant in the environment (~4.7%) classified in the order Pelagibacterales (SAR11) but from an unnamed genus according to GTDB-tk v2.1.1[46]. The order Pelagibacterales is widely recognized as one of the most abundant yet challenging to cultivate microorganisms in the ocean surface waters due to their nutritional requirements, which has been attributed to their small genome size and low biosynthetic potential[47,48]. In our study, the Pelagibacterales MAG possessed an estimated genome size of 0.93 Mbp and grew in a community of at least 10 species which had larger genome sizes, some up to 6 Mbp. Both the large novel diversity cultivated and their high abundance in nature highlight the potential of model communities to capture hard-to-cultivate yet abundant microbes from the environment, offering a promising future for microbial cultivation.

Species strictly found growing in groups are found more frequently in the environment.

In our microbial model communities, a total of 72 species were identified. The prevalence of species among the model communities varied greatly, from some detected only once (e.g., sc_576 and sc_639, names of species-clusters are found in **Table S4** and **Figure 3**) to some found in over 40 model communities (e.g., sc_011 and sc_014). Fifteen species (21%) were present in more than 10 model communities, collectively accounting for approximately 63% of the total MAGs (330 out of 527). Distinct patterns were observed across species, with 4 species found exclusively on their own (e.g., sc_035 and sc_281) and 48 species found only in groups (e.g., sc_102 and sc_177). Notably, 20 species were found both independently and in groups (e.g., sc_011 and sc_014). Most of the frequently retrieved MAGs showed this flexibility, growing independently or in groups (**Figure 3A** and **B**). However, our experiment did not systematically test if species only found on their own, could also grow in groups or if species found in groups only, could also grow alone. Nevertheless, when we examined presence patterns in the microbial model communities in relation to relative abundance across the Baltic Sea environmental sample used as inoculum, we see a trend for the species found both independently and in groups (**Figure 3C**). The total number of genomes found in the cultures per species increases as their relative abundance in the environment of origin increases.

On the other hand, the two most abundant species identified in the environment of origin belong

to the order Pelagibacterales (sc_139) and Methylacidiphilales (sc_121) and were found

exclusively in groups and cultivated only a few times. Moreover, for species that demonstrated

the ability to grow both independently and in groups, as well as those that exclusively grew in

groups, a significant positive correlation emerged between the total number of genomes per

species and their prevalence in the environment (**Figure 3D**). The relationship indicates that

species with higher environmental prevalence are more frequently cultured in laboratory

settings. Moreover, the correlation is further enhanced when species grow more easily, meaning

those found growing both independently and in groups. Nevertheless, the species with the

highest prevalence across all environmental samples, as well as the highest relative abundance

in the sample of origin, were found exclusively growing in groups.

Figure 3. Cultured genomes per species-cluster in relation to their observed growth condition (alone,
group, or both), relative abundance in the sample of origin, and prevalence in the Baltic Sea. (A) Barplot
displaying the total number of cultured genomes for each species-cluster. Color coding indicates
whether the genomes originate from single-genome-cultures (light orange) or multi-genome-cultures
(green). Species-clusters were further differentiated according to whether they grew exclusively on their
own ("Single" = light orange), exclusively in groups ("Group" = green), or both on their own as well
as in groups ("Both" = orange). (B) Boxplot showing the total number of cultured genomes per species-

cluster for these three growth categories. Letters indicate groups with significant differences (p < 0.05)
(Kruskall-Wallis-Test followed by a Dunn's post-hoc test with Bonferroni correction). (C) The dot plots
display the relationship between the total number of cultured genomes per species-cluster and relative
abundance in the sample of origin (D) as well as the prevalence of these species-clusters in all
environmental samples. Each data point represents one species-cluster. Linear regression models were
fitted to the data.

Auxotrophies and interdependencies are more common in MAGs from model communities with more than three species

We compared estimated genome sizes and biosynthetic potential among species found on their

own sometimes or always (singleton-capable) or strictly in groups (group-only). Although

there was no significant difference in estimated genome sizes between singleton-capable and

group-only species (**Figure 4A**), we observed a significant difference in average pathway

completeness for both amino acids and vitamin modules, with species found exclusively in

groups exhibiting a lower average pathway completeness (**Figure 4B**). We next examined

biosynthetic potential across all microbial model communities, categorizing them by their

number of species (one, two, three, or more than three). Here, we found that average pathway

completeness for amino acids and vitamins was significantly lower for model communities

with more than 3 species found together (**Figure 4C** and **D**). Amino acid biosynthesis potential

is in particular lower for species that exclusively grow in groups and are found in model

communities that contain more than 3 species (**Figure S8**).

We explored interdependencies among species growing together. For this, we used the stepwise

module framework in all MAGs from the 204 model communities, regardless of genome

completeness, as partial genomes still provide informative evidence of biosynthetic potential.

Model communities with more species have cumulatively all the necessary steps for the

biosynthesis of essential metabolites (**Figure 5**).

Figure 4. Relationship between genome size, biosynthesis pathway completeness, and growth strategy.
(A) Boxplot comparing the estimated genome size of species-cluster representative genomes (n = 72)
growing exclusively in groups (mixed-only) and capable of growing alone (singleton-capable). (B)
Boxplot comparing the average pathway completeness for amino acid and vitamin biosynthesis
modules between species-cluster representative genomes (n = 57, completeness >90%) growing always
in groups and those capable of growing alone (Wilcoxon rank-sum test). (C) Boxplot displaying the
average pathway completeness for amino acid biosynthesis and (D) vitamin biosynthesis across
genomes (n = 305, completeness >90%) in cultures containing different numbers of species. Letters
indicate groups with significant differences (p < 0.05) (Kruskal-Wallis test followed by Dunn's post-
hoc test with Bonferroni correction). (E) Heatmap showing the average biosynthetic module
completeness for amino acid and B vitamin modules across all 204 microbial model communities.
Values reflect the average presence of module steps (genes) across all genomes per culture (n = 527),
regardless of genome completeness. Modules are grouped by compound type, and microbial model
communities are ordered by richness. Gradient scale ranges from white (0 = incomplete) to black (1 =
complete).

Figure 5. Stepwise module framework analysis of amino acid and B vitamin biosynthetic pathways
across microbial model communities. Heatmap shows the distribution of individual biosynthetic
module steps of all genomes in the 204 microbial model communities. Each column represents a
specific KEGG module step (e.g., M000020_01) grouped by compound and module type. Rows
correspond to individual model communities, organized by the number of species they contain (one,
two, three, or more than three). Values indicate the proportion of genomes in each culture that encode
a specific module step, ranging from 0 (white) to 1 (black).

Ubiquitous and highly relatively abundant species tend to have smaller estimated genome sizes and more auxotrophies.

Overall, we observed a negative correlation between amino acid and vitamin average pathway completeness and average relative abundance for species growing in groups as well as for species growing alone (**Figure 6A** and **Figure S9**). When considering vitamin biosynthesis alone, we found a significant positive correlation between estimated genome size and average pathway completeness for those species growing exclusively in groups (**Figure 6B**). Our results suggest that species found exclusively in groups have higher average relative abundances and smaller genomes with reduced biosynthetic capacity, particularly for vitamins. Scaling the observations from model communities to all species from the BalticMAG catalog, we found that higher average relative abundances (**Figure 6C**) and smaller estimated genome sizes (**Figure 6D**) were linked to lower average completeness of biosynthetic pathways for amino acids and vitamins. Overall, prokaryotes with smaller estimated genome sizes, are more prevalent and abundant in the Baltic Sea and possess lower average biosynthetic potential for amino acids and vitamins (**Figure 6** and **Figure S10**). Specifically, we found that biosynthetic pathways for cysteine, proline, arginine, tryptophan, histidine, leucine, tyrosine, and phenylalanine, vitamin B_1, B_7, and B_{12} were more frequently absent in many species (**Figure S11**).

Figure 6. Link between pathway completeness, relative abundance, and estimated genome size of cultured species and the BalticMAG species catalog. (A) Correlation between average biosynthesis pathway completeness of amino acids and vitamin modules and their relative abundance of cultured species (n = 57, completeness >90%). (B) Correlation between the average biosynthesis completeness for vitamin modules and the estimated genome size of cultured species-clusters. Each data point

represents one species cluster (n = 57, completeness >90%), which either grew exclusively on their own
(light orange), exclusively in groups (green), or both on their own as well as in groups (orange). Panels
(C) and (D) replicate the analysis from panels (A) and (B) but include all species-clusters (n = 450,
completeness >90%) from both cultures and environmental samples. Correlation between estimated
genome size and average relative abundance and (F) correlation between estimated genome size and
prevalence of all species-clusters (n = 701). Data points are color-coded to represent species clusters
from cultures only (red), environments only (blue), and found in both (purple).
369

DISCUSSION

The field of microbiology has traditionally relied on the isolation of microorganisms from

nature in pure culture[49–54]. The practice has yielded foundational insights into microbial

physiology[55–57], metabolism[58–61], and ecological interactions such as mutualism[30,62,63], and

competition[64–66]. However, this reductionistic approach has limitations, particularly a strong

cultivation bias favoring microorganisms with larger genome sizes[43,67], with broader

biosynthetic potential[68], and greater independence. As a result, a substantial portion of the most

abundant microbes in natural environments remains uncultivated[69].

Here, we demonstrate the utility of microbial model communities established through dilution

cultivation from a Baltic Sea pelagic sample for obtaining a wide range of abundant and

biosynthetically limited microorganisms. Despite not extensively varying physicochemical

parameters, our method has provided principles that allow for the cultivation of groups of

organisms with the potential to showcase important ecological principles governing

microorganisms in nature. Our results suggest that by using high-throughput cultivation of

microbial model communities, relevant ecological principles can be better examined and that

by varying cultivation physicochemical parameters, such as adding catalase[70], changing light

regimes[71], or temperature[72], we will be able to cultivate a greater proportion of the abundant

microorganisms found in nature.

In particular, we found that species growing in microbial model communities composed of

more than three species exhibited lower biosynthetic potential for both amino acids and B

vitamins compared to species found growing in smaller groups or independently. The threshold

for species growing together (n > 3) indicates that at higher community complexities,

microorganisms with auxotrophies can form metabolic networks to support their nutritional

requirements[73]. Moreover, a previous study has also found a threshold of microbial diversity at

which competition and complementation saturate[32]. Specifically, they observed that beyond

approximately 26 taxa, further increases in diversity had little detectable impact on an overall

community function (respiration), suggesting a saturation of functional capacity. Although our

system looks at different types of functions, both studies observe a diversity threshold for

functional saturation, emphasizing that microbial communities likely achieve a balanced state

of interaction complexity beyond certain diversity levels, thereby optimizing ecological

efficiency.

Our findings also align with the Black Queen Hypothesis, which posits that certain functions,

particularly costly biosynthetic pathways, can be lost by some community members as long as

the production of the metabolite at the community level is sufficient to sustain the individual[74].

In this context, the reduced genome sizes observed in the abundant species from the model

communities and the environment likely reflect a genome streamlining process[75]. The findings

further align with the hypothesis that obligate co-existing microbes have evolved to rely on

their community for essential nutrients[76], potentially creating social networks that might bring

microbial community stability[11,77–79]. To support this, our stepwise module framework analysis

revealed that microbial model communities with more than three species collectively maintain

the biosynthetic potential for amino acids and vitamins.

Importantly, our findings also complement large-scale studies investigating the ecological

distribution of auxotrophies. A recent analysis of over 26000 representative bacterial genomes

across diverse environments found that auxotrophy is more common in host-associated

environments while relatively rare in aquatic and soil ecosystems. However, their analysis

draws a robust but strict boundary on how to categorize microorganisms in a binary model of

prototrophs and auxotrophs (more than 40% genes missing per pathway for auxotrophy)[13].

While this categorization has been widely used, it might also be hiding potential

interdependencies. In our study, we draw no boundary for defining auxotrophy. Instead, we

studied the pathways by observing their completeness and compared them across the different

species in the dataset. We believe this has the potential to reveal a wide spectrum of possibilities

where microorganisms might just need a precursor to complete biosynthesis[16].

Interestingly, we observe that in the Baltic Sea, a brackish environment where salinity is a

driver for diversity, ecology, and evolution of microorganisms[80], the most abundant and

prevalent taxa possess streamlined genomes with low biosynthetic pathway completeness, as

previously observed in another study from the Baltic Sea[81] . This pattern has also been reported

in freshwater[14] and marine[11] ecosystems, reinforcing the idea that streamlined, auxotrophic

microorganisms are widespread and ecologically successful when they reside in communities

capable of fulfilling their metabolic needs.

In summary, our study highlights the larger potential of microbial model communities in

bridging the gap between laboratory cultivation and ecological representation. By allowing

naturally co-occurring microorganisms to grow together as communities, we can capture some

of the most abundant taxa with the potential for biosynthetic interdependence. These findings

provide evidence that increasing community complexity is associated with reduced metabolic

autonomy, suggesting that microbial interdependencies are common and can emerge as a

natural consequence of a shared metabolic lifestyle. By cultivating groups rather than

individuals, we can access a more ecologically relevant understanding of how microbes

survive, interact, and evolve in nature.

MATERIALS AND METHODS

Sampling and sample processing

We collected an environmental sample from the surface layer of the Baltic Sea close to Askö

in the Trosa archipelago (Lat 58°48.20'N, Lon 17°37.42'E) in October 2021 (**Figure 1A**). The

sample was subsequently processed in the laboratory for DNA extraction and metagenomic

sequencing. Briefly, the water sample was filtered through a 0.1 µm membrane and used to

extract environmental DNA using either the FastDNA® SPIN Kit for Soil (MP Bio) and the

Dneasy PowerWater kit (Qiagen). Additionally, we filtered water through a 0.1 µm hollow fiber

cartridge (Cytiva) and used the filtrate as media to establish cultures with different starting

inoculum sizes.

Establishment of microbial model communities.

We used the dilution-to-extinction technique, as previously described[76], to achieve

approximately the desired starting number of cells in our cultures. The cultures were designed

in two ways: low inoculum size (LS) and high inoculum size (HS). The LS cultures included

hundreds of individual cultures in 96-well plates with a starting number of cells of 0 (control),

2, 6, 10, 15, 20, 30, 50, and 100 cells/well. The HS cultures were prepared in bottles with

starting numbers of cells of 0 (control), 200, 600, 1000, 1500, 2000, 3000, 5000, 10000,

100000, and 1000000 cells/bottle. All cultures were incubated with light/dark cycles (light

6:42, dark 18:32 each day) and 12.2 °C at light and 11.8 °C at dark for four weeks before further

processing. This regime was aimed to emulate the time of dark and light and temperature that

the cells experienced in nature. However, after the experiment had already run and samples

were sequenced, we learned that our light intensity was about five times higher than in nature.

Flow cytometry.

We used a CytoFLEX instrument manufactured by Beckman Coulter to process the

environmental sample and to calculate the cell count as the number of events/mL. Briefly, we

stained 50 µl of the samples with Syto13 at a final concentration of 0.025 mM. During the flow

cytometer acquisition process, we set the following parameters: FSC 2500, SSC 2500, and

FITC 800 with a flow rate of 60 µl/min.

MDA, library preparation, and sequencing.

Before sending our samples for metagenomic sequencing, we performed multiple displacement

amplification (MDA) on all cultures to increase the concentration of DNA[83]. The MDA

reaction consisted of 0.6 µL of culture and 4.4 µL of reaction mix using the Repli-g Single Cell

kit (Qiagen). After DNA amplification, we found that 315 of the cultures passed the

amplification threshold of the negative controls set in the MDA reaction. These cultures were

deemed positive and were sent for sequencing. The selected cultures and our environmental

sample extracted with two different DNA extraction methods were subjected to library

preparation using TruSeq PCRfree DNA library preparation kit (Illumina Inc.) followed by

metagenomic Illumina sequencing at the SNP&SEQ Platform at Uppsala University. This

sequencing technology utilized cluster generation and 150 cycles of paired-end sequencing on

an SP flow cell, employing the NovaSeq 6000 system with v1.5 sequencing chemistry

(Illumina Inc).

Genome resolved metagenomics pipeline.

We removed low-quality reads from the raw sequences using the software Trimmomatic

(v0.36)[84]   with   the   following   options: ILLUMINACLIP:TruSeq3-PE-

2.fa:2:30:10:2:keepBothReads   LEADING:3   TRAILING:3   SLIDINGWINDOW:4:15

MINLEN:50. We employed the MetaWRAP pipeline (v1.3.2)[85] to process our clean

metagenomic reads. First, clean reads were assembled in a single-sample assembly style using

the "metaWRAP_assembly" module with MegaHit (v1.1.3) for the environmental samples[86]

and with SPAdes (v3.15.3)[87] for the culture samples. The quality of the assemblies was assessed

with QUAST (v.5.0.2)[88]. Since the culture DNA was amplified using MDA, the read
information for coverage could not be used. For this reason, we included background
metagenomic data from previous projects in the Baltic Sea (**Figure 1A** and **Table S2**) for
binning. Subsequently, we used multiple-sample coverage binning to decrease the
contamination and increase the completeness of bins[89]. The reads were mapped against all
assemblies using the Input_POGENOM pipeline[90], which uses Bowtie2[91] with default
parameters. After mapping and obtaining the BAM files, the min coverage was calculated using
samtools (v1.9)[92]. Input_POGENOM estimates the coverage of a genome in a sample by only
mapping a subset of the reads, which includes the ones that reached a mean coverage of 20 and
mean breadth of 40. These coverage values for each sample were combined and processed with
the "metaWRAP_bining" module, which uses three metagenomic binning tools: metaBAT2[93],
maxBIN2[94], and CONCOCT[95]. We consolidated all the bins generated by these different tools
using the "metaWRAP_bin_refinement" module. Finally, we first classified the resulting bins
taxonomically with GTDB-tk v2.1.1[46]. Subsequently, we assessed the quality of the bins using
CheckM (v1.1.3)[96]. We considered bins as MAGs when they had a completeness of >45% and
a contamination of <10%, and these MAGs were included for further analysis.

Complementing with previously published MAGs.
The de-replicated MAGs obtained here were supplemented with 771 MAGs from an earlier
study[39] that were based on metagenomics data from three studies[39–41]. We de-replicated the
collection of MAGs to obtain species-cluster representatives using ANI > 95% with mOTUpan
(v0.3.2)[97], and selected the genome with the highest quality as the species-cluster
representative genome.

Relative abundance analysis.
To calculate the relative abundance of our 701 species-clusters in the BalticMAG catalog, we
employed the mapping tool Strobealign (v0.14.0)[98] , which aligns the short metagenomic reads
to our species-clusters collection using a high-speed indexing method (**Table S5**). Briefly, we
created three different indexes with different lengths (100, 125, and 150) for our 701 species-
clusters. We filtered out low-quality reads from our 112 environmental samples with
Trimmonatic (v0.36)[84] with the following options: ILLUMINACLIP:TruSeq3-PE-
2.fa:2:30:10:2:keepBothReads   LEADING:3   TRAILING:3   SLIDINGWINDOW:4:15
MINLEN:50. After this, we did a competitive mapping of all reads against our created index
to obtain the corresponding BAM file for each sample. We sorted the BAM files with the

Anvi'o platform (v7.1)[99] using the anvi-init-bam function, and we calculated the coverage of
each genome per sample with the function anvi-profile-blitz. We calculated each species-
cluster's relative abundance with the previously obtained output by dividing each genome's
mean coverage inner quartiles (i.e., q2q3_cov) by the overall sample mean coverage.
Moreover, we measured the "prevalence", defined as the frequency of each species-cluster
across samples. Specifically, it represents the proportion of samples in which a species-cluster
was detected with a relative abundance > 0.

Functional annotation of KEGG pathways (KOs).
We used the Anvi'o platform (v7.1)[100] to perform functional annotation of KEGG pathways.
Initially, we created a contigs database for each genome, which served as the basis for the
subsequent functional annotation steps. To annotate each genome, we ran the "anvi-run-kegg-
kofams" program, which searches for functions and metabolic pathways based on the KEGG
orthologs (KOs) database[101]. We then predicted the metabolic capabilities of each genome by
running the "anvi-estimate-metabolism" program[102]. We evaluated the completeness of various
metabolic modules in each genome, focusing on the biosynthetic modules for amino acids and
B vitamins (**Table S6**).

Statistical analyses.
All statistical analyses from this study were performed with R version (v4.4.0)[103] and
RStudio[104]. We used the Shapiro test to assess the normality of our data to be compared; if $p <$
0.05, we interpret this as not normally distributed[105]. Since our data was not normally
distributed, we employed a non-parametric test, such as the Wilcoxon test, to find differences
between pairs of groups (e.g., culture vs environment)[106]. We also performed a Kruskal-Wallis
test to compare different groups, and the p-values were corrected using the Benjamin-Hochberg
procedure[107]. Kruskal-Wallis tests were followed by Dunn's post hoc test for pairwise
comparisons[108], and significant differences are indicated by the different letters when $p < 0.05$.
All comparisons were reported to be significant if the p-values were < 0.05.

Data availability. The paired-end sequences of both environmental (n =2) and culture (n =
204) metagenomic samples from the Baltic Sea samples from this study, along with the
corresponding 827 MAGs (>45% completeness and <10% contamination), have been
deposited in NCBI under the BioProject ID PRJNA1134408. These data are not yet publicly
available but can be accessed during peer review via the following temporary link:

https://dataview.ncbi.nlm.nih.gov/object/PRJNA1134408?reviewer=neftpak1i4ufv6i9i7qj3o6
97e. All other metagenomes from the Baltic Sea (n = 110) were downloaded from public
repositories, and the metadata is included in **Table S2**, including their publication reference.

ACKNOWLEDGEMENTS

This work was funded by SciLifeLab and by the Swedish Research Council VR (grant 2022-
03077). The authors would like to acknowledge support from the Genomics infrastructure
services at Science for Life Laboratory in Uppsala. The MDA was done by Claudia Bergin at
the Microbial Single Cell facility, and sequencing was performed by the SNP&SEQ
Technology Platform in Uppsala. The facilities are part of the National Genomics Infrastructure
(NGI) Sweden and Science for Life Laboratory. The SNP&SEQ Platform is also supported by
the Swedish Research Council and the Knut and Alice Wallenberg Foundation. The authors
acknowledge support from SNIC/Uppsala Multidisciplinary Center for Advanced
Computational Science for access to the UPPMAX computational infrastructure as well as the
National Academic Infrastructure for Supercomputing in Sweden (NAISS). Computational
work and data handling were enabled by resources in the projects SNIC 2022/5-392, 2023/5-
126, and naiss2023-5-379 provided by the Swedish National Infrastructure for Computing
(SNIC) at UPPMAX, partially funded by the Swedish Research Council through a grant
agreement no. 2018-05973 and 2022-06725.

SLG conceptualized, designed, and supervised the research. FB and SLG performed the
cultivation work. AFA and LFD-Z provided the Baltic Sea MAGs that contributed to the
genomic catalog. AP-V, JED, LFD-Z, and SLG conducted bioinformatics analysis. AR-G
supported bioinformatic analysis. AP-V, AT, and SLG performed the data interpretation and
visualization. AP-V led the writing of the manuscript with input from SLG. All co-authors
contributed to the literature searches, participated in editing and reviewing the manuscript, and
approved the final version.

CONFLICTS OF INTERESTS

The authors declare no competing interests.

REFERENCES

1. Konopka, A. What is microbial community ecology? *The ISME Journal* **3**, 1223–1230 (2009).

2. Falkowski, P. G., Fenchel, T. & Delong, E. F. The Microbial Engines That Drive Earth's Biogeochemical Cycles. *Science* **320**, 1034–1039 (2008).

3. Gralka, M. Searching for Principles of Microbial Ecology Across Levels of Biological Organization. *Integrative And Comparative Biology* **63**, 1520–1531 (2023).

4. Herman, M. A. *et al.* A Unifying Framework for Understanding Biological Structures and Functions Across Levels of Biological Organization. *Integrative and Comparative Biology* **61**, 2038–2047 (2022).

5. O'Toole, G. A. We have a community problem. *J Bacteriol* e00073-24 (2024) doi:10.1128/jb.00073-24.

6. Benoit, G. *et al.* High-quality metagenome assembly from long accurate reads with metaMDBG. *Nat Biotechnol* **42**, 1378–1383 (2024).

7. Eren, A. M. & Banfield, J. F. Modern microbiology: Embracing complexity through integration across scales. *Cell* **187**, 5151–5170 (2024).

8. Méheust, R., Castelle, C. J., Jaffe, A. L. & Banfield, J. F. Conserved and lineage-specific hypothetical proteins may have played a central role in the rise and diversification of major archaeal groups. *BMC Biol* **20**, 154 (2022).

9. Hug, L. A. *et al.* A new view of the tree of life. *Nat Microbiol* **1**, 16048 (2016).

10. Anantharaman, K. *et al.* Thousands of microbial genomes shed light on interconnected biogeochemical processes in an aquifer system. *Nat Commun* **7**, 13219 (2016).

11. Giordano, N. *et al.* Genome-scale community modelling reveals conserved metabolic cross-feedings in epipelagic bacterioplankton communities. *Nat Commun* **15**, 2721 (2024).

12. Hessler, T. *et al.* Vitamin interdependencies predicted by metagenomics-informed network analyses and validated in microbial community microcosms. *Nat Commun* **14**, (2023).

13. Ramoneda, J., Jensen, T. B. N., Price, M. N., Casamayor, E. O. & Fierer, N. Taxonomic and environmental distribution of bacterial amino acid auxotrophies. *Nat Commun* **14**, (2023).

634   14.   Rodríguez-Gijón, A. *et al.* The ecological success of freshwater microorganisms is
635   mediated by streamlining and biotic interactions. Preprint at
636   https://doi.org/10.1101/2025.03.24.644981 (2025).

637   15.   Akashi, H. & Gojobori, T. Metabolic efficiency and amino acid composition in the
638   proteomes of Escherichia coli and Bacillus subtilis. *Proc. Natl. Acad. Sci. U.S.A.* **99**,
639   3695–3700 (2002).

640   16.   Garcia, S. L. *et al.* Auxotrophy and intrapopulation complementary in the 'interactome'
641   of a cultivated freshwater model community. *Mol Ecol* **24**, 4449–4459 (2015).

642   17.   Mee, M. T., Collins, J. J., Church, G. M. & Wang, H. H. Syntrophic exchange in
643   synthetic microbial communities. *Proc. Natl. Acad. Sci. U.S.A.* **111**, (2014).

644   18.   Paerl, R. W. *et al.* Prevalent reliance of bacterioplankton on exogenous vitamin B1 and
645   precursor availability. *Proc. Natl. Acad. Sci. U.S.A.* **115**, (2018).

646   19.   Ryback, B., Bortfeld-Miller, M. & Vorholt, J. A. Metabolic adaptation to vitamin
647   auxotrophy by leaf-associated bacteria. *The ISME Journal* **16**, 2712–2724 (2022).

648   20.   Sirithanakorn, C. & Cronan, J. E. Biotin, a universal and essential cofactor: synthesis,
649   ligation and regulation. *FEMS Microbiology Reviews* fuab003 (2021)
650   doi:10.1093/femsre/fuab003.

651   21.   Embree, M., Liu, J. K., Al-Bassam, M. M. & Zengler, K. Networks of energetic and
652   metabolic interactions define dynamics in microbial communities. *Proc. Natl. Acad. Sci.*
653   *U.S.A.* **112**, 15450–15455 (2015).

654   22.   Gómez-Consarnau, L. *et al.* Mosaic patterns of B-vitamin synthesis and utilization in a
655   natural marine microbial community. *Environmental Microbiology* **20**, 2809–2823
656   (2018).

657   23.   Wienhausen, G. *et al.* Ligand cross-feeding resolves bacterial vitamin B12 auxotrophies.
658   *Nature* (2024) doi:10.1038/s41586-024-07396-y.

659   24.   Oña, L. & Kost, C. Cooperation increases robustness to ecological disturbance in
660   microbial cross-feeding networks. *Ecology Letters* **25**, 1410–1420 (2022).

661   25.   Aziz, F. A. A. *et al.* Interspecies interactions are an integral determinant of microbial
662   community dynamics. *Front. Microbiol.* **6**, (2015).

663   26.   Goldford, J. E. *et al.* Emergent simplicity in microbial community assembly. (2018).

664   27.   Pacheco, A. R., Osborne, M. L. & Segrè, D. Non-additive microbial community
665   responses to environmental complexity. *Nat Commun* **12**, 2365 (2021).

666   28.   Bayer, B. *et al.* Metabolite release by nitrifiers facilitates metabolic interactions in the
667   ocean. *The ISME Journal* wrae172 (2024) doi:10.1093/ismejo/wrae172.

29. Grant, M. A. A., Kazamia, E., Cicuta, P. & Smith, A. G. Direct exchange of vitamin B12
is demonstrated by modelling the growth dynamics of algal–bacterial cocultures. *The*
ISME Journal **8**, 1418–1427 (2014).

30. Hillesland, K. L. & Stahl, D. A. Rapid evolution of stability and productivity at the
origin of a microbial mutualism. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 2124–2129 (2010).

31. Chang, C.-Y., Bajić, D., Vila, J. C. C., Estrela, S. & Sanchez, A. Emergent coexistence
in multispecies microbial communities. *Science* **381**, 343–348 (2023).

32. Yu, X., Polz, M. F. & Alm, E. J. Interactions in self-assembled microbial communities
saturate with diversity. *The ISME Journal* **13**, 1602–1617 (2019).

33. Hammarlund, S. P., Gedeon, T., Carlson, R. P. & Harcombe, W. R. Limitation by a
shared mutualist promotes coexistence of multiple competing partners. *Nat Commun* **12**,
619 (2021).

34. Pande, S. *et al.* Fitness and stability of obligate cross-feeding interactions that emerge
upon gene loss in bacteria. *ISME J* **8**, 953–962 (2014).

35. Garcia, S. L. *et al.* Model Communities Hint at Promiscuous Metabolic Linkages
between Ubiquitous Free-Living Freshwater Bacteria. *mSphere* **3**, e00202-18 (2018).

36. Ratzke, C., Barrere, J. & Gore, J. Strength of species interactions determines
biodiversity and stability in microbial communities. *Nat Ecol Evol* **4**, 376–383 (2020).

37. Steen, A. D. *et al.* High proportions of bacteria and archaea across most biomes remain
uncultured. *ISME J* **13**, 3126–3130 (2019).

38. Garcia, S. Mixed cultures as model communities: hunting for ubiquitous
microorganisms, their partners, and interactions. *Aquat. Microb. Ecol.* **77**, 79–85 (2016).

39. Alneberg, J. *et al.* Ecosystem-wide metagenomic binning enables prediction of
ecological niches from genomes. *Commun Biol* **3**, (2020).

40. Alneberg, J. *et al.* BARM and BalticMicrobeDB, a reference metagenome and interface
to meta-omic data for the Baltic Sea. *Sci Data* **5**, (2018).

41. Larsson, J. *et al.* Picocyanobacteria containing a novel pigment gene cluster dominate
the brackish water Baltic Sea. *The ISME Journal* **8**, 1892–1903 (2014).

42. Pacheco-Valenciana, A., Garcia, S. L., Dharamshi, J. E., Delgado-Zambrano, L. F. &
Andersson, A. F. The BalticMAG catalog.
https://doi.org/10.17044/scilifelab.28746086.v1 (2025).

43. Rodríguez-Gijón, A. *et al.* A Genomic Perspective Across Earth's Microbiomes Reveals
That Genome Size in Archaea and Bacteria Is Linked to Ecosystem Type and Trophic
Strategy. *Front. Microbiol.* **12**, 761869 (2022).

44. Herlemann, D. P. *et al.* Transitions in bacterial communities along the 2000 km salinity
gradient of the Baltic Sea. *ISME J* **5**, 1571–1579 (2011).

45. Rodríguez-Gijón, A. *et al.* Linking prokaryotic genome size variation to metabolic
potential and environment. *ISME COMMUN.* **3**, 25 (2023).

46. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927
(2020).

47. Giovannoni, S. J. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. *Annu.*
Rev. Mar. Sci. **9**, 231–255 (2017).

48. Herlemann, D. P. R., Woelk, J., Labrenz, M. & Jürgens, K. Diversity and abundance of
"Pelagibacterales" (SAR11) in the Baltic Sea salinity gradient. *Systematic and Applied*
Microbiology **37**, 601–604 (2014).

49. Gich, F., Schubert, K., Bruns, A., Hoffelner, H. & Overmann, J. Specific Detection,
Isolation, and Characterization of Selected, Previously Uncultured Members of the
Freshwater Bacterioplankton Community. *Appl Environ Microbiol* **71**, 5908–5919
(2005).

50. Janssen, P. H., Yates, P. S., Grinton, B. E., Taylor, P. M. & Sait, M. Improved
Culturability of Soil Bacteria and Isolation in Pure Culture of Novel Members of the
Divisions *Acidobacteria* , *Actinobacteria* , *Proteobacteria* , and *Verrucomicrobia*. *Appl*
Environ Microbiol **68**, 2391–2396 (2002).

51. Kaeberlein, T., Lewis, K. & Epstein, S. S. Isolating 'Uncultivable' Microorganisms in
Pure Culture in a Simulated Natural Environment. *Science* **296**, 1127–1129 (2002).

52. Könneke, M. *et al.* Isolation of an autotrophic ammonia-oxidizing marine archaeon.
Nature **437**, 543–546 (2005).

53. Rappé, M. S., Connon, S. A., Vergin, K. L. & Giovannoni, S. J. Cultivation of the
ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**, 630–633 (2002).

54. Schut, F. *et al.* Isolation of Typical Marine Bacteria by Dilution Culture: Growth,
Maintenance, and Characteristics of Isolates under Laboratory Conditions. *Appl Environ*
Microbiol **59**, 2150–2160 (1993).

55. Mella-Flores, D. *et al.* Prochlorococcus and Synechococcus have Evolved Different
Adaptive Mechanisms to Cope with Light and UV Stress. *Front. Microbio.* **3**, (2012).

56. Orr, P. T. & Jones, G. J. Relationship between microcystin production and cell division
rates in nitrogen-limited Microcystis aeruginosa cultures. *Limnology & Oceanography*
43, 1604–1614 (1998).

57. Sohm, J. A., Edwards, B. R., Wilson, B. G. & Webb, E. A. Constitutive Extracellular
Polysaccharide (EPS) Production by Specific Isolates of Crocosphaera watsonii. *Front.*
Microbio. **2**, (2011).

58. Fischer, F., Zillig, W., Stetter, K. O. & Schreiber, G. Chemolithoautotrophic metabolism
of anaerobic extremely thermophilic archaebacteria. *Nature* **301**, 511–513 (1983).

59. Leschine, S. B., Holwell, K. & Canale-Parola, E. Nitrogen Fixation by Anaerobic
Cellulolytic Bacteria. *Science* **242**, 1157–1159 (1988).

60. Zehnder, A. J. & Brock, T. D. Methane formation and methane oxidation by
methanogenic bacteria. *J Bacteriol* **137**, 420–432 (1979).

61. Zhang, T., Shi, X.-C., Ding, R., Xu, K. & Tremblay, P.-L. The hidden
chemolithoautotrophic metabolism of *Geobacter sulfurreducens* uncovered by
adaptation to formate. *The ISME Journal* **14**, 2078–2089 (2020).

62. Belzer, C. *et al.* Microbial Metabolic Networks at the Mucus Layer Lead to Diet-
Independent Butyrate and Vitamin B_{12} Production by Intestinal Symbionts. *mBio* **8**,
e00770-17 (2017).

63. Kazamia, E. *et al.* Mutualistic interactions between vitamin B_{12} -dependent algae and
heterotrophic bacteria exhibit regulation. *Environmental Microbiology* **14**, 1466–1476
(2012).

64. Chodkowski, J. L. & Shade, A. Bioactive exometabolites drive maintenance competition
in simple bacterial communities. *mSystems* e00064-24 (2024)
doi:10.1128/msystems.00064-24.

65. Harrison, F., Paul, J., Massey, R. C. & Buckling, A. Interspecific competition and
siderophore-mediated cooperation in *Pseudomonas aeruginosa*. *The ISME Journal* **2**,
49–55 (2008).

66. Rao, D., Webb, J. S. & Kjelleberg, S. Competitive Interactions in Mixed-Species
Biofilms Containing the Marine Bacterium *Pseudoalteromonas tunicata*. *Appl Environ*
Microbiol **71**, 1729–1736 (2005).

67. Han, K. *et al.* Extraordinary expansion of a Sorangium cellulosum genome from an
alkaline milieu. *Sci Rep* **3**, 2101 (2013).

68. Bentkowski, P., Van Oosterhout, C. & Mock, T. A Model of Genome Size Evolution for
Prokaryotes in Stable and Fluctuating Environments. *Genome Biol Evol* **7**, 2344–2351
(2015).

69. Lloyd, K. G., Steen, A. D., Ladau, J., Yin, J. & Crosby, L. Phylogenetically Novel
Uncultured Microbial Cells Dominate Earth Microbiomes. *mSystems* **3**, e00055-18
(2018).

70. Kim, S., Kang, I., Seo, J.-H. & Cho, J.-C. Culturing the ubiquitous freshwater
actinobacterial acI lineage by supplying a biochemical 'helper' catalase. *ISME J* **13**,
2252–2263 (2019).

71. Bialevich, V., Zachleder, V. & Bišová, K. The Effect of Variable Light Source and Light
Intensity on the Growth of Three Algal Species. *Cells* **11**, 1293 (2022).

72. Jiang, L. & Morin, P. J. Temperature fluctuation facilitates coexistence of competing
species in experimental microbial communities. *Journal of Animal Ecology* **76**, 660–668
(2007).

73. Johnson, W. M. *et al.* Auxotrophic interactions: a stabilizing attribute of aquatic
microbial communities? *FEMS Microbiology Ecology* **96**, fiaa115 (2020).

74. Morris, J. J., Lenski, R. E. & Zinser, E. R. The Black Queen Hypothesis: Evolution of
Dependencies through Adaptive Gene Loss. *mBio* **3**, e00036-12 (2012).

75. Giovannoni, S. J., Cameron Thrash, J. & Temperton, B. Implications of streamlining
theory for microbial ecology. *ISME J* **8**, 1553–1565 (2014).

76. Sokolovskaya, O. M., Shelton, A. N. & Taga, M. E. Sharing vitamins: Cobamides unveil
microbial interactions. *Science* **369**, (2020).

77. Kost, C., Patil, K. R., Friedman, J., Garcia, S. L. & Ralser, M. Metabolic exchanges are
ubiquitous in natural microbial communities. *Nat Microbiol* **8**, 2244–2252 (2023).

78. Zelezniak, A. *et al.* Metabolic dependencies drive species co-occurrence in diverse
microbial communities. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 6449–6454 (2015).

79. Zengler, K. & Zaramela, L. S. The social network of microorganisms — how
auxotrophies shape complex communities. *Nat Rev Microbiol* **16**, 383–390 (2018).

80. Jurdzinski, K. T. *et al.* Large-scale phylogenomics of aquatic bacteria reveal molecular
mechanisms for adaptation to salinity. *Sci. Adv.* **9**, eadg2059 (2023).

81. Hugerth, L. W. *et al.* Metagenome-assembled genomes uncover a global brackish
microbiome. *Genome Biol* **16**, 279 (2015).

82. Garcia, S. L., McMahon, K. D., Grossart, H.-P. & Warnecke, F. Successful enrichment
of the ubiquitous freshwater acI Actinobacteria. *Environmental Microbiology Reports* **6**,
21–27 (2014).

83. Blanco, L. *et al.* Highly Efficient DNA Synthesis by the Phage ϕ 29 DNA Polymerase.
Journal of Biological Chemistry **264**, 8935–8940 (1989).

84.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

803   sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

804   85.   Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for

805   genome-resolved metagenomic data analysis. *Microbiome* **6**, 158 (2018).

806   86.   Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-

807   node solution for large and complex metagenomics assembly via succinct *de Bruijn*

808   graph. *Bioinformatics* **31**, 1674–1676 (2015).

809   87.   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications

810   to Single-Cell Sequencing. *Journal of Computational Biology* **19**, 455–477 (2012).

811   88.   Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for

812   genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

813   89.   Mattock, J. & Watson, M. A comparison of single-coverage and multi-coverage

814   metagenomic binning reveals extensive hidden contamination. *Nat Methods* **20**, 1170–

815   1173 (2023).

816   90.   Sjöqvist, C., Delgado, L. F., Alneberg, J. & Andersson, A. F. Ecologically coherent

817   population structure of uncultivated bacterioplankton. *The ISME Journal* **15**, 3034–3049

818   (2021).

819   91.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*

820   *Methods* **9**, 357–359 (2012).

821   92.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

822   2078–2079 (2009).

823   93.   Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient

824   genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

825   94.   Wu, Y.-W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning

826   algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**,

827   605–607 (2016).

828   95.   Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat*

829   *Methods* **11**, 1144–1146 (2014).

830   96.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:

831   assessing the quality of microbial genomes recovered from isolates, single cells, and

832   metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

833   97.   Buck, M., Mehrshad, M. & Bertilsson, S. mOTUpan: a robust Bayesian approach to

834   leverage metagenome-assembled genomes for core-genome estimation. *NAR Genomics*

835   *and Bioinformatics* **4**, lqac060 (2022).

836   98.   Sahlin, K. Strobealign: flexible seed size enables ultra-fast and accurate read alignment.
837   *Genome Biol* **23**, 260 (2022).

838   99.   Eren, A. M. *et al.* Anvi'o: an advanced analysis and visualization platform for 'omics
839   data. *PeerJ* **3**, e1319 (2015).

840   100.   Eren, A. M. *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat*
841   *Microbiol* **6**, 3–6 (2020).

842   101.   Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM
843   and adaptive score threshold. *Bioinformatics* **36**, 2251–2252 (2020).

844   102.   Veseli, I. *et al.* Microbes with higher metabolic independence are enriched in human gut
845   microbiomes under stress. Preprint at https://doi.org/10.7554/eLife.89862.2 (2024).

846   103.   R Core Team. R: A Language and Environment for Statistical Computing. R Foundation
847   for Statistical Computing (2024).

848   104.   RStudio Team. RStudio: Integrated Development for R. (2024).

849   105.   Royston, J. P. An Extension of Shapiro and Wilk's W Test for Normality to Large
850   Samples. *Applied Statistics* **31**, 115 (1982).

851   106.   Bauer, D. F. Constructing Confidence Sets Using Rank Statistics. *Journal of the*
852   *American Statistical Association* **67**, 687–690 (1972).

853   107.   Kruskal, W. H. & Wallis, W. A. Use of Ranks in One-Criterion Variance Analysis.
854   *Journal of the American Statistical Association* **47**, 583–621 (1952).

855   108.   Dunn, O. J. Multiple Comparisons Using Rank Sums. *Technometrics* **6**, 241–252 (1964).

856