Biting time of day in malaria mosquitoes is modulated by nutritional status

Catherine E. Oke[1], Samuel S. C. Rund[3,4], Maxwell G. Machani[5], Abdul Rahim Mohammed Sabtiu[6], Yaw Akuamoah-Boateng[6], Yaw A. Afrane[6], Sarah E. Reece[1,2]

[1]Institute of Ecology and Evolution and [2]Institute of Immunology and Infection Research, School of Biological Sciences, University of Edinburgh, Charlotte Auerbach Road, Edinburgh, UK; [3]Center for Research Computing and [4]Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, 46556, USA; [5]Entomology Department, Centre for Global Health Research, Kenya Medical Research Institute, Kisumu, Kenya; [6]Department of Medical Microbiology, University of Ghana Medical School, College of Health Sciences, University of Ghana, Accra, Ghana.

Corresponding author:

Catherine E. Oke, Email: catherine.oke@ed.ac.uk

Abstract

Background and objectives: Vector-borne disease transmission follows daily rhythms because the transmission of pathogens occur at the time of day vectors forage for blood. Insecticide-treated bed nets significantly reduce malaria transmission by interrupting the host seeking behaviour of *Anopheles* spp. mosquitoes, yet residual transmission is an increasing problem. Biting when humans are unprotected by bed nets is thought to be a driver of residual transmission, but why mosquitoes are shifting their biting rhythms is poorly understood. We test whether food availability, which mediates activity and foraging rhythms across diverse animal taxa, influences the time of day that mosquitoes bite.

Methodology: We varied the amount of blood and sucrose that female *Anopheles gambiae s.l.* received, and used human-mimic traps in a semi-field system to test the hypothesis that low resources cause host seeking to occur at earlier and later times of day.

Results: Nutritional resources determine both the likelihood and time of day that host seeking occurs. Specifically, low-resourced mosquitoes were 2-3 fold more likely to host seek overall, and 5-10 fold more likely to host seek at an earlier time of day than well-resourced mosquitoes, which predominantly sought a host in the second half of the night time.

Conclusions and Implications: By driving plasticity in biting time of day, mosquito nutritional condition is an underappreciated contributor to residual malaria transmission. Understanding the drivers of biting time of day variation, and their impacts on parasite development, is crucial for the future success of vector control tools and controlling malaria transmission.

Lay summary

We reveal that the nutritional status of mosquitoes influences the time of day that they seek a human to bite. Specifically, poorly-fed mosquitoes attempt to bite humans in the early evening when they are unlikely to be protected by bed nets, which could lead to increased malaria transmission.

Background and Objectives

The transmission of vector-borne diseases (VBDs), such as malaria and dengue, relies on the foraging rhythms of insect vectors, which take up pathogens from an infected vertebrate host during a blood meal and transmit them to a new host during a subsequent blood-feeding event [1,2]. For example, *Plasmodium* parasites, the causative agent of malaria, are vectored between humans by members of the *Anopheles* genus of mosquitoes, which are generally nocturnally active and preferentially bite between the hours of 11pm and 4am (*i.e.* the 'classical' time window) [3–5]. Control strategies, such as insecticide-treated bed nets (ITNs) and indoor residual spraying (IRS), exploit rhythmicity in foraging and resting behaviours, respectably, to bring mosquitoes into contact with a lethal insecticide. Bed nets are one of the most effective methods to curb the spread of malaria [6,7], averting 68% of deaths since 2000 [8]. Despite this success, residual transmission occurs, in part due to the evolution of physiological insecticide resistance (e.g. biochemical and morphological modifications) and behavioural resistance in vector populations [9]. In particular, millions of clinical malaria cases are predicted to be occurring annually due mosquitoes altering their biting time of day [10].

There are increasing reports suggesting that *Anopheles* spp. are shifting their biting time of day to earlier in the evening ('early' biting) or later in the morning ('late' biting) when humans are unprotected by bed nets [4,5,11–15]. While shifts in biting time of day may promote transmission simply due to increased access to hosts, vector foraging at non-classical times of day affects multiple parameters of parasite fitness and transmission (often expressed as the basic reproductive number, R_0 [16]) in complex and potentially opposing ways [1]. For example, studies with animal models of malaria reveal that mosquitoes are less susceptible to *Plasmodium* infection at night [17], likely due to rhythmicity in mosquitos' immune defences [18], but parasites are intrinsically more infective to mosquitoes at night, and both mosquito and parasite rhythms interact to influence parasite productivity within mosquitoes and onwards transmission potential [17,19]. There are also increased risks of desiccation and predation to mosquitoes foraging at suboptimal times of day. Further, energetic costs could result from rhythmic metabolic processes being misaligned with food intake [1]. Therefore, shifts in biting time of day are likely to impact on vectorial capacity and the success of vector control tools in hard to intuit, but important, manners [6].

While *Anopheles* spp. biting rhythms are weakly heritable, variation in biting time of day is predominantly driven by non-genetic factors [20]. This suggests that mosquitoes use phenotypic plasticity to adjust their foraging behaviour in response to environmental conditions [21], such as limited access to blood meals. However, the extent to which mosquitos can shift biting time of day and the range of potential environmental drivers are poorly understood. Variation in physiological condition and resource availability can cause animals to alter their foraging rhythms [22] in adaptive (i.e. fitness enhancing) ways, especially in resource-limiting conditions [23]. For example, low food availability promotes a shift to diurnal foraging patterns in small nocturnal rodents, which minimises energy loss by remaining in burrows during particularly cold nights [24,25]. Similarly, mosquitoes may garner greater benefits from shifting biting time of day when they are in poor nutritional condition. Such context-dependent variation in the costs and benefits of shifts in biting time may explain why biting times have seemingly not

significantly changed in some *Anopheles* spp. populations despite high ITN use [26,27], and why the evolution of behavioural resistance varies across populations in manners that correlate with the abundance of sources of nutrition for mosquitoes [28].

The rhythmic feeding ecology of anopheline mosquitoes [13,29,30] is under circadian clock control [31] and revolves around the need for blood and sugar meals. Female mosquitoes utilise proteins and lipids in blood for egg production, and sugar sources power flight and the accumulation of energy reserves [32]. Females take a blood meal after mating, beginning their first gonotrophic cycle of egg development and oviposition, seeking more blood approximately every 3 days to fuel subsequent gonotrophic cycles [33]. The type and abundance of nutritional resources available to mosquitoes varies across habitats and seasons [32,34], and vector control tools such as ITNs can limit blood meal availability [35]. In response, mosquitoes exhibit adaptive plasticity in multiple feeding behaviours; for example, failure to acquire a blood meal leads to increased sugar feeding [30] to promote survival until future biting opportunities. Malaria parasites rely on mosquito host seeking for transmission [36]; they are taken up during a blood meal and become ready to transmit to a new host after approximately 10-14 days, during a subsequent blood meal. However, despite the importance of mosquito foraging rhythms for malaria transmission and evidence that physiological condition can alter foraging rhythms in other taxa, the effect of resource availability on mosquito biting time of day is unknown.

Here, we perturb the nutritional resources provided to adult mosquitoes to test whether they respond by plastically altering the time of day that host seeking occurs. We simulated scenarios in which mosquitoes have taken an initial blood meal (simulating the point at which malaria parasites would be acquired), and then: (i) were unsuccessful at acquiring a subsequent blood meal and only have access to a sugar-poor environment, (ii) were unsuccessful at acquiring a subsequent blood meal but have access to a sugar-rich environment, and (iii) successfully acquired a subsequent blood meal as well as having access to a sugar-rich environment. We reveal that mosquitoes with access to additional blood meals and/or high levels of sugar host seek during the classical night-time biting window and into the early morning (i.e. late biting), but when sugar and blood are limited, mosquitoes are more likely to host seek in the evening (i.e. early biting) when humans are unlikely to be protected by ITNs. We also supplement our data with preliminary field collections that suggest biting time of day of wild-caught adult mosquitoes may correlate with some aspects of nutritional condition. We discuss the potential outcomes of shifts in biting time of day in response to resource availability for understanding residual transmission patterns, and for parasite fitness.

Methodology

We performed a semi-field experiment to assess if mosquito nutritional status impacts host seeking and biting time of day. We reared wild-caught larvae, and maintained blood fed female *Anopheles gambiae s.l.* mosquitoes from the F_2 generation on diets that varied in nutritional resources, before releasing them into an enclosed semi-field system with traps mimicking human odour. The traps were

programmed to separately capture mosquitoes biting at early (evening), classical (during the night) and late (morning) times of day.

Mosquitoes

Experimental mosquitoes were F_2 progeny of wild-caught *Anopheles gambiae s.l.* larvae, collected from three sites across the Greater Accra region of Ghana (Tuba, 5° 30' 47" N 0° 23' 16" W; Teshie, 5° 35′ 0″ N, 0° 6′ 0″ W; East Legon, 5° 38' 16.39" N, 0° 9' 40.33" W) in Dec 2023-Jan 2024 (Supplementary Fig. S1). We maintained all generations of infection-free mosquitoes under standard rearing conditions (26±2°C, 80% relative humidity with a 12L:12D cycle) with blood provided via direct-feeding [37]. We fed both F_0 and F_1 generations *ad libitum* with 10% sucrose solution in water, and provided them with soaked cotton wool in petri dishes for egg laying. On day 1 post-emergence of the F_2 generation, we transferred female mosquitoes to three experimental cages (n=100-150 per cage) and allocated each cage to one of three feeding treatments: (i) 0.5% sucrose in water (0.5% suc), (ii) 10% sucrose in water (10% suc) or (iii) 10% sucrose in water plus an additional blood meal on day 6 post-emergence (10% suc + bm). We maintained all treatment groups on their feeding treatments from day 1 post-emergence until the host seeking assay. In addition, to mimic mosquito life history in nature, we provided all groups with an initial blood meal on day 3 post-emergence and the opportunity to lay eggs. On day 9-10 post-emergence, we transferred mosquitoes from each cage to paper cups (n=70-100 per cup) and provided them with water only for six hours prior to assaying biting time of day. We reared six batches of mosquitoes to conduct a total of six semi-field system releases.

Semi-field host seeking experiment

We conducted mosquito behavioural experiments in January-February 2024 at the University of Ghana in an enclosed semi-field system (the "MalariaSphere", [38,39]). The average (± SEM) minimum and maximum daily temperatures during the experimental releases were 25.8±0.6°C and 34.5±0.4°C respectively [40]. The enclosure of 5.8 x 4.2 x 2.8m was covered in an insect-proof screen to prevent mosquito escape and/or entry from the external environment. We placed three BG-Sentinel traps (Biogents, Germany) in the enclosure, with an MB5 human-mimic lure developed to attract anopheline mosquitoes (Biogents, Germany) and a CO_2 flow of approximately 21g/hour, to mimic human scent and breath. We programmed the traps to turn on at 6pm-10pm to capture 'early' biters, 11pm-3am to capture 'classical' biters and 4am-8am to capture 'late' biters (Fig. 1). CO_2 flow was controlled with BG-CO_2 timers (Biogents, Germany), ensuring CO_2 was only emitted when the respective trap was switched on.

Within the MalariaSphere, released mosquitoes could access resting sites (four clay pots around the enclosure). We did not provide water or food sources because this would have confounded the experimental treatment groups. We colour-marked mosquitoes from each feeding treatment using aerosolised fluorescent powder (FTX series, Swada London), applied following Machani et al [38]. We rotated the colour of each treatment group across sequential releases to ensure any differential impacts

of colour on mosquito behaviour did not confound treatment groups. We carried out six releases over a period of three weeks, with mosquitoes being released into the enclosure between 5pm-5:30pm, just prior to dusk at 6pm (n=235-300 mosquitoes per release). A period of at least 48 hours between each release and checking all resting sites with a Prokopack aspirator ensured that any remaining untrapped mosquitoes had been removed or died before the next release. We collected mosquitoes from traps at 8am the day after the release, then cold-anaesthetised and counted them.

Figure 1. The semi-field behavioural assay set-up. Schematic diagram of the set-up (A), featuring four resting sites for mosquitoes, and traps baited with human odour which were programmed to turn on at 6pm-10pm, 11pm-3am and 4am-8pm respectively. These windows correspond to approx. ZT12-ZT16, ZT17-21, and ZT22-ZT2 respectively, where Zeitgeber time (ZT) defines the hours since lights on and ZT0/24 is dawn. All traps were linked to a CO_2 source, which was split to three timers ensuring the CO_2 was only flowing when the trap that was switched on. Photographs show the semi-field facility (B) and the behavioural assay set-up (C).

Nutrition assays

Immediately prior to each release, we collected a subset of mosquitoes from each feeding treatment group (n=14-19 per group, total n=48) to confirm our resource perturbations impacted nutritional status. These mosquitoes were frozen at -20°C for lipid, glycogen and total sugar analyses using modified Van Handel protocols [41–43]. Firstly, we removed a wing from each mosquito to quantify body size. Wing length is a commonly used proxy for adult body size because dry weight and wing length are highly correlated [44]. We then lysed individual mosquitoes in 100μl 2% sodium sulphate and added 750μl of 1:2 chloroform:methanol. We centrifuged samples at 12000 rpm for 3 minutes; the supernatant was

used for lipid and total sugars analysis, and the precipitate was used for glycogen analysis. We conducted lipid, glycogen and total sugars assays following Oke et al [45].

Statistical analysis

We used R v. 4.1.3 to perform data analysis. We confirmed that adult body size did not differ across the treatments using a linear model with wing length, mosquito release batch and their interaction as main effects (Supplementary Fig. S2). We then analysed nutritional content (lipid, glycogen, total sugars) of individual mosquitoes using linear mixed models with feeding treatment as a main effect and mosquito release batch as a random effect, accounting for any within treatment variation in body size. To investigate how feeding treatments impacted the overall proportion of mosquitoes caught across each release session, we used binomial generalised linear mixed models (glmm) with mosquito release batch as a random effect and feeding treatment, trap time and their interaction as main effects. We estimated proportion of caught mosquitoes and relative odds ratios (OR) ± SE from models using the *emmeans* [46] package. We square root transformed nutrition data to meet assumptions of normality and homogeneity of variance. We minimised all models using likelihood ratio tests and AICc for non-nested models, and confirmed model assumptions using the *easystats* [47] package. To account for differences between mosquito release batches, we present estimated marginal means ± SEM (*emmeans* package), predicted from models. Whenever the minimised model contained a significant effect of feeding treatment, we conducted post hoc pairwise comparisons using the Tukey method with the *emmeans* package.

Ethics statement

The study protocols for mosquito colony blood feeds and human landing catches (HLCs) (see Discussion and Supplementary Methods) were ethically reviewed and approved by the Ghana Health Service Ethics Review Committee (GHS-ERC: 021/07/23). Volunteers involved in colony blood feeds were tested for malaria prior to direct-feeding, and the same person fed the mosquitoes throughout the experiment. Permission to conduct HLCs in study sites was obtained from the District Assembly Representative, and written consent was obtained from all collectors. Collectors had access to malaria prophylaxis and did not report any adverse events.

Results

Resource availability perturbs mosquito nutritional status

Mosquitoes fed on the lowest concentration of sucrose (0.5%) had significantly lower lipid levels (55.5±17.8µg) (χ^2_2=17.6, p < 0.001), approximately 2.5-fold lower than those fed with 10% sucrose (138±31.4µg) and 3.3-fold lower those fed an additional blood meal (182±29.6µg) (Fig. 2A, Table 1). A similar pattern was observed for total sugars: mosquitoes fed with 0.5% sucrose had the lowest total

sugar levels (85.6±69.9µg) (χ^2_2=30.8, p < 0.001), approximately 5-fold lower than those given 10% sucrose (420±158µg) and an additional blood meal (439±156µg) (Fig. 2B, Table 1). Finally, mosquitoes fed with 0.5% sucrose had the lowest glycogen levels (105±64.2µg) (χ^2_2=21.7, p < 0.001), approximately 7-fold lower than those given 10% sucrose (754±187µg), with mosquitoes given an additional blood meal exhibiting intermediate levels (481±128µg) (Fig. 2C, Table 1).

Figure 2. Concentrations (µg) per mosquito of lipids (A), total free sugars (B) and glycogen (C) in individual mosquitoes under differing feeding treatments on day 10 post-emergence. Nutritional perturbations began on day 1 post-emergence and all treatment groups received a blood meal on day 3 post-emergence (with an additional blood meal given to the 10% suc + bm group on day 6). Data presented are estimated marginal means ± SEM.

Table 1. Post hoc pairwise comparisons for nutritional content of mosquitoes provided with different feeding treatments. Significant p-values (<0.05) are highlighted in bold, and borderline p-values (0.05<p<0.09) are italicised and underlined.

		Test statistic	p-value
Lipids	0.5% suc – 10% suc	t = 2.75	**0.02**
	0.5% suc – 10% suc + bm	t = -4.54	**<0.001**
	10% suc – 10% suc + bm	t = -1.16	0.48
Total sugars	0.5% suc – 10% suc	t = 5.13	**<0.001**
	0.5% suc – 10% suc + bm	t = -6.13	**<0.001**
	10% suc – 10% suc + bm	t = -0.21	0.98
Glycogen	0.5% suc – 10% suc	t = 4.66	**<0.001**
	0.5% suc – 10% suc + bm	t = -3.69	**0.002**
	10% suc – 10% suc + bm	t = 1.55	0.28

Resource availability affects host seeking tendency and biting time of day

Across all six releases and trapping times of day, 32.2% of mosquitoes (506/1569) were caught overall. The likelihood of host seeking correlated negatively with the level of nutritional resources provided (χ^2_2=65.8, p < 0.001) (Fig. 3A). Specifically, mosquitoes fed 0.5% sucrose were 2-fold more likely to be trapped than those fed 10% sucrose (odds ratio (OR): 2.08±0.28) and 3-fold more likely than those fed an additional blood meal (OR: 2.97±0.42). Mosquitoes fed only 10% sucrose were 1.4-fold more likely to be trapped than those fed an additional blood meal (OR: 1.43±0.21).

We also found that nutritional resources differentially affected the time of day of host seeking (χ^2_4=95.4, p < 0.001). The majority of early biting mosquitoes were from the 0.5% sucrose group, whereas mosquitoes with more resources were more likely to be caught in subsequent traps (Fig. 3B). Specifically, mosquitoes fed 0.5% sucrose were approximately 5- and 10-fold more likely to be caught in the early biting window (20±2.9%) than those fed with 10% sucrose (3.7±0.9%) and an additional blood meal (1.6±0.5%), respectively. Mosquitoes fed 10% sucrose with or without an additional blood meal followed similar temporal patterns, with approximately 4-fold more of these mosquitoes being trapped during the classical (10% sucrose: 11±2.0%, 10% sucrose + bm: 9.0±1.7%) and late biting windows (10% sucrose: 13±2.3%, 10% sucrose + bm: 11±2.0%) than during the early biting window (10% sucrose: 3.7±0.9%, 10% sucrose + bm: 1.6±0.5%).

Figure 3. The proportion of released mosquitoes from each feeding treatment that were trapped across the entire night (A) and that were caught in the evening (early, 6pm-10pm), classical night time biting window (11pm-3am) and morning (late, 4am-8am) biting (B). Data presented are estimated marginal means ± SEM.

Discussion

By altering sugar provision and access to blood meals, and using traps that mimic human breath and scent, we find that resource availability modulates the time of day that mosquitoes search for a blood meal. Mosquitoes on a low sucrose diet had significantly lower levels of lipids, glycogen and sugar, and were 5-10 fold more likely to host seek in the evening (early) than mosquitoes on a high sucrose diet with or without a blood meal (Fig. 2, 3B). In contrast, better-resourced mosquitoes were most likely to host seek during the classical biting time window and into the morning (Fig. 2, 3B). We also find that as resources increased, the overall tendency to host seek decreased; mosquitoes on a low sucrose diet were 2-fold more likely to be caught compared to mosquitoes given a higher concentration of sucrose, and 3-fold more likely than those with access to 10% sucrose and an additional blood meal (Fig. 3A).

Foraging at unusual times of day can be detrimental because it exposes organisms to rhythmic environmental risks, such as the active phase of predators and suboptimal temperature/humidity. Furthermore, circadian rhythms anticipate rhythmic events such as feeding and prepare metabolism accordingly, so foraging at unanticipated times of day may cause suboptimal digestion and nutrition acquisition [1]. However, when resource availability is low, finding food becomes more critical for survival, outweighing the environmental risks and physiological costs [22,25]. For example, a poorly-resourced mosquito may need to become active earlier in the night because they do not have the energy reserves to be able to wait until the classical time of day to forage for blood. In addition, a mosquito that has been unsuccessful at foraging throughout the night may have used significant energy, causing it continue to forage later in the morning because it does not have sufficient reserves to survive until the subsequent night. Our results are consistent with our hypothesis for early biting because poorly-resourced mosquitoes with low nutritional reserves are more likely to host seek, predominantly in the early evening when temperatures are higher and humidity lower, and risk being killed because human hosts are alert. Although we did not quantify survival, we did not encounter any live mosquitoes fed on a 0.5% sucrose diet whilst clearing the semi-field systems the morning after each release, but regularly encountered live mosquitoes fed on higher sucrose solutions and/or an additional blood meal. Overall, this suggests that early biting is an adaptive temporal shift, which may be necessary in regions which are sugar-poor and hosts are not easily accessible due to high ITN use, and/or where larvae are poorly nourished.

That better-resourced mosquitoes on high sucrose diets (with or without a blood meal) had higher nutritional reserves and waited until the classical time window to host seek, when environmental risks are likely to be lower, also supports our hypotheses. However, we also found that better-resourced mosquitoes were just as likely as poorly resourced mosquitoes to be trapped later into the morning. Such late biting by well-fed mosquitoes was unexpected but may be due to the lack of water/food sources provided in the semi-field enclosures, forcing untrapped mosquitoes to host seek to mitigate dehydration caused by the rise in temperature and the reduction in humidity in the morning [48,49]. Alternatively, this finding may be explained by our hypothesis that well-resourced mosquitoes which had failed to blood feed (*i.e.* had not entered a trap) during the night have become resource-depleted,

leading to an increased propensity to continue host seeking at unusual times of day. Regardless of the explanation(s), our results reveal that adult nutritional condition drives variation in biting time of day.

To probe the ecological relevance of our experimental results, we also undertook a trapping session for wild mosquitoes (see Supplementary Information). While the nutritional content of the 66 *Anopheles* spp. mosquitoes we caught between 6pm and 7am varied, levels were not significantly different to the profiles of low-resourced (0.5% sucrose) experimental mosquitoes (Supplementary Fig. S3, Table S1). Our low sample size precludes comparing nutritional reserves of wild mosquitoes caught host seeking at different times of day, but suggests that adult nutritional content does not correlate with body size (a product of larval condition) (Fig. S4). Thus, catching lowly resourced mosquitoes at the time of host seeking is consistent with our experimental results revealing that adult nutrition influences biting rhythms. Furthermore, like previous studies across Africa [3,50,51], we found a large proportion (approximately 20%) of 'late' biting occurring in the morning when humans are unprotected by bed nets. This pattern has become more pronounced since the widespread use of ITNs. Because biting time shifts have a genetic component [20], and universal coverage of ITNs is approximately 75% in Ghana [52], late biting could be an evasion strategy evolving in the mosquito population where the larval collections for our experiment were carried out, rather than a plastic behavioural change. However, further work is needed to ascertain the extent to which biting time of day is genetically determined and/or due to behavioural plasticity in response to variation in resource availability. For example, modelling suggests that early biting is more likely to evolve in sugar-poor environments than sugar-rich environments [28]. However, in areas of fluctuating resource availability or ITN coverage, plastic genotypes that can switch between early, classical, or late biting depending on their environmental conditions should have higher fitness [21].

Both our semi-field experiment and preliminary field data demonstrate significant variation in biting time of day. Parasite fitness is a culmination of the parameters contributing to R_0, many of which are directly related to within-vector parasite development, vector survival and vector behaviour [16]. Intuition suggests that early biting by poorly-resourced mosquitoes could sustain malaria transmission. However, the overall impact of shifts in biting time of day for malaria transmission is difficult to predict, because of potential synergistic and antagonistic interactions between R_0's constituent parameters. For example, while early biting causes greater contact between vectors and hosts in regions with high ITN coverage, low-resourced mosquitoes have reduced survival [45]. Thus, while early biters are more likely to obtain a blood meal, their poor nutritional state could curtail onward transmission if vectors do not survive long enough for malaria parasites to complete their development [53,54]. Furthermore, if early biters are more likely to be low-resourced mosquitoes, there are conflicting outcomes for parasites. While early biting is more suitable for parasite establishment in the mosquito due to lower night-time temperatures [55] and higher infectivity than in the day time [1,17,19], parasite development is slower and less productive in low-resourced mosquitoes [45,56,57]. In addition, disruption to foraging rhythms reduces fitness in other insects [58], suggesting that blood feeding at a non-classical time of day may be suboptimal for mosquitoes. The breakdown of blood meal contents leads to high levels of reactive oxygen species (ROS), and mosquitoes have an anticipatory rhythmic detoxification mechanisms for

coping with oxidative stress [59]. Thus, if blood feeding at an unusual time of day causes detoxification mechanisms and metabolic processes to be mismatched to the time of day blood is digested, mosquitoes may experience higher oxidative stress and/or less productive digestion of nutrients, both of which would reduce parasite fitness. While transmission at a non-classical time of day is clearly better for parasites than no transmission at all in areas of ITN use, recognising that transmission is time of day dependent is necessary for accurately predicting epidemiology, the risk of clinical malaria cases, and the trajectory of parasite evolution.

Conclusions and Implications

Our results suggest that plasticity in biting time of day, driven by variation in adult mosquito nutritional condition, is an underappreciated contributor to residual malaria transmission. Addressing the drivers of variation in biting time of day and the overall impact on malaria transmission is critical for assessing and improving the efficacy of current and future vector control tools harnessing mosquito foraging behaviours. While this is especially important for malaria, where up to 30% of biting is reported to occur when humans are unprotected by bed nets and is predicted to lead to millions of additional clinical cases per year [3,10], we also expect our findings to apply to other vector-borne diseases that rely on vector foraging rhythms for between-host transmission.

Funding

This work was funded by a Varley-Gradwell Travelling Fellowship in Insect Ecology (2023-2024) to C.E.O, and by a Wellcome Trust Investigator Award (202769/Z/16/Z) and Wellcome Trust Discovery Award (227904/Z/23/Z) to S.E.R. Research equipment and travel funds for M.G.M. and S.S.C.R. were funded by an Initiation Grant from the University of Notre Dame, Notre Dame Research office. Y. A. A. is funded by the National Institute for Health (D43 TW 011513). The funders had no role or influence on the design of this study, data collection, analyses, and interpretation of the data collected, as well as in writing this manuscript.

Conflict of interest: We declare no conflict of interest.

Acknowledgements

We thank Anisa Abdulai for experimental and technical assistance, and the serology lab team at the Noguchi Memorial Institute for Medical Research for kindly letting us use their facilities.

References

1. Rund SSC, O'Donnell AJ, Gentile JE *et al.* Daily rhythms in mosquitoes and their consequences for malaria transmission. *Insects* 2016;**7**:1–20.

2. Westwood ML, O'Donnell AJ, de Bekker C *et al.* The evolutionary ecology of circadian rhythms in infection. *Nature Ecology and Evolution* 2019;**3**:552–60.

3. Sangbakembi-ngounou C, Costantini C, Longo-pendy NM *et al.* Diurnal biting of malaria mosquitoes in the Central African Republic indicates residual transmission may be " out of control ." *Proceedings of the National Academy of Sciences* 2022;**119**:e2104282119.

4. Thomsen EK, Koimbu G, Pulford J *et al.* Mosquito behavior change after distribution of bednets results in decreased protection against malaria exposure. *Journal of Infectious Diseases* 2017;**215**:790–7.

5. Moiroux N, Gomez MB, Pennetier C *et al.* Changes in anopheles funestus biting behavior following universal coverage of long-lasting insecticidal nets in benin. *Journal of Infectious Diseases* 2012;**206**:1622–9.

6. Oke CE, Ingham VA, Walling CA *et al.* Vector control: agents of selection on malaria parasites? *Trends in Parasitology* 2022;**38**:890–903.

7. World Health Organisation. *World Malaria Report 2023*. Geneva, 2023.

8. Bhatt S, Weiss DJ, Cameron E *et al.* The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. *Nature* 2015;**526**:207–11.

9. Huijben S, Paaijmans KP. Putting evolution in elimination: Winning our ongoing battle with evolving malaria mosquitoes and parasites. *Evolutionary Applications* 2018;**11**:415–30.

10. Sherrard-Smith E, Skarp JE, Beale AD *et al.* Mosquito feeding behavior and how it influences residual malaria transmission across Africa. *Proceedings of the National Academy of Sciences of the United States of America* 2019;**116**:15086–96.

11. Russell TL, Beebe NW, Cooper RD *et al.* Successful malaria elimination strategies require interventions that target changing vector behaviours. *Malaria Journal* 2013;**12**:1–5.

12. Sougoufara S, Diédhiou SM, Doucouré S *et al.* Biting by Anopheles funestus in broad daylight after use of long-lasting insecticidal nets: A new challenge to malaria elimination. *Malaria Journal* 2014;**13**:1–7.

13. Yohannes M, Boelee E. Early biting rhythm in the afro-tropical vector of malaria, Anopheles arabiensis, and challenges for its control in Ethiopia. *Medical and Veterinary Entomology* 2012;**26**:103–5.

14. Cooke MK, Kahindi SC, Oriango RM *et al.* "A bite before bed": Exposure to malaria vectors outside the times of net use in the highlands of western Kenya. *Malaria Journal* 2015;**14**:1–15.

15. Carrasco D, Lefèvre T, Moiroux N *et al.* Behavioural adaptations of mosquito vectors to insecticide control. *Current Opinion in Insect Science* 2019;**34**:48–54.

16. Smith DL, Battle KE, Hay SI *et al.* Ross, Macdonald, and a theory for the dynamics and control of mosquito-transmitted pathogens. *PLoS Pathogens* 2012;**8**, DOI: 10.1371/journal.ppat.1002588.

17. Schneider P, Rund SSC, Smith NL *et al.* Adaptive periodicity in the infectivity of malaria gametocytes to mosquitoes. *Proceedings of the Royal Society B: Biological Sciences* 2018;**285**, DOI: 10.1098/rspb.2018.1876.

18. Murdock CC, Moller-Jacobs LL, Thomas MB. Complex environmental drivers of immunity and resistance in malaria mosquitoes. *Proceedings of the Royal Society B: Biological Sciences* 2013;**280**, DOI: 10.1098/rspb.2013.2030.

19. Pigeault R, Caudron Q, Nicot A *et al.* Timing malaria transmission with mosquito fluctuations. *Evolution Letters* 2018;**2**:378–89.

20. Govella NJ, Johnson PCD, Killeen GF *et al.* Heritability of biting time behaviours in the major African malaria vector Anopheles arabiensis. *Malaria Journal* 2023;**22**:1–9.

21. Pigliucci M. Evolution of phenotypic plasticity: Where are we going now? *Trends in Ecology and Evolution* 2005;**20**:481–6.

22. Van Der Veen DR, Riede SJ, Heideman PD *et al.* Flexible clock systems: adjusting the temporal programme. *Phil Trans R Soc B* 2017;**372**:20160254.

23. Van Der Vinne V, Riede SJ, Gorter JA *et al.* Cold and hunger induce diurnality in a nocturnal mammal. *Proc Natl Acad Sci USA* 2014;**111**:15256–60.

24. Van Der Vinne V, Gorter JA, Riede SJ *et al.* Diurnality as an energy-saving strategy: energetic consequences of temporal niche switching in small mammals. *Journal of Experimental Biology* 2015;**218**:2585–93.

25. Van Der Vinne V, Tachinardi P, Riede SJ *et al.* Maximising survival by shifting the daily timing of activity. Ghalambor C (ed.). *Ecology Letters* 2019;**22**:2097–102.

26. Sougoufara S, Harry M, Doucouré S *et al.* Shift in species composition in the Anopheles gambiae complex after implementation of long-lasting insecticidal nets in Dielmo, Senegal. *Medical and veterinary entomology* 2016;**30**:365–8.

27. Kreppel KS, Viana M, Main BJ *et al.* Emergence of behavioural avoidance strategies of malaria vectors in areas of high LLIN coverage in Tanzania. *Scientific Reports* 2020;**10**:1–11.

28. Stone C, Chitnis N, Gross K. Environmental influences on mosquito foraging and integrated vector management can delay the evolution of behavioral resistance. *Evolutionary Applications* 2016;**9**:502–17.

29. Yee WL, Foster WA. Diel Sugar-Feeding and Host-Seeking Rhythms in Mosquitoes (Diptera: Culicidae) Under Laboratory Conditions. *Journal of Medical Entomology* 1992;**29**:784–91.

30. Gary RE, Foster WA. Diel timing and frequency of sugar feeding in the mosquito Anopheles gambiae, depending on sex, gonotrophic state and resource availability. *Medical and Veterinary Entomology* 2006;**20**:308–16.

31. Rund SSC, Bonar NA, Champion MM *et al.* Daily rhythms in antennal protein and olfactory sensitivity in the malaria mosquito Anopheles gambiae. *Sci Rep* 2013;**3**:2494.

32. Barredo E, Degennaro M. Not Just from Blood: Mosquito Nutrient Acquisition from Nectar Sources. *Trends in Parasitology* 2020;**36**:473–84.

33. Mitchell SN, Catteruccia F. Anopheline reproductive biology: Impacts on vectorial capacity and potential avenues for malaria control. *Cold Spring Harbor Perspectives in Medicine* 2017;**7**:14.

34. Gu W, Müller G, Schlein Y *et al.* Natural plant sugar sources of Anopheles mosquitoes strongly impact malaria transmission potential. *PLoS ONE* 2011;**6**, DOI: 10.1371/journal.pone.0015996.

35. Killeen GF, Smith TA, Ferguson HM *et al.* Preventing childhood malaria in Africa by protecting adults from mosquitoes with insecticide-treated nets. *PLoS Medicine* 2007;**4**:1246–58.

36. Shaw WR, Marcenac P, Catteruccia F. Plasmodium development in Anopheles: a tale of shared resources. *Trends in Parasitology* 2022;**38**:124–35.

37. Harrington LC, Foy BD, Bangs MJ. Considerations for Human Blood-Feeding and Arthropod Exposure in Vector Biology Research: An Essential Tool for Investigations and Disease Control. *Vector-Borne and Zoonotic Diseases* 2020;**20**:807–16.

38. Machani MG, Ochomo E, Amimo F *et al.* Behavioral responses of pyrethroid resistant and susceptible Anopheles gambiae mosquitoes to insecticide treated bed net. *PLoS ONE* 2022;**17**:e0266420.

39. Osoro JK, Machani MG, Ochomo E *et al.* Insecticide resistant Anopheles gambiae have enhanced longevity but reduced reproductive fitness and a longer first gonotrophic cycle. *Scientific Reports* 2022:1–7.

40. Valer GB. Ogimet. 2024.

41. Van Handel E, Day JF. Assay of lipids, glycogen and sugars in individual mosquitoes: correlations with wing length in field-collected Aedes vexans. *Journal of the American Mosquito Control Association* 1988;**4**:549–50.

42. Van Handel E. Rapid determination of glycogen and sugars in mosquitoes. *Journal of the American Mosquito Control Association* 1985;**1**:299–301.

43. Van Handel E. Rapid determination of total lipids in mosquitoes. *Journal of the American Mosquito Control Association* 1985:302–4.

44. Lehmann T, Dalton R, Kim EH *et al.* Genetic contribution to variation in larval development time, adult size, and longevity of starved adults of Anopheles gambiae. *Infection, Genetics and Evolution* 2006;**6**:410–6.

45. Oke CE, O'Donnell AJ, Schneider P *et al.* Plasticity in malaria parasite development: mosquito resources influence vector-to-host transmission potential. *Front Malar* 2024;**2**:1481816.

46. Lenth RV. emmeans: Estimated Marginal Means, aka Least-Squares Means. 2023.

47. Makowski D, Ben-Shachar M, Lüdecke D. The {easystats} collection of R packages. 2020.

48. Hagan RW, Didion EM, Rosselot AE *et al.* Dehydration prompts increased activity and blood feeding by mosquitoes. *Scientific Reports* 2018;**8**:1–12.

49. Lin S, Senapati B, Tsao CH. Neural basis of hunger-driven behaviour in Drosophila. *Open Biology* 2019;**9**, DOI: 10.1098/rsob.180259.

50. Odero JI, Abong'o B, Moshi V *et al.* Early morning anopheline mosquito biting, a potential driver of malaria transmission in Busia County, western Kenya. *Malaria Journal* 2024;**23**:1–13.

51. Akuoko OK, Dhikrullahi SB, Hinne IA *et al.* Biting behaviour and the spatio-temporal dynamics of malaria vectors in different ecological zones in Ghana. *Parasites & Vectors* 2022;**17**.

52. Afagbedzi SK, Alhassan Y, Kenu E *et al.* Universal coverage and utilization of free long-lasting insecticidal nets for malaria prevention in Ghana: a cross-sectional study. *Front Public Health* 2023;**11**:1140604.

53. Ohm JR, Baldini F, Barreaux P *et al.* Rethinking the extrinsic incubation period of malaria parasites. *Parasites and Vectors* 2018;**11**:178.

54. Shaw WR, Holmdahl I, Itoe M *et al.* Multiple blood feeding in mosquitoes shortens the Plasmodium falciparum incubation period and increases malaria transmission potential. *PLoS Pathogens* 2020;**16**:e1009131.

55. Suh E, Grossman MK, Waite JL *et al.* The influence of feeding behaviour and temperature on the capacity of mosquitoes to transmit malaria. *Nature Ecology and Evolution* 2020;**4**:940–51.

56. Costa G, Gildenhard M, Eldering M *et al.* Non-competitive resource exploitation within mosquito shapes within-host malaria infectivity and virulence. *Nature Communications* 2018;**9**:1–11.

57. Habtewold T, Sharma AA, Wyer CAS *et al.* Plasmodium oocysts respond with dormancy to crowding and nutritional stress. *Scientific Reports* 2021;**11**:1–10.

58. Gill S, Le HD, Melkani GC *et al.* Time-restricted feeding attenuates age-related cardiac decline in Drosophila. *Science* 2015;**347**:1265–9.

59. Rund SSC, Hou TY, Ward SM *et al.* Genome-wide profiling of diel and circadian gene expression in the malaria vector Anopheles gambiae. *Proceedings of the National Academy of Sciences of the United States of America* 2011;**108**, DOI: 10.1073/pnas.1100584108.