Biting time of day in malaria mosquitoes is modulated by nutritional status

Catherine E. Oke[1], Samuel S. C. Rund[3,4], Maxwell G. Machani[5], Abdul Rahim Mohammed Sabtiu[6],

Yaw Akuamoah-Boateng[6], Yaw A. Afrane[6], Sarah E. Reece[1,2]

[1]Institute of Ecology and Evolution and [2]Institute of Immunology and Infection Research, School of

Biological Sciences, University of Edinburgh, Edinburgh, UK; [3]Center for Research Computing and

[4]Department of Biological Sciences, University of Notre Dame, Notre Dame, USA; [5]Entomology

Department, Centre for Global Health Research, Kenya Medical Research Institute, Kisumu, Kenya;

[6]Department of Medical Microbiology, University of Ghana Medical School, College of Health

Sciences, University of Ghana, Accra, Ghana.

Corresponding author:

Catherine E. Oke, Email: catherine.oke@ed.ac.uk

Abstract

Background: Transmission of vector-borne pathogens follows daily rhythms, occurring at the time of day that vectors forage for blood. Control measures such as insecticide-treated bed nets (ITNs) have been particularly successful for reducing malaria transmission, because they exploit the nocturnal biting behaviour of the *Anopheles* spp. that vector malaria. However, shifts in biting behaviour to earlier or later hours when people are unprotected can undermine the efficacy of ITNs. Despite the implications for malaria transmission, the mechanisms underlying these shifts remain poorly understood. Because food availability mediates activity and foraging rhythms, and ITNs block access to human blood as a food source, we hypothesized that nutritional deprivation could cause mosquitoes to shift their biting behaviour to earlier or later in the diel cycle.

Methods: We provided female *Anopheles gambiae s.l.* mosquitoes with a blood meal on day 3 post-emergence, and access to one of three feeding treatments that varied in nutritional resources: (i) 0.5% sucrose, (ii) 10% sucrose, or (iii) 10% sucrose plus an additional blood meal on day 6. We released mosquitoes into a semi-field system on day 10 with human-mimic traps to investigate how food availability impacted the time of day that mosquitoes host seek.

Results: Nutritional resources determine both the likelihood and time of day that host seeking occurs. Specifically, low-resourced mosquitoes were 2-3 fold more likely to host seek overall, and 5-10 fold more likely to host seek at an earlier time of day than well-resourced mosquitoes (fed 10% sucrose with and without an additional blood meal), which predominantly sought a host in the second half of the night time.

Conclusions: We reveal that mosquito nutritional condition drives plasticity in biting time of day, suggesting it is an underappreciated contributor to residual malaria transmission. Furthermore, our results suggest that targeting mosquito nutrition (e.g. sugar-baited traps) could influence mosquito behaviour in ways that affect the success of ITNs. More broadly, understanding the drivers of biting time of day variation is crucial for the future success of vector control tools and controlling malaria transmission.

Key words:

Anopheles, transmission, vector behaviour, daily rhythms, food availability

Background

The transmission of vector-borne diseases (VBDs), such as malaria and dengue, relies on the

foraging rhythms of insect vectors, which take up pathogens from an infected vertebrate host during a

blood meal and transmit them to a new host during a subsequent blood-feeding event (1,2). For

example, *Plasmodium* parasites, the causative agent of malaria, are vectored between humans by

Anopheles spp. mosquitoes, which are generally nocturnally active and preferentially bite between the

hours of 11pm and 4am (*i.e.* the 'classical' time window) (3–5). Insecticide-treated bed nets (ITNs)

exploit these rhythmic foraging behaviours, and are consequently one of the most effective methods

for curbing the spread of malaria (6,7), averting 68% of deaths since 2000 (8). Despite this success,

residual transmission occurs, in part due to the evolution of physiological insecticide resistance (e.g.

biochemical and morphological modifications) and behavioural resistance in vector populations (9). In

particular, millions of clinical malaria cases are predicted to be occurring annually due to mosquitoes

altering the time of day they bite (10).

There are increasing reports that *Anopheles* spp. are shifting their biting time of day to earlier in the

evening ('early' biting) or later in the morning ('late' biting) when humans are unprotected by ITNs

(4,5,11–15), undermining the success of ITNs, especially in high-coverage regions. However, the

extent to which mosquitos can shift biting time of day and the range of potential drivers are poorly

understood. *Anopheles* spp. biting rhythms are weakly heritable, so shifts in biting time of day are

expected to be predominantly driven by phenotypic plasticity (16). Phenotypic plasticity evolves to

enable organisms to alter (via epigenetic mechanisms, for example) aspects of phenotype, including

behaviours, in manners that maximise fitness in response to environmental variation (17).

Furthermore, variation in physiological condition and resource limitation can cause animals to alter

their foraging rhythms (18) in adaptive (i.e. fitness enhancing) ways (19). For example, low food

availability promotes a shift to diurnal foraging patterns in small nocturnal rodents, which minimises

energy loss by remaining in burrows during particularly cold nights (20,21). Similarly, mosquitoes may

garner greater benefits from shifting biting time of day when they are in poor nutritional condition,

because the fitness benefits associated with acquiring a blood meal at suboptimal times of day

outweigh the increased risks of desiccation and predation. Such context-dependent variation in the

costs and benefits of plastic shifts in biting time may explain why biting times have seemingly not

significantly changed in some *Anopheles* spp. populations despite high ITN use (22,23), and why the

evolution of behavioural resistance varies across populations in manners that correlate with the

abundance of sources of nutrition for mosquitoes (24).

Mosquitoes rely on blood and sugar meals for reproduction and survival, with female mosquitoes

utilising proteins in blood for egg production, and using sugar sources to accumulate energy reserves

for powering energetically demanding processes, such as flight (25). Their feeding ecology is under

circadian clock control (26), with females seeking blood predominantly at night and approximately

every 3 days to fuel gonotrophic cycles of egg development and oviposition (27), and feeding on

sugar sources in between these cycles to promote survival until future biting opportunities (28,29).

Mosquitoes are likely to vary in their nutritional status because the type and abundance of sugar

sources available to mosquitoes varies across habitats and seasons (25,30), and vector control tools

such as ITNs can limit access to protein rich blood meals (31). In response to variation in food

sources, mosquitoes exhibit plasticity in multiple feeding behaviours. If a preferred host is not found,

mosquitoes are more likely to feed on sugar (32) or choose an alternative host (23), and will feed on

low-sugar plant tissues when sugar-rich plants are unavailable (33). However, despite the importance

of mosquito foraging rhythms for malaria transmission and that physiological condition alters foraging

rhythms in other taxa, the effect of nutritional resource availability on mosquito biting time of day is

unknown.

Here, we perturb the nutritional resources provided to adult mosquitoes to test whether nutritional

constraints cause a shift in biting time of day. Using a semi-field system, we simulated scenarios in

which mosquitoes have taken an initial blood meal (simulating the point at which malaria parasites

would be acquired) and then resided in environments that provide varying access to further blood

meals and sugar concentrations. Our three scenarios represented mosquitos that: (i) were

unsuccessful at acquiring a subsequent blood meal and only have access to low levels of sugar (0.5%

sucrose), (ii) were unsuccessful at acquiring a subsequent blood meal but have access to high levels

of sugar (10% sucrose), and (iii) successfully acquired a subsequent blood meal and have access to

high levels of sugar (10% sucrose plus an additional blood meal). We hypothesised that nutritionally

deprived mosquitoes, provided with limited sugar and blood, would be more likely to host seek and

shift their biting behaviour to earlier or later in the diel cycle, compared to better-resourced

mosquitoes. We also supplemented our data with a preliminary field collection to investigate if biting

time of day of wild-caught adult mosquitoes correlates with nutritional condition. We discuss the

potential outcomes of shifts in biting time of day in response to resource availability for understanding

residual transmission patterns, and how this may impact parasite development and transmission

between hosts.

Methods

We performed a semi-field experiment to assess if mosquito nutritional status impacts host seeking at

different times of day. We reared wild-caught larvae, and maintained blood fed female *Anopheles*

gambiae s.l. mosquitoes from the F_2 generation on diets that varied in nutritional resources, before

releasing them into an enclosed semi-field system with traps mimicking human odour. The three traps

were programmed to separately capture mosquitoes biting at early (evening), classical (during the

night) and late (morning) times of day. In addition, we collected wild adult *Anopheles* mosquitoes over

the course of one night, to test for correlations between mosquito nutritional status and the time of day

they attempted to bite.

Study sites

For the semi-field experiment, we collected *Anopheles gambiae s.l.* larvae from three urban sites

across the Greater Accra region of Ghana (Tuba, 5° 30' 47" N 0° 23' 16" W; Teshie, 5° 35′ 0″ N, 0° 6′

0″ W; East Legon, 5° 38' 16.39" N, 0° 9' 40.33" W) during the dry season in Dec 2023-Jan 2024

(Additional file 1, Fig. S1). Adult *An. gambiae s.l.* collections were conducted in Teshie during the dry

season in February 2024. The most abundant malaria vector species in these three sites are *An.*

gambiae s.s. and *An. coluzzii*, which both have a high Human Biting Index and predominantly bite

between 10pm and 4am (34,35). We transported larvae to the insectary at the Department of Medical

Microbiology, University of Ghana Medical School, Accra, and raised them to adults. We conducted

mosquito behavioural experiments in January-February 2024 in an enclosed semi-field system (the

"MalariaSphere", (38,39)) at the University of Ghana campus in Legon, Accra. The average (± SEM)

minimum and maximum daily temperatures during the experimental releases were 25.8±0.6°C and

34.5±0.4°C respectively (40). The enclosure was 5.8 x 4.2 x 2.8m and covered in an insect-proof

screen to prevent mosquito escape and/or entry from the external environment.

Semi-field host seeking experiment

To ensure sufficient sample sizes, we conducted semi-field experiments using the F_2 progeny of wild-

caught *An. gambiae s.l.* larvae (Fig 1A). We pooled collected larvae and then transferred them into

large tubs of water. We kept larvae at similar densities throughout the rearing procedure and fed them

on TetraMin Baby fish food. The emerged adults formed the F_0 generation, which we maintained in

cages under standard rearing conditions (26±2°C, 80% relative humidity with a 12L:12D cycle), with

ad libitum access to 10% sucrose solution in water. We provided a blood meal via direct-feeding (36),

and placed soaked cotton wool in petri dishes into the cages for egg laying. We reared the resulting

F_1 generation larvae and adults under the same conditions, and their eggs were used to produce the

experimental F_2 generation. On day 1 post-emergence of F_2 adults, we randomly allocated female

mosquitoes to three experimental cages (n=100-150 per cage) and allocated each cage to one of

three feeding treatments (Fig.1A): (i) 0.5% sucrose in water (0.5% suc), (ii) 10% sucrose in water

(10% suc) or (iii) 10% sucrose in water plus an additional blood meal on day 6 post-emergence (10%

suc + bm). We maintained all treatment groups on their respective sucrose solutions *ad libitum* from

day 1 post-emergence until the host seeking experiment. In addition, to mimic mosquito life history in

nature, we provided all groups with an initial blood meal on day 3 post-emergence. We gave

mosquitoes the opportunity to lay eggs after all blood meals. On day 9-10 post-emergence, we

transferred mosquitoes from each cage to paper cups (n=70-100 per cup), where they were housed

for six hours with access to water only prior to assaying biting time of day. We reared six batches of

mosquitoes to conduct a total of six semi-field system releases.

We placed three BG-Sentinel traps (Biogents, Germany) in the semi-field enclosure, with an MB5

human-mimic lure developed to attract anopheline mosquitoes (Biogents, Germany) and a CO_2 flow

of approximately 21g/hour, to mimic human scent and breath. We programmed each trap to turn on at

either 6pm-10pm to capture 'early' biters, 11pm-3am to capture 'classical' biters, or 4am-8am to

capture 'late' biters (Fig. 1B-D). We chose these timings because mosquitoes in this region

predominantly bite between 11pm and 4am, but biting can occur from approximately 6pm and persist

into the morning beyond 6am (34). Thus, these time windows also correspond to when people are

typically indoors and protected by ITNs (classical), or are outdoors and unprotected (early and late)

(37). CO_2 flow was controlled with BG-CO_2 timers (Biogents, Germany), ensuring CO_2 was only

emitted when the respective trap was switched on.

Within the MalariaSphere, released mosquitoes could access resting sites (four clay pots around the

enclosure). We did not provide water or food sources because this would have confounded the

experimental treatment groups. We colour-marked mosquitoes from each feeding treatment using

aerosolised fluorescent powder (FTX series, Swada London), applied following Machani et al (38).

We rotated the colour of each treatment group across sequential releases to ensure any differential

impacts of colour on mosquito behaviour did not confound treatment groups. We carried out six

releases over a period of three weeks, with mosquitoes being released into the enclosure between

5pm-5:30pm, just prior to dusk at 6pm (n=235-300 mosquitoes per release). A period of at least 48

hours between each release and checking all resting sites with a Prokopack aspirator ensured that

any remaining untrapped mosquitoes in the MalariaSphere had been removed or died before the next

release. We collected mosquitoes from the traps, at 8am the day after the release, then cold-

anaesthetised and counted the number caught in each trap.

Figure 1. The experimental design for the semi-field behavioural experiment. Schematic diagrams of

the entire experiment, from wild larvae collection to release of mosquitoes reared under the three

feeding treatments into the semi-field enclosure (A), and the semi-field assay set-up (B), featuring four

resting sites for mosquitoes, and traps baited with human odour which were programmed to turn on at

6pm-10pm, 11pm-3am and 4am-8pm respectively. These windows correspond to approximately

ZT12-ZT16, ZT17-ZT21, and ZT22-ZT2 respectively, where Zeitgeber time (ZT) defines the hours

since lights on and ZT0/24 is dawn. Each trap was linked to the CO_2 source, using separate BG-CO_2

timers to ensure the CO_2 was only flowing when the corresponding trap that was switched on.

Photographs show the semi-field facility (C) and the behavioural assay set-up (D).

Wild mosquito collections

In addition to the experiment, we also conducted a preliminary investigation into correlations between

biting time of day and nutritional content in wild mosquitoes, by collecting adult mosquitoes (n=95)

overnight between 6pm and 7am using the human landing catch (HLC) technique. This technique

involves trained human collectors exposing their lower legs and collecting mosquitoes that land on

them before they bite, and is the gold standard for investigating human exposure to disease vectors

(39). Every hour, we snap-froze collected mosquitoes on dry ice and identified specimens to genus

level according to morphological characteristics (40), noting the number of *Anopheles* spp.

mosquitoes collected per hourly bin. We placed individual anopheline mosquitoes in 1.5ml

microcentrifuge tubes and stored them on dry ice, transferring them to a -20°C freezer after the

overnight collection period had ended.

Nutrition assays

Immediately prior to each semi-field release in the experiment, we collected a subset of mosquitoes

from each feeding treatment group (n=14-19 per group, total n=48) to confirm our resource

perturbations impacted nutritional status. We also quantified mosquito nutritional status in individual

wild-caught mosquitoes (n=66). Mosquitoes were frozen at -20°C for lipid, glycogen and total sugar

analyses using modified Van Handel protocols (41–43). Firstly, we removed a wing from each

mosquito to quantify body size. Wing length is a commonly used proxy for adult body size because

dry weight and wing length are highly correlated (44), and is included in statistical analyses to ensure

any size variation does not confound differences in nutritional content. We lysed individual

mosquitoes in 100µl 2% sodium sulphate and added 750µl of 1:2 chloroform:methanol. We

centrifuged samples at 12000 rpm for 3 minutes; the supernatant was used for lipid and total sugars

analysis, and the precipitate was used for glycogen analysis. We conducted lipid, glycogen and total

sugars assays following Oke et al (45). For wild-caught mosquitoes, we used 250µl supernatant for

total sugars analysis.

Statistical analysis

We used R v. 4.1.3 to perform data analysis. We confirmed that adult body size did not differ across

the treatments using a linear model with wing length, mosquito release batch and their interaction as

main effects (Additional file 1, Fig. S2). To analyse nutritional content (lipid, glycogen, total sugars) of

individual mosquitoes, we used linear mixed models with feeding treatment as a main effect and

mosquito release batch as a random effect, accounting for any within treatment variation in body size.

We used binomial generalised linear mixed models (glmm) with mosquito release as a random effect

to account for any between-release variation to investigate how feeding treatment impacted the

overall proportion of mosquitoes caught across the night, with feeding treatment as a main effect, and

how feeding treatment impacted the proportion of mosquitoes caught across the different biting time

of day windows, with feeding treatment, trap time and their interaction as main effects. We estimated

proportion of caught mosquitoes and relative odds ratios (OR) ± SE from models using the *emmeans*

(46) package. We square root transformed nutrition data to meet assumptions of normality and

homogeneity of variance. We minimised all models using likelihood ratio tests and AICc for non-

nested models, and confirmed model assumptions using the *easystats* (47) package. To account for

differences between mosquito release batches, we present estimated marginal means ± SEM

(*emmeans* package), predicted from models. Whenever the minimised model contained a significant

effect of feeding treatment, we conducted post hoc pairwise comparisons using the Tukey method

with the *emmeans* package.

To investigate how nutritional content between field-caught and lab-reared mosquitoes varied, we

used linear models to compare between the field-caught, 0.5% suc, 10% suc and 10% suc + bm

groups. We performed a chi-square test to investigate if the number of field-caught mosquitoes

differed throughout the night by grouping hourly bins into 'early' biters (6pm-11pm), 'classical' biters

(11pm-4am) and 'late' biters (4am-7am). Using linear models, we analysed the nutritional content

(lipid, glycogen, total sugars) of individual wild-caught mosquitoes with biting time of day as an

unordered factor, and assessed the correlation between wing length and nutritional content. Nutrition

data were square root transformed to meet assumptions of normality and homogeneity of variance.

We conducted post hoc pairwise comparisons using the Tukey method with the *emmeans* package to

compare nutritional content between field-caught and lab-reared mosquitoes.

Results

Resource availability perturbs mosquito nutritional status

First, we confirmed that mosquitoes fed on the lowest concentration of sucrose (0.5%) had lower

nutritional reserves than mosquitoes fed 10% sucrose with or without an additional blood meal.

Mosquitoes fed on 0.5% sucrose had significantly lower lipid levels (55.5±17.8µg), approximately 2.5-

fold lower than those fed with 10% sucrose (138±31.4µg) and 3.3-fold lower than those fed 10%

sucrose plus an additional blood meal (182±29.6µg) (χ^2_2=17.6, p < 0.001; Fig. 2A, Table 1). A similar

pattern was observed for total sugars: mosquitoes fed with 0.5% sucrose had the lowest total sugar

levels (85.6±69.9µg), approximately 5-fold lower than those given 10% sucrose (420±158µg) and

10% sucrose plus an additional blood meal (439±156µg) (χ^2_2=30.8, p < 0.001; Fig. 2B, Table 1).

Finally, mosquitoes fed with 0.5% sucrose had the lowest glycogen levels (105±64.2µg),

approximately 7-fold lower than those given 10% sucrose (754±187µg), with mosquitoes given 10%

sucrose plus an additional blood meal exhibiting intermediate levels (481±128µg) (χ^2_2=21.7, p <

0.001; Fig. 2C, Table 1).

Figure 2. Concentrations (µg) per mosquito of lipids (A), total free sugars (B) and glycogen (C) in

individual mosquitoes under differing feeding treatments on day 10 post-emergence. Nutritional

perturbations began on day 1 post-emergence and all treatment groups received a blood meal on day

3 post-emergence (with an additional blood meal given to the 10% suc + bm group on day 6). Data

presented are estimated marginal means ± SEM.

Table 1. Post hoc pairwise comparisons for nutritional content of mosquitoes provided with

different feeding treatments. Significant p-values (<0.05) are highlighted in bold.

		Test statistic	*p-value*
Lipids	0.5% suc – 10% suc	*t* = 2.75	**0.02**
	0.5% suc – 10% suc + bm	*t* = -4.54	**<0.001**
	10% suc – 10% suc + bm	*t* = -1.16	0.48
Total sugars	0.5% suc – 10% suc	*t* = 5.13	**<0.001**
	0.5% suc – 10% suc + bm	*t* = -6.13	**<0.001**
	10% suc – 10% suc + bm	*t* = -0.21	0.98
Glycogen	0.5% suc – 10% suc	*t* = 4.66	**<0.001**
	0.5% suc – 10% suc + bm	*t* = -3.69	**0.002**
	10% suc – 10% suc + bm	*t* = 1.55	0.28

Resource availability affects host seeking tendency and biting time of day

Across all six releases and trapping times of day, the total mosquitoes trapped ranged from 21% to

49%, with a mean of 32.4% of mosquitoes caught in the traps. The likelihood of host seeking at any

time of day correlated negatively with the level of nutritional resources provided (χ^2_2=65.8, p < 0.001;

Fig. 3A). Specifically, mosquitoes fed 0.5% sucrose were 2-fold more likely to be trapped than those

fed 10% sucrose (odds ratio (OR): 2.08±0.28) and 3-fold more likely than those fed 10% sucrose plus

an additional blood meal (OR: 2.97±0.42). Mosquitoes fed only 10% sucrose were 1.4-fold more likely

to be trapped than those fed 10% sucrose plus an additional blood meal (OR: 1.43±0.21).

We also found that nutritional resources differentially affected the time of day of host seeking

(interaction between treatment and trapping time: χ^2_4=95.4, p < 0.001; Fig 3B). The majority of early

biting mosquitoes were from the 0.5% sucrose group, whereas mosquitoes with more resources were

more likely to be caught at classic and late times of day. Specifically, mosquitoes fed 0.5% sucrose

were approximately 5- and 10-fold more likely to be caught in the early biting window (20±2.9%) than

those fed with 10% sucrose (3.7±0.9%) and 10% sucrose plus an additional blood meal (1.6±0.5%),

respectively. At the classical and late trapping times of day, the proportion of mosquitoes fed 0.5%

sucrose caught was similar to mosquitoes fed 10% sucrose with or without an additional blood meal.

Mosquitoes fed 10% sucrose with or without an additional blood meal followed similar temporal

patterns across all trapping times of day, with approximately 4-fold more of these mosquitoes being

trapped during the classical (10% sucrose: 11±2.0%, 10% sucrose + bm: 9.0±1.7%) and late biting

windows (10% sucrose: 13±2.3%, 10% sucrose + bm: 11±2.0%) than during the early biting window

(10% sucrose: 3.7±0.9%, 10% sucrose + bm: 1.6±0.5%).

Figure 3. The proportion of released mosquitoes from each feeding treatment that were trapped

across the entire night (A) and that were caught in the evening (early, 6pm-10pm), classical night time

biting window (11pm-3am) and morning (late, 4am-8am) biting (B). Data presented are estimated

marginal means ± SEM.

Biting time distribution and nutritional status of wild host-seeking mosquitoes

We conducted a preliminary collection of wild host seeking *An. gambiae s.l.* mosquitoes to assess

whether mosquito nutritional status correlates with biting time of day. We caught a total of 95 wild host

seeking mosquitoes during our overnight collection, with a peak in biting between 12am-1am. We

grouped hourly bins into 'early' biters (6pm-11pm), 'classical' biters (11pm-4am) and 'late' biters (4am-

7am) and found that there were differences in the number of mosquitoes caught across these

windows. Specifically, 68% of mosquitoes were caught during the classical biting window, compared

to 7% and 24% caught during the evening and morning windows respectively, confirming that this

population predominantly bite during the 'classical' time of day (χ^2_2= 45.0, p < 0.001; Fig 4).

Figure 4. The proportion of wild adult *An. gambiae s.l.* mosquitoes caught by HLC collections across

hourly bins between 6pm and 7am. Early, classical, and late bins are indicated by shaded regions

from left to right. To highlight how the pattern changes over the course of the night, data points are

connected by an X-spline.

Although wild mosquitoes varied in their nutritional content, we found no differences between lipid

($F_{9,56}$ = 0.75, p = 0.66), sugars ($F_{9,56}$ = 1.06, p = 0.40) or glycogen ($F_{9,56}$ = 0.40, p= 0.93) contents

across hourly bins (Additional file 1, Fig S3). However, we found that wild host-seeking mosquitoes

had similar nutritional reserves to our experimental mosquitoes fed a 0.5% sucrose diet (estimated

marginal means for field-caught mosquito lipids: 39.7±5.5μg, sugars: 65.2±13μg, and glycogen:

51.4±11μg), but had significantly lower reserves than mosquitoes fed 10% sucrose and 10% sucrose

plus an additional blood meal (lipid: $F_{3,110}=24.8$, $p<0.001$; sugars: $F_{3,110}=25.9$, $p<0.001$; glycogen:

$F_{3,110}=58.1$, $p<0.001$; Table 2).

Table 2. Post hoc pairwise comparisons of the nutritional content of field-caught mosquitoes

versus. lab-reared mosquitoes on different feeding treatments. Significant p-values (<0.05) are

highlighted in bold.

		Test statistic	*p-value*
Lipids	field – 0.5% suc	$t = -1.26$	0.59
	field – 10% suc	$t = -4.83$	**<0.001**
	field – 10% suc + bm	$t = -7.93$	**<0.001**
Total sugars	field – 0.5% suc	$t = -1.14$	0.67
	field – 10% suc	$t = -4.96$	**<0.001**
	field – 10% suc + bm	$t = -8.07$	**<0.001**
Glycogen	field – 0.5% suc	$t = -1.97$	0.21
	field – 10% suc	$t = -11.3$	**<0.001**
	field – 10% suc + bm	$t = -8.87$	**<0.001**

Discussion

By altering sugar provision and access to blood meals, and using traps that mimic human breath and

scent, we find that resource availability modulates the time of day that *Anopheles gambiae s.l.*

mosquitoes in Ghana search for a blood meal. Mosquitoes on a low sucrose diet had significantly

lower levels of lipids, glycogen and sugar, and were 5-10 fold more likely to host seek in the evening

(early) than mosquitoes on a high sucrose diet with or without a blood meal (Fig. 2, 3B). In contrast,

better-resourced mosquitoes were most likely to host seek during the classical biting time window and

into the morning (Fig. 2, 3B). We also find that as resources increased, the overall tendency to host

seek decreased; mosquitoes on a low sucrose diet were 2-fold more likely to be caught compared to

mosquitoes given a higher concentration of sucrose, and 3-fold more likely than those with access to

10% sucrose and an additional blood meal (Fig. 3A). Finally, to probe the ecological relevance of our

experimental results, we also undertook a trapping session for wild host-seeking *An. gambiae s.l.*

mosquitoes. While the nutritional content of wild mosquitoes varied, we found that levels were similar

to the profiles of our low-resourced experimental mosquitoes fed 0.5% sucrose (Fig. 4). Given that

mosquito flight during host seeking is energetically consuming (48), this suggests that our

experimental feeding treatments resulted in mosquitoes with nutritional contents within an ecologically

realistic range.

Foraging at unusual times of day can be detrimental because it exposes organisms to rhythmic

environmental risks, such as the active phase of predators and suboptimal temperature/humidity, and

misalignment between metabolic and other circadian rhythms may impair digestion and metabolism

(1). However, under nutritional stress, finding food becomes more critical for survival, outweighing the

environmental risks and physiological costs (18,21). For example, a poorly-resourced mosquito may

begin host seeking earlier in the night because their lack of energy reserves reduce the chance of

surviving until later at night (or the morning, depending on whether ITNs limit access to hosts at

night). Our results are consistent with this hypothesis because poorly-resourced mosquitoes with low

nutritional reserves are more likely to host seek, predominantly in the early evening when

temperatures are higher and humidity is lower, and they risk being killed because human hosts are

alert. Although we did not quantify survival, we did not encounter any live mosquitoes fed on a 0.5%

sucrose diet whilst clearing the semi-field systems the day after each release, but regularly

encountered live mosquitoes fed on higher sucrose solutions and/or an additional blood meal. This

indicates that mosquitoes fed 0.5% sucrose were low-resourced and did not have sufficient reserves

to survive until morning. Overall, our results suggest that early biting could be an adaptive plastic

temporal shift to maximise fitness in sugar-poor regions and where hosts are not easily accessible

due to high ITN use.

Better-resourced mosquitoes on high sucrose diets (with or without an additional blood meal) had

higher nutritional reserves and waited until the classical time window to host seek, in alignment with

their circadian physiology and when environmental risks are likely to be lower, which also supports

our hypothesis. However, we also found that better-resourced mosquitoes were just as likely as

poorly resourced mosquitoes to be trapped later in the morning. Such late biting by well-fed

mosquitoes was unexpected but may be due to the lack of water/food sources provided in the semi-

field enclosures, forcing untrapped mosquitoes to host seek to mitigate dehydration caused by the

rise in temperature and the reduction in humidity in the morning (49,50). Alternatively, well-resourced

mosquitoes which had failed to locate a host (*i.e.* had not entered a trap) during the night may have

become resource-depleted because flight is energetically demanding (48), leading to an increased

propensity to continue host seeking because they do not have sufficient energy to survive until the

subsequent night. Regardless of the explanation(s), our results reveal that adult nutritional condition

drives variation in biting time of day.

Our semi-field system allowed mosquitoes to experience ecologically realistic temperature and

humidity fluctuations and provided sufficient flight space, but excluded other ecological factors,

including predation and access to alternative food sources. Thus, to fully investigate how resource

availability impacts biting time, future studies should consider adding plants of varying sugar contents

and alternative blood hosts into semi-field systems. However, these omissions were necessary in our

experiment to isolate the effect of adult nutritional status on biting behaviour. To probe the ecological

relevance of our experimental results, we also undertook a preliminarily trapping session for wild host-

seeking *Anopheles gambiae* s.l. mosquitoes. We found that they had similar nutritional profiles to our

low-resourced mosquitoes fed on 0.5% sucrose, suggesting that our experimental perturbations

represented natural conditions. The low sample size for wild caught mosquitoes precludes comparing

nutritional reserves between host seeking times of day but does suggest that adult nutritional content

does not correlate with body size (a product of larval condition) (Additional file 1, Fig. S4). Thus,

catching lowly resourced mosquitoes at the time of host seeking is consistent with our experimental

results revealing that adult nutrition influences biting rhythms. Further field studies are necessary to

build on our findings, including sampling host-seeking mosquitoes across multiple sites, both indoors

and outdoors, throughout a 24-hour period, examining their nutritional status, and investigating when

mosquitoes become active.

Like previous studies across Africa (3,35,51), we found a large proportion of wild mosquitoes

(approximately 20%) host seeking during the morning when humans are unprotected by ITNs. This

pattern of 'late' biting has become more pronounced since the widespread use of ITNs. Because

biting time shifts have a genetic component (16), and universal coverage of ITNs is approximately

75% in Ghana (52), late biting could be an evasion strategy evolving in the mosquito population

where the larval collections for our experiment were carried out, rather than an entirely plastic

behavioural change. Further work is needed to ascertain the extent to which biting time of day is

genetically determined and/or due to behavioural plasticity, because plasticity itself evolves and can

facilitate or constrain evolution that genetically segregates mosquito populations into early, classical

or late biters. Modelling suggests that early biting is more likely to evolve in sugar-poor environments

than sugar-rich environments, for example, due to differences in the abundance and diversity of

plants (24). However, in areas of fluctuating resource availability or ITN coverage, plastic genotypes

that can switch between early, classical, or late biting depending on their environmental conditions

should have higher fitness (17).

Both our semi-field experiment and preliminary field data reveal significant variation in biting time of

day, and intuition suggests that early biting by poorly-resourced mosquitoes could sustain malaria

transmission. However, the overall impact of shifts in mosquito biting time of day for malaria

transmission is difficult to predict, because of complex and potentially opposing effects on both

mosquito and parasite fitness. While early biting causes greater contact between vectors and hosts in

regions with high ITN coverage, malaria parasite development is less productive in low-resourced

mosquitoes, and mosquito survival is reduced (45). Thus, the poor nutritional state of early biters

could curtail onward transmission through lower parasite numbers (53) and because vectors may not

survive long enough for malaria parasites to complete their development (54,55). In addition, parasite

establishment in mosquitoes is impacted by daily temperature rhythms (56) and parasite infectivity

and mosquito susceptibility vary between day and night (1,57,58). Disruption to foraging rhythms also

reduces fitness in other insects (59), and in mosquitoes could lead to mismatch between metabolic

processes and blood digestion (60). The net outcome for malaria transmission of these myriad and

potentially synergistic and antagonistic effects of biting time of day is challenging to predict. However,

recognising that transmission is time of day dependent is necessary for accurately predicting

epidemiology, the risk of clinical malaria cases, and the trajectory of parasite evolution.

Conclusions

Our results suggest that plasticity in biting time of day, driven by variation in adult mosquito nutritional

condition, may be an underappreciated contributor to residual malaria transmission. In particular,

nutritional stress in mosquito populations may lead to shifts in biting behaviour, reducing ITN efficacy.

This highlights that mosquito nutrition and its influence on the timing of biting behaviour should be

considered when implementing and assessing vector control strategies, especially those which target

mosquito foraging behaviour, such as ITNs and sugar-baited traps. More broadly, investigating the

drivers of variation in biting time of day and the overall impact on pathogen transmission is critical for

assessing and improving the efficacy of current and future vector control tools. This is especially

important for malaria, where up to 30% of biting is reported to occur when humans are unprotected by

ITNs and is predicted to lead to millions of additional clinical cases per year (3,10). However, we also

expect our findings to apply to other vector-borne diseases that rely on vector foraging rhythms for

between-host transmission.

Declarations

Ethics approval and consent to participate

The study protocols for mosquito colony blood feeds and human landing catches (HLCs) were

ethically reviewed and approved by the Ghana Health Service Ethics Review Committee (GHS-ERC:

021/07/23). Volunteers involved in colony blood feeds were tested for malaria prior to direct-feeding,

and the same person fed the mosquitoes throughout the experiment. Permission to conduct HLCs in

study sites was obtained from the District Assembly Representative, and written consent was

obtained from all collectors. Collectors had access to malaria prophylaxis and did not report any

adverse events.

Consent for publication

Not applicable.

Availability of data and materials

The datasets supporting the conclusions of this article are available in the Edinburgh DataShare

repository and can be accessed with [https://doi.org/LINK AVAILABLE UPON ACCEPTANCE].

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by a Varley-Gradwell Travelling Fellowship in Insect Ecology (2023-2024) to

C.E.O, and by a Wellcome Trust Investigator Award (202769/Z/16/Z) and Wellcome Trust Discovery

Award (227904/Z/23/Z) to S.E.R. Research equipment and travel funds for M.G.M. and S.S.C.R. were

funded by an Initiation Grant from the University of Notre Dame, Notre Dame Research office. Y.A. A.

is funded by the National Institute for Health (D43 TW 011513). The funders had no role or influence

on the design of this study, data collection, analyses, and interpretation of the data collected, as well

as in writing this manuscript.

Authors' contributions

Conceptualisation: CEO, SSCR, YAA, SER. Methodology: CEO, SSCR, MGM, YAA, SER.

Investigation: CEO, SSCR, MGM, ARMS, YAB. Analysis: CEO. Writing-original draft: CEO, SER.

Writing-review and editing: CEO, SSCR, MGM, YAA, SER. All authors read and approved the final

manuscript.

Acknowledgements

We thank Anisa Abdulai for experimental and technical assistance, and the serology lab team at the

Noguchi Memorial Institute for Medical Research for kindly letting us use their facilities.

References

1.   Rund SSC, O'Donnell AJ, Gentile JE, Reece SE. Daily rhythms in mosquitoes and their
consequences for malaria transmission. Insects. 2016;7(2):1–20.

2.   Westwood ML, O'Donnell AJ, de Bekker C, Lively CM, Zuk M, Reece SE. The evolutionary
ecology of circadian rhythms in infection. Nature Ecology and Evolution. 2019;3(4):552–60.

3.   Sangbakembi-ngounou C, Costantini C, Longo-pendy NM, Ngoagouni C, Akone-ella O, Rahola
N. Diurnal biting of malaria mosquitoes in the Central African Republic indicates residual
transmission may be " out of control ." Proceedings of the National Academy of Sciences.
2022;119(21):e2104282119.

4.   Thomsen EK, Koimbu G, Pulford J, Jamea-Maiasa S, Ura Y, Keven JB, et al. Mosquito behavior
change after distribution of bednets results in decreased protection against malaria exposure.
Journal of Infectious Diseases. 2017;215(5):790–7.

5.   Moiroux N, Gomez MB, Pennetier C, Elanga E, Djènontin A, Chandre F, et al. Changes in
anopheles funestus biting behavior following universal coverage of long-lasting insecticidal nets in
benin. Journal of Infectious Diseases. 2012;206(10):1622–9.

6.   Oke CE, Ingham VA, Walling CA, Reece SE. Vector control: agents of selection on malaria
parasites? Trends in Parasitology. 2022;38(10):890–903.

7.   World Health Organisation. World Malaria Report 2023. Geneva; 2023.

8.   Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria
control on Plasmodium falciparum in Africa between 2000 and 2015. Nature.
2015;526(7572):207–11.

9.   Huijben S, Paaijmans KP. Putting evolution in elimination: Winning our ongoing battle with
evolving malaria mosquitoes and parasites. Evolutionary Applications. 2018;11(4):415–30.

10. Sherrard-Smith E, Skarp JE, Beale AD, Fornadel C, Norris LC, Moore SJ, et al. Mosquito feeding
behavior and how it influences residual malaria transmission across Africa. Proceedings of the
National Academy of Sciences of the United States of America. 2019;116(30):15086–96.

11. Russell TL, Beebe NW, Cooper RD, Lobo NF, Burkot TR. Successful malaria elimination
strategies require interventions that target changing vector behaviours. Malaria Journal.
2013;12(1):1–5.

12. Sougoufara S, Diédhiou SM, Doucouré S, Diagne N, Sembène PM, Harry M, et al. Biting by
Anopheles funestus in broad daylight after use of long-lasting insecticidal nets: A new challenge
to malaria elimination. Malaria Journal. 2014;13(1):1–7.

13. Yohannes M, Boelee E. Early biting rhythm in the afro-tropical vector of malaria, Anopheles
arabiensis, and challenges for its control in Ethiopia. Medical and Veterinary Entomology.
2012;26(1):103–5.

14. Cooke MK, Kahindi SC, Oriango RM, Owaga C, Ayoma E, Mabuka D, et al. "A bite before bed":
Exposure to malaria vectors outside the times of net use in the highlands of western Kenya.
Malaria Journal. 2015;14(1):1–15.

15. Carrasco D, Lefèvre T, Moiroux N, Pennetier C, Chandre F, Cohuet A. Behavioural adaptations of
mosquito vectors to insecticide control. Current Opinion in Insect Science. 2019;34:48–54.

16. Govella NJ, Johnson PCD, Killeen GF, Ferguson HM. Heritability of biting time behaviours in the
major African malaria vector Anopheles arabiensis. Malaria Journal. 2023;22(1):1–9.

17. Pigliucci M. Evolution of phenotypic plasticity: Where are we going now? Trends in Ecology and
Evolution. 2005;20(9):481–6.

18. Van Der Veen DR, Riede SJ, Heideman PD, Hau M, Van Der Vinne V, Hut RA. Flexible clock
systems: adjusting the temporal programme. Phil Trans R Soc B. 2017 Nov
19;372(1734):20160254.

19. Van Der Vinne V, Riede SJ, Gorter JA, Eijer WG, Sellix MT, Menaker M, et al. Cold and hunger
induce diurnality in a nocturnal mammal. Proc Natl Acad Sci USA. 2014 Oct 21;111(42):15256–
60.

20. Van Der Vinne V, Gorter JA, Riede SJ, Hut RA. Diurnality as an energy-saving strategy: energetic
consequences of temporal niche switching in small mammals. Journal of Experimental Biology.
2015 Aug 1;218(16):2585–93.

21. Van Der Vinne V, Tachinardi P, Riede SJ, Akkerman J, Scheepe J, Daan S, et al. Maximising
survival by shifting the daily timing of activity. Ghalambor C, editor. Ecology Letters. 2019
Dec;22(12):2097–102.

22. Sougoufara S, Harry M, Doucouré S, Sembène PM, Sokhna C. Shift in species composition in
the Anopheles gambiae complex after implementation of long-lasting insecticidal nets in Dielmo,
Senegal. Medical and veterinary entomology. 2016;30(3):365–8.

23. Kreppel KS, Viana M, Main BJ, Johnson PCD, Govella NJ, Lee Y, et al. Emergence of
behavioural avoidance strategies of malaria vectors in areas of high LLIN coverage in Tanzania.
Scientific Reports. 2020;10(1):1–11.

24. Stone C, Chitnis N, Gross K. Environmental influences on mosquito foraging and integrated
vector management can delay the evolution of behavioral resistance. Evolutionary Applications.
2016;9(3):502–17.

25. Barredo E, Degennaro M. Not Just from Blood: Mosquito Nutrient Acquisition from Nectar
Sources. Trends in Parasitology. 2020;36(5):473–84.

26. Rund SSC, Bonar NA, Champion MM, Ghazi JP, Houk CM, Leming MT, et al. Daily rhythms in
antennal protein and olfactory sensitivity in the malaria mosquito Anopheles gambiae. Sci Rep.
2013 Aug 29;3(1):2494.

27. Mitchell SN, Catteruccia F. Anopheline reproductive biology: Impacts on vectorial capacity and
potential avenues for malaria control. Cold Spring Harbor Perspectives in Medicine.
2017;7(12):14.

28. Foster WA. Behavioural ecology of plant-mosquito relations. In: Ignell R, Lazzari CR, Lorenzo
MG, Hill SR, editors. Sensory ecology of disease vectors [Internet]. The Netherlands:
Wageningen Academic Publishers; 2022. p. 171–234. Available from:
https://doi.org/10.3920/978-90-8686-932-9

29. Ebrahimi B, Jackson BT, Guseman JL, Przybylowicz CM, Stone CM, Foster WA. Alteration of
plant species assemblages can decrease the transmission potential of malaria mosquitoes.
Journal of Applied Ecology. 2018;55(2):841–51.

30. Gu W, Müller G, Schlein Y, Novak RJ, Beier JC. Natural plant sugar sources of Anopheles
mosquitoes strongly impact malaria transmission potential. PLoS ONE. 2011;6(1).

31. Killeen GF, Smith TA, Ferguson HM, Mshinda H, Abdulla S, Lengeler C, et al. Preventing
childhood malaria in Africa by protecting adults from mosquitoes with insecticide-treated nets.
PLoS Medicine. 2007;4(7):1246–58.

32. Gary RE, Foster WA. Diel timing and frequency of sugar feeding in the mosquito Anopheles
gambiae, depending on sex, gonotrophic state and resource availability. Medical and Veterinary
Entomology. 2006;20(3):308–16.

33. Müller G, Schlein Y. Plant tissues: the frugal diet of mosquitoes in adverse conditions. Medical
Vet Entomology. 2005 Dec;19(4):413–22.

34. Sabtiu ARM, Sr IK, Mfum COA, Akuamoah-Boateng Y, Doe RT, Boadu EN, et al. Malaria
transmission risk in the city of Accra, Ghana: vector behavior and distribution [Internet]. 2025
[cited 2025 July 2]. Available from: https://www.researchsquare.com/article/rs-6008722/v1

35. Akuoko OK, Dhikrullahi SB, Hinne IA, Mohammed AR, Mfum Owusu-Asenso C, Coleman S, et al.
Biting behaviour and the spatio-temporal dynamics of malaria vectors in different ecological
zones in Ghana. Parasites & Vectors [Internet]. 2024;17(16). Available from:
https://doi.org/10.21203/rs.3.rs-1975878/v1

36. Harrington LC, Foy BD, Bangs MJ. Considerations for Human Blood-Feeding and Arthropod
Exposure in Vector Biology Research: An Essential Tool for Investigations and Disease Control.
Vector-Borne and Zoonotic Diseases. 2020 Nov 1;20(11):807–16.

37. Nzioki I, Machani MG, Onyango SA, Kabui KK, Githeko AK, Ochomo E, et al. Current
observations on shifts in malaria vector biting behavior and changing vulnerability to malaria
transmission in contrasting ecosystems in Western Kenya. Parasites & Vectors. 2023;1–19.

38. Machani MG, Ochomo E, Amimo F, Mukabana WR, Githeko AK, Yan G, et al. Behavioral
responses of pyrethroid resistant and susceptible Anopheles gambiae mosquitoes to insecticide
treated bed net. PLoS ONE. 2022;17(4):e0266420.

39. Gimnig JE, Walker ED, Otieno P, Kosgei JK, Olang GB, Ombok MO, et al. Incidence of Malaria
among Mosquito Collectors Conducting Human Landing Catches in Western Kenya. The
American Journal of Tropical Medicine and Hygiene. 2013 Feb 6;88(2):301–8.

40. Gillies MT, Coetzee M. A Supplement to the Anophelinae of Africa South of the Sahara.
Publications of the South African Institute for Medical Research. 1987;55:1–143.

41. Van Handel E, Day JF. Assay of lipids, glycogen and sugars in individual mosquitoes:
correlations with wing length in field-collected Aedes vexans. Journal of the American Mosquito
Control Association. 1988;4(4):549–50.

42. Van Handel E. Rapid determination of glycogen and sugars in mosquitoes. Journal of the
American Mosquito Control Association. 1985;1:299–301.

43. Van Handel E. Rapid determination of total lipids in mosquitoes. Journal of the American
Mosquito Control Association. 1985;(1):302–4.

44. Lehmann T, Dalton R, Kim EH, Dahl E, Diabate A, Dabire R, et al. Genetic contribution to
variation in larval development time, adult size, and longevity of starved adults of Anopheles
gambiae. Infection, Genetics and Evolution. 2006 Sept;6(5):410–6.

45. Oke CE, O'Donnell AJ, Schneider P, Reece SE. Plasticity in malaria parasite development:
mosquito resources influence vector-to-host transmission potential. Front Malar. 2024 Nov
1;2:1481816.

46. Lenth RV. emmeans: Estimated Marginal Means, aka Least-Squares Means [Internet]. 2023.
Available from: https://cran.r-project.org/package=emmeans

47. Makowski D, Ben-Shachar M, Lüdecke D. The {easystats} collection of R packages [Internet].
2020. Available from: https://easystats.github.io/easystats/

48. Kaufmann C, Briegel H. Flight performance of the malaria vectors Anopheles gambiae and
Anopheles atroparvus. Journal of Vector Ecology. 2004;29(1):140–53.

49. Hagan RW, Didion EM, Rosselot AE, Holmes CJ, Siler SC, Rosendale AJ, et al. Dehydration
prompts increased activity and blood feeding by mosquitoes. Scientific Reports. 2018;8(1):1–12.

50. Lin S, Senapati B, Tsao CH. Neural basis of hunger-driven behaviour in Drosophila. Open
Biology. 2019;9(3).

51. Odero JI, Abong'o B, Moshi V, Ekodir S, Harvey SA, Ochomo E, et al. Early morning anopheline
mosquito biting, a potential driver of malaria transmission in Busia County, western Kenya.
Malaria Journal. 2024;23(1):1–13.

52. Afagbedzi SK, Alhassan Y, Kenu E, Malm K, Bandoh DAB, Peprah NY, et al. Universal coverage
and utilization of free long-lasting insecticidal nets for malaria prevention in Ghana: a cross-
sectional study. Front Public Health. 2023 May 25;11:1140604.

53. Aleshnick M, Ganusov VV, Nasir G, Yenokyan G, Sinnis P. Experimental determination of the
force of malaria infection reveals a non-linear relationship to mosquito sporozoite loads. PLoS
Pathogens. 2020;16(5):e1008181.

54. Ohm JR, Baldini F, Barreaux P, Lefevre T, Lynch PA, Suh E, et al. Rethinking the extrinsic
incubation period of malaria parasites. Parasites and Vectors. 2018;11:178.

55. Shaw WR, Holmdahl I, Itoe M, Werling K, Marquette M, Paton D, et al. Multiple blood feeding in
mosquitoes shortens the Plasmodium falciparum incubation period and increases malaria
transmission potential. PLoS Pathogens. 2020;16(12):e1009131.

56. Suh E, Grossman MK, Waite JL, Dennington NL, Sherrard-Smith E, Churcher TS, et al. The
influence of feeding behaviour and temperature on the capacity of mosquitoes to transmit
malaria. Nature Ecology and Evolution. 2020;4(7):940–51.

57. Pigeault R, Caudron Q, Nicot A, Rivero A, Gandon S. Timing malaria transmission with mosquito
fluctuations. Evolution Letters. 2018;2(4):378–89.

58. Schneider P, Rund SSC, Smith NL, Prior KF, O'Donnell AJ, Reece SE. Adaptive periodicity in the
infectivity of malaria gametocytes to mosquitoes. Proceedings of the Royal Society B: Biological
Sciences. 2018;285(1888).

59. Gill S, Le HD, Melkani GC, Panda S. Time-restricted feeding attenuates age-related cardiac
decline in Drosophila. Science. 2015;347(6227):1265–9.

60. Rund SSC, Hou TY, Ward SM, Collins FH, Duffield GE. Genome-wide profiling of diel and
circadian gene expression in the malaria vector Anopheles gambiae. Proceedings of the National
Academy of Sciences of the United States of America. 2011;108(32).

Additional files

Additional file 1 (.pdf)

Supplementary Figures.

Figure S1 is a map of larval and adult collection sites, Figure S2 is the wing length of lab-reared

mosquitoes under different feeding treatments, Figure S3 provides information about the nutritional

contents of wild adult mosquitoes collected at different times of day, and Figure S4 is the correlation

between wing length and nutritional content in wild adult mosquitoes.