Repeated blood-brain barrier opening using low-intensity pulsed ultrasound mitigates amyloid pathology

Michael Canney[1], Guillaume Bouchoux[1], Alexandre Carpentier[2], Pierre De Rossi[1*]

[1]Carthera, Lyon, France

[2]Sorbonne Université, AP-HP, Hôpitaux Universitaires La Pitié Salpêtrière - Charles Foix, Service de Neurochirurgie, Paris, France

*Corresponding author: pierre.derossi@carthera.eu

Abstract

Background: The delivery of large molecules to the pathological brain is one of the main obstacles in the development of disease-modifying drugs. This is partly due to the presence of the blood-brain barrier (BBB), which blocks the free passage of lipophobic molecules and those larger than 400 Da. One strategy to bypass this natural barrier is to use low intensity pulsed ultrasound (LIPU) to oscillate circulating micro-sized microbubbles which then exert mechanical stress on the vessel walls. This procedure allows for temporary disruption of the BBB and enhanced local delivery of therapeutics from the blood to the brain parenchyma. In this study, the effect of repeated BBB opening on neuroinflammation in a healthy mouse model was first explored followed by the effect of repeated opening on amyloid pathology in a model for Alzheimer's disease.

Methods: A cohort of wildtype mice was used to determine the effect of a single BBB opening session mediated by ultrasound/microbubbles (US/MB) on the inflammatory profile, using RT-Q-PCR on brain extracts at 2-, 4-, 8- and 15-days post opening. A second cohort of ARTE10 mice, a mouse model for amyloid pathology, was treated with a different sequence of repeated US/MB mediated BBB opening to explore the effect on the pathology. Tissues were also analyzed for immune cell infiltration, microglia and astrocyte activation, and inflammatory response.

Results: Our results demonstrate that the opening of the BBB leads to a mild inflammatory response in wild-type animals. However, repeated opening of the BBB in the ARTE10 model resulted in a mild decrease in amyloid pathology, along with a mild increase of growth factor.

Conclusions: Altogether, this study suggests that sonication is not only a safe method to deliver therapeutics to the brain but could also have synergistic effects in the treatment of neurodegenerative diseases.

Introduction

Alzheimer's disease (AD) is a neurodegenerative disorder that slowly affects memory, thinking and behavior, associated with progressive neurodegeneration. AD is characterized by two molecular hallmarks: the extracellular aggregation of amyloid-beta in plaques and the intracellular aggregation of hyperphosphorylated tau, resulting in tau tangles. These progressive pathological features have been hypothesized to lead to the resulting cognitive deterioration observed in patients.

While the last two decades of research have uncovered several risk factors for AD and led to the discovery of new pathways that may be responsible for disease onset, most pharmaceutical development has been focused on developing therapies targeting tau and amyloid. The hypothesis behind this strategy is that the removal of these pathological aggregates will lead to restoration of normal neuronal function and delay or stop the progression of the disease.

Recently, the first regulatory approvals have been obtained for antibody-based therapies targeting amyloid for the treatment of AD, despite only showing modest results in Phase 3 clinical trials[1]. However, one of the main obstacles that limit the efficacy of these antibody treatments targeting amyloid is the difficulty in bypassing the blood-brain barrier (BBB). Normally, only 0.01%[2] of the administered antibody reaches the brain. If higher parenchymal concentrations could be reached, we could observe a better clearance of the plaques associated with a better efficacy of these treatments, as recently demonstrated by the last results for the phase 2 clinical trial for the Trontinemab[3].

To overcome the BBB, ultrasound (US) technology can be used to temporarily disrupt the BBB and increase delivery of therapeutics. This method uses the combination of circulating/oscillating microbubbles (MB) administered intravenously and stimulated by low intensity pulsed ultrasound (LIPU) to disrupt the tight junctions of the BBB, allowing enhanced entry of molecules, ranging from small molecule therapeutics to large antibodies into the parenchyma[4–21]. This strategy has been used to enhance delivery of therapeutics into the CNS of mouse models for AD[4,7,9]. In addition, a recent clinical trial by Rezai et al. 2024[15] has shown promising results for delivering the FDA-approved anti-amyloid therapy aducanumab in Alzheimer's patients using US/MB with a clinical ultrasound system. A 32% reduction in SUVR on ^{18}F-florbetaben PET scans was observed after 26 weeks in the regions that received treatment to open the BBB. These clinical results demonstrate the potential efficacy gained by the delivery of antibody therapies using ultrasound.

US-mediated BBB opening alone, without concomitant drug administration has also been

shown to have beneficial effects, potentially by altering the immune microenvironment in

Alzheimer's mouse models and in patients[4,22–30], with improved outcomes when coupled

to anti-amyloid therapies[4,31–35]. Moreover, it was shown that opening of the BBB can also

potentiate a diverse range of beneficial biological effects including neurogenesis[36–38]. In

amyloid models of the disease, repeated BBB opening decreased the number of amyloid

deposits in the brain[22,24,27–30,35] and improved cognition[22,24,27,30]. Similarly, in tau

models of the disease, repeated BBB opening was able to decrease tau pathology[39,40]

and improve cognition[39]. Finally, in patients with AD, repeated opening of the BBB using

US was shown to be safe[41–44] and efficient to mitigate the amyloid burden in the

brain[43,44].

While several studies have already explored the effect of BBB opening using US/MB on the

inflammatory profile of the brain at short term, evidence is still lacking on the lasting effects

of BBB opening in the healthy brain as well as its potential as a therapeutic approach for

neurodegenerative diseases. In this pilot study, we explored the effect of BBB opening in

wildtype mice and in a mouse model for amyloid pathology (ARTE10) on the inflammatory

profile of the brain and its effect on neuropathological hallmarks.

Experimental Procedures

Mouse models

Twenty (20) male C57BL/6J mice of approximately 17 weeks of age were obtained from The

Jackson Laboratory (Bar Harbor, ME). Thirty-six (36) one-year-old females of the ARTE10

mouse model were used as a model for amyloid pathology and obtained from Taconic.

Sonication

The following ultrasound parameters were used: 120 seconds using a 1-MHz 25,000-cycle

burst at a 1 Hz pulse repetition frequency and an acoustic pressure of 0.3 MPa (Carthera,

Lyon, France). To prepare for sonication, the hair on the head of all animals was removed

using depilatory cream. At the time of sonication, mice were injected intraperitoneally with 5

µL/gram of 90 mg/kg ketamine / 10 mg/kg xylazine (Zetamine: VetOne, catalogue No.

AH202C6)/xylazine (Rompun: Bayer, catalogue No. AH200W5). Lumason microbubbles

(Bracco Diagnostics, catalogue No. 20A108A) were injected into the retro-orbital sinus (r.o.)

at a dose of 150 µL/mouse and mice underwent a single sonication event. Mice were given

10 mg/kg atipamezole (Rivertidine; Modern Veterinary Therapeutics, Cat #023575) i.p. within

20 min of ketamine administration to reverse the anesthesia. After 15 min, blood was

collected for PK.

Experimental Design for WT animals

Animals were weighed on study days 0, 1, 7, and 14. A pre-sonication blood sample was

collected on study day 0, as a baseline reference. On study day 1, the animals received a

single sonication event. All animals were bled by retro-orbital blood drawn for plasma

(EDTA) on study day 0 (baseline; 120 µL of plasma/mouse). Plasma samples were stored

frozen at –80°C for analysis of S100. The animals were euthanized for necropsy and

terminal sample collection 1 day, 3 days, 7 days, or 14 days post-sonication.

Experimental Design for ARTE10 animals

Animals were weighed and clinical observations were recorded on D0, D7, D14, D21, D28,

and D49. A pre-sonication blood sample was collected on study day 0. Groups 1 and 2 did

not receive any sonication (control groups). Groups 3 and 5 (sparse sonication groups, 4

sonication total) received repeated sonication every 2 weeks for 8 weeks with respective

sacrifice 1 week and 1 month after the last sonication procedure. Group 4 (dense sonication

group, 8 sonication total) received repeated sonication twice a week for 4 weeks and were

sacrificed 1 week after the last sonication. Group 6 (very sparse sonication group, 2

sonication total) received repeated sonication every 4 weeks for 8 weeks and were sacrificed

1 week after the last sonication.

Postmortem brain tissue collection

Brains were halved along the longitudinal fissure. Right brain hemispheres were fixed in 4%

PFA, then transferred to 40% sucrose/saline the next day and stored at 4°C for histology.

Left brain hemispheres were collected into RNase/DNase free clean tubes and stored at

–80°C for gene expression analysis. Cerebrospinal fluid (CSF) samples were collected and

whole blood was collected and processed to plasma (EDTA, 300 µL/mouse). Plasma and

CSF samples (5-10 µL/mouse) were stored at –80°C for analysis.

S100-β ELISA Methods

150   Plasma samples were analyzed for S100-β using a Mouse Protein S100-β /S100 Beta

151   ELISA Kit (CUSABio, catalogue No. CSB-EL020643MO, lot No. J16147391). All reagents,

152   working standards, and samples were prepared as directed in the kit instructions.

Real-Time Quantitative Polymerase Chain Reaction (RT-qPCR) Methods

For WT animals: RT-qPCR analysis was performed on left brain hemispheres from
all animals. Tissue samples (approx. 30 mg each) were weighed out and homogenized in
RLT buffer on a FastPrep-24 5G. RNA was isolated using a Qiagen RNeasy Mini kit (No.
74106) and the manufacturer-recommended protocol, and RNA was reverse transcribed into
cDNA using the ThermoFisher High-Capacity cDNA kit (ThermoFisher, No. 4368814). cDNA
samples were then analyzed using Taqman primers against *cyc1*, *cxcl1*, *ccl2*, *Il-1β*, *Il-6*, *tnf-*
a, *Icam*, *Il-10*, *tgf-β*, *Il-13*, *bdnf*, *nf-κb*, *p-Selectin*, *e-Selectin*, GM-CSF, and G-CSF. All
primers were selected from validated primer sets from ThermoFisher Scientific.

For ARTE10 cohort: Multiplex qPCR master mix was prepared using TaqMan
Multiplex Master Mix (ThermoFisher, Cat #4484263) according to manufacturer's
specifications for 96-well plates. Samples were run in triplicate for the following genes: *Il-1β,*
Il-6, Il-10, tnf-α, bdnf, ccl2, Icam1

Histological Processing
Mouse hemibrains were embedded in paraffin, and brain sections were mounted on glass
slides and stained with hematoxylin and eosin (H&E) using standard methods. For
immunohistochemical (IHC) staining of mouse hemibrains, staining was conducted on the
Leica Bond RXm platform using standard chromogenic methods. For antibodies requiring
heat induced epitope retrieval (HIER), slides were heated in a pH9
ethylenediaminetetraacetic acid (EDTA)-based buffer at 94 °C for 25 min.

Antibody Information.

Target	Host	Clone	Vendor	Cat #
Iba1	Rabbit	EPR16588	AbCam	ab178846PUR
GFAP	Rabbit	EPR1034Y	AbCam	ab68428
Aβ	Rabbit	D12B2	Cell Signaling	9888
CD3	Rabbit	SP7	AbCam	ab16669

179   **Staining Parameters.**

Antibody	Antigen Retrieval	Dilution	Chromogen

Iba1	pH9	1:40,000	DAB
GFAP	pH9	1:200	DAB
Aβ	pH9	1:200	DAB
CD3	pH9	1:100	DAB

Aβ Image Analysis

IHC-stained glass slides (n = 62) labelled for Aβ were scanned using an Aperio AT2 whole
slide scanner with a maximum objective of 20x. Image analysis was conducted on Aβ IHC-
stained digital slide images using Visiopharm® software. Six coronal sections of the brain
per animal (across 2 slides) were annotated manually as individual regions of interest
(ROIs): rostral 1 and 2, middle 1 and 2, and caudal 1 and 2. IHC-positive area and total area
within each ROI were used to calculate percent positive area for Aβ immunolabeling. The
calculations for percentage positive area (%) are provided below:

$$Percent\ Positive\ Area\ (\%) = Positive\ Area\ (\mu m2)/Total\ Area\ (\mu m2)\ x\ 100$$

For further analysis a deep learning algorithm trained to detect tissue was run on each slide
to define the region of interest (ROI). Deep learning algorithms (DLA) use convolutional
neural networks (CNNs) that can be trained to identify specific objects within the tissue
based on morphological patterns, RGB signatures, texture, size, and spatial relationships.
This algorithm was developed using the U-Net model. A second DLA trained to detect
vascular endothelium trained on slides stained with β-amyloid IHC was used to define
vessels. These vessels were then outlined into a separate ROI. β-amyloid immunolabeling
was detected using a final DLA. β-amyloid signal was assessed as total, plaque and non-
plaque. Total β-amyloid burden was defined by the sum of the areas of plaques and non-
plaques. For the frontal cortex segmentation, data were reported as percent area of total β-
amyloid, percent area of plaques, number of plaques, and average plaque size. These data
were reported from the total brain area, including all 6 sections of the brain including
vasculature, as well as frontal cortex as defined by the Allen Brain Atlas. Three of the 6
sections had frontal cortex present.

CD3, GFAP, and IBA1 Image Analysis

CD3, GFAP, and IBA1 immunolabeling were detected using thresholding techniques. All
three thresholds were developed individually based on Visiopharm's proprietary HDAB filter,
which isolates the specific intensity and colorimetric signature associated with positive DAB-

based immunolabeling. Total brain area was defined as the sum of the non-frontal cortex
brain area and the frontal cortex. CD3 quantification is reported as positive cell density/mm2;
GFAP and IBA1 quantification are reported as positive percentage area.
214

Statistical Analyses
Statistical analyses were performed on the data generated from this study using GraphPad
Prism 10.03 for Windows (GraphPad Software, Inc.). Details of each analysis can be found
in the figure legend presenting the data. Statistical significance was set at $p < 0.05$.
Considering the sample size of our groups, data showing a statistical trend are also
presented in this study.
221

Results

223 **Unique BBB disruption changes the molecular inflammatory signaling profile for up to**
224 **15 days post-sonication**

225 To better understand the effects of mechanical opening of the BBB by US/MB, we
226 investigated blood-based markers and inflammatory markers following a single session of
227 ultrasound to disrupt the BBB (experimental plan, Fig 1A).
228

No significant changes in body weight were observed, however with a trend toward weight
loss in the first 24h post treatment (ANOVA p=0.06277, D0 vs D1 p=0.0023) (Fig S1A).
Treated mice showed a trend towards an increased in red blood cell count 15 days post
opening (MEA p=0.0773, group1 vs Group4 p=0.0366) (Fig 1B), however no significant
effect was recorded on the number of circulating white cells (Fig 1C). We used the S100b
marker to monitor the opening of the BBB post procedure and its level overtime. As
expected, we observed a drop of the S100b level over time reaching the lowest level 8 days
after BBB opening (LS mean = 16.00), becoming significant only at 15d post sonication (LS
mean 16.41) (MEA p=0.0063, post sonication vs terminal 15 days p= 0.0115) (Fig S1B,
measured by ELISA).
239

Once the safety of BBB opening was established, we performed RT-Q-PCR to explore the
level of different inflammatory markers in the brain at several time points [2-, 4-, 8- and 15-
days post opening] post BBB opening. Interestingly, the only significant increase compared
to baseline, group 0 without BBB opening, was an increase in *Il-6* mRNA levels, (MEA
p<0.0001, Group 0 vs 1: p=0.0065, Group 0 vs 3 p=0.0003, Group 0 vs 4 p<0.0001). It is

important to note that *Il-6* increase was correlated with an increase of red blood cells (Fig
1B)[45]. Notably, all the other factors measured were decreased, either transiently, or with a
prolonged effect, up to 15 days post treatment.

Inflammatory chemokines *Cxcl1* and *Ccl2* as well as inflammatory cytokines *Il-1b*, *Il-10* and
Tnf-a were decreased. While most of the chemokines decreased over 15 days post
treatment, *Tnf-a* was only decreased in the first 48h of the protocol, with a recovery to
baseline by day 4. Adhesion molecules *Icam1* and *E-selectin* were also decreased, with a
recovery of *E-selectin* by day 15. The growth factors *Tgfb* and *Bdnf* were found to be
significantly reduced after a single sonication. Finally, the transcription factor *Nfkb* was
significantly decreased post sonication. Altogether, these results suggested that the opening
of the BBB with the selected parameters did not lead to a pro-inflammatory response, or any
significant changes to other physiological metrics in a healthy mouse model.

Different schedules of repeated BBB opening did not impact physiological output

Next, the effect of different sonication protocols was explored in a mouse model for AD that
develops amyloid deposition (ARTE10). We tested different regimens of US/MB mediated
BBB opening going from twice a week to once a month with sacrifice and profiling of different
markers at one week or one month after the last sonication (Fig.2A). We first investigated
the effect of repeated sonication on the general health and survival of the mice. Our data
showed no statistical difference between the groups (MEA, p=0.6513), with however a
significant difference between timepoints (MEA, p=0.0459, D21 vs D28 p=0.0280), as it was
described for the WT cohort. (Fig. 2B). A few mice had weight reduction following the first
session of the sonication but recovered in the following week(s).

Overall, our data suggests that LIPU-mediated BBB opening is a safe procedure that allows
repetitive treatment. Altogether, these data demonstrated the low level of toxicity of a
repeated protocol of sonication, as well as the structural plasticity of the BBB allowing
repeated mechanical disruption of its structure.

Sonication frequency does not trigger inflammation in the brain over time

Using the post-mortem tissues from the ARTE10 cohort, we then investigated the effect of
repeated BBB opening on inflammatory markers of the brain at the cellular and the
molecular level. CD3, GFAP and Iba1 markers were used to explore T-cell infiltration, gliosis
and microglial activation respectively (Fig. 3A). We used an automated pipeline of analysis,
able to recognize the cells/markers of interest to extract the number of cells per mm^2 or the
percentage of occupancy (see Supp Fig2. A-C showing the quantification mask for the 3

markers). The obtained results showed that repeated sonication did not lead to an increase
of CD3+ cell infiltration (Fig3B and C). A statistical trend toward an increase in astrogliosis
was observed between Group 1 and Group 3 (ANOVA p<0.0001, Group 1 vs Group 3
P=0.0918) (Fig3D and F), as previously reported[46] and consistent with the level of S100b
observed. In all the other conditions, as well as for all the conditions for Iba1, no difference
between groups was detected (Fig3D-F). A few CD3+ cells were observed in the vicinity of
amyloid deposits. As expected, GFAP+ and Iba+ cells tend to accumulate around the
amyloid deposits. However, to explore how sonication stimulates the relocalization of
inflammatory cells around the pathological aggregates, to possibly participate in their
elimination, more in-depth studies will be required, to explore the cellular mechanisms
triggered by LIPU-mediated BBB opening.
291
We generated pilot results on the mRNA expression of few inflammatory markers (as
explored in Fig. 1). Interestingly, our data showed a decrease in the proinflammatory
cytokines in all conditions, as was observed in wildtype mice (Fig. 1). Opposite to what was
observed in the single BBB opening protocol, repeated opening of the BBB was associated
with an increase of *bdnf* in most treatments, however not statistically tested due to limited
number of data (Supp fig 2D). This brings the possibility that the repeated mechanical
opening of the BBB could favor the local release of growth factors, as it was observed in
other studies[36–38].These data would require further investigation, but they provide an
interesting first look at the molecular changes occurring during a repeated sonication
regimen, which could provide additional benefits for the patients.

Monthly repeated sonication shows promising results on pathological hallmarks
Finally, we assessed the effect of repeated sonication on amyloid deposits in the ARTE10
model (Fig4A). The ARTE10 mouse model is characterized by the presence of amyloid and
mineralized deposits in the brain (Supp fig 4A), a noticeable loss of hippocampal neurons
(Suppl Fig4B), and the presence of amyloid deposition around the brain vasculature (Suppl
Fig3C). Consistently across all analyses, our data showed a significant reduction of the
amyloid load in the very sparse sonication regimen (Group 6), consisting of a monthly
sonication, compared to the other sonication regimens (Group 3 and 4, respectively
p=0.0216 and 0.0437), with more regular sonication regimen. While not significant, it is worth
noting that Group 6 also showed a slight reduction in the plaque area distribution compared
to the non-sonicated control (Group 6 vs Group1, p=0.019, q=0.0385). Indeed, if analyzed
separately, the average size of amyloid deposits was significantly reduced, while the
percentage of amyloid positive area showed a tendency to decrease under the monthly
sonication regimen (Supp Fig4D, p=0.1143, q=0.1154). The treatment did not have any

effect on the overall number of deposits. Taken together, these data confirm the results
previously shown that ultrasound alone can have beneficial effects in the treatment of
Alzheimer's pathology.

Discussion

Altogether, this short pilot study showed that LIPU is a safe method to open the BBB,
allowing repeated use with potential benefits in a pathological context. We previously
demonstrated in a small cohort of nine patients with mild AD that seven repeated, bi-monthly
LIPU-mediated BBB openings reduced the amyloid burden by 6.6% in the brain region
targeted by the implantable ultrasound system used in the study[44]. The results from this
preclinical study are in agreement and point to a modest effect of ultrasound alone on
amyloid pathology. Further analyses of the different types of amyloid aggregates showed
that repeated sonication triggers a reduction in the amyloid deposit size and amyloid
distribution in the brain. These results are encouraging and point out the potential for
ultrasound-mediated BBB opening as a therapeutic tool by itself. Further research should
focus on finding the best synergistic effects between a specific therapeutic compound and
an optimized sonication regimen to improve the outcome for patients.
332
Several studies have already explored the effect of sonication on amyloid pathology in AD
models (see tables 4 and 5). However, the experimental designs between these different
studies, including this one, make it complicated to grasp the real value of sonication in this
pathology. In comparison to these models, ARTE10 model presents an amyloid deposition at
3 months, synaptic loss at 3 months, and a cognitive impairment at 12 months[47]. This
model presents similarities with the TgCRND8 models. However, in the previously published
studies, the animals were treated at a younger age (4.5mo in Jordao et al, 2010, 4mo in
Jordao et al 2013, 6 to 8 mo in Burgess et al, 2014a and b, 7 mo in Poon et al, 2018). Our
study explored the effect of repeated sonication in 1yo mice, where the amyloid deposition is
already highly established and with a heavy load of amyloid deposits. 3TG and APP/PS1dE9
models were used at later stages, but the rate of amyloid deposition appears slower than in
the ARTE10 mouse model (8mo in Shen et al, 2020; 2.5mo, 1yo and 2yo in Noel et al, 2023
and 14mo in Karakatsani et al, 2023, 12 and 14 mo in Hsu et al, 2018 and 16, 24 and 30 mo
in Sun et al, 2021). To focus on comparable studies, we will only consider studies which
used repeated sonication protocol to show the effect on the amyloid pathology[22,24,27–
30,35]. Despite these experimental differences, similar effects were recorded, including our
study: 1) Repeated sonication consistently triggers a decrease in the amyloid load across all

models[22,24,27–30,35]; 2) A significant improvement of the cognitive
performance[22,24,27,30], 3) An increased microglial activity around the deposits[22,29].

Our study demonstrated that in both a wildtype and a mouse model for AD with amyloid
pathology (ARTE10), repeated LIPU-mediated BBB opening does not affect the general
physiology of the mouse. However, we observed that a single sonication led to an increase
of *Il-6*, while most of the other factors were either unchanged or slightly decreased (Fig.1D).
Il-6 is classically described as a pro-inflammatory cytokine with several key functions in the
central nervous system but also shows anti-inflammatory functions in specific contexts[48].
Il-6 has been shown to play a major role in activating microglia and favoring
microvasculature repair after injury[49]. In brain diseases, *Il-6* expression was found to be
increased in traumatic brain injury (TBI) as well as some initial steps of Alzheimer's
disease[50–52], however with conflicting associated functions. While the increase of *Il-6*
levels after TBI was associated with anti-inflammatory and pro-survival effects, sustained
elevated levels of *Il-6* were shown to be detrimental by increasing neuronal excitability
leading to potential seizures[48]. Moreover, *Il-6* stimulates angiogenesis following a stroke
episode, highlighting its potent role in stimulating vasculature repair after vascular
injuries[53]. In addition, *Il-6* is known to increase in response to tissue injuries and to induce
hematopoiesis (increase in red blood cell, Fig1B)[45]. In our experiments, we hypothesized
that the increase in *Il-6* was tightly correlated to the vasculature repair process, without
triggering a pathological chronic inflammatory response. Indeed, other cytokines and
chemokines were unchanged or decreased after BBB opening, pointing out the unique role
of *Il-6* in this context.

Other factors like the *Tnf-a*, another known partner involved in pathogenesis of AD[54],
showed a downregulation post BBB-opening. Interestingly, in the context of TBI, *Il-6* inhibits
the production of *Tnf-a* as an anti-inflammatory process, as observed in our data[48]. *Tnf-a*
plays a central role in neuroinflammation[55], which reinforces the observation that LIPU-
mediated BBB opening does not trigger pathological inflammation. Similarly to *Tnf-a*, *Il-1b* is
associated with a myriad of negative effects on the brain, in regard to seizures, a role in the
development of different psychiatric disorders and neurodegenerative diseases[56–58].
More importantly for the BBB, the inhibition of the *Il-1b* pathway showed a positive effect on
the brain vascularization[59]. Counter intuitively, we observed a decrease in level of *Il-10*, an
anti-inflammatory cytokine of the brain, with important roles in the repair process of the
brain[60] and limiting pathological inflammation[61]. This suggests that LIPU-mediated BBB-
opening is not a major traumatic event in the brain and only triggers transient changes in
inflammatory markers in the brain. Our results showed a transitory effect on astrocytic

activation (Fig. 3F). These results are in accordance with recent publications, showing a
transient activation of astrocytes and microglia in the first hours post sonication, with a
normalization to baseline after few days[46,62–64].

In this direction, the decrease in *nfkb* expression confirms the minimal impact of our BBB
opening strategy on neuroinflammation[65,66]. Similarly, Ccl2, a factor known for disrupting
the BBB integrity[67–69], as well as favorizing the infiltration of leukocytes and monocytes to
the brain parenchyma[67,70], was found decreased post sonication, reinforcing the idea that
LIPU-mediated BBB opening is a mild disruption of the BBB itself, and does not trigger
pathways observed in stroke- or infection-mediated BBB disruptions. Our study also showed
a transient decrease in surface adhesion molecule expression *Icam* and *E-Selectin*. *E-*
Selectin levels were recovered 8d post opening, while *Icam* took longer to recover. In a
recent publication, it was shown that repeated sonication schedules could be favorable for
the vasculature, by stimulating higher level of adhesion molecule VE-Cadherin [71]. To really
understand the effect of repeated sonication on the vasculature, especially in the context of
neurodegenerative diseases, more in-depth studies will be required in the future, using high
spatial resolution techniques such as in vivo 2-photon microscopy and spatial
transcriptomics and proteomics. These studies could bring evidence of the beneficial effect
of sonication in stimulating the repair process of the vasculature, and reperfusion of the
brain[62].

Our study offers a unique timeline into the changes happening post BBB opening. Several
studies attempted similar quantifications, however with a shorter timeline going up to 24h
post sonication, versus 15 days in our study[72–74]. The data obtained 24h post
sonication[72,73] showed a significant decrease in the magnitude of changes observed in
the different factors, reaching similar amplitudes as what was observed in our study (0.5 to
1-fold change). Compared to these studies, our data showed that there are no negative long-
term changes in induced neuroinflammation post sonication, confirming the normalizing level
of all analyzed cytokines and chemokines in the previous studies. These results are a
testimony to the transitory aspect of the US/MB-mediated BBB opening. It is however
interesting to compare the data generated in our wildtype cohort, where a single BBB
opening was performed, to the data obtained in our ARTE10 cohort, where different
repeated sonication regimens were applied. Two main differences were observed between
the two cohorts: The absence of *Il-6* increase and increase in *bdnf* expression in the
repeated sonication cohort compared to the single sonication cohort. Increased *bdnf* was
already observed post sonication[72] and suggested as the underlying mechanism
responsible for stimulating neurogenesis in other studies[27,36–38]. For example, in Kovacs

et al., *bdnf* levels were observed to increase around 24h post sonication. Our study showed that this increase might be transient, as we did not observe a detectable increase at 48h post-sonication or could be an effect of the much stronger sonication parameters used to disrupt the BBB in the Kovacs et al. study. However, as we observed an increase of *bdnf* levels after repeated sonication, it suggests that regular opening of the BBB could allow sustained levels of BDNF overtime, which could in turn have a beneficial effect, potentially through increased neurogenesis or higher neuronal excitability and synaptic homeostasis[75]. However, as observed for groups 3 and 4, higher levels of BDNF did not correlated with decrease of amyloid pathology. It is now well documented that neuronal activity correlates with amyloid beta secretion in the intercellular spaces[76–78]. These results would suggest that LIPU-mediated BBB opening might efficiently regulate amyloid beta pathology within a certain range of stimulation.

Cerebral amyloid angiopathy (CAA), characterized by the amyloid deposition around the blood vessel, is found in most patients presenting Alzheimer's disease. The accumulation at the vessel is believed to originate from incomplete clearance of the amyloid from the brain. Previous work has established that accumulation of amyloid in the blood vessels progressively replaces all the different tissues forming the arterial wall[79]. Accumulation of amyloid in the arterial wall affects blood vessel integrity and function, resulting in cerebral infarct or hemorrhage[80]. Post sonication, it was shown that vessels containing amyloid were less susceptible to diameter changes during US/MB-mediated BBB opening protocol, suggesting a more rigid structure. In addition, in a mouse model for AD, small vessels were most likely to display leakage post BBB opening in their littermate controls[26]. Unfortunately, the presence of CAA in patients with AD could be the reason for ineffective anti-amyloid immunotherapy and observed adverse events leading to microhemorrhage in patients[81,82]. In our study, the ARTE10 model also presents some level of CAA, with notable accumulation of amyloid positive staining around the blood vessels (Supp Fig. 3C, Supp fig.4). Targeted analysis did not show any beneficial effect of repeated sonication on the vascular amyloid (Fig 4). However, it is important to note that postmortem brain imaging did not reveal the presence of increased hemorrhage following sonication. This suggests that sonication can still be a safe and potent strategy to efficiently deliver therapies against amyloid to the brain [23]. As mentioned earlier, few studies have already explored this possibility[4,31–33,35], including during a clinical trial[15]. All studies reported a beneficial effect of therapeutic delivery to the brain and a reduction of the pathology in the different models used (See tables 1 and 2).

Conclusion

Altogether, this study brings additional evidence to the field that repeated sonication to disrupt the BBB is a safe procedure, which could bring positive therapeutic effects in the context of neurodegenerative diseases by both enhancing drug delivery and stimulating mild, transient inflammatory responses. More in-depth analyses are required to understand the molecular and cellular mechanisms at play. However, it is worth considering the potential therapeutic effect of ultrasound mediated BBB opening in the future, and to explore the synergic effects of their combination to disease modifying therapeutics, which could lead to improved outcomes for patients.

Acknowledgements

The authors acknowledge Inotiv (former Bolder BioPATH) for their work on this study as well as Dallas Tissue Research for their help with image analysis. We want to thank Prof. Benoit Delatour for his input on the paper. This study was fully funded by Carthera (Lyon, France).

Conflict of interest

M.C., G.B., and P.D.R. are employees of Carthera SA. A.C. is the founder of Carthera SA. All authors have approved the submitted and revised versions of the manuscript.

Contributions

M.C. and A.C. designed and supervised the study. M.C. and G.B. developed the preclinical set-up for BBB opening, G.B. provided technical expertise. P.D.R. performed data curation, formal analyses and statistical analyses. M.C. and P.D.R prepared the original manuscript and the revised manuscript. All authors attest that no generative AI or AI-assisted technologies were used in the writing process.

Data Availability statement

Data presented in this paper are available on request to the corresponding author.

488   **Table 4: List of the previous mouse models used in combination with Ultrasound.**
489   **Four models have been used to explore the effect of sonication on the amyloid**
490   **pathology:**
491

Mouse model	Strain characteristics	Pathological hallmarks	Publications
3TG model	Mix model for amyloid and tau pathologies Single mutations in APP, MAPT, PSEN2	• Cognitive impairment at 4 months, • LTP deficits at 6 months, • Amyloid deposition after 6 months, • Tau pathology after 1 year	[22–24,83]
TgCRND8 model	Pure amyloid model Double mutation in APP	• Amyloid deposition at 3 months • Associated with cognitive decline • Changes in LTP at 6 months	[4,25–28,84]
5xFAD model	Pure amyloid model Triple mutation inAPP and double mutation in PSEN1	• Early amyloid deposition around 2 months • Synaptic loss after 4 months • Associated with cognitive impairments • Changes in LTP and neuronal loss at 6 months	*[29,30,85]*
APP/PS1dE9 model	Pure amyloid model Single mutations in APP and PSEN1	• Early LTP impairment at 3 months • Synaptic loss at 4 months • Amyloid deposition at 6 months • Cognitive impairment around 1 year	*[33,35,86]*

493 **Table 5: Previous studies using ultrasound in mouse models for Alzheimer's Disease**
494 **and their output.**

Study	Model used	Age of the model	Sonication treatment	Associated therapeutics	Effect on the pathology	Effect on behavior
Shen et al, 2020[22]	3TG	8 mo	Repeated, twice a week for six weeks	none	• Decreased amyloid burden in Cortex, CA and amygdala • Reduced Tau phosphorylation • Reduced branching and process length in microglia • Axonal protection	Improvement in Y maze, Morris water maze and step down avoidance test
Noel et al, 2023[23]	3TG	2.5mo 1 yo 2 yo	One treatment	none	Opening volume slightly higher in 3TG compared to WT	NA
Karakatsani et al, 2023[24]	3TG	14 mo	Repeated, once a week for four weeks	none	>50% reduction in number and 42% in size	Improvement of memory (Morris water maze)
Jordao et al, 2013[25]	TgCRND8	4 mo	One treatment	none	• Reduced deposits in cortex • IgG & IgM to penetrate infiltration • Increased in Iba1 and GFAP • Increased distal localization of microglia and astrocytes • Increased neuronal area • Increased colocalization between glial cells and amyloid deposits	NA
Burgess et al, 2014a[27]	TgCRND8	7 mo	Repeated, once a week for 3 weeks, bilateral	none	• Decreased in deposit size (~20%) and deposit number (~19%) • Increased in DCX positive neurons (188% for nTG, 252% for Tg) • Increased in dendrite length and branching	Improvement at the Y maze
Burgess et al, 2014b[26]	TgCRND8	6-8 mo	One treatment	none	• Correlation of vessel diameter and leakage probability • Vessels displaying leakage are significantly smaller in Tg mice • Vessels coated with amyloid show less	NA

					changes in diameter	
Poon et al, 2017[28]	TgCRND8	7 mo	Repeated, once a week for 5 weeks	none	• Decreased number of amyloid deposits • Decrease amyloid deposit area	NA
Bobola et al, 2020[29]	5xFAD	6 mo	Repeated, daily for five days	none	• Decreased amyloid burden by ~50% compared to control • Increased microglia colocalization with amyloid deposits compared to control	NA
Lee et al, 2020[30]	5xFAD	4 mo	Repeated, once a week for six weeks	none	• Decrease of both amyloid deposits area and deposit area • Increased CSF abeta	Improvement in working memory
Jordao et al, 2010[4]	TgCRND8	4.5 mo	One treatment	BAM10 antibody	Slight decrease in Abeta deposits	NA
Alecou et al, 2017[31]	Rabbit, high cholesterol diet	Not disclosed	Repeated, three treatments	BC-10 (sigma), anti Amyloid-beta antibody	Original amyloid deposit density 200/cm2, decreased to 78/cm2 after treatment	NA
Hsu et al, 2018[35]	APP/PS1dE9	12-14 mo	Repeated, once a week for five weeks	GSK-3 Inhibitor	Significant decrease of the deposit size in hippocampus, including with FUS alone	NA
Sun et al, 2021[33]	APP/PS1dE9	16 mo, 24-30 mo	Three treatments for 16mo cohort, one treatment for 24-30mo cohort	anti-pGlu3 (07/2a)	• Decreased hippocampal plaque deposition in APP/PS1dE9 mice co-treated with anti-pGlu3 Aβ mAb and FUS-BBBD • Age-related hippocampal synaptic degeneration is reduced in APP/PS1dE9 mice after treatment with 07/2a and FUS-BBBD • Altered plaque-associated microglia and macrophage activation in aged APP/PS1dE9 after treatment with 07/2a and FUS-BBBD	No significant effects in fear learning and memory or locomotor activity with anti-pGlu3 Aβ mAb or FUS-BBBD
Deng et al, 2021[34]	APP/PS1dE9	10 mo	One treatment	IV injection of US-HA-Exo	Decreased deposit area and occupancy	NA
Antoniou et al, 2024[32]	5xFAD	5 mo	One treatment	Amyloid Protein (1–40) antibody produced in rabbit whole antiserum, A8326	Allow for non-invasive antibody delivery to amyloid deposits	NA

References

[1]   Yadollahikhales G, Rojas JC. Anti-Amyloid Immunotherapies for Alzheimer's Disease: A 2023 Clinical Update. Neurotherapeutics 2023;20:914. https://doi.org/10.1007/S13311-023-01405-0.

[2]   Zhao P, Zhang N, An Z. Engineering antibody and protein therapeutics to cross the blood–brain barrier. Antib Ther 2022;5:311. https://doi.org/10.1093/ABT/TBAC028.

[3]   Trontinemab Data Strengthen Hope for Brain Shuttles | ALZFORUM n.d. https://www.alzforum.org/news/conference-coverage/trontinemab-data-strengthen-hope-brain-shuttles (accessed February 24, 2025).

[4]   Jordão JF, Ayala-Grosso CA, Markham K, Huang Y, Chopra R, McLaurin JA, et al. Antibodies Targeted to the Brain with Image-Guided Focused Ultrasound Reduces Amyloid-β Plaque Load in the TgCRND8 Mouse Model of Alzheimer's Disease. PLoS One 2010;5:e10549. https://doi.org/10.1371/JOURNAL.PONE.0010549.

[5]   Burgess A, Huang Y, Querbes W, Sah DW, Hynynen K. Focused ultrasound for targeted delivery of siRNA and efficient knockdown of Htt expression. Journal of Controlled Release 2012;163:125–9. https://doi.org/10.1016/J.JCONREL.2012.08.012.

[6]   Park J, Aryal M, Vykhodtseva N, Zhang YZ, McDannold N. Evaluation of permeability, doxorubicin delivery, and drug retention in a rat brain tumor model after ultrasound-induced blood-tumor barrier disruption. Journal of Controlled Release 2017;250:77–85. https://doi.org/10.1016/j.jconrel.2016.10.011.

[7]   Nisbet RM, Van Der Jeugd A, Leinenga G, Evans HT, Janowicz PW, Götz J. Combined effects of scanning ultrasound and a tau-specific single chain antibody in a tau transgenic mouse model. Brain 2017;140:1220–30. https://doi.org/10.1093/BRAIN/AWX052.

[8]   Karakatsani ME, Kugelman T, Ji R, Murillo M, Wang S, Niimi Y, et al. Unilateral Focused Ultrasound-Induced Blood-Brain Barrier Opening Reduces Phosphorylated Tau from The rTg4510 Mouse Model. Theranostics 2019. https://doi.org/10.7150/thno.28717.

[9]   Janowicz PW, Leinenga G, Götz J, Nisbet RM. Ultrasound-mediated blood-brain barrier opening enhances delivery of therapeutically relevant formats of a tau-specific antibody. Sci Rep 2019. https://doi.org/10.1038/s41598-019-45577-2.

[10]   Meng Y, Pople CB, Huang Y, Jones RM, Ottoy J, Goubran M, et al. Putaminal Recombinant Glucocerebrosidase Delivery with Magnetic Resonance–Guided Focused Ultrasound in Parkinson's Disease: A Phase I Study. Movement Disorders 2022;37:2134–9. https://doi.org/10.1002/MDS.29190.

[11]   Trinh D, Nash J, Goertz D, Hynynen K, Bulner S, Iqbal U, et al. Microbubble drug conjugate and focused ultrasound blood brain barrier delivery of AAV-2 SIRT-3. Drug Deliv 2022;29:1176–83. https://doi.org/10.1080/10717544.2022.2035855.

[12]   Blesa J, Pineda-Pardo JA, Inoue KI, Gasca-Salas C, Balzano T, López-González Del Rey N, et al. BBB opening with focused ultrasound in nonhuman primates and Parkinson's disease patients: Targeted AAV vector delivery and PET imaging. Sci Adv 2023;9. https://doi.org/10.1126/sciadv.adf4888.

[13]   Shen Y, Zhang J, Xu Y, Sun S, Chen K, Chen S, et al. Ultrasound–enhanced brain delivery of edaravone provides additive amelioration on disease progression in an

ALS mouse model. Brain Stimul 2023;16:628–41.
https://doi.org/10.1016/J.BRS.2023.03.006.

[14] Arrieta VA, Gould A, Kim KS, Habashy KJ, Dmello C, Vázquez-Cervantes GI, et al.
Ultrasound-mediated delivery of doxorubicin to the brain results in immune modulation
and improved responses to PD-1 blockade in gliomas. Nature Communications 2024
15:1 2024;15:1–19. https://doi.org/10.1038/s41467-024-48326-w.

[15] Rezai AR, D'Haese P-F, Finomore V, Carpenter J, Ranjan M, Wilhelmsen K, et al.
Ultrasound Blood–Brain Barrier Opening and Aducanumab in Alzheimer's Disease.
New England Journal of Medicine 2024;390:55–62.
https://doi.org/10.1056/NEJMOA2308719/SUPPL_FILE/NEJMOA2308719_DATA-
SHARING.PDF.

[16] Chen H, Yang GZX, Getachew H, Acosta C, Sierra Sánchez C, Konofagou EE.
Focused ultrasound-enhanced intranasal brain delivery of brain-derived neurotrophic
factor. Scientific Reports 2016 6:1 2016;6:1–8. https://doi.org/10.1038/srep28599.

[17] Stavarache MA, Petersen N, Jurgens EM, Milstein ER, Rosenfeld ZB, Ballon DJ, et al.
Safe and stable noninvasive focal gene delivery to the mammalian brain following
focused ultrasound. J Neurosurg 2019;130:989–98.
https://doi.org/10.3171/2017.8.JNS17790.

[18] Hynynen K, McDannold N, Vykhodtseva N, Jolesz FA. Noninvasive MR Imaging–
guided Focal Opening of the Blood-Brain Barrier in Rabbits1.
Https://DoiOrg/101148/Radiol2202001804 2001;220:640–6.
https://doi.org/10.1148/RADIOL.2202001804.

[19] Beccaria K, Canney M, Goldwirt L, Fernandez C, Adam C, Piquet J, et al. Opening of
the blood-brain barrier with an unfocused ultrasound device in rabbits. J Neurosurg
2013;119:887–98. https://doi.org/10.3171/2013.5.JNS122374.

[20] Horodyckid C, Canney M, Vignot A, Boisgard R, Drier A, Huberfeld G, et al. Safe
long-term repeated disruption of the blood-brain barrier using an implantable
ultrasound device: a multiparametric study in a primate model. J Neurosurg
2017;126:1351–61. https://doi.org/10.3171/2016.3.JNS151635.

[21] Sabbagh A, Beccaria K, Ling X, Marisetty A, Ott M, Caruso H, et al. Opening of the
blood–brain barrier using low-intensity pulsed ultrasound enhances responses to
immunotherapy in preclinical glioma models. Clinical Cancer Research
2021;27:4325–37. https://doi.org/10.1158/1078-0432.CCR-20-
3760/672085/AM/OPENING-OF-THE-BLOOD-BRAIN-BARRIER-USING-LOW.

[22] Shen Y, Hua L, Yeh CK, Shen L, Ying M, Zhang Z, et al. Ultrasound with
microbubbles improves memory, ameliorates pathology and modulates hippocampal
proteomic changes in a triple transgenic mouse model of Alzheimer's disease.
Theranostics 2020;10:11794–819. https://doi.org/10.7150/THNO.44152.

[23] Noel RL, Gorman SL, Batts AJ, Konofagou EE. Getting ahead of Alzheimer's disease:
early intervention with focused ultrasound. Front Neurosci 2023;17.
https://doi.org/10.3389/fnins.2023.1229683.

[24] Karakatsani ME, Ji R, Murillo MF, Kugelman T, Kwon N, Lao YH, et al. Focused
ultrasound mitigates pathology and improves spatial memory in Alzheimer's mice and
patients. Theranostics 2023;13:4102–20. https://doi.org/10.7150/THNO.79898.

[25] Jordão JF, Thévenot E, Markham-Coultes K, Scarcelli T, Weng YQ, Xhima K, et al.
Amyloid-β plaque reduction, endogenous antibody delivery and glial activation by
brain-targeted, transcranial focused ultrasound. Exp Neurol 2013;248:16–29.
https://doi.org/10.1016/J.EXPNEUROL.2013.05.008.

[26] Burgess A, Nhan T, Moffatt C, Klibanov AL, Hynynen K. Analysis of focused
ultrasound-induced blood-brain barrier permeability in a mouse model of Alzheimer's
disease using two-photon microscopy. Journal of Controlled Release 2014;192:243–
8. https://doi.org/10.1016/j.jconrel.2014.07.051.

[27] Burgess A, Dubey S, Yeung S, Hough O, Eterman N, Aubert I, et al. Alzheimer
Disease in a Mouse Model: MR Imaging–guided Focused Ultrasound Targeted to the
Hippocampus Opens the Blood-Brain Barrier and Improves Pathologic Abnormalities
and Behavior. Https://DoiOrg/101148/Radiol14140245 2014;273:736–45.
https://doi.org/10.1148/RADIOL.14140245.

[28] Poon C, McMahon D, Hynynen K. Noninvasive and targeted delivery of therapeutics
to the brain using focused ultrasound. Neuropharmacology 2017;120:20–37.

[29] Bobola MS, Chen L, Ezeokeke CK, Olmstead TA, Nguyen C, Sahota A, et al.
Transcranial focused ultrasound, pulsed at 40 Hz, activates microglia acutely and
reduces Aβ load chronically, as demonstrated in vivo. Brain Stimul 2020;13:1014–23.
https://doi.org/10.1016/J.BRS.2020.03.016.

[30] Lee Y, Choi Y, Park EJ, Kwon S, Kim H, Lee JY, et al. Improvement of glymphatic-
lymphatic drainage of beta-amyloid by focused ultrasound in Alzheimer's disease
model. Sci Rep 2020;10. https://doi.org/10.1038/S41598-020-73151-8.

[31] Alecou T, Giannakou M, Damianou C. Amyloid β Plaque Reduction With Antibodies
Crossing the Blood-Brain Barrier, Which Was Opened in 3 Sessions of Focused
Ultrasound in a Rabbit Model. Journal of Ultrasound in Medicine 2017:n/a--n/a.
https://doi.org/10.1002/jum.14256.

[32] Antoniou A, Stavrou M, Evripidou N, Georgiou E, Kousiappa I, Koupparis A, et al.
FUS-mediated blood–brain barrier disruption for delivering anti-Aβ antibodies in
5XFAD Alzheimer's disease mice. J Ultrasound 2024;27:251–62.
https://doi.org/10.1007/S40477-023-00805-4/FIGURES/7.

[33] Sun T, Shi Q, Zhang Y, Power C, Hoesch C, Antonelli S, et al. Focused ultrasound
with anti-pGlu3 Aβ enhances efficacy in Alzheimer's disease-like mice via recruitment
of peripheral immune cells. J Control Release 2021;336:443–56.
https://doi.org/10.1016/J.JCONREL.2021.06.037.

[34] Deng Z, Wang J, Xiao Y, Li F, Niu L, Liu X, et al. Ultrasound-mediated augmented
exosome release from astrocytes alleviates amyloid-β-induced neurotoxicity.
Theranostics 2021;11:4351–62. https://doi.org/10.7150/THNO.52436.

[35] Hsu PH, Lin YT, Chung YH, Lin KJ, Yang LY, Yen TC, et al. Focused Ultrasound-
Induced Blood-Brain Barrier Opening Enhances GSK-3 Inhibitor Delivery for Amyloid-
Beta Plaque Reduction. Scientific Reports 2018 8:1 2018;8:1–9.
https://doi.org/10.1038/s41598-018-31071-8.

[36] Scarcelli T, Jordão JF, O'reilly MA, Ellens N, Hynynen K, Aubert I. Stimulation of
Hippocampal Neurogenesis by Transcranial Focused Ultrasound and Microbubbles in
Adult Mice. Brain Stimul 2014;7:304–7. https://doi.org/10.1016/J.BRS.2013.12.012.

[37] Mooney SJ, Shah K, Yeung S, Burgess A, Aubert I, Hynynen K. Focused Ultrasound-
Induced Neurogenesis Requires an Increase in Blood-Brain Barrier Permeability.
PLoS One 2016;11:e0159892. https://doi.org/10.1371/JOURNAL.PONE.0159892.

[38] Mooney SJ, Nobrega JN, Levitt AJ, Hynynen K. Antidepressant effects of focused
ultrasound induced blood-brain-barrier opening. Behavioural Brain Research
2018;342:57–61. https://doi.org/10.1016/j.bbr.2018.01.004.

[39] Pandit R, Leinenga G, Götz J. Repeated ultrasound treatment of tau transgenic mice
clears neuronal tau by autophagy and improves behavioral functions. Theranostics
2019;9:3754–67. https://doi.org/10.7150/THNO.34388.
[40] Géraudie A, Riche M, Lestra T, Trotier A, Dupuis L, Mathon B, et al. Effects of Low-
Intensity Pulsed Ultrasound-Induced Blood-Brain Barrier Opening in P301S Mice
Modeling Alzheimer's Disease Tauopathies. Int J Mol Sci 2023;24.
https://doi.org/10.3390/IJMS241512411.
[41] Meng Y, MacIntosh BJ, Shirzadi Z, Kiss A, Bethune A, Heyn C, et al. Resting state
functional connectivity changes after MR-guided focused ultrasound mediated blood-
brain barrier opening in patients with Alzheimer's disease. Neuroimage 2019.
https://doi.org/10.1016/j.neuroimage.2019.06.060.
[42] Rezai AR, Ranjan M, D'Haese PF, Haut MW, Carpenter J, Najib U, et al. Noninvasive
hippocampal blood–brain barrier opening in Alzheimer's disease with focused
ultrasound. Proc Natl Acad Sci U S A 2020;117:9180–2.
https://doi.org/10.1073/pnas.2002571117.
[43] D'Haese P-F, Ranjan M, Song A, Haut MW, Carpenter J, Dieb G, et al. β-Amyloid
Plaque Reduction in the Hippocampus After Focused Ultrasound-Induced Blood–
Brain Barrier Opening in Alzheimer's Disease. Front Hum Neurosci 2020;14:422.
https://doi.org/10.3389/fnhum.2020.593672.
[44] Epelbaum S, Burgos N, Canney M, Matthews D, Houot M, Santin MD, et al. Pilot
study of repeated blood-brain barrier disruption in patients with mild Alzheimer's
disease with an implantable ultrasound device. Alzheimers Res Ther 2022;14:1–13.
[45] Tanaka T, Narazaki M, Kishimoto T. IL-6 in Inflammation, Immunity, and Disease.
Cold Spring Harb Perspect Biol 2014;6:16295–6.
https://doi.org/10.1101/CSHPERSPECT.A016295.
[46] Angolano C, Hansen E, Ajjawi H, Nowlin P, Zhang Y, Thunemann N, et al.
Characterization of focused ultrasound blood-brain barrier disruption effect on
inflammation as a function of treatment parameters. Biomedicine & Pharmacotherapy
2025;182:117762. https://doi.org/10.1016/J.BIOPHA.2024.117762.
[47] ARTE10 | ALZFORUM n.d. https://www.alzforum.org/research-models/arte10
(accessed November 27, 2024).
[48] Kummer KK, Zeidler M, Kalpachidou T, Kress M. Role of IL-6 in the regulation of
neuronal development, survival and function. Cytokine 2021;144:155582.
https://doi.org/10.1016/J.CYTO.2021.155582.
[49] Choi BR, Johnson KR, Maric D, McGavern DB. Monocyte-derived IL-6 programs
microglia to rebuild damaged brain vasculature. Nature Immunology 2023 24:7
2023;24:1110–23. https://doi.org/10.1038/s41590-023-01521-1.
[50] Clarke JR, Lyra e Silva NM, Figueiredo CP, Frozza RL, Ledo JH, Beckman D, et al.
Alzheimer-associated Aβ oligomers impact the central nervous system to induce
peripheral metabolic deregulation. EMBO Mol Med 2015;7:190–210.
https://doi.org/10.15252/EMMM.201404183/SUPPL_FILE/EMMM201404183.REVIE
WER_COMMENTS.PDF.
[51] Lyra e Silva NM, Gonçalves RA, Pascoal TA, Lima-Filho RAS, Resende E de PF,
Vieira ELM, et al. Pro-inflammatory interleukin-6 signaling links cognitive impairments
and peripheral metabolic alterations in Alzheimer's disease. Translational Psychiatry
2021 11:1 2021;11:1–15. https://doi.org/10.1038/s41398-021-01349-z.
[52] Ooi SZY, Spencer RJ, Hodgson M, Mehta S, Phillips NL, Preest G, et al. Interleukin-6
as a prognostic biomarker of clinical outcomes after traumatic brain injury: a

systematic review. Neurosurg Rev 2022;45:3035. https://doi.org/10.1007/S10143-
022-01827-Y.

[53] Gertz K, Kronenberg G, Kälin RE, Baldinger T, Werner C, Balkaya M, et al. Essential
role of interleukin-6 in post-stroke angiogenesis. Brain 2012;135:1964–80.
https://doi.org/10.1093/brain/aws075.

[54] Lourenco M V., Clarke JR, Frozza RL, Bomfim TR, Forny-Germano L, Batista AF, et
al. TNF-α mediates PKR-dependent memory impairment and brain IRS-1 inhibition
induced by Alzheimer's β-amyloid oligomers in mice and monkeys. Cell Metab
2013;18:831–43. https://doi.org/10.1016/j.cmet.2013.11.002.

[55] Gonzalez Caldito N. Role of tumor necrosis factor-alpha in the central nervous
system: a focus on autoimmune disorders. Front Immunol 2023;14.
https://doi.org/10.3389/fimmu.2023.1213448.

[56] Shaftel SS, Griffin WST, Kerry KM. The role of interleukin-1 in neuroinflammation and
Alzheimer disease: An evolving perspective. J Neuroinflammation 2008;5:1–12.
https://doi.org/10.1186/1742-2094-5-7/FIGURES/1.

[57] Hewett SJ, Jackman NA, Claycomb RJ. Interleukin-1β in Central Nervous System
Injury and Repair. Eur J Neurodegener Dis 2012;1:195.

[58] Tsai SJ. Effects of interleukin-1beta polymorphisms on brain function and behavior in
healthy and psychiatric disease conditions. Cytokine Growth Factor Rev 2017;37:89–
97. https://doi.org/10.1016/J.CYTOGFR.2017.06.001.

[59] Fetsko AR, Sebo DJ, Budzynski LB, Scharbarth A, Taylor MR. IL-1β disrupts the
initiation of blood-brain barrier development by inhibiting endothelial Wnt/β-catenin
signaling. IScience 2024;27.
https://doi.org/10.1016/J.ISCI.2024.109651/ASSET/D30C3404-1CE2-4F00-A09D-
579FF0CEE190/MAIN.ASSETS/GR3_LRG.JPG.

[60] Garcia JM, Stillings SA, Leclerc JL, Phillips H, Edwards NJ, Robicsek SA, et al. Role
of interleukin-10 in acute brain injuries. Front Neurol 2017;8:253134.
https://doi.org/10.3389/FNEUR.2017.00244/BIBTEX.

[61] Shemer A, Scheyltjens I, Frumer GR, Kim JS, Grozovski J, Ayanaw S, et al.
Interleukin-10 Prevents Pathological Microglia Hyperactivation following Peripheral
Endotoxin Challenge. Immunity 2020;53:1033-1049.e7.
https://doi.org/10.1016/J.IMMUNI.2020.09.018/ATTACHMENT/384895DF-BBA9-
40F0-A26B-52E33311F1B4/MMC3.PDF.

[62] Balzano T, Pineda-Pardo JA, Esteban-García N, López-Aguirre M, Reinares-
Sebastián A, Trigo-Damas I, et al. Temporal dynamics of neurovascular unit changes
following blood-brain barrier opening in the putamen of non-human primates. Journal
of Controlled Release 2025;377:116–26.
https://doi.org/10.1016/J.JCONREL.2024.11.019.

[63] Guo Y, Lee H, Kim C, Park C, Yamamichi A, Chuntova P, et al. Ultrasound frequency-
controlled microbubble dynamics in brain vessels regulate the enrichment of
inflammatory pathways in the blood-brain barrier. Nature Communications 2024 15:1
2024;15:1–17. https://doi.org/10.1038/s41467-024-52329-y.

[64] Rigollet S, Rome C, Ador T, Dumont E, Pichon C, Delalande A, et al. FUS-mediated
BBB opening leads to transient perfusion decrease and inflammation without acute or
chronic brain lesion. Theranostics 2024;14:4147–60.
https://doi.org/10.7150/THNO.96721.

[65] Liu T, Zhang L, Joo D, Sun SC. NF-κB signaling in inflammation. Signal Transduction
and Targeted Therapy 2017 2:1 2017;2:1–9. https://doi.org/10.1038/sigtrans.2017.23.

[66] Xu G, Dong F, Su L, Tan ZX, Lei M, Li L, et al. The role and therapeutic potential of
nuclear factor κB (NF-κB) in ischemic stroke. Biomedicine & Pharmacotherapy
2024;171:116140. https://doi.org/10.1016/J.BIOPHA.2024.116140.
[67] Roberts TK, Eugenin EA, Lopez L, Romero IA, Weksler BB, Couraud PO, et al. CCL2
disrupts the adherens junction: implications for neuroinflammation. Laboratory
Investigation 2012 92:8 2012;92:1213–33. https://doi.org/10.1038/labinvest.2012.80.
[68] Schellenberg AE, Buist R, Del Bigio MR, Toft-Hansen H, Khorooshi R, Owens T, et al.
Blood-brain barrier disruption in CCL2 transgenic mice during pertussis toxin-induced
brain inflammation. Fluids Barriers CNS 2012;9:1–9. https://doi.org/10.1186/2045-
8118-9-10/FIGURES/8.
[69] Guo F, Xu D, Lin Y, Wang G, Wang F, Gao Q, et al. Chemokine CCL2 contributes to
BBB disruption via the p38 MAPK signaling pathway following acute intracerebral
hemorrhage. FASEB J 2020;34:1872–84. https://doi.org/10.1096/FJ.201902203RR.
[70] Dimitrijevic OB, Stamatovic SM, Keep RF, Andjelkovic A V. Effects of the chemokine
CCL2 on blood-brain barrier permeability during ischemia-reperfusion injury. J Cereb
Blood Flow Metab 2006;26:797–810. https://doi.org/10.1038/SJ.JCBFM.9600229.
[71] Liu Z, Zhang H, Lu K, Chen L, Zhang Y, Xu Z, et al. Low-intensity pulsed ultrasound
modulates disease progression in the SOD1G93A mouse model of amyotrophic
lateral sclerosis. Cell Rep 2024;43:114660.
https://doi.org/10.1016/j.celrep.2024.114660.
[72] Kovacsa ZI, Kima S, Jikariaa N, Qureshia F, Miloa B, Lewisa BK, et al. Disrupting the
blood-brain barrier by focused ultrasound induces sterile inflammation. Proc Natl Acad
Sci U S A 2017;114:E75–84.
https://doi.org/10.1073/PNAS.1614777114/SUPPL_FILE/PNAS.201614777SI.PDF.
[73] McMahon D, Hynynen K. Acute Inflammatory Response Following Increased Blood-
Brain Barrier Permeability Induced by Focused Ultrasound is Dependent on
Microbubble Dose. Theranostics 2017;7:3989. https://doi.org/10.7150/THNO.21630.
[74] Chang JW, Wu MT, Song WS, Yang FY. Ultrasound Stimulation Suppresses LPS-
Induced Proinflammatory Responses by Regulating NF-κB and CREB Activation in
Microglial Cells. Cerebral Cortex 2020;30:4597–606.
https://doi.org/10.1093/CERCOR/BHAA062.
[75] Rauti R, Cellot G, D'Andrea P, Colliva A, Scaini D, Tongiorgi E, et al. BDNF impact on
synaptic dynamics: extra or intracellular long-term release differently regulates
cultured hippocampal synapses. Molecular Brain 2020 13:1 2020;13:1–16.
https://doi.org/10.1186/S13041-020-00582-9.
[76] Cirrito JR, Yamada KA, Finn MB, Sloviter RS, Bales KR, May PC, et al. Synaptic
Activity Regulates Interstitial Fluid Amyloid-β Levels In Vivo. Neuron 2005;48:913–22.
https://doi.org/10.1016/J.NEURON.2005.10.028.
[77] Li X, Uemura K, Hashimoto T, Nasser-Ghodsi N, Arimon M, Lill CM, et al. Neuronal
activity and secreted amyloid β lead to altered amyloid β precursor protein and
presenilin 1 interactions. Neurobiol Dis 2013;50:127–34.
https://doi.org/10.1016/J.NBD.2012.10.002.
[78] Kazim SF, Seo JH, Bianchi R, Larson CS, Sharma A, Wong RKS, et al. Neuronal
Network Excitability in Alzheimer's Disease: The Puzzle of Similar versus Divergent
Roles of Amyloid β and Tau. ENeuro 2021;8. https://doi.org/10.1523/ENEURO.0418-
20.2020.
[79] Keable A, Fenna K, Yuen HM, Johnston DA, Smyth NR, Smith C, et al. Deposition of
amyloid β in the walls of human leptomeningeal arteries in relation to perivascular

drainage pathways in cerebral amyloid angiopathy. Biochim Biophys Acta
2016;1862:1037–46. https://doi.org/10.1016/J.BBADIS.2015.08.024.
[80] Smith EE, Greenberg SM. β-Amyloid, blood vessels, and brain function. Stroke
2009;40:2601–6.
https://doi.org/10.1161/STROKEAHA.108.536839/ASSET/29E98A53-936C-43BB-
B576-D1DB972A2999/ASSETS/GRAPHIC/87FF2.JPEG.
[81] Sveikata L, Charidimou A, Viswanathan A. Vessels Sing Their ARIAs: The Role of
Vascular Amyloid in the Age of Aducanumab. Stroke 2022;53:298–302.
https://doi.org/10.1161/STROKEAHA.121.036873/ASSET/15751578-D8F8-4698-
9BE4-2841D7284378/ASSETS/GRAPHIC/STROKEAHA.121.036873.FIG01.JPG.
[82] Taylor X, Clark IM, Fitzgerald GJ, Oluoch H, Hole JT, DeMattos RB, et al. Amyloid-β
(Aβ) immunotherapy induced microhemorrhages are associated with activated
perivascular macrophages and peripheral monocyte recruitment in Alzheimer's
disease mice. Mol Neurodegener 2023;18. https://doi.org/10.1186/S13024-023-
00649-W.
[83] TauPS2APP | ALZFORUM n.d. https://www.alzforum.org/research-models/taups2app
(accessed November 27, 2024).
[84] TgCRND8 | ALZFORUM n.d. https://www.alzforum.org/research-models/tgcrnd8
(accessed November 27, 2024).
[85] 5xFAD (B6SJL) | ALZFORUM n.d. https://www.alzforum.org/research-models/5xfad-
b6sjl (accessed November 27, 2024).
[86] APPswe/PSEN1dE9 (C57BL6) | ALZFORUM n.d. https://www.alzforum.org/research-
models/appswepsen1de9-c57bl6 (accessed November 27, 2024).
803
804

Figure Legends:

Figure 1: Physiological impact of a single session of ultrasound mediated BBB-opening: A) Schematic summarizing the key steps of the protocol. Briefly, wildtype mice were first weighed and a blood sample was collected before undergoing a single sonication procedure. Four groups of mice were followed with terminal points at 2-, 4-, 8- and 15-days post sonication. During this time, mouse weights were monitored. At sacrifice, a blood sample was collected, and brains were harvested for *ex vivo* analyses. **B)** Graphic plots representing the complete blood count, each dot representing one animal. Red blood and White blood cells number were counted in the final collected blood sample (number of cells expressed in nx10^3 for white blood cells and nx10^6 for red blood cells, left Y axis). Hemoglobin levels and Mean corpuscular hemoglobin concentration (MCHC) was measured (in ng/dl, right Y axis). 2-way ANOVA with post hoc Tukey's multiple comparison test, difference between groups $F_{(3, 12)} = 2.923$, p=0.0773 **C)** Graphic plots representing the complete blood count, each dot representing one animal. Number of neutrophils, lymphocytes, monocytes, eosinophils and basophils were counted for each group (in cells per microliter). 2-way ANOVA with post hoc Tukey's multiple comparison test, difference between groups $F_{(12, 48)} = 0.8972$, P=0.5560 **D)** Heatmap representing the RT-Q-PCR quantification performed on extracted brains for different markers and normalized by the baseline group (group 0). The legend on the left indicates the level of increase or decrease compared to the Mean of the group 0. Colors are based on the Median. P value in each cell indicates the significance compared to the baseline group 0, mixed effect analysis with post hoc dunnett's multiple comparison test, difference between groups $F_{(4, 18)} = 25.68$, p<0.0001.

Figure 2: Physiological data on the ARTE10 cohort following repeated sonication protocols: A) Schematic summarizing the experimental plan for the different groups (1 to 6), the sonication schedule and sacrificing date following the last sonication session. **B)** Line graph representing the evolution of the median weight for each group (bars represent the range). No significance was observed between groups, mixed effects analysis at 1 week: $F_{(3, 20)} = 0.5542$, p=0.6513, at 1 month: $F_{(1, 10)} = 3.214$, p=0.1033.

Figure 3: Inflammatory cellular landscape of the brain following repeated sonication procedures: A) Representative image showing the presence of CD3+, GFAP+ and Iba1+ cells around an amyloid deposit, labelled with a b-amyloid antibody. **B)** Representative images showing the presence of T-cells (CD3+) present in the brain parenchyma of the different groups. **C)** Box plots showing median, 25 and 75 percentiles and min/max values,

along with individual data point for each animal, for the density of T-cells (CD3+) present in
the brain parenchyma of the different group. No significant differences between groups were
recorded, 2-way ANOVA with post hoc Tukey's multiple comparisons test, difference
between groups F (5, 25) = 1.285, P=0.3016. **D)** Representative images showing the
occupancy of astrocytes (GFAP+) in the brain parenchyma of the different groups. **E)**
Representative images showing the occupancy of microglial (Iba1+) in the brain parenchyma
of the different groups. **F)** Box plot showing median, 25 and 75 percentiles and min/max
values, along with individual data point for each animal, for the total occupancy for
astrocytes (purple bars) and microglia (yellow bars) in the brain parenchyma of the different
groups. No significant differences between groups were recorded, 2-way ANOVA with post
hoc Tukey's multiple comparisons test, difference between groups F (5, 25) = 1.280,
P=0.3038.

Figure 4: Decrease of the amyloid load following monthly sonication regimen: A)
Representative images of the hippocampal region isolated from immunohistochemistry
staining of b-amyloid on coronal sections. **B)** Box plot showing median, 25 and 75
percentiles and min/max values, along with individual data point for each animal
representing the total amyloid load (in percentage of positive area) for each group. Each dot
represents an independent animal from the group. One-way ANOVA with post hoc Tukey's
multiple comparisons test, F(3, 16)=0.04014, p=0.0163. **C)** Heatmap representing the level
of amyloid deposition in the brain at different locations from rostral to caudal. Pseudo
coloring represents the Median of each group. 2-way ANOVA, difference between groups F
(5, 25) = 3.323, P=0.0195. D) Box plot showing median, 25 and 75 percentiles and min/max
values, along with individual data point for each animal, for the total amyloid positive area
(red boxes), amyloid deposits (green boxes) and vascular amyloid deposition (yellow boxes).
2-way ANOVA with post hoc Tukey's multiple comparisons test, difference between groups
F (5, 25) = 4.317, P=0.0057.

Figure S1: Health check and BBB opening markers: A) line graph representing the
evolution of the mouse weight for each group. Each line represents an individual mouse.
Group 1 (blue) and group 2 (orange) only have 2 measurements, with a respective sacrifice
at day 2 and day 4 post sonication. Mixed effect analysis with post hoc Sidak's multiple
comparisons test, difference between timepoints F (0.1883, 1.569) = 5.678, P=0.06277, D0
vs D1 p=0.0023, **B)** Arrow plot representing the blood level of S100β in samples 15min post
sonication and at time of sacrifice (24h, 48h, 8d and 15d post sonication). Direction of the
arrow indicates the evolution of the level (ie downward arrow shows decreasing level,
upward arrow shows increasing level). Mixed effect analysis with post hoc Sidak's multiple

comparisons test, difference between timepoints F (1, 12) = 10.91, P=0.0063, post
sonication vs terminal 15 days p= 0.0115.

Supplemental Figure 2: automated image analysis pipeline and inflammatory marker
expression post repeated sonication protocol: A-C) Representative example of the
automated pipeline allowing detection of CD3+ cell **(A)**, GFAP **(B)** and Iba1 **(C)**. Positive
cells are highlighted in red, allowing the quantification of the number of CD3+ per mm2 and
the surface of occupancy for GFAP and Iba1 positive cells. **D)** heatmap representing the
pilot RT-Q-PCR dataset collected at each timepoint and exploring the level of inflammatory
markers post repeated sonication. No statistical analyses were performed due to the lack of
independent data available.

Supplemental Figure 3: Pathology of the ARTE10 and supplemental analyses for the
effect of a monthly sonication regimen: A) Representative images showing different types
of pathological deposition, with the amyloid deposits (left) and the mineralized deposits
(right, M). **B)** Representative images showing loss of hippocampal neurons in the cellular
layer of the dentate gyrus. Arrows point at the area showing loss of neuronal cell bodies.
H=hippocampus, C=cortex, M=molecular layer, G=granular layer, P=polymorphic layer. **C)**
Representative images highlighting the presence of amyloid deposition around the brain
vasculature. Arrows point at the vascular deposition, observed on serial sections. **D)** Side by
side comparison of group 1 (control) and group 6 (very sparse sonication), highlighting the
effect of the monthly sonication. Number of amyloid deposits (per mm2), percentage of
amyloid positive area and amyloid deposits area was compared between the two groups.
For all the comparisons, Group1 and Group6 were compared using nonparametric multiple t-
test Mann-Whitney. Graphical representations show median, 25 and 75 percentiles and
min/max values, along with individual data point for each animal, and statistical values
showing calculated q-value. Percentage of b-amyloid deposit area, total, group 1 vs group 6,
p=0.1143, q=0.1154; Frontal cortex, group 1 vs group 6, p=0.1143, q=0.1154. Average
deposits area, total, group 1 vs group 6, p=0.019, q=0.0385; Frontal cortex, group 1 vs group
6, p=0.1143, q=0.1154.

Supplemental Figure 4: Artificial intelligence quantification of amyloid pathology:
Representative images showing the total amyloid load by immunohistochemistry and pseudo
color from the automated quantification pipeline, as well as the vascular amyloid deposition
by immunohistochemistry and pseudo color from the automated quantification pipeline.

Supplemental Figures 5 to 10: Full brain scans of each groups, respectively Group 1 –
SuppFig5, Group 2 – Supp Fig 6, Group 3 – Suppl Fig 7, Group 4 – Suppl Fig 8, Group 5 –
Suppl Fig 9, Group 6 – Suppl Fig 10

Table 1: Row Statistics for the heatmap (Fig1D): Table summarizing for each group the
mean, SD and N for the different markers analyzed. Triplicates were averaged per mouse.

Table 2: Row Statistics for the heatmap (Supp Fig3D): Table summarizing for each group
the mean, SD and N for the different markers analyzed. As several data were missing, we
analyzed Nplicates from different mice, but did not express the data per mouse for each
group.

Table 3: Row Statistics for the heatmap (Fig4C): Table summarizing for each group the
mean, SD and N for the different locations analyzed. Triplicates were averaged per mouse.

Table 4: List of the previous mouse models used in combination with Ultrasound.
Four models have been used to explore the effect of sonication on the amyloid
pathology

Table 5: Previous studies using ultrasound in mouse models for Alzheimer's Disease
and their output.

A CD3 immunostaining

B GFAP immunostaining

C IBA1 immunostaining

D

A Control Group 1: Amyloid deposit | Control Group 1: Mineralized deposit

B Control Group 1: Loss of Neurons | Control Group 1: Loss of Neurons (zoom)

C Control Group 1: Blood Vessels | Control Group 1: Vascular Amyloid

D Control Group 1

Very Sparse Sonication Group 6

	Cxcl1			Ccl2			Il1β			Il6			Il10			Tnfα		
	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N
No BBB opening	99.9999999	6.39638142	7	99.9999999	3.30325326	7	100	3.31684234	7	100	2.21083873	7	100	3.18333499	6	100	4.48916116	7
2d post BBB opening	84.1701903	10.3300847	4	85.6161992	1.64140685	4	86.7289878	3.47666786	4	111.936937	3.37861543	4	87.399131	7.02388338	4	91.7602996	2.72942346	4
4d post BBB opening	97.4483204	6.56072308	4	84.1702549	12.1639782	4	78.9595572	9.26497926	4	105.457855	4.13181313	4	82.3091248	6.81195551	3	88.21005	8.0498684	4
8d post BBB opening	89.3910031	5.07905368	4	87.3472594	2.49080473	4	83.644939	3.23633328	4	107.984697	3.04039091	4	90.5234844	1.38846255	3	93.8540366	2.48773547	4
15d post BBB opening	86.4928354	15.2681285	4	85.6976609	1.47683189	4	85.7602802	3.27355627	4	111.461805	1.61981849	4	94.1858059	0.41330404	3	92.5977944	3.56796066	4

	Icam1			E-selectin			Tgfβ			Bdnf			Nfkb1		
	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N
No BBB opening	100	4.98481868	7	100	4.3340069	7	100	2.96480229	7	100	5.51132486	7	100	16.2264476	7
2d post BBB opening	64.494044	8.9441538	4	71.9571264	10.3749887	4	59.3623128	8.09670815	4	73.480663	11.9242656	4	67.6649466	12.1936331	4
4d post BBB opening	72.1662778	4.4780801	4	81.2168297	5.42643355	3	69.5786084	5.15336776	4	90.0636196	6.90450951	4	80.9743212	7.56250881	3
8d post BBB opening	75.8412134	3.1642435	4	87.3489353	5.27094089	4	69.2846143	6.4246666	4	83.0026788	7.85192778	4	76.69554	6.88676532	4
15d post BBB opening	75.5403414	2.01769987	4	84.0422981	9.53785609	4	65.4136917	1.61721509	4	75.7366483	3.326295	4	72.6962772	1.64031966	4

	CCl2			IL-1b			IL-6			TNF			ICAM1			Bdnf1		
	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N
Control Group 1	100	5.64080851	4	100	23.7174991	9	100	15.1470505	7	100	21.8173811	17	100	8.05468398	15	100	8.81438848	18
Sparse Sonication Group 3	105.928533	11.7733734	4	47.8831499	21.7464705	7	49.7764048	9.02670181	6	61.5928659	20.0734448	12	106.820794	20.358028	12	109.067975	62.3874537	15
Dense Sonication Group 4	107.146724	17.2155197	4	48.0101609	24.9578515	13	38.9232127	26.2442241	13	55.6141276	19.1764132	14	128.623785	9.59131353	11	187.925718	20.7368079	15
Very Sparse Sonication Group 6				54.5088907	33.2651908	6	27.9876435	17.2127141	2	46.1715867	23.6504337	8				118.01857	48.1988862	12
Control group 2	100	8.40236401	10	99.9999999	20.9397726	9	100	21.4582402	8	100	19.975462	18	100	10.631707	18	100	26.8140186	13
Sparse Sonication Group 5	83.537182	4.02692703	3	84.0437464	16.0663535	6	61.907884	23.5723261	8	69.7287926	16.8768758	14	103.320948	13.6820944	7	167.101224	21.7228397	15

	Rostral 1			Rostral 2			Middle 1			Middle 2			Caudal 1			Caudal 2		
	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N
Control Group 1	45.0633333	8.84636347	6	26.4366667	5.52090089	6	18.72	4.97570899	6	20.235	5.27793804	6	1.53666667	0.94167227	6	1.08666667	0.80988065	6
Sparse Sonication Group 3	51.554	8.37815194	5	32.4	5.06450886	5	25.356	6.96379422	5	25.132	2.81310682	5	3.19	1.59284337	5	1.252	0.37432606	5
Dense Sonication Group 4	52.242	5.79775991	5	31.006	8.54236091	5	24.794	2.25200577	5	22.134	4.6122424	5	1.708	0.53895269	5	1.134	0.45429066	5
Very Sparse Sonication Group 6	43.505	6.25032532	4	20.665	5.3789807	4	15.905	5.51050209	4	13.1075	2.96266068	4	1.235	1.42348165	4	0.7475	0.38836194	4
Control group 2	52.1633333	8.39945157	6	34.6483333	3.78678448	6	23.5783333	6.79734482	6	25.915	4.65206513	6	2.49833333	1.73152438	6	0.985	0.54309299	6
Sparse Sonication Group 5	57.806	6.49661681	5	30.46	5.54458745	5	27.006	3.18426287	5	19.918	4.69592057	5	2.856	0.92829413	5	1.406	0.28866936	5