Title Page:

Cellular and molecular underpinnings of functional networks

in the human brain

Guozheng Feng[1], Jiayu Chen[1], Jing Sui[2], Vince D. Calhoun[1*]

[1]Tri-Institutional Center for Translational Research in Neuroimaging and Data

Science (TReNDS), Georgia State University, Georgia Institute of Technology, and

Emory University, Atlanta, GA, USA;

[2]State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern

Institute for Brain Research, Beijing Normal University, Beijing, China.

[*]**Corresponding To:**

Vince D. Calhoun, PhD, Tri-Institutional Center for Translational Research in

Neuroimaging and Data Science (TReNDS), Georgia State University, Georgia

Institute of Technology, and Emory University, Atlanta, GA, USA; E-mail:

vcalhoun@gatech.edu.

Abstract

Understanding how cellular and molecular architecture underpins the large-scale organization of human brain function is a central challenge in neuroscience. By integrating transcriptomic (microarray data and single-nucleus RNA-sequencing [sn-RNA] data), molecular imaging, and neuroimaging datasets, we observed spatial correspondences suggesting that the distributions of diverse cell types, neurotransmitter systems, and mitochondrial phenotypes are aligned with intrinsic connectivity networks (ICNs)—the macroscale scaffolding of brain function. These associations extend beyond local correspondence to reflect network-level structure: inter-ICN similarity networks derived from cellular and molecular profiles significantly recapitulate both static and dynamic patterns of functional network connectivity (FNC), mirroring the established division of ICNs into canonical functional domains. Importantly, these cellular and molecular profiles not only colocalize with ICNs and FNCs but also appear to support their role as intermediaries linking microscale biological substrates to cognitive function. Mediation analyses reveal that specific ICNs statistically mediate the relationship between microscale cell-type architecture and domain-specific cognitive and behavioral processes. Moreover, FNCs capture the mediating pathways linking cell-type and neurotransmitter similarity networks to cognitive network organization. Taken together, our findings suggest that the brain's functional architecture shows systematic associations with cellular and molecular organization, which may act as a biological constraint guiding functional network formation and contribute to the neural basis of cognition.

Keywords: cell types, intrinsic connectivity network, functional networks, cognitive function, transcriptomic, cellular and molecular regulation

Introduction

Advances in study of brain networks with resting-state functional MRI (rs-fMRI) have significantly deepened our understanding of both typical and atypical neural activity[1-6]. Among various analytic approaches, decomposition-based independent component analysis (ICA) enables the reliable extraction of intrinsic connectivity networks (ICNs), which reflect coherent patterns of spontaneous brain activity within distinct functions[7-9]. Each ICN comprises a weight map paired with an associated time course (TC), capturing within-component connectivity in the blood oxygenation level-dependent (BOLD) signal[10]. Correlating these TCs yields between-component relationships, typically summarized as functional network connectivity (FNC)[11]. The macroscopic functional brain networks emerge from the dynamic interplay of distributed neural circuit architectures, whose region-specific implementations—distinct cellular compositions and specialized input–output motifs—necessitate tailored neuromodulatory regulation and mitochondrial bioenergetics to sustain network-specific information processing and integration[12-21]. Despite growing evidence for the relevance of these cellular and molecular factors, their specific contributions to the organization of functional networks remain largely uncharacterized.

The growing availability of high-resolution cellular and molecular atlases of the human brain—such as microarray-based transcriptional data from the Allen Human Brain Atlas (AHBA)[22], single-nucleus RNA sequencing (snRNA-seq) datasets from discrete brain regions[21,23,24], and positron emission tomography (PET) imaging of diverse neurotransmitter systems[25]—has created new opportunities to bridge microscale biology with macroscale brain organization. Previous work established spatial associations between the distribution of interneuron-linked genes and regional differences in fMRI signal variability[26,27]. The low-dimensional manifold organizing fMRI activity shows opposing associations with regional densities of facilitatory versus inhibitory neuromodulatory receptors[28]. In parallel, specific mitochondrial phenotypes selectively align with distinct fMRI signal metrics (e.g., maximum BOLD, regional homogeneity, entropy)[21]. Beyond spatial alignment, molecular similarity further supports the organization of functional networks: functionally coupled regions tend to share gene-expression signatures[29-31], and distinct cell-type distributions differentially align with the cortex's principal unimodal-to-transmodal functional gradient[32]. Concordantly, neurotransmitter receptor and transporter similarity across regions

correlates with functional connectivity, with brain areas exhibiting similar receptor profiles showing stronger co-activation[25]. Although most prior studies have focused on individual neurobiological scales, observed correlations among neurobiological scales[25,33] suggest that multi-scale factors are likely to synergistically contribute to shape functional network phenotypes. One illustrative case is cholinergic neurotransmission primarily modulates neuronal excitability, regulates presynaptic transmitter release, and orchestrates the firing of neuronal populations[34-36]. Additionally, developmental patterns of cortical thickness are better captured by models that integrate the spatial distributions of several systems (e.g., cell-type composition, neurotransmitter receptor/transporter densities), with neurotransmitter markers showing relatively greater scale-level relative importance[37]. Together, integrating diverse cellular and molecular atlases provides a multi-scale perspective for revealing the biological foundations of intrinsic functional networks.

The human brain supports a wide range of cognitive and behavioral functions through large-scale neural circuits shaped by its underlying cellular and molecular architecture[17]. At the macroscale, ICNs, as the core organizational units of functional architecture, along with their static and dynamic FNCs, demonstrate heterogeneous contributions to diverse cognitive abilities[38,39]. At the microscale, specific inhibitory neuron subtypes are selectively preserved in individuals who maintain high cognitive performance in late life, implicating cell-type-specific influences on cognitive resilience[40]. The spatial distribution of neurotransmitter receptors differentiates intrinsic and extrinsic cognitive modes[25]. Animal studies further underscore the role of molecular processes in cognition and behavior: mitochondrial function has been shown to impact both cognitive capacity[41,42] and behavioral regulation[43,44], while distinct neurotransmitter receptor systems have been associated with specific cognitive domains[45]. Despite these converging findings, the mechanisms through which ICNs and FNCs mediate the relationship between microscale biological substrates and cognitive and behavior abilities remain poorly understood. Uncovering this linkage is essential for a deeper understanding of how cellular and molecular systems scaffold the brain's functional architecture and enable complex cognition.

Here, motivated by the above evidence, we aim to probe the cellular and molecular bases of the spatial and connectional architecture of ICA-derived ICNs from a multi-scale perspective. For this purpose, we integrate spatial maps of cell types, neuromodulatory receptor/transporter systems,

and mitochondrial phenotypes. We hypothesize that ICN spatial maps colocalize with these

microscale architectures—such that networks are differentially enriched for several biological

features that support network-specific information processing and integration. We further

hypothesize that between-region similarity network derived from these cellular and molecular

features is associated with FNC, consistent with an organizing role for these similarity networks.

Moreover, multimodal combinations of neurobiological features explain more ICN/FNC variance

than any single system, consistent with potential synergistic effects among distinct microscale

processes. Finally, we posit that ICNs and FNCs provide the macroscale substrates that statistically

link microscale neurobiology to cognitive/behavioral maps.

Results

We analyzed 53 reproducible of ICN spatial maps and their corresponding FNC (*Supplementary*

Fig. S1 and Fig. 1A) derived from the Neuromark_fMRI_1.0 template[7]. The NeuroMark template

was based on components that replicated across two large-scale datasets: the Genomics Superstruct

Project (GSP; $n = 1,005$; mean age $= 21.54 \pm 2.96$ years; 411 males)[46] and the Human Connectome

Project (HCP; $n = 823$; mean age $= 28.79 \pm 3.68$ years; 356 males)[47]. To investigate the cellular and

molecular architecture underlying ICNs, we systematically examined the spatial and connectional

correspondence between these functional networks and a diverse sets of cell-type[23],

neurotransmitter[25], and mitochondrial[21] maps (Fig. 1B). Extending this analysis, we incorporated

123 cognitive and behavioral probabilistic maps from Neurosynth[48] to assess whether functional

networks (ICNs and FNCs) mediate the relationship between microscale biological substrates and

spatial and connectional variations in cognitive and behavioral functions.

Cell-type associations of ICNs. To investigate the cellular architecture underlying ICNs, we

related postmortem gene expression–based estimates of cell-type distribution to the spatial maps of

44 cortical ICNs. Bulk tissue transcriptomic data from AHBA[22] were processed using an established

harmonization pipeline[49-51]. Cell-type–specific transcriptional signatures were derived from

snRNA-seq of eight cortical areas[23], enabling the imputation of 24 cell-type abundances across

AHBA samples. These included nine excitatory glutamatergic neurons (e.g., L2/3 IT, L4IT, L6IT),

nine inhibitory GABAergic interneurons (e.g., Chandelier, PVALB, SST), and six non-neuronal cell

types (e.g., Astro, Endo, OPC), each characterized by distinct laminar locations, developmental

origins, morphology, and projection targets[23,52]. Voxel-level spatial distributions of cell-type

abundance were correlated with ICN spatial maps using Spearman's r correlation. Statistical

significance was determined using permutation-based two-sided Moran test[53] (1,000 iterations),

with false discovery rate (FDR) correction applied across 24×44 comparisons.

Seven ICNs exhibited at least one significant association with the imputed abundance of

specific cell-types (Fig. 2A). These associations included both positive and negative correlations,

indicating that cell-type gradients may support the spatial organization of ICNs. IC14 (precentral

gyrus network) in the sensorimotor network (SM) domain was positively associated with L5ET (r

$(107) = 0.48$, $q < 0.001$). IC25 (middle temporal gyrus network) in the visual network (VIS) was

positively associated with L2/3IT ($r (140) = 0.30$, $q < 0.001$). In the cognitive-control network (CC)

domain, IC27 (insula network) was significantly associated with L6ITCar3 ($r (127) = 0.58$, $q <$

0.001), and IC28 (superior medial frontal gyrus network) was negatively associated with L4IT (r

$(123) = -0.58$, $q < 0.001$), IC31 (middle frontal gyrus network) was positively associated with

LAMP5 ($r (110) = 0.49$, $q < 0.001$), IC40 (inferior frontal gyrus network) was negatively associated

with L5/6NP ($r (88) = -0.40$, $q < 0.001$). IC45 (anterior cingulate cortex network) in the default

mode domain was positively associated with SST ($r (139) = 0.32$, $q < 0.001$) and negative

associations with OPC ($r (139) = -0.25$, $q < 0.001$). The spatial distributions for all 24 cell types are

presented in Fig. 2B. A complete summary of cell-type–ICN associations is provided in *Extended*

Data Table 1. Collectively, these findings indicate that particular ICNs show specific spatial

correspondences with distinct cell types.

Neurotransmitter and mitochondrial associations of ICNs. To examine the molecular

architecture underlying ICNs, we assessed the spatial alignment between ICN maps and

neurotransmitter systems using PET-derived estimates of receptor and transporter densities. We

compiled 19 neurotransmitter maps encompassing nine neurotransmitter systems, aggregated from

PET imaging data across more than 1,200 healthy individuals[25]. These maps included serotonergic

(5-HT_{1A}, 5-HT_{1B}, 5-HT_{2A}, 5-HT_4, 5-HT_6, 5-HTT), dopaminergic (D_1, D_2, DAT), noradrenergic

(NET), histaminergic (H_3), cholinergic ($\alpha_4\beta_2$, M_1, VAChT), cannabinoid (CB_1), opioid (MOR),

glutamatergic (NMDA, $mGluR_5$), and GABAergic ($GABA_A$) systems.

For each ICN, we constructed multilinear regression models using the full set of receptor and transporter densities as predictors. Statistical significance was assessed using a one-sided Moran test (1,000 permutations) to account for spatial autocorrelation[53], FDR correction applied across 53 comparisons. Sixteen ICNs exhibited significant model fits (adjusted r^2: 0.43–0.86, $q < 0.05$, Fig. 3A), indicating that distinct combinations of neurotransmitter systems spatially align with specific ICN architecture; effects were densest in SC and AUD domains. To determine the relative contribution of individual neurotransmitter systems, we applied dominance analysis, which quantifies the proportion of variance in each ICN map attributable to each receptor/transporter in the model[54]. Dominance scores were normalized by the model's total adjusted r^2, enabling comparison across ICNs (Fig. 3B). Statistical significance of neurotransmitter-specific dominance was assessed using one-sided Moran test (1,000 permutations) to account for spatial autocorrelation[53], FDR correction applied across 19×53 comparisons. Variance contributions of neurotransmitter receptors were heterogeneous across ICNs. For example, in CC domain, the $\alpha_4\beta_2$ receptor made a prominent contribution to IC31 (middle frontal gyrus network, contribution = 26%, $q < 0.001$), while CB_1 receptor made a prominent contribution to IC37 (Hippocampus network, contribution = 19%, $q < 0.001$). In DM domain, 5-HT_{1B} receptor showed significant influence on IC48 (precuneus network, contribution = 25%, $q < 0.001$). A complete summary of neurotransmitter–ICN associations is provided in *Extended Data Table 2*.

In parallel, we examined the association between mitochondrial phenotypes and ICNs. Six mitochondrial maps (complex I [CI], complex II [CII], complex IV [CIV], mitochondrial density [MitoD], tissue respiratory capacity [TRC], and mitochondrial respiratory capacity [MRC])[21] were regressed onto ICN spatial maps, FDR correction applied across 53 comparisons for adjusted r^2 and 6×53 comparisons for dominance. We observed significant model fits for IC2 (adjusted $r^2 = 0.25$, $p < 0.001$) and IC4 (caudate network, adjusted $r^2 = 0.24$, $p < 0.001$), both located in SC domain (*Supplementary Fig. S2A*). Within these models, MitoD, and CIV each made significant relative contributions (*Supplementary Fig. S2B*). A complete summary of mitochondrion–ICN associations is provided in *Extended Data Table 3*. These findings suggest that mitochondrial energy metabolism may contribute to the functional architecture of specific ICNs in SC domain.

To investigate the synergy and relative contributions of distinct neurobiological scales to ICNs,

we examined associations between multi-scale markers (cell types [21, after collinearity attenuation,

Supplementary Fig. S3; see Methods], neurotransmitters [19], and mitochondria [6]) and ICNs.

Fourteen ICNs exhibited significant model fits (adjusted r^2: 0.62–0.99, $q < 0.05$; *Supplementary Fig.*

S4A). In 13 of these 14 ICNs, multi-scale models outperformed single-scale models ($q < 0.05$;

Supplementary Fig. S4B), underscoring the synergistic value of integrating distinct neurobiological

scales. We further observed a consistent trend of synergistic contributions from markers across

scales ($p < 0.05$, *Supplementary Fig. S4A*). Among these, neurotransmitter systems consistently

accounted for the largest overall relative contributions across scales (*Supplementary Fig. S4B*).

Cellular and molecular similarity networks relate to functional connectivity. Having

established that ICNs exhibit distinct spatial associations with cellular and molecular features, we

next asked whether ICNs that are similarly composed at the cellular and molecular levels also

exhibit stronger functional interactions. To this end, we computed similarity networks for cell-type,

mitochondrial, neurotransmitter, and combined molecular fingerprints by correlating the mean

spatial profiles of each ICN pair using Spearman's correlation (Fig. 4A). Across all modalities,

cellular and molecular similarities were significantly greater among ICNs within the same

functional domain than between different domains, even after regressing out Euclidean distance and

spatial overlap (Dice coefficient) from the similarity matrices ($p < 0.001$, ranksum test; $N_{within} = 230$

edges, $N_{between} = 716$ edges for cell-type; $N_{within} = 246$ edges, $N_{between} = 1,132$ edges for others, Fig.

4B). We then assessed the relationship between cellular/molecular similarity and functional

connectivity, as indexed by the group-level FNC from the HCP dataset (Fig. 4C). We observed that

cell-type similarity (Spearman's r (946) = 0.19, $p < 0.001$), neurotransmitter similarity (Spearman's

r (1378) = 0.27, $p < 0.001$), and the combined similarity (Spearman's r (1378) = 0.46, $p < 0.001$)

networks were all significantly correlated with FNC strength (Fig. 4D). At the regional level,

correlations were predominantly positive, with the strongest associations observed in nodes located

within the SM and VIS domains (Fig. 4E, *Extended Data Table 4*). To further investigate the

dynamic aspects of this relationship, we identified five discrete states of dynamic FNC

(*Supplementary Fig. S5A*) and examined their correspondence with cellular and molecular similarity

networks. Results consistently demonstrated robust associations between FNC states and similarity

networks, both at the global (*Supplementary Fig. S5B–F*) and regional levels (*Supplementary Fig.*

S6A–E). Together, these results suggest that ICNs with similar cellular and molecular compositions are more likely to be functionally coupled, independent of spatial proximity, supporting the idea that inter-regional communication is constrained by a common microscale architecture.

Identify cellular and molecular structural associations with functional domain. To test whether cellular and molecular profiles reflect the functional domain architecture of ICNs, we constructed similarity networks based on cell-type, neurotransmitter, mitochondrion and combination fingerprints, while regressing out the effects of Euclidean distance and Dice similarity from the network matrices. We reduced network connectivity features using diffusion embedding[55], and projected each similarity network into a three-dimensional space (cumulative variance explained: 57% for cell type, 51% for neurotransmitter, 54% for mitochondrion, and 50% for the combined profile). Component-wise explained variance is reported in *Supplementary Fig. S7*. Comparisons of dimensionality reduction methods are shown in *Supplementary Fig. S8.* We then applied k-means to cluster features of cellular/molecular similarity network. The ground-truth labels were derived from the functional domain assignments of ICNs, and clustering performance was quantified using the adjusted rand index (ARI), with significance assessed via a null generated from 1,000 label permutations. Clustering based on cell-type (ARI = 0.34, $p < 0.001$, 5 clusters; Fig. 5A), neurotransmitter (ARI = 0.34, $p < 0.001$, 7 clusters; Fig. 5B), mitochondrion (ARI = 0.26, $p < 0.001$, 7 clusters; Fig. 5C), and combined profiles (ARI = 0.26, $p < 0.001$, 7 clusters; Fig. 5D) all showed significant alignment with functional domain structures. These findings indicate that the spatial organization of functional domains is supported by coherent cellular and molecular architectures.

ICNs mediate spatial associations between cellular/molecular architecture and cognitive function. To examine whether and how ICNs mediate the relationship between microscale cellular and molecular organization and macroscale cognitive architecture, we leveraged probabilistic meta-analytic activation maps from Neurosynth (https://github.com/neurosynth/neurosynth)[48]. These maps encode the likelihood that a given cognitive term is associated with activation at each voxel, based on over 15,000 published fMRI studies. A total of 123 cognitive and behavioral terms were selected based on their functional annotations in the Cognitive Atlas[56]. These term-wise maps showed significant spatial correspondence with specific ICNs (*Supplementary Fig. S9*). For cellular/molecular maps and ICNs that showed significant spatial correspondence in the primary

analyses, we then conducted post-hoc mediation analyses using the PROCESS toolbox in R. In each model, the cellular/molecular map served as the predictor, the cognitive probability map as the outcome, and the ICN as the mediator. FDR correction applied across 123×30 comparisons.

These analyses revealed significant mediation pathways (*Extended Data Table 5*), predominantly involving 7 ICNs, 10 cellular/molecular maps, and 123 cognitive and behavior functions. Among cortical ICNs (Fig. 6), the insula network (IC27; L6ITCar3-coupled) showed the broadest and strongest mediation spectrum (mediation effect β: 0.21–0.57), spanning language (speech perception/production, naming), sensory integration (multisensory, categorization), executive control (cognitive control, inhibition, decision making), and emotion-related processes (reward anticipation, anxiety, fear). The precentral gyrus network (IC14; L5ET-coupled) exhibited weaker mediation for executive/memory functions but stronger effects on perceptual attention and movement/motor control (mediation effect β: 0.10–0.23). The superior medial frontal gyrus network (IC28; L4IT-coupled) showed modest effects focused on planning and attention, with additional links to movement (mediation effect β: 0.11–0.19). Finally, the anterior cingulate cortex network (IC45; SST-coupled) preferentially mediated affective and motivational processes, including fear, anxiety, mood regulation, and reinforcement learning (mediation effect β: 0.13–0.21). By comparison, neurotransmitter and mitochondrial markers primarily mediated subcortical ICN–cognition associations (*Extended Data Table 5*). Overall, these spatial association–mediation results indicate that specific cellular/molecular markers interface with functional networks to provide indirect links to cognitive and behavioral phenotypes.

FNCs mediate the connectional associations between cellular and molecular architecture and cognitive function. To further investigate whether inter-ICN functional connectivity (static and dynamic FNCs) mediates the relationship between cellular/molecular similarity and cognitive similarity, we first averaged the cognitive probabilistic maps across the 53 ICNs to obtain ICN-level cognitive features. We then constructed cognitive similarity networks by computing Spearman's correlation across ICN pairs. Partial Spearman's correlations were performed to assess associations between cellular/molecular similarity networks and the cognitive similarity network, as well as between static and dynamic FNC states and the cognitive similarity network. To account for spatial confounds, Euclidean distance and Dice similarity were included as covariates in all correlation and

mediation analyses. We found that the cell-type, neurotransmitter, and combined similarity networks

were significantly associated with the cognitive similarity network ($p < 0.001$, *Supplementary Fig.*

S10). Both static FNC and dynamic FNC states showed significant associations with the cognitive

similarity network ($p < 0.001$, *Supplementary Fig. S11*). Mediation analyses at the network level

treated the cellular or molecular similarity network as the predictor, the cognitive similarity network

as the outcome, and the FNC (static or dynamic) as the mediator. The results revealed that static

FNC partially mediated the relationship between the cell-type similarity network and the cognitive

similarity network (path *ab*: $p = 0.002$, $\beta = 0.04$, 95% CI = [0.01, 0.06]; path *c'*: $p < 0.001$, $\beta = 0.31$,

95% CI = [0.26, 0.35]; path *c*: $p < 0.001$, $\beta = 0.31$, 95% CI = [0.30, 0.40], Fig. 7A). Static FNC also

mediated the link between the neurotransmitter similarity network and cognitive similarity network

(path *ab*: $p < 0.001$, $\beta = 0.05$, 95% CI = [0.03, 0.06]; path *c'*: $p < 0.001$, $\beta = 0.21$, 95% CI = [0.16,

0.25]; path *c*: $p < 0.001$, $\beta = 0.25$, 95% CI = [0.21, 0.29], Fig. 7B), as well as between the

combination similarity network and the cognitive similarity network (path *ab*: $p < 0.001$, $\beta = 0.08$,

95% CI = [0.07, 0.10]; path *c'*: $p < 0.001$, $\beta = 0.36$, 95% CI = [0.32, 0.41]; path *c*: $p < 0.001$, $\beta =$

0.45, 95% CI = [0.40, 0.49], Fig. 7D). No significant mediation was detected for the mitochondrial

similarity network (Fig. 7C). Mediation by dynamic FNC states was also significant as shown in

Supplementary Fig. S12. Specifically, states 2–5 mediated the relationship between cell-type and

cognitive similarity; states 3–5 mediated the neurotransmitter–cognition link; states 1, 3–5 mediated

the mitochondrion–cognition link. All states significantly mediated the association between

combination–cognition link. Together, these findings demonstrate that static and dynamic FNCs

provide indirect pathways linking cellular/molecular architecture to cognitive organization, with

dynamic FNC offering additional state-specific routes.

Discussion

By integrating transcriptomic, molecular, and neuroimaging data, we observed spatial

associations between the brain's large-scale functional architecture and its underlying cellular and

molecular organization. Spatial profiles of diverse cell types, neurotransmitter systems, and

mitochondrial markers show selective alignment with ICNs, providing biologically grounded

support for functional segregation. These relationships extend beyond local overlap: similarity

networks constructed from molecular and cellular features approximated patterns of functional

connectivity and captured core ICN functional domains. Importantly, ICNs statistically served as

intermediaries in the associations between molecular architecture and cognitive processes,

positioning them as critical intermediaries linking microscale biology to behavior. Together, these

findings suggest that cellular and molecular organization may impose constraints on the formation

and function of large-scale brain networks.

rs-fMRI investigations of brain networks have substantially expanded the frontiers of cognitive

neuroscience and psychiatric research. Among data-driven approaches, ICA leverages the hidden

spatiotemporal structure of the BOLD signal to separate meaningful neural signals from noise, while

preserving subject-level variability[57] and yielding improved signal-to-noise ratios[7]. Critically,

unlike parcellation-based approaches, ICA enables the delineation of distinct patterns of functional

activity, thereby yielding ICN maps[58]. Each ICN comprises brain voxels that co-vary over time,

capturing within-component connectivity. Functional connectivity between ICNs (FNC) can

subsequently be derived by quantifying temporal correlations across the ICA time series. To

facilitate cross-study reproducibility, methodological advances have led to the fully automated

spatially constrained ICA framework (NeuroMark) as a template based on 1828 young adult

participants in the GSP and HCP datasets[7]. This approach has identified 53 reliable ICNs, which

have since served as functional templates in studies of lifespan development[8], individual differences

in cognition[38,59], and a range of neuropsychiatric disorders[60,61]. These large-scale ICNs are thought

to arise from the coordinated activity of underlying neural circuits, yet the extent to which they are

shaped by microscale cellular and molecular substrates remains poorly understood. Elucidating the

cellular and molecular architecture underlying ICNs and their connectivity patterns (FNCs) could

offer a foundational neurobiological annotation of functional network organization. Such insights

are crucial for bridging levels of analysis from genes and cells to circuits and cognition.

The macroscale functional network organization of the brain is increasingly understood to

reflect underlying microscale cellular and molecular architecture[18,26,32]. To explore this hypothesis,

we systematically integrated multiple cellular and molecular datasets and quantified their spatial

associations with ICNs. We observe spatial coupling between the distributions of select cell types

and ICNs, consistent with shared structural and computational motifs. For example, L5ET neurons

constitute the cortex's principal corticofugal output, routing columnar computations to thalamus,

brainstem, and spinal targets[62,63]; their alignment with the precentral gyrus network (IC14) supports

involvement in action control and motor execution circuits[64]. By contrast, L6ITCar3 neurons

possess extensive intracortical axon projections that enable cross-areal integration[63,65], providing a

cellular substrate for the insula network's (IC27) broad role as a switching hub across externally

oriented and internally oriented systems[66]. SST interneurons in layers 2/3 and 5 elaborate axons into

layer 1 and thereby preferentially suppress inputs arriving through layer 1, i.e., gate top-down

signals at apical dendrites[67,68] consistent with anterior cingulate cortex (IC45) computations, which

integrate cost–benefit and performance signals to guide the allocation of control[69]. Interestingly, the

anatomical cores of the ICNs showing cellular associations lie at key waypoints along the cortex's

principal unimodal-to-transmodal gradient[55]—anchored by IC14 (precentral gyrus) at the unimodal

sensorimotor pole and extending toward IC28 (superior medial frontal gyrus) and IC45 (anterior

cingulate) near the transmodal apex—while IC27 (insula) acts as switching hub between large-scale

systems for external attention and self-referential cognition[70]. The cell types that capture ICN spatial

variability likewise traverse this gradient, echoing evidence that functional gradients parallel

systematic shifts in cellular composition[32]. Neurotransmitter receptor/transporter maps and

mitochondrial phenotypes most strongly predicted ICN variation in SC domain, consistent with the

high SC expression of dopaminergic and cholinergic markers (e.g., D1/D2[71,72], DAT[73], NET[74],

VAChT[75]) and with the energetic specializations of subcortical circuits[12]. In contrast, within the

cortex, neurotransmitter maps were more closely associated with ICNs situated along the non-

waypoints of the principal gradient, seemingly reflecting scale-level specificity in the

neurobiological determinants of ICN organization. Although speculative, cell-type maps, which

index laminar architecture and compositional differences[23,52,76,77] preferentially provided structural

substrates for ICNs anchored at the waypoints of the gradient. By comparison, neurotransmitter

maps, which capture modulatory sensitivity and synaptic communication[12,25,28,78,79], appeared to

preferentially support the dynamic switching of networks positioned at non-waypoints along the

gradient. Moreover, multi-scale models that combined neurobiological scales explained more ICN

variance than any single scale, suggesting that functional networks may be shaped by the synergistic

interactions among distinct cellular and molecular processes. This pattern is consistent with

independent reports that neuromodulatory systems disproportionately shape patterns of brain

structural development[37]. Notably, unlike prior parcellation-based spatial correlation approaches, the ICN-based framework offers a functionally segregated and spatially specific lens for cellular and molecular annotation. ICNs capture voxel-wise coactivation patterns associated with independent functional processes, enabling a finer-grained and functionally dedicated mapping between brain function and neurobiology. This approach advances beyond prior transcriptomic–functional studies by allowing the annotation of individual functional components, rather than broad cortical territories, with targeted cellular and molecular features.

Extending these findings, we observed that ICNs assigned to the same functional domain exhibit significantly greater similarity in their cellular and molecular profiles—across cellular, neurotransmitter, and mitochondrial modalities—compared to ICNs from different domains. These similarities were paralleled by stronger functional co-activation, suggesting that shared microscale composition may facilitate coordinated activity between regions. This supports and expands upon prior work linking transcriptional profiles[29-31,80] and neurotransmitter systems[25] to macroscale brain networks. Clustering similarity networks derived from cellular and molecular fingerprints revealed a non-random alignment with canonical functional domain assignments. Specifically, ICNs were grouped in a manner that significantly recapitulated known functional subdivisions, suggesting that the brain's cellular and molecular landscape may, in part, mirror the organization of functional networks revealed by group-average fMRI data. These results reinforce the notion that domain-specific functional architectures are supported by cell-type and neurotransmitter-specific features. Notably, this molecular-functional alignment echoes anatomical observations from histological analyses, where homogeneous cellular composition is observed within local territories, and sharp transitions often demarcate adjacent cortical areas[81]. Interestingly, ICNs within the VIS and SM domains showed higher molecular and functional coupling than those within CC and DM domains. These domain-specific molecular architectures may also help explain observed correlation patterns of white matter networks[82] and structure–function network coupling[18,83], highlighting their role in shaping both network identity and inter-network communication. From a molecular–structural perspective, this finding supports the view that the association cortex may escape the strong structural constraints imposed by early sensory–motor cascades, thereby enabling higher-order cognitive functions beyond simple sensorimotor exchange[84].

Microscale evidence implicates cellular and molecular systems sculpting cognitive capacities[40,85-87]. Yet, the extent to which macroscale ICNs and FNCs mediate the relationship between cellular and molecular architecture and cognition remains poorly understood. Meta-analytic cognitive atlases (e.g., Neurosynth) combine text mining, meta-analysis, and machine learning to generate probabilistic mappings between cognitive terms and neural activity patterns[48]. Building on this resource, our significant spatial association–based mediation results indicate that a set of mesoscale ICNs acts as putative interfaces through which cortical cell-type gradients relate to cognitive representations. This pattern accords with established functional accounts: the precentral gyrus network (IC14) aligns with motor circuitry and corticofugal output[88]; the insula network (IC27) supports switching between externally oriented and internally oriented/self-referential systems[70]; and the anterior cingulate cortex network (IC45) integrates cognitive control with affective/motivational processes[69,89]. SC-related ICNs anchored mediation pathways linking neurotransmitter receptor/transporter markers and mitochondrial phenotypes to broad cognitive functions, underscoring the central role of SC circuits in neuromodulatory regulation and energetic support for cognition[12,90-92]. These exploratory findings extend prior work by anchoring these functional profiles in plausible cellular architecture. From a network connectivity perspective, we observed significant associations among cellular/molecular similarity networks, static and dynamic FNCs, and the cognitive similarity network. These findings suggest that FNC—particularly its dynamic aspects—may act as a critical bridge linking microscale biological architecture to macroscale cognitive function. The observed mediation effects suggest that the alignment between brain biology and behavior is not merely correlative but may be systematically organized through the brain's dynamic communication architecture. In particular, the mediating role of dynamic FNC highlights how transient, state-dependent connectivity patterns may encode biologically meaningful variations relevant to higher-order cognition. Together, these findings move beyond simple correlations between cell types and networks or networks and cognition. Instead, they delineate a mediational pathway wherein macroscale functional networks serve as mechanistic conduits linking molecular-scale organization to behavioral phenotypes. This integrative model not only strengthens the biological plausibility of structure–function–cognition relationships but also provides a principled framework for understanding the neurobiological substrates of individual cognitive

differences.

While our findings offer novel insights into the cellular and molecular foundations of ICNs,

several limitations should be acknowledged. In this study, we derived imputed spatial cellular maps

by cross-platform integration of snDrop-seq with AHBA microarray data and evaluated two

deconvolution algorithms (CIBERSORTx[93] and BayesPrism[94]), which showed overall moderate

concordance (*Supplementary Fig. S13*). Given the early stage of this field, we adopted the

CIBERSORTx-derived estimates as our primary analytic input for continuity with prior work[26,32],

and because a recent human-brain benchmarking study reported comparatively higher accuracy for

this method[95]. Nevertheless, algorithm-dependent biases highlight the need for further data

generation and head-to-head methodological evaluations tailored to this question. Second, the

cognitive density maps used here are meta-analytic, inheriting heterogeneity from the underlying

literature and potentially underrepresenting the multidimensionality of cognitive representations.

Accordingly, our associational and mediation results should be interpreted as indirect, exploratory

inferences rather than causal claims. Third, the current analyses focus on healthy young adults,

leaving generalizability across development, aging, and neuropsychiatric conditions unresolved; the

cellular and molecular maps are donor-based, which may obscure individual-level variability related

to age, sex, and other demographics; and brain-wide coverage remains uneven—particularly in

subcortical and cerebellar territories and in regions with sparse receptor mapping. Looking forward,

expanding spatially resolved molecular datasets (especially in these under-characterized areas) and

integrating emerging individual-level multimodal resources will be essential to build more complete

and personalized models of molecular–functional coupling.

In summary, this study offers a multi-scale framework linking molecular and cellular

architecture to large-scale brain function by integrating transcriptomic and neurotransmitter maps

with functional networks. We show that ICNs appear not only spatially aligned with, but also

structurally constrained by underlying biological features—constraints that extend beyond local

correspondence to reflect network-level structure. Moreover, we identify specific ICNs that mediate

the translation of molecular substrates into cognitive functions, suggesting a potential bridge from

microscale biology to macroscale behavior. These findings advance our understanding of how

functional brain networks emerge from biological structure and laying the groundwork for

integrative models that link molecular neuroscience, functional imaging, and cognition.

Materials and methods

Participants and imaging preprocessing. A total of 823 healthy young adults were selected from HCP (mean age = 28.79 ± 3.68 years; 356 males)[47] (http://www.humanconnectomeproject.org/data) dataset. Comprehensive demographic information and imaging acquisition protocols are described in reference[47]. We used preprocessed rs-fMRI data (http://www.humanconnectomeproject.org/data). The HCP preprocessing pipeline included spatial distortion correction, motion correction, echo planar imaging (EPI) distortion correction, registration to standard Montreal Neurological Institute (MNI) space, and intensity normalization. Subsequently, the data were resampled to $3 \times 3 \times 3$ mm³ isotropic voxels and spatially smoothed using a Gaussian kernel with a full width at half maximum (FWHM) of 6 mm. The detailed quality control procedure for all the preprocessed data can be found in the reference[7].

ICNs. We utilized 53 ICNs from the Neuromark_fMRI_1.0[7], which are highly reproducible and biologically meaningful across independent ICA analysis of multiple large N cohorts following standard criteria for selecting ICNs. Each ICN represents a spatially independent component reflecting coherent brain activity. These ICNs were grouped into seven functional domains based on anatomical and functional priors: subcortical network (SC, 5 ICNs), auditory network (AUD, 2 ICNs), sensorimotor network (SM, 9 ICNs), visual network (VIS, 9 ICNs), cognitive-control network (CC, 17 ICNs), default-mode network (DM, 7 ICNs), and cerebellar network (CB, 4 ICNs).

FNC. The preprocessed HCP data were decomposed into 53 subject-specific independent components and their associated TCs using a spatially constrained single-subject ICA method, with the Neuromark_fMRI_1.0 template serving as spatial priors (available via GIFT at http://trendscenter.org/software/gift)[7]. To minimize residual noise in the ICN TCs, four additional postprocessing steps were applied: (1) detrending to remove linear, quadratic, and cubic trends; (2) outlier detection and removal; (3) multiple regression of six head motion parameters (three translations and three rotations) along with their temporal derivatives; and (4) band-pass filtering (0.01–0.15 Hz). Pearson's correlations were computed between the denoised ICN time courses to generate individual-level FNC matrices, which were then averaged to obtain a group-level FNC

matrix. Dynamic FNC was estimated using a sliding-window approach, segmenting the TCs into

overlapping temporal windows, as described previously[38]. For each segment, we computed

covariance from the regularized precision matrix[96] using the graphical LASSO method[97] with an L1

penalty to promote sparsity. The regularization parameter (λ) was optimized for each scan by

maximizing the log-likelihood on held-out data[98]. Window-wise FNC estimates were concatenated

to yield a subject-level dynamic FNC representation, capturing temporal variations in ICN

connectivity. To reduce redundancy, windows were subsampled along the temporal axis. Exemplar

windows were selected based on local maxima in FNC variance across connections. These

exemplars were clustered using k-means, repeated 500 times with random centroid initializations to

ensure stability. The optimal number of states was determined via the elbow criterion[99]. The final

centroids were used to assign all windows to discrete states. The fractional rate was quantified as

the percentage of windows assigned to each state. For each state, dynamic FNCs were averaged

across all windows assigned to that state to generate group-level state-specific FNC patterns.

AHBA microarray data processing. Human brain gene expression data from 3,702 bulk tissue

samples across six postmortem brains were obtained from the AHBA dataset (http:// human.brain-

map.org/)[22]. Raw data were processed at the sample level using the abagen toolbox

(https://abagen.readthedocs.io/en/stable)[49,50], following a standardized and widely recommended

pipeline[49]. Probe reannotation was performed based on data provided in reference[49], and probes

lacking Entrez Gene IDs were excluded. To ensure reliability, only probes that exceeded background

expression levels in at least 30% of all tissue samples were retained. Among these, the probe with

the highest differential stability was selected to represent each gene. After filtering and selection, a

total of 16,383 genes were retained. Within each donor, gene expression values were first z-scored

across genes within each sample, and then z-scored across all tissue samples.

Cell-type deconvolution. Single-nucleus droplet-based sequencing (snDrop-seq) data[23] were

obtained from the CellxGene-Census (https://cellxgene.cziscience.com/collections/d17249d2-

0e6e-4500-abb8-e6c93fa1ac6f). Cell-type deconvolution was performed following the pipeline

described previously[32]. Gene identifiers were harmonized by matching Entrez Gene IDs between

the snDrop-seq and AHBA datasets, retaining only shared genes. To reduce computational load and

enhance specificity, we limited the gene set to reliable marker genes consistently identified across

eight cortical regions, as reported previously[23]. Transcriptionally similar cell types were grouped according to the groupings across area subclasses[23,52] to mitigate collinearity in downstream analyses. The aggregated snDrop-seq data across all eight cortical areas were used as a reference for cell-type abundance deconvolution on each AHBA bulk-tissue sample via CIBERSORTx (https://cibersortx.stanford.edu/)[93], a strategy also adopted in similar studies[26,32]. This resulted in estimated abundances for 24 distinct cell types across cortical samples, including L2/3IT, L4IT, L6IT, L6ITCar3, L5IT, L5ET, L5/6NP, L6CT, L6b, Chandelier, PVALB, SST, SSTCHODL, LAMP5, SNCG, PAX6, LAMP5LHX6, VIP, Astro, Endo, OPC, VLMC, Micro/PVM, and Oligo. For each AHBA donor, estimated cell-type abundances were mapped into MNI space at $3 \times 3 \times 3$ mm³ isotropic resolution using the corresponding MNI coordinates of the tissue samples. Samples mapped to the same voxel were averaged. To maximize sample retention, averaging across donors was performed only for overlapping samples, yielding a final set of 1,569 samples. The resulting group-level cell-type maps were visually consistent with parcellation-level maps reported previously[32].

Molecular atlas. We included two sets of molecular maps representing neurotransmitter systems and mitochondrial phenotypes. The neurotransmitter receptor and transporter density maps were estimated via 19 PET-derived tracer images from https://github.com/netneurolab/hansen_receptors[25]. The analyzed receptors and transporters included 5-HT$_{1A}$, 5-HT$_{1B}$, 5-HT$_{2A}$, 5-HT$_4$, 5-HT$_6$, 5-HTT, D$_1$, D$_2$, DAT, NET, H$_3$, α$_4$β$_2$, M$_1$, VAChT, CB$_1$, MOR, NMDA, mGluR$_5$, and GABA$_A$. Six mitochondrial phenotype images were obtained from https://neurovault.org/collections/16418/[21]. These phenotypes included CI, CII, CIV, MitoD, TRC, and MRC. All molecular images were coregistered to MNI space with $3 \times 3 \times 3$ mm³ isotropic resolution, matching the resolution of the ICN and gene expression data to ensure consistent spatial alignment across modalities.

Cognitive probabilistic measures. We utilized probabilistic maps from Neurosynth (https://github.com/neurosynth/neurosynth)[48] to obtain voxel-wise cognitive and behavioral density maps. These maps were generated from meta-analyses of over 15,000 published fMRI studies and quantify the probability that a given cognitive term is reported in a study, conditional on the presence of activation at a specific voxel. In total, 123 cognitive and behavioral terms were selected based on

the Cognitive Atlas[56]. All cognitive association maps were resampled and aligned to MNI space

with $3 \times 3 \times 3$ mm³ isotropic resolution, consistent with the spatial resolution used for functional,

molecular, and gene expression data, ensuring cross-modality comparability.

Linking ICNs with cellular and molecular atlases. To investigate the potential cellular and

molecular underpinnings of individual ICNs, we assessed the spatial associations between each ICN

map and the cellular and molecular atlases. For each ICN, we applied a threshold of $|z| > 3$ (three

standard deviations; a conventional choice to emphasize spatial specificity in ICA maps[100]) to

exclude weakly activated regions and to focus on spatially specific patterns. Only voxels above the

threshold were included in the voxelwise spatial correlation analyses. Because the cellular atlas is

cortex-limited and discrete sampled, we implemented a local neighborhood averaging approach: for

each voxel in the cellular atlas that overlapped with the thresholded ICN map, we extracted ICN

values ($|z| > 3$) within a 3-mm radius and computed their mean, yielding a matched spatial

representation that preserved localization while reducing noise from sampling sparsity. Accordingly,

ICN-to-cell-type associations were computed only within cortical regions. Spatial similarity

between ICN maps and cell-type distributions was quantified using Spearman's correlation. To

account for spatial autocorrelation, we performed a two-sided Moran test[53] with 1,000 permutations.

For each ICN spatial map, the spatial weights matrix was defined by Euclidean distance proximity.

Moran spectral randomization was then used to generate 1,000 spatially autocorrelated surrogate

maps that matched the observed degree of spatial dependence (Moran's I). Significance was

determined by comparing observed statistics to null distributions, with FDR correction applied

across 24×44 comparisons. For the neurotransmitter and mitochondrial atlases, we employed

multivariate linear regression models[25] to estimate the contribution of each molecular map to ICN

spatial variance. ICN intensity maps were treated as the outcomes, and the mitochondrial or

neurotransmitter profiles were entered as predictors. The adjusted r^2 was used to quantify the model

fit. To evaluate the relative importance of each feature, we used the relaimpo R package, which

partitions the explained variance across predictors[54]. The significance of adjusted r^2 and relative

importance was determined by one-sided Moran test[53] with 1,000 permutations. The p-values of

relative importance were corrected for (19×53 for neurotransmitter; 6×53 for mitochondrion)

comparisons using the FDR method. Robustness was assessed at alternative thresholds of 2.58 (two-

sided $p = 0.01$) and 3.29 (two-sided $p = 0.001$), yielding consistent correlation results of spatial

similarity across cell-type (*Supplementary Fig. S14*), neurotransmitter (*Supplementary Fig. S15*),

and mitochondrion (*Supplementary Fig. S16*).

To investigate the synergy and relative contributions of distinct neurobiological scales to ICNs,

we fitted a multi-scale model (ICN ~ all cell-type, neurotransmitter, and mitochondrion predictors)

and single-scale models (one system at a time). To mitigate collinearity among cell-type

proportional data while preserving interpretability, we applied factor analysis (minimum residuals,

promax oblique rotation)[37]. We retained all unrotated factors that explained ≥1% of variance, and

we named factors by assigning each original atlas to the factor on which it loaded most strongly.

Oblique rotation was chosen because intercorrelations are expected among non-independent

biological processes and cell populations. To align modalities spatially, neurotransmitter and

mitochondrial maps were resampled to the discrete cortical sampling of the cell-type data using the

local neighborhood averaging approach; the same sampled inputs were used when fitting the single-

system models to ensure fair comparisons. Model improvement was assessed with *F*-tests

comparing the multi-scale model to each single-scale model, with FDR applied across 3×44 + 2×9

comparisons. Scale-level relative contributions were defined as the sum of contributions from all

sub-markers within each scale.

Associations between FNC and cellular/molecular similarity networks. We averaged cellular

and molecular features across the 53 ICNs ($|z| > 3$) and constructed feature similarity networks for

cell types, neurotransmitter receptor densities, mitochondrial phenotypes, and combined molecular

fingerprints using Spearman's correlation between ICNs. Robustness was assessed at alternative

thresholds of 2.58 (two-sided $p = 0.01$) and $z = 3.29$ (two-sided $p = 0.001$), yielding consistent

results (*Supplementary Fig. S17*). To assess whether these similarity networks recapitulate

functional organization, we performed a rank-sum test comparing within-domain versus between-

domain connections. Importantly, to control for spatial autocorrelation, we regressed out spatial

proximity (both Euclidean distance and Dice similarity) between ICN spatial maps from all

connectivity matrices. Next, we assessed the global and regional alignment between the FNC and

each cellular/molecular similarity network using partial Spearman's correlation, again controlling

for spatial proximity. For regional correlation analyses, multiple comparisons were controlled using

FDR across the tested ICNs, 44 for cell-type analyses and 53 for neurotransmitter, mitochondrial, and combined models.

Structural associations between ICNs and cellular/molecular architecture. To investigate whether the functional domain structure of ICNs is underpinned by cellular and molecular organization, we first constructed cellular and molecular similarity networks while controlling for spatial autocorrelation (Euclidean distance and Dice similarity) across ICNs. We then applied a diffusion embedding method[55] to denoise and reduce the similarity network into a low-dimensional manifold representation. A three-component solution was selected via the elbow criterion, which preserved global structure while mitigating the influence of noise or local fluctuations on subsequent clustering. Effectiveness was verified by comparison with principal component analysis (PCA, first three components) and with analyses conducted without dimensionality reduction. We then applied k-means clustering to group the projected features. The number of clusters was determined based on the number of known functional domains: 5 clusters for the cell-type similarity network and 7 clusters for the other similarity networks. The cluster labels derived from this unsupervised embedding were compared to the true functional domain labels of the ICNs. The ARI was used to quantify the accuracy of the clustering, and 1,000 label permutations were performed to assess the statistical significance of the results.

Mediation analysis. For cellular/molecular maps and ICNs that showed significant spatial correspondence in the primary analyses, we then conducted mediation analyses using the PROCESS toolbox in R. In these analyses, the cellular/molecular maps served as the predictor, the cognitive probabilistic measures were the outcome, and the ICN acted as the mediator. The significance of the mediation was estimated by a bias-corrected bootstrap approach with 1,000 random samplings. FDR was applied to correct the p-values for 123×30 comparisons for mediation analysis. To further investigate whether inter-ICN functional connectivity (FNC and dynamic FNC) mediates the relationship between cellular/molecular similarity and cognitive similarity, we first averaged the probabilistic values of each cognitive map across the 53 ICNs ($|z| > 3$) to obtain a single feature vector for each cognitive term. We then constructed cognitive similarity networks by computing Spearman's correlation between ICNs. Partial Spearman's correlations were performed to assess associations between cellular/molecular similarity networks and the cognitive similarity network,

as well as between (static) FNC or dynamic FNC states and the cognitive similarity network. To

account for spatial confounds, Euclidean distance and Dice similarity were included as covariates

in all correlation and mediation analyses. Mediation analyses were conducted using the PROCESS,

with the cellular/molecular similarity network specified as the predictor, the cognitive similarity

network as the outcome, and FNC or dynamic FNC as the mediator. The 95% confidence intervals

were estimated using a bias-corrected bootstrap with 1,000 resamples. For the dynamic FNC models,

multiple comparisons were controlled using FDR across 5×4 tests.

Figure Legends

Fig. 1 | ICNs and cellular and molecular process. A. Fifty-three robust and replicable ICNs were identified and organized into seven canonical functional domains using Neuromark_fMRI_1.0[7], derived from rs-fMRI data of 1,828 young adults across the GSP and HCP cohorts. Each ICN comprises brain voxels that co-vary over time, capturing intra-network coherence. Functional connectivity between ICNs (FNC) was derived by quantifying temporal correlations between ICN TCs, reflecting inter-network coupling. **B.** A diverse set of cellular and molecular maps derived from transcriptomic data (microarray and snRNA-seq) and molecular imaging data (PET), illustrate regional variations in cellular composition, mitochondrial metabolism, and neurotransmitter signaling, providing a multiscale context for interpreting ICN organization.

Fig. 2. | Cell-type associations of ICNs. Imputed spatial distributions of 24 cortical cell types were derived by integrating bulk tissue transcriptomic data from the AHBA[22] with snRNA-seq from eight cortical regions[23,52]. Voxel-wise abundance maps were correlated with the spatial patterns of 44 cortical ICNs using Spearman's correlation. Statistical significance was determined using a two-sided Moran test[53] (1,000 iterations), with FDR correction applied across 24×44 comparisons. **A.** Eight significant cell type–ICN associations (positive and negative). **B.** Cortical distribution maps of the 24 cell types.

Fig. 3. | Neurotransmitter associations of ICNs. A multilinear regression framework was used to evaluate the spatial correspondence between 53 ICNs and the distribution of 19 neurotransmitter receptors and transporters, as estimated by PET imaging[25]. Dominance analysis was performed to quantify the unique variance in each ICN map attributable to individual receptor/transporter densities[54]. Statistical significance was assessed using one-sided Moran test (1,000 permutations) to account for spatial autocorrelation[53], with FDR correction applied across 53 adjusted r^2 comparisons and 19×53 relative contribution comparisons. **A.** Sixteen ICNs demonstrated significant model fits $(0.43 \leqslant$ adjusted $r^2 \leqslant 0.86)$. Box plots show the median and interquartile range (IQR; 25–75%), with whiskers indicating 1.5× IQR from the first or third quartile. **B.** Dominance analysis showed

655   that variability across ICNs was selectively explained by distinct receptor/transporter profiles.

Fig. 4. | Cellular and molecular similarity networks relate to functional connectivity. A. Similarity networks were constructed for cell-type composition, mitochondrial metabolism, neurotransmitter receptor density, and an integrated molecular profile by computing pairwise Spearman correlations between the spatial fingerprints of all ICN pairs. **B.** Within-domain versus between-domain similarity was compared across these networks using a rank-sum test, with confounding effects of spatial proximity (Euclidean distance and Dice similarity) regressed out. Box plots show the median and interquartile range (IQR; 25–75%), with whiskers indicating 1.5× IQR from the first or third quartile. **C.** Mean FNC. Global (**D**) and regional (**E**) associations between

666   FNC and cellular/molecular similarity networks were evaluated using partial Spearman's

667   correlation, controlling for spatial proximity. In panel **E**, filled bars indicate statistically significant

668   associations ($q < 0.05$), whereas unfilled bars denote non-significant results. Note: r_1 denotes the

669   correlation without controlling for spatial proximity; r_2 denotes the correlation after controlling for

670   spatial proximity. **: $p < 0.01$; ***: $p < 0.001$.

Fig. 5. | Structural associations between ICNs and cellular/molecular architecture. Based on constructed cellular and molecular similarity networks controlling for spatial proximity across ICNs, we applied a diffusion embedding method[55] to project these similarity networks into a shared three-dimensional space. K-means clustering was then performed on the embedded features, with the number of clusters determined by the number of established functional domains: five clusters for the cell-type similarity network (**A**), and seven for the neurotransmitter (**B**), mitochondrial (**C**), and combined (**D**) networks. The labels derived from this unsupervised clustering were compared to the true functional domain labels of the ICNs. The ARI was used to quantify the accuracy of the clustering, and 1,000 label permutations were performed to assess the statistical significance of the results.

Fig. 6 | ICNs mediate the spatial associations between cellular and molecular architecture and cognitive function. We used 123 probabilistic cognitive activation maps derived from meta-analyses in Neurosynth[48]. For cellular/molecular maps and ICNs that showed significant spatial correspondence in the primary analyses, mediation analyses were conducted using the PROCESS toolbox in R, with cellular/molecular maps as predictors, ICNs as mediators, and cognitive maps as outcomes. Edges in the diagram represent significant mediation pathways ($q < 0.05$), with color indicating the magnitude of the mediation effect.

Fig. 7 | FNC mediate the connectional associations between cellular and molecular architecture and cognitive function. Based on 123 probabilistic cognitive activation maps derived from meta-analyses in Neurosynth[48], a cognitive similarity network was constructed by computing pairwise Spearman correlations between the spatial fingerprints of all ICN pairs. Mediation analyses at the network level treated cellular/molecular similarity as the predictor, FNC as the mediator, and cognitive similarity as the outcome. Significant mediation effects were observed for cell-type (A), neurotransmitter (B), and combination (C) similarity networks, whereas the mitochondrion similarity network (D) did not yield a significant mediation path.

Data and Code availability

The HCP dataset is publicly available at http://www.humanconnectomeproject.org/data. The

Neuromark_fMRI_1.0 template is available at http://trendscenter.org/software/gift. The AHBA data

can be accessed at http:// human.brain-map.org/ and the sn-RNA dataset is available at

https://cellxgene.cziscience.com/collections/d17249d2-0e6e-4500-abb8-e6c93fa1ac6f.

Neurotransmitter PET imaging can be found at https://github.com/netneurolab/hansen_receptors

and mitochondrial phenotype images can be found at https://neurovault.org/collections/16418/.

SPM12 is available at http://www.fil.ion.ucl.ac.uk/spm/ , and the GIFT toolbox can be accessed at

http://trendscenter.org/software/gift. The resulting data and code relevant to this study can be

accessed through the following links: https://trendscenter.org/data/,

https://github.com/trendscenter/ICNs-annotation and https://github.com/FelixFengCN/ICNs-

annotation. Other codes of this study can be obtained from the corresponding author.

Acknowledgement

The authors thank all the volunteers for their participation in the study and anonymous reviewers for their insightful comments and suggestions. This work was supported by NSF Grant #2112455 and NIH grant #R01MH123610. This work used transcriptomic data from the Allen Institute for Brain Science. Neuroimaging data were provided by the HCP, WU-Minn Consortium (principal investigators: D.V. Essen and K. Ugurbil; 1U54MH091657), funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research, and by the McDonnell Center for Systems Neuroscience at Washington University. The content of this manuscript solely reflects the views of the authors and may not reflect the views of the Allen Institute, NIH or the HCP consortium investigators.

Financial Disclosures

There are no conflicts of interest, including any financial, personal, or other relationships with people or organizations, for any of the authors related to the work described in the article.

References

References

1. Smith, S.M._, et al._ Functional connectomics from resting-state fMRI. _Trends Cogn Sci_ **17**,
666-682 (2013).

2. Zuo, X.N._, et al._ Human Connectomics across the Life Span. _Trends Cogn Sci_ **21**, 32-45
(2017).

3. Perovnik, M., Rus, T., Schindlbeck, K.A. & Eidelberg, D. Functional brain networks in the
evaluation of patients with neurodegenerative disorders. _Nat Rev Neurol_ **19**, 73-90 (2023).

4. Fornito, A., Zalesky, A. & Breakspear, M. The connectomics of brain disorders. _Nat Rev_
Neurosci **16**, 159-172 (2015).

5. Calhoun, V. Dynamic Interactivity Signatures of Mental Health Risk. _Biological Psychiatry_
97, S24 (2025).

6. Calhoun, V.D., Miller, R., Pearlson, G. & Adalı, T. The chronnectome: time-varying
connectivity networks as the next frontier in fMRI data discovery. _Neuron_ **84**, 262-274
(2014).

7. Du, Y._, et al._ NeuroMark: An automated and adaptive ICA based pipeline to identify
reproducible fMRI markers of brain disorders. _Neuroimage Clin_ **28**, 102375 (2020).

8. Fu, Z._, et al._ Searching Reproducible Brain Features using NeuroMark: Templates for
Different Age Populations and Imaging Modalities. _Neuroimage_ **292**, 120617 (2024).

9. Iraji, A._, et al._ Identifying canonical and replicable multi-scale intrinsic connectivity
networks in 100k+ resting-state fMRI datasets. _Hum Brain Mapp_ **44**, 5729-5748 (2023).

10. Erhardt, E.B., Allen, E.A., Damaraju, E. & Calhoun, V.D. On network derivation, classification,
and visualization: a response to Habeck and Moeller. _Brain Connect_ **1**, 1-19 (2011).

11. Joel, S.E., Caffo, B.S., van Zijl, P.C. & Pekar, J.J. On the relationship between seed-based
and ICA-based measures of functional connectivity. _Magn Reson Med_ **66**, 644-657 (2011).

12. Shine, J.M. Neuromodulatory Influences on Integration and Segregation in the Brain.
Trends Cogn Sci **23**, 572-583 (2019).

13. Zhang, D. & Raichle, M.E. Disease and the brain's dark energy. _Nat Rev Neurol_ **6**, 15-28
(2010).

14. Cossart, R. & Garel, S. Step by step: cells with multiple functions in cortical circuit assembly.
Nat Rev Neurosci **23**, 395-410 (2022).

15. Vanderhaeghen, P. & Polleux, F. Developmental mechanisms underlying the evolution of
human cortical circuits. _Nat Rev Neurosci_ **24**, 213-232 (2023).

16. Paquola, C., Amunts, K., Evans, A., Smallwood, J. & Bernhardt, B. Closing the mechanistic
gap: the value of microarchitecture in understanding cognitive networks. _Trends Cogn Sci_
26, 873-886 (2022).

17. Lee, A.T., Chang, E.F., Paredes, M.F. & Nowakowski, T.J. Large-scale neurophysiology and
single-cell profiling in human neuroscience. _Nature_ **630**, 587-595 (2024).

18. Bazinet, V., Hansen, J.Y. & Misic, B. Towards a biologically annotated brain connectome.
Nat Rev Neurosci (2023).

19. Hansen, J.Y. & Misic, B. Integrating and interpreting brain maps. _Trends Neurosci_ **48**, 594-
607 (2025).

20. Palomero-Gallagher, N. & Zilles, K. Cortical layers: Cyto-, myelo-, receptor- and synaptic
architecture in human cortical areas. _Neuroimage_ **197**, 716-741 (2019).

21. Mosharov, E.V., *et al.* A human brain map of mitochondrial respiratory capacity and
diversity. *Nature* (2025).

22. Hawrylycz, M.J., *et al.* An anatomically comprehensive atlas of the adult human brain
transcriptome. *Nature* **489**, 391-399 (2012).

23. Jorstad, N.L., *et al.* Transcriptomic cytoarchitecture reveals principles of human neocortex
organization. *Science* **382**, eadf6812 (2023).

24. Lake, B.B., *et al.* Integrative single-cell analysis of transcriptional and epigenetic states in
the human adult brain. *Nature biotechnology* **36**, 70-80 (2018).

25. Hansen, J.Y., *et al.* Mapping neurotransmitter systems to the structural and functional
organization of the human neocortex. *Nat Neurosci* **25**, 1569-1581 (2022).

26. Anderson, K.M., *et al.* Transcriptional and imaging-genetic association of cortical
interneurons, brain function, and schizophrenia risk. *Nat Commun* **11**, 2889 (2020).

27. Anderson, K.M., *et al.* Convergent molecular, cellular, and cortical neuroimaging
signatures of major depressive disorder. *Proc Natl Acad Sci U S A* **117**, 25138-25149
(2020).

28. Shine, J.M., *et al.* Human cognition involves the dynamic integration of neural activity and
neuromodulatory systems. *Nat Neurosci* **22**, 289-296 (2019).

29. Anderson, K.M., *et al.* Gene expression links functional networks across cortex and striatum.
Nat Commun **9**, 1428 (2018).

30. Richiardi, J., *et al.* BRAIN NETWORKS. Correlated gene expression supports synchronous
activity in brain networks. *Science* **348**, 1241-1244 (2015).

31. Krienen, F.M., Yeo, B.T., Ge, T., Buckner, R.L. & Sherwood, C.C. Transcriptional profiles of
supragranular-enriched genes associate with corticocortical network architecture in the
human brain. *Proc Natl Acad Sci U S A* **113**, E469-478 (2016).

32. Zhang, X.H., *et al.* The cell-type underpinnings of the human functional cortical
connectome. *Nat Neurosci* **28**, 150-160 (2025).

33. Hansen, J.Y., *et al.* Local molecular and global connectomic contributions to cross-
disorder cortical abnormalities. *Nat Commun* **13**, 4682 (2022).

34. Picciotto, M.R., Higley, M.J. & Mineur, Y.S. Acetylcholine as a neuromodulator: cholinergic
signaling shapes nervous system function and behavior. *Neuron* **76**, 116-129 (2012).

35. Kawai, H., Lazar, R. & Metherate, R. Nicotinic control of axon excitability regulates
thalamocortical transmission. *Nat Neurosci* **10**, 1168-1175 (2007).

36. Marder, E. Neuromodulation of neuronal circuits: back to the future. *Neuron* **76**, 1-11
(2012).

37. Lotter, L.D., *et al.* Regional patterns of human cortex development correlate with
underlying neurobiology. *Nat Commun* **15**, 7987 (2024).

38. Fu, Z., Sui, J., Iraji, A., Liu, J. & Calhoun, V.D. Cognitive and psychiatric relevance of dynamic
functional connectivity states in a large (N > 10,000) children population. *Molecular
Psychiatry* **30**, 402-413 (2024).

39. Zhi, D., *et al.* Triple Interactions Between the Environment, Brain, and Behavior in Children:
An ABCD Study. *Biol Psychiatry* **95**, 828-838 (2024).

40. Mathys, H., *et al.* Single-cell atlas reveals correlates of high cognitive function, dementia,
and resilience to Alzheimer's disease pathology. *Cell* **186**, 4365-4385 e4327 (2023).

41. Hara, Y., *et al.* Presynaptic mitochondrial morphology in monkey prefrontal cortex

correlates with working memory and is improved with estrogen treatment. *Proc Natl Acad*
Sci U S A **111**, 486-491 (2014).

42.   Sharpley, M.S., *et al.* Heteroplasmy of mouse mtDNA is genetically unstable and results in
altered behavior and cognition. *Cell* **151**, 333-343 (2012).

43.   Hollis, F., *et al.* Mitochondrial function in the brain links anxiety with social subordination.
Proc Natl Acad Sci U S A **112**, 15486-15491 (2015).

44.   Gebara, E., *et al.* Mitofusin-2 in the Nucleus Accumbens Regulates Anxiety and
Depression-like Behaviors Through Mitochondrial and Neuronal Actions. *Biol Psychiatry*
89, 1033-1044 (2021).

45.   Noudoost, B. & Moore, T. The role of neuromodulators in selective attention. *Trends Cogn*
Sci **15**, 585-591 (2011).

46.   Holmes, A.J., *et al.* Brain Genomics Superstruct Project initial data release with structural,
functional, and behavioral measures. *Sci Data* **2**, 150031 (2015).

47.   Van Essen, D.C., *et al.* The WU-Minn Human Connectome Project: an overview.
Neuroimage **80**, 62-79 (2013).

48.   Yarkoni, T., Poldrack, R.A., Nichols, T.E., Van Essen, D.C. & Wager, T.D. Large-scale
automated synthesis of human functional neuroimaging data. *Nat Methods* **8**, 665-670
(2011).

49.   Arnatkeviciute, A., Fulcher, B.D. & Fornito, A. A practical guide to linking brain-wide gene
expression and neuroimaging data. *Neuroimage* **189**, 353-367 (2019).

50.   Markello, R.D., *et al.* Standardizing workflows in imaging transcriptomics with the abagen
toolbox. *Elife* **10**(2021).

51.   Arnatkeviciute, A., Markello, R.D., Fulcher, B.D., Misic, B. & Fornito, A. Toward Best Practices
for Imaging Transcriptomics of the Human Brain. *Biol Psychiatry* **93**, 391-404 (2023).

52.   Jorstad, N.L., *et al.* Comparative transcriptomics reveals human-specific cortical features.
Science **382**, eade9516 (2023).

53.   Vos de Wael, R., *et al.* BrainSpace: a toolbox for the analysis of macroscale gradients in
neuroimaging and connectomics datasets. *Commun Biol* **3**, 103 (2020).

54.   Azen, R. & Budescu, D.V. The dominance analysis approach for comparing predictors in
multiple regression. *Psychol Methods* **8**, 129-148 (2003).

55.   Margulies, D.S., *et al.* Situating the default-mode network along a principal gradient of
macroscale cortical organization. *Proc Natl Acad Sci U S A* **113**, 12574-12579 (2016).

56.   Poldrack, R.A., *et al.* The cognitive atlas: toward a knowledge foundation for cognitive
neuroscience. *Frontiers in neuroinformatics* **5**, 17 (2011).

57.   Yu, Q., *et al.* Comparing brain graphs in which nodes are regions of interest or
independent components: A simulation study. *J Neurosci Methods* **291**, 61-68 (2017).

58.   Xu, J., *et al.* Functional network overlap as revealed by fMRI using sICA and its potential
relationships with functional heterogeneity, balanced excitation and inhibition, and
sparseness of neuron activity. *PLoS One* **10**, e0117029 (2015).

59.   Fu, Z., Liu, J., Salman, M.S., Sui, J. & Calhoun, V.D. Functional connectivity uniqueness and
variability? Linkages with cognitive and psychiatric problems in children. *Nature Mental*
Health **1**, 956-970 (2023).

60.   Yan, W., *et al.* A Brainwide Risk Score for Psychiatric Disorder Evaluated in a Large
Adolescent Population Reveals Increased Divergence Among Higher-Risk Groups Relative

to Control Participants. *Biol Psychiatry* **95**, 699-708 (2024).

61.   Du, Y., *et al.* Evidence of shared and distinct functional and structural brain signatures in
schizophrenia and autism spectrum disorder. *Commun Biol* **4**, 1073 (2021).

62.   Kim, E.J., Juavinett, A.L., Kyubwa, E.M., Jacobs, M.W. & Callaway, E.M. Three Types of
Cortical Layer 5 Neurons That Differ in Brain-wide Connectivity and Function. *Neuron* **88**,
1253-1267 (2015).

63.   A multimodal cell census and atlas of the mammalian primary motor cortex. *Nature* **598**,
86-102 (2021).

64.   Economo, M.N., *et al.* Distinct descending motor cortex pathways and their roles in
movement. *Nature* **563**, 79-84 (2018).

65.   Peng, H., *et al.* Morphological diversity of single neurons in molecularly defined cell types.
Nature **598**, 174-181 (2021).

66.   Sridharan, D., Levitin, D.J. & Menon, V. A critical role for the right fronto-insular cortex in
switching between central-executive and default-mode networks. *Proc Natl Acad Sci U S*
A **105**, 12569-12574 (2008).

67.   Urban-Ciecko, J. & Barth, A.L. Somatostatin-expressing neurons in cortical networks. *Nat*
Rev Neurosci **17**, 401-409 (2016).

68.   Riedemann, T. & Sutor, B. Cell-Type-Specific Effects of Somatostatin on Synaptic
Transmission in the Anterior Cingulate Cortex. *J Neurosci* **44**(2024).

69.   Shenhav, A., Botvinick, M.M. & Cohen, J.D. The expected value of control: an integrative
theory of anterior cingulate cortex function. *Neuron* **79**, 217-240 (2013).

70.   Menon, V. & Uddin, L.Q. Saliency, switching, attention and control: a network model of
insula function. *Brain Struct Funct* **214**, 655-667 (2010).

71.   Surmeier, D.J., Carrillo-Reid, L. & Bargas, J. Dopaminergic modulation of striatal neurons,
circuits, and assemblies. *Neuroscience* **198**, 3-18 (2011).

72.   Bolam, J.P., Hanley, J.J., Booth, P.A. & Bevan, M.D. Synaptic organisation of the basal
ganglia. *J Anat* **196 (Pt 4)**, 527-542 (2000).

73.   Leroy, C., *et al.* Assessment of 11C-PE2I binding to the neuronal dopamine transporter in
humans with the high-spatial-resolution PET scanner HRRT. *J Nucl Med* **48**, 538-546
(2007).

74.   Moriguchi, S., *et al.* PET Quantification of the Norepinephrine Transporter in Human Brain
with (S,S)-(18)F-FMeNER-D(2). *J Nucl Med* **58**, 1140-1145 (2017).

75.   Albin, R.L., *et al.* Regional vesicular acetylcholine transporter distribution in human brain:
A [(18) F]fluoroethoxybenzovesamicol positron emission tomography study. *The Journal*
of comparative neurology **526**, 2884-2897 (2018).

76.   Tasic, B., *et al.* Shared and distinct transcriptomic cell types across neocortical areas.
Nature **563**, 72-78 (2018).

77.   Hodge, R.D., *et al.* Conserved cell types with divergent features in human versus mouse
cortex. *Nature* **573**, 61-68 (2019).

78.   Goulas, A., *et al.* The natural axis of transmitter receptor distribution in the human cerebral
cortex. *Proc Natl Acad Sci U S A* **118**(2021).

79.   Brezina, V. Beyond the wiring diagram: signalling through complex neuromodulator
networks. *Philos Trans R Soc Lond B Biol Sci* **365**, 2363-2374 (2010).

80.   Hawrylycz, M., *et al.* Canonical genetic signatures of the adult human brain. *Nat Neurosci*

18, 1832-1844 (2015).

81.   Brodmann, K. Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien
dargestellt auf Grund des Zellenbaues. (Barth, 1909).

82.   Feng, G.*, et al.* Longitudinal development of the human white matter structural
connectome and its association with brain transcriptomic and cellular architecture.
Communications Biology **6**(2023).

83.   Feng, G.*, et al.* Spatial and temporal pattern of structure–function coupling of human brain
connectome with development. *eLife* **13**(2024).

84.   Buckner, R.L. & Krienen, F.M. The evolution of distributed association networks in the
human brain. *Trends Cogn Sci* **17**, 648-665 (2013).

85.   Zou, Y.*, et al.* Cell-type-specific optogenetic fMRI on basal forebrain reveals functional
network basis of behavioral preference. *Neuron* **112**, 1342-1357.e1346 (2024).

86.   Hansen, J.Y.*, et al.* Mapping gene transcription and neurocognition across human
neocortex. *Nat Hum Behav* **5**, 1240-1250 (2021).

87.   Robbins, T.W. & Arnsten, A.F. The neuropsychopharmacology of fronto-executive
function: monoaminergic modulation. *Annu Rev Neurosci* **32**, 267-287 (2009).

88.   Dum, R.P. & Strick, P.L. The origin of corticospinal projections from the premotor areas in
the frontal lobe. *J Neurosci* **11**, 667-689 (1991).

89.   Bush, G., Luu, P. & Posner, M.I. Cognitive and emotional influences in anterior cingulate
cortex. *Trends Cogn Sci* **4**, 215-222 (2000).

90.   Rangaraju, V., Lauterbach, M. & Schuman, E.M. Spatially Stable Mitochondrial
Compartments Fuel Local Translation during Plasticity. *Cell* **176**, 73-84.e15 (2019).

91.   Hwang, K., Bertolero, M.A., Liu, W.B. & D'Esposito, M. The Human Thalamus Is an
Integrative Hub for Functional Brain Networks. *J Neurosci* **37**, 5594-5607 (2017).

92.   Haber, S.N. & Knutson, B. The reward circuit: linking primate anatomy and human imaging.
Neuropsychopharmacology **35**, 4-26 (2010).

93.   Newman, A.M.*, et al.* Determining cell type abundance and expression from bulk tissues
with digital cytometry. *Nature biotechnology* **37**, 773-782 (2019).

94.   Chu, T., Wang, Z., Pe'er, D. & Danko, C.G. Cell type and gene expression deconvolution
with BayesPrism enables Bayesian integrative analysis across bulk and single-cell RNA
sequencing in oncology. *Nat Cancer* **3**, 505-517 (2022).

95.   Huuki-Myers, L.A.*, et al.* Benchmark of cellular deconvolution methods using a multi-
assay reference dataset from postmortem human prefrontal cortex. *bioRxiv* (2024).

96.   Smith, S.M.*, et al.* Network modelling methods for FMRI. *Neuroimage* **54**, 875-891 (2011).

97.   Friedman, J., Hastie, T. & Tibshirani, R. Sparse inverse covariance estimation with the
graphical lasso. *Biostatistics* **9**, 432-441 (2008).

98.   Allen, E.A.*, et al.* Tracking whole-brain connectivity dynamics in the resting state. *Cereb*
Cortex **24**, 663-676 (2014).

99.   Fu, Z.*, et al.* Characterizing dynamic amplitude of low-frequency fluctuation and its
relationship with dynamic functional connectivity: An application to schizophrenia.
Neuroimage **180**, 619-631 (2018).

100.   Correa, N.M., Li, Y.O., Adalı, T. & Calhoun, V.D. Canonical Correlation Analysis for Feature-
Based Fusion of Biomedical Imaging Modalities and Its Application to Detection of
Associative Networks in Schizophrenia. *IEEE J Sel Top Signal Process* **2**, 998-1007 (2008).