Title Page:

Cellular and molecular associations with intrinsic brain

organization

Guozheng Feng[1], Jiayu Chen[1], Jing Sui[2], Vince D. Calhoun[1*]

[1]Tri-Institutional Center for Translational Research in Neuroimaging and Data

Science (TReNDS), Georgia State University, Georgia Institute of Technology, and

Emory University, Atlanta, GA, USA;

[2]State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern

Institute for Brain Research, Beijing Normal University, Beijing, China.

[*]**Corresponding To:**

Vince D. Calhoun, PhD, Tri-Institutional Center for Translational Research in

Neuroimaging and Data Science (TReNDS), Georgia State University, Georgia

Institute of Technology, and Emory University, Atlanta, GA, USA; E-mail:

vcalhoun@gatech.edu.

Abstract

Understanding how cellular and molecular architecture underpins the large-scale organization of human brain function is a central challenge in neuroscience. By integrating transcriptomic (microarray and single-nucleus RNA-sequencing), molecular imaging, and neuroimaging datasets, we observe spatial correspondences indicating that the distributions of diverse cell types, neurotransmitter systems, and mitochondrial phenotypes align with intrinsic connectivity networks (ICNs). These associations extend beyond local correspondence to reflect network-level structure: inter-ICN similarity networks derived from cellular and molecular profiles recapitulate static and dynamic patterns of functional network connectivity (FNC), mirroring canonical functional domains. Mediation analyses reveal that specific ICNs mediate the relationship between microscale cell-type architecture and domain-specific cognitive processes, while FNCs capture mediating pathways linking cell-type and neurotransmitter similarity networks to cognitive organization. Together, our findings show that the brain's functional architecture systematically aligns with cellular and molecular organization, which may constrain functional network formation and contribute to the neural basis of cognition.

Keywords: cell types, intrinsic connectivity network, functional networks, cognitive function, transcriptomic, cellular and molecular regulation

Introduction

Advances in study of brain networks with resting-state functional MRI (rs-fMRI) have significantly deepened our understanding of both typical and atypical neural activity[1-6]. Among various analytic approaches, decomposition-based independent component analysis (ICA) enables the reliable extraction of intrinsic connectivity networks (ICNs), which reflect coherent patterns of spontaneous brain activity within distinct functions[7-9]. Each ICN comprises a spatial weight map paired with an associated time course (TC), capturing within-component connectivity in the blood oxygenation level-dependent (BOLD) signal[10]. Correlating these TCs yields between-component relationships, typically summarized as functional network connectivity (FNC)[11]. The macroscopic functional brain networks emerge from the dynamic interplay of distributed neural circuit architectures, whose region-specific implementations—distinct cellular compositions and specialized input–output motifs—necessitate tailored neuromodulatory regulation and mitochondrial bioenergetics to sustain network-specific information processing and integration[12-21]. Despite growing evidence for the relevance of these cellular and molecular factors, their specific contributions to the organization of functional networks remain largely uncharacterized.

The growing availability of high-resolution cellular and molecular atlases of the human brain— such as microarray-based transcriptional data from the Allen Human Brain Atlas (AHBA)[22], single-nucleus RNA sequencing (snRNA-seq) datasets from discrete brain regions[21,23,24], and positron emission tomography (PET) imaging of diverse neurotransmitter systems[25]—has created new opportunities to bridge microscale biology with macroscale brain organization. Previous work established spatial associations between the distribution of interneuron-linked genes and regional differences in fMRI signal variability[26,27]. The low-dimensional manifold organizing fMRI activity shows opposing associations with regional densities of facilitatory versus inhibitory neuromodulatory receptors[28]. In parallel, specific mitochondrial phenotypes selectively align with distinct fMRI signal metrics (e.g., maximum BOLD, regional homogeneity, entropy)[21]. Beyond spatial alignment, molecular similarity further supports the organization of functional networks: functionally coupled regions tend to share gene-expression signatures[29-31], and distinct cell-type distributions differentially align with the cortex's principal unimodal-to-transmodal functional gradient[32]. Concordantly, neurotransmitter receptor and transporter similarity across regions

correlates with functional connectivity, with brain areas exhibiting similar receptor profiles showing

stronger co-activation[25]. Although most prior studies have focused on individual neurobiological

scales, observed correlations among neurobiological scales[25,33] suggest that multi-scale factors are

likely to synergistically contribute to shape functional network phenotypes. One illustrative case is

cholinergic neurotransmission primarily modulates neuronal excitability, regulates presynaptic

transmitter release, and orchestrates the firing of neuronal populations[34-36]. Additionally,

developmental patterns of cortical thickness are better captured by models that integrate the spatial

distributions of several systems (e.g., cell-type composition, neurotransmitter receptor/transporter

densities), with neurotransmitter markers showing relatively greater scale-level relative

importance[37]. Together, integrating diverse cellular and molecular atlases provides a multi-scale

perspective for revealing the biological foundations of intrinsic functional networks.

The human brain supports a wide range of cognitive and behavioral functions through large-

scale neural circuits shaped by its underlying cellular and molecular architecture[17]. At the

macroscale, ICNs, as the core organizational units of functional architecture, along with their static

and dynamic FNCs, demonstrate heterogeneous contributions to diverse cognitive abilities[38,39]. At

the microscale, specific inhibitory neuron subtypes are selectively preserved in individuals who

maintain high cognitive performance in late life, implicating cell-type-specific influences on

cognitive resilience[40]. The spatial distribution of neurotransmitter receptors differentiates intrinsic

and extrinsic cognitive modes[25]. Animal studies further underscore the role of molecular processes

in cognition and behavior: mitochondrial function has been shown to impact both cognitive

capacity[41,42] and behavioral regulation[43,44], while distinct neurotransmitter receptor systems have

been associated with specific cognitive domains[45]. Despite these converging findings, the

mechanisms through which ICNs and FNCs mediate the relationship between microscale biological

substrates and cognitive and behavior abilities remain poorly understood. Uncovering this linkage

is essential for a deeper understanding of how cellular and molecular systems scaffold the brain's

functional architecture and enable complex cognition.

Here, motivated by the above evidence, we aim to probe the cellular and molecular bases of

the spatial and connectional architecture of ICA-derived ICNs from a multi-scale perspective. For

this purpose, we integrate spatial maps of cell types, neuromodulatory receptor/transporter systems,

and mitochondrial phenotypes. We hypothesize that ICN spatial maps colocalize with these

microscale architectures—such that networks are differentially enriched for several biological

features that support network-specific information processing and integration. We further

hypothesize that between-region similarity network derived from these cellular and molecular

features is associated with FNC, consistent with an organizing role for these similarity networks.

Moreover, multimodal combinations of neurobiological features explain more ICN/FNC variance

than any single system, consistent with potential synergistic effects among distinct microscale

processes. Finally, we posit that ICNs and FNCs provide the macroscale substrates that statistically

link microscale neurobiology to cognitive/behavioral maps.

Results

We analyzed 53 reproducible of ICN spatial maps and their corresponding FNC (*Supplementary*

Fig. S1 and Fig. 1a) derived from the Neuromark_fMRI_1.0 template[7]. The NeuroMark template

was based on components that replicated across two large-scale datasets: the Genomics Superstruct

Project (GSP; $n = 1,005$; mean age = 21.54 ± 2.96 years; 411 males)[46] and the Human Connectome

Project (HCP; $n = 823$; mean age = 28.79 ± 3.68 years; 356 males)[47]. To investigate the cellular and

molecular architecture underlying ICNs, we systematically examined the spatial and connectional

correspondence between these functional networks and a diverse sets of cell-type[23],

neurotransmitter[25], and mitochondrial[21] maps (Fig. 1b). Extending this analysis, we incorporated

123 cognitive and behavioral probabilistic maps from Neurosynth[48] to assess whether functional

networks (ICNs and FNCs) mediate the relationship between microscale biological substrates and

spatial and connectional variations in cognitive and behavioral functions.

Cell-type associations of ICNs. To investigate the cellular architecture underlying ICNs, we

related postmortem gene expression–based estimates of cell-type distribution to the spatial maps of

44 cortical ICNs. Bulk tissue transcriptomic data from AHBA[22] were processed using an established

harmonization pipeline[49-51]. Cell-type–specific transcriptional signatures were derived from

snRNA-seq of eight cortical areas[23], enabling the imputation of 24 cell-type abundances across

AHBA samples. These included nine excitatory glutamatergic neurons (e.g., L2/3 IT, L4IT, L6IT),

nine inhibitory GABAergic interneurons (e.g., Chandelier, PVALB, SST), and six non-neuronal cell

types (e.g., Astro, Endo, OPC), each characterized by distinct laminar locations, developmental

origins, morphology, and projection targets[23,52]. Voxel-level spatial distributions of cell-type

abundance were correlated with ICN spatial maps using Spearman's r correlation. Statistical

significance was determined using permutation-based two-sided Moran test[53] (1,000 iterations),

with false discovery rate (FDR) correction applied across 24×44 comparisons.

Seven ICNs exhibited at least one significant association with the imputed abundance of

specific cell-types (Fig. 2a). These associations included both positive and negative correlations,

indicating that cell-type gradients may support the spatial organization of ICNs. IC14 (precentral

gyrus network) in the sensorimotor network (SM) domain was positively associated with L5ET (r

$(107) = 0.48$, $q < 0.001$). IC25 (middle temporal gyrus network) in the visual network (VIS) was

positively associated with L2/3IT ($r (140) = 0.30$, $q < 0.001$). In the cognitive-control network (CC)

domain, IC27 (insula network) was significantly associated with L6ITCar3 ($r (127) = 0.58$, $q <$

0.001), and IC28 (superior medial frontal gyrus network) was negatively associated with L4IT (r

$(123) = -0.58$, $q < 0.001$), IC31 (middle frontal gyrus network) was positively associated with

LAMP5 ($r (110) = 0.49$, $q < 0.001$), IC40 (inferior frontal gyrus network) was negatively associated

with L5/6NP ($r (88) = -0.40$, $q < 0.001$). IC45 (anterior cingulate cortex network) in the default

mode domain was positively associated with SST ($r (139) = 0.32$, $q < 0.001$) and negative

associations with OPC ($r (139) = -0.25$, $q < 0.001$). The spatial distributions for all 24 cell types are

presented in Fig. 2b. Collectively, these findings indicate that particular ICNs show specific spatial

correspondences with distinct cell types.

Neurotransmitter and mitochondrial associations of ICNs. To examine the molecular

architecture underlying ICNs, we assessed the spatial alignment between ICN maps and

neurotransmitter systems using PET-derived estimates of receptor and transporter densities. We

compiled 19 neurotransmitter maps encompassing nine neurotransmitter systems, aggregated from

PET imaging data across more than 1,200 healthy individuals[25]. These maps included serotonergic

($5-HT_{1A}$, $5-HT_{1B}$, $5-HT_{2A}$, $5-HT_4$, $5-HT_6$, 5-HTT), dopaminergic (D_1, D_2, DAT), noradrenergic

(NET), histaminergic (H_3), cholinergic ($\alpha_4\beta_2$, M_1, VAChT), cannabinoid (CB_1), opioid (MOR),

glutamatergic (NMDA, $mGluR_5$), and GABAergic ($GABA_A$) systems.

For each ICN, we constructed multilinear regression models using the full set of receptor and

transporter densities as predictors. Statistical significance was assessed using a one-sided Moran

test (1,000 permutations) to account for spatial autocorrelation[53], FDR correction applied across 53

comparisons. Sixteen ICNs exhibited significant model fits (adjusted r^2: 0.43–0.86, $q < 0.05$, Fig.

3a), indicating that distinct combinations of neurotransmitter systems spatially align with specific

ICN architecture; effects were densest in SC and AUD domains. To determine the relative

contribution of individual neurotransmitter systems, we applied dominance analysis, which

quantifies the proportion of variance in each ICN map attributable to each receptor/transporter in

the model[54]. Dominance scores were normalized by the model's total adjusted r^2, enabling

comparison across ICNs (Fig. 3b). Statistical significance of neurotransmitter-specific dominance

was assessed using one-sided Moran test (1,000 permutations) to account for spatial

autocorrelation[53], FDR correction applied across 19×53 comparisons. Variance contributions of

neurotransmitter receptors were heterogeneous across ICNs. For example, in CC domain, the $\alpha_4\beta_2$

receptor made a prominent contribution to IC31 (middle frontal gyrus network, contribution = 26%,

$q < 0.001$), while CB_1 receptor made a prominent contribution to IC37 (Hippocampus network,

contribution = 19%, $q < 0.001$). In DM domain, $5\text{-}HT_{1B}$ receptor showed significant influence on

IC48 (precuneus network, contribution = 25%, $q < 0.001$).

In parallel, we examined the association between mitochondrial phenotypes and ICNs. Six

mitochondrial maps (complex I [CI], complex II [CII], complex IV [CIV], mitochondrial density

[MitoD], tissue respiratory capacity [TRC], and mitochondrial respiratory capacity [MRC])[21] were

regressed onto ICN spatial maps, FDR correction applied across 53 comparisons for adjusted r^2 and

6×53 comparisons for dominance. We observed significant model fits for IC2 (adjusted $r^2 = 0.25$, p

< 0.001) and IC4 (caudate network, adjusted $r^2 = 0.24$, $p < 0.001$), both located in SC domain

(*Supplementary Fig. S2a*). Within these models, MitoD, and CIV each made significant relative

contributions (*Supplementary Fig. S2b*). These findings suggest that mitochondrial energy

metabolism may contribute to the functional architecture of specific ICNs in SC domain.

To investigate the synergy and relative contributions of distinct neurobiological scales to ICNs,

we examined associations between multi-scale markers (cell types [21, after collinearity attenuation,

Supplementary Fig. S3; see Methods], neurotransmitters [19], and mitochondria [6]) and ICNs.

Fourteen ICNs exhibited significant model fits (adjusted r^2: 0.62–0.99, $q < 0.05$; *Supplementary Fig.*

S4a). In 13 of these 14 ICNs, multi-scale models outperformed single-scale models ($q < 0.05$; *Supplementary Fig. S4b*), underscoring the synergistic value of integrating distinct neurobiological scales. We further observed a consistent trend of synergistic contributions from markers across scales ($p < 0.05$, *Supplementary Fig. S4a*). Among these, neurotransmitter systems consistently accounted for the largest overall relative contributions across scales (*Supplementary Fig. S4b*).

Cellular and molecular similarity networks relate to functional connectivity. Having established that ICNs exhibit distinct spatial associations with cellular and molecular features, we next asked whether ICNs that are similarly composed at the cellular and molecular levels also exhibit stronger functional interactions. To this end, we computed similarity networks for cell-type, mitochondrial, neurotransmitter, and combined molecular fingerprints by correlating the mean spatial profiles of each ICN pair using Spearman's correlation (Fig. 4a). Across all modalities, cellular and molecular similarities were significantly greater among ICNs within the same functional domain than between different domains, even after regressing out Euclidean distance and spatial overlap (Dice coefficient) from the similarity matrices ($p < 0.001$, ranksum test; $N_{within} = 230$ edges, $N_{between} = 716$ edges for cell-type; $N_{within} = 246$ edges, $N_{between} = 1,132$ edges for others, Fig. 4b). We then assessed the relationship between cellular/molecular similarity and functional connectivity, as indexed by the group-level FNC from the HCP dataset (Fig. 4c). We observed that cell-type similarity (Spearman's r (946) = 0.19, $p < 0.001$), neurotransmitter similarity (Spearman's r (1378) = 0.27, $p < 0.001$), and the combined similarity (Spearman's r (1378) = 0.46, $p < 0.001$) networks were all significantly correlated with FNC strength (Fig. 4d). At the regional level, correlations were predominantly positive, with the strongest associations observed in nodes located within the SM and VIS domains (Fig. 4e). To further investigate the dynamic aspects of this relationship, we identified five discrete states of dynamic FNC (*Supplementary Fig. S5a*) and examined their correspondence with cellular and molecular similarity networks. Results consistently demonstrated robust associations between FNC states and similarity networks, both at the global (*Supplementary Fig. S5b–f*) and regional levels (*Supplementary Fig. S6a–e*). Together, these results suggest that ICNs with similar cellular and molecular compositions are more likely to be functionally coupled, independent of spatial proximity, supporting the idea that inter-regional communication is constrained by a common microscale architecture.

Identify cellular and molecular structural associations with functional domain. To test whether cellular and molecular profiles reflect the functional domain architecture of ICNs, we constructed similarity networks based on cell-type, neurotransmitter, mitochondrion and combination fingerprints, while regressing out the effects of Euclidean distance and Dice similarity from the network matrices. We reduced network connectivity features using diffusion embedding[55], and projected each similarity network into a three-dimensional space (cumulative variance explained: 57% for cell type, 51% for neurotransmitter, 54% for mitochondrion, and 50% for the combined profile). Component-wise explained variance is reported in *Supplementary Fig. S7*. Comparisons of dimensionality reduction methods are shown in *Supplementary Fig. S8.* We then applied k-means to cluster features of cellular/molecular similarity network. The ground-truth labels were derived from the functional domain assignments of ICNs, and clustering performance was quantified using the adjusted rand index (ARI), with significance assessed via a null generated from 1,000 label permutations. Clustering based on cell-type (ARI = 0.34, $p < 0.001$, 5 clusters; Fig. 5a), neurotransmitter (ARI = 0.34, $p < 0.001$, 7 clusters; Fig. 5b), mitochondrion (ARI = 0.26, $p < 0.001$, 7 clusters; Fig. 5c), and combined profiles (ARI = 0.26, $p < 0.001$, 7 clusters; Fig. 5d) all showed significant alignment with functional domain structures. These findings indicate that the spatial organization of functional domains is supported by coherent cellular and molecular architectures.

ICNs mediate spatial associations between cellular/molecular architecture and cognitive function. To examine whether and how ICNs mediate the relationship between microscale cellular and molecular organization and macroscale cognitive architecture, we leveraged probabilistic meta-analytic activation maps from Neurosynth (https://github.com/neurosynth/neurosynth)[48]. These maps encode the likelihood that a given cognitive term is associated with activation at each voxel, based on over 15,000 published fMRI studies. A total of 123 cognitive and behavioral terms were selected based on their functional annotations in the Cognitive Atlas[56]. These term-wise maps showed significant spatial correspondence with specific ICNs (*Supplementary Fig. S9*). For cellular/molecular maps and ICNs that showed significant spatial correspondence in the primary analyses, we then conducted post-hoc mediation analyses using the PROCESS toolbox in R. In each model, the cellular/molecular map served as the predictor, the cognitive probability map as the outcome, and the ICN as the mediator. FDR correction applied across 123×30 comparisons.

These analyses revealed significant mediation pathways, predominantly involving 7 ICNs, 10

cellular/molecular maps, and 123 cognitive and behavior functions. Among cortical ICNs (Fig. 6),

the insula network (IC27; L6ITCar3-coupled) showed the broadest and strongest mediation

spectrum (mediation effect β: 0.21–0.57), spanning language (speech perception/production,

naming), sensory integration (multisensory, categorization), executive control (cognitive control,

inhibition, decision making), and emotion-related processes (reward anticipation, anxiety, fear). The

precentral gyrus network (IC14; L5ET-coupled) exhibited weaker mediation for executive/memory

functions but stronger effects on perceptual attention and movement/motor control (mediation effect

β: 0.10–0.23). The superior medial frontal gyrus network (IC28; L4IT-coupled) showed modest

effects focused on planning and attention, with additional links to movement (mediation effect β:

0.11–0.19). Finally, the anterior cingulate cortex network (IC45; SST-coupled) preferentially

mediated affective and motivational processes, including fear, anxiety, mood regulation, and

reinforcement learning (mediation effect β: 0.13–0.21). By comparison, neurotransmitter and

mitochondrial markers primarily mediated subcortical ICN–cognition associations (Source Data

Table S1). Overall, these spatial association–mediation results indicate that specific

cellular/molecular markers interface with functional networks to provide indirect links to cognitive

and behavioral phenotypes.

FNCs mediate the connectional associations between cellular and molecular architecture

and cognitive function. To further investigate whether inter-ICN functional connectivity (static and

dynamic FNCs) mediates the relationship between cellular/molecular similarity and cognitive

similarity, we first averaged the cognitive probabilistic maps across the 53 ICNs to obtain ICN-level

cognitive features. We then constructed cognitive similarity networks by computing Spearman's

correlation across ICN pairs. Partial Spearman's correlations were performed to assess associations

between cellular/molecular similarity networks and the cognitive similarity network, as well as

between static and dynamic FNC states and the cognitive similarity network. To account for spatial

confounds, Euclidean distance and Dice similarity were included as covariates in all correlation and

mediation analyses. We found that the cell-type, neurotransmitter, and combined similarity networks

were significantly associated with the cognitive similarity network ($p < 0.001$, *Supplementary Fig.*

S10). Both static FNC and dynamic FNC states showed significant associations with the cognitive

similarity network ($p < 0.001$, *Supplementary Fig. S11*). Mediation analyses at the network level

treated the cellular or molecular similarity network as the predictor, the cognitive similarity network

as the outcome, and the FNC (static or dynamic) as the mediator. The results revealed that static

FNC partially mediated the relationship between the cell-type similarity network and the cognitive

similarity network (path *ab*: $p = 0.002$, $\beta = 0.04$, 95% CI = [0.01, 0.06]; path *c'*: $p < 0.001$, $\beta = 0.31$,

95% CI = [0.26, 0.35]; path *c*: $p < 0.001$, $\beta = 0.31$, 95% CI = [0.30, 0.40], Fig. 7a). Static FNC also

mediated the link between the neurotransmitter similarity network and cognitive similarity network

(path *ab*: $p < 0.001$, $\beta = 0.05$, 95% CI = [0.03, 0.06]; path *c'*: $p < 0.001$, $\beta = 0.21$, 95% CI = [0.16,

0.25]; path *c*: $p < 0.001$, $\beta = 0.25$, 95% CI = [0.21, 0.29], Fig. 7b), as well as between the

combination similarity network and the cognitive similarity network (path *ab*: $p < 0.001$, $\beta = 0.08$,

95% CI = [0.07, 0.10]; path *c'*: $p < 0.001$, $\beta = 0.36$, 95% CI = [0.32, 0.41]; path *c*: $p < 0.001$, $\beta =$

0.45, 95% CI = [0.40, 0.49], Fig. 7d). No significant mediation was detected for the mitochondrial

similarity network (Fig. 7c). Mediation by dynamic FNC states was also significant as shown in

Supplementary Fig. S12. Specifically, states 2–5 mediated the relationship between cell-type and

cognitive similarity; states 3–5 mediated the neurotransmitter–cognition link; states 1, 3–5 mediated

the mitochondrion–cognition link. All states significantly mediated the association between

combination–cognition link. Together, these findings demonstrate that static and dynamic FNCs

provide indirect pathways linking cellular/molecular architecture to cognitive organization, with

dynamic FNC offering additional state-specific routes.

Discussion

By integrating transcriptomic, molecular, and neuroimaging data, we observed spatial

associations between the brain's large-scale functional architecture and its underlying cellular and

molecular organization. Spatial profiles of diverse cell types, neurotransmitter systems, and

mitochondrial markers show selective alignment with ICNs, providing biologically grounded

support for functional segregation. These relationships extend beyond local overlap: similarity

networks constructed from molecular and cellular features approximated patterns of functional

connectivity and captured core ICN functional domains. Importantly, ICNs statistically served as

intermediaries in the associations between molecular architecture and cognitive processes,

positioning them as critical intermediaries linking microscale biology to behavior. Together, these

findings suggest that cellular and molecular organization may impose constraints on the formation

and function of large-scale brain networks.

rs-fMRI investigations of brain networks have substantially expanded the frontiers of cognitive

neuroscience and psychiatric research. Among data-driven approaches, ICA leverages the hidden

spatiotemporal structure of the BOLD signal to separate meaningful neural signals from noise, while

preserving subject-level variability[57] and yielding improved signal-to-noise ratios[7]. Critically,

unlike parcellation-based approaches, ICA enables the delineation of distinct patterns of functional

activity, thereby yielding ICN maps[58]. Each ICN comprises brain voxels that co-vary over time,

capturing within-component connectivity. Functional connectivity between ICNs (FNC) can

subsequently be derived by quantifying temporal correlations across the ICA time series. To

facilitate cross-study reproducibility, methodological advances have led to the fully automated

spatially constrained ICA framework (NeuroMark) as a template based on 1828 young adult

participants in the GSP and HCP datasets[7]. This approach has identified 53 reliable ICNs, which

have since served as functional templates in studies of lifespan development[8], individual differences

in cognition[38,59], and a range of neuropsychiatric disorders[60,61]. These large-scale ICNs are thought

to arise from the coordinated activity of underlying neural circuits, yet the extent to which they are

shaped by microscale cellular and molecular substrates remains poorly understood. Elucidating the

cellular and molecular architecture underlying ICNs and their connectivity patterns (FNCs) could

offer a foundational neurobiological annotation of functional network organization. Such insights

are crucial for bridging levels of analysis from genes and cells to circuits and cognition.

The macroscale functional network organization of the brain is increasingly understood to

reflect underlying microscale cellular and molecular architecture[18,26,32]. To explore this hypothesis,

we systematically integrated multiple cellular and molecular datasets and quantified their spatial

associations with ICNs. We observe spatial coupling between the distributions of select cell types

and ICNs, consistent with shared structural and computational motifs. For example, L5ET neurons

constitute the cortex's principal corticofugal output, routing columnar computations to thalamus,

brainstem, and spinal targets[62,63]; their alignment with the precentral gyrus network (IC14) supports

involvement in action control and motor execution circuits[64]. By contrast, L6ITCar3 neurons

possess extensive intracortical axon projections that enable cross-areal integration[63,65], providing a

cellular substrate for the insula network's (IC27) broad role as a switching hub across externally oriented and internally oriented systems[66]. SST interneurons in layers 2/3 and 5 elaborate axons into layer 1 and thereby preferentially suppress inputs arriving through layer 1, i.e., gate top-down signals at apical dendrites[67,68] consistent with anterior cingulate cortex (IC45) computations, which integrate cost–benefit and performance signals to guide the allocation of control[69]. Interestingly, the anatomical cores of the ICNs showing cellular associations lie at key waypoints along the cortex's principal unimodal-to-transmodal gradient[55]—anchored by IC14 (precentral gyrus) at the unimodal sensorimotor pole and extending toward IC28 (superior medial frontal gyrus) and IC45 (anterior cingulate) near the transmodal apex—while IC27 (insula) acts as switching hub between large-scale systems for external attention and self-referential cognition[70]. The cell types that capture ICN spatial variability likewise traverse this gradient, echoing evidence that functional gradients parallel systematic shifts in cellular composition[32]. Neurotransmitter receptor/transporter maps and mitochondrial phenotypes most strongly predicted ICN variation in SC domain, consistent with the high SC expression of dopaminergic and cholinergic markers (e.g., D1/D2[71,72], DAT[73], NET[74], VAChT[75]) and with the energetic specializations of subcortical circuits[12]. In contrast, within the cortex, neurotransmitter maps were more closely associated with ICNs situated along the non-waypoints of the principal gradient, seemingly reflecting scale-level specificity in the neurobiological determinants of ICN organization. Although speculative, cell-type maps, which index laminar architecture and compositional differences[23,52,76,77] preferentially provided structural substrates for ICNs anchored at the waypoints of the gradient. By comparison, neurotransmitter maps, which capture modulatory sensitivity and synaptic communication[12,25,28,78,79], appeared to preferentially support the dynamic switching of networks positioned at non-waypoints along the gradient. Moreover, multi-scale models that combined neurobiological scales explained more ICN variance than any single scale, suggesting that functional networks may be shaped by the synergistic interactions among distinct cellular and molecular processes. This pattern is consistent with independent reports that neuromodulatory systems disproportionately shape patterns of brain structural development[37]. Notably, unlike prior parcellation-based spatial correlation approaches, the ICN-based framework offers a functionally segregated and spatially specific lens for cellular and molecular annotation. ICNs capture voxel-wise coactivation patterns associated with

independent functional processes, enabling a finer-grained and functionally specific mapping

between brain function and neurobiology. This approach advances beyond prior transcriptomic–

functional studies by allowing the annotation of individual functional components, rather than broad

cortical territories, with targeted cellular and molecular features.

Extending these findings, we observed that ICNs assigned to the same functional domain

exhibit significantly greater similarity in their cellular and molecular profiles—across cellular,

neurotransmitter, and mitochondrial modalities—compared to ICNs from different domains. These

similarities were paralleled by stronger functional co-activation, suggesting that shared microscale

composition may facilitate coordinated activity between regions. This supports and expands upon

prior work linking transcriptional profiles[29-31,80] and neurotransmitter systems[25] to macroscale brain

networks. Clustering similarity networks derived from cellular and molecular fingerprints revealed

a non-random alignment with canonical functional domain assignments. Specifically, ICNs were

grouped in a manner that significantly recapitulated known functional subdivisions, suggesting that

the brain's cellular and molecular landscape may, in part, mirror the organization of functional

networks revealed by group-average fMRI data. These results reinforce the notion that domain-

specific functional architectures are supported by cell-type and neurotransmitter-specific features.

Notably, this molecular-functional alignment echoes anatomical observations from histological

analyses, where homogeneous cellular composition is observed within local territories, and sharp

transitions often demarcate adjacent cortical areas[81]. Interestingly, ICNs within the VIS and SM

domains showed higher molecular and functional coupling than those within CC and DM domains.

These domain-specific molecular architectures may also help explain observed correlation patterns

of white matter networks[82] and structure–function network coupling[18,83], highlighting their role in

shaping both network identity and inter-network communication. From a molecular–structural

perspective, this finding supports the view that the association cortex may escape the strong

structural constraints imposed by early sensory–motor cascades, thereby enabling higher-order

cognitive functions beyond simple sensorimotor exchange[84].

Microscale evidence implicates cellular and molecular systems sculpting cognitive

capacities[40,85-87]. Yet, the extent to which macroscale ICNs and FNCs mediate the relationship

between cellular and molecular architecture and cognition remains poorly understood. Meta-

analytic cognitive atlases (e.g., Neurosynth) combine text mining, meta-analysis, and machine

learning to generate probabilistic mappings between cognitive terms and neural activity patterns[48].

Building on this resource, our significant spatial association–based mediation results indicate that a

set of mesoscale ICNs acts as putative interfaces through which cortical cell-type gradients relate to

cognitive representations. This pattern accords with established functional accounts: the precentral

gyrus network (IC14) aligns with motor circuitry and corticofugal output[88]; the insula network (IC27)

supports switching between externally oriented and internally oriented/self-referential systems[70];

and the anterior cingulate cortex network (IC45) integrates cognitive control with

affective/motivational processes[69,89]. SC-related ICNs anchored mediation pathways linking

neurotransmitter receptor/transporter markers and mitochondrial phenotypes to broad cognitive

functions, underscoring the central role of SC circuits in neuromodulatory regulation and energetic

support for cognition[12,90-92]. These exploratory findings extend prior work by anchoring these

functional profiles in plausible cellular architecture. From a network connectivity perspective, we

observed significant associations among cellular/molecular similarity networks, static and dynamic

FNCs, and the cognitive similarity network. These findings suggest that FNC—particularly its

dynamic aspects—may act as a critical bridge linking microscale biological architecture to

macroscale cognitive function. The observed mediation effects suggest that the alignment between

brain biology and behavior is not merely correlative but may be systematically organized through

the brain's dynamic communication architecture. In particular, the mediating role of dynamic FNC

highlights how transient, state-dependent connectivity patterns may encode biologically meaningful

variations relevant to higher-order cognition. Together, these findings move beyond simple

correlations between cell types and networks or networks and cognition. Instead, they delineate a

mediational pathway wherein macroscale functional networks serve as mechanistic conduits linking

molecular-scale organization to behavioral phenotypes. This integrative model not only strengthens

the biological plausibility of structure–function–cognition relationships but also provides a

principled framework for understanding the neurobiological substrates of individual cognitive

differences.

While our findings offer insights into the cellular and molecular foundations of ICNs, several

limitations should be acknowledged. In this study, we derived imputed spatial cellular maps by

cross-platform integration of snDrop-seq with AHBA microarray data and evaluated two

deconvolution algorithms (CIBERSORTx[93] and BayesPrism[94]), which showed overall moderate

concordance (*Supplementary Fig. S13*). Given the early stage of this field, we adopted the

CIBERSORTx-derived estimates as our primary analytic input for continuity with prior work[26,32],

and because a recent human-brain benchmarking study reported comparatively higher accuracy for

this method[95]. Nevertheless, algorithm-dependent biases highlight the need for further data

generation and head-to-head methodological evaluations tailored to this question. Second, the

cognitive density maps used here are meta-analytic, inheriting heterogeneity from the underlying

literature and potentially underrepresenting the multidimensionality of cognitive representations.

Accordingly, our associational and mediation results should be interpreted as indirect, exploratory

inferences rather than causal claims. Third, the current analyses focus on healthy young adults,

leaving generalizability across development, aging, and neuropsychiatric conditions unresolved; the

cellular and molecular maps are donor-based, which may obscure individual-level variability related

to age, sex, and other demographics; and brain-wide coverage remains uneven—particularly in

subcortical and cerebellar territories and in regions with sparse receptor mapping. Looking forward,

expanding spatially resolved molecular datasets (especially in these under-characterized areas) and

integrating emerging individual-level multimodal resources will be essential to build more complete

and personalized models of molecular–functional coupling.

In summary, this study offers a multi-scale framework linking molecular and cellular

architecture to large-scale brain function by integrating transcriptomic and neurotransmitter maps

with functional networks. We show that ICNs appear not only spatially aligned with, but also

structurally constrained by underlying biological features—constraints that extend beyond local

correspondence to reflect network-level structure. Moreover, we identify specific ICNs that mediate

the translation of molecular substrates into cognitive functions, suggesting a potential bridge from

microscale biology to macroscale behavior. These findings advance our understanding of how

functional brain networks emerge from biological structure and laying the groundwork for

integrative models that link molecular neuroscience, functional imaging, and cognition.

Methods

Participants and imaging preprocessing. This study complies with all relevant ethical regulations. All analyses were performed using publicly available, de-identified human brain datasets[7,22,23,25,47,48], each of which was obtained with ethical approval in the original studies. No new human or animal subjects were involved in this work, and therefore additional institutional ethics approval was not required. Information on participants' sex, age, consent procedures, and compensation is available in the original publications. As this study was conducted at the group and atlas levels, sex- or gender-specific analyses were not applicable.

A total of 823 healthy young adults were selected from HCP (mean age = 28.79 ± 3.68 years; 356 males) (http://www.humanconnectomeproject.org/data) dataset[47]. Comprehensive demographic information and imaging acquisition protocols are described in reference[47]. We used preprocessed rs-fMRI data (http://www.humanconnectomeproject.org/data). The HCP preprocessing pipeline included spatial distortion correction, motion correction, echo planar imaging (EPI) distortion correction, registration to standard Montreal Neurological Institute (MNI) space, and intensity normalization. Subsequently, the data were resampled to $3 \times 3 \times 3$ mm³ isotropic voxels and spatially smoothed using a Gaussian kernel with a full width at half maximum (FWHM) of 6 mm. The detailed quality control procedure for all the preprocessed data can be found in the reference[7].

ICNs. We utilized 53 ICNs from the Neuromark_fMRI_1.0[7], which are highly reproducible and biologically meaningful across independent ICA analysis of multiple large N cohorts following standard criteria for selecting ICNs. Each ICN represents a spatially independent component reflecting coherent brain activity. These ICNs were grouped into seven functional domains based on anatomical and functional priors: subcortical network (SC, 5 ICNs), auditory network (AUD, 2 ICNs), sensorimotor network (SM, 9 ICNs), visual network (VIS, 9 ICNs), cognitive-control network (CC, 17 ICNs), default-mode network (DM, 7 ICNs), and cerebellar network (CB, 4 ICNs).

FNC. The preprocessed HCP data were decomposed into 53 subject-specific independent components and their associated TCs using a spatially constrained single-subject ICA method, with the Neuromark_fMRI_1.0 template serving as spatial priors (available via GIFT at http://trendscenter.org/software/gift)[7]. To minimize residual noise in the ICN TCs, four additional postprocessing steps were applied: (1) detrending to remove linear, quadratic, and cubic trends; (2) outlier detection and removal; (3) multiple regression of six head motion parameters (three

translations and three rotations) along with their temporal derivatives; and (4) band-pass filtering

(0.01–0.15 Hz). Pearson's correlations were computed between the denoised ICN time courses to

generate individual-level FNC matrices, which were then averaged to obtain a group-level FNC

matrix. Dynamic FNC was estimated using a sliding-window approach, segmenting the TCs into

overlapping temporal windows[38]. For each segment, we computed covariance from the regularized

precision matrix[96] using the graphical LASSO method[97] with an L1 penalty to promote sparsity. The

regularization parameter (λ) was optimized for each scan by maximizing the log-likelihood on held-

out data[98]. Window-wise FNC estimates were concatenated to yield a subject-level dynamic FNC

representation, capturing temporal variations in ICN connectivity. To reduce redundancy, windows

were subsampled along the temporal axis. Exemplar windows were selected based on local maxima

in FNC variance across connections. These exemplars were clustered using k-means, repeated 500

times with random centroid initializations to ensure stability. The optimal number of states was

determined via the elbow criterion[99]. The final centroids were used to assign all windows to discrete

states. The fractional rate was quantified as the percentage of windows assigned to each state. For

each state, dynamic FNCs were averaged across all windows assigned to that state to generate group-

level state-specific FNC patterns.

AHBA microarray data processing. Human brain gene expression data from 3,702 bulk tissue

samples across six postmortem brains were obtained from the AHBA dataset (http:// human.brain-

map.org/)[22]. Raw data were processed at the sample level using the abagen toolbox

(https://abagen.readthedocs.io/en/stable)[49,50], following a standardized and widely recommended

pipeline[49]. Probe reannotation was performed based on data provided in reference[49], and probes

lacking Entrez Gene IDs were excluded. To ensure reliability, only probes that exceeded background

expression levels in at least 30% of all tissue samples were retained. Among these, the probe with

the highest differential stability was selected to represent each gene. After filtering and selection, a

total of 16,383 genes were retained. Within each donor, gene expression values were first z-scored

across genes within each sample, and then z-scored across all tissue samples.

Cell-type deconvolution. Single-nucleus droplet-based sequencing (snDrop-seq) data[23] were

obtained from the CellxGene-Census (https://cellxgene.cziscience.com/collections/d17249d2-

0e6e-4500-abb8-e6c93fa1ac6f). Cell-type deconvolution was performed following the pipeline

described previously[32]. Gene identifiers were harmonized by matching Entrez Gene IDs between

the snDrop-seq and AHBA datasets, retaining only shared genes. To reduce computational load and

enhance specificity, we limited the gene set to reliable marker genes consistently identified across

eight cortical regions, as reported previously[23]. Transcriptionally similar cell types were grouped

according to the groupings across area subclasses[23,52] to mitigate collinearity in downstream

analyses. The aggregated snDrop-seq data across all eight cortical areas were used as a reference

for cell-type abundance deconvolution on each AHBA bulk-tissue sample via CIBERSORTx

(https://cibersortx.stanford.edu/)[93], a strategy also adopted in similar studies[26,32]. This resulted in

estimated abundances for 24 distinct cell types across cortical samples, including L2/3IT, L4IT,

L6IT, L6ITCar3, L5IT, L5ET, L5/6NP, L6CT, L6b, Chandelier, PVALB, SST, SSTCHODL,

LAMP5, SNCG, PAX6, LAMP5LHX6, VIP, Astro, Endo, OPC, VLMC, Micro/PVM, and Oligo.

For each AHBA donor, estimated cell-type abundances were mapped into MNI space at $3 \times 3 \times 3$

mm³ isotropic resolution using the corresponding MNI coordinates of the tissue samples. Samples

mapped to the same voxel were averaged. To maximize sample retention, averaging across donors

was performed only for overlapping samples, yielding a final set of 1,569 samples. The resulting

group-level cell-type maps were visually consistent with parcellation-level maps reported

previously[32].

Molecular atlas. We included two sets of molecular maps representing neurotransmitter systems

and mitochondrial phenotypes. The neurotransmitter receptor and transporter density maps were

estimated   via   19   PET-derived   tracer   images   from

https://github.com/netneurolab/hansen_receptors[25]. The analyzed receptors and transporters

included 5-HT$_{1A}$, 5-HT$_{1B}$, 5-HT$_{2A}$, 5-HT$_4$, 5-HT$_6$, 5-HTT, D$_1$, D$_2$, DAT, NET, H$_3$, α$_4$β$_2$, M$_1$, VAChT,

CB$_1$, MOR, NMDA, mGluR$_5$, and GABA$_A$. Six mitochondrial phenotype images were obtained

from https://neurovault.org/collections/16418/[21]. These phenotypes included CI, CII, CIV, MitoD,

TRC, and MRC. All molecular images were coregistered to MNI space with $3 \times 3 \times 3$ mm³ isotropic

resolution, matching the resolution of the ICN and gene expression data to ensure consistent spatial

alignment across modalities.

Cognitive probabilistic measures. We utilized probabilistic maps from Neurosynth

(https://github.com/neurosynth/neurosynth)[48] to obtain voxel-wise cognitive and behavioral density

maps. These maps were generated from meta-analyses of over 15,000 published fMRI studies and

quantify the probability that a given cognitive term is reported in a study, conditional on the presence

of activation at a specific voxel. In total, 123 cognitive and behavioral terms were selected based on

the Cognitive Atlas[56]. All cognitive association maps were resampled and aligned to MNI space

with $3 \times 3 \times 3$ mm³ isotropic resolution, consistent with the spatial resolution used for functional,

molecular, and gene expression data, ensuring cross-modality comparability.

Linking ICNs with cellular and molecular atlases. To investigate the potential cellular and

molecular underpinnings of individual ICNs, we assessed the spatial associations between each ICN

map and the cellular and molecular atlases. For each ICN, we applied a threshold of $|z| > 3$ (three

standard deviations; a conventional choice to emphasize spatial specificity in ICA maps[100]) to

exclude weakly activated voxels and to focus on the core or hub voxels that contribute most strongly

to the spatially specific network pattern. Only voxels above the threshold were included in the

voxelwise spatial correlation analyses. Because the cellular atlas is cortex-limited and discrete

sampled, we implemented a local neighborhood averaging approach: for each voxel in the cellular

atlas that overlapped with the thresholded ICN map, we extracted ICN values ($|z| > 3$) within a 3-

mm radius and computed their mean, yielding a matched spatial representation that preserved

localization while reducing noise from sampling sparsity. Accordingly, ICN-to-cell-type

associations were computed only within cortical regions. Spatial similarity between ICN maps and

cell-type distributions was quantified using Spearman's correlation. To account for spatial

autocorrelation, we performed a two-sided Moran test[53] with 1,000 permutations. For each ICN

spatial map, the spatial weights matrix was defined by Euclidean distance proximity. Moran spectral

randomization was then used to generate 1,000 spatially autocorrelated surrogate maps that matched

the observed degree of spatial dependence (Moran's I). Significance was determined by comparing

observed statistics to null distributions, with FDR correction applied across 24×44 comparisons. For

the neurotransmitter and mitochondrial atlases, we employed multivariate linear regression

models[25] to estimate the contribution of each molecular map to ICN spatial variance. ICN intensity

maps were treated as the outcomes, and the mitochondrial or neurotransmitter profiles were entered

as predictors. The adjusted r^2 was used to quantify the model fit. To evaluate the relative importance

of each feature, we used the relaimpo R package, which partitions the explained variance across

predictors[54]. The significance of adjusted r^2 and relative importance was determined by one-sided

Moran test[53] with 1,000 permutations. The *p*-values of relative importance were corrected for (19×

53 for neurotransmitter; 6×53 for mitochondrion) comparisons using the FDR method. Robustness

was assessed at alternative thresholds of 2.58 (two-sided $p = 0.01$) and 3.29 (two-sided $p = 0.001$),

yielding consistent correlation results of spatial similarity across cell-type (*Supplementary Fig. S14*),

neurotransmitter (*Supplementary Fig. S15*), and mitochondrion (*Supplementary Fig. S16*).

To investigate the synergy and relative contributions of distinct neurobiological scales to ICNs,

we fitted a multi-scale model (ICN ~ all cell-type, neurotransmitter, and mitochondrion predictors)

and single-scale models (one system at a time). To mitigate collinearity among cell-type

proportional data while preserving interpretability, we applied factor analysis (minimum residuals,

promax oblique rotation)[37]. We retained all unrotated factors that explained ≥1% of variance, and

we named factors by assigning each original atlas to the factor on which it loaded most strongly.

Oblique rotation was chosen because intercorrelations are expected among non-independent

biological processes and cell populations. To align modalities spatially, neurotransmitter and

mitochondrial maps were resampled to the discrete cortical sampling of the cell-type data using the

local neighborhood averaging approach; the same sampled inputs were used when fitting the single-

system models to ensure fair comparisons. Model improvement was assessed with *F*-tests

comparing the multi-scale model to each single-scale model, with FDR applied across 3×44 + 2×9

comparisons. Scale-level relative contributions were defined as the sum of contributions from all

sub-markers within each scale.

Associations between FNC and cellular/molecular similarity networks. We averaged cellular

and molecular features across the 53 ICNs ($|z| > 3$) and constructed feature similarity networks for

cell types, neurotransmitter receptor densities, mitochondrial phenotypes, and combined molecular

fingerprints using Spearman's correlation between ICNs. Robustness was assessed at alternative

thresholds of 2.58 (two-sided $p = 0.01$) and $z = 3.29$ (two-sided $p = 0.001$), yielding consistent

results (*Supplementary Fig. S17*). To assess whether these similarity networks recapitulate

functional organization, we performed a two-sided rank-sum test comparing within-domain versus

between-domain connections. Importantly, to control for spatial autocorrelation, we regressed out

spatial proximity (both Euclidean distance and Dice similarity) between ICN spatial maps from all

connectivity matrices. Next, we assessed the global and regional alignment between the FNC and each cellular/molecular similarity network using two-sided partial Spearman's correlation, again controlling for spatial proximity. For regional correlation analyses, multiple comparisons were controlled using FDR across the tested ICNs, 44 for cell-type analyses and 53 for neurotransmitter, mitochondrial, and combined models.

Structural associations between ICNs and cellular/molecular architecture. To investigate whether the functional domain structure of ICNs is underpinned by cellular and molecular organization, we first constructed cellular and molecular similarity networks while controlling for spatial autocorrelation (Euclidean distance and Dice similarity) across ICNs. We then applied a diffusion embedding method[55] to denoise and reduce the similarity network into a low-dimensional manifold representation. A three-component solution was selected via the elbow criterion, which preserved global structure while mitigating the influence of noise or local fluctuations on subsequent clustering. Effectiveness was verified by comparison with principal component analysis (PCA, first three components) and with analyses conducted without dimensionality reduction. We then applied k-means clustering to group the projected features. The number of clusters was determined based on the number of known functional domains: 5 clusters for the cell-type similarity network and 7 clusters for the other similarity networks. The cluster labels derived from this unsupervised embedding were compared to the true functional domain labels of the ICNs. The ARI was used to quantify the accuracy of the clustering, and 1,000 label permutations were performed to assess the statistical significance of the results using a one-sided permutation test.

Mediation analysis. For cellular/molecular maps and ICNs that showed significant spatial correspondence in the primary analyses, we then conducted mediation analyses using the PROCESS toolbox in R. In these analyses, the cellular/molecular maps served as the predictor, the cognitive probabilistic measures were the outcome, and the ICN acted as the mediator. The significance of the mediation was estimated by a bias-corrected two-sided bootstrap approach with 1,000 random samplings. FDR was applied to correct the p-values for 123×30 comparisons for mediation analysis. To further investigate whether inter-ICN functional connectivity (FNC and dynamic FNC) mediates the relationship between cellular/molecular similarity and cognitive similarity, we first averaged the probabilistic values of each cognitive map across the 53 ICNs ($|z| > 3$) to obtain a single feature

vector for each cognitive term. We then constructed cognitive similarity networks by computing

Spearman's correlation between ICNs. Two-sided partial Spearman's correlations were performed

to assess associations between cellular/molecular similarity networks and the cognitive similarity

network, as well as between (static) FNC or dynamic FNC states and the cognitive similarity

network. To account for spatial confounds, Euclidean distance and Dice similarity were included as

covariates in all correlation and mediation analyses. Mediation analyses were conducted using the

PROCESS, with the cellular/molecular similarity network specified as the predictor, the cognitive

similarity network as the outcome, and FNC or dynamic FNC as the mediator. The 95% confidence

intervals were estimated using a bias-corrected two-sided bootstrap with 1,000 resamples. For the

dynamic FNC models, multiple comparisons were controlled using FDR across 5×4 tests.

Data Availability

The HCP dataset is publicly available at http://www.humanconnectomeproject.org/data. The Neuromark_fMRI_1.0 template is available at http://trendscenter.org/software/gift. The AHBA data can be accessed at http://human.brain-map.org/ and the sn-RNA dataset is available at https://cellxgene.cziscience.com/collections/d17249d2-0e6e-4500-abb8-e6c93fa1ac6f. Neurotransmitter PET imaging can be found at https://github.com/netneurolab/hansen_receptors and mitochondrial phenotype images can be found at https://neurovault.org/collections/16418/. Derived data generated in this study have been deposited at https://trendscenter.org/data/ and https://github.com/FelixFengCN/ICNs-annotation. Source data are provided with this paper.

Code Availability

SPM12 is available at http://www.fil.ion.ucl.ac.uk/spm/ , and the GIFT toolbox can be accessed at http://trendscenter.org/software/gift. AHBA sample preprocessing was performed using the abagen toolbox (https://abagen.readthedocs.io/en/stable). Single-nucleus RNA-seq data processing and cell-type imputation were conducted with CIBERSORTx (https://cibersortx.stanford.edu/). Relative importance analysis was carried out using the relaimpo R package (https://cran.r-project.org/web/packages/relaimpo/index.html). Moran test and diffusion embedding analyses were implemented via BrainSpace (https://brainspace.readthedocs.io/en/latest/index.html). Mediation analyses were performed using the PROCESS R package (https://search.r-project.org/CRAN/refmans/bruceR/html/PROCESS.html). Custom analysis codes used in this study are available at https://github.com/FelixFengCN/ICNs-annotation[101].

.

References

1. Smith, S.M., *et al.* Functional connectomics from resting-state fMRI. *Trends Cogn Sci* **17**, 666-682 (2013).

2. Zuo, X.N., *et al.* Human Connectomics across the Life Span. *Trends Cogn Sci* **21**, 32-45 (2017).

3. Perovnik, M., Rus, T., Schindlbeck, K.A. & Eidelberg, D. Functional brain networks in the evaluation of patients with neurodegenerative disorders. *Nat Rev Neurol* **19**, 73-90 (2023).

4. Fornito, A., Zalesky, A. & Breakspear, M. The connectomics of brain disorders. *Nat Rev Neurosci* **16**, 159-172 (2015).

5. Calhoun, V. Dynamic Interactivity Signatures of Mental Health Risk. *Biological Psychiatry* **97**, S24 (2025).

6. Calhoun, V.D., Miller, R., Pearlson, G. & Adalı, T. The chronnectome: time-varying connectivity networks as the next frontier in fMRI data discovery. *Neuron* **84**, 262-274 (2014).

7. Du, Y., *et al.* NeuroMark: An automated and adaptive ICA based pipeline to identify reproducible fMRI markers of brain disorders. *Neuroimage Clin* **28**, 102375 (2020).

8. Fu, Z., *et al.* Searching Reproducible Brain Features using NeuroMark: Templates for Different Age Populations and Imaging Modalities. *Neuroimage* **292**, 120617 (2024).

9. Iraji, A., *et al.* Identifying canonical and replicable multi-scale intrinsic connectivity networks in 100k+ resting-state fMRI datasets. *Hum Brain Mapp* **44**, 5729-5748 (2023).

10. Erhardt, E.B., Allen, E.A., Damaraju, E. & Calhoun, V.D. On network derivation, classification, and visualization: a response to Habeck and Moeller. *Brain Connect* **1**, 1-19 (2011).

11. Joel, S.E., Caffo, B.S., van Zijl, P.C. & Pekar, J.J. On the relationship between seed-based and ICA-based measures of functional connectivity. *Magn Reson Med* **66**, 644-657 (2011).

12. Shine, J.M. Neuromodulatory Influences on Integration and Segregation in the Brain. *Trends Cogn Sci* **23**, 572-583 (2019).

13. Zhang, D. & Raichle, M.E. Disease and the brain's dark energy. *Nat Rev Neurol* **6**, 15-28 (2010).

14. Cossart, R. & Garel, S. Step by step: cells with multiple functions in cortical circuit assembly. *Nat Rev Neurosci* **23**, 395-410 (2022).

15. Vanderhaeghen, P. & Polleux, F. Developmental mechanisms underlying the evolution of human cortical circuits. *Nat Rev Neurosci* **24**, 213-232 (2023).

16. Paquola, C., Amunts, K., Evans, A., Smallwood, J. & Bernhardt, B. Closing the mechanistic gap: the value of microarchitecture in understanding cognitive networks. *Trends Cogn Sci* **26**, 873-886 (2022).

17. Lee, A.T., Chang, E.F., Paredes, M.F. & Nowakowski, T.J. Large-scale neurophysiology and single-cell profiling in human neuroscience. *Nature* **630**, 587-595 (2024).

18. Bazinet, V., Hansen, J.Y. & Misic, B. Towards a biologically annotated brain connectome. *Nat Rev Neurosci* (2023).

19. Hansen, J.Y. & Misic, B. Integrating and interpreting brain maps. *Trends Neurosci* **48**, 594-607 (2025).

20. Palomero-Gallagher, N. & Zilles, K. Cortical layers: Cyto-, myelo-, receptor- and synaptic architecture in human cortical areas. *Neuroimage* **197**, 716-741 (2019).

690   21.   Mosharov, E.V., *et al.* A human brain map of mitochondrial respiratory capacity and
691   diversity. *Nature* (2025).
692   22.   Hawrylycz, M.J., *et al.* An anatomically comprehensive atlas of the adult human brain
693   transcriptome. *Nature* **489**, 391-399 (2012).
694   23.   Jorstad, N.L., *et al.* Transcriptomic cytoarchitecture reveals principles of human neocortex
695   organization. *Science* **382**, eadf6812 (2023).
696   24.   Lake, B.B., *et al.* Integrative single-cell analysis of transcriptional and epigenetic states in
697   the human adult brain. *Nature biotechnology* **36**, 70-80 (2018).
698   25.   Hansen, J.Y., *et al.* Mapping neurotransmitter systems to the structural and functional
699   organization of the human neocortex. *Nat Neurosci* **25**, 1569-1581 (2022).
700   26.   Anderson, K.M., *et al.* Transcriptional and imaging-genetic association of cortical
701   interneurons, brain function, and schizophrenia risk. *Nat Commun* **11**, 2889 (2020).
702   27.   Anderson, K.M., *et al.* Convergent molecular, cellular, and cortical neuroimaging
703   signatures of major depressive disorder. *Proc Natl Acad Sci U S A* **117**, 25138-25149
704   (2020).
705   28.   Shine, J.M., *et al.* Human cognition involves the dynamic integration of neural activity and
706   neuromodulatory systems. *Nat Neurosci* **22**, 289-296 (2019).
707   29.   Anderson, K.M., *et al.* Gene expression links functional networks across cortex and striatum.
708   *Nat Commun* **9**, 1428 (2018).
709   30.   Richiardi, J., *et al.* BRAIN NETWORKS. Correlated gene expression supports synchronous
710   activity in brain networks. *Science* **348**, 1241-1244 (2015).
711   31.   Krienen, F.M., Yeo, B.T., Ge, T., Buckner, R.L. & Sherwood, C.C. Transcriptional profiles of
712   supragranular-enriched genes associate with corticocortical network architecture in the
713   human brain. *Proc Natl Acad Sci U S A* **113**, E469-478 (2016).
714   32.   Zhang, X.H., *et al.* The cell-type underpinnings of the human functional cortical
715   connectome. *Nat Neurosci* **28**, 150-160 (2025).
716   33.   Hansen, J.Y., *et al.* Local molecular and global connectomic contributions to cross-
717   disorder cortical abnormalities. *Nat Commun* **13**, 4682 (2022).
718   34.   Picciotto, M.R., Higley, M.J. & Mineur, Y.S. Acetylcholine as a neuromodulator: cholinergic
719   signaling shapes nervous system function and behavior. *Neuron* **76**, 116-129 (2012).
720   35.   Kawai, H., Lazar, R. & Metherate, R. Nicotinic control of axon excitability regulates
721   thalamocortical transmission. *Nat Neurosci* **10**, 1168-1175 (2007).
722   36.   Marder, E. Neuromodulation of neuronal circuits: back to the future. *Neuron* **76**, 1-11
723   (2012).
724   37.   Lotter, L.D., *et al.* Regional patterns of human cortex development correlate with
725   underlying neurobiology. *Nat Commun* **15**, 7987 (2024).
726   38.   Fu, Z., Sui, J., Iraji, A., Liu, J. & Calhoun, V.D. Cognitive and psychiatric relevance of dynamic
727   functional connectivity states in a large (N > 10,000) children population. *Molecular
728   Psychiatry* **30**, 402-413 (2024).
729   39.   Zhi, D., *et al.* Triple Interactions Between the Environment, Brain, and Behavior in Children:
730   An ABCD Study. *Biol Psychiatry* **95**, 828-838 (2024).
731   40.   Mathys, H., *et al.* Single-cell atlas reveals correlates of high cognitive function, dementia,
732   and resilience to Alzheimer's disease pathology. *Cell* **186**, 4365-4385 e4327 (2023).
733   41.   Hara, Y., *et al.* Presynaptic mitochondrial morphology in monkey prefrontal cortex

734		correlates with working memory and is improved with estrogen treatment. *Proc Natl Acad*
735		*Sci U S A* **111**, 486-491 (2014).
736	42.	Sharpley, M.S._, et al._ Heteroplasmy of mouse mtDNA is genetically unstable and results in
737		altered behavior and cognition. *Cell* **151**, 333-343 (2012).
738	43.	Hollis, F._, et al._ Mitochondrial function in the brain links anxiety with social subordination.
739		*Proc Natl Acad Sci U S A* **112**, 15486-15491 (2015).
740	44.	Gebara, E._, et al._ Mitofusin-2 in the Nucleus Accumbens Regulates Anxiety and
741		Depression-like Behaviors Through Mitochondrial and Neuronal Actions. *Biol Psychiatry*
742		**89**, 1033-1044 (2021).
743	45.	Noudoost, B. & Moore, T. The role of neuromodulators in selective attention. *Trends Cogn*
744		*Sci* **15**, 585-591 (2011).
745	46.	Holmes, A.J._, et al._ Brain Genomics Superstruct Project initial data release with structural,
746		functional, and behavioral measures. *Sci Data* **2**, 150031 (2015).
747	47.	Van Essen, D.C._, et al._ The WU-Minn Human Connectome Project: an overview.
748		*Neuroimage* **80**, 62-79 (2013).
749	48.	Yarkoni, T., Poldrack, R.A., Nichols, T.E., Van Essen, D.C. & Wager, T.D. Large-scale
750		automated synthesis of human functional neuroimaging data. *Nat Methods* **8**, 665-670
751		(2011).
752	49.	Arnatkeviciute, A., Fulcher, B.D. & Fornito, A. A practical guide to linking brain-wide gene
753		expression and neuroimaging data. *Neuroimage* **189**, 353-367 (2019).
754	50.	Markello, R.D._, et al._ Standardizing workflows in imaging transcriptomics with the abagen
755		toolbox. *Elife* **10**(2021).
756	51.	Arnatkeviciute, A., Markello, R.D., Fulcher, B.D., Misic, B. & Fornito, A. Toward Best Practices
757		for Imaging Transcriptomics of the Human Brain. *Biol Psychiatry* **93**, 391-404 (2023).
758	52.	Jorstad, N.L._, et al._ Comparative transcriptomics reveals human-specific cortical features.
759		*Science* **382**, eade9516 (2023).
760	53.	Vos de Wael, R._, et al._ BrainSpace: a toolbox for the analysis of macroscale gradients in
761		neuroimaging and connectomics datasets. *Commun Biol* **3**, 103 (2020).
762	54.	Azen, R. & Budescu, D.V. The dominance analysis approach for comparing predictors in
763		multiple regression. *Psychol Methods* **8**, 129-148 (2003).
764	55.	Margulies, D.S._, et al._ Situating the default-mode network along a principal gradient of
765		macroscale cortical organization. *Proc Natl Acad Sci U S A* **113**, 12574-12579 (2016).
766	56.	Poldrack, R.A._, et al._ The cognitive atlas: toward a knowledge foundation for cognitive
767		neuroscience. *Frontiers in neuroinformatics* **5**, 17 (2011).
768	57.	Yu, Q._, et al._ Comparing brain graphs in which nodes are regions of interest or
769		independent components: A simulation study. *J Neurosci Methods* **291**, 61-68 (2017).
770	58.	Xu, J._, et al._ Functional network overlap as revealed by fMRI using sICA and its potential
771		relationships with functional heterogeneity, balanced excitation and inhibition, and
772		sparseness of neuron activity. *PLoS One* **10**, e0117029 (2015).
773	59.	Fu, Z., Liu, J., Salman, M.S., Sui, J. & Calhoun, V.D. Functional connectivity uniqueness and
774		variability? Linkages with cognitive and psychiatric problems in children. *Nature Mental*
775		*Health* **1**, 956-970 (2023).
776	60.	Yan, W._, et al._ A Brainwide Risk Score for Psychiatric Disorder Evaluated in a Large
777		Adolescent Population Reveals Increased Divergence Among Higher-Risk Groups Relative

to Control Participants. *Biol Psychiatry* **95**, 699-708 (2024).

61.   Du, Y.*, et al.* Evidence of shared and distinct functional and structural brain signatures in
schizophrenia and autism spectrum disorder. *Commun Biol* **4**, 1073 (2021).

62.   Kim, E.J., Juavinett, A.L., Kyubwa, E.M., Jacobs, M.W. & Callaway, E.M. Three Types of
Cortical Layer 5 Neurons That Differ in Brain-wide Connectivity and Function. *Neuron* **88**,
1253-1267 (2015).

63.   A multimodal cell census and atlas of the mammalian primary motor cortex. *Nature* **598**,
86-102 (2021).

64.   Economo, M.N.*, et al.* Distinct descending motor cortex pathways and their roles in
movement. *Nature* **563**, 79-84 (2018).

65.   Peng, H.*, et al.* Morphological diversity of single neurons in molecularly defined cell types.
Nature **598**, 174-181 (2021).

66.   Sridharan, D., Levitin, D.J. & Menon, V. A critical role for the right fronto-insular cortex in
switching between central-executive and default-mode networks. *Proc Natl Acad Sci U S*
A **105**, 12569-12574 (2008).

67.   Urban-Ciecko, J. & Barth, A.L. Somatostatin-expressing neurons in cortical networks. *Nat*
Rev Neurosci **17**, 401-409 (2016).

68.   Riedemann, T. & Sutor, B. Cell-Type-Specific Effects of Somatostatin on Synaptic
Transmission in the Anterior Cingulate Cortex. *J Neurosci* **44**(2024).

69.   Shenhav, A., Botvinick, M.M. & Cohen, J.D. The expected value of control: an integrative
theory of anterior cingulate cortex function. *Neuron* **79**, 217-240 (2013).

70.   Menon, V. & Uddin, L.Q. Saliency, switching, attention and control: a network model of
insula function. *Brain Struct Funct* **214**, 655-667 (2010).

71.   Surmeier, D.J., Carrillo-Reid, L. & Bargas, J. Dopaminergic modulation of striatal neurons,
circuits, and assemblies. *Neuroscience* **198**, 3-18 (2011).

72.   Bolam, J.P., Hanley, J.J., Booth, P.A. & Bevan, M.D. Synaptic organisation of the basal
ganglia. *J Anat* **196 (Pt 4)**, 527-542 (2000).

73.   Leroy, C.*, et al.* Assessment of 11C-PE2I binding to the neuronal dopamine transporter in
humans with the high-spatial-resolution PET scanner HRRT. *J Nucl Med* **48**, 538-546
(2007).

74.   Moriguchi, S.*, et al.* PET Quantification of the Norepinephrine Transporter in Human Brain
with (S,S)-(18)F-FMeNER-D(2). *J Nucl Med* **58**, 1140-1145 (2017).

75.   Albin, R.L.*, et al.* Regional vesicular acetylcholine transporter distribution in human brain:
A [(18) F]fluoroethoxybenzovesamicol positron emission tomography study. *The Journal*
of comparative neurology **526**, 2884-2897 (2018).

76.   Tasic, B.*, et al.* Shared and distinct transcriptomic cell types across neocortical areas.
Nature **563**, 72-78 (2018).

77.   Hodge, R.D.*, et al.* Conserved cell types with divergent features in human versus mouse
cortex. *Nature* **573**, 61-68 (2019).

78.   Goulas, A.*, et al.* The natural axis of transmitter receptor distribution in the human cerebral
cortex. *Proc Natl Acad Sci U S A* **118**(2021).

79.   Brezina, V. Beyond the wiring diagram: signalling through complex neuromodulator
networks. *Philos Trans R Soc Lond B Biol Sci* **365**, 2363-2374 (2010).

80.   Hawrylycz, M.*, et al.* Canonical genetic signatures of the adult human brain. *Nat Neurosci*

822   **18**, 1832-1844 (2015).

823   81.   Brodmann, K. Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien
824   dargestellt auf Grund des Zellenbaues. (Barth, 1909).

825   82.   Feng, G*., et al.* Longitudinal development of the human white matter structural
826   connectome and its association with brain transcriptomic and cellular architecture.
827   *Communications Biology* **6**(2023).

828   83.   Feng, G*., et al.* Spatial and temporal pattern of structure–function coupling of human brain
829   connectome with development. *eLife* **13**(2024).

830   84.   Buckner, R.L. & Krienen, F.M. The evolution of distributed association networks in the
831   human brain. *Trends Cogn Sci* **17**, 648-665 (2013).

832   85.   Zou, Y*., et al.* Cell-type-specific optogenetic fMRI on basal forebrain reveals functional
833   network basis of behavioral preference. *Neuron* **112**, 1342-1357.e1346 (2024).

834   86.   Hansen, J.Y*., et al.* Mapping gene transcription and neurocognition across human
835   neocortex. *Nat Hum Behav* **5**, 1240-1250 (2021).

836   87.   Robbins, T.W. & Arnsten, A.F. The neuropsychopharmacology of fronto-executive
837   function: monoaminergic modulation. *Annu Rev Neurosci* **32**, 267-287 (2009).

838   88.   Dum, R.P. & Strick, P.L. The origin of corticospinal projections from the premotor areas in
839   the frontal lobe. *J Neurosci* **11**, 667-689 (1991).

840   89.   Bush, G., Luu, P. & Posner, M.I. Cognitive and emotional influences in anterior cingulate
841   cortex. *Trends Cogn Sci* **4**, 215-222 (2000).

842   90.   Rangaraju, V., Lauterbach, M. & Schuman, E.M. Spatially Stable Mitochondrial
843   Compartments Fuel Local Translation during Plasticity. *Cell* **176**, 73-84.e15 (2019).

844   91.   Hwang, K., Bertolero, M.A., Liu, W.B. & D'Esposito, M. The Human Thalamus Is an
845   Integrative Hub for Functional Brain Networks. *J Neurosci* **37**, 5594-5607 (2017).

846   92.   Haber, S.N. & Knutson, B. The reward circuit: linking primate anatomy and human imaging.
847   *Neuropsychopharmacology* **35**, 4-26 (2010).

848   93.   Newman, A.M*., et al.* Determining cell type abundance and expression from bulk tissues
849   with digital cytometry. *Nature biotechnology* **37**, 773-782 (2019).

850   94.   Chu, T., Wang, Z., Pe'er, D. & Danko, C.G. Cell type and gene expression deconvolution
851   with BayesPrism enables Bayesian integrative analysis across bulk and single-cell RNA
852   sequencing in oncology. *Nat Cancer* **3**, 505-517 (2022).

853   95.   Huuki-Myers, L.A*., et al.* Benchmark of cellular deconvolution methods using a multi-
854   assay reference dataset from postmortem human prefrontal cortex. *bioRxiv* (2024).

855   96.   Smith, S.M*., et al.* Network modelling methods for FMRI. *Neuroimage* **54**, 875-891 (2011).

856   97.   Friedman, J., Hastie, T. & Tibshirani, R. Sparse inverse covariance estimation with the
857   graphical lasso. *Biostatistics* **9**, 432-441 (2008).

858   98.   Allen, E.A*., et al.* Tracking whole-brain connectivity dynamics in the resting state. *Cereb*
859   *Cortex* **24**, 663-676 (2014).

860   99.   Fu, Z*., et al.* Characterizing dynamic amplitude of low-frequency fluctuation and its
861   relationship with dynamic functional connectivity: An application to schizophrenia.
862   *Neuroimage* **180**, 619-631 (2018).

863   100.   Correa, N.M., Li, Y.O., Adalı, T. & Calhoun, V.D. Canonical Correlation Analysis for Feature-
864   Based Fusion of Biomedical Imaging Modalities and Its Application to Detection of
865   Associative Networks in Schizophrenia. *IEEE J Sel Top Signal Process* **2**, 998-1007 (2008).

866   101.   Feng, G. FelixFengCN/ICNs-annotation: v1.0.0 (v1.0.0). *Zenodo* (2025).

Acknowledgement

The authors thank all the volunteers for their participation in the study and anonymous reviewers for their insightful comments and suggestions. This work was supported by the National Science Foundation (NSF; Grant #2112455 to V.D.C.) and the National Institutes of Health (NIH; Grants #R01MH123610 and #R01MH136665 to V.D.C.). This work used transcriptomic data from the Allen Institute for Brain Science. Neuroimaging data were provided by the HCP, WU-Minn Consortium (principal investigators: D.V. Essen and K. Ugurbil; 1U54MH091657), funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research, and by the McDonnell Center for Systems Neuroscience at Washington University. The content of this manuscript solely reflects the views of the authors and may not reflect the views of the Allen Institute, NIH or the HCP consortium investigators.

Author Contributions

G.F. and V.D.C. designed the research. G.F. conducted the research. J.C., J.S., and V.D.C. provided analytic support. G.F. wrote the manuscript and made figures. All authors edited the manuscript. V.D.C. was the project administrator.

Competing Interests

There are no conflicts of interest, including any financial, personal, or other relationships with people or organizations, for any of the authors related to the work described in the article.

Figures

Fig. 1 | ICNs and cellular and molecular process. a Fifty-three robust and replicable ICNs were identified and organized into seven canonical functional domains using Neuromark_fMRI_1.0[7], derived from rs-fMRI data of 1,828 young adults across the GSP and HCP cohorts. Each ICN comprises brain voxels that co-vary over time, capturing intra-network coherence. Functional connectivity between ICNs (FNC) was derived by quantifying temporal correlations between ICN TCs, reflecting inter-network coupling. **b** A diverse set of cellular and molecular maps derived from transcriptomic data (microarray and snRNA-seq) and molecular imaging data (PET), illustrate regional variations in cellular composition, mitochondrial metabolism, and neurotransmitter signaling, providing a multiscale context for interpreting ICN organization.

Fig. 2. | Cell-type associations of ICNs. Imputed spatial distributions of 24 cortical cell types were derived by integrating bulk tissue transcriptomic data from the AHBA[22] with snRNA-seq from eight cortical regions[23,52]. Voxel-wise abundance maps were correlated with the spatial patterns of 44 cortical ICNs using Spearman's correlation. Statistical significance was determined using a two-sided Moran test[53] (1,000 iterations), with FDR correction applied across 24×44 comparisons. **a** Eight significant cell type–ICN associations (positive and negative). **b** Cortical distribution maps of the 24 cell types. Abbreviations: SC, subcortical network; AUD, auditory network; SM, sensorimotor network; VIS, visual network; CC, cognitive-control network; DM, default-mode network; CB, cerebellar network. Source data are provided as a Source Data file.

Fig. 3. | Neurotransmitter associations of ICNs. A multilinear regression framework was used to evaluate the spatial correspondence between 53 ICNs and the distribution of 19 neurotransmitter receptors and transporters, as estimated by PET imaging[25]. Dominance analysis was performed to quantify the unique variance in each ICN map attributable to individual receptor/transporter densities[54]. Statistical significance was assessed using one-sided Moran test (1,000 permutations) to account for spatial autocorrelation[53], with FDR correction applied across 53 adjusted r^2 comparisons and 19×53 dominance comparisons. **a** Sixteen ICNs demonstrated significant model fits (0.43 ≤ adjusted r^2 ≤ 0.86). Box plots show the median and interquartile range (IQR; 25–75%), with whiskers indicating 1.5× IQR from the first or third quartile. **b** Dominance analysis showed that

variability across ICNs was selectively explained by distinct receptor/transporter profiles.

Abbreviations: SC, subcortical network; AUD, auditory network; SM, sensorimotor network; VIS,

visual network; CC, cognitive-control network; DM, default-mode network; CB, cerebellar network.

Source data are provided as a Source Data file.

Fig. 4. | Cellular and molecular similarity networks relate to functional connectivity. a

Similarity networks were constructed for cell-type composition, mitochondrial metabolism,

neurotransmitter receptor density, and an integrated molecular profile by computing pairwise

Spearman correlations between the spatial fingerprints of all ICN pairs. **b** Within-domain versus

between-domain similarity was compared across these networks using a two-sided rank-sum test

(N_{within} = 230 edges, $N_{between}$ = 716 edges for cell-type; N_{within} = 246 edges, $N_{between}$ = 1,132 edges

for others), with confounding effects of spatial proximity (Euclidean distance and Dice similarity)

regressed out. Box plots show the median and interquartile range (IQR; 25–75%), with whiskers

indicating 1.5× IQR from the first or third quartile. **c** Mean FNC. Global (**d**) and regional (**e**)

associations between FNC and cellular/molecular similarity networks were evaluated using two-

sided partial Spearman's correlation, controlling for spatial proximity. In panel **e**, filled bars indicate

statistically significant associations ($q < 0.05$), whereas unfilled bars denote non-significant results.

Note: r_1 and p_1 denote the correlation coefficient and p-value without controlling for spatial

proximity; r_2 and p_2 denote those after controlling for spatial proximity. Abbreviations: SC,

subcortical network; AUD, auditory network; SM, sensorimotor network; VIS, visual network; CC,

cognitive-control network; DM, default-mode network; CB, cerebellar network. Source data are

provided as a Source Data file.

Fig. 5. | Structural associations between ICNs and cellular/molecular architecture. Based on

constructed cellular and molecular similarity networks controlling for spatial proximity across ICNs,

we applied a diffusion embedding method[55] to project these similarity networks into a shared three-

dimensional space. K-means clustering was then performed on the embedded features, with the

number of clusters determined by the number of established functional domains: five clusters for

the cell-type similarity network (**a**), and seven for the neurotransmitter (**b**), mitochondrial (**c**), and

combined (**d**) networks. The labels derived from this unsupervised clustering were compared to the

true functional domain labels of the ICNs. The adjusted rand index (ARI) was used to quantify the

accuracy of the clustering, and 1,000 label permutations were performed to assess the statistical

significance of the results using a one-sided permutation test. Abbreviations: SC, subcortical

network; AUD, auditory network; SM, sensorimotor network; VIS, visual network; CC, cognitive-

control network; DM, default-mode network; CB, cerebellar network. Source data are provided as

a Source Data file.

Fig. 6 | ICNs mediate the spatial associations between cellular and molecular architecture and

cognitive function. We used 123 probabilistic cognitive activation maps derived from meta-

analyses in Neurosynth[48]. For cellular/molecular maps and ICNs that showed significant spatial

correspondence in the primary analyses, mediation analyses were conducted using the PROCESS

toolbox in R, with cellular/molecular maps as predictors, ICNs as mediators, and cognitive maps as

outcomes. Edges in the diagram represent significant mediation pathways ($q < 0.05$), with color

indicating the magnitude of the mediation effect. Abbreviations: L4IT, layer 4 intratelencephalic

neurons; L6ITCar3, layer 6 intratelencephalic Car3-expressing neurons; L5ET, layer 5

extratelencephalic neurons; SST, somatostatin interneurons. Source data are provided as a Source

Data file.

Fig. 7 | FNC mediate the connectional associations between cellular and molecular

architecture and cognitive function. Based on 123 probabilistic cognitive activation maps derived

from meta-analyses in Neurosynth[48], a cognitive similarity network was constructed by computing

pairwise Spearman correlations between the spatial fingerprints of all ICN pairs. Mediation analyses

at the network level treated cellular/molecular similarity as the predictor, FNC as the mediator, and

cognitive similarity as the outcome. To account for spatial confounds, Euclidean distance and Dice

similarity were included as covariates in mediation analyses. The 95% confidence intervals were

estimated using a bias-corrected two-sided bootstrap with 1,000 resamples. Significant mediation

effects were observed for cell-type (**a**), neurotransmitter (**b**), and combination (**c**) similarity

networks, whereas the mitochondrion similarity network (**d**) did not yield a significant mediation

path. Source data are provided as a Source Data file.