Continuous theta-burst stimulation of the prefrontal cortex in the macaque monkey: no behavioral evidence for within-target inhibition or neural evidence for cross-hemisphere disinhibition

Running head: cTBS in macaque monkey prefrontal cortex

Sebastian J Lehmann[1] and Brian D Corneil[1-3],

[1]Department of Physiology & Pharmacology, Western University, London, ON, Canada, N6A 5B7
[2]Department of Psychology, Western University, London, ON, Canada, N6A 5B7
[3]Robarts Research Institute, Western University, London, ON, Canada, N6A 5B7

Corresponding Author: Brian D Corneil, bcorneil@uwo.ca

Number of pages: 48
Number of figures: 8
Number of tables: 1

Keywords: TMS, continuous theta burst stimulation, pre-frontal cortex, macaque

Highlights

- cTBS widely used in lab and clinic to rebalance activity across cortex
- We tested this in a monkey model, delivering cTBS to the prefrontal cortex
- No behavioural evidence for inhibition of brain area targeted by cTBS
- No evidence for disinhibition of spiking activity in mirroring, contralateral cortex
- Results question key assumptions about how cTBS influences network activity

Acknowledgements: This work was supported by Operating Grants from the Canadian Institutes of Health Research (CIHR; MOP-123247, MOP-142317, PJT-186160). We thank Dr. Julio Martinez-Trujillo for surgical expertise during the insertion of the Utah arrays.

Abstract

Background

Continuous theta-burst stimulation (cTBS) can perturb neural activity and behavior by inducing effects that persist beyond the relatively short stimulation period. Although widely used in basic research and clinical settings, there lacks an understanding of the neurophysiological and behavioural effects of cTBS.

Objectives/Hypothesis

Two assumptions motivating the use of cTBS are that it will i) inhibit neural activity in the targeted area, and ii) consequently disinhibit neural activity in the mirroring region in the contralateral cortex. Here, we test these assumptions in the oculomotor system of healthy rhesus macaques.

Methods:

In two macaques, we delivered cTBS between blocks of trials where they performed a delayed pro-/anti-saccade task, delivered cTBS to the right PFC (areas 8Ar and 46, which includes the frontal eye fields; 32 *cTBS-PFC* sessions), to the air as a SHAM control (27 *cTBS-SHAM* sessions), or to the nearby primary motor cortex as a brain control (21 *cTBS-M1* sessions). Across these different types of sessions, we compared changes in oculomotor behavior (reaction times, error rates, peak saccade velocity), and changes in neural activity recorded from the left, contralateral PFC.

Results:

Despite multiple lines of evidence consistent with TMS influencing neural activity in the cTBS-PFC and cTBS-M1 sessions, we found no behavioral evidence for inhibition of the right PFC in the cTBS-PFC sessions, nor any evidence for contralateral disinhibition in the left PFC.

Conclusions:

Our results call into question some of the fundamental assumptions underlying the application of cTBS.

Introduction

Transcranial magnetic stimulation (TMS) is one of the most common forms of non-invasive brain stimulation. TMS is widely used in both basic research and clinical scenarios, offering a reliable means to manipulate cortical activity in a targeted and non-invasive manner. Repetitive forms of TMS such as theta burst stimulation (TBS) are particularly intriguing, as they induce neural effects that persist for some time after its application. For example, continuous theta-burst stimulation (cTBS) is thought to transiently decrease neural activity in the targeted area. Such effects were first reported in primary motor cortex (M1, Huang et al., 2005, 2017), but cTBS is now routinely applied to any number of brain areas (Franca *et al.*, 2006; Rahnev *et al.*, 2013; Hanlon *et al.*, 2015; Valchev *et al.*, 2015; Blumberger *et al.*, 2018; Strzalkowski *et al.*, 2019). Doing so assumes that the effects of cTBS-M1 will generalize to other brain areas, and that such effects will modulate activity within anatomically and functionally connected regions. Indeed, one common assumption motivating its use is that cTBS-induced inhibition of the targeted brain area will increase, via disinhibition, activity in the mirroring, callosal targets (Mochizuki *et al.*, 2007; Stefan *et al.*, 2008; Suppa *et al.*, 2008, 2016; Lefaucheur *et al.*, 2020). The idea that repetitive TMS can 'rebalance' activity across hemispheres via ipsilateral inhibition and contralateral disinhibition have motivated its exploration as a treatment mode following stroke (Zheng, Liao and Xia, 2015; Li *et al.*, 2016; Long *et al.*, 2018; Vink *et al.*, 2023) or intractable depression (George *et al.*, 2010; Li *et al.*, 2014; Kang *et al.*, 2016; Brunoni *et al.*, 2017; Theleritis *et al.*, 2017).

Despite widespread use there remains a gap in knowledge in the precise neurophysiological effects of cTBS both in the targeted area and in interconnected areas, and a series of recent results in humans have questioned the reliability of effects of repetitive TMS protocols (Hamada *et al.*, 2013; Hordacre *et al.*, 2017; Perellón-Alfonso *et al.*, 2018; Boucher *et al.*, 2021; Magnuson *et al.*, 2023). Recent consensus statements (Edwards *et al.*, 2024) emphasize the importance of animal models in helping bridge the gap in knowledge. We and others (Gerits *et al.*, 2011; Mueller *et al.*, 2014; Balan *et al.*, 2017; Romero *et al.*, 2019, 2022; Pouget *et al.*, 2020; Klink *et al.*, 2021; Lehmann and

Corneil, 2022; De Lima-Pardini *et al.*, 2023) have argued that the non-human primate

(NHP) macaque offers an excellent animal model, given homologies in cortical

microstructure, gyrification, and the anatomy and function of distributed brain networks to

those found in humans. The oculomotor system can offer a particularly useful network to

study in this animal model, given similarities in retinal structure and oculomotor repertoire

to humans, and the extensive work in the rhesus macaque that has detailed patterns of

neural activity associated with the performance of complex behavioral tasks (Lehmann

and Corneil, 2022). Further, the macaque oculomotor system has been studied with many

causal techniques, such as microstimulation (Robinson and Fuchs, 1969; Robinson,

1972; Bruce *et al.*, 1985), cryogenic (Johnston, Lomber and Everling, 2016; Peel *et al.*,

2017, 2020; Dash *et al.*, 2018) or pharmacological inactivation (Sommer and Tehovnik,

1997; Dias and Segraves, 1999), or permanent lesions (Schiller, Sandell and Maunsell,

1987; Kunimatsu *et al.*, 2015; Adam *et al.*, 2020), providing a grounding for how non-

invasive brain stimulation techniques like cTBS should influence brain and behavior.

The focus of this current study is on the prefrontal cortex (PFC), and specifically the areas

rostral to and including the anterior bank of the arcuate sulcus that includes the frontal

eye fields (FEF) and adjacent area 8Ar/46. This area has been extensively targeted with

TMS in humans (for a review, see Vernet et al., 2014) and in macaques (Gerits *et al.*,

2011; Valero-Cabre *et al.*, 2012; Gu and Corneil, 2014; Mueller *et al.*, 2014; Balan *et al.*,

2017). These frontal areas are reciprocally connected anatomically with their callosal

target (Schwartz and Goldman-Rakic, 1984), and neurophysiological results in NHPs

support the notion of crossed-hemisphere inhibition (Schlag, Dassonville and Schlag-

Rey, 1998; Cohen *et al.*, 2010; Ramezanpour *et al.*, 2024). The PFC of the rhesus

macaque therefore offers an excellent platform to test the two assumptions that cTBS-

PFC will i) inhibit neural activity in the targeted area, and ii) consequently disinhibit neural

activity in the mirroring, contralateral PFC (Fig. 1A).

To test these predictions, we investigated the behavioral and neurophysiological effects

of cTBS-PFC in two healthy rhesus macaques performing a memory-guided saccade task

that required them to remember the rule to generate a pro- (look toward) or anti- (look

away from) a flashed target upon disappearance of a central fixation point. Successfully
executing this task requires working memory for rule maintenance, stimulus encoding,
cognitive control, and visuomotor integration; neural correlates of all of these features
have been reported in area 8Ar (Wallis, Anderson and Miller, 2001; Everling and
DeSouza, 2005; Backen, Treue and Martinez-Trujillo, 2018; Khanna, Scott and Smith,
2020). We used a block design, comparing behavior and neural activity before and after
cTBS-PFC to that observed with intervening bouts of cTBS delivered either to the air
(termed cTBS-SHAM, to control for acoustic effects) or cTBS delivered to the nearby hand
area of motor cortex (termed cTBS-M1, serving as a brain control that does not target the
oculomotor network but controls for tactile sensations arising from TMS). The hypothesis
that cTBS inhibits neural activity in the targeted PFC predicts increased error rates, and
longer reaction times and reduced velocities for contralateral saccades; such effects are
observed with other forms of temporary inactivation of this area (Sommer and Tehovnik,
1997; Dias and Segraves, 1999; Peel *et al.*, 2014). These behavioral effects should also
be greater for more cognitively-demanding anti-saccades (Pierrot-Deseilligny *et al.*, 1991;
Pierrot-Deseilligny *et al.*, 2003; Peel *et al.*, 2017). The hypothesis that cTBS-mediated
inhibition disinhibits the contralateral PFC predicts increased PFC activity after cTBS in
the opposite hemisphere. To test this prediction, we recorded neural activity from the PFC
contralateral to the side of cTBS via a chronically-implanted array.

Methods

Subjects and surgical procedures
The experiments were performed in two male adult rhesus macaques (*Macaca mulatta*,
animals Bu and Gr, 7 and 9 years old, weighing ca. 11 and 9 kg, respectively). Animal
housing, training, surgical, and experimental procedures were approved by the Animal
Use Subcommittee of the University of Western Ontario Council on Animal Care, and
were conducted in accordance with the Canadian Council on Animal Care policy on the
use of laboratory animals, which conforms to the guidelines laid down by the National
Institutes of Health regarding the care and use of animals for experimental procedures.
The NHPs' health and weight were monitored daily. Each macaque underwent two
surgeries. In the first surgery, a head post was implanted to restrain head motion during

training and recording sessions, along with several fiduciary markers for neuro-navigation
embedded in the acrylic implant. The implant was secured to the skull using titanium and
ceramic screws, with the ceramic screws placed in the anterior part of the skull where
TMS would be delivered. The thickness of the acrylic above the areas of interest for
delivery of TMS was limited to ~5 mm to ensure a minimal distance to the cortical surface
(for more details see Gu and Corneil, 2014). Animals then underwent structural MRI
scans to confirm cortical landmarks and locate the embedded fiduciary markers, serving

164
Fig. 1. Methods A. cTBS is expected to inhibit neural activity in the targeted right PFC and disinhibit activity
in the contralateral left PFC, potentially impacting behavioral markers such as error rate, saccadic reaction
time and peak velocity, and spiking activity in the contralateral PFC. **B.** Two rhesus macaques performed
a delayed pro- and anti-saccade task with a working memory component, requiring saccades toward or
away from a target appearing on the left or right, resulting in four conditions. **C.** cTBS was delivered in
separate daily sessions to either the right PFC (red), right M1 (blue), or as a SHAM control (black, 2 cm
above the right hemisphere). Neural activity was recorded from multielectrode arrays in the left PFC-8Ar
area, rostral to the arcuate sulcus and caudal to the principal sulcus. Individual positions for monkeys Bu
(opaque) and Gr (grey) are shown in the magnified section. **D.** In a blocked design, the monkeys first
completed 200 trials (50 per condition) of the task (PRE-cTBS), followed by cTBS delivery, and then
immediately completed a second set of 200 trials (POST-cTBS). *cTBS: continuous theta burst stimulation;*
SRT: saccadic reaction time; PFC: prefrontal cortex; M1: primary motor cortex; 8Ar: rhesus macaque area
8Ar; AS: arcuate sulcus; PS: principal sulcus.

for surgical planning and the use of neuro-navigation techniques for precise positioning

of the TMS coil (described in detail below). In a second surgery, a multi-electrode Utah

array (Blackrock Microsystems, 96 channels, electrode length 1.5mm, 10-by-10 design

with 400 micrometer inter-electrode spacing) was implanted in the left (contralateral to

the side of cTBS) prefrontal cortex, caudal to the posterior end of the principal sulcus and

anterior to the principal sulcus (see Fig. 1C for array locations in both animals, for

additional technical details see Bullock *et al.,* 2017). Recordings were started after a

recovery period of 2 to 3 weeks.

Behavioural Task. In the time between the two surgeries, the macaques were trained to

comfortably sit upright in an individually adjusted primate chair, with their head restrained,

facing a board of LEDs while sitting in a dark, secluded experimental room. They were

trained to perform a delayed pro- and anti-saccade task, which required them to hold in

working memory the rule to look either toward (*pro-saccade*) or away from (*anti-saccade*)

an upcoming peripheral stimulus (Fig.1B), and then apply this rule when the peripheral

stimulus was presented. The task was controlled via custom written real-time LabView

programs (NI-PXI controller, National Instruments, 1kHz sampling rate), and the position

of the left eye was measured by a remote eye tracking system (ETL-200, 120 Hz sampling

rate, ISCAN Inc, USA). Each trial began with the presentation of a central orange LED,

which the animal had to fixate for a variable period of 500-700 ms, with a tolerance of ~3

deg radius. The color of this central LED then changed briefly for 300 ms to either red or

green, which cued the monkey to plan for an upcoming pro-saccade (red instructional

cue) or anti-saccade (green instructional cue). The monkey had to maintain central

fixation during this time. The central LED then changed back to orange for a period of

700-900 ms, and during this time the animal had to remember the rule for the current

trials while maintaining fixation on the central target. The central LED was then turned off

and at the same time a peripheral red target was turned on either 15 degrees of visual

angle to the left or right. The monkey was rewarded with a small amount of fluid if they

executed the correct pro- or anti-saccade within 500 ms of target appearance, and

remained for 200ms within the target window ~(5 deg radius). With this experimental

structure, there are four unique trial types (pro- vs anti-, stimulus presentation left or right),

which were presented pseudo-randomly interleaved. Monkeys Bu and Gr performed ~9
to 11 trials per minute, respectively. At this pace, the animals completed a block of 200
correct trials in ~15 to 25 min, depending on their daily performance. The amount of
reward was kept constant within a given experimental session.

As illustrated in Fig 1D, data on any given day were collected within an experimental
session that consisted of ~200 correctly executed trials before cTBS (~50 of each unique
trial type, termed 'PRE-cTBS'), followed by delivery of cTBS, followed immediately by
another ~200 correctly executed trials (~50 of each unique trial type, termed 'POST-
cTBS'). cTBS always consisted of 600 pulses of TMS, delivered in bursts of 3 pulses at
50 Hz, with an inter-burst frequency of 5 Hz (40 seconds in total). cTBS was delivered to
one of three targets on any given recording day, these being: the prefrontal cortex (cTBS-
PFC), the hand area of motor cortex (cTBS-M1, serving as a brain control, and a control
for tactile sensations and acoustic effects), or 2 cm above the head (cTBS-SHAM, which
controls for acoustic effects). In both monkeys, we initially alternated sessions of cTBS-
PFC and -SHAM, introducing cTBS-M1 sessions as a second control only after collecting
data from ~10 cTBS-PFC and cTBS-SHAM sessions (e.g., the majority of cTBS-M1 data
in monkeys Bu and Gr were collected ~17 or ~5 weeks after the start of data collection;
see Fig. S1 for a summary of the timing of the recording sessions). Our rationale was to
prioritize data collection from cTBS-PFC and cTBS-SHAM, in case of device failure.

Neuronavigation and Transcranial Magnetic Stimulation. To ensure consistent
positioning of the TMS coil, we used a neuronavigation system (cortEXplore, Austria)
along with structural MRI scans. For each session, the fiduciary markers embedded within
the acrylic implants were used to register the monkey's head to a 3D model obtained from
the MRI scans. This allowed us to precisely track and position the TMS coil relative to the
cortical target areas in real-time.

Biphasic pulses of TMS were applied via a MagStim Rapid Transcranial Magnetic
Stimulator with a figure-eight coil originally designed for peripheral nerve stimulation (25
mm inner coil radius; MagStim, UK) and previously used for a variety of NHP studies

(Amaya *et al.*, 2010; Gerits *et al.*, 2011; Valero-Cabre *et al.*, 2012; Gu and Corneil, 2014;
Romero *et al.*, 2019, 2022). At the beginning of each session, the coil was first positioned
above the hand area of the right primary motor cortex (M1) to confirm the accuracy of
neuronavigation, and to confirm the repeatability of the effects of single-pulse TMS across
days. Before the start of data collection, 5-10 single pulses of TMS were delivered to M1
at an output setting of 30% (monkey Bu) or 32% (monkey Gr), which reliably evoked
visible thumb twitches in the contralateral (left) hand on between 80-100% of attempts
(see supplementary video S1 for demonstration). After this, the TMS coil was either kept
in place for cTBS-M1 sessions, or adjusted for targeting PFC (~8mm rostral to M1, using
neuronavigation), where single pulses at the same intensity were confirmed to evoke
either none or less than 20% hand twitches. We estimate that the output setting used for
cTBS was ~110% relative to resting motor threshold, and the observation of qualitatively
different evoked responses from the -PFC vs -M1 locations confirms a degree of focality
that is consistent with previous reports (Gu and Corneil, 2014). For cTBS-SHAM control
sessions, the coil was positioned above the acrylic implant, placing it on a 2cm plastic
spacer which was in secure contact with the acrylic implant; in this way, TMS pulses in
cTBS-SHAM sessions also delivered a mechanical sensation. In all cases, the coil was
locked in place by a clamp anchored to the head post. During cTBS-delivery, the TMS
coil was cooled with a combination of air suction and iced water in order to prevent
overheating. To avoid potential long-term accumulation of physiological effects, cTBS-
PFC or cTBS-M1 sessions were never repeated on sequential days, and we always
followed a day with cTBS-PFC with either a cTBS-SHAM (usually) or cTBS-M1
(occasionally in Monkey Bu only) session, or a day where data were not collected
(Supplementary Fig 1). Thus, there was a minimum of 48 hours between repeated cTBS-
PFC or cTBS-M1 sessions.

Data acquisition and analysis. Neural activity was recorded using a 128-channel
Omniplex D neural data acquisition system (Plexon Inc). Neural signals were acquired
and digitized at the headstage (16 bit resolution, Plexon DigiAmp), sampled at 40 kHz per
channel, with bandpass filtering applied online (300–8000 Hz). All data were stored for
offline spike sorting using principal component analysis techniques (Offline Sorter, Plexon

Inc), the resulting spike time information was then imported into Matlab and aligned to

behavioral events. Analog data (including horizontal and vertical eye movement traces

and stimulus presentation) was recorded at 1kHz resolution. Data were analyzed offline

using custom-written scripts in MATLAB (MathWorks). Saccade onset (saccadic reaction

time, relative to target stimulus onset) and offset were detected by applying a velocity

criterion (>50 degrees per second), trials with eye blinks or unstable fixation were

removed from the database. Peak saccadic velocity was defined as the maximum velocity

between saccade on and offset.

Spiking activity. In monkey Gr, we isolated 3979 units with an average of 94.7 ± 15.2

units per session (range 63 to 118), while in monkey Bu we isolated 873 units with an

average of 25.7 ± 8.2 units per session (range 12 to 40). The spiking activity of each

recorded neuron was aligned to three task events on a trial-by-trial basis: rule cue onset,

target onset, and saccade onset. Peristimulus time histograms (PSTHs) were computed

using a causal smoothing kernel based on a gamma distribution (Baumann et al. 2009).

Neurons were included for subsequent analyses if they exhibited an average firing rate

of at least 1 Hz across all correct trials in at least one of those alignments. To account for

variability in overall firing rates across neurons, each unit's firing rate was normalized by

dividing by the maximum average firing rate observed in any of the four task conditions

(L-pro, R-pro, L-anti, R-anti). Task modulation and selectivity for each epoch were

assessed using one-way ANOVAs ($p < 0.01$), comparing spike rates within specific

analysis windows for the following condition pairs: (1) pro- vs. anti-saccades during the

rule epoch (200–300 ms after cue onset), (2) left vs. right target locations during the visual

response epoch (75–150 ms after target onset), and (3) leftward vs. rightward saccades

during the peri-saccadic epoch (−50 to +50 ms around saccade onset).

Statistical analyses. Behavioral analyses of saccade parameters were conducted in

Matlab (MathWorks Inc.), while statistical analyses were performed using linear mixed

models (LMMs) in Jamovi (v 2.3.28). LMMs were run for behavioral and neural data

combined from both monkeys to quantify main and interaction effects; the Satterthwaite

method was used to estimate degrees of freedom and p-values. For significant

interactions, post-hoc comparisons were made using Bonferroni-adjusted p-values to

correct for multiple comparisons. LMMs assessed the effects of cTBS location (cTBS-
PFC, -SHAM, -M1), trial type (pro- vs. anti-), saccade direction (left vs. right, for the visual
response and saccade epochs), and monkey, with sessions (for behavioral analyses) and
unit IDs (for neuronal spike rate analyses) included as random effects. In the case of the
more fine-grained assessment of short-term effects on saccadic reaction time, we also
assessed the factor block (pre vs post cTBS delivery).

Data visualization. Final figures were generated using standard matlab plotting functions
and the "gramm data visualization toolbox" (Morel, 2018). Parts of the Methods figure
(Fig. 1) was created using "biorender.com".

Results

Two rhesus macaques performed an eye movement task in which the instruction to
execute a pro- or anti-saccade had to be memorized until the appearance of the visual
target, which served as a "go" signal for saccade execution (Fig. 1B, see *Methods* for
details). After ~200 correct trials (i.e., ~50 per condition), 600 pulses of cTBS were
delivered to either PFC, M1 (serving as "brain-control"), or above the head (SHAM-
control), followed by a second block of the same number of trials. Sessions were limited
to one per day, recorded across a total of 23 (monkey Gr) and 13 weeks (monkey Bu),
totalling 80 sessions where cTBS was delivered either to the right PFC (17 and 15
sessions for monkey Gr or Bu, respectively), right M1 (11/10), or to the air (a SHAM
control, 16/11).

In the following, we first examine saccade error rate, reaction time, and peak velocity to
assess whether there are behavioural signatures of inhibition following cTBS to the right
PFC. For each of these behavioural metrics, we reasoned that bilateral saccade
performance could be influenced if cTBS disrupted the encoding or maintenance of the
rule to execute a pro- or anti-saccade. Alternatively, unilateral saccade performance could
be influenced if cTBS to the right FEF disrupted the processing of leftward visual stimuli
(L-pro or R-anti), and/or the generation of leftward saccades (L-pro or L-anti). Our
hypothesis predicted that such effects would be specific for cTBS-PFC but not cTBS-M1

or cTBS-SHAM, and greater on anti- vs pro-saccade trials. For these behavioural
analyses, we compare the pre-post changes observed with cTBS-PFC to those observed
for cTBS-M1 and cTBS-SHAM. Following these behavioural analyses, we present an
analysis of spiking activity recorded from the left PFC, segregating our analysis of neural
activity in segments associated with rule maintenance, target onset, and saccade
execution.

For brevity, we report only main and interaction effects involving cTBS location. We
observed the expected pattern of results for the other factors (e.g., higher error rates,
longer reaction times, slower peak velocities for anti- vs pro-saccades), so present these
results in the Supplementary Information unless they bear on the interpretation of the
effects of cTBS location.

Behavioral analyses
Error rates. We derived error rates for pro- and anti-saccades in each direction, doing so
separately for the pre- and post-cTBS intervals for each of the three areas targeted by
cTBS (Fig. 2A&B; red and green colors depict data for pro- and anti-saccades before
cTBS respectively, with darker shades for saccades to the right; grey bars show the post-
cTBS intervals). As expected from the additional complexity of remembering and
executing an anti-saccade, error rates in both monkeys before cTBS were ~5-15% higher
on anti- vs pro-saccade trials. The change in error rate across cTBS can be appreciated
by comparing these colored bars to the grey bars immediately to the right.

Our specific interest is how performance changes across cTBS, as a function of cTBS
location. For every session, we therefore derived the change in error rate (Rate$_{post}$ -
Rate$_{pre}$) across cTBS (Fig. 2C&D). Applying the linear mixed model analyses, we found
that there was no main effect for cTBS location ($F(2,74) = 0.52$, $p = 0.60$; PFC = 2.26% ±
0.65, SHAM = 1.34% ± 0.72, M1 = 1.51% ± 0.80). The only significant effects involving
cTBS location emerged in higher-order interactions. However, post-hoc analyses
indicated that these effects were primarily driven by differences in saccade type rather
than cTBS location. The significant cTBS location × direction × saccade type interaction

Fig.2. Effect of cTBS on error rates. In these and ensuing plots, the colours indicate task type (red: pro-saccades, green: anti-saccades), while lightness indicates saccade direction (light: leftwards, dark: rightwards). **A,B**. Error rates before (coloured bars) and after (grey bars) cTBS for both monkeys, averaged across sessions (error bars: SEM). **C,D**. Changes in error rate after cTBS delivery (positive values denote an increase in error rate after cTBS). Individual dots indicate the error rate change in a given session, while bar plots represent average change across sessions (error bars: SEM). *PFC: prefrontal cortex; M1: primary motor cortex; SEM: standard error of the mean.*

(p = 0.004) was driven by a contrast between PFC/right/pro and SHAM/right/anti trials (p = 0.039; all other comparisons p > 0.085). Similarly, the cTBS location × saccade type × monkey interaction (p < 0.04) was driven by differences within Monkey Gr: SHAM/pro vs. SHAM/anti (p < 0.001), PFC/pro vs. PFC/anti (p < 0.042), and PFC/pro vs. SHAM/anti (p < 0.003).

Importantly, the observed changes in error rates across cTBS-PFC resembled those seen across cTBS-SHAM and cTBS-M1. Thus, there was no evidence for changes in error rate that could solely be attributed to cTBS-PFC, nor any evidence for the predicted state-dependent disruption in the animals' ability to perform anti-saccades.

Saccadic Reaction Time. We present the data for saccadic reaction time (RT) in a similar manner, first showing the data recorded before and after cTBS broken down by cTBS

location and monkey (Fig. 3A,B), and then by representing the change in RT across cTBS

locations (Fig. 3C,D). RT data are only shown for correctly-executed trials. The expected

increase in RTs for anti-saccades was more pronounced in monkey Gr (Fig. 3A) than

monkey Bu (Fig. 3B). RTs tended to decrease after cTBS In monkey Bu (Fig 3B), but a

similar trend was not apparent in monkey Gr (Fig. 3A).

Fig.3. Effects of cTBS on saccadic reaction time (RT). Same general format as Fig. 2. **A,B**. RTs before
and after cTBS-delivery for both monkeys, averaged across sessions. **C,D**. RT changes following cTBS
delivery, relative to baseline. Black markers shifted to the right of each bar graph indicate individual
sessions with significant RT changes for that condition on that day (1-way ANOVA, uncorrected), while
red/green markers shifted to the left of each bar graph indicate sessions without significance. *PFC:*
prefrontal cortex; M1: primary motor cortex.

As with error rate, there was no main effect of cTBS location on the change in RT across

cTBS (F(2,74) = 0.52, p = 0.60; PFC = -3.43ms ± 1.27; SHAM = -1.98ms ± 1.40; M1 = -

4.02ms ± 1.56). There were also no significant two-way interactions involving cTBS

location (all p > 0.05). Furthermore, post-hoc analysis of a significant interaction effect for

cTBS location × direction × monkey (p = 0.03) revealed it was not driven by cTBS location

but by a significant difference between monkeys (PFC/right/monkeyGr vs

PFC/right/monkeyBu, difference = -11.64ms, SE = 2.79, t(108) = -4.17, p = 0.004,

Bonferroni-corrected). Thus, we observed little no evidence for our prediction that cTBS-

PFC should selectively lead to increased RTs, finding in fact that RTs in monkey Bu
consistently decreased after cTBS regardless of stimulation location.

Sessions with significant RT changes across cTBS. Recent studies suggest that the
effects of cTBS can vary across days within the same individual, likely due to state-
dependent factors (Boucher *et al.,* 2021; Ozdemir *et al.,* 2021). We therefore assessed
whether the proportion of sessions with significant within-session RT changes depended
on cTBS location. Significant RT changes are indicated by black markers lying above or
below the x-axis in Fig. 3C–D, representing sessions with increased or decreased
averaged RT post-cTBS, respectively (see Table 1 for details).

In monkey Gr, significant RT changes were rare (12%, 14%, and 7% of sessions for
cTBS-PFC, SHAM, and M1, respectively), with a trend toward more effects in anti-
saccades, but no association with cTBS location (Chi-Square test, $\chi^2(2, N = 176) = 1.38$,
$p = 0.50$). In contrast, monkey Bu showed more frequent RT changes (32%, 23%, and
25% for cTBS-PFC, SHAM, and M1), although though the slightly higher proportion
following PFC stimulation did not reach significance ($\chi^2(2, N = 144) = 1.15$, $p = 0.56$).

	monkey Gr (Fig 3C) *No of sessions sign / ns (% sign fractions)*						**monkey Bu** (Fig 3D) *sign / ns (% sign fractions)*				
	Lpro	*Rpro*	*Lanti*	*Ranti*	*total*		*Lpro*	*Rpro*	*Lanti*	*Ranti*	*total*
PFC	1 / 16 (6%)	1 / 16 (6%)	3 / 14 (18%)	3 / 14 (18%)	**8 / 60 (12%)**		6 / 9 (40%)	5 / 10 (33%)	4 / 11 (27%)	4 / 11 (27%)	**19 / 41 32%**
SHAM	2 / 14 (13%)	2 / 14 (13%)	1 / 15 (6%)	4 / 12 (25%)	**9 / 55 (14%)**		1 / 10 (9%)	2 / 9 (18%)	4 / 7 (36%)	3 / 8 (27%)	**10 / 34 23%**
M1	0 / 11 (0%)	0 / 11 (0%)	2 / 9 (18%)	1 / 10 (9%)	**3 / 41 (7%)**		2 / 8 (20%)	2 / 8 (20%)	2 / 8 (20%)	4 / 6 (40%)	**10 / 30 25%**

421
422 **Table 1.** Proportion of sessions showing significant or non-significant changes in saccadic reaction time
423 (RT) following cTBS, along with corresponding rounded percentages of significant sessions. *PFC:*
424 *prefrontal cortex; M1: primary motor cortex; sign: significant; ns: not significant*
425

Exploring cTBS-induced short-term changes in saccadic RT. The behavioural effects

of cTBS may last for only a few minutes (Huang *et al.*, 2005). Figure 4A&B (top rows)

shows the trends of single-trial RTs relative to the time of cTBS delivery ("trial zero"),

covering the 200 trials within the PRE- and POST-sessions (recall that trial types were

pseudo-randomly interleaved, thus there were 50 trials of each type before and after

cTBS; the colored lines represent data from the three different cTBS locations). For both

monkeys, RTs tended to be shorter immediately after cTBS delivery, regardless of cTBS

location. These changes were smaller for monkey Gr (ranging around -5ms across

conditions, Fig 4A, top row) than monkey Bu (ranging around -20ms, Fig 4B, top row).

Further, RTs tended to be more stable through time in monkey Gr, whereas RTs in

monkey Bu tended to increase through both the PRE-cTBS and POST-cTBS intervals.

Such increasing RTs through time are apparent for all cTBS locations, and all trial types.

To analyze these short-term effects, we examined the distribution of RTs immediately

before or after cTBS, only including 25 trials of each condition (opaque grey boxes in the

upper row of subplots in Fig. 4), which span ~10 minutes of data.

The bottom rows of Fig. 4A&B depict individual RTs from the +/- 25 trials before (light

colors) or after (dark colors) cTBS delivery, across all conditions. To analyze short-term

effect of cTBS location on saccadic RT, our linear mixed model included the factor block

(pre- and post-cTBS), in addition to the factors cTBS-location, trial direction, trial type,

and monkey.

Saccadic reaction times tended to be shorter in the post-cTBS vs pre-cTBS blocks (factor

block, $F(1,15921) = 558.2$, $p < 0.001$; PFC/pre/cTBS 204ms +/- 1.3, PFC/post 190ms +/-

1.3, SHAM/pre 204ms +/- 1.4, SHAM/post 192ms +/- 1.3, M1/pre 202ms +/- 1.4, M1/post

188ms +/- 1.4), regardless of cTBS location or saccade condition. We found a significant

main effect for cTBS location on RT in general ($F(2,8063) = 6.98$, $p <0.001$; PFC = 197ms

± 1.2; SHAM = 198ms ± 1.3; M1 = 195ms ± 1.3). However, post-hoc tests revealed

significant differences between SHAM and M1 conditions (difference 2.7ms +/-0.7,

$p<0.001$ Bonferroni corrected), while the differences between PFC and M1 (diff 1.5 ms,

$p=0.14$) as well as PFC and SHAM (diff 1.22ms, $p=0.23$) did not reach significance. Most

Fig.4. Short-term effects of cTBS on reaction time. Top row (A,B): RT trends over time relative to cTBS delivery (trial 0), shown as mean ± 95% CI across all sessions and cTBS locations (colors), for monkey Gr (A) and monkey Bu (B). Shaded grey areas demarcate the 25 trials before (light) and after (dark grey) cTBS. Bottom row (A,B): Colored dots represent individual RTs within the pre- and post-cTBS windows, with light and dark shades indicating trials before and after cTBS, respectively. Large black dots and error bars show the median and 95th percentile range. *PFC: prefrontal cortex; M1: primary motor cortex; RT: reaction time; ms: milliseconds.*

importantly, since our primary focus was on site-specific effects on RT following cTBS delivery, we observed no interaction effect between cTBS location and block (F(2,15921) =1.9, p=0.15), indicating that the change in RT following cTBS was not influenced by cTBS location. Post-hoc comparisons following the significant cTBS location × direction × monkey interaction (F(2,15921) = 8.96, p<0.001) revealed that most significant effects

were due to differences between saccade directions and monkeys. However, we
identified three significant effects related to cTBS location: PFC/left/monkeyGr vs
M1/left/monkeyGr, p<0.01; SHAM/left/monkeyGr vs M1/left/monkeyGr, p=0.001;
PFC/left/monkeyBu vs SHAM/left/monkeyBu, p=0.023. Similarly, post-hoc comparisons
of the significant cTBS location × direction × type × monkey interaction (F(2,15921) =
3.17, p=0.042) revealed that the effect was largely driven by task-related features -
specifically, strong main effects and lower-order interactions involving the factors direction
and monkey. These interaction effects suggest that cTBS location contributed to the
observed differences under specific conditions, but was not the primary driver of the
interactions. Notably, neither the significant lower- nor higher-order interaction effects
involved the factor block, which was the primary factor of interest based on our
hypotheses.

In summary, we conducted a thorough analysis of the change in RTs across cTBS
location, analyzing average changes in RT, the proportion of significant within-session
changes in RT, and the time course of RTs. We did not find any evidence that changes
in RT following cTBS delivery could be specifically attributed to cTBS-PFC, and did not
observe any evidence for the predicted increase in RT that should have resulted from a
hypothesized cTBS-mediated inhibition of the PFC.

Saccadic peak velocity. We conducted similar analyses of peak saccade velocity, and
did not find any evidence for a main or interaction effects of cTBS-location (all p>0.09),
nor any consistent trend for a decrease in saccade velocity predicted by cTBS-mediated
inhibition of the right PFC. A figure (Supplementary Fig. S2) and further statistical
analyses for task related effects on saccade velocity are included in the Supplementary
Information.

Analyses of neural spiking activity
Description of functional tuning before cTBS. Despite the absence of behavioral
effects that conform to our predictions, cTBS-PFC may influence neural activity in more
subtle ways, perhaps below the threshold required for a behavioral change or by inducing
both excitatory and inhibitory effects. To explore this, we analyzed neural activity from

Utah arrays chronically implanted in the frontal cortex contralateral to the cTBS site. Our
analysis assesses changes in neural activity induced by cTBS within a session,
comparing activity relative to the baseline preceding cTBS. Given the stability of unit
recordings over time with Utah arrays (Sponheim *et al.*, 2021), many units were likely
recorded across multiple sessions.

A large proportion of units exhibited task modulation, and we focused on neural activity
during: (1) the "rule epoch" following rule instruction, (2) the "visual response epoch"
shortly after target onset, and (3) the "peri-saccade epoch" around saccade execution.
Figure 5A-C illustrates three example neurons exhibiting tuned activity in these three time
periods before cTBS, with the shaded areas illustrating the analysis interval. The neuron
in Figure 5A exhibited higher activity for anti-saccades (green) compared to pro-saccades
(red), which persisted even after removal of the instruction cue, consistent with previous
reports of rule maintenance in PFC (Wallis, Anderson and Miller, 2001; Everling and
DeSouza, 2005). The neuron in Figure 5B responded more strongly to visual targets in
the contralateral field, with essentially identical responses on R-pro and L-anti trials. The
neuron in Figure 5C exhibited higher activity before rightward (contralateral) saccades,
with similar increases for R-pro and R-anti trials.

Figure 5D-F shows population tuning properties for all neurons recorded in the two
monkeys, based on their activity in these epochs (dark points indicate neurons where
activity was significantly modulated during the respective epoch using a 1-way ANOVA,
$p<0.01$). Of the 3979 neurons recorded in monkey Gr (left column in Fig. 5D-F), 6% of
neurons exhibited rule-tuning (4% for anti-saccades), 32% exhibited visual tuning (21%
for contralateral targets), and 36% were tuned for saccade direction (20% for
contralateral saccades). Out of the 873 neurons recorded in monkey Bu (Fig. 5D-F, right),
only 9 units (1%) were rule-tuned, 13% exhibited visual tuning (10% for contralateral
targets), and 15% were tuned for saccade direction (8% for contralateral saccades).
These preferences for contralateral tuning to visual targets and more balanced contra-
vs- ipsilateral tuning to saccade direction resembles previous findings in area 8Ar (Bullock
et al., 2017).

Fig. 5. Task specific activity: single unit examples and population tuning. A–C Three examples of task-modulated neurons recorded in area 8Ar, showing a unit preferentially active for anti-saccades (A, aligned to cue onset, horizontal bars depict time of rule instructions), a unit with a visual response to contralateral targets (B, R-pro & L-anti, aligned to target onset), and a unit exhibiting greater activity before contralateral saccades (C, R-pro & R-anti, aligned to saccade onset). Shaded grey areas denote the analysis epochs. **D–F** Population tuning across both monkeys (monkey Gr, left; monkey Bu, right), showing the distribution of tuning directions and the percentage of neurons significantly modulated during each epoch (D: rule epoch, E: visual response epoch, F: saccade epoch). Points represent normalized firing rates of individual neurons (in arbitrary units), with darker shades indicating a significant modulation (1-way-ANOVA, p<0.01), and colors denoting tuning direction. Histograms display the distribution of tuning directions for significantly modulated units, while pie charts indicate the proportion of tuned units relative to the total recorded population in each monkey.

Assessment of changes in task-related activity over time. Neural activity in the PFC
can vary during a session due to fluctuations in slow-varying factors like attention,
arousal, and motivation (Cowley *et al.*, 2020). Given our blocked design, such longer-
timescale variations may mask any changes induced by cTBS. Thus, we first examined
changes in task-related activity across the entire session, and used this to define the
interval over which we analyzed changes induced by cTBS. Figure 6 shows the average
activity of significantly task-modulated units in each epoch, aligned to the time of cTBS
delivery (trial 0). In both monkeys, we generally observed a gradual decrease in neural
activity, particularly in the visual (Fig. 6B, E) and saccade (Fig 6C, F) epochs; the
exception is the general increase in the activity of neurons encoding the anti-saccade rule
in monkey Gr (Fig. 6A, right panel). Over the timeframe of the 200 trials before and after
cTBS, these slow fluctuations could change normalized neural activity by ~10-20%,
regardless of cTBS location. To reduce the influence of these non-cTBS-specific, time-
dependent changes while preserving sufficient trial numbers, we focused subsequent
statistical analyses on the 25 trials immediately before and after cTBS (highlighted in grey
in Fig. 6; these intervals span ~10 minutes each). We also conducted the same analysis
across the full set of trials (not shown), but this did not provide any additional insights
beyond what is reported below.

Population analysis of cTBS effects on spike rates. Figure 7 shows the change in
average spike rate across cTBS for every neuron that exhibited significant tuning during
the various epochs of the task. Given that cTBS was delivered to the other hemisphere,
contralateral disinhibition predicted increased neural activity following cTBS-PFC, i.e., the
red distributions in Fig. 7 should shift upward. For the rule epoch, the linear mixed model
included the factors cTBS location and trial type (pro- vs. anti-saccades, i.e., the only
available information at that epoch), while analyses for the visual and saccade responses
also include saccade direction.

For neurons tuned during the rule epoch (Fig. 7A, D), the linear mixed model revealed no
evidence for a main effect of cTBS location, nor any interaction effects including cTBS
location on firing rate change (all $p > 0.27$).

Fig.6 Normalized spiking activity of task modulated neurons over time. Average normalized spike
rates for all significantly task-modulated units, aligned to the time of cTBS delivery (trial 0), shown separately
for monkey Gr (A–C) and monkey Bu (D–F). Each row corresponds to a different task epoch: rule (A, D),
visual (B, E), and saccade (C, F). For the rule epoch, neural activity is grouped by saccade type (pro vs.
anti); for the visual and saccade epochs, all four task conditions are displayed. Line colors indicate cTBS
location (PFC, SHAM, M1), shaded colored areas represent SEM. Grey shaded areas indicate the time
windows used for subsequent statistical analyses of cTBS-induced effects. Y-axis scales (in arbitrary units)
vary between epochs and monkeys to best illustrate condition-specific trends. *PFC: prefrontal cortex; M1:
primary motor cortex; SEM: standard error of the mean; a.u.: arbitrary units.*

During the visual epoch (Fig. 7B, E), there was no main effect of cTBS location ($p = 0.52$),

but we found a significant interaction between cTBS location and saccade type ($F(2,4119)$

$= 18.2$, $p < 0.001$). Specifically, the cTBS-location effects on rate change were due to

differences between PFC and M1 (PFC/pro vs. M1/pro, $p < 0.001$, mean difference = 0.08

$\pm$ 0.02; PFC/anti vs. M1/anti, $p = 0.005$, mean difference = -0.07 $\pm$ 0.02) as well as

between SHAM and M1 (SHAM/pro vs. M1/pro, $p = 0.001$, difference = 0.08 $\pm$ 0.02; all

Bonferroni-corrected). Subsequently, post-hoc analyses of the significant interaction

involving cTBS location, saccade type, and monkey revealed that these differences were

driven by effects in Monkey Bu (PFC/pro/Bu vs. M1/pro/Bu, $p < 0.001$, mean difference =

0.17 $\pm$ 0.04; M1/pro/Bu vs. SHAM/pro/Bu, $p = 0.001$, difference = 0.16 $\pm$ 0.04).

For the saccade epoch (Fig. 7C,F), the main effect of cTBS location approached

significance ($F(2,1659) = 2.94$, $p = 0.053$), and post-hoc comparison revealed a small but

significant difference between PFC and M1 ($t(1659) = -2.42$, $p\_Bonferroni = 0.046$; mean

difference = -0.02 $\pm$ 0.01), while PFC vs. SHAM ($p = 0.90$) and SHAM vs. M1 ($p = 0.74$)

were not significant. Furthermore, post-hoc analyses of the significant interactions

between cTBS location and monkey ($F(2,1659) = 3.89$, $p = 0.021$) and cTBS location,

direction, type, and monkey ($F(2,4977) = 3.80$, $p < 0.022$) indicated that these effects were

largely driven by differences between monkeys Bu and Gr ($t(1659) = 3.81$, $p < 0.001$, mean

difference = 0.03 $\pm$ 0.01).

In summary, cTBS appeared to induce only minimal effects in the population of task-

modulated activity, with no consistent effect of cTBS location across both monkeys. We

also found no evidence for the predicted increase in PFC activity following cTBS-PFC.

Session-by-session analysis of cTBS effects on spike rates. Finally, we analyzed the

possibility that the neural effects of cTBS may change across days, perhaps being more

effective on some days than others. We therefore repeated the above analyses after

averaging the changes in spike rate across cTBS across all units recorded on a given day

(Figure 8). We observed no main ($F(2,42) = 1.20$, $p = 0.31$) or interaction effects (all $p >$

0.49) of cTBS location on neural activity during the rule epoch. We did observe a

Fig. 7. Effects of cTBS on normalized spike rates of individual neurons. Differences in spiking (post–
pre cTBS, in arbitrary units) for all significantly task-modulated neurons in each epoch, shown for monkey
Gr (A–C) and monkey Bu (D–F). Rows correspond to task epochs: rule (A, D), visual (B, E), and saccade

(C, F). For the rule epoch, data are grouped by saccade type (pro vs. anti); for the visual and saccade
epochs, all four task conditions are shown. Data points represent rate changes of individual neurons
(positive values denote an increase in activity after cTBS); colors indicate cTBS location (PFC, SHAM, M1).
Bars show group medians; black error bars represent SEM. *PFC: prefrontal cortex; M1: primary motor*
cortex; SEM: standard error of the mean; a.u.: arbitrary units.

significant interaction between cTBS location and saccade type during the visual epoch

(Fig. 8B,E; $F_{(2,192)} = 7.95$, $p < 0.001$), which was partially driven by a significant

difference between SHAM and M1 prosaccades (difference = 0.07 ± 0.02, $p = 0.029$,

Bonferroni-corrected). Additionally, a significant interaction between cTBS location,

saccade type, and monkey ($F_{(2,192)} = 7.09$, $p = 0.001$) was due in part to differences in

prosaccades for Monkey Bu following cTBS over M1 (M1/pro/Bu vs. PFC/pro/Bu,

difference = 0.11 ± 0.03, $p = 0.045$; M1/pro/Bu vs. SHAM/pro/Bu, difference = 0.14 ± 0.03,

$p = 0.007$). Finally, during the saccade epoch, we found a weak but significant interaction

for cTBS location x direction x monkey ($F_{(2,272)} = 3.19$, $p = 0.043$) where post-hoc results

did not survive Bonferroni correction.

Taken together, analyses across both individual neurons and session-level averages

revealed no evidence for any consistent influence of cTBS to the PFC on spiking activity

in the contralateral PFC. While we observed some significant interaction effects during

the visual and saccade epochs, these effects were not consistent across monkeys, often

did not survive corrections for multiple comparisons, and varied considerably in both

absolute magnitude and directionality. We did not see any evidence for disinhibition

following cTBS-PFC when compared to the cTBS-SHAM and cTBS-M1 controls. Broadly

speaking, the absence of disinhibitory effects aligns with the lack of any behavioural

changes after cTBS.

Fig. 8. Effects of cTBS on normalized spike rates averaged within sessions. Same format as Fig 8, except that each data point shows the change in spiking activity of significantly task-modulated, averaged

651 within a given session (i.e., one data point per session containing tuned units). *PFC: prefrontal cortex; M1:*
652 *primary motor cortex; SEM: standard error of the mean; a.u.: arbitrary units.*
653

Discussion

We investigated the behavioural and neural effects of delivering cTBS to the PFC of two
healthy rhesus macaques performing an oculomotor task. Our study tested the
hypothesis that cTBS will inhibit the PFC, and disinhibit the mirroring, contralateral PFC.
We did not find any reliable behavioral or neural evidence consistent with this hypothesis,
and in fact many of our observations went in the direction opposite to these predictions.
These are effectively negative results, and we report them in the context of recent reviews
that have emphasized the importance of reporting such results for the field (Héroux,
Taylor and Gandevia, 2015; Ziemann and Siebner, 2015). Work in animal models, and
NHPs in particular, do not afford the opportunity for extensive pilots, and in this Discussion
we will consider a number of methodological choices that, in retrospect, may have been
suboptimal. Our results contribute to a growing series of studies that question the
fundamental assumptions of notion of cTBS-mediated inhibition and consequent
disinhibition of callosal targets, and emphasize the challenges associated with assessing
changes in neural activity across intermediate timescales given the known fluctuations in
PFC activity.
670

Choice of behavioural task, and rationale for using a repetitive mode of TMS
We trained our animals to perform an inter-mixed pro- and anti-saccade task that required
them to first encode and then remember this instruction in order to correctly complete the
trial following peripheral target presentation. The memory component increases the
complexity of the task, which is reflected in the elevated pro-saccade error rate
particularly in monkey Bu (Fig. 2B) compared to when the instruction cue persists for the
entire trial (Everling, Dorris and Munoz, 1998). We chose this task given the excellent
understanding of the neural correlates of task performance (Munoz and Everling, 2004),
and since anti-saccades have been used in other studies delivering TMS to the FEF in
macaques (Valero-Cabre *et al.*, 2012) and humans (Olk *et al.*, 2006; Beynel *et al.*, 2014;
Cameron, Riddle and D'Esposito, 2015). Our choice of task is validated by PFC activity

reflecting both rule maintenance (Wallis, Anderson and Miller, 2001; Everling and
DeSouza, 2005), and the visuo-motor transformation (Bullock *et al.*, 2017; Khanna, Scott
and Smith, 2020). The interpretation of our results are bolstered through comparisons to
results in similar tasks that modulate PFC activity through other means to change
behaviour and/or neural activity (Wegener, Johnston and Everling, 2008; Phillips,
Johnston and Everling, 2011; Pouget *et al.*, 2020).

Another common task used to assess how a given intervention may change neural activity
or behaviour is the stimulus-onset asynchrony (SOA) task, which staggers the
presentation time of diametrically-opposed targets (Balan *et al.*, 2017; Adam, Johnston
and Everling, 2019; Kubanek *et al.*, 2020). The influence of a given intervention can be
assessed by shifts in the timing at which the subject would look to either target with equal
likelihood. This is a simpler task than the one we employed, and behavioural shifts in the
point of equal selection can often be explained by the changes in RTs to single left or
right targets (Rincon-Gonzalez *et al.*, 2016). Given the unchanged RTs of either pro- or
anti-saccades, it seems unlikely that behavioural effects would have been observed in
the SOA task.

A rTMS study necessitates blocked "before-vs-after" analyses. The application of cTBS
clearly influenced multiple aspects of oculomotor behaviour and PFC activity, but this
influence was largely the same regardless of cTBS target. Simply delivering cTBS likely
influences the animal's r, which temporarily shortens RTs (Fig. 4) and elevates the visual
and motor responses (Fig. 7). The dynamics of such changes complicate the identification
of effects attributable to cTBS-PFC, as any such changes would have to ride on top of
the already substantial changes induced by non-specific arousal. Other groups have
reported behavioral and neural changes in NHPs across rTMS that exceed that
attributable to non-specific arousal (Romero *et al.*, 2019, 2022), but in no cases did we
observe results from cTBS-PFC that differed substantially from both cTBS-M1 and cTBS-
SHAM. rTMS protocols in humans may be more effective when the targeted area is
actively engaged in a task (for a review, see Beynel *et al.*, 2019), but the effectiveness

and directionality of this approach has not been sufficiently established to warrant the
additional training required for experimental animals.

Finally, teaching animals the pro-/anti- saccade rule required progressive stages of
training. A nuance in the current dataset is that monkey Bu had a longer training history
than monkey Gr before the start of data collection, and in both monkeys the cTBS-M1
sessions were acquired after a substantial number of cTBS-PFC and -SHAM sessions
(see Methods, Supplementary Fig 1). This difference in training history likely explains why
monkey Gr's anti-saccade error rates before cTBS were generally lower in the M1
sessions compared to the PFC and SHAM sessions (Fig 2). The importance of training
history is also apparent in the work by Gerits and colleagues (2011), wherein saccadic
RTs decreased considerably by the time data from cTBS-M1 was collected. But even if
we discount the cTBS-M1 data from monkey Gr, the change in error rates across cTBS
were remarkably similar for the PFC vs SHAM conditions, and neither exhibited the
increase predicted from the hypothesis that cTBS will inhibit neural activity within the
targeted area.

Choice of TMS parameters and controls
rTMS protocol. We decided to use cTBS rather than 1-Hz rTMS for both pragmatic and
theoretical reasons. Delivery of 600 cTBS pulses requires only ~40s, whereas an
equivalent number of pulses at 1-Hz requires 10 minutes. Given the before-vs-after nature
of our analyses, we opted for a briefer time for rTMS delivery. There is also evidence for
superior effects of cTBS than 1-Hz rTMS, although such effects may not last as long (Di
Lazzaro *et al.*, 2010). There is also emerging evidence that delivering more rTMS pulses
in a single bout of cTBS results in longer lasting effects (Wischnewski and Schutter,
2015), but doing so would have overheated the TMS coil.

Interleaving cTBS sessions Our decision to not deliver cTBS to PFC or M1 on
sequential days was motivated by a desire to avoid possible cumulative effects of cTBS
(see Valero-Cabré, Pascual-Leone and Rushmore, 2008). Separating sessions where
cTBS is delivered to the same brain area by at least 48 hours permits within-session

analysis unconfounded by the history of what was delivered the day before. However, in
Monkey Bu we acknowledge that cTBS was delivered to the PFC 24 hours after M1 on a
subset of sessions (or vice versa; see Supplementary Fig 1). While we think the degree
to which cTBS-M1 influenced PFC activity is fairly minimal (see below), it is possible that
this subset of sessions in Monkey Bu featured some degree of a cumulative effect. We
avoided cumulative effects in Monkey Gr, and the lack of influence of cTBS across both
animals suggests that possible cumulative effects are not having a major impact on our
results. This interleaved approach may have decreased the likelihood of seeing cTBS
effects, particularly given recent trends advocating for more TMS pulses within a given
day, and across sequential days (Lefaucheur *et al.*, 2014, 2020). The decision to
interleave cTBS sessions is also influenced by the limited number of NHPs, the potential
confounds with training history (see above), and the possibility of device failure or
decreasing yields over time inherent with chronically-implanted recording arrays. It is also
possible that cTBS effects would have been realized had we delivered multiple bouts of
cTBS on a given day (e.g., 1200 pulses total; (Gerits *et al.*, 2011; Balan *et al.*, 2017)),
although we note that work in macaques has reported an influence of cTBS on brain
activity and behaviour with only 300 individual pulses during 20 s of cTBS (Merken *et al.*,
2021; Romero *et al.*, 2022).

TMS intensity. In humans, a typical pulse intensity for cTBS to M1 is 80% of resting
motor threshold (Di Lazzaro *et al.*, 2005; Huang *et al.*, 2005; for a review see Fitzgerald,
Fountain and Daskalakis, 2006). The absence of any effects in our study contrasts with
reports of cTBS changing grasping behaviour and neural excitability in the macaque
parietal cortex (Merken *et al.,* 2021; Romero *et al.*, 2022), and on saccade behaviour and
intra- and inter-hemispheric functional connectivity in the macaque FEF (Gerits *et al.*,
2011; Balan *et al.*, 2017). These studies also delivered cTBS at an intensity of ~80% of
resting motor threshold, whereas we used higher pulse intensities above the resting motor
threshold to resemble that used in the DLPFC for treatment of intractable depression
(Lefaucheur *et al.*, 2014; Van Rooij *et al.*, 2024). Dose–response relationships in TMS
are often nonlinear (Bergmann and Hartwigsen, 2020), and the tradeoff between pulse
intensity and effect focality is particularly pertinent in smaller brains, but we view it as

unlikely that a higher pulse intensity explains why the observed behavioral or neural
results of cTBS did not conform to predictions.

We are also confident that cTBS of the PFC or M1 influenced brain activity. We did not
record neural activity in the PFC targeted by cTBS, but we confirmed on cTBS-PFC
sessions that the reliable and distinct contralateral thumb twitches evoked by single pulse
TMS to M1 were less pronounced and less frequent for single-pulse TMS to PFC,
consistent with a high degree of focality. Many aspects of our implant design and pulse
intensity resemble that used in our previous study, wherein single pulses of TMS over a
widespread area of frontal cortex, including the PFC location we targeted, induced a feed-
forward neck muscle response (Gu and Corneil, 2014). In retrospect, obtaining such
objective markers to single-pulse TMS-PFC would have provided further insights about
the effects of cTBS. For example, Romero and colleagues (2022) showed that cTBS to
the parietal cortex reduced the neural response to single-pulse TMS. Despite such a
reduction, cTBS to the parietal cortex did not influence the feedforward response to a
visual stimulus, paralleling our finding of the lack of any influence of cTBS-PFC on the
visual response in the contralateral hemisphere.

Choice of controls. We included both a SHAM and a brain (M1) control to control for
non-specific effects of cTBS delivery, including tactile and auditory sensations, and
changes in arousal. This SHAM control is not ideal, as we did not have access to a true
sham coil. Instead, our SHAM control relies on the additional physical separation induced
by moving the TMS coil a further 2 cm above the acrylic, which itself lies ~5 mm above
the intact skull. While this 2 cm distance may seem modest relative to the electric field
estimates of standard figure-8 coils used in human research (Roth *et al.*, 2007; Deng,
Lisanby and Peterchev, 2013), the 25 mm figure-8 coil used here has a higher degree of
focality but consequently smaller depth of penetration (Deng, Lisanby and Peterchev,
2013). Thus, while we cannot exclude the possibility that our SHAM condition completely
eliminates the possibility of influencing PFC activity, the degree of activation will certainly
be less than that achieved by cTBS-PFC.

Considerations of focality also pertain to the suitability of our brain control, where we
delivered cTBS to M1. While the PFC and M1 cTBS locations were separated by ~8mm,
distinct neck muscle responses can be evoked from these locations (Gu and Corneil,
2014). However, given that our stimulation intensity is above resting motor threshold, is it
possible that both cTBS-PFC and cTBS-M1 influenced the oculomotor network? While
we cannot exclude this possibility, our results provide no consistent evidence for this
across both NHP, as we observed no evidence for any 'dosing' like effects, wherein the
behavioral or neural effects of cTBS-M1 could have been greater than cTBS-SHAM, given
closer proximity of the former to the PFC.

Finally, an alternative approach may have been to use an intensity control (e.g., delivering
cTBS-PFC at ~80% rather than ~110% of motor threshold) to investigate potential dose-
dependent effects. While a reasonable approach, the controls we did use may actually
serve as more extreme versions of this type of control, if indeed such controls induced
unintended influences on the PFC. Regardless, all of these possibilities must be
interpreted in light of the absence of clear differences between cTBS applied to the PFC,
M1, and SHAM conditions, and the lack of evidence for any behavioral or neural effects
that conformed to the predictions arising from our hypotheses.

Assessment window for behavioural and neural effects. Is it possible that our
analyses examined intervals that were either too soon or too late after cTBS? The original
report showed that the decrease in cortical excitability following cTBS peaks ~5-10
minutes later (Huang *et al.*, 2005), but decreases certainly begin within the first few
minutes. More recent reviews have focused on the duration of such effects on neural
excitability, which exceed 20 minutes or more depending on the number of cTBS pulses
delivered (Thut and Pascual-Leone, 2010; Wischnewski and Schutter, 2015). In the
macaque, the work from the Janssen lab (after 300 pulses of cTBS) show more variability
in the timing of peak effects after cTBS to the parietal cortex, with such effects peaking
~20-30 min after cTBS (Merken *et al.*, 2021; Romero *et al.*, 2022). However, even in this
work, reliable changes in behaviour or neural excitability begin to appear soon after cTBS,
even if the peak effect is delayed. Our presentation of saccadic RT (Fig. 4) or neural

activity (Fig. 6) through time emphasizes that the time-dependent effects were common

to all cTBS conditions. Finally, we note that mechanisms often invoked to explain the

delayed nature of the effects of cTBS, such as a delayed increase in GABA levels (Stagg

et al., 2009) in the stimulated area and an associated decrease in the contralesional

hemisphere (Matsuta *et al.*, 2022) should have altered saccade behaviour in predictable

ways, given the effects of delivery of GABA agonists or antagonists to the FEF (Dias,

Kiesau and Segraves, 1995). For example, increased GABA in the stimulated (rightward)

hemisphere should have increased the RT and decreased the peak velocity of leftward

saccades, whereas decreased GABA in the opposite (leftward) hemisphere should have

decreased the RT and increased the peak velocity of rightward saccades. We found no

evidence of this in our cTBS-PFC data.

Further technical considerations. The TMS coil used for this study has been widely

employed in previous NHP research (e.g., Amaya *et al.*, 2010; Gerits *et al.*, 2011; Valero-

Cabre *et al.*, 2012; Gu and Corneil, 2014; Romero *et al.*, 2019, 2022; Merken *et al.*, 2021).

It is big enough for efficient delivery of cTBS while small enough given the spatial

constraints imposed by cranial implants. The cranial implant was optimized to avoid field

distortion and minimize the distance from coil to skull by incorporating a thin acrylic layer

(~5 mm; for details, see (Gu and Corneil, 2014)). Precise and repeatable coil placement

was ensured via neuronavigation, which was further confirmed via the consistency of the

evoked thumb twitches from single-pulse TMS-M1 at the start of every session; coil

orientation has been shown to be a relatively negligible factor for this type of coil

(Alekseichuk *et al.*, 2019). Overall, it is unlikely that any of these technical considerations

explained the lack of cTBS-PFC effects. Finally, we delivered biphasic pulses, while some

evidence is showing that monophasic pulses induce stronger and longer-lasting effects

(Tings *et al.*, 2005; De Lima-Pardini *et al.*, 2023); this may be a parameter to modify in

future studies.

No selective influence of cTBS-PFC on spiking activity in the mirroring,
contralateral PFC

Neural activity was not recorded from the stimulated area, due to concerns that TMS
would negatively impact the stability or functioning of the chronically-implanted Utah
array. However, the effects of non-invasive brain stimulation are not limited to the targeted
area, and the possibility that cTBS-mediated inhibition can be used to induce balanced
or compensatory excitation in the mirroring, contralateral target has clear value for basic
and clinical research. One message of our manuscript is the lack of evidence for this in
terms of task-relevant spiking activity in the contralateral PFC in healthy macaques. This
absence is all the more surprising considering findings in macaques that show that very
similar cTBS protocols can alter the excitability of the targeted area to single-pulse TMS
in the parietal cortex (Romero *et al.*, 2019), and functional connectivity in the frontal
cortex, including to the mirroring, contralateral target (Balan *et al.*, 2017). In our study,
cTBS-PFC may not have induced a strong enough perturbation for changes in task-
related neural activity to stand out from the already substantial fluctuations in neural
activity in this area. Further, there may well be other perhaps more sensitive measures of
neural activity (e.g., local field potentials, and/or spike-field coherence) that do change
selectively with cTBS-PFC; such measures will be explored in a future manuscript.

PFC may also not be an ideal location to test the hypothesis of cross-hemisphere
disinhibition (Pascual-Leone, Walsh and Rothwell, 2000). There is some evidence that
the oculomotor network is fairly resilient to perturbations, perhaps because animals
execute visually-guided saccades during the inter-trial interval; thus, it may only be in the
case of more potent or widespread lesions that behavioural deficits emerge, and even
then any deficits can still be surprisingly task-dependent (Peel *et al.*, 2017, 2020; Vaidya
et al., 2019). It is also possible that effects would have been seen had cTBS been
combined with a pre-existing lesion, and our failure to observe cTBS-PFC effects in
healthy animals may not generalize to animal models of clinical conditions, such as
stroke. Finally, a recent meta-review (Kirkovski *et al.*, 2023) on cTBS protocols
emphasized that cTBS-PFC effects are more variable than cTBS-M1, perhaps due to
inherent variability in both task-related activity and in PFC activity more generally (Cowley

et al., 2020; Khanna, Scott and Smith, 2020). Such variability in PFC activity may thus be

a doubled-edge sword that influences both the ability of cTBS to influence outcome

measures, and the ability to differentiate such outcome measures from noise.

Finally, and inherent to many NHP studies, our study's sample size of only two NHPs

limits the generalizability of the results and may not capture individual variability in cTBS

response. It is possible that the two NHPs in this study were simply non-responders. While

non-responder rates in NHPs have not been documented to our knowledge, non-

responder rates in humans to TBS range between 30% to 50% (López-Alonso *et al.*,

2014; Huang *et al.*, 2017; Boucher *et al.*, 2021; Ozdemir *et al.*, 2021). If these rates

generalize to NHPs, then the chances that both our NHPs are non-responders is ~10%

to 25%. However, evidence in humans from M1 suggests that a given individual may

respond to TBS on some days but not others (Perellón-Alfonso *et al.*, 2018; Boucher *et*

al., 2021). Therefore, the limitations of our small sample size is counterbalanced by

repeated sampling of the same subject across many days. Finally, analyses motivated by

the possibility of the effectiveness of cTBS on some days but not others failed to reveal

anything unique for cTBS-PFC (Fig 4, Table 1).

Conclusions

Our findings offer no support for the widely held assumptions that cTBS, at least of the

PFC in a healthy brain, will inhibit neural activity within the stimulated area, and

consequently disinhibit neural activity in the mirroring, contralateral target. Our results

challenge the simple 'rebalancing' logic often applied for cTBS in the healthy brain, and

are in line with recent work showing limited reliability and reproducibility after cTBS of M1.

More work is needed to better understand the effects of repetitive modes of TMS,

including cTBS, on brain and behaviour in human and animal models.

Supplementary Materials

Fig. S1. Timeline of Recording Sessions and Stimulation Site Mapping for Monkeys Bu (A) and Gr (B).
Dot colors indicate cTBS target location; numbers between dots reflect the number of days between
sessions. Dates follow the format: [MonkeyIDYYMMDD]. *PFC: prefrontal cortex; M1: primary motor cortex.*

Fig. S2 Effects of cTBS on saccadic peak velocity. Same general format as Figs. 2 and 3. **A,B**. Peak velocities before and after cTBS-delivery or both monkeys, averaged across sessions. **C,D**. Change in peak velocity following cTBS delivery, relative to baseline. Black or coloured markers shifted to the right or left denote individual sessions with significant on non-significant changes in peak velocity, respectively (1-way ANOVA, uncorrected). *PFC: prefrontal cortex; M1: primary motor cortex;*

Task-related effects identified by Linear Mixed Model Analyses

This section lists the results from our Linear Mixed Model (LMM) analyses for task-related effects, excluding the factor cTBS location.

Error Rate changes (Figure 2) Our analysis of error rates revealed a significant main effect of Saccade Type ($F_{(1,222)}=8.3$, $p=0.004$) and a significant interaction between Saccade Type and Monkey ($F_{(1,222)}=8.7$, $p=0.004$).

Reaction time differences (Figure 3) For saccadic reaction times (RTs), we found significant main effects of Saccade Type ($F_{(1,222)}=42.6$, $p<0.001$) and Monkey ($F_{(1,74)}=17.9$, $p<0.001$). We also observed several significant interactions for Saccade Direction and Monkey ($F_{(1,222)}=6.2$, $p=0.013$), Saccade Type and Monkey ($F_{(1,222)}=20.5$, $p<0.001$), as well as a significant three-way interaction among Saccade Direction, Saccade Type, and Monkey: $F_{(1,222)}=6.2$, $p=0.013$.

Short-term reaction time differences (+/- 25 trial, Figure 4) The LMM on RTs within the +/- 25 trial window identified significant main effects for Block ($F_{(1,15921)}=558.2$, $p<0.001$), Saccade Type ($F_{(1,15921)}=3397.6$, $p<0.001$), Saccade Direction ($F_{(1,15921)}=195.2$, $p<0.001$), and Monkey ($F_{(1,495)}=377.7$, $p<0.001$).

Numerous significant interactions were also found:

- Monkey x Block: $F_{(1,15921)}=264.8$, $p<0.001$
- Monkey x Direction: $F_{(1,15921)}=20.0$, $p<0.001$
- Monkey x Type: $F_{(1,15921)}=1507.6$, $p<0.001$
- Monkey x Block x Direction: $F_{(1,15921)}=16.6$, $p<0.001$
- Block x Direction x Type: $F_{(1,15921)}=3.8$, $p<0.05$
- Block x Direction x Type x Monkey: $F_{(1,15921)}=24.3$, $p<0.001$

Peak Velocities (Figure S1) For peak saccade velocities, we observed significant main effects of Saccade Direction ($F_{(1,222)}=4.9$, $p=0.029$), Saccade Type ($F_{(1,222)}=4.4$, $p=0.037$), and Monkey ($F_{(1,74)}=5.6$, $p=0.021$). Significant two-way interactions were found between Monkey and Saccade Direction ($F_{(1,222)}=14.1$, $p<0.001$) and Monkey and Saccade Type ($F_{(1,222)}=10.4$, $p=0.001$), as well as a significant three-way interaction among Monkey, Saccade Direction, and Saccade Type ($F_{(1,222)}=10.8$, $p=0.001$).

Spike Rate Difference, Tuned Units (Figure 7)

- **Cue epoch:** The analysis of firing rates for tuned units revealed no significant effects during the cue epoch.

985   - **Visual epoch:** During the visual epoch, we found a significant interaction between
986   Saccade Direction and Saccade Type (F(1,4119)=6.7, p=0.01).
987   - **Saccade epoch:** For the saccade epoch, there was a significant main effect of
988   Monkey (F(1,1659)=14.5, p<0.001).
989   -

990 **Spike Rate Difference, Sessions (Figure 8)**

991   - **Cue epoch:** In the session-level analysis, we found a significant main effect of
992   Monkey during the cue epoch (F(1,42)=5.61, p=0.023).
993   - **Visual epoch:** No significant effects during the visual epoch.
994   - **Saccade epoch:** During the saccade epoch, a significant main effect of Monkey

1000
1001

Supplementary Video S1. Single-pulse TMS over the hand area of right M1 in awake monkey Bu reliably
evoked highly localized contra-lateral thumb twitches. The example shown in this video confirms the
suitability of the neuronavigation approach prior to cTBS (see methods). The single pulses in this video
(audible as distinct clicks) were delivered at 30% of stimulator output, matching the intensity used for cTBS
over the same site (M1) or PFC in subsequent recording sessions. Note the specificity of the motor
response to the thumb, irrespective of hand and arm posture. *PFC: prefrontal cortex; M1: primary motor*
cortex;

References

Adam, R. *et al.* (2020) 'Functional reorganization during the recovery of contralesional target selection deficits after prefrontal cortex lesions in macaque monkeys', *NeuroImage*, 207, p. 116339. Available at: https://doi.org/10.1016/j.neuroimage.2019.116339.

Adam, R., Johnston, K. and Everling, S. (2019) 'Recovery of contralesional saccade choice and reaction time deficits after a unilateral endothelin-1-induced lesion in the macaque caudal prefrontal cortex', *Journal of Neurophysiology*, 122(2), pp. 672–690. Available at: https://doi.org/10.1152/jn.00078.2019.

Alekseichuk, I. *et al.* (2019) 'Comparative modeling of transcranial magnetic and electric stimulation in mouse, monkey, and human', *NeuroImage*, 194, pp. 136–148. Available at: https://doi.org/10.1016/j.neuroimage.2019.03.044.

Amaya, F. *et al.* (2010) 'Transcranial magnetic stimulation and PAS-induced cortical neuroplasticity in the awake rhesus monkey', *Clinical Neurophysiology*, 121(12), pp. 2143–2151. Available at: https://doi.org/10.1016/j.clinph.2010.03.058.

Backen, T., Treue, S. and Martinez-Trujillo, J.C. (2018) 'Encoding of Spatial Attention by Primate Prefrontal Cortex Neuronal Ensembles', *eneuro*, 5(1), p. ENEURO.0372-16.2017. Available at: https://doi.org/10.1523/ENEURO.0372-16.2017.

Balan, P.F. *et al.* (2017) 'Selective TMS-induced modulation of functional connectivity correlates with changes in behavior', *NeuroImage*, 149, pp. 361–378. Available at: https://doi.org/10.1016/j.neuroimage.2017.01.076.

Bergmann, T.O. and Hartwigsen, G. (2020) 'Inferring Causality from Noninvasive Brain Stimulation in Cognitive Neuroscience', *Journal of Cognitive Neuroscience*, pp. 1–29. Available at: https://doi.org/10.1162/jocn_a_01591.

Beynel, L. *et al.* (2014) 'What saccadic eye movements tell us about TMS-induced neuromodulation of the DLPFC and mood changes: a pilot study in bipolar disorders', *Frontiers in Integrative Neuroscience*, 8. Available at: https://doi.org/10.3389/fnint.2014.00065.

Beynel, L. *et al.* (2019) 'Effects of online repetitive transcranial magnetic stimulation (rTMS) on cognitive processing: A meta-analysis and recommendations for future studies', *Neuroscience & Biobehavioral Reviews*, 107, pp. 47–58. Available at: https://doi.org/10.1016/j.neubiorev.2019.08.018.

Blumberger, D.M. *et al.* (2018) 'Effectiveness of theta burst versus high-frequency repetitive transcranial magnetic stimulation in patients with depression (THREE-D): a randomised non-inferiority trial', *The Lancet*, 391(10131), pp. 1683–1692. Available at: https://doi.org/10.1016/S0140-6736(18)30295-2.

Boucher, P.O. *et al.* (2021) 'Sham-derived effects and the minimal reliability of theta burst stimulation', *Scientific Reports*, 11(1), p. 21170. Available at: https://doi.org/10.1038/s41598-021-98751-w.

Bruce, C.J. *et al.* (1985) 'Primate frontal eye fields. II. Physiological and anatomical correlates of
electrically evoked eye movements', *Journal of Neurophysiology*, 54(3), pp. 714–734. Available
at: https://doi.org/10.1152/jn.1985.54.3.714.

Brunoni, A.R. *et al.* (2017) 'Repetitive Transcranial Magnetic Stimulation for the Acute
Treatment of Major Depressive Episodes: A Systematic Review With Network Meta-analysis',
JAMA Psychiatry, 74(2), p. 143. Available at: https://doi.org/10.1001/jamapsychiatry.2016.3644.

Bullock, K.R. *et al.* (2017) 'Visual and presaccadic activity in area 8Ar of the macaque monkey
lateral prefrontal cortex', *Journal of Neurophysiology*, 118(1), pp. 15–28. Available at:
https://doi.org/10.1152/jn.00278.2016.

Cameron, I.G.M., Riddle, J.M. and D'Esposito, M. (2015) 'Dissociable Roles of Dorsolateral
Prefrontal Cortex and Frontal Eye Fields During Saccadic Eye Movements', *Frontiers in Human
Neuroscience*, 9. Available at: https://doi.org/10.3389/fnhum.2015.00613.

Cohen, J.Y. *et al.* (2010) 'Cooperation and Competition among Frontal Eye Field Neurons
during Visual Target Selection', *The Journal of Neuroscience*, 30(9), pp. 3227–3238. Available
at: https://doi.org/10.1523/JNEUROSCI.4600-09.2010.

Cowley, B.R. *et al.* (2020) 'Slow Drift of Neural Activity as a Signature of Impulsivity in Macaque
Visual and Prefrontal Cortex', *Neuron*, 108(3), pp. 551-567.e8. Available at:
https://doi.org/10.1016/j.neuron.2020.07.021.

Dash, S. *et al.* (2018) 'Frontal Eye Field Inactivation Reduces Saccade Preparation in the
Superior Colliculus but Does Not Alter How Preparatory Activity Relates to Saccades of a Given
Latency', *eneuro*, 5(2), p. ENEURO.0024-18.2018. Available at:
https://doi.org/10.1523/ENEURO.0024-18.2018.

De Lima-Pardini, A.C. *et al.* (2023) 'Transcranial magnetic stimulation in non-human primates: A
systematic review', *Neuroscience & Biobehavioral Reviews*, 152, p. 105273. Available at:
https://doi.org/10.1016/j.neubiorev.2023.105273.

Deng, Z.-D., Lisanby, S.H. and Peterchev, A.V. (2013) 'Electric field depth–focality tradeoff in
transcranial magnetic stimulation: Simulation comparison of 50 coil designs', *Brain Stimulation*,
6(1), pp. 1–13. Available at: https://doi.org/10.1016/j.brs.2012.02.005.

Di Lazzaro, V. *et al.* (2005) 'Theta-burst repetitive transcranial magnetic stimulation suppresses
specific excitatory circuits in the human motor cortex: Theta-burst repetitive transcranial
magnetic stimulation', *The Journal of Physiology*, 565(3), pp. 945–950. Available at:
https://doi.org/10.1113/jphysiol.2005.087288.

Di Lazzaro, V. *et al.* (2010) 'The effects of motor cortex rTMS on corticospinal descending
activity', *Clinical Neurophysiology*, 121(4), pp. 464–473. Available at:
https://doi.org/10.1016/j.clinph.2009.11.007.

Dias, E.C., Kiesau, M. and Segraves, M.A. (1995) 'Acute activation and inactivation of macaque
frontal eye field with GABA-related drugs', *Journal of Neurophysiology*, 74(6), pp. 2744–2748.
Available at: https://doi.org/10.1152/jn.1995.74.6.2744.

Dias, E.C. and Segraves, M.A. (1999) 'Muscimol-Induced Inactivation of Monkey Frontal Eye

Field: Effects on Visually and Memory-Guided Saccades', *Journal of Neurophysiology*, 81(5),
pp. 2191–2214. Available at: https://doi.org/10.1152/jn.1999.81.5.2191.

Edwards, J.D. *et al.* (2024) 'A translational roadmap for transcranial magnetic and direct current
stimulation in stroke rehabilitation: Consensus-based core recommendations from the third
stroke recovery and rehabilitation roundtable', *International Journal of Stroke*, 19(2), pp. 145–
157. Available at: https://doi.org/10.1177/17474930231203982.

Everling, S. and DeSouza, J.F.X. (2005) 'Rule-dependent Activity for Prosaccades and
Antisaccades in the Primate Prefrontal Cortex', *Journal of Cognitive Neuroscience*, 17(9), pp.
1483–1496. Available at: https://doi.org/10.1162/0898929054985455.

Everling, S., Dorris, M.C. and Munoz, D.P. (1998) 'Reflex Suppression in the Anti-Saccade Task
Is Dependent on Prestimulus Neural Processes', *Journal of Neurophysiology*, 80(3), pp. 1584–
1589. Available at: https://doi.org/10.1152/jn.1998.80.3.1584.

Fitzgerald, P., Fountain, S. and Daskalakis, Z. (2006) 'A comprehensive review of the effects of
rTMS on motor cortical excitability and inhibition', *Clinical Neurophysiology*, 117(12), pp. 2584–
2596. Available at: https://doi.org/10.1016/j.clinph.2006.06.712.

Franca, M. *et al.* (2006) 'Effects of theta burst stimulation protocols on phosphene threshold',
Clinical Neurophysiology, 117(8), pp. 1808–1813. Available at:
https://doi.org/10.1016/j.clinph.2006.03.019.

George, M.S. *et al.* (2010) 'Daily Left Prefrontal Transcranial Magnetic Stimulation Therapy for
Major Depressive Disorder: A Sham-Controlled Randomized Trial', *Archives of General*
Psychiatry, 67(5), p. 507. Available at: https://doi.org/10.1001/archgenpsychiatry.2010.46.

Gerits, A. *et al.* (2011) 'Transcranial magnetic stimulation of macaque frontal eye fields
decreases saccadic reaction time', *Experimental Brain Research*, 212(1), pp. 143–152.
Available at: https://doi.org/10.1007/s00221-011-2710-3.

Gu, C. and Corneil, B.D. (2014) 'Transcranial Magnetic Stimulation of the Prefrontal Cortex in
Awake Nonhuman Primates Evokes a Polysynaptic Neck Muscle Response That Reflects
Oculomotor Activity at the Time of Stimulation', *Journal of Neuroscience*, 34(44), pp. 14803–
14815. Available at: https://doi.org/10.1523/JNEUROSCI.2907-14.2014.

Hamada, M. *et al.* (2013) 'The Role of Interneuron Networks in Driving Human Motor Cortical
Plasticity', *Cerebral Cortex*, 23(7), pp. 1593–1605. Available at:
https://doi.org/10.1093/cercor/bhs147.

Hanlon, C.A. *et al.* (2015) 'What goes up, can come down: Novel brain stimulation paradigms
may attenuate craving and craving-related neural circuitry in substance dependent individuals',
Brain Research, 1628, pp. 199–209. Available at: https://doi.org/10.1016/j.brainres.2015.02.053.

Héroux, M.E., Taylor, J.L. and Gandevia, S.C. (2015) 'The Use and Abuse of Transcranial
Magnetic Stimulation to Modulate Corticospinal Excitability in Humans', *PLOS ONE*. Edited by
F. Tremblay, 10(12), p. e0144151. Available at: https://doi.org/10.1371/journal.pone.0144151.

Hordacre, B. *et al.* (2017) 'Variability in neural excitability and plasticity induction in the human
cortex: A brain stimulation study', *Brain Stimulation*, 10(3), pp. 588–595. Available at:

https://doi.org/10.1016/j.brs.2016.12.001.

Huang, Y.-Z. *et al.* (2005) 'Theta Burst Stimulation of the Human Motor Cortex', *Neuron*, 45(2),
pp. 201–206. Available at: https://doi.org/10.1016/j.neuron.2004.12.033.

Huang, Y.-Z. *et al.* (2017) 'Plasticity induced by non-invasive transcranial brain stimulation: A
position paper', *Clinical Neurophysiology*, 128(11), pp. 2318–2329. Available at:
https://doi.org/10.1016/j.clinph.2017.09.007.

Johnston, K., Lomber, S.G. and Everling, S. (2016) 'Unilateral deactivation of macaque
dorsolateral prefrontal cortex induces biases in stimulus selection', *Journal of Neurophysiology*,
115(3), pp. 1468–1476. Available at: https://doi.org/10.1152/jn.00563.2015.

Kang, J.I. *et al.* (2016) 'Frontostriatal Connectivity Changes in Major Depressive Disorder After
Repetitive Transcranial Magnetic Stimulation: A Randomized Sham-Controlled Study', *The*
Journal of Clinical Psychiatry, 77(09), pp. e1137–e1143. Available at:
https://doi.org/10.4088/JCP.15m10110.

Khanna, S.B., Scott, J.A. and Smith, M.A. (2020) 'Dynamic shifts of visual and saccadic signals
in prefrontal cortical regions 8Ar and FEF', *Journal of Neurophysiology*, 124(6), pp. 1774–1791.
Available at: https://doi.org/10.1152/jn.00669.2019.

Kirkovski, M. *et al.* (2023) 'A systematic review of the neurobiological effects of theta-burst
stimulation (TBS) as measured using functional magnetic resonance imaging (fMRI)', *Brain*
Structure and Function, 228(3–4), pp. 717–749. Available at: https://doi.org/10.1007/s00429-
023-02634-x.

Klink, P.C. *et al.* (2021) 'Combining brain perturbation and neuroimaging in non-human
primates', *NeuroImage*, 235, p. 118017. Available at:
https://doi.org/10.1016/j.neuroimage.2021.118017.

Kubanek, J. *et al.* (2020) 'Remote, brain region–specific control of choice behavior with
ultrasonic waves', *Science Advances*, 6(21), p. eaaz4193. Available at:
https://doi.org/10.1126/sciadv.aaz4193.

Kunimatsu, J. *et al.* (2015) 'Application of radiosurgical techniques to produce a primate model
of brain lesions', *Frontiers in Systems Neuroscience*, 9. Available at:
https://doi.org/10.3389/fnsys.2015.00067.

Lefaucheur, J.-P. *et al.* (2014) 'Evidence-based guidelines on the therapeutic use of repetitive
transcranial magnetic stimulation (rTMS)', *Clinical Neurophysiology*, 125(11), pp. 2150–2206.
Available at: https://doi.org/10.1016/j.clinph.2014.05.021.

Lefaucheur, J.-P. *et al.* (2020) 'Evidence-based guidelines on the therapeutic use of repetitive
transcranial magnetic stimulation (rTMS): An update (2014–2018)', *Clinical Neurophysiology*,
131(2), pp. 474–528. Available at: https://doi.org/10.1016/j.clinph.2019.11.002.

Lehmann, S.J. and Corneil, B.D. (2022) 'Completing the puzzle: Why studies in non-human
primates are needed to better understand the effects of non-invasive brain stimulation',
Neuroscience & Biobehavioral Reviews, 132, pp. 1074–1085. Available at:
https://doi.org/10.1016/j.neubiorev.2021.10.040.

Li, C.-T. *et al.* (2014) 'Efficacy of prefrontal theta-burst stimulation in refractory depression: a
randomized sham-controlled study', *Brain*, 137(7), pp. 2088–2098. Available at:
https://doi.org/10.1093/brain/awu109.

Li, J. *et al.* (2016) 'Effects of different frequencies of repetitive transcranial magnetic stimulation
on the recovery of upper limb motor dysfunction in patients with subacute cerebral infarction',
Neural Regeneration Research, 11(10), p. 1584. Available at: https://doi.org/10.4103/1673-
5374.193236.

Long, H. *et al.* (2018) 'Effects of combining high- and low-frequency repetitive transcranial
magnetic stimulation on upper limb hemiparesis in the early phase of stroke', *Restorative
Neurology and Neuroscience*, 36(1), pp. 21–30. Available at: https://doi.org/10.3233/RNN-
170733.

López-Alonso, V. *et al.* (2014) 'Inter-individual Variability in Response to Non-invasive Brain
Stimulation Paradigms', *Brain Stimulation*, 7(3), pp. 372–380. Available at:
https://doi.org/10.1016/j.brs.2014.02.004.

Magnuson, J. *et al.* (2023) 'Neuromodulatory effects and reproducibility of the most widely used
repetitive transcranial magnetic stimulation protocols', *PLOS ONE*. Edited by F. Tremblay,
18(6), p. e0286465. Available at: https://doi.org/10.1371/journal.pone.0286465.

Matsuta, H. *et al.* (2022) 'Continuous theta-burst stimulation to the sensorimotor cortex affects
contralateral gamma-aminobutyric acid level and resting-state networks', *PLOS ONE*. Edited by
B.A. Murphy, 17(8), p. e0272268. Available at: https://doi.org/10.1371/journal.pone.0272268.

Merken, L. *et al.* (2021) 'Behavioral effects of continuous theta-burst stimulation in macaque
parietal cortex', *Scientific Reports*, 11(1), p. 4511. Available at: https://doi.org/10.1038/s41598-
021-83904-8.

Mochizuki, H. *et al.* (2007) 'Hemoglobin concentration changes in the contralateral hemisphere
during and after theta burst stimulation of the human sensorimotor cortices', *Experimental Brain
Research*, 180(4), pp. 667–675. Available at: https://doi.org/10.1007/s00221-007-0884-5.

Mueller, J.K. *et al.* (2014) 'Simultaneous transcranial magnetic stimulation and single-neuron
recording in alert non-human primates', *Nature Neuroscience*, 17(8), pp. 1130–1136. Available
at: https://doi.org/10.1038/nn.3751.

Munoz, D.P. and Everling, S. (2004) 'Look away: the anti-saccade task and the voluntary control
of eye movement', *Nature Reviews Neuroscience*, 5(3), pp. 218–228. Available at:
https://doi.org/10.1038/nrn1345.

Olk, B. *et al.* (2006) 'Modulation of Antisaccades by Transcranial Magnetic Stimulation of the
Human Frontal Eye Field', *Cerebral Cortex*, 16(1), pp. 76–82. Available at:
https://doi.org/10.1093/cercor/bhi085.

Ozdemir, R.A. *et al.* (2021) 'Reproducibility of cortical response modulation induced by
intermittent and continuous theta-burst stimulation of the human motor cortex', *Brain
Stimulation*, 14(4), pp. 949–964. Available at: https://doi.org/10.1016/j.brs.2021.05.013.

Pascual-Leone, A., Walsh, V. and Rothwell, J.C. (2000) 'Transcranial magnetic stimulation in

cognitive neuroscience – virtual lesion, chronometry, and functional connectivity', *Current*
Opinion in Neurobiology, 10(2), pp. 232–237. Available at: https://doi.org/10.1016/S0959-
4388(00)00081-7.

Peel, T.R. *et al.* (2014) 'Bilateral saccadic deficits following large and reversible inactivation of
unilateral frontal eye field', *Journal of Neurophysiology*, 111(2), pp. 415–433. Available at:
https://doi.org/10.1152/jn.00398.2013.

Peel, T.R. *et al.* (2017) 'Frontal Eye Field Inactivation Diminishes Superior Colliculus Activity,
But Delayed Saccadic Accumulation Governs Reaction Time Increases', *The Journal of*
Neuroscience, 37(48), pp. 11715–11730. Available at:
https://doi.org/10.1523/JNEUROSCI.2664-17.2017.

Peel, T.R. *et al.* (2020) 'Frontal eye field inactivation alters the readout of superior colliculus
activity for saccade generation in a task-dependent manner', *Journal of Computational*
Neuroscience [Preprint]. Available at: https://doi.org/10.1007/s10827-020-00760-7.

Perellón-Alfonso, R. *et al.* (2018) 'Similar effect of intermittent theta burst and sham stimulation
on corticospinal excitability: A 5-day repeated sessions study', *European Journal of*
Neuroscience, 48(4), pp. 1990–2000. Available at: https://doi.org/10.1111/ejn.14077.

Phillips, J.M., Johnston, K. and Everling, S. (2011) 'Effects of Anterior Cingulate
Microstimulation on Pro- and Antisaccades in Nonhuman Primates', *Journal of Cognitive*
Neuroscience, 23(2), pp. 481–490. Available at: https://doi.org/10.1162/jocn.2010.21482.

Pierrot-Deseilligny, C. *et al.* (2003) 'Decisional role of the dorsolateral prefrontal cortex in ocular
motor behaviour', *Brain*, 126(6), pp. 1460–1473. Available at:
https://doi.org/10.1093/brain/awg148.

Pierrot-Deseilligny, Ch. *et al.* (1991) 'CORTICAL CONTROL OF REFLEXIVE VISUALLY-
GUIDED SACCADES', *Brain*, 114(3), pp. 1473–1485. Available at:
https://doi.org/10.1093/brain/114.3.1473.

Pouget, P. *et al.* (2020) 'Neuronavigated Repetitive Transcranial Ultrasound Stimulation Induces
Long-Lasting and Reversible Effects on Oculomotor Performance in Non-human Primates',
Frontiers in Physiology, 11, p. 1042. Available at: https://doi.org/10.3389/fphys.2020.01042.

Rahnev, D. *et al.* (2013) 'Continuous theta burst transcranial magnetic stimulation reduces
resting state connectivity between visual areas', *Journal of Neurophysiology*, 110(8), pp. 1811–
1821. Available at: https://doi.org/10.1152/jn.00209.2013.

Ramezanpour, H. *et al.* (2024) 'Dynamics of Saccade Trajectory Modulation by Distractors:
Neural Activity Patterns in the Frontal Eye Field', *The Journal of Neuroscience*, p.
e0635242024. Available at: https://doi.org/10.1523/JNEUROSCI.0635-24.2024.

Rincon-Gonzalez, L. *et al.* (2016) 'Decisions in motion: vestibular contributions to saccadic
target selection', *Journal of Neurophysiology*, 116(3), pp. 977–985. Available at:
https://doi.org/10.1152/jn.01071.2015.

Robinson, D.A. (1972) 'Eye movements evoked by collicular stimulation in the alert monkey',
Vision Research, 12(11), pp. 1795–1808. Available at: https://doi.org/10.1016/0042-

6989(72)90070-3.

Robinson, D.A. and Fuchs, A.F. (1969) 'Eye movements evoked by stimulation of frontal eye
fields.', *Journal of Neurophysiology*, 32(5), pp. 637–648. Available at:
https://doi.org/10.1152/jn.1969.32.5.637.

Romero, M.C. *et al.* (2019) 'Neural effects of transcranial magnetic stimulation at the single-cell
level', *Nature Communications*, 10(1), p. 2642. Available at: https://doi.org/10.1038/s41467-019-
10638-7.

Romero, M.C. *et al.* (2022) 'Neural effects of continuous theta-burst stimulation in macaque
parietal neurons', *eLife*, 11, p. e65536. Available at: https://doi.org/10.7554/eLife.65536.

Roth, Y. *et al.* (2007) 'Three-Dimensional Distribution of the Electric Field Induced in the Brain
by Transcranial Magnetic Stimulation Using Figure-8 and Deep H-Coils', *Journal of Clinical
Neurophysiology*, 24(1), pp. 31–38. Available at:
https://doi.org/10.1097/WNP.0b013e31802fa393.

Schiller, P.H., Sandell, J.H. and Maunsell, J.H. (1987) 'The effect of frontal eye field and
superior colliculus lesions on saccadic latencies in the rhesus monkey', *Journal of
Neurophysiology*, 57(4), pp. 1033–1049. Available at: https://doi.org/10.1152/jn.1987.57.4.1033.

Schlag, J., Dassonville, P. and Schlag-Rey, M. (1998) 'Interaction of the Two Frontal Eye Fields
Before Saccade Onset', *Journal of Neurophysiology*, 79(1), pp. 64–72. Available at:
https://doi.org/10.1152/jn.1998.79.1.64.

Schwartz, M.L. and Goldman-Rakic, P.S. (1984) 'Callosal and intrahemispheric connectivity of
the prefrontal association cortex in rhesus monkey: Relation between intraparietal and principal
sulcal cortex', *Journal of Comparative Neurology*, 226(3), pp. 403–420. Available at:
https://doi.org/10.1002/cne.902260309.

Sommer, M.A. and Tehovnik, E.J. (1997) 'Reversible inactivation of macaque frontal eye field',
Experimental Brain Research, 116, pp. 229–249.

Sponheim, C. *et al.* (2021) 'Longevity and reliability of chronic unit recordings using the Utah,
intracortical multi-electrode arrays', *Journal of Neural Engineering*, 18(6), p. 066044. Available
at: https://doi.org/10.1088/1741-2552/ac3eaf.

Stagg, C.J. *et al.* (2009) 'Polarity-Sensitive Modulation of Cortical Neurotransmitters by
Transcranial Stimulation', *The Journal of Neuroscience*, 29(16), pp. 5202–5206. Available at:
https://doi.org/10.1523/JNEUROSCI.4432-08.2009.

Stefan, K. *et al.* (2008) 'Theta-burst stimulation: Remote physiological and local behavioral after-
effects', *NeuroImage*, 40(1), pp. 265–274. Available at:
https://doi.org/10.1016/j.neuroimage.2007.11.037.

Strzalkowski, N.D.J. *et al.* (2019) 'Both 50 and 30 Hz continuous theta burst transcranial
magnetic stimulation depresses the cerebellum', *The Cerebellum*, 18(2), pp. 157–165. Available
at: https://doi.org/10.1007/s12311-018-0971-0.

Suppa, A. *et al.* (2008) 'Theta burst stimulation induces after-effects on contralateral primary

motor cortex excitability in humans', *The Journal of Physiology*, 586(18), pp. 4489–4500.
Available at: https://doi.org/10.1113/jphysiol.2008.156596.

Suppa, A. *et al.* (2016) 'Ten Years of Theta Burst Stimulation in Humans: Established
Knowledge, Unknowns and Prospects', *Brain Stimulation*, 9(3), pp. 323–335. Available at:
https://doi.org/10.1016/j.brs.2016.01.006.

Theleritis, C. *et al.* (2017) 'Two Versus One High-Frequency Repetitive Transcranial Magnetic
Stimulation Session per Day for Treatment-Resistant Depression: A Randomized Sham-
Controlled Trial', *The Journal of ECT*, 33(3), pp. 190–197. Available at:
https://doi.org/10.1097/YCT.0000000000000387.

Thut, G. and Pascual-Leone, A. (2010) 'A Review of Combined TMS-EEG Studies to
Characterize Lasting Effects of Repetitive TMS and Assess Their Usefulness in Cognitive and
Clinical Neuroscience', *Brain Topography*, 22(4), pp. 219–232. Available at:
https://doi.org/10.1007/s10548-009-0115-4.

Vaidya, A.R. *et al.* (2019) 'Lesion Studies in Contemporary Neuroscience', *Trends in Cognitive*
Sciences, 23(8), pp. 653–671. Available at: https://doi.org/10.1016/j.tics.2019.05.009.

Valchev, N. *et al.* (2015) 'cTBS delivered to the left somatosensory cortex changes its functional
connectivity during rest', *NeuroImage*, 114, pp. 386–397. Available at:
https://doi.org/10.1016/j.neuroimage.2015.04.017.

Valero-Cabre, A. *et al.* (2012) 'Frontal Non-Invasive Neurostimulation Modulates Antisaccade
Preparation in Non-Human Primates', *PLoS ONE*, 7(6), p. 11.

Valero-Cabré, A., Pascual-Leone, A. and Rushmore, R.J. (2008) 'Cumulative sessions of
repetitive transcranial magnetic stimulation (rTMS) build up facilitation to subsequent TMS-
mediated behavioural disruptions', *European Journal of Neuroscience*, 27(3), pp. 765–774.
Available at: https://doi.org/10.1111/j.1460-9568.2008.06045.x.

Vernet, M. *et al.* (2014) 'Frontal eye field, where art thou? Anatomy, function, and non-invasive
manipulation of frontal regions involved in eye movements and associated cognitive operations',
Frontiers in Integrative Neuroscience, 8(66). Available at:
https://doi.org/10.3389/fnint.2014.00066.

Vink, J.J.T. *et al.* (2023) 'Continuous Theta-Burst Stimulation of the Contralesional Primary
Motor Cortex for Promotion of Upper Limb Recovery After Stroke: A Randomized Controlled
Trial', *Stroke*, 54(8), pp. 1962–1971. Available at:
https://doi.org/10.1161/STROKEAHA.123.042924.

Wallis, J.D., Anderson, K.C. and Miller, E.K. (2001) 'Single neurons in prefrontal cortex encode
abstract rules', *Nature*, 411(6840), pp. 953–956. Available at: https://doi.org/10.1038/35082081.

Wegener, S.P., Johnston, K. and Everling, S. (2008) 'Microstimulation of monkey dorsolateral
prefrontal cortex impairs antisaccade performance', *Experimental Brain Research*, 190(4), pp.
463–473. Available at: https://doi.org/10.1007/s00221-008-1488-4.

Wischnewski, M. and Schutter, D.J.L.G. (2015) 'Efficacy and Time Course of Theta Burst
Stimulation in Healthy Humans', *Brain Stimulation*, 8(4), pp. 685–692. Available at:

https://doi.org/10.1016/j.brs.2015.03.004.

Zheng, C., Liao, W. and Xia, W. (2015) 'Effect of combined low-frequency repetitive transcranial
magnetic stimulation and virtual reality training on upper limb function in subacute stroke: a
double-blind randomized controlled trail', *Journal of Huazhong University of Science and*
Technology [Medical Sciences], 35(2), pp. 248–254. Available at:
https://doi.org/10.1007/s11596-015-1419-0.

Ziemann, U. and Siebner, H.R. (2015) 'Inter-subject and Inter-session Variability of Plasticity
Induction by Non-invasive Brain Stimulation: Boon or Bane?', *Brain Stimulation*, 8(3), pp. 662–
663. Available at: https://doi.org/10.1016/j.brs.2015.01.409.