Mapping the Architecture of Protein Complexes in *Arabidopsis* Using Cross-Linking Mass

Spectrometry

Cao Son Trinh[1,2]*, Ruben Shrestha[1]*, William C Conner[1], Andres V Reyes[1,2], Sumudu S

Karunadasa[1,2], Grace Liu[1], Ken Hu[1], Shou-Ling Xu[1,2]

[1]Department of Plant Biology, Carnegie Institution for Science, Stanford, California, USA;

[2]Carnegie Mass Spectrometry Facility, Carnegie Institution for Science, Stanford, California, USA

*These authors contributed equally.

Corresponding author: Shou-Ling Xu, slxu@stanford.edu

Classification: Research Reports

Keywords: cross-linking mass spectrometry (XL-MS), structural analysis, protein-protein

interaction network mapping, gene annotation.

Abstract

Capturing molecular machines in action is essential to understanding the architecture of protein complexes, cellular regulation, and gene function. Here, we present a robust cross-linking mass spectrometry (XL-MS) platform that uses the PhoX cross-linker to map proteome-wide interactions in *Arabidopsis thaliana* under semi-native conditions. Using whole-cell lysates, chloroplasts, and nuclei, we identified 47,119 unique cross-links, including 3,527 inter-protein cross-links. We then constructed a high-confidence protein-protein interaction (PPI) network containing 1,229 proteins and 1,446 PPIs. In silico validation using STRING confirmed 637 heteromeric interactions with a combined confidence score of at least 0.4, including 532 interactions with a score of at least 0.7. The remaining interactions are novel. Our dataset provides direct experimental evidence of physical interactions and defines residue-level binding interfaces. XL-MS resolved the spatial topology of structurally challenging protein complexes, such as the chloroplast 70S ribosome stalks, and transient intermediates involved in photosystem repair. XL-MS also revealed both conserved and distinct interaction patterns across photosystem complexes, ribosomes, and chromatin-associated proteins. Notably, the data suggest that the ribosome undergoes ubiquitination and is regulated by the 26S proteasome. This finding is further supported by *in vivo* diGly proteomic analysis. Among the novel interactions uncovered is a metabolite enzyme that interacts with histones, which could potentially contribute to histone modification. This work establishes XL-MS as a powerful tool for advancing the functional annotation of genomes and the understanding of the architecture of protein complexes and cellular regulation in plants, providing a valuable resource for the plant research community.

Significance Statement

Understanding how proteins interact within living cells is essential to deciphering cellular architecture and function. However, capturing native protein-protein interactions (PPIs) on a global scale has proven technically challenging. Here, we present a proteome-wide cross-linking mass spectrometry (XL-MS) platform that can systematically map direct PPIs in plant cells without requiring transgenic manipulation. This approach reveals thousands of interactions spanning major subcellular compartments and resolves the architecture of critical protein assemblies, such as photosystems, ribosomes, and chromatin-associated connectivity. We mapped both known and novel interactions. This comprehensive dataset advances protein annotation and provides a valuable resource for investigating the dynamic organization of the plant proteome.

Introduction

Understanding how molecular machines operate within cells - and capturing them in action - is essential to unraveling the principles of life and gene function. A powerful tool for this purpose is cross-linking mass spectrometry (XL-MS), an emerging high-throughput approach that enables the capture of protein complexes in their native context across diverse organisms (1, 2). Unlike affinity purification mass spectrometry (AP-MS) and proximity labeling mass spectrometry (PL-MS) (3-5), XL-MS does not require antibodies or genetically tagged transgenes. Instead, it relies on chemical cross-linkers that covalently link spatially proximal residues within or between proteins in solution, followed by protease digestion and mass spectrometric identification of the resulting cross-linked peptides. In addition to generating a "parts list" of interacting proteins, XL-MS provides direct evidence of physical interactions and precisely maps contact sites at the amino acid level, providing unparalleled structural insight into macromolecular assemblies.

Despite its promise, large-scale proteome-wide XL-MS - such as in whole-cell lysates, isolated organelles, or *in vivo* systems - remains challenging due to the inherent low abundance and heterogeneity of cross-linked peptides and the complexity of data analysis (1, 6). In plants, additional barriers further complicate its application, including rigid cell walls and large central vacuoles that dilute cross-linkers, as well as secondary metabolites that can chemically interfere with cross-linking chemistry (7, 8). Recent pioneering studies in Arabidopsis, algae, and soybean have demonstrated the potential of XL-MS in plant systems (7, 9, 10), but its overall implementation remains limited.

To address these challenges, we developed a cross-linking strategy optimized for proteome-wide mapping in complex Arabidopsis samples using the enrichable cross-linker PhoX. PhoX is a phosphonic acid-based reagent that, like BS3 and DSSO, contains NHS esters and

primarily targets lysine residues (11). Its phosphonic acid handle allows efficient enrichment by

immobilized metal affinity chromatography (IMAC). With a maximum range of 35Å - consisting

of a 20 Å Cα-Cα distance from its rigid structure and an additional 15 Å to account for protein

flexibility - PhoX is well suited for high confidence, large-scale structural proteomics (12). To

reduce sample complexity and improve cross-linking efficiency, we used subcellular

fractionations. To further improve the detection of the low abundance cross-linked peptides, we

performed extensive high-pH reversed-phase HPLC fractionation after IMAC enrichment.

Together, these measures allowed for comprehensive proteome-wide XL-MS mapping.

We identified a total of 47,119 unique cross-linked peptide pairs from cross-linked cell

lysates, chloroplasts, and nuclei, including 43,592 intra-protein (self-link) and 3,527 inter-protein

cross-links. These results enabled us to construct a PPI network comprising 1,229 proteins and

1,446 interactions, of which 1,390 are heteromeric and 56 are homomultimeric. STRING analysis,

which catalogs only heteromeric PPIs, validated 637 interactions with a combined confidence

score of at least 0.4. The rest represent previously uncharacterized interactions. Our dataset

provides experimental evidence of direct physical binding and defines protein binding interfaces

with amino acid resolution for both known and novel interactions.

Beyond interaction mapping, our XL-MS data provide structural and functional insights

into key cellular processes. By resolving flexible or transient components within the 70S

chloroplast ribosome and photosystem complexes, our results complement existing cryo-EM

structures. XL-MS also enabled a comprehensive mapping of the photosystem, 80S ribosome, and

histone interactomes within heterogeneous protein populations, revealing structural biology and

capturing both known and novel interactions. We identified a metabolic O-acyltransferase that

interacts with histones, suggesting a potential new mechanism of chromatin regulation via post-

translational modification. Together, these results improve protein annotation, shed light on

protein complex architectures, and provide as a valuable resource for the broader research

community - laying the foundation for future discoveries.

Results

Efficient cross-linking workflow for complex protein samples identifies numerous cross-links

in Arabidopsis

To gain structural insights and visualize molecular machines in their native cellular context within

Arabidopsis cells, we performed XL-MS (see Methods) on fractionated seedling samples,

including cell lysates and chloroplasts from light-grown seedlings and nuclei from dark-grown

seedlings. Using the PhoX cross-linker (Fig. 1A), we established a robust pipeline optimized for

proteome-wide detection of low-abundance cross-linked peptides in complex Arabidopsis

samples.

Optimization of the PhoX concentration showed a shift to higher molecular weights at 2-5

mM (red arrows, Fig. S1A-D), indicating successful cross-linking, with no additional

improvement at higher concentrations. Based on these results, we used 2 and 5 mM PhoX for

subsequent experiments (see Methods). After cross-linking, we precipitated the proteins with

phenol, digested them with trypsin, and enriched the cross-linked peptides using $GaCl_3$-based

IMAC (13). We then fractionated the enriched peptides by high-pH reversed-phase

chromatography, analyzed them by LC-MS/MS, and identified cross-links using pLINK2 (14).

High-resolution, high-accuracy data for both MS1 (precursor ions) and MS2 (fragment ions) were

acquired on an Orbitrap Eclipse mass spectrometer to ensure reliable cross-link identification.

To optimize data acquisition, we compared the standard XL workflow with real-time

spectral library search (RTLS) (15), both targeting highly charged precursors (+3 and above).

RTLS uses a survey MS2 scan to reduce mono-link detection. In cell lysates, RTLS reduced mono-

and loop-link identifications by ~50% and ~31%, respectively (Fig. S2), but also significantly

reduced intra-link (43%) and inter-link (40%) identifications. Because of this trade-off, we used

the standard XL workflow to analyze cross-linked chloroplast and nuclear samples.

In total, we identified 47,119 unique cross-linked peptide pairs (intra- and inter-links)

representing 5,759 proteins. Specifically, we detected 25,795, 17,090, and 19,612 cross-links from

the nucleus, chloroplast, and cell lysate, respectively (Fig. 1B and. S3, Table S1). Subcellular

localization analysis using SUBA5 revealed that the cross-linked proteins were distributed in

various cellular compartments, including the nucleus, plastid, cytosol, mitochondria and other

organelles (Fig. 1C, Table S2).

To assess the quality of cross-linked peptide identification, we examined the identified

cross-links (Table S1) and found consistent peptide pairs mapping to the same sites or regions. For

example, in ribosomal subunits, we detected site-specific cross-links with variable miscleavages,

such as those between uL15x-K92 and eL13z-K166/K172 (Fig. S4A). Conserved region cross-

links were also observed, such as VHA-E3-K49 with both VHA-B1-K437 and VHA-B2-K438

(Fig. S4B), and TOP1A-K672 with ERD14-K38 and COR47-K46 (Fig. S4C). These independent

and consistent observations underscore the robustness and reliability of our results.

Validation of intra- and inter-protein cross-links by structural mapping

To evaluate the validity of the cross-links, we first used ComMap (16) to map intra-links to high-

resolution protein structures available in the Protein Data Bank (PDB) and predicted structures by

AlphaFold. Specifically, 721 Arabidopsis proteins had high-resolution structures in the PDB, and

3,975 were available through AlphaFold (Table S1). A total of 21,981 cross-links were

successfully mapped, with 94% of the PDB-mapped cross-links and 89% of the AlphaFold-

mapped cross-links satisfying the 35 Å distance constraint (Fig. 2A-B).

To evaluate both intra- and inter-protein cross-links, we mapped 174 cross-links onto the

L_8S_8 Rubisco assembly modeled from the Arabidopsis L_2S_2 crystal structure (PDB:5IU0) (17). Of

these, 169 cross-links were mapped to the large (479 aa) and small (181 aa) subunits, while five

could not be mapped because the PDB structure lacks the N-terminal region (residues 1–11) (Fig.

2C). The majority (126, 74.6%) of mapped cross-links fell within the expected $C\alpha$-$C\alpha$ distance of

≤35 Å; the remaining 43 (25.4%) exceeded this range, suggesting regions of structural flexibility

(Fig. 2D–E). Similar mapping results were obtained using the hornwort Rubisco cryo-EM structure

(Fig. S5A-B), further supporting the validity of our cross-linking data.

To assess whether cross-linked lysine residues are positioned within predicted interaction

interfaces, we examined several PPIs using available structural data. The cross-link between

ACINUS-K633 and SR45-K154 maps near the RSB domain of ACINUS (aa 599-631) and within

the RRM domain of SR45 in the ASAP core complex (PDB:4A8X) (18-20) (Fig. 2F, Fig. S6A).

Similarly, the cross-link between SR45-K154 and PININ-K274 aligns with the same RRM domain

and the predicted RSB domain of PININ (Fig. 2F, Fig. S6B). For PRK-K217 and CP12.1-K65, the

cross-link localizes to the interaction interface in the GAPDH-CP12-PRK complex (PDB:6KEZ)

(21) with a $C\alpha$-$C\alpha$ distance of 13.3 Å (Fig. 2G-H). These results confirm that the cross-linked

residues are located at or within a few amino acids of the interaction interfaces, consistent with

previous observations (22).

XL-MS systematically identifies direct protein-protein interactions and reveals known and

novel complexes

We imported 3,527 inter-protein cross-links into Cytoscape (23), revealing 1,446 PPIs

among 1,229 proteins, including 1,390 heteromeric and 56 homomultimeric interactions (Fig. 3A,

Table S1). Proteins were classified using the PANTHER protein class tool, and nodes were

colored accordingly (24). Many interacting proteins shared the same classification, supporting the

reliability of our PPI map. Notably, 457 of the 1,210 proteins involved in heteromeric interactions

lacked PANTHER annotations, contributing to 819 interactions - 369 between unclassified

proteins and 450 between classified and unclassified proteins.

We also incorporated STRING confidence scores (25) into the network and color-coded

the edges accordingly (Fig. 3A). These scores combine evidence from eight sources (Fig.3B) (25).

For heteromeric PPIs, 38.2% of PPIs had high STRING confidence: 388 scored $\geq$ 0.9 and 144

scored 0.7-0.9. The remainder included 105 medium (0.4-0.7), 144 low (<0.4), and 609 with no

STRING score (Fig.3B). The PANTHER and STRING results indicate that both known and novel

PPIs were captured, highlighting the power of systematic XL-MS to reveal previously

uncharacterized protein complexes and pathways.

Even among PPIs with good STRING combined scores, many lack experimental support

for direct or indirect interactions in Arabidopsis or homologs in other species. Among the

heteromeric PPIs identified, only 28.9% (n = 402) have experimental support (score $\geq$ 0.4), while

71.1% have low (<0.4) or no scores (Fig. 3B). Our XL-MS data provide evidence for their direct

interactions and contact sites. For example, ORP3B (AT3G09300, oxysterol-binding protein) and

PLDRP1 (AT5G39570, PLD-regulated protein 1) have a combined score of 0.602 but a low

experimental score (0.055); we identified a cross-link between ORP3B-K7 and PLDRP1-K193

(Fig. 3C), suggesting a common role in lipid trafficking and signaling. Similarly, NRPA2

(AT1G29940, RNA polymerase predicted to localize to both the nucleus and chloroplasts) and

uL1c (AT3G63490, chloroplast 50S ribosomal protein) have a combined score of 0.588 but no

experimental support; a cross-link between NRPA2-K394 and uL1c-K127 suggests a direct

interaction and potential coupling of chloroplast transcription and translation (Fig. 3D).

XL-MS resolves the topology of chloroplast 70S ribosome stalk regions inaccessible by Cryo-EM

We mapped the topologies of several protein complexes (Fig. S7A-G), including the flexible stalk regions of the chloroplast 70S ribosome (Fig. 4A-C), which have remained difficult to resolve due to the dynamic nature of proteins such as uL1c, uL10c, uL11c, and bL12c (26, 27). In total, we identified 424 cross-links across 34 ribosomal subunits of the 70S ribosome, including 352 intraprotein, 64 heteromeric, and 8 homomultimeric cross-links. Notably, many intra- and interprotein cross-links involved these flexible stalk proteins. When cross-links were mapped to spinach 70S ribosome structures (PDB: 5MMM and 5X8P; Fig. S8A-B), most cross-links involving uL1c, uL10c, uL11c, and bL12c could not be precisely assigned to specific locations within the stalk regions because these proteins are either missing or only partially resolved in the cryo-EM structures.

Arabidopsis bL12c has a plant-specific disordered N-terminus, followed by conserved structured N-terminal (NTD) and C-terminal (CTD) domains connected by a flexible hinge (Fig. 4B). Essential for translation, bL12c uses hinge-mediated mobility to recruit specific translation factors(28). XL-MS identified interactions between bL12c (the two highly homologous proteins AT3G27850/bL12cx and AT3G27830/bL12cz) and several ribosomal core proteins, as well as the elongation factor CPEF-G, mainly through its structured CTD (Fig. 4C). Although bL12c dimers are typically mediated by the NTD (29), XL-MS revealed unexpected homodimer cross-links within the CTD - the same region involved in translation factor binding (Fig. 4C). This suggests that CTD dimerization may serve as an interface for the recruitment of interaction partners. Taken together, these results highlight both conserved and plant-specific features of bL12c and demonstrate the power of XL-MS in resolving dynamic aspects of ribosomal architecture.

XL-MS reveals dynamic connectivity and unique features in photosystem complexes

Photosystems I and II (PSI and PSII) and their associated light-harvesting complexes (LHCI and LHCII) are essential for light energy harvesting and carbon fixation, but they require continuous repair, especially under high and fluctuating light conditions (30-32). Although progress has been made in understanding certain steps of the PSII repair (33, 34), many aspects, including the detailed mechanism, identification of repair intermediates, and specific contributing factors, remain unresolved.

Using XL-MS, we identified extensive interactions involving the PSII repair factor Psb27 (AT1G03600) (Fig. 5A). Unlike its cyanobacterial counterpart, which functions in both assembly and repair, Arabidopsis Psb27 appears to be specialized for repair. In contrast, the less homologous Psb27-H1(AT1G05385) is involved in assembly. Notably, Arabidopsis Psb27 lacks a lipid anchor and is structurally more flexible than the cyanobacterial version (35) (Fig. S9A). A partial structure from Chlamydomonas (PDB: 6KAC) captured residues 91-114 of Psb27 (36). Our XL-MS data revealed that Psb27 interacts with both the PSII reaction center and the oxygen-evolving complex (OEC), involving K93 (as in 6KAC) and a second site at K168 (Fig. 5A, Fig. S9B, S10A-H). These distance constraints (Fig. S9B) suggest that Psb27 engages PSII in a more dynamic and flexible manner in plants.

XL-MS also revealed extensive interactions between PSI, PSII, LHCI, LHCII, ATP synthase, ATP-dependent proteases (FtsH, SPPA) and transient proteins involved in state transitions, complex assembly and repair (Fig. 5B). Notably, chlorophyll biosynthesis and catabolic proteins (e.g., ACCELERATED CELL DEATH 1 (ACD1), magnesium protoporphyrin IX O-methyltransferase (CHLM), FLUORESCENT IN BLUE LIGHT (FLU)) were found to directly associate with LHCII, suggesting coordinated chlorophyll delivery and reactive oxygen

species (ROS) control during LHCII assembly. In addition, many assembly and repair factors were

found to associate with LHCII, indicating their readiness for rapid engagement during dynamic

photosystem remodeling.

XL-MS of ribosome complexes reveals conserved ribosome-associated factors

We identified extensive interactions involving the 80S ribosome, a complex of 60S and

40S subunits that synthesizes nuclear-encoded proteins. Ribosomes assemble in the nucleolus,

localize to the cytoplasm, and associate with the endoplasmic reticulum (ER). As a dynamic

macromolecular machine, the ribosome coordinates translation with numerous proteins and RNAs.

Errors in translation that produce misfolded proteins, or defects in the ribosome itself, can trigger

ribosome-associated quality control pathways (37). Using XL-MS, we mapped detailed structural

information within this highly heterogeneous complex, capturing not only intra- and inter-subunit

contacts (Fig. S11A-D), but also interactions with proteins involved in ribosome biogenesis,

translation, folding, quality control, stress response, and other regulatory pathways (Fig. 6A).

These interactions were resolved at the amino acid level, providing detailed insight into the

architecture of the functional interface (Fig. 6B, Table S4).

While many interactions showed good STRING scores (≥0.4), our analysis also revealed

numerous novel interactions (green edges) between specific protein pairs. Although homologs of

these proteins have been studied in other organisms for their ribosome-related functions, their

specific roles in plants remain largely uncharacterized. Our XL-MS data suggest conserved

functions: for example, the UTP7 homologs AT3G10530 and NOG1-1 may contribute to ribosome

biogenesis (38, 39), whereas AT5G50840 (α-taxilin-like) and ILITHYIA(ILA) may be involved

in protein folding or translation (40, 41). MBF1A, a transcriptional coactivator, may be involved

in stress responses at stalled ribosomes, similar to its role in yeast (42) . Other candidates include

a methyltransferase potentially involved in rRNA processing and polysome assembly (43, 44) , a

RIO kinase associated with ribosome biogenesis (45, 46), and PARP1, which may regulate

translation via ADP-ribosylation (47) (Fig. 6C). By mapping these interactions to specific

ribosomal proteins, our XL-MS dataset provides a valuable resource for future functional studies

of ribosome-associated factors in plants.

We identified several interactors related to ubiquitination, including two E3 ligases (EBF1

and AT4G02760) and one deubiquitinase UBP14. In addition, we detected direct interactions

between ribosomes and 26S proteasome components, such as RPN8A and PAD1 (Fig.6A). Given

the need for active ribosome turnover to maintain quality control and these XL interaction data,

we hypothesized that ribosome proteins undergo *in vivo* ubiquitination as a regulatory mechanism.

To test this hypothesis, we conducted a diGly ubiquitin proteomic analysis (48), which revealed

the *in vivo* ubiquitination of 37 large and 27 small ribosomal subunits (Fig. 6A, Table S4). These

findings suggest that the ribosomes are ubiquitinated *in vivo* and may be targeted for degradation

by the 26S proteosome.

Histone interactome reveals conserved and novel chromatin-associated interactions

Our XL-MS analysis revealed extensive interactions between histones and both known and

novel interactors, providing new insights into the networks involved in gene expression and

chromatin regulation (Fig. 7A). We identified several nuclear complexes associated with

chromatin, including nucleosome assembly proteins (NAPs), chromosome structure maintenance

proteins (SMCs), nuclear RNA polymerase components (NRPs), chromatin regulators (PWWP

DOMAIN PROTEIN 3(PDP3), MULTICOPY SUPPRESSOR OF IRA1 4(MSI4)), transcriptional

activator and acetylation reader (AT2G36410 and ACETYLATED INTERACTING PROTEIN

1(ACIP1)), and subunits of the PAF complex (VERNALIZATION INDEPENDENCE 2 and 4

(VIP2, VIP4)) (Fig. 7A).

Although many proteins are known to influence chromatin architecture and gene regulation

in plants, their direct interactions with histones and the nature of their contact sites remain largely

uncharacterized. Using XL-MS, we identified a diverse array of proteins that interact with histone,

including chaperones such as SUVH1/3-interacting DNAJ domain-containing protein 2(SDJ2) and

NUCLEOLIN 1(NUC-L1)), chromatin remodeling factors such as EARLY BOLTING IN SHORT

DAYS(EBS), SPLAYED(SYD) and SPIKE1(SPK1)), transcription factors such as SQUAMOSA

PROMOTER BINDING PROTEIN-LIKE 3(SPL3), co-regulators such as SIN3-LIKE 6(SNL6),

DNA repair proteins such as MUTS HOMOLOG 2 (MSH2), and structural proteins such as HIGH-

MOBILITY GROUP BOX 7 (HMGB7). Notably, we found that MOS7, a nucleoporin

homologous to the human NUP88, cross-linked with several histones. This provides direct

evidence of nuclear envelope-histone interactions that may contribute to chromatin organization

in plants (49). In addition, we identified histone interactions with several enzymatic regulators,

including the kinase MLK4, the poly(ADP-ribose) polymerase PARP1 (both of which were cross-

linked to H1), and the ubiquitin-like protein NEDD8 (which was cross-linked to H2B and H3).

Surprisingly, XL-MS identified four distinct cross-links between histone H2B and the O-

acyltransferase WSD4 (Fig. 7A and S12A-E). Members of the WSD family, including WSD4, are

traditionally associated with lipid metabolism and function as wax ester synthase/acyl-CoA-

dependent O-acetyltransferases (50, 51). However, WSD4 is preferentially expressed in roots,

which is unusual because wax deposition is uncommon in these tissues (51). This suggests that

WSD4 may have an alternative function unrelated to wax biosynthesis. Notably, TAIR predicts

that WSD4 is localized in nucleus. To test whether WSD4 co-localizes with histones, we co-

expressed mCherry-WSD4 and GFP-HTB11 in *Nicotiana benthamiana*, and observed overlapping

nuclear signals, as indicated by the arrows in Fig. 7B. Together, the XL-MS and co-localization

data raise the intriguing possibility that WSD4 may catalyze histone acylation, a hypothesis that

warrants further investigation (Fig.7C).

Taken together, these findings reveal previously uncharacterized histone-associated

proteins and enzymatic activities in plants, providing new insights into chromatin biology and

laying the groundwork for future studies of plant gene regulation.

Discussion

Systematic mapping of PPIs has greatly advanced our understanding of cellular function,

but plant PPI networks remain largely unexplored (5, 52). While comprehensive maps exist for

humans and other model organisms (53-55), similar efforts are just beginning in plants (56-58).

This gap represents a critical opportunity, as plants are central to addressing global challenges such

as food security, climate change, and sustainability. Expanding proteome-wide PPI mapping and

structural biology studies in plants will accelerate our mechanistic understanding of cellular

processes and drive advances in agriculture and biotechnology.

We developed a systems biology approach to capture molecular machines in action by

mapping PPIs and their interfaces using XL-MS. To address the inherent challenges of XL-MS in

complex biological samples, we established an optimized pipeline that integrates the PhoX

crosslinker, subcellular fractionation, and high-pH reversed-phase HPLC offline fractionation.

This workflow enabled the identification of over 47,000 crosslinks. The high quality of the data

was confirmed by mapping the intralinks onto experimentally determined structures in the PDB

and predicted models from AlphaFold.

Our XL-MS analysis revealed the topological organization of multiple protein complexes,

including the 70S ribosome, 26S proteasome, photosystems I and II, ATP synthases, and

chromatin-associated factors. We demonstrate that XL-MS is highly effective in resolving spatial

relationships within dynamic and heterogenous assemblies. Notably, we mapped flexible regions of the 70S ribosome stalk and established connectivity for components that are missing or only partially resolved in the existing cryo-EM structures. This includes flexible elements such as BL12c, for which we discovered unexpected C-terminal domain (CTD) dimerization within the translation factor binding region – suggesting that CTD dimerization may serve as an interface for recruiting interaction partners. Additionally, we mapped interactions involving transient repair proteins in PSII. Together, these results highlight the unique strength of XL-MS in complementing cryo-EM by providing structural insights into conformationally dynamic regions, underscoring the value of our approach in advancing in situ structural biology.

Our XL-MS analysis identified 1,446 high-confidence PPIs among 1,229 proteins, establishing a comprehensive interaction map in *Arabidopsis*. Functional categorization using PANTHER and network analysis via STRING analysis revealed that these interactions span a wide range of biological processes. Notably, 637 heteromeric PPIs had a combined STRING confidence score ≥ 0.4, including 532 with score ≥ 0.7, underscoring the robustness of our interaction mapping. In addition, we identified 56 homomultimeric PPIs, which are not represented in STRING, as the database catalogs only heteromeric interactions. Recent studies suggest that dimerization is essential for the function of many proteins (59, 60).

Further validation of our dataset is provided by genetic evidence: mutations in several newly identified interacting pairs result in shared or opposing phenotypic outcomes. For example, the chromatin remodeling protein EBS promotes early flowering, whereas its interactor SYD, a component of the chromatin structure remodeling complex, delays flowering when mutated(61, 62). In another example, loss-of-function mutations in both FLK (a flowering locus KH domain protein) and its interactor KHZ1 (a CCCH zinc finger and KH domain protein) result in late

flowering (63, 64). These phenotype correlations support the functional relevance of the identified

interactions and highlight the potential of XL-MS-based interaction maps to accelerate gene

discovery.

In addition, the interaction between ribosomal proteins and E3 ligases, deubiquitinases,

and 26S proteasome components, such as RPN8A and PAD1, suggests that ribosomes may be

regulated by ubiquitin-mediated pathways. Our ubiquitin diGly proteomics analysis supports this

hypothesis by identifying the *in vivo* ubiquitination of 37 large and 27 small ribosomal subunits.

These identifications were of high confidence, with many di-Gly peptides detected in both light

(^{14}N) and heavy (^{15}N) forms (65, 66). Together with the XL-MS data, these findings imply the

existence of a potential ribosome quality control mechanism similar to those pathways described

in yeast (67). The exact role of E3 ligases and deubiquitinases remains to be confirmed

functionally. EBF1 has been shown to regulate ethylene signaling by targeting EIN3/EIL1 for

degradation (47); however, its ribosome-related functions have yet to be explored. UBP14 is

essential for development in Arabidopsis but dispensable in yeast (48).

XL-MS uncovered a wealth of novel PPIs, with over half not previously annotated in the

STRING database. For example, we identified both well-established and unreported PPIs

involving ribosomal proteins and histones (see Figs. 6A and 7A), expanding our understanding of

these conserved molecular machines across species (68, 69). We also uncovered a broad array of

plant-specific PPIs, particularly within chloroplasts (Fig. 5B), reflecting unique features of the

plant cellular machinery.

Among the histone interactors, we identified several enzymatic regulators, including

MLK4 and PARP1 (cross-linked to H1), and NEDD8 (cross-linked to H2B and H3). Although

PARP1 and NEDD8 are known histone modifiers in other systems, their roles in plants remain

undefined (70-73). Previous studies have shown that MLK4 phosphorylates H2A and regulate

flowering (74); thus, it may have broader functions in histone regulation.

We identified WSD4, an O-acyltransferase, as a potential novel histone regulator. Four

independent cross-links were identified between WSD4 and H2B, and their interaction was further

supported by nuclear co-localization. Although O-acyltransferase are usually linked with lipid

metabolism(50), their function in protein modification remains largely unknown. Given the known

examples of protein acylation by metabolic enzymes and the regulatory role of lysine acylation(75-

78), we propose that WSD4 may catalyze histone acylation. This would reveal a previously

unrecognized chromatin regulatory mechanism.

Despite decades of study, many *Arabidopsis* genes remain uncharacterized (79). Our work

demonstrates that XL-MS efficiently maps PPIs under semi-native conditions, accelerating gene

discovery and functional annotation. In addition, the extensive intra-protein cross-links generated

in this study can aid in developing structural prediction tools such as AlphaFold. This dataset is a

valuable resource for Arabidopsis functional genomics, and this approach can be applied to other

species, particularly crops. Further developing this technique will advance comparative proteomics

and deepen our understanding of fundamental biological processes.

Acknowledgement

This work was supported by the National Institutes of Health (NIGMS R01GM135706 and

S10OD030441 to S-L.X.) and by the Carnegie endowment fund for the Carnegie Mass

Spectrometry Facility. We thank Prof. Laura Helen Gunn for providing the structure of

Arabidopsis Rubisco L8S8, and Drs. Mike Trnka and Robert Chalkley (UCSF) and Dr. Rosa Viner

(Thermo Fisher Scientific) for technical assistance. We also thank the Hong Qiao lab (UT Austin)

for the nuclear isolation protocol and the Adrien Burlacot lab for the chloroplast isolation protocol.

We thank Dr. Ajeet Chaudhary and Prof. Arthur Grossman for their critical reading of the

manuscript.

Author contributions

Data Availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the

PRIDE partner repository with the dataset accession numbers PXD066234 and PXD066291. All other data

is available from the corresponding author on reasonable request.

Supplemental Table

Table S1: Summary of cross-linked peptide, including intra- and inter- protein cross-links, PPI annotations

based on STRING and mapping of intra-cross-links to known structures from PDB and predicted structure

from AlphaFold.

Table S2: Summary of cross-linked protein information and SUBA annotation.

Table S3: Analysis of protein classification using PANTHER.

Table S4: Detected cross-links for ribosomal and histones PPIs, and a list of ribosome-related ubiquitinated

peptides.

Table S5: Names, accession numbers and description used in the figures and a glossary.

Material and methods

Plant growth condition

Arabidopsis thaliana wild-type (Col-0) seeds were sterilized and sown on ½ Murashige and

Skoog (MS) agar square plates supplemented with 0.6% (w/v) phytoblend and lined with nylon

fabric. After stratification at 4°C for 3 days, the plates were transferred to growth conditions.

Light-grown seedlings: plants were vertically grown in a growth chamber at 22°C under a

16-hour light / 8-hour dark cycle for 10-12 days, then flash frozen in liquid nitrogen and stored at

-80°C for crosslinking experiments on cell lysates and isolated chloroplasts. Dark-grown

seedlings: plants were grown horizontally at 22°C in complete darkness for 3 days in a growth

chamber, then flash frozen in liquid nitrogen and stored at -80°C for nucleus isolation. Dark growth

was used to reduce Rubisco content.

Frozen Arabidopsis tissue was cryo-ground to a fine powder. Different amounts of tissue

were used for cross-linking experiments: 6 g from 10 d seedlings for cell lysates, 195 g from 12d

seedlings for chloroplasts, and 32 g (replicate 1 with PhoX) or 14 g (replicate 2 with tbu-PhoX)

from 3d dark-grown seedlings for nuclei.

Fraction of plant seedling tissues for cell lysate, chloroplast and nucleus

Cell lysates: Homogenized 10 d old light growth tissues were extracted using MOPS buffer [100

mM MOPS pH 7.6, 150 mM NaCl, 1% Nonidet P-40, 1 mM PMSF, 1x Roche protease inhibitor]

at a tissue:buffer ratio of 1:1 (g/mL, w/v). The samples were centrifuged at 20,000 g for 15 min at

4°C, and the supernatant was then filtered twice with Miracloth and used for the cross-linking

reaction.

Chloroplast isolation: Frozen, homogenized tissue powder from 12 d old seedlings was

mixed with a grinding buffer similar to (80), using HEPES instead of Tricine for crosslinker

compatibility. Briefly, 5 g of tissue powder was ground in a mortar and pestle with 30ml of

grinding buffer [20mM HEPES-KOH pH 8.0, 330mM sorbitol, 5 mM EDTA, 5 mM EGTA, 5 mM

MgCl$_2$, and 10 mM NaHCO$_3$] at 4°C, and under green illumination only. The resuspended

homogenate was filtered twice through two layers of Miracloth. 30 ml of the filtrate was carefully

transferred onto 10 ml of 40% Percoll [20 mM HEPES-KOH pH 8.0, 330 mM sorbitol, 40% v/v

Percoll, 2.5 mM EDTA, 5 mM MgCl$_2$, and 10 mM NaHCO$_3$] in a 50ml centrifuge tube, and

centrifuged at 2,600g for 10 min at 4°C. The bottom pellet, enriched in intact chloroplasts, was

collected and resuspended in 30 ml of wash buffer [20 mM HEPES-KOH pH 8.0, 330 mM sorbitol,

2.5 mM EDTA, 2 mM MgCl$_2$, and 10 mM NaHCO$_3$], and centrifuged at 2,600g for 3 min at 4°C.

Each pellet derived from 5 g of tissue was washed twice with 4.5 ml of 1x PBS supplemented with

MgCl$_2$ [137 mM NaCl, 2.7 mM KCl, 10 mM Na$_2$HPO$_4$, 1.8 mM KH$_2$PO$_4$, and 2.5 mM MgCl$_2$]

and then resuspended in 1mL 1x PBS supplemented with MgCl$_2$ for the cross-linking reaction.

This process was repeated across multiple batches, with a total of 195 g of tissue used for cross-

linking.

Nuclear isolation: The method for nuclear isolation was adapted from (81) with slight

modification. Powdered dark-growth Arabidopsis tissue of approximately 2 g was completely

homogenized with lysis buffer [20 mM HEPES (pH 7.4), 20 mM KCl, 2 mM EDTA, 2.5 mM

MgCl$_2$, 25% glycerol, 250 mM sucrose, 1 mM dithiothreitol, 1 mM phenylmethylsulfonyl fluoride

(PMSF), 1× protease inhibitor cocktail, and 0.2% formaldehyde] at tissue:buffer ratio of 1:5 (g/mL,

w/v). The homogenate was then filtered through a 40 μm cell strainer and centrifuged at 4°C,

1,500g for 10 min. The pellet was retained as the nuclear fraction. The nuclear pellet was then

washed 5 times with 5 mL of nuclear washing buffer [20 mM HEPES (pH 7.4), 2.5 mM MgCl$_2$,

25% glycerol, 0.2% Triton X-100, 1 mM PMSF, and 1× protease inhibitor cocktail] and

centrifuged at 4°C, 1,500g for 10 min to collect the pellets after each wash. The washed nuclear

pellet was then resuspended in 0.5 mL of nuclear resuspension buffer 1 [20 mM HEPES (pH 7.4),

10 mM MgCl$_2$, 250 mM sucrose, 0.2% Triton X-100, 1 mM PMSF, and 1× protease inhibitor

cocktail] and carefully laid on top of nuclear resuspension buffer 2 [20 mM HEPES (pH 7.4), 10

mM MgCl$_2$, 1.7 M sucrose, 0.2% Triton X-100, 1 mM PMSF, and 1× protease inhibitor cocktail],

then centrifuged at 4°C,16,000g for 45 min. The supernatant was carefully removed, and the

nuclear pellet was resuspended in 0.5 mL 1 X PBS (pH 7.5) buffer for crosslinking reaction. The

intactness and purity of the nuclear pellet was assessed by DAPI staining under a Leica SP8

confocal microscope. Multiple samples were prepared in parallel, and final isolated nuclei were

pooled before crosslinking.

Cross-linking reaction

PhoX or tBU-PhoX cross-linker was freshly prepared at a stock concentration of 100 mM in

DMSO for each cross-linking experiment (supplementary Fig. S3). A working concentration of 2-

5 mM cross-linker was added to buffer solution containing whole cell lysate, isolated chloroplasts

or isolated nuclei and incubated for 2 h at 4°C (dark condition for chloroplast cross-linking) with

end-to-end rotation. The cross-linking reaction was quenched by adding Tris pH 8.0 to a final

concentration of 20 mM and subjected to protein extraction.

Protein extraction, digestion, IMAC enrichment and RPLC fractionation

Protein extraction from the cross-linked samples, digestion to peptides, immobilized metal affinity

chromatography (IMAC) enrichment, and high pH reverse HPLC fractionation followed the same

procedure as described in (13). Multiple batches of PhoX-cross-linked peptides from isolated intact

chloroplasts were combined before IMAC enrichment. More information can be found in

supplementary Fig. S3. Multiple batches of samples were combined prior to high pH reversed-

phase HPLC offline fractionation (RPLC) on Vanquish Flex HPLC, and fractions were collected

for further LC-MS/MS analysis.

To generate selected proteomic libraries, total peptides from cell lysate and chloroplast samples

were analyzed by LC-MS/MS. Chloroplast peptides were further fractionated into eight fractions

using the Pierce High pH Reverse-Phase Peptide Fractionation Kit.

diGly proteomics analysis to identify ubiquitinated proteins *in vivo*

Metabolically labeled Arabidopsis plants with ^{14}N and ^{15}N isotope were harvested, followed by

protein extraction and trypsin digestion as previously described (13, 65). Ubiquitinated peptides

were enriched using the PTMScan® Ubiquitin Remnant Motif (K-ε-GG) Kit (Cell Signaling

Technology) according to the manufacturer's instructions. The enriched peptides were then

analyzed by LC-MS/MS.

LC-MS/MS analysis

Cross-linked peptides were analyzed by liquid chromatography-tandem mass spectrometry (LC-

MS/MS) using an Easy-nLC 1200 ultra-performance liquid chromatography (UPLC) system

(Thermo Fisher) coupled to an Orbitrap Eclipse Tribrid quadrupole Orbitrap mass spectrometer

(Thermo Fisher). Peptides were first loaded onto a C18 trap column (Acclaim PepMap 100 C18)

and then separated on an analytical C18 column (Aurora Series, 25 cm × 75 μm ID, Ion Opticks).

The flow rate was set to 300 nL/min and the separation was performed over a 120-minute gradient.

Peptides were eluted using a gradient from 3% to 28% solvent B over 106 minutes, followed by

an increase from 28% to 44% over 15 minutes and a final wash at 90% solvent B for 14 minutes.

Solvent A consisted of 0.1% formic acid and solvent B consisted of 80% (v/v) acetonitrile with

0.1% (v/v) formic acid.

Data was acquired on an Orbitrap Eclipse mass spectrometer. Full MS scans were acquired

over a mass-to-charge (m/z) range of 375–1600 at a resolution of 120,000, with an automatic gain

control (AGC) target of 200,000, a maximum injection time of 50 ms, normalized AGC target set

509 to 50%, and RF lens set to 30%. The most intense multiply charged precursors (charge states 2–8)

510 were selected for fragmentation with a cycle time of 3 seconds. MS/MS scans were acquired at a

511 resolution of 15,000 with an AGC target of 5×10^4, maximum injection time of 22 ms, isolation

512 window of 1.4 m/z, and normalized AGC target set to 100%. Fragmentation was performed using

513 higher-energy collisional dissociation (HCD) with a normalized collision energy (NCE) of 27.

514 Dynamic exclusion was enabled for 30 seconds. For cross-linked peptide detection, charge states

515 3 to 8 were used.

516 For Real-time library search (RTLS) acquisition (15), a low-resolution MS2 survey scan

517 performed in the ion trap over an m/z range of 120–500 was included. The resulting MS2 peaks

518 were matched against a custom spectral library containing signature peak patterns characteristic of

519 mono-links and cross-links (signature m/z values: 201.1231, 215.1387, 312.0632, and 377.1269,

520 each with characteristic relative intensities for mono- or cross-linked species). Only precursor ions

521 whose survey MS2 spectra matched the cross-link pattern using the "Similarity Search" function

522 were selected for high-resolution MS2 acquisition.

523 **Data search using pLINK2 and Protein Prospector**

524 Liner peptide searches were searched using Protein Prospector against Arabidopsis TAIR database

525 as described in (65). Precursor and fragment mass tolerance were set up as 10 and 20 ppm.

526 Crosslinked peptide searches were performed using pLink 2.3.11 with the following

527 parameters: crosslinker set to PhoX (209.972 Da); peptide length range: 6-60 amino acids;

528 enzyme: trypsin with up to 3 missed cleavages. Fixed modification: Cys carbamidomethylation;

529 variable modifications: protein N-terminal acetylation and Met oxidation. Precursor and fragment

530 mass tolerances were set to 5 ppm and 20 ppm, respectively.

MS2 spectra were searched against the TAIR 10 protein database (35,386 entries), as well

as selective database for cell lysate (7,835 entries) and chloroplast (9,897 entries), as described in

Supplementary Fig.S3. Search results were filtered at 5% FDR at the peptide spectrum matches

(PSMs) level. FDRs for monolinks, loop-links, intra- and inter-protein cross-links were calculated

separately using pLINK2. To compare two acquisition methods (XL-Standard [Standard] and

Real-time library search [RTLS]), data were acquired on the whole cell lysate XL samples, search

against TAIR10 database, and directly compared.

Data compilation and reporting

Intra- and inter-protein cross-links were compiled from multiple experiments (Supplementary Fig.

S3) and reported separately (Supplementary Table S1). The total number of cross-linked peptide

pairs was determined by counting unique lysine–lysine (K–K) residue pairs along with their

corresponding backbone sequences. Peptides with identical backbone sequences and the same K–

K pair were collapsed into a single count, regardless of charge state or peptide backbone

modifications (e.g., oxidation).

Subcellular localization analysis of proteins using SUBA5

The subcellular localization of cross-linked proteins was analyzed using SUBA5

(https://suba.live/). Accession IDs of proteins identified in both intra- and inter-protein cross-links

were submitted as input. Only the localization consensus data were used for classification. Proteins

assigned to multiple compartments were counted once in each corresponding localization category

when calculating total counts.

PANTHER protein class analysis and mapping intralinks to PDB and AlphaFold structures

using ComMap

Arabidopsis gene identifiers (ATG accession numbers) were used for protein class analysis

using PANTHER™ Protein Class tool (24), which contains 210 classification terms (version 19.0,

released 2024-06-20).

To map the results of intralinks against experimentally determined structures from the

Protein Data Bank (PDB) and predicted structures from AlphaFold protein structure database, the

protein accession IDs were first converted to UniProt IDs. The intralink information was then

mapped to available structures using ComMap, a tool that enables large-scale structure-based

mapping by integrating XL-MS data with existing structural models and performing distance

calculations (16).

Crosslinking data visualization

Cytoscape visualization: The entire cross-link-based protein-protein interaction (PPI)

network was visualized in Cytoscape (version 3.10, https://cytoscape.org/) using inter-protein

cross-link data, including heteromeric and homomultimeric cross-links. Nodes were colored

according to protein class information from the PANTHER classification system, and edges were

colored according to the STRING interaction score (https://string-db.org/).

xiVIEW: PPIs, including both inter- and intra-protein cross-links, were visualized using

xiVIEW (https://xiview.org/) (82). PPIs were categorized via the built-in STRING data loader

(score cutoff 0.4). Cross-linked peptides supporting the same lysine pair were consolidated into a

unique K-K linkage, collapsing redundancy (e.g. due to miscleavage events) and reporting each

linkage only once.

Mapping of crosslinks to high-resolution structures of proteins/complexes. Intra- and inter-

crosslinks were superimposed on PDB or AlphaFold structures using the built-in function of

xiVIEW, which also calculated the corresponding crosslink distances within the high-resolution

structures. Alignments of protein orthologs were performed and visualized using PyMOL

(https://www.pymol.org/).

Co-localization assay

Co-localization was assessed in *Nicotiana Benthamian* leaves using transient co-

expression. WSD4 was cloned into a modified pMDC43 Gateway binary vector containing an N-

terminal mCherry tag to generate 35S::mCherry-WSD4, and HTB11 was introduced to pGWB6

gateway vector to generate 35S::sGFP-HTB11. Agrobacterium-mediated infiltration was

performed, and fluorescence signals were visualized three days post-infiltration using a Leica SP8

confocal microscope.

Reference

1. Yu C & Huang L (2018) Cross-Linking Mass Spectrometry: An Emerging Technology for
Interactomics and Structural Biology. *Anal Chem* 90(1):144-165.
2. Piersimoni L, Kastritis PL, Arlt C, & Sinz A (2022) Cross-Linking Mass Spectrometry for
Investigating Protein Conformations and Protein-Protein Interactions horizontal line A
Method for All Seasons. *Chem Rev* 122(8):7500-7531.
3. Michaelis AC*, et al.* (2023) The social and structural architecture of the yeast protein
interactome. *Nature* 624(7990):192-200.
4. Xu SL, Shrestha R, Karunadasa SS, & Xie PQ (2023) Proximity Labeling in Plants. *Annu*
Rev Plant Biol 74:285-312.
5. Greenblatt JF, Alberts BM, & Krogan NJ (2024) Discovery and significance of protein-
protein interactions in health and disease. *Cell* 187(23):6501-6517.
6. Lee K & O'Reilly FJ (2023) Cross-linking mass spectrometry for mapping protein
complex topologies in situ. *Essays Biochem* 67(2):215-228.
7. Zhu X*, et al.* (2016) In planta chemical cross-linking and mass spectrometry analysis of
protein structure and interaction in Arabidopsis. *Proteomics* 16(13):1915-1927.
8. Jayachandran B, Parvin TN, Alam MM, Chanda K, & Mm B (2022) Insights on Chemical
Crosslinking Strategies for Proteins. *Molecules* 27(23).
9. Smythers AL, Iannetta AA, & Hicks LM (2021) Crosslinking mass spectrometry unveils
novel interactions and structural distinctions in the model green alga Chlamydomonas
reinhardtii. *Mol Omics* 17(6):917-928.
10. Dai S*, et al.* (2023) Capturing the hierarchically assorted modules of protein-protein
interactions in the organized nucleome. *Mol Plant* 16(5):930-961.
11. Steigenberger B, Pieters RJ, Heck AJR, & Scheltema RA (2019) PhoX: An IMAC-
Enrichable Cross-Linking Reagent. *ACS Cent Sci* 5(9):1514-1522.
12. Jiang PL*, et al.* (2022) A Membrane-Permeable and Immobilized Metal Affinity
Chromatography (IMAC) Enrichable Cross-Linking Reagent to Advance In Vivo Cross-
Linking Mass Spectrometry. *Angew Chem Int Ed Engl* 61(12):e202113937.

13.   Xu S-L, *et al.* (2017) Proteomic analysis reveals O-GlcNAc modification on proteins with
614   key regulatory functions in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* 114(8):E1536-
615   E1543.

616   14.   Chen ZL, *et al.* (2019) A high-speed search engine pLink 2 with systematic evaluation
617   for proteome-scale identification of cross-linked peptides. *Nat Commun* 10(1):3404.

618   15.   Ruwolt M, *et al.* (2023) Real-Time Library Search Increases Cross-Link Identification
619   Depth across All Levels of Sample Complexity. *Anal Chem* 95(12):5248-5255.

620   16.   Zhang W, *et al.* (2023) ComMap: a software to perform large-scale structure-based
621   mapping for cross-linking mass spectrometry. *Bioinformatics* 39(2).

622   17.   Valegard K, *et al.* (2018) Structural and functional analyses of Rubisco from arctic
623   diatom species reveal unusual posttranslational modifications. *J Biol Chem*
624   293(34):13033-13043.

625   18.   Murachelli AG, Ebert J, Basquin C, Le Hir H, & Conti E (2012) The structure of the ASAP
626   core complex reveals the existence of a Pinin-containing PSAP complex. *Nat Struct Mol*
627   *Biol* 19(4):378-386.

628   19.   Bi Y, *et al.* (2021) Arabidopsis ACINUS is O-glycosylated and regulates transcription and
629   alternative splicing of regulators of reproductive transitions. *Nat Commun* 12(1):945.

630   20.   Shrestha R, *et al.* (2025) Next-Generation Mapping of the ACINUS-Mediated Alternative
631   Splicing Machinery and Its Regulation by O-glycosylation in Arabidopsis. . *BioRxiv*.

632   21.   Yu A, *et al.* (2020) Photosynthetic Phosphoribulokinase Structures: Enzymatic
633   Mechanisms and the Redox Regulation of the Calvin-Benson-Bassham Cycle. *Plant Cell*
634   32(5):1556-1573.

635   22.   Wheat A, *et al.* (2021) Protein interaction landscapes revealed by advanced in vivo
636   cross-linking-mass spectrometry. *Proc Natl Acad Sci U S A* 118(32).

637   23.   Shannon P, *et al.* (2003) Cytoscape: a software environment for integrated models of
638   biomolecular interaction networks. *Genome Res* 13(11):2498-2504.

639   24.   Mi H, *et al.* (2021) PANTHER version 16: a revised family classification, tree-based
640   classification tool, enhancer regions and extensive API. *Nucleic Acids Res* 49(D1):D394-
641   D403.

642   25.   Szklarczyk D, *et al.* (2021) The STRING database in 2021: customizable protein-protein
643   networks, and functional characterization of user-uploaded gene/measurement sets.
644   *Nucleic Acids Res* 49(D1):D605-D612.

645   26.   Ahmed T, Yin Z, & Bhushan S (2016) Cryo-EM structure of the large subunit of the
646   spinach chloroplast ribosome. *Sci Rep* 6:35793.

647   27.   Bieri P, Leibundgut M, Saurer M, Boehringer D, & Ban N (2017) The complete structure
648   of the chloroplast 70S ribosome in complex with translation factor pY. *EMBO J*
649   36(4):475-486.

650   28.   Younkin AD, Gregory ST, & O'Connor M (2020) Alterations in the ribosomal protein bL12
651   of E. coli affecting the initiation, elongation and termination of protein synthesis.
652   *Biochimie* 175:173-180.

653   29.   Liljas A & Sanyal S (2018) The enigmatic ribosomal stalk. *Q Rev Biophys* 51:e12.

654   30.   Komenda J, Sobotka R, & Nixon PJ (2024) The biogenesis and maintenance of PSII:
655   Recent advances and current challenges. *Plant Cell* 36(10):3997-4013.

656   31.   Chen H, *et al.* (2006) A Psb27 homologue in Arabidopsis thaliana is required for efficient
657   repair of photodamaged photosystem II. *Plant Mol Biol* 61(4-5):567-575.

658   32.   Wei L, *et al.* (2010) LPA19, a Psb27 homolog in Arabidopsis thaliana, facilitates D1
659   protein precursor processing during PSII biogenesis. *J Biol Chem* 285(28):21391-21398.

660   33.   Li A, *et al.* (2024) Structural basis for an early stage of the photosystem II repair cycle in
661   Chlamydomonas reinhardtii. *Nat Commun* 15(1):5211.

662   34.   Weisz DA, *et al.* (2019) A novel chlorophyll protein complex in the repair cycle of
663   photosystem II. *Proc Natl Acad Sci U S A* 116(43):21907-21913.

35. Xingxing C*, et al.* (2018) Crystal structure of Psb27 from Arabidopsis thaliana
determined at a resolution of 1.85 A. *Photosynth Res* 136(2):139-146.
36. Sheng X*, et al.* (2019) Structural insight into light harvesting for photosystem II in green
algae. *Nat Plants* 5(12):1320-1330.
37. Martinez-Seidel F, Beine-Golovchuk O, Hsieh YC, & Kopka J (2020) Systematic Review
of Plant Ribosome Heterogeneity and Specialization. *Front Plant Sci* 11:948.
38. Chaker-Margot M (2018) Assembly of the small ribosomal subunit in yeast: mechanism
and regulation. *RNA* 24(7):881-891.
39. Lee S*, et al.* (2017) The small GTPase, nucleolar GTP-binding protein 1 (NOG1), has a
novel role in plant innate immunity. *Sci Rep* 7(1):9260.
40. Yoshida K*, et al.* (2005) Interaction of the taxilin family with the nascent polypeptide-
associated complex that is involved in the transcriptional and translational processes.
Genes Cells 10(5):465-476.
41. Faus I*, et al.* (2018) Arabidopsis ILITHYIA protein is necessary for proper chloroplast
biogenesis and root development independent of eIF2alpha phosphorylation. *J Plant*
Physiol 224-225:173-182.
42. Kim KQ*, et al.* (2024) Multiprotein bridging factor 1 is required for robust activation of the
integrated stress response on collided ribosomes. *Mol Cell* 84(23):4594-4611 e4599.
43. Tafforeau L*, et al.* (2013) The complexity of human ribosome biogenesis revealed by
systematic nucleolar screening of Pre-rRNA processing factors. *Mol Cell* 51(4):539-551.
44. Chen J*, et al.* (2024) The Ptch/SPOUT1 methyltransferase deposits an m(3)U
modification on 28S rRNA for normal ribosomal function in flies and humans. *Sci Adv*
10(50):eadr1743.
45. Ghandadi M, Dobi A, & Malhotra SV (2024) A role for RIO kinases in the crosshair of
cancer research and therapy. *Biochim Biophys Acta Rev Cancer* 1879(3):189100.
46. Messling JE*, et al.* (2022) Targeting RIOK2 ATPase activity leads to decreased protein
synthesis and cell death in acute myeloid leukemia. *Blood* 139(2):245-255.
47. Kim DS, Challa S, Jones A, & Kraus WL (2020) PARPs and ADP-ribosylation in RNA
biology: from RNA expression and processing to protein translation and proteostasis.
Genes Dev 34(5-6):302-320.
48. Udeshi ND, Mertins P, Svinkina T, & Carr SA (2013) Large-scale identification of
ubiquitination sites by mass spectrometry. *Nat Protoc* 8(10):1950-1960.
49. Van de Vosse DW, Wan Y, Wozniak RW, & Aitchison JD (2011) Role of the nuclear
envelope in genome organization and gene expression. *Wiley Interdiscip Rev Syst Biol*
Med 3(2):147-166.
50. Li F*, et al.* (2008) Identification of the wax ester synthase/acyl-coenzyme A:
diacylglycerol acyltransferase WSD1 required for stem wax ester biosynthesis in
Arabidopsis. *Plant Physiol* 148(1):97-107.
51. Patwari P*, et al.* (2019) Surface wax esters contribute to drought tolerance in
Arabidopsis. *Plant J* 98(4):727-744.
52. Vazquez A, Flammini A, Maritan A, & Vespignani A (2003) Global protein function
prediction from protein-protein interaction networks. *Nat Biotechnol* 21(6):697-700.
53. Giurgiu M*, et al.* (2019) CORUM: the comprehensive resource of mammalian protein
complexes-2019. *Nucleic Acids Res* 47(D1):D559-D563.
54. Luck K*, et al.* (2020) A reference map of the human binary protein interactome. *Nature*
580(7803):402-408.
55. Burke DF*, et al.* (2023) Towards a structurally resolved human protein interaction
network. *Nat Struct Mol Biol* 30(2):216-225.
56. Brandao MM, Dantas LL, & Silva-Filho MC (2009) AtPIN: Arabidopsis thaliana protein
interaction network. *BMC Bioinformatics* 10:454.

57. Herrera-Ubaldo H*, et al.* (2023) The protein-protein interaction landscape of transcription
factors during gynoecium development in Arabidopsis. *Mol Plant* 16(1):260-278.
58. Yang X*, et al.* (2020) PlaPPISite: a comprehensive resource for plant protein-protein
interaction sites. *BMC Plant Biol* 20(1):61.
59. Balch S*, et al.* (2025) Cryo-EM structure of human telomerase dimer reveals H/ACA
RNP-mediated dimerization. *Science* 389(6756):eadr5817.
60. Dang DT (2022) Molecular Approaches to Protein Dimerization: Opportunities for
Supramolecular Chemistry. *Front Chem* 10:829312.
61. Yang Z*, et al.* (2018) EBS is a bivalent histone reader that regulates floral phase
transition in Arabidopsis. *Nat Genet* 50(9):1247-1253.
62. Shu J*, et al.* (2021) Genome-wide occupancy of Arabidopsis SWI/SNF chromatin
remodeler SPLAYED provides insights into its interplay with its close homolog BRAHMA
and Polycomb proteins. *Plant J* 106(1):200-213.
63. Lim MH*, et al.* (2004) A new Arabidopsis gene, FLK, encodes an RNA binding protein
with K homology motifs and regulates flowering time via FLOWERING LOCUS C. *Plant
Cell* 16(3):731-740.
64. Yan Z, Shi H, Liu Y, Jing M, & Han Y (2020) KHZ1 and KHZ2, novel members of the
autonomous pathway, repress the splicing efficiency of FLC pre-mRNA in Arabidopsis. *J
Exp Bot* 71(4):1375-1386.
65. Shrestha R*, et al.* (2022) 15N Metabolic Labeling Quantification Workflow in Arabidopsis
Using Protein Prospector. *Front. Plant Sci.* 13:832562.
66. Reyes AV, Shrestha R, Baker PR, Chalkley RJ, & Xu S-L (2022) Application of Parallel
Reaction Monitoring in 15N Labeled Samples for Quantification. *Front. Plant Sci.*
13:832585.
67. Matsuo Y*, et al.* (2017) Ubiquitination of stalled ribosome triggers ribosome-associated
quality control. *Nat Commun* 8(1):159.
68. Simsek D*, et al.* (2017) The Mammalian Ribo-interactome Reveals Ribosome Functional
Diversity and Heterogeneity. *Cell* 169(6):1051-1065 e1018.
69. Peng Y, Markov Y, Goncearenco A, Landsman D, & Panchenko AR (2021) Human
Histone Interaction Networks: An Old Concept, New Trends. *J Mol Biol* 433(6):166684.
70. Zhang S, Yu Q, Li Z, Zhao Y, & Sun Y (2024) Protein neddylation and its role in health
and diseases. *Signal Transduct Target Ther* 9(1):85.
71. Alemasova EE & Lavrik OI (2019) Poly(ADP-ribosyl)ation by PARP1: reaction
mechanism and regulatory proteins. *Nucleic Acids Res* 47(8):3811-3827.
72. Messner S*, et al.* (2010) PARP1 ADP-ribosylates lysine residues of the core histone
tails. *Nucleic Acids Res* 38(19):6350-6362.
73. Nosella ML*, et al.* (2024) Poly(ADP-ribosyl)ation enhances nucleosome dynamics and
organizes DNA damage repair components within biomolecular condensates. *Mol Cell*
84(3):429-446 e417.
74. Su Y*, et al.* (2017) Phosphorylation of Histone H2A at Serine 95: A Plant-Specific Mark
Involved in Flowering Time Regulation and H2A.Z Deposition. *Plant Cell* 29(9):2197-
2213.
75. Sabari BR, Zhang D, Allis CD, & Zhao Y (2017) Metabolic regulation of gene expression
through histone acylations. *Nat Rev Mol Cell Biol* 18(2):90-101.
76. Goudarzi A (2019) The recent insights into the function of ACAT1: A possible anti-cancer
therapeutic target. *Life Sci* 232:116592.
77. Liu Y*, et al.* (2022) Mechanisms and inhibition of Porcupine-mediated Wnt acylation.
Nature 607(7920):816-822.
78. Nile AH & Hannoush RN (2016) Fatty acylation of Wnt proteins. *Nat Chem Biol* 12(2):60-
69.

79. Xue B & Rhee SY (2023) Status of genome function annotation in model organisms and
crops. *Plant Direct* 7(7):e499.
80. Casazza AP, Tarantino D, & Soave C (2001) Preparation and functional characterization
of thylakoids from Arabidopsis thaliana. *Photosynth Res* 68(2):175-180.
81. Shao Z*, et al.* (2024) Nuclear pyruvate dehydrogenase complex regulates histone
acetylation and transcriptional regulation in the ethylene response. *Sci Adv*
10(30):eado2825.
82. Combe CW, Graham M, Kolbowski L, Fischer L, & Rappsilber J (2024) xiVIEW:
Visualisation of Crosslinking Mass Spectrometry Data. *J Mol Biol* 436(17):168656.
773

Figure 1. Efficient cross-linking workflow for complex protein samples identifies numerous cross-links in Arabidopsis.

(**A**) Schematic of the workflow using PhoX or tBu-PhoX to covalently cross-link spatially proximal lysine residues in Arabidopsis subcellular fractions (lysate, chloroplast, nucleus), followed by digestion, IMAC enrichment, high pH reversed-phase HPLC fractionation, and LC-MS/MS. Data were analyzed using pLINK2, Cytoscape, and xiVIEW.

(**B**) Summary of unique intra- and inter-protein cross-links from each subcellular fraction. Intra-links connect non-overlapping peptides within a protein; inter-links include heteromeric and homomultimeric cross-links.

(**C**) Subcellular localization of cross-linked proteins based on SUBA5, with redundancy of proteins assigned to multiple compartments.

Figure 2. Validation of intra- and inter-protein cross-links by structural mapping.
(**A**-**B**) Histogram of Euclidean Cα-Cα distances of intra-protein cross-links mapped to
Arabidopsis structures from PDB (A) or AlphaFold (B), showing 94% and 89% within the 35Å
constraint, respectively.
(**C**-**E**) Cross-links mapped to the Arabidopsis Rubisco L_8S_8 complex modeled from the L_2S_2
structure (PDB:5IU0). RBCL is shown in cyan and RBCS is shown in pink. Self(intra),
heteromeric, and homomultimeric links are shown in purple, green, and orange, respectively.
The circular view (C) and structural mapping (D-E) show that 74.56% of the cross-links are
within 35Å.

(**F**) Cross-links between SR45 and ACINUS or PININ occur at known interaction interfaces involving the RRM domain of SR45 and the RSB domain of ACINUS and PININ.
(**G-H**) Intra- and inter-protein cross-links between PRK and CP12.1 mapped onto the GAPDH-CP12-PRK complex (PDB:6KEZ), aligning with the known interaction interface.

Figure 3. XL-MS identifies direct protein-protein interactions and reveals known and novel complexes.

(**A**) XL-MS identified an Arabidopsis interactome comprising 1,446 PPIs, including 1,390 heteromeric and 56 homomultimeric interactions among 1,229 proteins. Nodes represent proteins, colored by PANTHER protein class; edges represent interactions, colored by the combined STRING confidence score. Of the total PPIs, 637 PPIs have a combined confidence score of at least 0.4, including 532 with a score of at least 0.7.

(**B**) Summary of the STRING score, showing the contribution from eight evidence sources. XL-MS provides experimental support for 71% of the heteromeric PPIs that either have low confidence scores (<0.4) or lack prior experimental evidence.

(**C-D**) Examples of specific PPIs detected by XL-MS that complement low confidence or missing experimental STRING scores: the ORP3B-PLDRP1 interaction (cross-linked at ORP3B-K7/PLDRP1-K193) and the NRPA2-uL1c interaction (NRPA2-K394/uL1c-K127).

Figure 4. XL-MS resolves the topology of chloroplast 70S ribosome stalk regions inaccessible by cryo-EM.

(**A**) XL-MS identifies flexible stalk proteins (bL12c, uL1c, uL10c, uL11c), with cross-links color-coded by STRING confidence.

(**B**) Domain comparison between Arabidopsis bL12c and *E. coli* bL12. Homodimer interfaces are found at the C-terminus of bL12c.

(**C**) Cross-links mapped onto the predicted bL12c structure (AF-P36212-F1-v4) showing lysines K64, K137, K148, and K151 interacting with ribosomal subunits and the elongation factor CPEF-G. Nodes are color-coded according to annotation category.

Figure 5. XL-MS reveals dynamic connectivity and unique features in photosystem complexes.

(**A**) PSII topology showing cross-links color-coded by STRING confidence. Notable PSB27 cross-links (K93, K168) highlight interactions with PSII core and OEC components.

(**B**) XL-MS uncovers extensive interactions among PSI, PSII, LHCI, LHCII, ATPase, FtsH/SPPA proteases, and transient proteins involved in state transitions, assembly, repair, and chlorophyll metabolism.

A Translation initiation and elongation — Protein folding/QC/Gatekeeper — Biogenesis related — PTM

B

Interactor	Lys(K)	Ribosome	Lys(K)
MEE49	22	uL29z	35
RNC1	85	uL4z	289
AT5G19300	3	eS1z	19
NOG1	532	eL31z	58
UBP14	154	eS31z	48
UTP7	453	eS30z	3
UTP7	451	eS30z	20
RGGA	255	eS31γ	119
RGGA	266	uS19u	65
RGGB	245	eS31γ	119
RGGB	256	uS19u	65
RGGC	42	eL8z	250
NUC-L1	79	uL23y	32
NUC-L2	228	eL8γ	116
EIF2	429	eL24z	130
AT1G30455	169	eL24y	155
EIF3A	568	uL24z	41
EIF3A	568	uL24z	130
MBF1A	19	eS30z	53
NACα-like 5	92	uL29y	43
NACβ	18	eL31z	58
NACβ	18	eL31z	61
NACα-like 4	95	uL29y	43
TAXILIN	339	eL31γ	38
SRP72	579	uL29y	77
ILA	793	eS8z	152
ILA	793	uS19u	65
AT1G06690	85	uL23z	32
AT1G06690	85	eL8z	227
THIOREDOXIN	229	eL38z	29
PDI 2-1	73	uL18z	279
HIP2	161	uL18z	240
PAD1	243	eL24y	155
RPN8A	40	eS1z	27
EBF1	365	uL15x	118
AT4G02760	7	uL18z	254
MPK11	38	uL4z	289
AT1G07550	77	eL27z	47
RIOK2	13	eS30z	3
PARP1	61	eL43y	7
AT1G29350	5	uS15z	43
AT3G07760	16	eS31z	6
H1.2	217	uL23z	24

C

Detected	Description/Function	Human homolog	Ribosome related function (based on human/yeast homolog)	Ribosome related function in plant
AT5G19300	methytransferase C9orf114	SPOUT1	Ribosome synthesis trans-Acting factor	uncharacterized
PARP1	ADP-ribosylation	PARPs (17)	Polysome formation	uncharacterized
RIOK2	Kinase/ATPase	RIOKs(3)	40S maturation	uncharacterized
MPK11	Kinase	MPKs	uncharacterized	uncharacterized
UBP14	De-ubiquitinase	UBPs	Maintain Ribosome assembly	uncharacterized
EBF11	F-box E3 ligase	FBXL20	uncharacterized	uncharacterized
AT4G02760	F box E3 ligase with LRR	FBXL2-like	uncharacterized	uncharacterized

Figure 6. XL-MS of ribosome complexes reveals conserved ribosome-associated factors.
(**A**) XL-MS identified extensive interactions within ribosomal subunits as well as with proteins involved in ribosome biogenesis, translation, folding, quality control, stress response, and other regulatory pathways. Ribosomal proteins are overlaid with ubiquitination data; red nodes indicate ubiquitinated proteins.
(**B**) Table showing covalently cross-linked lysine residues between ribosomal proteins, reflecting spatial proximity. Protein labels are color-coded according to the functional categories shown in (A).
(**C**) Table of XL-MS-identified ribosome-associated interactors with potential post-transcriptional or post-translational regulatory functions, many of which remain uncharacterized in plants.

Figure 7. Histone interactome reveals conserved and novel chromatin-associated interactions.

(**A**) XL-MS reveals extensive interactions between histones and both known and previously uncharacterized proteins, including components of chromatin-associated complexes. Enzymatic regulators are highlighted in red.

(**B**) Visualization of the co-localization of GFP-HTB11 and mCherry-WSD4 in *N.benthamiana.* Green: GFP-HTB11; red: mCherry-WSD4. Overlapping signals are indicated by arrows, with a zoom-in view highlighted in the inset. Scale bar=10μm.

(**C**) XL-MS identifies interaction between histones and PARP1, NEDD8 and WSD4, suggesting their potential roles in histone modifications and polynucleosome organization. Multiple cross-links between histone H2B proteins and WSD4, an O-acyltransferase, support a possible function for WSD4 in histone acylation.

Supplementary Figure S1: Optimization of the PhoX cross-linker concentration for XL-MS.
(**A-D**) Different concentrations of PhoX cross-linker (0 to 10 mM) were tested on cell lysates from light (A, C, D) or dark grown seedlings (B). Increased protein cross-linking, indicated by the accumulation of higher molecular weight proteins (red arrow), was observed with increasing cross-linker concentration. However, no additional visible cross-linking was detected at higher concentrations (7-10 mM).

Supplementary Figure S2: **Standard XL acquisition outperforms Real-Time Library Search (RTLS) acquisition.**

Data were acquired on an Orbitrap Eclipse mass spectrometer using PhoX-enriched cross-linked peptides from Arabidopsis cell lysate samples, then searched against the entire Arabidopsis protein database. RTLS acquisition significantly reduced the identification of monolinks, where one NHS ester group is cross-linked to a peptide while the other is hydrolyzed, by approximately 50%, and loop links, which are cross-links within a single peptide, by about 31%. However, RTLS also substantially decreased the detection of inter-peptide cross-links, including both intraprotein (intralinks) and interprotein (interlinks) cross-links, by 43% and 40%, respectively.

Supplementary Fig. S3: Summary of XL-MS experiments.
Key details of sample preparation, data acquisition using the Orbitrap Eclipse mass spectrometer, and database searching using pLINK2 are provided. A summary of the unique cross-linked peptide residue pairs identified across experiments is also included.

Supplementary Fig. S4. Representative cross-links supporting the high confidence of XL-MS analysis.

(**A**) Multiple cross-links were identified between the ribosomal proteins uL15x and eL13z, including peptide pairs with different degrees of miscleavage cross-linked at identical sites (e.g., uL15x-K92 to eL13z-K166/K172).

(**B**) Cross-links map conserved regions, with the ATPase subunit VHA-E3 (K49) cross-linked to both VHA-B1 (K437) and VHA-B2 (K438).

(**C**) Similar cross-linking patterns were observed between homologous interactions, as shown by DNA topoisomerase 1 alpha (TOP1A-K672) cross-linked to two dehydrin proteins, ERD14-K38 and COR47-K46.

Supplementary Figure S5. Mapping of Rubisco cross-links on a high-resolution orthogonal structure of hornwort Rubisco.
Cross-links are color-coded by type: purple for self (intra)links, green for heteromeric (inter)links, and orange for homomultimeric links.
(**A**) Cross-links were mapped onto the high-resolution structure of L8S8 heterotetrameric Rubisco from hornwort (PDB: 9CHZ). RBCL subunits are shown in cyan and RBCS subunits in pink.
(**B**) Distance distribution of cross-links mapped onto Rubisco 9CHZ, showing that 71.6% of the identified cross-links fall within the expected spatial constraint of 35 Å.

Supplementary Figure S6. High-confidence identifications of cross-linked peptides between ACINUS–SR45 and PININ–SR45.

b-ions are shown in green and y-ions in yellow. A series of fragment ions, including those from α- and β-peptides, as well as cross-linker-containing ions, were detected, supporting the reliability of the cross-link identifications.

(**A**) Spectrum of the cross-linked peptide between ACINUS-K633 and SR45-K154.

(**B**) Spectrum of the cross-linked peptide between PININ-K274 and SR45-K154.

Supplementary Fig. S7. Topological mapping of multiple protein complexes by XL-MS.
Proteins are shown as rectangles scaled to their respective lengths. Cross-links are color-coded based on STRING score confidence thresholds, highlighting both known and novel interactions.

(**A**) Mapping of 26S proteasome subunits, highlighting interaction interfaces between RPN7 and DSS1(I) and DSS1(V). DSS1 proteins are particularly challenging to map in cryo-EM due to their small size, intrinsic disorder, and dynamic interaction properties.

(**B**) Mapping of photosystem I (PSI) showing connectivity between different subunits.

(**C-E**) Mapping of different ATPase complexes, including chloroplast, V-type, and mitochondrial ATPases.

(**F**) Mapping of the prefoldin complex.

(**G**) Mapping of the phosphoenolpyruvate carboxylase (PEPC) complex.

Supplementary Figure S8. Mapping of cross-links in the 70S ribosome complex onto high-resolution structures of spinach 70S ribosomes (PDB: 5MMM and 5X8P).
Cross-links are color-coded by type: purple for intralinks, green for heteromeric inter-protein cross-links, and orange for homomultimeric inter-protein cross-links.
(**A**) Structural mapping of 70S ribosome cross-links on the spinach cryo-EM structure (PDB: 5MMM), which lacks bL12 and other flexible proteins in the stalk region.
(**B**) Distance distribution of the cross-links mapped in (**A**). Of the 424 total cross-links detected, 276 were mapped (263 self-links and 13 heteromeric), with 241 falling within the 35 Å spatial constraint.
(**C**) Structural mapping of 70S ribosome cross-links onto an alternative spinach cryo-EM structure (PDB: 5X8P), which also lacks most of the flexible proteins in the stalk region.
(**D**) Distance distribution of the cross-links mapped in (**C**). Of the 424 cross-links detected, 234 were mapped (224 self-links and 10 heteromeric), with 206 falling within the 35 Å constraint.

Supplementary Figure S9. Mapping of PSB27 cross-links onto a high-resolution structure of 6KAC from Chlamydomonas.

(**A**) Superimposed AlphaFold structures of PSB27 homologs reveal unique features of Arabidopsis PSB27. While it contains a conserved C-terminal four-helix bundle, Arabidopsis PSB27 lacks a lipid anchor and shows divergence in sequence and surface charge.

(**B**) Cross-links between Arabidopsis PSB27 and PSII core/oxygen-evolving complex (OEC) subunits mapped onto the Chlamydomonas PSII structure (PDB: 6KAC), which captures residues 91-114 of PSB27. The distance constraints suggest a more dynamic interaction between PSB27 and the PSII complex.

Supplementary Figure S10. Mapping of PSII complex cross-links onto several available high-resolution structures. A total of 581 cross-links were mapped, including 504 self-links

and 77 heteromeric links. Cross-links are colored according to the type of cross-link (purple for self (intra) link; green for heteromeric (inter) link; orange for homomultimeric link). The distribution of cross-link distances from the integrated cross-link model mapping is shown alongside the structural mapping.

(**A-B**) Structure mapping of PSII cross-links onto the Chlamydomonas PSII-LHCII supercomplex (PDB: 6KAC), which includes the PSII reaction center, PsbO, and a short sequence of Psb27.

(**C-D**) Structure mapping of PSII cross-links onto the Arabidopsis PSII-LHCII supercomplex, (PDB: 5MDX) which represents mature PSII, containing the PSII reaction center and PsbO but lacking Psb27.

(**E-F**) Cross-links mapped onto the cyanobacterial PSII assembly intermediate structure (PDB: 7NHP), which includes the PSII reaction center together with Psb27 and Psb28. Cyanobacteria have a single copy of the PSB27 gene. Fewer cross-links were mapped onto this assembly intermediate.

(**G-H**) Cross-links mapped onto the *Chlamydomonas* PSII assembly intermediate structure (PDB: 8R2I), representing a native assembly intermediate containing the PSII reaction center and Psb27. Similarly, fewer cross-links were mapped to this structure.

Supplementary Figure S11. Topology of the 80S ribosomal complex.
(**A**) 2D topological arrangement of the 40S and 60S ribosomal subunits based on their connectivity. Links between proteins are colored according to the confidence thresholds of the STRING score.
(**B**) Summary of cross-links and protein-protein interaction (PPI) information from (A).
(**C**) 80S ribosome cross-links mapped onto the high-resolution structure of the tobacco 80S ribosome (PDB: 8B2L). The large ribosomal subunit is colored orange and the small ribosomal subunit is colored red. Links are color-coded according to link type: purple for self-links (intralinks), green for heteromeric links (interlinks), and orange for homomultimeric links.
(**D**) Distribution of cross-link distances from the integrated cross-link model in (C).

Supplementary Fig. S12. XL-MS identifies WSD4 as a novel histone interactor that associates with H2B family members.

(**A**) Cross-links were detected between WSD4 (an O-acyltransferase) and several H2B family members, including HTB1, HTB9, HTB3, and HTB11.

(**B**-**E**) High-confidence cross-link spectra reveal interactions between WSD4 and H2B family members. Series of fragment ions-including those from α- and β-peptides, as well as cross-linker-containing ions-were detected, supporting the reliability of the cross-link identifications.