Mapping Architecture of Protein complexes in Arabidopsis using XL-MS

Cao Son Trinh [1,2]*, Ruben Shrestha[1]*, William C Conner[1], Andres V Reyes[1,2], Sumudu S

Karunadasa[1,2], Grace Liu[1], Ken Hu[1], Shou-Ling Xu[1,2]

[1]Department of Plant Biology, Carnegie Institution for Science, Stanford, California, USA;

[2]Carnegie Mass Spectrometry Facility, Carnegie Institution for Science, Stanford, California,

USA

*These authors contributed equally.

Corresponding author: Shou-Ling Xu, slxu@stanford.edu

Abstract

Capturing molecular machines in action is essential for understanding protein complex

architecture, cellular regulation, and gene function. Here, we establish a cross-linking mass

spectrometry (XL-MS) platform using the PhoX cross-linker for proteome-wide interaction

mapping in Arabidopsis under semi-native conditions. From cell lysates, chloroplasts, and

nuclei, we identified 47,119 unique cross-links, including 3,527 inter-protein cross-links, and

constructed a high-confidence protein-protein interaction (PPI) network containing 1,229

proteins and 1,446 PPIs. In silico analysis (STRING) confirmed 637 heteromeric interactions,

with the remainder representing novel interactions. Our dataset provides experimental evidence

for direct interactions and defines binding interfaces. XL-MS resolved the topology of

challenging proteins and revealed both conserved and unique interactions, as exemplified in

photosystem complexes, ribosomes, and histone interactomes. By characterizing the histone

machinery, we identified a potential novel histone modifying enzyme. This work advances

protein annotation and provides new insights into cellular organization and function.

Introduction

Understanding how molecular machines operate within cells - and capturing them in action - is essential to unraveling the principles of life and gene function. A powerful tool for this purpose is cross-linking mass spectrometry (XL-MS), an emerging high-throughput approach that enables the capture of protein complexes in their native context across diverse organisms (1, 2). Unlike affinity purification mass spectrometry (AP-MS) and proximity labeling mass spectrometry (PL-MS) (3-5), XL-MS does not require antibodies or genetically tagged transgenes. Instead, it relies on chemical cross-linkers that covalently link spatially proximal residues within or between proteins in solution, followed by protease digestion and mass spectrometric identification of the resulting cross-linked peptides. In addition to generating a "parts list" of interacting proteins, XL-MS provides direct evidence of physical interactions and precisely maps contact sites at the amino acid level, providing unparalleled structural insight into macromolecular assemblies.

Despite its promise, large-scale proteome-wide XL-MS - such as in whole-cell lysates, isolated organelles, or *in vivo* systems - remains challenging due to the inherent low abundance and heterogeneity of cross-linked peptides and the complexity of data analysis (1, 6). In plants, additional barriers further complicate its application, including rigid cell walls and large central vacuoles that dilute cross-linkers, as well as secondary metabolites that can chemically interfere with cross-linking chemistry (7, 8). Recent pioneering studies in Arabidopsis, algae, and soybean have demonstrated the potential of XL-MS in plant systems (7, 9, 10), but its overall implementation remains limited.

To address these challenges, we developed a cross-linking strategy optimized for proteome-wide mapping in complex Arabidopsis samples using the enrichable cross-linker

PhoX. PhoX is a phosphonic acid-based reagent that, like BS3 and DSSO, contains NHS esters

and primarily targets lysine residues (11). Its phosphonic acid handle allows efficient enrichment

by immobilized metal affinity chromatography (IMAC). With a maximum range of 35Å -

consisting of a 20 Å Cα-Cα distance from its rigid structure and an additional 15 Å to account for

protein flexibility - PhoX is well suited for high confidence, large-scale structural proteomics

(12). To reduce sample complexity and improve cross-linking efficiency, we used subcellular

fractionations. To further improve the detection of the low abundance cross-linked peptides, we

performed extensive high-pH reversed-phase HPLC after IMAC enrichment. Together, these

measures allowed for comprehensive proteome-wide XL-MS mapping.

From cell lysates, chloroplasts, and nuclei, we identified a total of 47,119 unique cross-

linked peptide pairs, including 43,592 intra-protein (self-link) and 3,527 inter-protein cross-links.

These data allowed the construction of a PPI network comprising 1,229 proteins and 1,446

interactions, including 1,390 heteromeric and 56 homomultimeric interactions. STRING analysis

validated 637 heteromeric PPIs (combined confidence score ≥ 0.4), while the remainder

represent previously uncharacterized interactions. Our dataset provides experimental evidence

for direct physical binding and defines protein binding interfaces at amino acid resolution for

both known and novel interactions.

Beyond interaction mapping, our XL-MS data provide structural and functional insights

into key cellular processes. By resolving flexible or transient components within the 70S

chloroplast ribosome and photosystem complexes, our results complement existing cryo-EM

structures. XL-MS also enabled a comprehensive mapping of the photosystem, 80S ribosome,

and histone interactomes within heterogeneous protein populations, revealing structural biology

and capturing both known and novel interactions. In particular, we identified a metabolic O-

acyltransferase that interacts with histones, suggesting a potential new mechanism of chromatin

regulation via post-translational modification. Together, these results improve protein annotation,

shed light on protein complex architectures, and provide as a valuable resource for the broader

research community - laying the foundation for future discoveries.

Results

Efficient cross-linking workflow for complex protein samples identifies numerous cross-

links in Arabidopsis

To gain structural insights and visualize molecular machines in their native cellular context

within Arabidopsis cells, we performed cross-linking mass spectrometry (XL-MS) (see Methods)

on fractionated seedling samples, including cell lysates and chloroplasts from light-grown

seedlings and nuclei from dark-grown seedlings. Using the PhoX cross-linker (Fig. 1A), we

established a robust pipeline optimized for proteome-wide detection of low-abundance cross-

linked peptides in complex Arabidopsis samples.

Optimization of the PhoX concentration showed a shift to higher molecular weights at 2-

5 mM (red arrows, Fig. S1A-D), indicating successful cross-linking, with no additional

improvement at higher concentrations. Based on these results, we used 2 and 5 mM PhoX for

subsequent experiments (see Methods). After cross-linking, we precipitated the proteins with

phenol, digested them with trypsin, and enriched the cross-linked peptides using $GaCl_3$-based

IMAC (13). We then fractionated the enriched peptides by high-pH reversed-phase

chromatography, analyzed them by LC-MS/MS, and identified cross-links using pLINK2 (14).

High-resolution, high-accuracy data for both MS1 (precursor ions) and MS2 (fragment ions)

were acquired on an Orbitrap Eclipse mass spectrometer to ensure reliable cross-link

identification.

To optimize data acquisition, we compared the standard XL workflow with real-time

spectral library search (RTLS) (15), both targeting highly charged precursors (+3 and above).

RTLS uses a survey MS2 scan to reduce mono-link detection. In cell lysates, RTLS reduced

mono- and loop-link identifications by ~50% and ~31%, respectively (Fig. S2), but also

significantly reduced intra-link (43%) and inter-link (40%) identifications. Because of this trade-

off, we used the standard XL workflow to analyze cross-linked chloroplast and nuclear samples.

In total, we identified 47,119 unique cross-linked peptide pairs (intra- and inter-links)

representing 5,759 proteins. Specifically, we detected 25,795, 17,090, and 19,612 cross-links

from the nucleus, chloroplast, and cell lysate, respectively (Fig. 1B and. S3, Table S1).

Subcellular localization analysis using SUBA5 revealed that the cross-linked proteins were

distributed in various cellular compartments, including the nucleus, plastid, cytosol,

mitochondria and other organelles (Fig. 1C, Table S2).

To assess the quality of cross-linked peptide identification, we examined the identified

cross-links (Table S1) and found consistent peptide pairs mapping to the same sites or regions.

For example, in ribosomal subunits, we detected site-specific cross-links with variable

miscleavages, such as those between uL15x-K92 and eL13z-K166/K172 (Fig. S4A). Conserved

region cross-links were also observed, including VHA-E3-K49 with both VHA-B1-K437 and

VHA-B2-K438 (Fig. S4B), and TOP1A-K672 with ERD14-K38 and COR47-K46 (Fig. S4C).

These independent and consistent observations underscore the robustness and reliability of our

results.

Validation of intra- and inter-protein cross-links by structural mapping

To evaluate the validity of the cross-links, we first used ComMap (16) to map intra-links to high-

resolution protein structures available in the Protein Data Bank (PDB) and predicted structures

by AlphaFold. Specifically, 721 Arabidopsis proteins had high-resolution structures in the PDB,

and 3,975 were available through AlphaFold (Table S1). A total of 21,981 cross-links were

successfully mapped, with 94% of the PDB-mapped cross-links and 89% of the AlphaFold-

mapped cross-links satisfying the 35 Å distance constraint (Fig. 2A-B).

To evaluate both intra- and inter-protein cross-links, we mapped 174 cross-links onto the

L_8S_8 Rubisco assembly modeled from the Arabidopsis L_2S_2 crystal structure (PDB:5IU0) (17).

Of these, 169 cross-links were mapped to the large (479 aa) and small (181 aa) subunits, while

five were localized to the N-terminal residues (1–11) of RBCL, which are not present in the PDB

structure (Fig. 2C). The majority (126, 74.6%) of mapped cross-links fell within the expected

Cα-Cα distance of ≤35 Å; the remaining 43 (25.4%) exceeded this range, suggesting regions of

structural flexibility (Fig. 2D–E). Similar mapping results were obtained using the hornwort

Rubisco cryo-EM structure, which yielded similar results (Fig. S5A-B), further supporting the

validity of our cross-linking data.

To assess whether cross-linked lysine residues are positioned within predicted interaction

interfaces, we examined several PPIs using available structural data. The cross-link between

ACINUS-K633 and SR45-K154 maps near the RSB domain of ACINUS (aa 599-631) and

within the RRM domain of SR45 in the ASAP core complex (PDB:4A8X) (18-20) (Fig. 2F, Fig.

S6A). Similarly, the cross-link between SR45-K154 and PININ-K274 aligns with the same RRM

domain and the predicted RSB domain of PININ (Fig. 2F, Fig. S6B). For PRK-K217 and

CP12.1-K65, the cross-link localizes to the interaction interface in the GAPDH-CP12-PRK

complex (PDB:6KEZ) (21) with a Cα-Cα distance of 13.3 Å (Fig. 2G-H). These results confirm

that the cross-linked residues are located at or within a few amino acids of the interaction

interfaces, consistent with previous observations (22).

XL-MS systematically identifies direct protein-protein interactions and reveals known and novel complexes

We imported 3,527 inter-protein cross-links into Cytoscape (23), revealing 1,446 PPIs among 1,229 proteins, including 1,390 heteromeric and 56 homomultimeric interactions (Fig. 3A, Table S1). Proteins were classified using the PANTHER protein class tool, and nodes were colored accordingly (24). Many interacting proteins shared the same classification, supporting the reliability of our PPI map. Notably, 457 of the 1,210 proteins involved in heteromeric interactions lacked PANTHER annotations, contributing to 819 interactions - 369 between unclassified proteins and 450 between classified and unclassified proteins.

We also incorporated STRING confidence scores (25) into the network and color-coded the edges accordingly (Fig. 3A). These scores combine evidence from eight sources (Fig.3B) (25). For heteromeric PPIs, 38.2% of PPIs had high STRING confidence: 388 scored $\geq$ 0.9 and 144 scored 0.7-0.9. The remainder included 105 medium (0.4-0.7), 144 low (<0.4), and 609 with no STRING score (Fig.3B). The PANTHER and STRING results indicate that both known and novel PPIs were captured, highlighting the power of systematic XL-MS to reveal previously uncharacterized protein complexes and pathways.

Even among PPIs with good STRING combined scores, many lack experimental support for direct or indirect interactions in Arabidopsis or homologs in other species. Among the heteromeric PPIs identified, only 28.9% (n = 402) have experimental support (score $\geq$ 0.4), while 71.1% have low or no scores (Fig. 3B). Our XL-MS data provide evidence for their direct interactions. For example, ORP3B (AT3G09300, oxysterol-binding protein) and PLDRP1 (AT5G39570, PLD-regulated protein 1) have a combined score of 0.602 but a low experimental score (0.055); we identified a cross-link between ORP3B-K7 and PLDRP1-K193 (Fig. 3C),

suggesting a common role in lipid trafficking and signaling. Similarly, NRPA2 (AT1G29940,

RNA polymerase predicted to localize to both the nucleus and chloroplasts) and uL1c

(AT3G63490, chloroplast 50S ribosomal protein) have a combined score of 0.588 but no

experimental support; a cross-link between NRPA2-K394 and uL1c-K127 suggests a direct

interaction and potential coupling of chloroplast transcription and translation (Fig. 3D).

XL-MS reveals unique topology of chloroplast 70S ribosome stalk regions

Using XL-MS, we mapped the topologies of several protein complexes (Fig. S7A-G),

including the flexible stalk regions of the chloroplast 70S ribosome (Fig. 4A-C), which have

remained difficult to resolve due to the dynamic nature of proteins such as uL1c, uL10c, uL11c,

and bL12c (26, 27). In total, we identified 424 cross-links across 34 ribosomal subunits of the

70S ribosome, including 352 intraprotein, 64 heteromeric, and 8 homomultimeric cross-links.

Notably, many intra- and interprotein cross-links involved these flexible stalk proteins. Although

cross-links were mapped to spinach 70S ribosome structures (PDB: 5MMM and 5X8P; Fig.

S8A-B), most cross-links involving uL1c, uL10c, uL11c, and bL12c could not be precisely

assigned to specific locations within the stalk regions because these proteins are either missing or

only partially resolved in the cryo-EM structures.

Arabidopsis bL12c has a plant-specific disordered N-terminus, followed by conserved

structured N-terminal (NTD) and C-terminal (CTD) domains connected by a flexible hinge (Fig.

4B). Essential for translation, bL12c uses hinge-mediated mobility to recruit specific translation

factors. XL-MS identified interactions between bL12c (the two highly homologous proteins

AT3G27850/bL12cx and AT3G27830/bL12cz) and several ribosomal core proteins, as well as

the elongation factor CPEF-G, mainly through its structured CTD (Fig. 4C). Although bL12c

dimers are typically mediated by the NTD (28), XL-MS revealed unexpected homodimer cross-

links within the CTD - the same region involved in translation factor binding (Fig. 4C). This

suggests that CTD dimerization may serve as an interface for the recruitment of interaction

partners. Taken together, these results highlight both conserved and plant-specific features of

bL12c and demonstrate the power of XL-MS in resolving dynamic aspects of ribosomal

architecture.

XL-MS reveals dynamic connectivity and unique features in photosystem complexes

Photosystems I and II (PSI and PSII) and their associated light-harvesting complexes

(LHCI and LHCII) are essential for light energy harvesting and carbon fixation, but they require

continuous repair, especially under high and fluctuating light conditions (29-31). Although

progress has been made in understanding certain steps of the PSII repair (32, 33), many aspects,

including the detailed mechanism, identification of repair intermediates, and specific

contributing factors, remain unresolved.

Using XL-MS, we identified extensive interactions involving the PSII repair factor Psb27

(AT1G03600) (Fig. 5A). Unlike its cyanobacterial counterpart, which functions in both assembly

and repair, Arabidopsis Psb27 appears to be specialized for repair. In contrast, the less

homologous Psb27-H1(AT1G05385) is involved in assembly. Notably, Arabidopsis Psb27 lacks

a lipid anchor and is structurally more flexible than the cyanobacterial version (34) (Fig. S9A). A

partial structure from Chlamydomonas (PDB: 6KAC) captured residues 91-114 of Psb27 (35).

Our XL-MS data revealed that Psb27 interacts with both the PSII reaction center and the oxygen-

evolving complex (OEC), involving K93 (as in 6KAC) and a second site at K168 (Fig. 5A, Fig.

S9B, S10A-H). These distance constraints (Fig. S9B) suggest that Psb27 engages PSII in a more

dynamic and flexible manner in plants.

XL-MS also revealed extensive interactions between PSI, PSII, LHCI, LHCII, ATP

synthase, ATP-dependent proteases (FtsH, SPPA) and transient proteins involved in state

transitions, complex assembly and repair (Fig. 5B). Notably, chlorophyll biosynthesis and

catabolic proteins (e.g., ACD1, CHLM, FLU) were found in close association with LHCII,

suggesting coordinated chlorophyll delivery and reactive oxygen species (ROS) control during

LHCII assembly. In addition, many assembly and repair factors were closely associated with

LHCII, indicating their readiness for rapid engagement during dynamic photosystem remodeling.

XL-MS of ribosome complexes reveals conserved ribosome-associated factors

We identified extensive interactions involving the 80S ribosome, a complex of 60S and

40S subunits that synthesizes nuclear-encoded proteins. Ribosomes assemble in the nucleolus,

localize to the cytoplasm, and associate with the endoplasmic reticulum (ER). As a dynamic

macromolecular machine, the ribosome coordinates translation with numerous proteins and

RNAs, while translation errors can generate misfolded proteins that trigger ribosome-associated

quality control pathways (36). Using XL-MS, we mapped detailed structural information within

this heterogeneous complex, capturing not only intra- and inter-subunit contacts (Fig. S11A-D),

but also interactions with proteins involved in ribosome biogenesis, translation, folding, quality

control, stress response, and other regulatory pathways (Fig. 6A). These interactions were

resolved at the amino acid level, providing detailed insight into the architecture of the functional

interface (Fig. 6B, Table S4).

While many interactions showed good STRING scores (≥ 0.4), our analysis also revealed

numerous novel interactions (green edges) between specific protein pairs. Although homologs of

these proteins have been studied in other organisms for their ribosome-related functions, their

specific roles in plants remain largely uncharacterized. Our XL-MS data suggest conserved

functions: for example, the UTP7 homologs AT3G10530 and NOG1-1 may contribute to

ribosome biogenesis (37, 38), whereas AT5G50840 (α-taxilin-like) and ILITHYIA(ILA) may be

involved in protein folding or translation (39, 40). MBF1A, a transcriptional coactivator, may be

involved in stress responses at stalled ribosomes, similar to its role in yeast (41) . Other

candidates include a methyltransferase potentially involved in rRNA processing and polysome

assembly (42, 43) , a RIO kinase associated with ribosome biogenesis (44, 45), and PARP1,

which may regulate translation via ADP-ribosylation (46) (Fig. 6C). By mapping these

interactions to specific ribosomal proteins, our XL-MS dataset provides a valuable resource for

future functional studies of ribosome-associated factors in plants.

We examined ubiquitination-related interactors, including the E3 ligases EBF1 and

AT4G02760, and the deubiquitinase UBP14. However, we did not identify homologs of known

ribosome-targeting E3 ligases from other species. EBF1 regulates ethylene signaling by

targeting EIN3/EIL1 for degradation (47), whereas UBP14 is essential for development in

Arabidopsis but dispensable in yeast (48). Given the active turnover of defective ribosomes, we

confirmed the *in vivo* ubiquitination of 37 large and 27 small ribosomal subunits by ubiquitin

diGly proteomic analysis (Fig. 6A, Table S4). Together with observed interactions with the 26S

proteasome components RPN8A, PAD1 (Fig.6A), these results suggest the ribosome undergoes

ubiquitination and may be targeted for degradation by the 26S proteosome.

Histone interactome reveals conserved and novel chromatin-associated interactions

Our XL-MS analysis revealed extensive interactions between histones and both known and novel

interactors, providing new insights into the networks involved in gene expression and chromatin

regulation (Fig. 7A). We identified several nuclear complexes associated with chromatin,

including nucleosome assembly proteins (NAPs), chromosome structure maintenance proteins

(SMCs), nuclear RNA polymerase components (NRPs), chromatin regulators (PDP3, MSI4),

transcriptional activator and acetylation reader (AT2G36410 and ACIP1), and subunits of the

PAF complex (VIP2, VIP4) (Fig. 7A).

256   Major histone interactors include chaperones (SDJ2, NUC-L1), chromatin remodeling

factors (EBS, SYD, SPK1), a transcription factor (SPL3), co-regulators (SNL6), DNA repair

proteins (MSH2), and structural proteins such as HMGB7. While many of these proteins are

known to influence chromatin architecture and gene regulation, their direct interactions with

histones and their roles in plant systems remained largely unexplored. Notably, MOS7, a

nucleoporin homologous to human NUP88, cross-linked with several histones, providing direct

evidence for nuclear envelope-histone interactions that may contribute to chromatin organization

in plants (49).

264   In addition, we identified histone interactions with several enzymatic regulators,

including the kinase MLK4 and the poly(ADP-ribose) polymerase PARP1 (both cross-linked to

H1), and the ubiquitin-like protein NEDD8 (cross-linked to H2B and H3). While the roles of

PARP1 and NEDD8 in histone regulation have not been defined in plants, their involvement in

histone modification has been well documented in other organisms (50-53). MLK4, previously

shown to phosphorylate H2A to regulate flowering (54), was found to interact with H1,

suggesting an expanded role for MLK4 in histone regulation that warrants further investigation.

271   Strikingly, we also identified WSD4, an O-acyltransferase annotated as a wax ester

synthase/acyl coenzyme and predicted by TAIR to localize to the nucleus (Fig. 7A-B). Although

O-acyltransferases are typically associated with lipid metabolism (55), WSD4 was found to

cross-link with four H2B family members - HTB1, HTB9, HTB3 and HTB11- via four

independent cross-linked peptides (Fig. S12A-E). These data suggest that WSD4 may catalyze

histone acylation, possibly on serine or threonine residues, rather than the conventional lysine

residues modified by other acylation proteins.

Taken together, these findings reveal previously uncharacterized histone-associated

proteins and enzymatic activities in plants, providing new insights into chromatin biology and

laying the groundwork for future studies of plant gene regulation.

Discussion

Systematic mapping of PPIs has greatly advanced our understanding of cellular function,

but plant PPI networks remain largely unexplored (5, 56). While comprehensive maps exist for

humans and other model organisms (57-59), similar efforts are just beginning in plants (60-62).

This gap represents a critical opportunity, as plants are central to addressing global challenges

such as food security, climate change, and sustainability. Expanding proteome-wide PPI

mapping and structural biology studies in plants will accelerate our mechanistic understanding of

cellular processes and drive advances in agriculture and biotechnology.

We developed a systems biology approach to capture molecular machines in action by

mapping protein-protein interactions (PPIs) and their interfaces using XL-MS. To overcome the

challenges of XL-MS in complex biological samples, we established an effective pipeline

integrating the PhoX crosslinker, subcellular fractionation, and high-pH reversed-phase HPLC

fractionation, enabling the identification of over 47,000 crosslinks. PDB and AlphaFold mapping

confirmed the quality of the data.

Our XL-MS analysis identified 1,446 high-confidence PPIs among 1,229 proteins,

providing a comprehensive map of protein interactions in Arabidopsis. Functional categorization

and STRING analysis revealed that these interactions span diverse biological processes, with

54.2% unannotated in STRING, demonstrating the ability of XL-MS to uncover novel networks.

The dataset also revealed the topological organization of protein complexes, including the 70S

ribosome, 26S proteasome, photosystems I and II, ATPases, and chromatin-associated factors.

XL-MS was particularly effective in resolving spatial relationships within dynamic protein

assemblies, such as mapping flexible proteins in the 70S ribosome stalk regions and transient

repair proteins in PSII.

XL-MS allows the detection of both known and novel PPIs in heterogeneous populations,

greatly expanding our understanding of conserved molecular machines such as ribosome and

histones (63, 64). While many of the identified interactors have characterized roles in other

species, their functions in plants remain unexplored. We discovered several ribosome-associated

E3 ligases and confirmed the *in vivo* ubiquitination of most ribosomal proteins, linking ribosome

function to ubiquitination-mediated regulation. In addition, our analysis revealed extensive plant-

specific PPIs, particularly in chloroplasts, highlighting unique aspects of the plant cellular

machinery.

Our XL-MS analysis revealed WSD-4, an O-acyltransferase, as a novel factor potentially

involved in histone regulation. WSD-4 forms cross-links with several members of the histone

H2B family and is predicted to localize in the nucleus. This family of O-acyltransferases is

typically involved in lipid metabolism (55); their role in protein modification has not been

reported. Interestingly, short-chain lysine acylations have been implicated in the regulation of

gene expression (65). Given the emerging evidence that metabolic enzymes such as ACAT1

(acetyl-CoA acetyltransferase 1) (66) and PORCN (membrane-bound O-acyltransferase with

protein-serine O-palmitoleoytransferase activity) (67, 68) also modify proteins, we propose that

WSD-4 may catalyze histone acylation, targeting serine or threonine residues. This provides a

novel mechanism of chromatin regulation, suggesting that WSD-4 works in concert with other

histone modifying enzymes to influence histone dynamics, chromatin structure, and gene

expression.

Despite Arabidopsis being a well-studied model organism, many of its genes remain

uncharacterized (69). This study highlights how XL-MS can map PPIs in semi-native conditions

and accelerate functional annotation, providing new insights into cellular organization. While

focused on Arabidopsis, our approach has broad applicability and provides a valuable resource

for comparative proteomic and structural studies across species. These findings improve protein

annotation, enhance understanding of protein complex architectures, and lay the foundation for

future discoveries.

Acknowledgement

This work was supported by the National Institutes of Health (NIGMS R01GM135706 and

S10OD030441 to S-L.X.) and by the Carnegie endowment fund for the Carnegie Mass

Spectrometry Facility. We thank Prof. Laura Helen Gunn for providing the structure of

Arabidopsis Rubisco L8S8, and Drs. Mike Trnka and Robert Chalkley (UCSF) and Dr. Rosa

Viner (Thermo Fisher Scientific) for technical assistance. We also thank the Hong Qiao lab (UT

Austin) for the nuclear isolation protocol and the Adrien Burlacot lab for the chloroplast isolation

protocol. We thank Dr. Ajeet Chaudhary and Prof. Arthur Grossman for their critical reading of

the manuscript.

Author contributions

S.L.X conceived the project and supervised the study. C.S.T. and R.S. designed and performed

most of the XL-MS experiments. R.S. developed the initial XL workflow. W.C.C performed

chloroplast XL-MS. S.K. performed ubiquitination enrichment. C.S.T and S.L.X analyzed all

the MS data and prepared figures, with assistance from A.V.R, G.L. and K.H. C.S.T and S.L.X

interpreted the data and wrote the manuscript with feedback from all authors.

Data Availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the

PRIDE partner repository with the dataset accession numbers. All other data is available from the

corresponding author on reasonable request.

Supplemental Table

Table S1: Summary of cross-linked peptide, including intra- and inter- protein cross-links, PPI

annotations based on STRING and mapping of intra-cross-links to known structures from PDB and

predicted structure from AlphaFold.

Table S2: Summary of cross-linked protein information and SUBA annotation.

Table S3: Analysis of protein classification using PANTHER.

Table S4: Detected cross-links for ribosomal and histones PPIs, and a list of ribosome-related

ubiquitinated peptides.

Table S5: Names, accession numbers and description used in the figures and a glossary.

Material and methods

Plant growth condition

Arabidopsis thaliana wild-type (Col-0) seeds were sterilized and sown on ½ Murashige

and Skoog (MS) agar square plates supplemented with 0.6% (w/v) phytoblend and lined with

nylon fabric. After stratification at 4°C for 3 days, the plates were transferred to growth

conditions.

Light-grown seedlings: plants were vertically grown in a growth chamber at 22°C under a

16-hour light / 8-hour dark cycle for 10-12 days, then flash frozen in liquid nitrogen and stored at

-80°C for crosslinking experiments on cell lysates and isolated chloroplasts. Dark-grown

seedlings: plants were grown horizontally at 22°C in complete darkness for 3 days in a growth

chamber, then flash frozen in liquid nitrogen and stored at -80°C for nucleus isolation. Dark

growth was used to reduce Rubisco content.

Frozen Arabidopsis tissue was cryo-ground to a fine powder using a Retsch MM400

Mixer Mill for downstream processing. Different amounts of tissue were used for cross-linking

experiments: 6 g from 10 d seedlings for cell lysates, 195 g from 12d seedlings for chloroplasts,

and 32 g (replicate 1 with PhoX) or 14 g (replicate 2 with tbu-PhoX) from 3d dark-grown

seedlings for nuclei.

Fraction of plant seedling tissues for cell lysate, chloroplast and nucleus

Cell lysates: Homogenized 10 d old light growth tissues were extracted using MOPS buffer [100

mM MOPS pH 7.6, 150 mM NaCl, 1% Nonidet P-40, 1 mM PMSF, 1x Roche protease inhibitor]

at a tissue:buffer ratio of 1:1 (g/mL, w/v). The samples were centrifuged at 20,000 g for 15 min

at 4°C, and the supernatant was then filtered twice with Miracloth and used for the cross-linking

reaction.

Chloroplast isolation: Frozen, homogenized tissue powder from 12 d old seedlings was

mixed with a grinding buffer similar to (70), using HEPES instead of Tricine for crosslinker

compatibility. Briefly, 5 g of tissue powder was ground in a mortar and pestle with 30ml of

grinding buffer [20mM HEPES-KOH pH 8.0, 330mM sorbitol, 5 mM EDTA, 5 mM EGTA, 5

mM $MgCl_2$, and 10 mM $NaHCO_3$] at 4°C, and under green illumination only. The resuspended

homogenate was filtered twice through two layers of Miracloth. 30 ml of the filtrate was

carefully transferred onto 10 ml of 40% Percoll [20 mM HEPES-KOH pH 8.0, 330 mM sorbitol,

40% v/v Percoll, 2.5 mM EDTA, 5 mM $MgCl_2$, and 10 mM $NaHCO_3$] in a 50ml centrifuge tube,

and centrifuged at 2,600g for 10 min at 4°C. The bottom pellet, enriched in intact chloroplasts,

was collected and resuspended in 30 ml of wash buffer [20 mM HEPES-KOH pH 8.0, 330 mM

sorbitol, 2.5 mM EDTA, 2 mM $MgCl_2$, and 10 mM $NaHCO_3$], and centrifuged at 2,600g for 3

min at 4°C. Each pellet derived from 5 g of tissue was washed twice with 4.5 ml of 1x PBS

supplemented with $MgCl_2$ [137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4,

and 2.5 mM $MgCl_2$] and then resuspended in 1mL 1x PBS supplemented with $MgCl_2$ for the

cross-linking reaction. This process was repeated across multiple batches, with a total of 195 g

of tissue used for cross-linking.

Nuclear isolation: The method for nuclear isolation was adapted from (71) with slight

modification. Powdered dark-growth Arabidopsis tissue of approximately 2 g was completely

homogenized with lysis buffer [20 mM HEPES (pH 7.4), 20 mM KCl, 2 mM EDTA, 2.5 mM

$MgCl_2$, 25% glycerol, 250 mM sucrose, 1 mM dithiothreitol, 1 mM phenylmethylsulfonyl

fluoride (PMSF), 1× protease inhibitor cocktail, and 0.2% formaldehyde] at tissue:buffer ratio of

1:5 (g/mL, w/v). The homogenate was then filtered through a 40 μm cell strainer and centrifuged

at 4°C, 1,500g for 10 min. The pellet was retained as the nuclear fraction. The nuclear pellet was

then washed 5 times with 5 mL of nuclear washing buffer [20 mM HEPES (pH 7.4), 2.5 mM

$MgCl_2$, 25% glycerol, 0.2% Triton X-100, 1 mM PMSF, and 1× protease inhibitor cocktail] and

centrifuged at 4°C, 1,500g for 10 min to collect the pellets after each wash. The washed nuclear

pellet was then resuspended in 0.5 mL of nuclear resuspension buffer 1 [20 mM HEPES (pH

7.4), 10 mM $MgCl_2$, 250 mM sucrose, 0.2% Triton X-100, 1 mM PMSF, and 1× protease

inhibitor cocktail] and carefully laid on top of nuclear resuspension buffer 2 [20 mM HEPES (pH

7.4), 10 mM $MgCl_2$, 1.7 M sucrose, 0.2% Triton X-100, 1 mM PMSF, and 1× protease inhibitor

cocktail], then centrifuged at 4°C,16,000g for 45 min. The supernatant was carefully removed,

and the nuclear pellet was resuspended in 0.5 mL 1 X PBS (pH 7.5) buffer for crosslinking

reaction. The intactness and purity of the nuclear pellet was assessed by DAPI staining under a

Leica SP8 confocal microscope. Multiple samples were prepared in parallel, and final isolated

nuclei were pooled before crosslinking.

Cross-linking reaction

PhoX or tBU-PhoX cross-linker was freshly prepared at a stock concentration of 100 mM in

DMSO for each cross-linking experiment (supplementary Fig. S3). A working concentration of

2-5 mM cross-linker was added to buffer solution containing whole cell lysate, isolated

chloroplasts or isolated nuclei and incubated for 2 h at 4°C (dark condition for chloroplast cross-

linking) with end-to-end rotation on a hulamixer. The cross-linking reaction was quenched by

adding Tris pH 8.0 to a final concentration of 20 mM and subjected to protein extraction.

Protein extraction, digestion, IMAC enrichment and RPLC fractionation

Protein extraction from the cross-linked samples, digestion to peptides, immobilized metal

affinity chromatography (IMAC) enrichment, and high pH reverse HPLC fractionation followed

the same procedure as described in (13). Multiple batches of PhoX-cross-linked peptides from

isolated intact chloroplasts were combined before IMAC enrichment. More information can be

found in supplementary Fig. S3. Multiple batches of samples were combined prior to high pH

reversed-phase HPLC fractionation (RPLC).

To generate selected proteomic libraries, flow-through peptides were retained in the

IMAC enrichment process for cell lysate and chloroplast samples. The chloroplast flow-through

peptides were further fractionated using the Pierce High pH Reverse-Phase Peptide Fractionation

Kit and collected in 8 fractions, measured by LC/MS/MS. The selected cell lysate database

contained 7835 abundant proteins, and the selected chloroplast database contained 9798 proteins.

Identification of ubiquitinated proteins by enrichment of ubiquitinated peptides

Proteins were extracted as described in Xu et al. (2017), followed by enrichment using the

PTMScan® Ubiquitin Remnant Motif (K-ε-GG) Kit (Cell Signaling Technology) according to

the manufacturer's instructions. The resulting peptides were then analyzed by LC-MS/MS.

LC-MS/MS analysis

Cross-linked peptides were analyzed by liquid chromatography-tandem mass spectrometry (LC-

MS/MS) using an Easy-nLC 1200 ultra-performance liquid chromatography (UPLC) system

(Thermo Fisher) coupled to an Orbitrap Eclipse Tribrid quadrupole Orbitrap mass spectrometer

(Thermo Fisher). Peptides were first loaded onto a C18 trap column (Acclaim PepMap 100 C18)

and then separated on an analytical C18 column (Aurora Series, 25 cm × 75 µm ID, Ion

Optics). The flow rate was set to 300 nL/min and the separation was performed over a 120-

minute gradient. Peptides were eluted using a gradient from 3% to 28% solvent B over 106

minutes, followed by an increase from 28% to 44% over 15 minutes and a final wash at 90%

solvent B for 14 minutes. Solvent A consisted of 0.1% formic acid and solvent B consisted of

80% (v/v) acetonitrile with 0.1% (v/v) formic acid.

Data was acquired on an Orbitrap Eclipse mass spectrometer. Full MS scans were

acquired over a mass-to-charge (m/z) range of 375–1600 at a resolution of 120,000, with an

automatic gain control (AGC) target of 200,000, a maximum injection time of 50 ms, normalized

AGC target set to 50%, and RF lens set to 30%. The most intense multiply charged precursors

(charge states 2–8) were selected for fragmentation with a cycle time of 3 seconds. MS/MS scans

were acquired at a resolution of 15,000 with an AGC target of 5×10^4, maximum injection time

of 22 ms, isolation window of 1.4 m/z, and normalized AGC target set to 100%. Fragmentation

was performed using higher-energy collisional dissociation (HCD) with a normalized collision

energy (NCE) of 27. Dynamic exclusion was enabled for 30 seconds. For cross-linked peptide

detection, charge states 3 to 8 were used.

Real-time library search (RTLS) was used to reduce mono-link identifications. RTLS

included a low-resolution MS2 survey scan performed in the ion trap over an m/z range of 120–

500. The resulting MS2 peaks were matched against a custom spectral library containing

signature peak patterns characteristic of mono-links and cross-links (signature m/z values:

201.1231, 215.1387, 312.0632, and 377.1269, each with characteristic relative intensities for

mono- or cross-linked species). Only precursor ions whose survey MS2 spectra matched the

cross-link pattern using the "Similarity Search" function were selected for high-resolution MS2

acquisition. The RTLS method increases the overall cycle time compared to standard XL-MS

acquisition, resulting in a lower overall identification rate.

Data search using pLINK2 and Protein Prospector

Liner peptide searches were searched using Protein Prospector against Arabidopsis TAIR

database as described in (72). Precursor and fragment mass tolerance were set up as 10 and 20

ppm.

Crosslinked peptide searches were performed using pLink 2.3.11 with the following

parameters: crosslinker set to PhoX (209.972 Da); peptide length range: 6-60 amino acids;

enzyme: trypsin with up to 3 missed cleavages. Fixed modification: Cys carbamidomethylation;

variable modifications: protein N-terminal acetylation and Met oxidation. Precursor and

fragment mass tolerances were set to 5 ppm and 20 ppm, respectively.

MS2 spectra were searched against the TAIR10_pep_20101214, selective cell lysate

(7835) or selective chloroplast (9897) databases as described in Supplementary Fig.S3. Results

were filtered at 5% FDR for cell lysate and nuclear data sets and 1% FDR for chloroplast data

sets, all at the peptide spectrum matches (PSMs) level. FDRs for monolinks, loop-links, intra-

and inter-protein cross-links were calculated separately by pLINK2. To compare two acquisition

methods (XL-Standard [Standard] vs. Real-time library search [RTLS]), data were acquired on

the whole cell lysate XL samples, the search was against TAIR10_pep_20101214 database and

compared.

Data compilation and reporting.

Intra- and inter-protein cross-links were compiled from multiple experiments (Supplementary

Fig. S3) and reported separately (Supplementary Table S1). The total number of cross-linked

peptide pairs was determined by counting unique lysine–lysine (K–K) residue pairs along with

their corresponding backbone sequences. Peptides with identical backbone sequences and the

same K–K pair were collapsed into a single count, regardless of charge state or peptide backbone

modifications (e.g., oxidation).

Subcellular localization analysis of proteins using SUBA5

The subcellular localization of cross-linked proteins was analyzed using SUBA5

(https://suba.live/). Accession IDs of proteins identified in both intra- and inter-protein cross-

links were submitted as input. Only the localization consensus data were used for classification.

Proteins assigned to multiple compartments were counted once in each corresponding

localization category when calculating total counts.

PANTHER protein class analysis and mapping intralinks to PDB and AlphaFold

structures using ComMap

Arabidopsis gene identifiers (ATG accession numbers) were used for protein class analysis using

PANTHER™ Protein Class tool, which contains 210 classification terms (version 19.0, released

2024-06-20).

To map the results of intralinks against experimentally determined structures from the Protein

Data Bank (PDB) and predicted structures from AlphaFold, the protein accession IDs were first

converted to UniProt IDs. The intralink information was then mapped to available structures

using ComMap, a tool that enables large-scale structure-based mapping by integrating XL-MS

data with existing structural models and performing distance calculations (16).

Crosslinking data visualization

Cytoscape visualization: The entire cross-link-based protein-protein interaction (PPI) network

was visualized in Cytoscape (version 3.10, https://cytoscape.org/) using inter-protein cross-link

data, including heteromeric and homomultimeric cross-links. Nodes were colored according to

protein class information from the PANTHER classification system, and edges were colored

according to the STRING interaction score (https://string-db.org/).

Xiview: Protein–protein interactions (PPIs), including both inter- and intra-protein cross-

links, were visualized using XiView (https://xiview.org/) (Combe et al., 2024). PPIs were

categorized via the built-in STRING data loader (score cutoff 0.4). Cross-linked peptides

supporting the same lysine pair were consolidated into a unique K-K linkage, collapsing

redundancy (e.g. due to miscleavage events) and reporting each linkage only once.

Mapping of crosslinks to high-resolution structures of proteins/complexes. Intra- and

inter-crosslinks were superimposed on PDB or AlphaFold structures using the built-in function

of XiView, which also calculated the corresponding crosslink distances within the high-

resolution structures. Alignments of protein orthologs were performed and visualized using

Pymol (https://www.pymol.org/).

Figure legend

Reference

1.   Yu C & Huang L (2018) Cross-Linking Mass Spectrometry: An Emerging Technology for
Interactomics and Structural Biology. *Anal Chem* 90(1):144-165.
2.   Piersimoni L, Kastritis PL, Arlt C, & Sinz A (2022) Cross-Linking Mass Spectrometry for
Investigating Protein Conformations and Protein-Protein Interactions horizontal line A
Method for All Seasons. *Chem Rev* 122(8):7500-7531.
3.   Michaelis AC*, et al.* (2023) The social and structural architecture of the yeast protein
interactome. *Nature* 624(7990):192-200.
4.   Xu SL, Shrestha R, Karunadasa SS, & Xie PQ (2023) Proximity Labeling in Plants. *Annu
Rev Plant Biol* 74:285-312.
5.   Greenblatt JF, Alberts BM, & Krogan NJ (2024) Discovery and significance of protein-
protein interactions in health and disease. *Cell* 187(23):6501-6517.
6.   Lee K & O'Reilly FJ (2023) Cross-linking mass spectrometry for mapping protein
complex topologies in situ. *Essays Biochem* 67(2):215-228.
7.   Zhu X*, et al.* (2016) In planta chemical cross-linking and mass spectrometry analysis of
protein structure and interaction in Arabidopsis. *Proteomics* 16(13):1915-1927.
8.   Jayachandran B, Parvin TN, Alam MM, Chanda K, & Mm B (2022) Insights on Chemical
Crosslinking Strategies for Proteins. *Molecules* 27(23).
9.   Smythers AL, Iannetta AA, & Hicks LM (2021) Crosslinking mass spectrometry unveils
novel interactions and structural distinctions in the model green alga Chlamydomonas
reinhardtii. *Mol Omics* 17(6):917-928.
10.   Dai S*, et al.* (2023) Capturing the hierarchically assorted modules of protein-protein
interactions in the organized nucleome. *Mol Plant* 16(5):930-961.
11.   Steigenberger B, Pieters RJ, Heck AJR, & Scheltema RA (2019) PhoX: An IMAC-
Enrichable Cross-Linking Reagent. *ACS Cent Sci* 5(9):1514-1522.
12.   Jiang PL*, et al.* (2022) A Membrane-Permeable and Immobilized Metal Affinity
Chromatography (IMAC) Enrichable Cross-Linking Reagent to Advance In Vivo Cross-
Linking Mass Spectrometry. *Angew Chem Int Ed Engl* 61(12):e202113937.
13.   Xu S-L*, et al.* (2017) Proteomic analysis reveals O-GlcNAc modification on proteins with
key regulatory functions in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* 114(8):E1536-
E1543.
14.   Chen ZL*, et al.* (2019) A high-speed search engine pLink 2 with systematic evaluation
for proteome-scale identification of cross-linked peptides. *Nat Commun* 10(1):3404.
15.   Ruwolt M*, et al.* (2023) Real-Time Library Search Increases Cross-Link Identification
Depth across All Levels of Sample Complexity. *Anal Chem* 95(12):5248-5255.
16.   Zhang W*, et al.* (2023) ComMap: a software to perform large-scale structure-based
mapping for cross-linking mass spectrometry. *Bioinformatics* 39(2).
17.   Valegard K*, et al.* (2018) Structural and functional analyses of Rubisco from arctic
diatom species reveal unusual posttranslational modifications. *J Biol Chem*
293(34):13033-13043.
18.   Murachelli AG, Ebert J, Basquin C, Le Hir H, & Conti E (2012) The structure of the ASAP
core complex reveals the existence of a Pinin-containing PSAP complex. *Nat Struct Mol
Biol* 19(4):378-386.
19.   Bi Y*, et al.* (2021) Arabidopsis ACINUS is O-glycosylated and regulates transcription and
alternative splicing of regulators of reproductive transitions. *Nat Commun* 12(1):945.
20.   Shrestha R*, et al.* (2025) Next-Generation Mapping of the ACINUS-Mediated Alternative
Splicing Machinery and Its Regulation by O-glycosylation in Arabidopsis. . *BioRxiv*.

21.   Yu A, *et al.* (2020) Photosynthetic Phosphoribulokinase Structures: Enzymatic
Mechanisms and the Redox Regulation of the Calvin-Benson-Bassham Cycle. *Plant Cell*
32(5):1556-1573.
22.   Wheat A, *et al.* (2021) Protein interaction landscapes revealed by advanced in vivo
cross-linking-mass spectrometry. *Proc Natl Acad Sci U S A* 118(32).
23.   Shannon P, *et al.* (2003) Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res* 13(11):2498-2504.
24.   Mi H, *et al.* (2021) PANTHER version 16: a revised family classification, tree-based
classification tool, enhancer regions and extensive API. *Nucleic Acids Res* 49(D1):D394-
D403.
25.   Szklarczyk D, *et al.* (2021) The STRING database in 2021: customizable protein-protein
networks, and functional characterization of user-uploaded gene/measurement sets.
Nucleic Acids Res 49(D1):D605-D612.
26.   Ahmed T, Yin Z, & Bhushan S (2016) Cryo-EM structure of the large subunit of the
spinach chloroplast ribosome. *Sci Rep* 6:35793.
27.   Bieri P, Leibundgut M, Saurer M, Boehringer D, & Ban N (2017) The complete structure
of the chloroplast 70S ribosome in complex with translation factor pY. *EMBO J*
36(4):475-486.
28.   Liljas A & Sanyal S (2018) The enigmatic ribosomal stalk. *Q Rev Biophys* 51:e12.
29.   Komenda J, Sobotka R, & Nixon PJ (2024) The biogenesis and maintenance of PSII:
Recent advances and current challenges. *Plant Cell* 36(10):3997-4013.
30.   Chen H, *et al.* (2006) A Psb27 homologue in Arabidopsis thaliana is required for efficient
repair of photodamaged photosystem II. *Plant Mol Biol* 61(4-5):567-575.
31.   Wei L, *et al.* (2010) LPA19, a Psb27 homolog in Arabidopsis thaliana, facilitates D1
protein precursor processing during PSII biogenesis. *J Biol Chem* 285(28):21391-21398.
32.   Li A, *et al.* (2024) Structural basis for an early stage of the photosystem II repair cycle in
Chlamydomonas reinhardtii. *Nat Commun* 15(1):5211.
33.   Weisz DA, *et al.* (2019) A novel chlorophyll protein complex in the repair cycle of
photosystem II. *Proc Natl Acad Sci U S A* 116(43):21907-21913.
34.   Xingxing C, *et al.* (2018) Crystal structure of Psb27 from Arabidopsis thaliana
determined at a resolution of 1.85 A. *Photosynth Res* 136(2):139-146.
35.   Sheng X, *et al.* (2019) Structural insight into light harvesting for photosystem II in green
algae. *Nat Plants* 5(12):1320-1330.
36.   Martinez-Seidel F, Beine-Golovchuk O, Hsieh YC, & Kopka J (2020) Systematic Review
of Plant Ribosome Heterogeneity and Specialization. *Front Plant Sci* 11:948.
37.   Chaker-Margot M (2018) Assembly of the small ribosomal subunit in yeast: mechanism
and regulation. *RNA* 24(7):881-891.
38.   Lee S, *et al.* (2017) The small GTPase, nucleolar GTP-binding protein 1 (NOG1), has a
novel role in plant innate immunity. *Sci Rep* 7(1):9260.
39.   Yoshida K, *et al.* (2005) Interaction of the taxilin family with the nascent polypeptide-
associated complex that is involved in the transcriptional and translational processes.
Genes Cells 10(5):465-476.
40.   Faus I, *et al.* (2018) Arabidopsis ILITHYIA protein is necessary for proper chloroplast
biogenesis and root development independent of eIF2alpha phosphorylation. *J Plant
Physiol* 224-225:173-182.
41.   Kim KQ, *et al.* (2024) Multiprotein bridging factor 1 is required for robust activation of the
integrated stress response on collided ribosomes. *Mol Cell* 84(23):4594-4611 e4599.
42.   Tafforeau L, *et al.* (2013) The complexity of human ribosome biogenesis revealed by
systematic nucleolar screening of Pre-rRNA processing factors. *Mol Cell* 51(4):539-551.

43.   Chen J*, et al.* (2024) The Ptch/SPOUT1 methyltransferase deposits an m(3)U
627   modification on 28S rRNA for normal ribosomal function in flies and humans. *Sci Adv*
628   10(50):eadr1743.
629   44.   Ghandadi M, Dobi A, & Malhotra SV (2024) A role for RIO kinases in the crosshair of
630   cancer research and therapy. *Biochim Biophys Acta Rev Cancer* 1879(3):189100.
631   45.   Messling JE*, et al.* (2022) Targeting RIOK2 ATPase activity leads to decreased protein
632   synthesis and cell death in acute myeloid leukemia. *Blood* 139(2):245-255.
633   46.   Kim DS, Challa S, Jones A, & Kraus WL (2020) PARPs and ADP-ribosylation in RNA
634   biology: from RNA expression and processing to protein translation and proteostasis.
635   *Genes Dev* 34(5-6):302-320.
636   47.   Binder BM*, et al.* (2007) The Arabidopsis EIN3 binding F-Box proteins EBF1 and EBF2
637   have distinct but overlapping roles in ethylene signaling. *Plant Cell* 19(2):509-523.
638   48.   Doelling JH, Yan N, Kurepa J, Walker J, & Vierstra RD (2001) The ubiquitin-specific
639   protease UBP14 is essential for early embryo development in Arabidopsis thaliana. *Plant*
640   *J* 27(5):393-405.
641   49.   Van de Vosse DW, Wan Y, Wozniak RW, & Aitchison JD (2011) Role of the nuclear
642   envelope in genome organization and gene expression. *Wiley Interdiscip Rev Syst Biol*
643   *Med* 3(2):147-166.
644   50.   Zhang S, Yu Q, Li Z, Zhao Y, & Sun Y (2024) Protein neddylation and its role in health
645   and diseases. *Signal Transduct Target Ther* 9(1):85.
646   51.   Alemasova EE & Lavrik OI (2019) Poly(ADP-ribosyl)ation by PARP1: reaction
647   mechanism and regulatory proteins. *Nucleic Acids Res* 47(8):3811-3827.
648   52.   Messner S*, et al.* (2010) PARP1 ADP-ribosylates lysine residues of the core histone
649   tails. *Nucleic Acids Res* 38(19):6350-6362.
650   53.   Nosella ML*, et al.* (2024) Poly(ADP-ribosyl)ation enhances nucleosome dynamics and
651   organizes DNA damage repair components within biomolecular condensates. *Mol Cell*
652   84(3):429-446 e417.
653   54.   Su Y*, et al.* (2017) Phosphorylation of Histone H2A at Serine 95: A Plant-Specific Mark
654   Involved in Flowering Time Regulation and H2A.Z Deposition. *Plant Cell* 29(9):2197-
655   2213.
656   55.   Li F*, et al.* (2008) Identification of the wax ester synthase/acyl-coenzyme A:
657   diacylglycerol acyltransferase WSD1 required for stem wax ester biosynthesis in
658   Arabidopsis. *Plant Physiol* 148(1):97-107.
659   56.   Vazquez A, Flammini A, Maritan A, & Vespignani A (2003) Global protein function
660   prediction from protein-protein interaction networks. *Nat Biotechnol* 21(6):697-700.
661   57.   Giurgiu M*, et al.* (2019) CORUM: the comprehensive resource of mammalian protein
662   complexes-2019. *Nucleic Acids Res* 47(D1):D559-D563.
663   58.   Luck K*, et al.* (2020) A reference map of the human binary protein interactome. *Nature*
664   580(7803):402-408.
665   59.   Burke DF*, et al.* (2023) Towards a structurally resolved human protein interaction
666   network. *Nat Struct Mol Biol* 30(2):216-225.
667   60.   Brandao MM, Dantas LL, & Silva-Filho MC (2009) AtPIN: Arabidopsis thaliana protein
668   interaction network. *BMC Bioinformatics* 10:454.
669   61.   Herrera-Ubaldo H*, et al.* (2023) The protein-protein interaction landscape of transcription
670   factors during gynoecium development in Arabidopsis. *Mol Plant* 16(1):260-278.
671   62.   Yang X*, et al.* (2020) PlaPPISite: a comprehensive resource for plant protein-protein
672   interaction sites. *BMC Plant Biol* 20(1):61.
673   63.   Simsek D*, et al.* (2017) The Mammalian Ribo-interactome Reveals Ribosome Functional
674   Diversity and Heterogeneity. *Cell* 169(6):1051-1065 e1018.
675   64.   Peng Y, Markov Y, Goncearenco A, Landsman D, & Panchenko AR (2021) Human
676   Histone Interaction Networks: An Old Concept, New Trends. *J Mol Biol* 433(6):166684.

677   65.   Sabari BR, Zhang D, Allis CD, & Zhao Y (2017) Metabolic regulation of gene expression
678   through histone acylations. *Nat Rev Mol Cell Biol* 18(2):90-101.
679   66.   Goudarzi A (2019) The recent insights into the function of ACAT1: A possible anti-cancer
680   therapeutic target. *Life Sci* 232:116592.
681   67.   Liu Y*, et al.* (2022) Mechanisms and inhibition of Porcupine-mediated Wnt acylation.
682   *Nature* 607(7920):816-822.
683   68.   Nile AH & Hannoush RN (2016) Fatty acylation of Wnt proteins. *Nat Chem Biol* 12(2):60-
684   69.
685   69.   Xue B & Rhee SY (2023) Status of genome function annotation in model organisms and
686   crops. *Plant Direct* 7(7):e499.
687   70.   Casazza AP, Tarantino D, & Soave C (2001) Preparation and functional characterization
688   of thylakoids from Arabidopsis thaliana. *Photosynth Res* 68(2):175-180.
689   71.   Shao Z*, et al.* (2024) Nuclear pyruvate dehydrogenase complex regulates histone
690   acetylation and transcriptional regulation in the ethylene response. *Sci Adv*
691   10(30):eado2825.
692   72.   Shrestha R*, et al.* (2022) 15N Metabolic Labeling Quantification Workflow in Arabidopsis
693   Using Protein Prospector. *Front. Plant Sci.* 13:832562.
694

Figure 1. Efficient cross-linking workflow for complex protein samples identifies numerous cross-links in Arabidopsis.

(**A**) Schematic of the workflow using PhoX or tBu-PhoX to covalently cross-link spatially proximal lysine residues in Arabidopsis subcellular fractions (lysate, chloroplast, nucleus), followed by digestion, IMAC enrichment, high pH reversed-phase HPLC fractionation, and LC-MS/MS. Data were analyzed using pLINK2, Cytoscape, and xiVIEW.

(**B**) Summary of unique intra- and inter-protein cross-links from each fraction. Intra-links connect non-overlapping peptides within a protein; inter-links include heteromeric and homomultimeric cross-links.

(**C**) Subcellular localization of cross-linked proteins based on SUBA5, with redundancy of proteins assigned to multiple compartments.

Figure 2. Validation of intra- and inter-protein cross-links by structural mapping.
(**A-B**) Histogram of Euclidean Cα-Cα distances of intra-protein cross-links mapped to Arabidopsis structures from PDB (A) or AlphaFold (B), showing 94% and 89% within the 35Å constraint, respectively.
(**C-E**) Cross-links mapped to the Arabidopsis Rubisco L_8S_8 complex modeled from the L_2S_2 structure (PDB:5IU0). RBCL is shown in cyan and RBCS is shown in pink. Self(intra), heteromeric, and homomultimeric links are shown in purple, green, and orange, respectively. The circular view (C) and structural mapping (D-E) show that 74.56% of the cross-links are within 35Å.

(**F**) Cross-links between SR45 and ACINUS or PININ occur at known interaction interfaces involving SR45 RRM and RSB domains.

(**G-H**) Intra- and inter-protein cross-links between PRK and CP12.1 mapped onto the GAPDH-CP12-PRK complex (PDB:6KEZ), aligning with the known interaction interface.

Figure 3. XL-MS identifies direct protein-protein interactions and reveals known and novel complexes.

(**A**) XL-MS identified an Arabidopsis interactome comprising 1,446 PPIs, including 1,390 heteromeric and 56 homomultimeric interactions among 1,229 proteins. Nodes represent proteins colored by PANTHER protein class; edges represent interactions colored by the combined STRING confidence score.

(**B**) The breakdown of the STRING analysis shows that more than 71% of the detected heteromeric PPIs are either of low confidence or have no experimental evidence.

(**C-D**) Examples of specific PPIs detected by XL-MS that complement low confidence or missing experimental STRING scores: the ORP3B-PLDRP1 interaction (cross-linked at ORP3B-K7/PLDRP1-K193) and the NRPA2-uL1c interaction (NRPA2-K394/uL1c-K127).

Figure 4. XL-MS reveals unique topology of chloroplast 70S ribosome stalk regions.

(**A**) XL-MS identifies flexible stalk proteins (bL12c, uL1c, uL10c, uL11c), with cross-links color-coded by STRING confidence. Homodimer interfaces are found at the C-terminus of bL12c.

(**B**) Domain comparison between Arabidopsis bL12c and E. coli bL12.

(**C**) Cross-links mapped onto the predicted bL12c structure (AF-P36212-F1-v4) showing lysines K64, K137, K148, and K151 interacting with ribosomal subunits and the elongation factor CPEF-G. Nodes are color-coded according to annotation category.

Figure 5. XL-MS reveals dynamic connectivity and unique features in photosystem complexes.

(**A**) PSII topology showing cross-links color-coded by STRING confidence. Notable PSB27 cross-links (K93, K168) highlight interactions with PSII core and OEC components.

(**B**) XL-MS uncovers extensive interactions among PSI, PSII, LHCI, LHCII, ATPase, FtsH/SPPA proteases, and transient proteins involved in state transitions, assembly, repair, and chlorophyll metabolism.

Figure 6. XL-MS of ribosome complexes reveals conserved ribosome-associated factors.
(**A**) XL-MS identified extensive interactions within ribosomal subunits as well as with proteins involved in ribosome biogenesis, translation, folding, quality control, stress response, and other regulatory pathways. Ribosomal proteins are overlaid with ubiquitination data; red nodes indicate ubiquitinated proteins.
(**B**) Table showing covalently cross-linked lysine residues between ribosomal proteins, reflecting spatial proximity. Protein labels are color-coded according to the functional categories shown in (A).
(**C**) Table of XL-MS-identified ribosome-associated interactors with potential post-transcriptional or post-translational regulatory functions, many of which remain uncharacterized in plants.

Figure 7. Histone interactome reveals conserved and novel chromatin-associated interactions.

(**A**) XL-MS reveals extensive interactions between histones and both known and previously uncharacterized proteins, including components of chromatin-associated complexes. Enzymatic regulators are highlighted in red.

(**B**) Schematic illustration of histone modifications and polynucleosome organization affected by PARylation and neddylation. A potential role is proposed for an O-acyltransferase - previously associated with metabolic processes - that cross-links with four members of the histone H2B family, suggesting its involvement in histone modification based on XL-MS interaction data.

Supplementary Figure S1: Optimization of the PhoX cross-linker concentration for XL-MS.

(**A-D**) Different concentrations of PhoX cross-linker (0 to 10 mM) were tested on cell lysates from light (A, C, D) or dark grown seedlings (B). Increased protein cross-linking, indicated by the accumulation of higher molecular weight proteins (red arrow), was observed with increasing cross-linker concentration. However, no additional visible cross-linking was detected at higher concentrations (7-10 mM).

Supplementary Figure S2: **Standard XL acquisition outperforms Real-Time Library Search (RTLS) acquisition.**

Data were acquired on the Orbitrap Eclipse mass spectrometer using PhoX-enriched cross-linked peptides from cell lysate samples and searched against the entire Arabidopsis library. While RTLS acquisition significantly reduced the identification of monolinks (where one NHS ester group is cross-linked to a peptide while the other is hydrolyzed) and loop links (cross-links within a single peptide), it also significantly reduced the detection of inter-peptide cross-links (pairs of peptides covalently linked by the cross-linker), including both intra- and inter-protein cross-links (intralinks and interlinks).

Supplementary Fig. S3: Summary of XL-MS experiments.
Key details of sample preparation, data acquisition using the Orbitrap Eclipse mass spectrometer, and database searching using pLINK2 are provided. A summary of the unique cross-linked peptide residue pairs identified across experiments is also included.

Supplementary Fig. S4. Representative cross-links supporting the high confidence of XL-MS analysis.

(**A**) Multiple cross-links were identified between the ribosomal proteins uL15x and eL13z, including peptide pairs with different degrees of miscleavage cross-linked at identical sites (e.g., uL15x-K92 to eL13z-K166/K172).

(**B**) Cross-links were mapped to conserved regions of homologous proteins; for example, the ATPase subunit VHA-E3 (K49) was cross-linked to both VHA-B1 (K437) and VHA-B2 (K438).

(**C**) Similar cross-linking patterns were observed between homologous interactions, as shown by DNA topoisomerase 1 alpha (TOP1A-K672) cross-linked to two dehydrin proteins, ERD14-K38 and COR47-K46.

Supplementary Figure S5. Mapping of Rubisco cross-links on a high-resolution orthogonal structure of hornwort Rubisco.

Cross-links are color-coded by type: purple for self (intra)links, green for heteromeric (inter)links, and orange for homomultimeric links.

(**A**) Cross-links were mapped onto the high-resolution structure of L8S8 heterotetrameric Rubisco from hornwort (PDB: 9CHZ). RBCL subunits are shown in cyan and RBCS subunits in pink.

(**B**) Distance distribution of cross-links mapped onto Rubisco 9CHZ, showing that 71.6% of the identified cross-links fall within the expected spatial constraint of 35 Å.

Supplementary Figure S6. High-confidence identifications of cross-linked peptides between ACINUS–SR45 and PININ–SR45.

b-ions are shown in green and y-ions in yellow. A series of fragment ions—including those from α- and β-peptides, as well as cross-linker-containing ions—were detected, supporting the reliability of the cross-link identifications.

(**A**) Spectrum of the cross-linked peptide between ACINUS-K633 and SR45-K154.

(**B**) Spectrum of the cross-linked peptide between PININ-K274 and SR45-K154.

Supplementary Fig. S7. Topological mapping of multiple protein complexes by XL-MS. Proteins are shown as rectangles scaled to their respective lengths.
Cross-links are color-coded based on STRING score confidence thresholds, highlighting both known and novel interactions.

(**A**) Mapping of 26S proteasome subunits, highlighting interaction interfaces between RPN7 and DSS1(I) and DSS1(V). DSS1 proteins are particularly challenging to map in cryo-EM due to their small size, intrinsic disorder, and dynamic interaction properties.

(**B**) Mapping of photosystem I (PSI) showing connectivity between different subunits.

(**C**-**E**) Mapping of different ATPase complexes, including chloroplast, V-type, and mitochondrial ATPases.

(**F**) Mapping of the prefoldin complex.

(**G**) Mapping of the phosphoenolpyruvate carboxylase (PEPC) complex.

Supplementary Figure S8. Mapping of cross-links in the 70S ribosome complex onto high-resolution structures of spinach 70S ribosomes (PDB: 5MMM and 5X8P).
Cross-links are color-coded by type: purple for intralinks, green for heteromeric inter-protein cross-links, and orange for homomultimeric inter-protein cross-links.
(**A**) Structural mapping of 70S ribosome cross-links on the spinach cryo-EM structure (PDB: 5MMM), which lacks bL12 and other flexible proteins in the stalk region.
(**B**) Distance distribution of the cross-links mapped in (A). Of the 424 total cross-links detected, 276 were mapped (263 self-links and 13 heteromeric), with 241 falling within the 35 Å spatial constraint.
(**C**) Structural mapping of 70S ribosome cross-links onto an alternative spinach cryo-EM structure (PDB: 5X8P), which also lacks most of the flexible proteins in the stalk region.
(**D**) Distance distribution of the cross-links mapped in (C). Of the 424 cross-links detected, 234 were mapped (224 self-links and 10 heteromeric), with 206 falling within the 35 Å constraint.

Supplementary Figure S9. Mapping of PSB27 cross-links onto a high-resolution structure of 6KAC from Chlamydomonas.

(**A**) Superimposed AlphaFold structures of PSB27 homologs reveal unique features of Arabidopsis PSB27. While it contains a conserved C-terminal four-helix bundle, Arabidopsis PSB27 lacks a lipid anchor and shows divergence in sequence and surface charge.

(**B**) Cross-links between Arabidopsis PSB27 and PSII core/OEC subunits mapped onto the Chlamydomonas PSII structure (PDB: 6KAC), which captures residues 91-114 of PSB27. The distance constraints suggest a more dynamic interaction between PSB27 and the PSII complex.

Supplementary Figure S10. Mapping of PSII complex cross-links onto several available high-resolution structures. A total of 581 cross-links were mapped, including 504 self-links

and 77 heteromeric links. Cross-links are colored according to the type of cross-link (purple for self (intra) link; green for heteromeric (inter) link; orange for homomultimeric link). The distribution of cross-link distances from the integrated cross-link model mapping is shown alongside the structural mapping.

(**A-B**) Structure mapping of PSII cross-links onto the Chlamydomonas PSII-LHCII supercomplex (PDB: 6KAC), which includes the PSII reaction center, PsbO, and a short sequence of Psb27.

(**C-D**) Structure mapping of PSII cross-links onto the Arabidopsis PSII-LHCII supercomplex, (PDB: 5MDX) which represents mature PSII, containing the PSII reaction center and PsbO but lacking Psb27.

(**E-F)** Structure mapping of PSII cross-links onto the cyanobacterial PSII assembly intermediate (PDB: 7NHP), which presents a PSII assembly intermediate containing the PSII reaction center, Psb27 and Psb28.

(**G-H**) Structure mapping of PSII cross-links onto the Chlamydomonas PSII assembly intermediate (PDB: 8R2I), which represents a native PSII assembly intermediate containing the PSII reaction center and Psb27.

Supplementary Figure S11. Topology of the 80S ribosomal complex.
(**A**) 2D topological arrangement of the 40S and 60S ribosomal subunits based on their connectivity. Links between proteins are colored according to the confidence thresholds of the STRING score.
(**B**) Summary of cross-links and protein-protein interaction (PPI) information from (A).
(**C**) 80S ribosome cross-links mapped onto the high-resolution structure of the tobacco 80S ribosome (PDB: 8B2L). The large ribosomal subunit is colored orange and the small ribosomal subunit is colored red. Links are color-coded according to link type: purple for self-links (intralinks), green for heteromeric links (interlinks), and orange for homomultimeric links.
(**D**) Distribution of cross-link distances from the integrated cross-link model in (C).

Supplementary Fig. S12. XL-MS identifies WSD4 as a novel histone interactor that associates with H2B family members.

(**A**) Cross-links were detected between WSD4 (an O-acyltransferase) and several H2B family members, including HTB1, HTB9, HTB3, and HTB11.

(**B**-**E**) High-confidence cross-link spectra reveal interactions between WSD4 and H2B family members. Series of fragment ions-including those from α- and β-peptides, as well as cross-linker-containing ions-were detected, supporting the reliability of the cross-link identifications.