Clade-wide proteome analysis shows widespread non-canonical Dcr proteins in fungi.

Lorena Melet[1,2,3], Jonathan Canan[1,2,5,6], Pablo Villalobos[4], Boris Vidal-Veuthey[1], Fabian

González-Toro[1], Ivana Orellana[1], J. Andrés Rivas-Pardo[1], Juan P. Cárdenas[1], Carol

Moraga[2,5,6], Víctor Castro-Fernandez[4], Nathan R. Johnson[1,2, *], Elena A. Vidal[1,2,3,*]

[1]Centro de Genómica y Bioinformática, Universidad Mayor, 8580745 Santiago, Chile.

[2]Agencia Nacional de Investigación y Desarrollo-Millennium Science Initiative Program,

Millennium Institute for Integrative Biology (iBio), 7500565 Santiago, Chile.

[3]Agencia Nacional de Investigación y Desarrollo-Millennium Science Initiative Program,

Millennium Nucleus in Data Science for Plant Resilience (Phytolearning), 8370186 Santiago,

Chile

[4]Departamento de Biología, Facultad de Ciencias, Universidad de Chile, 7800003 Santiago,

Chile.

[5]Instituto de Ciencias de la Ingeniería, Universidad de O'Higgins, 2841959 Rancagua, Chile.

[6]Centro UOH de Bioingeniería (CUBI), Universidad de O'Higgins, 2841959 Rancagua, Chile.

*Corresponding authors

Nathan R. Johnson (nathan.johnson@umayor.cl)

Elena A. Vidal (elena.vidal@umayor.cl)

Abstract

Dicer (Dcr) proteins are essential components of RNA interference (RNAi) pathways, yet

their evolutionary diversity and structural variation in fungi remain poorly understood. In this

work, we surveyed 1,593 fungal proteomes spanning eight phyla, identifying 2,917 putative

Dcr proteins. We found a predominance of non-canonical Dcr types, in which one or more

domains such as Piwi/Argonaute/Zwille (PAZ), Helicase or double-stranded RNA binding

domain failed to be detected, suggesting loss or sequence divergence. Phylogenetic

analyses showed lineage-specific patterns of domain retention and loss, with canonical Dcrs

restricted to members of the Mucoromycota and Basidiomycota. Despite high sequence

divergence, AlphaFold-based structural predictions revealed conserved L-shaped

architectures and PAZ-like folds in divergent Dcrs. Using Molecular simulations and

electrostatic analysis, we characterized a divergent fungal PAZ found in non-canonical Dcrs,

which maintains conserved motifs and RNA-binding surface properties found in canonical

PAZ domains. Our findings highlight the evolutionary plasticity and functional relevance of

non-canonical RNAi machinery in fungi.

Introduction

RNA interference (RNAi) is a conserved eukaryotic mechanism with diverse roles, including gene expression regulation (Lim et al., 2005), antiviral defense (Wilkins et al., 2005), genome stability (Volpe et al., 2002), and communication between cells and across species (Buck et al., 2014; Cai et al., 2018; Wang et al., 2016). RNAi is mediated by small RNA (sRNA) molecules broadly categorized as 18-30 nucleotides in length, which are generated through the cleavage of exogenous or endogenous double-stranded RNA (dsRNA) precursors. These sRNAs are loaded into RNA-induced silencing complexes, which use sequence complementarity to target nucleic acid sequences for transcriptional or post-transcriptional gene silencing (Martinez et al., 2002; Schwarz et al., 2003). Among the proteins responsible for processing sRNAs from dsRNAs, Dicer (Dcr) plays a central role, particularly in the small interfering RNA (siRNA) and microRNA (miRNA) pathways.

Numerous studies have elucidated the functions of Dcr domains in recognizing and processing their canonical substrates, including double-stranded RNAs and single-stranded hairpin precursors of miRNAs (pre-miRNAs). Structurally, Dcr proteins exhibit an L-shaped, modular, multidomain architecture that can be broadly divided into three regions along their three-dimensional organization: a base, a core (body), and a head. These regions reflect both domain composition and spatial-functional arrangement (Lau et al., 2009, 2012; Wang et al., 2009). The base (N-terminal region) contains the helicase domain, comprised of three subdomains: DExD/H-box, RESIII, and helicase C-terminal domains. These serve as the entry point for dsRNA substrates, recognizing and binding them before unwinding the RNA duplex to ensure proper positioning. Evidence shows that in some species and Dcr variants the helicase domain may or may not be involved in ATP hydrolysis, a function that has been related to recognition and processing of blunt-ended dsRNAs such as viral RNAs or endogenous dsRNAs (Sinha et al., 2015, 2018; Welker et al., 2011). The core (central region) includes the domain of unknown function (DUF283) and double-stranded RNA binding domain (dsRBD). DUF283 functions primarily as a protein–protein interaction module, selectively interacting with dsRBD-containing cofactors such as DRB4 and HYL1 in Arabidopsis (Hiraguri et al., 2005; H. Qin et al., 2010), or proteins such as ADAR1 in mammals (Ota et al., 2013). The dsRBD enhances the stability and specificity of enzyme-substrate interaction contributing to the fidelity of the cleavage reaction (Zhang, 2002). At the catalytic core, the tandem RNase IIIa and IIIb domains form a dimer that cleaves both strands of the dsRNA duplex. The head (C-terminal region) contains the Piwi/Argonaute/Zwille (PAZ) domain, which binds the 3' overhangs of dsRNA substrates, acting as an anchoring point that guides the cleavage site (Ma et al., 2004). In the three-dimensional structure of Dcr, the PAZ domain is connected to the RNase III catalytic center

by a connector α-helix. This helix acts as a molecular ruler, setting the distance between the
RNA-binding PAZ domain and the RNase III domains to define the length of the sRNAs
produced (~21–25 nt) (MacRae et al., 2006).

While this canonical domain architecture is conserved across most of the Dcr proteins in
animals and plants, reduced Dcr proteins have also been described. Remarkably, the Dcr
protein from the protist *Giardia intestinalis* consists solely of the tandem RNase III domains
connected through a connector helix to a platform region, yet remains fully competent in
cleaving dsRNA substrates (MacRae et al., 2006). Interestingly, many Dcr proteins in fungi
have been reported to lack identifiable Helicase, PAZ and/or dsRBD domains, suggesting a
widespread presence of non-canonical Dcrs (Choi et al., 2014; Drinnenberg et al., 2009; Hu
et al., 2013). This diversity positions fungal Dcrs as a valuable model for exploring how
ancestral RNase III enzymes evolved into the complex, multidomain Dcr proteins found in
higher eukaryotes.

Fungi comprise a large and diverse group of eukaryotic organisms, currently classified into at
least 12 major phyla (Li et al., 2021), with some recent taxonomic studies recognizing up to
18 or 19 distinct phyla based on phylogenomic evidence (James et al., 2020; Tedersoo et al.,
2018; Wijayawardene et al., 2024). Most described fungal species to date belong to the
higher fungi Ascomycota and Basidiomycota, which together form the clade Dikarya. Other
fungal phyla (e.g., Mucoromycota, Chytridiomycota, Zoopagomycota) represent a smaller
portion of known fungal diversity, though they are ecologically important and likely under-
sampled (James et al., 2020).

Multiple approaches have been used to identify RNAi-related proteins in fungi, ranging from
focused studies in groups of species (Drinnenberg et al., 2009; Hu et al., 2013; Huang, 2018)
to large-scale comparative surveys such as the FunRNA database, which catalogued RNAi
components across 131 fungal genomes (Choi et al., 2014). These studies have shown that
the number and architecture of Dcr proteins vary significantly across fungal species. While
many fungi encode two Dcr paralogs (Catalanotto et al., 2004; Kadotani et al., 2004), these
can differ in sequence and domain organization, which may reflect functional specialization
(Chang et al., 2012; Drinnenberg et al., 2009). However, there are also fungal lineages
where only a single Dcr has been identified (De Haro et al., 2009; Drinnenberg et al., 2009)
and others, including members of the Saccharomycotina and some Basidiomycota, where
Dcr genes appear to be absent altogether (Drinnenberg et al., 2009; Hu et al., 2013; Choi et
al., 2014). Notably, the apparent absence of Dcr domains, such as PAZ or dsRBD in some
fungal proteins may result from limitations in domain annotation pipelines rather than true
biological loss. For example, domains like PAZ may be highly divergent in fungi, and thus

escape identification by current probabilistic models based on protein alignments of animal and plant-derived PAZ domains (Kress et al., 2023). However, albeit divergent, PAZ-like folds have been reported in fungal Dcrs such as Dicer from *S. pombe* (Zapletal et al., 2023), hinting at tridimensional structure conservation, and at the importance of incorporating structural analyses to generate hypotheses on fungal Dcr function.

The number of publicly available annotated and reference genomes has expanded more than tenfold in the last years, presenting a prime opportunity to study Dcr diversity at a wide phylogenetic scale. Furthermore, integration of structure prediction methods like AlphaFold (Jumper et al., 2021) can help us determine whether structural conservation is widespread across fungal Dcrs despite apparent sequence diversity.

In this study, we present a comprehensive analysis of Dcr proteins across fungi, revealing widespread non-canonical architectures that often lack identifiable PAZ, Helicase, or dsRBD domains. Through a combination of phylogenetic analyses, structural modeling, and molecular docking predictions, we show that many of these divergent Dcrs retain key features for RNA processing, including structurally conserved PAZ-like regions. Our findings expand the understanding of Dcr diversity in fungi and suggest that RNAi functionality can be maintained despite extensive domain loss or divergence.

Results

Comprehensive identification of Dcr proteins in fungi

To comprehensively catalog fungal Dcr proteins, we retrieved all available fungal reference proteomes from the NCBI, obtaining 1,593 datasets representing 1,427 unique species (Supplementary Table 1). The selected proteomes were predicted from genomes labeled as 'reference' or 'annotated' in NCBI, excluding those classified as atypical. These genomes meet the criteria for reference-quality assemblies. Most of the selected proteomes exhibit high completeness, with 86.4% scoring over 90% in BUSCO completeness based on either the fungi-wide or phylum-specific datasets (Fig. 1A, Supplementary Table 1). The proteomes span a wide range of fungal organisms with a marked representation from the phyla Ascomycota (913 species, 64.1%) and Basidiomycota (333 species, 23.4%). However, non-Dikarya fungi also constitute a significant proportion of the proteomes, with over 172 species from Mucoromycota, Zoopagomycota, Chytridiomycota, Microsporidia, Blastocladiomycota, and Olpidiomycota (Supplementary Table 1). This dataset expands the taxonomic scope for the identification and comparative analysis of fungal Dcr proteins, as compared with previous studies (Huang, 2018; Hu et al., 2013; Choi et al., 2014).

To identify putative Dcr proteins, we screened the proteomes for canonical Dcr-associated
domains previously described in plants and animals. Domain searches were performed using
HMMER, following methodologies established for Dcr identification in fungi (Choi et al., 2014)
and other organisms (Mukherjee et al., 2013). A protein was classified as a putative Dcr if it
met two criteria: (1) the presence of two, tandem RNase III domains, essential for dsRNA
cleavage, and (2) at least one RNA-binding domain, including Helicase subdomains DExD/H-
box, ResIII or HeliC, DUF283, PAZ, or dsRBD. This approach identified 2,917 putative Dcr
proteins across 1,215 species from eight fungal phyla (Supplementary Table 2). To validate
these predictions, we used an independent domain annotation approach using InterProScan,
which showed 100% concordance with the HMMER-based results (Supplementary Table 2).

Interestingly, we found no evidence of Dcr proteins in 12% of BUSCO high-completeness
proteomes and 11% of BUSCO low-completeness proteomes, totaling 227 species (254
proteomes). This suggests that Dcr proteins may have been lost in up to 15% of the
surveyed fungal species. However, in proteomes with low completeness scores the absence
of Dcr should be interpreted with caution, as it may reflect incomplete gene models rather
than true gene loss. Among the proteomes with no detectable Dcrs we identified 127
Ascomycota species, of which 115 (91%) belong to the subphylum Saccharomycotina,
including members from the classes Dipodascomycetes, Saccharomycetes, Pichiomycetes
and Trigonopsidomycetes (Supplementary Table 3). This subphylum encompasses most
Ascomycete yeasts, several of which have previously been reported to lack a functional
RNAi machinery (Drinnenberg et al., 2009). In addition, species lacking Dcrs were also found
in Basidiomycota, Olpidiomycota, Blastocladiomycota, Chytridiomycota, Microsporidia,
Mucoromycota, and Zoopagomycota (Supplementary Table 3), suggesting that Dcr protein
loss has occurred multiple times independently across fungal lineages.

Widespread domain architecture variation and predominance of non-canonical Dcr types in
fungi

To characterize the domain composition of fungal Dcrs, we referred to the canonical domains
typically found in Dcr proteins (Fig. 1B). Using these domain features as a guide, we
classified the 2,917 identified Dcr proteins into six categories based on the presence or
absence of domains detected through HMMER analysis. The types of domains found across
the fungal dataset is summarized in Fig. 1C, revealing the predominance of non-canonical
forms and the high frequency of Dcr proteins where the PAZ and/or dsRBD domains could
not be detected. Proteins containing all expected domains were classified as canonical Dcrs,
representing a small subset of the dataset (35 proteins, 1.2%). Proteins with one or more

missing domains were designated as non-canonical Dcrs and further grouped according to
the specific patterns of domain loss. Non-canonical A includes proteins lacking only the
dsRBD domain (80 proteins, 2.7%), while non-canonical B consists of those missing the PAZ
domain but retaining all other domains (357 proteins, 12.2%). Non-canonical C comprises
proteins in which Helicase subdomains and dsRBD are absent (274 proteins, 9.4%). The
most abundant group, non-canonical D, consists of proteins lacking both the PAZ and
dsRBD domains (1,827 proteins, 62.6%). Finally, non-canonical E includes proteins in which
three or more domains could not be detected (344 proteins, 11.8%).

The PAZ domain, central to RNA binding and precise determination of sRNA length, was
undetectable in the vast majority of Dcr proteins, with 86.7% lacking a recognizable PAZ
domain. However, protein length analysis revealed that proteins without a detectable PAZ,
particularly those in the non-canonical B and D categories, are comparable in size to
canonical and non-canonical A proteins that do contain this domain (Fig. 1D). This suggests
that the PAZ domain may still be present but has undergone substantial sequence
divergence relative to the canonical PAZ (cPAZ) described in plants and animals, rendering it
undetectable by standard HMM-based methods. Consistent with this idea, the current PAZ
HMM matrix used in this analysis (PF02170/IPR3100) was constructed from Dcrs and
ARGONAUTE proteins mainly from plants and animals, with negligible representation of
fungal Dcrs. Similarly, the dsRBD domain, also involved in RNA binding, was frequently
undetected, with 84.7% of proteins missing this domain. Interestingly, proteins in the non-
canonical C and E categories, which often lack subdomains of the Helicase region as well as
the dsRBD domain, exhibited shorter lengths, suggesting that the absence of detectable
domains in these groups may in some cases result from protein truncation rather than
sequence divergence.

The distribution of Dcr categories varies substantially among fungal phyla (Fig. 1E).
Canonical Dcrs and non-canonical A proteins, both of which retain a cPAZ domain, are
enriched in Mucoromycota, a phylum composed of mychorrizal symbionts, plant and animal
pathogens, fungal parasites and decomposers of organic matter (Zhao et al., 2023). In
contrast, non-canonical C proteins, which also retain a cPAZ domain, were primarily found in
Basidiomycota. Non-canonical B, D, and E proteins, lacking a cPAZ domain, are most
prevalent in Ascomycota, with some also found in Basidiomycota. The number of dcr genes
per genome also varied by phylum; most Ascomycota species have two dcr genes (683
species, 68%) while Basidiomycota more frequently possess three (187 species, 49.5%).
Strikingly, Microsporidia, a group of obligate intracellular parasites, either lacked dcrs entirely
(31 species, 70.5%) or contained only a single dcr (12 species, 27.3%) (Fig. 1F), consistent
with previously described reductions in gene contents in this lineage (Jespersen et al., 2022).

We also analyzed the distribution of Dcr category combinations across fungal species (Fig.
1G). Most species contain non-canonical D Dcrs as the sole Dcr type (473 species), or in
combination with non-canonical B (221 species), C (189 species), or E (150 species).
Interestingly, canonical Dcrs were either found alone or in combination with non-canonical A
or C, both of which retain a cPAZ domain. These patterns underscore the diversity in both
the number and types of Dcr proteins across fungal taxa, suggesting lineage-specific
adaptations in the RNAi machinery, likely in response to differing ecological and/or
evolutionary pressures.

Clade-specific distribution of canonical and non-canonical Dcrs suggests independent
diversification of RNAi pathways in fungi

To investigate the phylogenetic relationships of fungal Dcr proteins, we constructed a
sequence similarity network, considering a cut-off of 80% coverage and 50% identity
between connected nodes. 2,636 Dcr proteins met these criteria and were clustered into 149
groups (Supplementary Table 2). Multiple sequence alignments were generated for each
cluster and integrated into a global alignment, which was used to construct a maximum-
likelihood phylogenetic tree, including outgroup sequences from plants and metazoans (Fig.
2A). The analysis revealed a set of well-supported clades that we defined as major Dcr
groups. These groups correspond to fungal taxonomic lineages, with Ascomycota sequences
separating into two major clades (As1 and As2) and Basidiomycota sequences into three
distinct clades (Ba1, Ba2, Ba3). In contrast, sequences from non-Dikarya fungi formed more
scattered groups, forming independent clades (Mu1, Mu2, Mi1, Bl1, Ch1), suggesting that
Dcr diversification in non-Dikarya followed an independent evolutionary trajectory. The
outgroup sequences formed a distinct branch, supporting the overall topology and
robustness of the tree (Fig. 2A).

To evaluate whether clade assignments relate to previously reported Dcr annotations, we
mapped proteins annotated as Dcrs in UniProtKB onto the phylogenetic tree (Fig. 2B). These
include proteins labeled as Dcr1, Dcr2, Dcr-like1 (Dcl1), Dcr-like2 (Dcl2), as well as others
annotated more generically as "Dicer" or "Dicer-like". We found that proteins annotated as
Dcr1 or Dcl1 are predominantly located within the Ascomycota As2 clade and the
Basidiomycota clades Ba2 and Ba3. Dcr1/Dcl1 proteins are also observed in the Mu1 and
Mu2 clades, as well as some identified as Dcr2/Dcl2 in Mu2. These ancient duplications
forming Ba2/3 and Mu1/2 point to an unclear resolution of the ancestral proteins and possible
paraphyletic naming. However, these are the exception. In general, Dcrs with similar
annotations tend to group within the same clades, suggesting that existing naming

conventions reflect underlying evolutionary relationships, even if their functional roles have
not been fully confirmed. Notably, most of the Dcrs identified in our analysis remain
unannotated in public databases (Fig. 2B, Supplementary Table 2), underscoring the
extensive under-representation of fungal Dcr diversity and abundance.

To further explore the relationship between phylogeny and domain architecture, we mapped
canonical and non-canonical classifications onto the phylogenetic tree. Canonical Dcrs
clustered primarily within the Mu1 clade, and to a lesser extent, in the Mu2 clade, consistent
with their enrichment in the Mucoromycota phylum (Fig. 1E, Fig. 2C). Non-canonical A and C
proteins, which, as canonical Dcrs, retain a cPAZ domain (Fig. 1C, Fig. 2D), also show a
restricted distribution, with non-canonical C primarily associated with the Ba2 clade and non-
canonical A with Ba3. In contrast, non-canonical B, D and E proteins, which lack a cPAZ
domain (Fig. 2D), were widely distributed across the tree (Fig. 2C). The restricted
phylogenetic distribution of cPAZ-containing Dcrs suggests this domain is not broadly
conserved across fungal lineages; rather, its retention appears to be lineage-specific and
may reflect convergent retention or loss events in different evolutionary contexts.

The helicase subdomains DExD/H-box, ResIII, and HeliC also showed differential
representation across fungal clades, although most Dcrs retained Helicases with all three
subdomains (Fig. 1C, Supplementary Fig. 1A). Notably, Dcrs lacking all helicase
components, or retaining only the HeliC subdomain, were mainly found in the Ba2 clade,
suggesting that Basidiomycota harbor both helicase-complete Dcrs (e.g., in Ba1 and Ba3)
and more divergent forms potentially lacking ATPase activity (Ba2). A similar pattern was
observed in Mucoromycota, with Mu1 Dcrs containing complete helicases and Mu2 Dcrs
showing partial loss of subdomains. Additional incomplete helicase architectures were also
found in the Microsporidian clade Mi1. In contrast, the dsRBD domain, another typical
component of canonical Dcrs, but rarely detected in our analysis, was present in proteins
found in diverse clades across the tree (Supplementary Fig. 1B). These results underscore
the evolutionary plasticity of fungal Dcrs, suggesting that distinct selective pressures have
driven domain loss, retention, or divergence across fungal lineages.

Fungal Dcrs have conserved structural features despite sequence divergence

The sequence similarity network and phylogenetic analyses suggest a high degree of
sequence divergence among fungal Dcr proteins, with several instances in which key
domains typically associated with Dcr function were not detected, likely due to either their
absence or low sequence conservation. To explore whether these proteins might

nonetheless retain conserved three-dimensional features, we generated structural predictions for all Dcr proteins using AlphaFold (Jumper et al., 2021). Protein models displaying a predicted Local Distance Difference Test (pLDDT) score greater than or equal to 70 were selected, and a structural similarity network was constructed, considering a template modeling (TM) score of ≥ 0.9 between connected nodes as a threshold. Among the selected models, the average pLDDT score was 75, supporting the overall reliability of the structural predictions. A total of 2,475 proteins met this criterion and were grouped into 26 clusters (Supplementary Table 2). Strikingly, nearly 95% of phylogenetically analyzed proteins are found within just four major structural clusters (Fig. 3A), suggesting core structural features are widely conserved. Each cluster showed a distinct representation of fungal phyla (Fig. 3A, B). Clusters 1 and 2 contain the majority of Dcr proteins, with cluster 1 including predominantly Ascomycota sequences and a smaller subset from Basidiomycota. This cluster harbors Dcrs with experimentally validated roles in sRNA processing such as *Botrytis cinerea* Dcr2 (Qin et al., 2023; Wang et al., 2016; Weiberg et al., 2013), *Aspergillus fumigatus* Dcr2 (Kelani et al., 2023) or *Trichoderma atroviride* Dcr2 (Carreras-Villaseñor et al., 2013; Enriquez-Felix et al., 2024; Villalobos-Escobedo et al., 2022). Cluster 2 is predominantly composed of Basidiomycota sequences, and includes *A. fumigatus* Dcr1, *T. atroviride* Dcr1 and *Neurospora crassa* Dcr1 (Lee et al., 2010). Clusters 3 and 4 include fewer proteins, mainly from Basidiomycota and Mucoromycota (cluster 3) or exclusively from Basidiomycota (cluster 4). The remaining smaller clusters consist of Ascomycota and Basidiomycota Dcrs and may represent functionally divergent paralogs present in these phyla (Fig. 3A, B).

The four major structural clades aligned well with the phylogenetic clades identified in our sequence-based analysis (Fig. 3C), indicating that despite extensive sequence divergence and domain variability, fungal Dcrs retain conserved structural architectures within evolutionary lineages. Cluster 1 contains sequences from clades As1 and Ba1, cluster 2 from As2, and cluster 3 from Ba2 and Mu1. Cluster 4 primarily contains Ba3, indicating that this is a unique lineage in Basidiomycota. Importantly, while proteins annotated as Dcr2 in UniProtKB majorly fall into Cluster 1, proteins annotated as Dcr1 are located into clusters 2, 3 and 4, indicating greater structural and likely functional diversity among Dcr1-labeled proteins. These findings underscore the need for systematic re-evaluation of Dcr annotations grounded in structural and evolutionary evidence rather than existing paralog labels.

The other cluster proteins (those falling outside the four major structural groups) may represent highly divergent Dcr variants, truncated sequences, or lineage-specific proteins with distinct structural features that deviate from the dominant structural folds. This group

contains members from several phylogenetic clades, particularly non-dikarya fungi, including

Mucoromycota and Zoopagomycota Dcrs from clade Mu2 (Fig. 2B, 3C). It also includes all

Dcrs from Blastocladiomycota clade Bl1, Miscrosporidia clade Mi1 and nearly all from

Chytridiomycota clade Ch1, indicating substantial structural divergence in Dcr proteins in

these non-Dikarya fungal lineages.

To further explore folding patterns, we examined the structure of the most connected

(central) node within each major structural cluster (Fig. 3B). These representative proteins

span different Dcr categories, with cluster 1 represented by a non-canonical D protein from

Aureobasidium namibiae (clade As1), cluster 2 represented by *Li*Dcl1, a non-canonical E

protein from *Lobaria immixta* (clade As2), cluster 3 represented by a non-canonical C protein

from *Claroideoglomus candidum* (clade Mu1), and cluster 4 represented by *Sc*Dcl1, a non-

canonical D protein from *Schizophyllum commune* (clade Ba3). Despite differences in

domain composition detected by HMMER analysis, AlphaFold models of these proteins show

broadly similar L-shaped Dcr architectures, comprising head, core and base regions.

Importantly, the structural models revealed PAZ-like folds in the head region of

representatives from clusters 1,2 and 4, even though cPAZ domains were not detectable at

the sequence level in these proteins. This suggests that the RNA-binding functionality

associated with PAZ might be structurally preserved in Dcrs despite substantial divergence,

particularly in the more widely distributed non-canonical Dcr types.

Ancestral state reconstruction supports an early origin and selective retention of the cPAZ

domain in Fungi

To gain deeper insights into the evolutionary history of the PAZ domain in fungi, we

performed an ancestral character estimation (ACE) analysis to determine whether the cPAZ

domain was present in the last common ancestor of fungal Dcrs (Fig. 4A). The ACE analysis

was based on the presence or absence of the cPAZ domain, mapped onto the Dcr

phylogeny. The reconstruction suggests that the common ancestor of fungal Dcrs most likely

harbored a cPAZ domain, supporting the hypothesis that this domain is an ancestral feature

of fungal Dcrs. Despite its ancestral origin, the phylogenetic distribution of cPAZ shows

strong lineage-specific conservation patterns (Fig. 4A), consistent with those observed in the

phylogenetic analysis (Fig. 2D) and structural clustering (Fig. 3A, C). The presence of the

cPAZ domain in these clades may reflect convergent retention under shared selective

pressures, or alternatively, could be the result of horizontal gene transfer (HGT) from

lineages such as plants or animals, where the cPAZ domain is nearly ubiquitous. Indeed,

nearly all Dcr proteins in metazoans (99.9% of Dcrs) and plants (100% of Dcrs) contain a
cPAZ domain in contrast to only 12.4% of fungal Dcrs (Fig. 4B). Interestingly, fungi are not
the only organisms lacking detectable cPAZ domains, with only 33.3% one of three protist
Dcrs possessing a cPAZ (Fig. 4B). This suggest that domain loss or divergence is not
exclusive to fungi. We evaluated the possibility of Dcr HGT from plants or animals to fungi,
however our analysis did not find evidence supporting these events (Fig. 4C). These findings
support a model in which the cPAZ was selectively retained in specific fungal lineages, while
its apparent absence in most fungal Dcrs is most likely due to widespread sequence
divergence rather than true domain loss. This interpretation is supported by the similar
protein lengths observed between Dcrs with and without a cPAZ (Fig. 1D) and by the
structural predictions showing a putative conserved PAZ-like fold in proteins lacking a
detectable cPAZ (Fig. 3C).

A sequence-divergent yet structurally conserved PAZ domain defines fungal Dcr diversity

We explored the possibility of a diverged fungal PAZ (fPAZ) in Dcr proteins lacking a
detectable cPAZ. To address this idea, we compared the sequence and structural features of
the cPAZ, present in canonical and non-canonical A and C Dcrs, with those of the putative
fPAZ, which should be found in non-canonical B, D and E proteins. In proteins without a
cPAZ, the fPAZ was defined as the region between the DUF and RNAse III domains. Using
MEME (Bailey et al., 2015), we identified four conserved motifs shared across both cPAZ
and fPAZ domains. These motifs, designated Logo-1 through Logo-4, represent conserved
sequence elements ordered from N- to C-terminal within the PAZ region (Fig. 5A). The logos
revealed moderate to high positional conservation of aminoacidic residues across the motifs.
Logo-1 displays highly conserved arginine (R) residues, and a conserved phenylalanine (F)
at position 8. These residues reflect the characteristic composition of the 3'-end binding
pockets in PAZ domains, where aromatic residues enable base stacking, and positive
residues contribute to electrostatic interactions with the RNA (Yan et al., 2003; Ma et al.,
2004). Logo-2, similar to Logo-1, contains conserved F and tyrosine (Y) residues, followed
by a stretch of basic residues, further supporting a conserved RNA-binding role. Logo-3
features several polar and charged residues, including K, D, N and Q, which are likely
involved in maintaining the OB-fold structure of the PAZ domain via hydrogen bonding and
overall stabilization. The presence of conserved K and N residues in the central region may
also contribute to interactions with the phosphate in the RNA. Logos 1 to 3 were strongly
associated with the cPAZ, being identified in over 80% of cPAZ-containing proteins, but in
fewer than 20% of the sequences containing only an fPAZ. In contrast, Logo-4 was more

broadly conserved across fungal Dcrs. It was present in 99.67% of the cPAZ sequences and
in 56.71% of the fPAZ sequences. This motif is enriched in hydrophobic amino acids (F, I, P,
A, L, M), consistent with a helical or turn-forming structure. Given its composition, Logo-4
likely localizes to the connector helix that links the PAZ to the first RNAse III domain, a region
well-known to be essential for maintaining the spatial configuration required for precise RNA
cleavage by Dcr (Tian et al., 2014). Mapping of these four motifs onto the three-dimensional
structure of a representative canonical PAZ domain confirmed their spatial clustering at the
RNA-binding interface (Supplementary Fig. 2). In particular, the motifs localize around the
region that interact with the 3′ end of the dsRNA, supporting their proposed role in RNA
anchoring and stabilization. The spatial proximity of Logo-1 and Logo-2 to the RNA entry
groove aligns with their enrichment in basic and aromatic residues, while Logo-4 occupies a
linker region adjacent to the RNase III domain, consistent with a role in domain coordination
and structural integrity. This structural mapping provides further support for the functional
relevance of these motifs in mediating RNA binding across diverse fungal Dcrs.

Next, we assessed the structural similarity of the cPAZ and divergent fPAZ domains. As a
reference, we used the cPAZ domain from a representative canonical Dcr (Dcl1 from *Absidia*
glauca). Structural comparisons revealed a high degree of similarity between this reference
and the cPAZ domains of non-canonical Dcrs A (Dcl1 from *Entomophthora muscae,* TM-
score of 0.821) and C (Dcl1 from *Dissophora ornata,* TM-score of 0.828). Although lower,
structural similarity scores between the reference cPAZ and the putative fPAZ domains found
in non-canonical Dcrs B, D and E (TM-scores of 0.678, 0.673 and 0.683, respectively), still
indicate substantial structural conservation. This pattern of conserved fold architecture
extends across the broader set of structural models analyzed (Fig. 5C).

Considering the high sequence divergence of fungal PAZ domains, and the under-
representation of fungal Dcrs in the current PAZ HMM model, we generated a custom HMM-
profile (FPAZ-HMM) based on a multiple sequence alignment of both cPAZ and structurally
inferred fPAZ domains from our dataset. Importantly, FPAZ-HMM was specifically trained to
recognize PAZ domains associated with Dicer proteins and does not detect PAZ domains
from Argonaute (AGO) proteins, as these were intentionally excluded from the alignment
used for model construction. In contrast, current general PAZ HMMs typically include PAZ
domains from both Dcr and AGO proteins.

We used this profile to search for PAZ domains in Dcr-annotated proteins in UniProtKB.
Compared to the current PAZ HMM, the FPAZ-HMM profile demonstrates markedly
improved sensitivity for detecting PAZ domains in fungal Dcrs, while maintaining comparable
performance in Metazoa and Plantae (Fig. 5D). These results support using of FPAZ-HMM

as a robust and lineage-inclusive tool for detecting Dcr PAZ domains across eukaryotes and

underscores the extensive sequence variability of PAZ domains across fungal clades.

Structural and functional evidence support conserved RNA-binding activity of fungal PAZ

domains

To gain insights into the functionality of the canonical and divergent fungal PAZ domains, we

analyzed structural models of full-length canonical and non-canonical Dcrs predicted in

complex with a pre-cleaved dsRNA substrate that was obtained from the PDB model of *D.*

melanogaster Dicer (pdb_00007w0f) (Figure 6A, B). In these models, the dsRNA (green)

interacts with the Dcr surface (grey), with key RNA-interacting residues highlighted in blue.

In Dcrs with cPAZ, such as those from *Puccinia striiformis, Cryptococcus neoformans* and

Absidia glauca, the PAZ domain and surrounding regions form a well-defined RNA-binding

pocket. Basic residues, particularly K and R were consistently positioned within 3 Å of the

dsRNA, forming a positively charged cleft that anchors the 3' end of the RNA duplex (Fig. 6A,

right). This configuration mirrors previously described PAZ-RNA interactions in plant and

animal systems.

Remarkably, similar interaction interfaces were observed in non-canonical Dcrs lacking a

canonical PAZ domain (cPAZ), but containing a structurally defined fungal-specific PAZ

(fPAZ), including Dcl1 and Dcl2 from *Botrytis cinerea* and the Dcr from *Cryptococcus*

neoformans. These proteins display conserved K and R residues positioned in close

proximity to the RNA substrate, supporting the idea that key electrostatic and structural

features necessary for RNA anchoring are retained, even in the absence of canonical

sequence motifs. To assess the conformational stability of these interactions, we conducted

molecular dynamics simulations on a representative set of Dcr–RNA complexes. Across all

variants analyzed—including canonical, non-canonical C, and fPAZ-containing non-canonical

B and D— RMSD trajectories stabilized within the first 10–20 nanoseconds, indicating that

the RNA-binding interface, particularly the PAZ or PAZ-like domain, maintains structural

integrity throughout the 50-nanosecond simulations (Supplementary Fig. 3). These results

support the functional relevance of the fPAZ fold as a stable RNA-interacting module across

divergent fungal Dcrs.

To further explore the functional potential of the fungal PAZ regions, we analyzed the

electrostatic surface distribution of representative Dcr proteins for which a function in sRNA

biogenesis has been evaluated using mutants and sRNA sequencing. These include Dcl1

and Dcl2 from *B. cinerea*, Dcr1 from *S. pombe*, Dcr1 from *S. japonicus*, and Dcl1 and Dcl2

from *P. oryzae* (Fig. 6C). In these non-canonical Dcrs, the dsRNA-binding pocket displayed a predominance of positively charged residues, consistent with a role in RNA anchoring (Fig. 6C). Additionally, we measured the distance between the fPAZ domain and the RNAse III catalytic center, finding it to be approximately 50-55 Å, comparable to that reported in human Dicer (Lau et al., 2012). These structural configurations support the model in which PAZ domains act as a molecular ruler, securing the 3'end of the RNA and enabling precise cleavage by the RNAse III domains.

Finally, we compared sRNA length profiles from wild-type and mutant strains with disrupted dcr genes (Fig. 6D). In all cases examined, dcr mutation resulted in clear losses of distinct sRNA size classes, as indicated by the disappearance of specific peaks. These results confirm that non-canonical Dcrs contribute to sRNA biogenesis and strongly suggest that the fPAZ domain plays a critical role in RNA processing, despite its sequence divergence from canonical PAZ domains.

Discussion

Understanding the structural diversity of Dcr proteins in fungi is essential, as these proteins are central components of RNAi, a mechanism critical for gene regulation, genome stability, defense and inter-species interactions. The dataset presented in this study represents, to our knowledge, the most extensive survey of fungal proteomes to date derived from annotated and reference-quality genomes, comprising 1,593 proteomes from 1,427 species across eight phyla and 44 taxonomic classes. While thousands of fungal genomes are publicly available, many are incomplete or lack standardized annotations, which limits their utility for comparative analyses of protein-coding genes such as Dcrs. Notably, our sampling includes a broader representation of non-Dikarya fungi compared to previous studies (Choi et al., 2014; Q. Huang, 2018), particularly within Mucoromycota and Microsporidia. This improvement reflects the increasing contribution of large-scale sequencing efforts such as the Joint Genome Institute's 1000 Fungal Genomes Project (https://mycocosm.jgi.doe.gov/mycocosm/home/1000-fungal-genomes). Despite this broader coverage, several phyla are still poorly represented in our dataset, including Zoopagomycota (20 proteomes), Chytridiomycota (19 proteomes), Blastocladiomycota (2 proteomes), and Olpidiomycota (1 proteome), while other phyla are entirely absent. These limitations highlight the continued need to incorporate non-Dikarya and ecologically diverse fungal lineages into genome sequencing initiatives, to enable a more complete understanding of fungal molecular diversity.

Our survey of fungal Dcrs identified proteins broadly across fungal clades in the dataset,
confirming that this machinery is widely retained across fungi, as others have found (Choi et
al., 2014). Nevertheless, we found a relevant part of fungal species (~10%) with no Dcr
identified in their proteomes, according to our pipeline. While the loss of Dcrs is well
established in budding yeasts within the Saccharomycotina and some Basidiomycota
lineages (Dang et al., 2011; Drinnenberg et al., 2009), broader surveys have shown that
such losses may also occur sporadically across the fungal kingdom (Billmyre et al., 2013;
Chang et al., 2012). However, the extent of Dcr loss in other fungal groups remains less
clear, largely due to the underrepresentation of non-Dikarya fungi in genomic datasets. In our
analysis, most proteomes lacking detectable Dcrs belonged to Saccharomycotina, consistent
with previous reports. However, we also identified apparent Dcr losses in non-Dikarya phyla.
While these absences require further validation, they raise the possibility that RNAi pathways
in fungi have evolved alternative mechanisms that operate independently of Dcr proteins.
This contrasts with animal and plant systems, were the loss of Dcr typically results in severe
developmental defects or lethality (Bernstein et al., 2003; Gasciolli et al., 2005) and suggests
that fungi might possess a more flexible RNA silencing machinery. Consistent with this idea,
evidence of Dcr-independent sRNA-processing by other members of the RNAseIII family has
been reported in fungi, as is the case of MRPL3 in *Neurospora crassa,* or R3B2 in *Mucor*
lusitanicus (Cánovas-Márquez et al., 2021; Lee et al., 2010)*.* Fungal genomes encode a
diverse array of RNases III which could play conceivable roles in RNAi. For example, our in-
depth analysis of RNAse III-containing proteins uncovered a small number of potential
DROSHA-like candidates. However, the general absence of the characteristic domain
architecture typically associated with DROSHA suggests that these proteins may be highly
divergent and thus require more targeted, in-depth analyses for definitive identification.

We found four major Dcr variants across fungi, but little is known about what the functional
characteristics of these groups are. Among Dikarya, Dcrs from multiple clusters are found in
organisms. This could relate to the radiation of RNAi functions in these organisms, as they
contain a diverse complement of Dcrs. This diversity may reflect evolutionary pressure to
fine-tune RNAi responses to different developmental or ecological contexts. In particular, the
presence of sequence-diverging PAZ domains seems to be widespread, at least in the
species represented in our dataset, with cPAZ domains limited number of species, mostly
within Mucoromycota and Basidiomycota. Considering that ACE analysis supports that cPAZ
was likely present in the last common ancestor of fungal Dcrs, this limited representation
implies lineage-specific retention, which could be attributed to selective pressures to maintain
a more precise or processive sRNA cleavage in these lineages. Both Mucoromycota and
Basidiomycota comprise species engaged in complex ecological interactions, including

mutualistic symbioses, such as Mycorrhizal associations, and pathogenic lifestyles. In these
contexts, sRNA-mediated communication between fungi and their plant hosts may play a
pivotal role in modulating host responses and facilitating successful colonization.

Interestingly, our phylogenetic analysis revealed that Dcr naming in databases have not been
consistently applied across fungal lineages. Proteins carrying common names (Dcr1/Dcl1 or
Dcr2/Dcl2) often appear in different clades, indicating that these labels do not accurately
reflect true evolutionary or functional relationships. For example, proteins annotated as
Dcr1/Dcl1 are distributed across Ba2 and Ba3 in Basidiomycota, which form separate
phylogenetic and structural clusters. Similar patterns are observed for other Dcr annotations.
These observations highlight that current Dcr naming conventions are largely disconnected
from the structural and phylogenetic diversity of fungal Dcr proteins, underscoring the need
for a more systematic, structure- and phylogeny-informed classification framework.

In fungi, the PAZ domain has traditionally been considered dispensable for Dcr function,
since various well-characterized fungal models, including *Schizosaccharomyces pombe,*
Neurospora crassa, *Trichoderma atroviride*, *Aspergillus fumigatus*, or *Botrytis cinerea,* harbor
non-canonical Dcrs that lack detectable PAZ domains but remain functional in sRNA
production (Carreras-Villaseñor et al., 2013; Drinnenberg et al., 2009; Enriquez-Felix et al.,
2024; Kelani et al., 2023; Lee et al., 2010; Villalobos-Escobedo et al., 2022; Weiberg et al.,
2013). However, our structural analyses challenge this notion. Many of these non-canonical
Dcrs retain a PAZ-like fold capable of anchoring dsRNA despite lacking sequence similarity
to canonical PAZ domains. This finding suggests that structural conservation, rather than
primary sequence similarity, may underline the preservation of PAZ functionality in fungi.

Detailed structural comparisons revealed strong conservation of R and K residues in the
binding pocket pointing to a conservation of the dsRNA-protein interface. However, we found
that the highly hydrophobic Logo-4 motif is much more conserved in fungal Dcrs. This hints
that the overall structure of the PAZ domain is more conserved among fungi, while the actual
binding interface could be more lineage specific (Logos-1, -2, -3). Consistent with previous
studies (Wei et al., 2021), we found that PAZ-like domains in fungal Dcrs conserve the R and
K residues critical for 3′ end binding. Amino acids like glutamic acid (E), R, and K are often
involved in ionic interactions and hydrogen bonding, essential for dsRNA binding. Y and F
contribute to hydrophobic interactions, stabilizing the RNA-protein complex. The identification
of these conserved residues and their role in ionic interactions, hydrogen bonding, and
hydrophobic stabilization strengthens the hypothesis that even non-canonical PAZ domains
can perform essential functions. Understanding these conserved motifs not only shed light on
the molecular mechanisms by which fungal Dcrs interact with dsRNA but also opens

avenues for functional validation and targeted manipulation. For instance, residues such as R and K, which mediate 3′ end binding, as well as other amino acids involved in electrostatic and hydrophobic interactions, are promising candidates for site-directed mutagenesis to assess their specific contribution to RNA binding and cleavage efficiency. These experiments would help clarify whether structurally conserved PAZ-like folds can fully compensate for the absence of canonical PAZ domains in non-canonical Dcrs. Moreover, identifying lineage-specific conservation patterns, such as the highly preserved hydrophobic Logo-4 motif, could inform the design of small molecule inhibitors or RNA mimetics that selectively disrupt dsRNA recognition in certain fungal groups. This may have practical implications for biotechnological or agricultural applications, such as developing of fungal RNAi-based tools for gene silencing or plant-pathogen interaction control. Ultimately, resolving how these motifs contribute to RNA processing in structurally divergent Dcrs will enhance our understanding of fungal RNAi flexibility and evolutionary innovation.

Our structural findings underscore a critical limitation of current protein annotation pipelines: the overreliance on sequence similarity to detect functional domains. Standard tools such as InterProScan or Pfam HMMs often fail to recognize divergent PAZ-like regions, leading to an underestimation of the true diversity of RNAi-related proteins in fungi. This bias not only affects gene annotations but also distorts evolutionary and functional interpretations. To provide an improved tool for PAZ detection in eukaryotes, we developed FPAZ-HMM, a custom profile trained on structurally validated PAZ-like domains from fungal Dcrs. Unlike conventional PAZ HMMs (PF02170), this model successfully detects both canonical and non-canonical PAZ folds, unveiling a broader and more realistic distribution of RNA-binding interfaces. Our findings align with recent evaluations showing that up to 50% of inferred domain losses in fungi may be artifactual, stemming from annotation inconsistencies or overly stringent cutoffs (Kress et al., 2023). Tools like FPAZ-HMM are thus essential to recover biologically meaningful domains that are missed by sequence-only approaches, particularly in non-model fungal lineages.

In sum, our work point to the necessity of a standardized classification system for Dcrs, one that incorporates domain architecture, structural similarity, and phylogenetic placement. Moving beyond binary presence/absence assessments toward integrated, multidimensional criteria will enhance cross-species comparability and improve functional predictions. This is especially critical in fungi, where domain plasticity and repeated loss events have shaped a highly heterogeneous Dcr landscape. Adopting such frameworks will be key to fully uncovering the evolutionary flexibility and functional innovation of fungal RNAi systems.

Conclusion

Our comprehensive survey of fungal Dcr proteins reveals a dynamic landscape shaped by recurrent domain losses, structural diversification, and lineage-specific retention of RNAi components. By integrating comparative genomics, structural modeling and molecular simulations, we show that many non-canonical Dcrs retain functionally conserved PAZ-like folds that are overlooked by sequence-based tools. These findings not only expand our understanding of RNAi evolution in fungi, but also underscore the need for structure-informed models and standardized nomenclature to better capture the functional diversity of this essential gene-silencing machinery.

Methods

Recovery and quality assessment of fungal proteomes

We retrieved a total of 1,593 annotated and reference fungal proteomes from the NCBI (data available up to November 2022), corresponding to 1,427 unique species and 172 duplicated species from different strains. We used the the Entrez esearch tool via its command line interface (Kans, 2013), with the query '"Fungi"[Organism] OR fungi[All Fields]) AND (fungi[filter] AND "representative genome"[filter] AND (all[filter] NOT anomalous[filter] AND all[filter] NOT partial[filter]) AND "has annotation"[Properties]', applying filters to select only representative genomes with available annotations, while excluding anomalous or partial assemblies.

We used BUSCO (Benchmarking Universal Single-Copy Ortholog assessment) (Seppey et al., 2019) v.5.3.2 to evaluate the completeness of proteomes. We first applied the universal_odb10 and fungi_odb10 dataset, followed by additional phylum-specific datasets when available, including asomcyota_odb10, basidiomycota_odb10, microsporidia_odb10, and mucoromycota_odb10. We retained all retrieved proteomes for posterior analyses, regardless of their completeness score in BUSCO. We made this decision to maximize the detection of Dcr proteins, ensuring that potentially relevant sequences from less complete proteomes were not excluded. For example, some fungal groups, such as Cryptomycota and Olpidiomycota, are known to include obligate parasites, which may require a specialized dataset to accurately assess completeness.

We labeled each proteome with its corresponding BUSCO completeness score, allowing its quality to be tracked throughout the subsequent analyses. This annotation ensured that for every Dcr protein identified in later steps, the completeness of its source proteome was

known. We retained this information to assess whether proteome completeness had any
impact on Dcr detection and to provide a transparent evaluation of the dataset composition.

Recovery of profile Hidden Markov Model and identification of putative Dcr proteins

We retrieved a set of profile Hidden Markov Models (HMMs) representing highly conserved
Dcr domains within eukaryotic species from the Interpro database (ebi.ac.uk/interpro). These
HMMs have been used in previous Dcr studies (Drinnenberg et al., 2009; Mukherjee et al.,
2013; Raman et al., 2017) and include PF00270/IPR011545 (representing the DEAD/DEAH
box helicase subdomain), PF04851/IPR006935 (representing ResIII helicase subdomain),
and PF00271/IPR001650 (representing Helicase C terminal helicase subdomain),
PF00636/IPR000999 (representing RNase III domain), PF02170/IPR003100 (representing
PAZ domain), PF03368/IPR005034 (representing DUF/Dicer dimerization domain), and
PF00035/IPR014720 (representing Double-stranded RNA binding motif domain). We
scanned the fungal proteomes the HMMER v.3.3.1 toolkit (Potter et al., 2018) to identify
putative Dcr proteins. The thresholds for detection of a domain were set at an E-value $\leq 10^{-3}$,
and only proteins containing at least one RNase III domain (PF00636/IPR000999) were
retained for further analysis.

InterProScan analysis

InterProScan v.5.44 (Jones et al., 2014) was used to scan protein sequences. All analyses
were conducted using default parameters.

In silico protein structure prediction

To predict the three-dimensional structures of the Dcr proteins, we used the local version of
Colabfold (Mirdita et al., 2022) v1.5.5, a Google Colab implementation of Alphafold2 (Jumper
et al., 2021). In the first step, multiple sequence alignments (MSA) were generated against
all available databases using the "DB load mode 3" setting, which preloads all sequence
databases into memory to speed up alignment searches. We executed this process using the
following command: colabfold_search --db-load-mode 3 --threads 54 dcr_sequences.fasta
database a3m_output. Subsequently, we predicted the protein structures using the same
software with the command: colabfold_batch --num-models 1 --num-relax 0 msas_input/

pdb_output/. To reduce computational time, we only generated one structural model per
protein, and no structural relaxation was performed. We selected protein models with a
predicted Local Distance Difference Test (pLDDT) score ≥70 for further analysis, ensuring
the inclusion of structures with high predicted accuracy for downstream applications.

Electrostatic Surface Calculations

To investigate the electrostatic surface distribution of selected Dicer proteins, we applied a
two-step computational workflow consisting of structure preparation and electrostatic
potential calculation.

Protein structure were preprocessed using PDB2PQR (Dolinsky et al., 2004), which assigns
atomic charges and radii according to the PARameters for Solvation Energy (PARSE) force
field and determines protonation states at pH 7.0 using PROPKA (Sondergaard et al., 2011;
Olsson et al., 2011). This preprocessing ensures consistent treatment of ionizable residues,
addition of hydrogens, and optimization of side-chain orientations for electrostatic
calculations.

Electrostatic potentials were then computed using the Adaptive Poisson–Boltzmann Solver
(APBS) (Baker et al., 2001). We performed the calculations using the linearized Poisson–
Boltzmann equation (LPBE), with the solute and solvent dielectric constants set to 2.0 and
78.54, respectively. We used a solvent probe radius of 1.4 Å, temperature of 298.15 K, and
ion-accessibility parameters (swin = 0.3 Å, sdens = 10.0). We set surface and charge
discretization methods to smol and spl2, respectively. We set the boundary condition to sdh
(single Debye–Hückel), and we automatically generated the electrostatic grid to encompass
each protein's structure with sufficient resolution.

Electrostatic potential maps were output in OpenDX format and used to visualize surface
charge distributions, with a focus on the double-stranded RNA binding regions. Total
electrostatic energies were also computed but not directly analyzed.

Generation of Protein Similarity Networks and Protein Structural Networks

We generated a sequence similarity network using the Sequence Similarity Network pipeline
v.1.0.0 (https://github.com/MiguelMSandin/SSNetworks). In brief, this pipeline performs a
local pairwise alignment of sequences using BLASTn to calculate similarities between all
sequences in the dataset. We processed the resulting pairwise similarities to remove
reciprocal hits and self-hits, ensuring a clean input for network construction. We generated

the network considering an identity threshold of 50% and a coverage of 80%. For the protein
structural network, we aligned the obtained protein structures in PDB format using mTM-align
(Dong et al., 2018). We generated the structural network considering TM-scores between
nodes of 90% or more. We used Cytoscape (Shannon et al., 2003) v.3.10.0 to visualize and
analyze the networks.

Phylogenetic tree generation

We performed multiple sequence alignments using T-Coffee (Notredame et al., 2000)
v13.45.0 with default parameters. Proteins corresponding to each of the 149 clusters
generated by the sequence similarity network analysis were aligned independently.
Subsequently, we carried out profile alignment for the 149 alignments to generate a global
alignment. For this, we used ClustalW (Thompson et al., 1994) v.2.1 with the -profile option.
The global protein alignments were trimmed using ClipKit (Steenwyk et al., 2020) v2.1.1 with
default parameters. We inferred the best-fit model and maximum-likelihood phylogenetic
trees from the trimmed sequence alignments using IQ-TREE (Minh et al., 2020) v2.2.5 with
the following parameters: -B 1000 -m MFP -nt AUTO.

Ancestral reconstruction

We used the fitmk function of the phytools (Revell, 2012) package to fit a discrete Markov
model for the ancestral reconstruction of the Dcr proteins, taking as input the phylogenetic
tree generated above. In this case, we used the "Equal Rates" (ER) model, which assumes
equal transition rates between states, with the following command: fitER =
fitMk(midroot_tree, PAZ_level, model = "ER"). The anc.recon function was used based on
the fitted model, and we performed ancestral reconstruction using the marginal method to
infer the most probable states at the ancestral nodes of the tree, using the command:
fit.marginal = ancr(fitER, type = "marginal").

Horizontal gene transfer analysis

To assess the potential occurrence of horizontal gene transfer (HGT) among fungal Dcrs, we
analyzed Dcr sequences using HGTector (Zhu et al., 2014) v2.0b3. The search module of
HGTector was executed with DIAMOND (Buchfink et al., 2015) against a predefined NR
reference database (40,310 genomes, retrieved in January 2023). We run the 'analyze'

command using default parameters, except for the --self parameter, which was set to 4751

(NCBI taxonomy ID for Fungi) to filter intra-fungal homologs.

Docking and molecular simulations

We performed docking of the Dcr-RNA complex using HDOCK (Yan et al., 2017) v1.0.

Protein models of Dcr, obtained from the previous ColabFold, were provided in PDB format,

and were used along with the dsRNA derived from the Drosophila Dicer2 PDB model

(7W0F). We performed *ab initio* docking, defining the residues in the receptor (Dcr) and

ligand (RNA) where preferential binding was expected.

For the molecular dynamics simulations, we used the CHARMM36 (J. Huang & MacKerell,

2013) force field to construct the protein structure file (PSF) for both protein and dsRNA. The

simulation workflow consisted of three stages. First, energy minimization was performed

using 20,000 steps to remove steric clashes and stabilize the initial system using a restriction

of 1 kcal/mol. Next, an NVT equilibration phase was conducted for 0.25 ns constraints

applied to the protein and dsRNA to maintain structural integrity during thermal adjustments

(1 kcal/mol). Finally, production simulations were carried out in two phases: 50 ns

simulations with dsRNA restrictions maintained, followed by an additional 50 ns simulations

with all constraints released to allow full system flexibility. Each molecular simulation was

performed in three independent replicates to assess the reproducibility and stability of the

results.

We generated all molecular dynamics steps using NAMD3 (Phillips et al., 2020). The

equations of motion were integrated with a 2-fs time step using the Verlet algorithm.

Langevin dynamics (damping coefficient of 1 ps) and the Nosé–Hoover Langevin piston

method was employed to maintain constant temperature (310 K) and pressure (1 atm). Long-

range electrostatic interactions were calculated using the Particle Mesh Ewald (PME)

method, and van der Waals forces were computed with a cutoff of 12 Å.

Motif discovery in protein sequences

We performed motif discovery using the Multiple Em for Motif Elicitation (MEME) tool v. 5.5.7

from the MEME suite (Bailey et al., 2015). We ran MEME locally with the options -protein -

mod zoops -nmotifs 4 -minw 6 -maxw 50. We used the obtained motifs to scan the

sequences with the Find Individual Motif Occurrences (FIMO) from the MEME suite. We ran

FIMO with default parameters.

Construction of a custom HMM profile for the PAZ domain

To generate a high-confidence HMM profile of the PAZ domain, we selected 2,452 PAZ-
containing regions from fungal Dicer proteins, including both canonical (cPAZ) and non-
canonical (fPAZ) representatives identified in our dataset. We first aligned sequences
separately using MAFFT (Katoh & Standley, 2013) v7.505 with the --auto option, visually
inspecting alignment quality to confirm the presence of informative blocks. Poorly aligned or
uninformative regions were trimmed using ClipKit v1.3.0 in kpi mode. We then merged the
resulting canonical and non-canonical alignments via profile-profile alignment using ClustalW
v2.1. This combined alignment was once again trimmed with ClipKit to remove potential
residual noise.

We used the resulting filtered alignment to build a PAZ domain-specific HMM profile using
hmmbuild from the HMMER suite v3.3. The final model consisted of 395 match states
(effective sequence number = 24.25; relative entropy per position = 0.59), indicating
moderate diversity and information content. Attempts to improve the entropy score using the
--symfrac 0.7 option did not yield significant differences.

We validated the sensitivity and specificity of the model using hmmsearch against several
curated datasets, including fungal Dicer proteins from our pipeline, annotated Dicers from
NCBI, and representative Dicer like proteins from plants and animals. As negative controls,
we included unrelated sequences such as the Argonaute protein from *Cardiosporidium*
cionae and *Arabidopsis thaliana* EIL1. The model reliably detected PAZ domains in *bona fide*
Dicers across taxa and showed no significant matches in negative controls, confirming its
performance.

Processing of sRNA-seq libraries

We performed a comprehensive search of publicly available sRNA-seq datasets in NCBI,
utilizing the NCBI-SRA tools, eDirect, and custom python scripts using fungal-related terms
(e.g., fungi, fungus, specific fungal genera) combined with sRNA-related keywords (e.g.,
miRNA, microRNA, milRNA, small RNA*, sRNA, siRNA, smRNA, epigen*, RNAi). We cross-
referenced the publications matching the search criteria with NCBI sample databases (SRA,
Bioprojects, Biosamples, and GEO datasets) to identify relevant datasets. We manually
reviewed the entries to identify BioProjects containing libraries likely including fungal sRNA-
seq data from wild-type or Dicer mutants. We processes the selected libraries using the
YASMA small RNA analysis suite (Johnson et al., 2025) (github.com/nateyjay/yasma). We

downloaded libraries using SRAtools. 3' adapters were identified using YASMA-adapter and
trimmed with cutadapt (Martin, 2011) with the following settings: cutadapt -a [adapter] --
minimum-length 15 --maximum-length 50 -O 4 --max-n 0 --trimmed-only -o [out_file] [file]. We
aligned trimmed sequences to the closest matching reference genome using YASMA-align,
which mimics the unique-weighting approach found in ShortStack3 (Johnson et al., 2016).
sRNAs were classified into general sizes using YASMA-size-profile.

Determination of molecular distances in protein structures

We used the Bio3D R library (Grant et al., 2021) to calculate the molecular distance between
the N-terminus of the PAZ domain and the N-terminus of the RNAse IIIa domain. The protein
structure, provided in PDB format, was imported to R using the read.pdb function and the
coordinates of the alpha-carbon atoms corresponding to the N-terminal residues of the PAZ
and RNAse IIIa domains were extracted. We calculated the Euclidean distance between
these coordinates in Angstroms using the xyz.dist function in Bio3D.

Data availability

Structural models of fungal Dicer-like (Dcr) proteins predicted by AlphaFold2, along with the
custom Hidden Markov Model (FPAZ.HMM) used for PAZ domain identification, have been
deposited in Figshare and are publicly accessible at: 10.6084/m9.figshare.28592981.
All other data supporting the findings of this study are available from the corresponding
authors upon reasonable request.

Acknowledgements

This work was funded by Agencia Nacional de Investigación y Desarrollo (ANID)-National
Fund for Scientific and Technological Development (FONDECYT) Grants 11220727 (to
N.R.J.), 11251927 (to C.M.), 1221667 (to V.C-F.), 3230600 (to P.V.), 11200209 (to J.P.C)
and 1221064 (to J.A.R-P); ANID-Millennium Science Initiative Program, Millennium Institute
for Integrative Biology iBio Grant ICN17_022 (to C.M., N.R.J. and E.A.V.); ANID-Millennium
Science Initiative Program, Millennium Nucleus in Data Science for Plant Resilience
Phytolearning Grant NCN2024_047 (to C.M. and E.A.V.). ANID-Vinculación Internacional
Grant FOVI230159 (to E.A.V.); ANID-Beca Doctoral 2021-21211564 (to B.V-V) and 2022-
21221055 (to I.O.); and Beca Doctoral Universidad Mayor (to L.M.).

The authors acknowledge the use of the High-Performance Computing UOH Laboratory (FIC
40059065-0) at Universidad de O'Higgins, Rancagua, for providing supercomputing
infrastructure support.

829
830 **Author contributions**

831 Conceptualization: L.M., N.R.J., E.A.V.; Investigation and formal analysis: L.M., J.C., P.V.,
832 B.V-V., F.G-T., I.O.; Supervision: J.A.R-P., J.P.C., C.M., V.C-F., N.R.J., E.A.V.; Writing-
833 Original Draft: L.M., N.R.J., E.A.V; Writing-Review & Editing: L.M., J.C., P.V., B.V-V., I.O.,
834 J.A.R-P., J.P.C., C.M., V.C-F., N.R.J., E.A.V.
835
836 **Competing interests**

The authors declare no competing interests.

References

1. Bailey, T. L., Johnson, J., Grant, C. E., & Noble, W. S. (2015). The MEME Suite. *Nucleic*
Acids Research, 43(W1), W39–W49. https://doi.org/10.1093/nar/gkv416

2. Baker, N. A., Sept, D., Joseph, S., Holst, M. J., & McCammon, J. A. (2001). Electrostatics
of nanosystems: Application to microtubules and the ribosome. *Proceedings of the*
National Academy of Sciences, 98(18), 10037–10041.
https://doi.org/10.1073/pnas.181342398

3. Bernstein, E., Kim, S. Y., Carmell, M. A., Murchison, E. P., Alcorn, H., Li, M. Z., Mills, A.
A., Elledge, S. J., Anderson, K. V., & Hannon, G. J. (2003). Dicer is essential for mouse
development. *Nature Genetics*, 35(3), 215–217. https://doi.org/10.1038/ng1253

4. Billmyre, R. B., Calo, S., Feretzaki, M., Wang, X., & Heitman, J. (2013). RNAi function,
diversity, and loss in the fungal kingdom. *Chromosome Research*, 21(6–7), 561–572.
https://doi.org/10.1007/s10577-013-9388-2

5. Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using
DIAMOND. *Nature methods*, 12(1), 59-60.

6. Buck, A. H., Coakley, G., Simbari, F., McSorley, H. J., Quintana, J. F., Le Bihan, T.,
Kumar, S., Abreu-Goodger, C., Lear, M., Harcus, Y., Ceroni, A., Babayan, S. A., Blaxter,
M., Ivens, A., & Maizels, R. M. (2014). Exosomes secreted by nematode parasites
transfer small RNAs to mammalian cells and modulate innate immunity. *Nature*
Communications, 5(1), 5488. https://doi.org/10.1038/ncomms6488

7. Cai, Q., Qiao, L., Wang, M., He, B., Lin, F.-M., Palmquist, J., Huang, S.-D., & Jin, H.
(2018). Plants send small RNAs in extracellular vesicles to fungal pathogen to silence
virulence genes. *Science*, 360(6393), 1126–1129. https://doi.org/10.1126/science.aar4142

8. Cánovas-Márquez, J. T., Falk, S., Nicolás, F. E., Padmanabhan, S., Zapata-Pérez, R.,
Sánchez-Ferrer, Á., Navarro, E., & Garre, V. (2021). A ribonuclease III involved in
virulence of Mucorales fungi has evolved to cut exclusively single-stranded RNA. *Nucleic*
Acids Research, 49(9), 5294–5307. https://doi.org/10.1093/nar/gkab238

9. Carreras-Villaseñor, N., Esquivel-Naranjo, E. U., Villalobos-Escobedo, J. M.,
Abreu-Goodger, C., & Herrera-Estrella, A. (2013). The RNAI machinery regulates growth
and development in the filamentous fungus Trichoderma atroviride. *Molecular*
Microbiology, 89(1), 96–112. https://doi.org/10.1111/mmi.12261

10.   Catalanotto, C., Pallotta, M., ReFalo, P., Sachs, M. S., Vayssie, L., Macino, G., &
870   Cogoni, C. (2004). Redundancy of the Two Dicer Genes in Transgene-Induced
871   Posttranscriptional Gene Silencing in Neurospora crassa. *Molecular and Cellular Biology*,
872   24(6), 2536–2545. https://doi.org/10.1128/MCB.24.6.2536-2545.2004

873   11.   Chang, S.-S., Zhang, Z., & Liu, Y. (2012). RNA Interference Pathways in Fungi:
874   Mechanisms and Functions. *Annual Review of Microbiology*, 66(1), 305–323.
875   https://doi.org/10.1146/annurev-micro-092611-150138

876   12.   Choi, J., Kim, K.-T., Jeon, J., Wu, J., Song, H., Asiegbu, F. O., & Lee, Y.-H. (2014).
877   funRNA: A fungi-centered genomics platform for genes encoding key components of
878   RNAi. *BMC Genomics*, 15(S9), S14. https://doi.org/10.1186/1471-2164-15-S9-S14

879   13.   Dang, Y., Yang, Q., Xue, Z., & Liu, Y. (2011). RNA Interference in Fungi: Pathways,
880   Functions, and Applications. *Eukaryotic Cell*, 10(9), 1148–1155.
881   https://doi.org/10.1128/EC.05109-11

882   14.   De Haro, J. P., Calo, S., Cervantes, M., Nicolás, F. E., Torres-Martínez, S., & Ruiz-
883   Vázquez, R. M. (2009). A Single dicer Gene Is Required for Efficient Gene Silencing
884   Associated with Two Classes of Small Antisense RNAs in Mucor circinelloides. *Eukaryotic
885   Cell*, 8(10), 1486–1497. https://doi.org/10.1128/EC.00191-09

886   15.   Dolinsky, T. J., Nielsen, J. E., McCammon, J. A., & Baker, N. A. (2004). PDB2PQR:
887   An automated pipeline for the setup of Poisson-Boltzmann electrostatics calculations.
888   *Nucleic Acids Research*, 32(Web Server), W665–W667.
889   https://doi.org/10.1093/nar/gkh381

890   16.   Dong, R., Pan, S., Peng, Z., Zhang, Y., & Yang, J. (2018). mTM-align: A server for
891   fast protein structure database search and multiple protein structure alignment. *Nucleic
892   Acids Research*. https://doi.org/10.1093/nar/gky430

893   17.   Drinnenberg, I. A., Weinberg, D. E., Xie, K. T., Mower, J. P., Wolfe, K. H., Fink, G. R.,
894   & Bartel, D. P. (2009). RNAi in Budding Yeast. *Science*, 326(5952), 544–550.
895   https://doi.org/10.1126/science.1176945

896   18.   Enriquez-Felix, E. E., Pérez-Salazar, C., Rico-Ruiz, J. G., Calheiros De Carvalho, A.,
897   Cruz-Morales, P., Villalobos-Escobedo, J. M., & Herrera-Estrella, A. (2024). Argonaute
898   and Dicer are essential for communication between Trichoderma atroviride and fungal
899   hosts during mycoparasitism. *Microbiology Spectrum*, 12(4), e03165-23.
900   https://doi.org/10.1128/spectrum.03165-23

19. Gasciolli, V., Mallory, A. C., Bartel, D. P., & Vaucheret, H. (2005). Partially Redundant
Functions of Arabidopsis DICER-like Enzymes and a Role for DCL4 in Producing trans-
Acting   siRNAs.   *Current   Biology*,   15(16),   1494–1500.
https://doi.org/10.1016/j.cub.2005.07.024

20. Grant, B. J., Skjærven, L., & Yao, X. (2021). The BIO3D packages for structural
bioinformatics. *Protein Science*, 30(1), 20–30. https://doi.org/10.1002/pro.3923

21. Hiraguri, A., Itoh, R., Kondo, N., Nomura, Y., Aizawa, D., Murai, Y., Koiwa, H., Seki,
M., Shinozaki, K., & Fukuhara, T. (2005). Specific interactions between Dicer-like proteins
and HYL1/DRB- family dsRNA-binding proteins in Arabidopsis thaliana. *Plant Molecular
Biology*, 57(2), 173–188. https://doi.org/10.1007/s11103-004-6853-5

22. Hu, Y., Stenlid, J., Elfstrand, M., & Olson, Å. (2013). Evolution of RNA interference
proteins dicer and argonaute in Basidiomycota. *Mycologia*, 105(6), 1489–1498.
https://doi.org/10.3852/13-171

23. Huang, J., & MacKerell, A. D. (2013). CHARMM36 all-atom additive protein force
field: Validation based on comparison to NMR data. *Journal of Computational Chemistry*,
34(25), 2135–2145. https://doi.org/10.1002/jcc.23354

24. Huang, Q. (2018). Evolution of Dicer and Argonaute orthologs in microsporidian
parasites.   Infection,   *Genetics   and   Evolution*,   65,   329–332.
https://doi.org/10.1016/j.meegid.2018.08.011

25. James, T. Y., Stajich, J. E., Hittinger, C. T., & Rokas, A. (2020). Toward a Fully
Resolved Fungal Tree of Life. *Annual Review of Microbiology*, 74(1), 291–313.
https://doi.org/10.1146/annurev-micro-022020-051835

26. Jespersen, N., Monrroy, L., & Barandun, J. (2022). Impact of Genome Reduction in
Microsporidia. In L. M. Weiss & A. W. Reinke (Eds.), Microsporidia (Vol. 114, pp. 1–42).
Springer International Publishing. https://doi.org/10.1007/978-3-030-93306-7_1

27. Johnson, N. R., Gonzalez-Toro, F., & Bernal Gomez, B. (2025). Class-agnostic
annotation of small RNAs balances sensitivity and specificity in diverse organisms.
Genomics. https://doi.org/10.1101/2025.01.25.634868

28. Johnson, N. R., Yeoh, J. M., Coruh, C., & Axtell, M. J. (2016). Improved placement of
multi-mapping small RNAs. *G3: Genes, Genomes, Genetics*, 6(7), 2103-2111.

29. Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,
Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A.,

Scheremetjew, M., Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: Genome-
scale protein function classification. *Bioinformatics*, 30(9), 1236–1240.
https://doi.org/10.1093/bioinformatics/btu031

30.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl,
S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., …
Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*,
596(7873), 583–589. https://doi.org/10.1038/s41586-021-03819-2

31.   Kadotani, N., Nakayashiki, H., Tosa, Y., & Mayama, S. (2004). One of the Two Dicer-
like Proteins in the Filamentous Fungi Magnaporthe oryzae Genome Is Responsible for
Hairpin RNA-triggered RNA Silencing and Related Small Interfering RNA Accumulation.
Journal of Biological Chemistry, 279(43), 44467–44474.
https://doi.org/10.1074/jbc.M408259200

32.   Kans, J. (2024). Entrez direct: E-utilities on the UNIX command line. In Entrez
programming utilities help [Internet]. National Center for Biotechnology Information (US).

33.   Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution*,
30(4), 772–780. https://doi.org/10.1093/molbev/mst010

34.   Kelani, A. A., Bruch, A., Rivieccio, F., Visser, C., Krüger, T., Weaver, D., Pan, X.,
Schäuble, S., Panagiotou, G., Kniemeyer, O., Bromley, M. J., Bowyer, P., Barber, A. E.,
Brakhage, A. A., & Blango, M. G. (2023). Disruption of the Aspergillus fumigatus RNA
interference machinery alters the conidial transcriptome. *RNA*, 29(7), 1033–1050.

35.   Kress, A., Poch, O., Lecompte, O., & Thompson, J. D. (2023). Real or fake?
Measuring the impact of protein annotation errors on estimates of domain gain and loss
events. *Frontiers in Bioinformatics*, 3, 1178926.
https://doi.org/10.3389/fbinf.2023.1178926

36.   Lau, P.-W., Guiley, K. Z., De, N., Potter, C. S., Carragher, B., & MacRae, I. J. (2012).
The molecular architecture of human Dicer. *Nature Structural & Molecular Biology*, 19(4),
436–440. https://doi.org/10.1038/nsmb.2268

37.   Lau, P.-W., Potter, C. S., Carragher, B., & MacRae, I. J. (2009). Structure of the
Human Dicer-TRBP Complex by Electron Microscopy. *Structure*, 17(10), 1326–1332.
https://doi.org/10.1016/j.str.2009.08.013

38.   Lee, H.-C., Li, L., Gu, W., Xue, Z., Crosthwaite, S. K., Pertsemlidis, A., Lewis, Z. A.,
Freitag, M., Selker, E. U., Mello, C. C., & Liu, Y. (2010). Diverse Pathways Generate
MicroRNA-like RNAs and Dicer-Independent Small Interfering RNAs in Fungi. *Molecular*
Cell, 38(6), 803–814. https://doi.org/10.1016/j.molcel.2010.04.005

39.   Li, Y., Steenwyk, J. L., Chang, Y., Wang, Y., James, T. Y., Stajich, J. E., Spatafora, J.
W., Groenewald, M., Dunn, C. W., Hittinger, C. T., Shen, X.-X., & Rokas, A. (2021). A
genome-scale phylogeny of the kingdom Fungi. *Current Biology*, 31(8), 1653-1665.e5.
https://doi.org/10.1016/j.cub.2021.01.074

40.   Lim, L. P., Lau, N. C., Garrett-Engele, P., Grimson, A., Schelter, J. M., Castle, J.,
Bartel, D. P., Linsley, P. S., & Johnson, J. M. (2005). Microarray analysis shows that some
microRNAs downregulate large numbers of target mRNAs. *Nature*, 433(7027), 769–773.
https://doi.org/10.1038/nature03315

41.   Ma, J.-B., Ye, K., & Patel, D. J. (2004). Structural basis for overhang-specific small
interfering RNA recognition by the PAZ domain. *Nature*, 429(6989), 318–322.
https://doi.org/10.1038/nature02519

42.   MacRae, I. J., Zhou, K., Li, F., Repic, A., Brooks, A. N., Cande, W. Z., Adams, P. D.,
& Doudna, J. A. (2006). Structural Basis for Double-Stranded RNA Processing by Dicer.
Science, 311(5758), 195–198. https://doi.org/10.1126/science.1121638

43.   Martinez, J., Patkaniowska, A., Urlaub, H., Lührmann, R., & Tuschl, T. (2002).
Single-Stranded Antisense siRNAs Guide Target RNA Cleavage in RNAi. *Cell*, 110(5),
563–574. https://doi.org/10.1016/S0092-8674(02)00908-X
986
987   44.   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
988   sequencing reads. EMBnet. journal, 17(1), 10-12.
989
45.   Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von
Haeseler, A., & Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for
Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution*, 37(5),
1530–1534. https://doi.org/10.1093/molbev/msaa015

46.   Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., & Steinegger, M.
(2022). ColabFold: Making protein folding accessible to all. *Nature Methods*, 19(6), 679–
682. https://doi.org/10.1038/s41592-022-01488-1

47.   Mukherjee, K., Campos, H., & Kolaczkowski, B. (2013). Evolution of Animal and Plant
Dicers: Early Parallel Duplications and Recurrent Adaptation of Antiviral RNA Binding in

Plants. *Molecular Biology and Evolution*, 30(3), 627–641. https://doi.org/10.1093/molbev/mss263

48. Notredame, C., Higgins, D. G., & Heringa, J. (2000). T-coffee: A novel method for fast and accurate multiple sequence alignment 1 1Edited by J. Thornton. *Journal of Molecular Biology*, 302(1), 205–217. https://doi.org/10.1006/jmbi.2000.4042

49. Olsson, Mats HM, Chresten R. Sondergaard, Michal Rostkowski, and Jan H. Jensen. "PROPKA3: consistent treatment of internal and surface residues in empirical pKa predictions." Journal of Chemical Theory and Computation 7, no. 2 (2011): 525-537. doi:10.1021/ct100578z

50. Ota, H., Sakurai, M., Gupta, R., Valente, L., Wulff, B.-E., Ariyoshi, K., Iizasa, H., Davuluri, R. V., & Nishikura, K. (2013). ADAR1 Forms a Complex with Dicer to Promote MicroRNA Processing and RNA-Induced Gene Silencing. *Cell*, 153(3), 575–589. https://doi.org/10.1016/j.cell.2013.03.024

51. Phillips, J. C., Hardy, D. J., Maia, J. D., Stone, J. E., Ribeiro, J. V., Bernardi, R. C., ... & Tajkhorshid, E. (2020). Scalable molecular dynamics on CPU and GPU architectures with NAMD. *The Journal of chemical physics*, 153(4).

52. Potter, S. C., Luciani, A., Eddy, S. R., Park, Y., Lopez, R., & Finn, R. D. (2018). HMMER web server: 2018 update. *Nucleic Acids Research*, 46(W1), W200–W204. https://doi.org/10.1093/nar/gky448

53. Qin, H., Chen, F., Huan, X., Machida, S., Song, J., & Yuan, Y. A. (2010). Structure of the Arabidopsis thaliana DCL4 DUF283 domain reveals a noncanonical double-stranded RNA-binding fold for protein–protein interaction. *RNA*, 16(3), 474–481. https://doi.org/10.1261/rna.1965310

54. Qin, S., Veloso, J., Baak, M., Boogmans, B., Bosman, T., Puccetti, G., Shi-Kunne, X., Smit, S., Grant-Downton, R., Leisen, T., Hahn, M., & Van Kan, J. A. L. (2023). Molecular characterization reveals no functional evidence for naturally occurring cross-kingdom RNA interference in the early stages of Botrytis cinerea –tomato interaction. *Molecular Plant Pathology*, 24(1), 3–15. https://doi.org/10.1111/mpp.13269

55. Raman, V., Simon, S. A., Demirci, F., Nakano, M., Meyers, B. C., & Donofrio, N. M. (2017). Small RNA functions are required for growth and development of Magnaporthe oryzae. *Molecular plant-microbe interactions*, *30*(7), 517-530.

1030   56.   Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology
1031   (and other things). *Methods in Ecology and Evolution*, 3(2), 217–223.
1032   https://doi.org/10.1111/j.2041-210X.2011.00169.x

1033   57.   Schwarz, D. S., Hutvágner, G., Du, T., Xu, Z., Aronin, N., & Zamore, P. D. (2003).
1034   Asymmetry in the Assembly of the RNAi Enzyme Complex. *Cell*, 115(2), 199–208.
1035   https://doi.org/10.1016/S0092-8674(03)00759-1

1036   58.   Seppey, M., Manni, M., & Zdobnov, E. M. (2019). BUSCO: assessing genome
1037   assembly and annotation completeness. *Gene prediction: methods and protocols*, 227-
1038   245.

1039   59.   Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N.,
1040   Schwikowski, B., & Ideker, T. (2003). Cytoscape: A Software Environment for Integrated
1041   Models of Biomolecular Interaction Networks. *Genome Research*, 13(11), 2498–2504.
1042   https://doi.org/10.1101/gr.1239303

1043   60.   Sinha, N. K., Iwasa, J., Shen, P. S., & Bass, B. L. (2018). Dicer uses distinct modules
1044   for recognizing dsRNA termini. *Science*, 359(6373), 329–334.
1045   https://doi.org/10.1126/science.aaq0921

1046   61.   Sinha, N. K., Trettin, K. D., Aruscavage, P. J., & Bass, B. L. (2015). Drosophila Dicer-
1047   2 Cleavage Is Mediated by Helicase- and dsRNA Termini-Dependent States that Are
1048   Modulated by Loquacious-PD. *Molecular Cell*, 58(3), 406–417.
1049   https://doi.org/10.1016/j.molcel.2015.03.012

1050   62.   Sondergaard, Chresten R., Mats HM Olsson, Michal Rostkowski, and Jan H. Jensen.
1051   "Improved Treatment of Ligands and Coupling Effects in Empirical Calculation and
1052   Rationalization of pKa Values." Journal of Chemical Theory and Computation 7, no. 7
1053   (2011): 2284-2295. doi:10.1021/ct200133y

1054   63.   Steenwyk, J. L., Buida, T. J., Li, Y., Shen, X.-X., & Rokas, A. (2020). ClipKIT: A
1055   multiple sequence alignment trimming software for accurate phylogenomic inference.
1056   *PLOS Biology*, 18(12), e3001007. https://doi.org/10.1371/journal.pbio.3001007

1057   64.   Tedersoo, L., Sánchez-Ramírez, S., Kõljalg, U., Bahram, M., Döring, M., Schigel, D.,
1058   May, T., Ryberg, M., & Abarenkov, K. (2018). High-level classification of the Fungi and a
1059   tool for evolutionary ecological analyses. *Fungal Diversity*, 90(1), 135–159.
1060   https://doi.org/10.1007/s13225-018-0401-0

1061   65.   Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: improving the
1062   sensitivity of progressive multiple sequence alignment through sequence weighting,

position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22(22), 4673–4680.

66. Tian, Y., Simanshu, D. K., Ma, J.-B., Park, J.-E., Heo, I., Kim, V. N., & Patel, D. J. (2014). A Phosphate-Binding Pocket within the Platform-PAZ-Connector Helix Cassette of Human Dicer. *Molecular Cell*, 53(4), 606–616. https://doi.org/10.1016/j.molcel.2014.01.003

67. Villalobos-Escobedo, J. M., Martínez-Hernández, J. P., Pelagio-Flores, R., González-De la Rosa, P. M., Carreras-Villaseñor, N., Abreu-Goodger, C., & Herrera-Estrella, A. H. (2022). Trichoderma atroviride hyphal regeneration and conidiation depend on cell-signaling processes regulated by a microRNA-like RNA. *Microbial Genomics*, *8*(10), 000869.

68. Volpe, T. A., Kidner, C., Hall, I. M., Teng, G., Grewal, S. I. S., & Martienssen, R. A. (2002). Regulation of Heterochromatic Silencing and Histone H3 Lysine-9 Methylation by RNAi. *Science*, 297(5588), 1833–1837. https://doi.org/10.1126/science.1074973

69. Wang, H.-W., Noland, C., Siridechadilok, B., Taylor, D. W., Ma, E., Felderer, K., Doudna, J. A., & Nogales, E. (2009). Structural insights into RNA processing by the human RISC-loading complex. *Nature Structural & Molecular Biology*, 16(11), 1148–1153. https://doi.org/10.1038/nsmb.1673

70. Wang, M., Weiberg, A., Lin, F.-M., Thomma, B. P. H. J., Huang, H.-D., & Jin, H. (2016). Bidirectional cross-kingdom RNAi and fungal uptake of external RNAs confer plant protection. *Nature Plants*, 2(10), 16151. https://doi.org/10.1038/nplants.2016.151

71. Weiberg, A., Wang, M., Lin, F.-M., Zhao, H., Zhang, Z., Kaloshian, I., Huang, H.-D., & Jin, H. (2013). Fungal Small RNAs Suppress Plant Immunity by Hijacking Host RNA Interference Pathways. *Science*, 342(6154), 118–123. https://doi.org/10.1126/science.1239705

72. Welker, N. C., Maity, T. S., Ye, X., Aruscavage, P. J., Krauchuk, A. A., Liu, Q., & Bass, B. L. (2011). Dicer's Helicase Domain Discriminates dsRNA Termini to Promote an Altered Reaction Mode. *Molecular Cell*, 41(5), 589–599. https://doi.org/10.1016/j.molcel.2011.02.005

73. Wijayawardene, N. N., Hyde, K. D., Mikhailov, K. V., Péter, G., Aptroot, A., Pires-Zottarelli, C. L., ... & Karpov, S. A. (2024). Classes and phyla of the kingdom Fungi. *Fungal diversity*, 1-165.

74.   Wilkins, C., Dishongh, R., Moore, S. C., Whitt, M. A., Chow, M., & Machaca, K.
(2005). RNA interference is an antiviral defence mechanism in Caenorhabditis elegans.
Nature, 436(7053), 1044–1047. https://doi.org/10.1038/nature03957

75.   Yan, Y., Zhang, D., Zhou, P., Li, B., & Huang, S.-Y. (2017). HDOCK: A web server for
protein–protein and protein–DNA/RNA docking based on a hybrid strategy. *Nucleic Acids*
Research, 45(W1), W365–W373. https://doi.org/10.1093/nar/gkx407

76.   Zapletal, D., Kubicek, K., Svoboda, P., & Stefl, R. (2023). Dicer structure and
function: Conserved and evolving features. *EMBO Reports*, 24(7), e57215.
https://doi.org/10.15252/embr.202357215

77.   Zhang, H. (2002). Human Dicer preferentially cleaves dsRNAs at their termini without
a requirement for ATP. *The EMBO Journal*, 21(21), 5875–5885.
https://doi.org/10.1093/emboj/cdf582

78.   Zhao, H., Nie, Y., Zong, T.-K., Wang, K., Lv, M.-L., Cui, Y.-J., Tohtirjap, A., Chen, J.-
J., Zhao, C.-L., Wu, F., Cui, B.-K., Yuan, Y., Dai, Y.-C., & Liu, X.-Y. (2023). Species
diversity, updated classification and divergence times of the phylum Mucoromycota.
Fungal Diversity, 123(1), 49–157. https://doi.org/10.1007/s13225-023-00525-4

79.   Zhu, Q., Kosoy, M., & Dittmar, K. (2014). HGTector: an automated method facilitating
genome-wide discovery of putative horizontal gene transfers. *BMC genomics*, 15, 1-18.

1113

1114

 Figures

1116

1117

Figure 1. Domain-based classification and distribution of Dicer-like proteins across fungal phyla. (**A**) Summary of the 1,593 annotated fungal proteomes analyzed for the identification of Dcr proteins, showing the percentage of low (<90%) and high (>90%) completeness proteomes according to BUSCO. (**B**) Representative domain architectures of known Dcr proteins from *Arabidopsis thaliana*, *Homo sapiens*, and *Schizosaccharomyces pombe*, illustrating the conserved domain composition: DEAD, ResIII, Helicase C (HeliC), DUF283, PAZ, RNase IIIa and IIIb, and dsRNA-binding domain (dsRBD). Protein lengths are shown in amino acids (aa). (**C**) Classification of all predicted fungal Dcr proteins into six structural categories based on the presence or absence of key domains. Canonical Dcrs retain a full complement of conserved domains, while non-canonical groups exhibit partial or divergent architectures. (**D**) Average protein length (in aa) per Dcr category. (**E**) Distribution of Dcr categories across eight fungal phyla, highlighting differences in domain conservation and structural diversity among lineages. (**F**) Frequency distribution of fungal species harboring one to seven Dcr copies, indicating variability in Dcr gene family expansion across the dataset. (**G**) Combinations of Dcr categories across fungal genomes. Each row shows a unique combination of canonical and non-canonical Dcrs (A–E), with colored squares indicating presence of a certain domain. Bars on the right represent the number of organisms per combination (in red).

1136

1137

1138

Figure 2. Phylogenetic and functional classification of fungal Dcr proteins.
(**A**) Maximum-likelihood phylogenetic tree showing the distribution of Dcr proteins across
eight fungal phyla, as well as representative outgroups from plants and metazoans. Colored
ovals highlight major clades identified within fungal lineages, labeled as As1–As2
(Ascomycota), Ba1–Ba3 (Basidiomycota), Mu1–Mu2 (Mucoromycota and Zoopagomycota),
and others, based on their phylogenetic position. (**B**) Annotation status of Dcr proteins
according to UniProtKB, distinguishing Dcr1/Dcl1 (red), Dcr2/Dcl2 (orange), unnumbered
Dcr/Dcl entries (cyan), and unnamed proteins (black). (**C**) Classification of fungal Dcrs into
canonical and five non-canonical categories (A–E), based on domain composition. (**D**)
Mapping of cPAZ domain presence (yellow) or absence (purple) across the phylogeny.

1149

1150

Figure 3. Structural Network of Proteins Revealing Functional Clusters Based on Folding Patterns. (**A**) Taxonomic composition of each structural cluster, shown as the number of proteins per fungal phylum. (**B**) Structural similarity network of predicted Dcr proteins, clustered based on TM-score ≥ 0.9. Each node represents a protein structure; edges indicate pairwise structural similarity above the threshold. Node colors match the clade classification defined in Fig. 2A (e.g., As1, As2, Ba1–Ba3, Mu1–Mu2), allowing direct comparison between structural and phylogenetic groupings. The most connected node (hub) from selected modules is shown with a red circle and its corresponding AlphaFold-predicted 3D structure is represented on the side. Proteins displayed correspond to representative members of non-canonical Dcr groups B, C, and E. Structural domains are colored as follows: PAZ (yellow), Helicase (green), RNase III (dark red/burgundy), DUF283 (cyan), and dsRBD (purple). Regions without domain predictions are shown in grey. Triangular nodes correspond to Dcr proteins with a canonical PAZ domain (cPAZ); oval nodes indicate non-canonical PAZ domains. (**C**) Phylogenetic tree of all Dcr proteins, with tips colored according to structural cluster membership, highlighting the correspondence between structural similarity and evolutionary lineage.

1171

1172
1173
Figure 4. Evolutionary reconstruction of the PAZ domain in fungal Dcr proteins. (**A**)
Phylogenetic tree of fungal Dcr proteins with ancestral state reconstruction for the canonical
PAZ domain. Pie charts at internal nodes represent the inferred probability of PAZ domain
presence (yellow) or absence (purple) using maximum likelihood estimation. Additional
metadata layers include the presence/absence of a canonical PAZ domain, phylogenetic
group (e.g., As1, Ba2, Mu1), and structural cluster as defined by the 3D similarity network in
Fig. 3. (**B**) Proportion of Dcr proteins with a canonical PAZ domain across four eukaryotic
kingdoms, Animalia, Fungi, Plantae, and Protista, highlighting variation in PAZ domain
conservation. (**C**) Horizontal gene transfer (HGT) analysis. No inter-kingdom HGT events
involving Dcr proteins were detected between fungi and plants, animals, or protists,
indicating that the PAZ domain in fungal Dcrs evolved independently from those in other
eukaryotic lineages.
1186

1187

Figure 5. Motif conservation and structural variability of PAZ domains in fungal Dicer-like proteins. (**A**) Conserved sequence motifs identified within the canonical PAZ domain using MEME Suite. Schematic representation (top) shows the approximate location of four conserved motifs (Logos 1–4) along the PAZ region. Sequence logos illustrate consensus motifs and their relative frequency in proteins where the PAZ domain was detected (PAZ-found) versus those where it was not (PAZ-not-found). (**B**) Structural comparison of PAZ domains across Dcr categories. The full-length AlphaFold model of a canonical Dcr (SAM06846.1) is shown on the left, with the PAZ domain highlighted in yellow. Right panel: Representative AlphaFold structural models of PAZ domains from non-canonical categories A–E, annotated with TM-score values and presence/absence of a recognizable cPAZ domain. PAZ regions in non-canonical Dcrs are color-coded according to their category (A–E), as shown in the figure. Structural alignment was performed using the PAZ domain of SAM06846.1 as reference to comparing the spatial conservation of PAZ-like regions across non-canonical Dcrs. (**C**) Boxplot of TM-scores comparing structural similarity of PAZ domains across canonical and non-canonical Dcr categories. Canonical domains show consistently higher structural conservation. (**D**) Sensitivity comparison of the canonical PAZ HMM (PAZ) versus the fungal-specific FPAZ-HMM (FPAZ) across Eukaryotic groups.

1205

Figure 6. Structural and functional characterization of RNA-binding interfaces in canonical and non-canonical fungal Dcr proteins. (**A–B**) Surface representations of models of full-length Dcr proteins (gray) in complex with dsRNA (green), showing the RNA-binding interfaces (blue). (**A**) Canonical Dcr proteins containing a canonical PAZ domain (cPAZ), grouped by structural group (Cl-3, Cl-5, and Cl-6) and domain architecture. (**B**) Non-canonical Dcrs possessing fungal-specific PAZ regions (fPAZ), grouped by structural group (Cl1, Cl2, Other, or NA). Representative residues involved in interaction with RNA (lysines and arginines) are shown in atomic detail, with the side chains located within 3 Å of the RNA. Red boxes highlight the cPAZ or fPAZ region that anchors the 3' end of the dsRNA. "Other" includes proteins belonging to minor structural clusters outside the four major groups. "NA" includes proteins that did not meet the clustering thresholds (≥90% TM-score similarity and pLDDT ≥70). (**C**) Electrostatic surface potential is shown for each model, with red indicating negative charge and blue indicating positive charge (scale in kT/e). The measured distance between the fPAZ region and the RNase III catalytic site (in Å) is shown, suggesting its possible role in determining the length of small RNA (sRNA). WT, solid lines, Dcr mutant, dashed lines. Light blue peaks indicate sRNA species lost in the mutant backgrounds.

1227 **Supplementary Figures**

1228

1229

1230

1231

1232

Supplementary Figure 1. Helicase and dsRBD composition of fungal Dcr proteins
mapped onto the phylogeny. (**A**) Distribution of helicase domains subcomponents (DEAD,
RESIII, and Helicase) across fungal Dcr proteins. Each tip in the phylogeny is colored on a
binary triplet indicating the presence or absence of helicase subdomains in the order: DEAD,
RESIII, and Helicase C (Heli_C). Green tips (111) indicate complete helicase domains, while
grey tips (000) correspond to complete absence of these subdomains. (**B**) Presence or
absence of the canonical dsRBD in fungal Dcr proteins. Presence (blue tips) or absence (no
visible tip) of the canonical dsRNA-binding domain (dsRBD) across the same phylogenetic
tree.

1242

1243

1244

1245

Supplementary Figure 2. Spatial localization of conserved PAZ motifs in a canonical
Dicer protein. Mapping of sequence logos (Logos 1–4) derived from conserved motifs
within the canonical PAZ domain onto the 3D structure of a Dicer protein from *Absidia*
glauca (SAM06846.1), modeled in complex with double-stranded RNA (dsRNA, magenta).
The protein backbone is shown in white cartoon representation. The core PAZ domain is
highlighted in yellow, while each conserved motif is colored individually: Logo 1 (red), Logo 2
(purple), Logo 3 (cyan), and Logo 4 (green). These motifs cluster at the interface between
the PAZ domain and the 3' end of the dsRNA, consistent with their proposed role in RNA
binding.
1255

1256
1257
Supplementary Figure 3. Structural stability of Dicer–RNA complexes during molecular dynamics simulations. Root mean square deviation (RMSD, in Å) over a 50-nanosecond (ns) molecular dynamics trajectory for six representative Dicer–RNA complexes. (**A**) Dcr proteins containing a canonical PAZ domain (cPAZ), (**B**) Fungal-specific PAZ domains (fPAZ) as defined in this study. Each plot shows the RMSD trajectories from three independent replicates, replicate 1 (blue), replicate 2 (red), and replicate 3 (green), with the black line indicating the mean RMSD across replicates. These values reflect the conformational stability of each complex over time.
1266