1   Title:

2   **Macromolecular Crowding-Affected Mobility of Bcnt/Cfdp1, a Disordered Component of**

3   **the Srcap Complex, from the Nucleus to Cytosol in Subcellular Fractionation**

Authors:

Shintaro Iwashita[1,2, †, *], Takehiro Suzuki[3, †], Kouichi Nishimura[4], Ayaka Onishi[1], Kyoka Tsukuda[2,5,¶], Yoshito Kakihara[6], Yoshimitsu Kiriyama[1,2], Yoshiharu Ohoka[1], Kentaro Nakashima[1,2*]

Authors and Affiliations

[1] Kagawa School of Pharmaceutical Sciences, Tokushima Bunri University, Kagawa, 769-2193, Japan[§]

[2] Institute of Neuroscience, Tokushima Bunri University, Kagawa, 769-2193, Japan[§]

[3] Biomolecular Characterization Unit, RIKEN Center for Sustainable Resource Science, Saitama 351-0198, Japan

[4] Graduate School of Biomedical Science, Tokushima University, Tokushima, 770-8503, Japan

[5] Graduate School of Pharmaceutical Sciences, Tokushima Bunri University, Kagawa, 769-2193, Japan[§]

[6] Division of Dental Pharmacology, Niigata University Graduate School of Medical and Dental Sciences, Niigata, 951-8514, Japan

[¶]Present address: Department of Pharmacy, Faculty of Pharmacy, Kindai University, Osaka, 577-8502, Japan

[†]These authors contributed equally to the analysis.

[§]Tokushima Bunri University at Sanuki City campus moved to Takamatsu City on 1st April 2025. New address: 8-53 Hamanocho, Takamatsu, Kagawa, 760-0011, Japan

* Correspondence: sanukishido1997@gmail.com & nkentaro@kph.bunri-u.ac.jp

Keywords:
Intracellular behavior, Nucleus-cytosol mobility, Western blot, Quantitative proteomic analysis, digitonin, Wnt signaling

ABSTRACT

Metazoan *Bucentaur/Craniofacial Development Protein 1* (*Bcnt/Cfdp1*) is crucial for embryonic

development, and its yeast homolog protein, SWC5, functions as part of the SWR1 complex in

histone exchange reactions. Bcnt/Cfdp1, an intrinsically disordered protein (IDP) subject to

various post-translational modifications, contains a putative nuclear localization sequence.

Nevertheless, its subcellular localization remains ambiguous. We investigated its intracellular

trafficking using diverse approaches. In C2C12 myoblasts, fluorescent protein-tagged

Bcnt/Cfdp1 was predominantly localized within the nucleus, preferentially in regions of low

DAPI density in myoblasts, and the localization persisted in differentiated multinucleated

myotubes. Conversely, detergent-based biochemical fractionation of C2C12 and HEK293T cells

indicated a substantial presence of Bcnt/Cfdp1 in cytoplasmic fractions. Quantitative proteomic

analysis of fractionated proteins revealed that six IDPs, including a PEST-containing nuclear

protein, demonstrated similar distribution patterns to Bcnt/Cfdp1; despite being annotated as

nuclear proteins, they were detected in cytoplasmic fractions. Direct treatment of both cell lines

with digitonin enabled the collection of released proteins in three steps to accurately isolate

cytoplasmic proteins. Bcnt/Cfdp1 exhibited distinct extraction patterns compared to cytoplasmic

soluble protein extracts in Ficoll/sucrose solution, while no differences were noted when cells

underwent digitonin treatment in an isotonic solution. Quantitative proteomic analysis of these

extracts highlighted the mobility or integrity of each protein within the cytoplasm.

Consequently, although Bcnt/Cfdp1 is essential within the nucleus, our findings suggest its

additional presence in the cytoplasm. We hypothesize that the elastic properties of Bcnt/Cfdp1

facilitate its translocation from the nucleus during subcellular fractionation, which is affected by

macromolecular crowding.

INTRODUCTION

Organisms have evolved cellular regulatory systems to maintain sophisticated homeostasis in

response to environmental stimuli, such as heat and radiation, and stimuli generated by

intracellular biosynthesis, such as oxidative stress. Cellular gene-dependent macromolecules

dynamically shuttle between the nucleus and cytoplasm during biosynthesis and are recruited to

appropriate cellular compartments with high efficiency in response to intracellular and

extracellular stimuli. Recent evidence has revealed that these macromolecular behavior

dynamics depend on the generated condensates or liquid-liquid phase separation (LLPS), which

allows the inclusion and exclusion of necessary and unnecessary components, respectively [1–

3].

Among the myriad cellular macromolecules, intrinsically disordered proteins (IDPs) and
proteins containing intrinsically disordered regions (IDRs) have attracted attention in the
context of condensate formation. IDPs and IDRs are frequently found as scaffolding proteins
with multivalent macromolecules and RNA in condensates, such as stress granules (SGs) and
processing bodies (PBs) in the cytoplasm and paraspeckles in the nucleus [4]. It is noteworthy
that the subtle structure changes in IDRs, such as modifications through
phosphorylation/dephosphorylation, have the potential to regulate condensate formation [3].
The low-density condensates have porous and permeable properties, demonstrated in SGs and
nuclear speckles [5], where most molecules move freely while the concentrations of related
molecules are high [6].
In the context of nuclear-cytoplasmic transport, the structure of the nuclear membrane,
particularly the nuclear pore complex (NPC), the only gate with bidirectional potential between
the cytoplasm and nucleus, has been intensively studied; IDP and IDR have also been found to
play crucial roles in this process [7]. The two types of nuclear transport mediated by NPCs are
classified as active and passive transport, depending on the requirement of receptors and energy.
The barrier allows ≤30–60 kDa molecules to pass through passive diffusion, showing its
entropy-driven nature rather than geometric constriction [8]. Notably, IDPs can diffuse through
the NPC without requiring energy or specific transport receptors, while folded protein transport
has been revealed to be size-dependent [9].
From yeast to humans, metazoan Bucentaur/Craniofacial development protein 1 (Bcnt/Cfdp1)[‡]
displays a high degree of structural disorder in the N-terminal region and an evolutionarily
conserved C-terminal amino acid sequence (BCNT domain) [10]. The yeast ortholog SWC5 is a
subunit of the SWR1 (SRCAP in humans) chromatin remodeling complex, SWR1-C (SRCAP-C
in humans), while BCNT/CFDP1 has often been treated as an orphan [11, 12]. This is partly due
to the failure to capture it as a component of SRCAP-C compared with other subunits [13]. The
N-terminal DEY/F motifs of SWC5 engage the canonical histone H2A–H2B [14], whereas the
C-terminal BCNT-C domain binds histone H3–H4 and DNA [15] and activates the ATPase of
SWR1 [16]. This dual function may contribute to the direction of the torsional force converting
the H2A–H2B dimer to H2A.Z–H2B [14–17].

[‡]The official name of *Bcnt/Cfdp1* is *Cfdp1*. However, genetic studies of disease [1] or transcriptome analysis [2] do not provide solid evidence for its involvement in craniofacial development, including mouse tooth development [3]. Therefore, the nomenclature of *Cfdp1* is misleading and needs to be more accurate. To maintain consistency with previous studies, the term *Bcnt/Cfdp1* rather than *Cfdp1* is used in this article.

SWC5 data, based on genetic information, provide clues for inferring the functional frame of

Bcnt/Cfdp1 from a causal perspective (https://www.yeastgenome.org/locus/S000000435) [18].

Although *SWR1* and ***SWC5*** are non-essential genes, they play crucial roles under stressful

conditions and have versatile functions. SWC5 protein abundance increases in response to DNA

replication stress, and each null strain has decreased or increased resistance to certain chemical

compounds, which, along with other factors, may affect their lifespan [18]. In addition, tight

control of the histone exchange reaction has been demonstrated to be essential for regulating

chromatin dynamics [19].

In contrast, *Bcnt/Cfdp1* is essential for metazoan development, as has been demonstrated for the

Drosophila homolog *Yeti* [20], zebrafish *cfdp1* [21, 22], and mouse *Cfdp1* [23],

[https://www.jax.org/strain/033834], suggesting its critical role in pluripotency and

differentiation. The following two lines of evidence for *Bcnt/Cfdp1* as a selected gene suggest

that it may have versatile functions in homeostasis, similar to *SWC5*. First, it contributes to the

persistence of Chinese hamster ovary cell cultures [24] and resistance to Marek's disease in

chickens [25]. Perturbation of *Bcnt/Cfdp1* in cells or tissues, whether through gain or loss, has

been demonstrated to modulate cell migration, cell cycle progression, and apoptosis/anti-

apoptosis [20-22, 26]. These phenotypes are preferentially associated with the Wnt/Notch/Akt

signaling pathway [27]. Notably, the Wnt signaling pathway is suppressed explicitly in

extraluminal mesenchymal cells of the valves of zebrafish Bcnt/Cfdp1-deficient embryos. This

defect leads to cardiac dysfunction [22]. Second, *BCNT/CFDP1* has been identified as a

prognostic marker for several types of cancers [28, 29]. In addition, Srcap-C has been

demonstrated to be involved in a wide range of biological processes, including differentiation

[30], DNA damage repair [31], and disease development [32].

We previously reported that *Bcnt/Cfdp1* knockdown at the genomic level in a mouse embryonic

stem (ES) cell line, Cfdp1-K1, had a significant impact on the expression of numerous genes,

with more than 4% of all genes upregulated or downregulated by more than two-fold [33].

Consistent with this result, RNAi depletion of BCNT/CFDP1 in HeLa cells has been reported to

have widespread effects on cellular functions [26]. Of note, this phenomenon may be

comparable to the natural genetic variation found in the fission yeast, *swc5*, likely due to the

insufficient functionality of the histone exchange reaction [34]. Given that the aforementioned

biological phenotypes induced by *Bcnt/Cfdp1* perturbations are preferentially related to the

Wnt/Notch/Akt signaling pathway, it seems reasonable to consider that they are indirectly or

partially due to perturbations in the Srcap-C transcriptional regulatory framework. This

consideration means that while we have gained insight into the function of Bcnt/Cfdp1, we still

need more information about its fundamental properties. Assessing the intracellular

trafficking of target proteins is essential for understanding fundamental physiological
networks and functions, such as cell cycle control [35], and clarifying the causes of
diseases [36].
Vertebrate Bcnt/Cfdp1 has typical putative nuclear localization sequences in the N-terminal
region, such as ^{50}GKKRKAQSIPARKRR^{64} of the human ortholog. However, its subcellular
localization remains ambiguous (https://www.proteinatlas.org/ENSG00000153774-
CFDP1/subcellular) mainly due to the lack of appropriate antibodies against Bcnt/Cfdp1 that
cannot be ignored. Recently, we overcame the issue of off-target bands and assigned the
Western blotting signals of mouse Bcnt/Cfdp1 using a Cfdp1-K1 cell extract as a potential
negative control [33, 37]. In light of these findings, herein, we investigate the intracellular
trafficking of endogenous Bcnt/Cfdp1 in the mouse myoblast cell line C2C12, a well-
established model of myocyte differentiation and multinucleated myotube formation, and in
HEK293T cells. We also discuss the causes of the conflicting results regarding the subcellular
distribution of Bcnt/Cfdp1 between different approaches, considering nuclear export trafficking
and macromolecular crowding. Furthermore, we address the Bcnt/Cfdp1 function in the Srcap-C
framework.

RESULTS
Observation of Bcnt/Cfdp1 in the Nucleus via Cytochemical Approaches
Assessing the subcellular trafficking of the target molecule is essential for clarifying its
functional framework, and examining the endogenous ones is imperative. However, no reliable
antibodies against Bcnt/Cfdp1 are currently available for immunocytochemistry. Previously, we
detected tag-free mouse Bcnt/Cfdp1 primarily in the nuclei of human HEK293T cells by
immunocytochemistry with two antibodies generated against two peptides of mouse
Bcnt/Cfdp1: one is specific to mice and located in the N-terminal region and the other is
common to mice and humans in the C-terminus (Figure 1A, modified from Nakashima *et al*.
[37]).
Here, we used the myoblast C2C12 cell line, a well-known model of myocyte differentiation
and multinucleated myotube formation, to analyze the dynamic behavior of Bcnt/Cfdp1 during
these processes. We expressed N-terminal FLAG- and fluorescent protein-tagged mouse
Bcnt/Cfdp1 (mCherry or enhanced green fluorescent protein (EGFP)) in C2C12 cells. We
examined its subcellular distribution using immunostaining with an anti-FLAG antibody. The
experimental results showed that, while the control fluorescent multiple cloning site (MCS) was
distributed throughout the cell, the fluorescent protein-tagged Bcnt/Cfdp1 was predominantly
localized in the nuclei (Figure 1B-1).

The density of the nuclear interior is recognized as a porous medium composed of the chromatin

network [6]. The chromatin density is heterogeneous; the inactive part, heterochromatin, is

denser, whereas the active part, euchromatin, is sparser. Using high-resolution imaging, we

observed mCherry Bcnt/Cfdp1 and DAPI in the nucleus of C2C12 myoblasts and found their

segregation pattern; mCherry Bcnt/Cfdp1 was present in the area with weaker DAPI signals

(Figure 1B-2).

To gain insights into the intranuclear dynamics of Bcnt/Cfdp1, we examined its potential

involvement in forming the condensates in the nucleus using the N-terminal BCNT/CFDP1

IDR. Nevertheless, the results were in vain (Supplementary Figure 1S, which will be discussed

later). Furthermore, the fluorescent protein-tagged Bcnt/Cfdp1 remained stable during

prolonged culture, and multiple signals were visible in the nuclei (Figure 1C). These results

indicated that fluorescent protein-tagged Bcnt/Cfdp1 was detected primarily in the nuclei of

C2C12 myoblasts and differentiated multinucleated myotubes.

Validity/Invalidity of Western Blot Monitoring in C2C12 Cell Subcellular Fractionation

To determine the subcellular distribution of endogenous Bcnt/Cfdp1 in cells, we investigated

Bcnt/Cfdp1 through the biochemical fractionation of C2C12 and HEK 293T cells, monitored by

Western blot analysis, followed by mass spectrometry. Our previous study demonstrated that

the Western blotting signal of Bcnt/Cfdp1 corresponds to a doublet of ~43 kDa (previously

described ~45 kDa) bands in mouse ES cell lines Cfdp1-K1 and mouse brain extracts using

A305-624A-M as an anti-CFDP1 antibody [37]. We confirmed that the ~43 kDa signal band

represents Bcnt/Cfdp1 by enriching the immunoreactive band from Cfdp1-K1 cells using two-

column chromatography followed by mass spectrometry (Supplementary Figure S2 and

Supplementary Table S1). Next, we validated the Western blot signal(s) of C2C12 myoblast

extracts obtained with A305-624A-M (Figure 2A-D). The treatment of an endonuclease-derived

siRNA (esiRNA) against mouse *Bcnt/Cfdp1* significantly decreased the signal of the ~43 kDa

band, but the effects on the lower-mobility band(s) were lesser (Figure 2B and 2C). To confirm

whether the lower-mobility band is derived from Bcnt/Cfdp1, which was previously referred to

as the "upper band" corresponding to its phosphorylated form [10, 33], we expressed and

isolated FLAG-tagged mouse Bcnt/Cfdp1 in C2C12 cells (Figure 2D and Supplementary Figure

S3). A small upper band appeared in C2C12 cells expressing FLAG-tagged Bcnt/Cfdp1, but its

phosphorylated peptide at serine 245 in the ~43 kDa band [33] was not captured (data not

shown). These results conclude that the upper band exists, but its signal region contains both

Bcnt/Cfdp1-derived molecule and off-target(s), which will be discussed later.

Compared to the cytoplasmic soluble fraction, the particulate fraction exhibited several off-

targets unaffected by esiRNA treatment (Figure 2C). In addition, during myotube

differentiation, the distinct 50 kDa signal gradually increased, whereas Bcnt/Cfdp1 levels

decreased in the later stages (Figure 2E and Supplementary Figure S4). These results indicate

that Bcnt/Cfdp1 can be effectively monitored in the cytoplasmic soluble fraction of C2C12

myoblasts by Western blotting. However, tracking the target in the nuclear fraction and

differentiated cell extracts, including the upper band(s), may not be feasible. Consequently, we

focused on the cytoplasmic soluble fraction.

Distribution of Bcnt/Cfdp1 by Subcellular Fractionation

We performed biochemical fractionation using a widely utilized kit from Thermo Scientific,

which may include different detergents in each step [38]. C2C12 or HEK293T cell suspensions

were serially fractionated according to the manufacturer's protocol, with additional washes at

each step. The five resulting fractions were subjected to Western blotting analysis using

antibodies against Rasa1 (p120 Ras GTPase activating protein) and Gapdh (glyceraldehyde-3-

phosphate dehydrogenase) as markers of the cytoplasmic extract, Vdac1 (voltage-dependent

anion channel 1) for the membrane extract, NonO (non-POU domain-containing octamer-bound

p54nrb) for the soluble nuclear extract, and HstH3 (histone H3) as a marker of the chromatin-

bound fraction. Nestin, a linker protein that interacts with the cytoskeleton, was also used and

detected in the cytoskeletal fraction of C2C12 cells and the cytoplasmic fraction of HEK293T

cells (Figure 3). Under these fractionation conditions, a substantial amount of the ~43 kDa band

was detected in the cytoplasmic fraction, while the nuclear fraction contained other bands (s) of

~50 kDa (Figure 4A). We subjected these five fractions of C2C12 myoblasts to quantitative

mass spectrometry to evaluate the fractionated preparations quantitatively. A total of 4,487

proteins were identified, of which 3,280 were eligible to evaluate relative distribution

(Supplementary Table S2). These included five biomarkers (Figure 4B), heterogeneous nuclear

ribonucleoproteins, and three paraspeckle proteins (Figure 4C). In sequential fractionation, the

components from the previous fractions were easily carried over into the following fraction.

Nevertheless, a significant amount of Bcnt/Cfdp1 was extracted in the first fraction, the

cytoplasmic fraction (Figures 4A and 4B). These results indicated that Bcnt/Cfdp1 is primarily

present in the cytoplasmic fraction of C2C12 cells using an available biochemical subcellular

fractionation kit and suggested that this is also the case for HEK293T cells (Figure 3, right

panel).

However, this result was in contrast to the above finding that the Bcnt/Cfdp1 was predominantly

observed in the nucleus using fluorescent protein labeling and immunocytochemical techniques

(Figures 1A–C). To gain insight into this intracellular Bcnt/Cfdp1 mobility during the

subcellular fractionation processes in C2C12 cells, we searched for proteins that showed similar

distribution to Bcnt/Cfdp1; the proteins annotated or reported to be present predominantly in the

nucleus are primarily detected in the cytoplasmic fraction. Of the 3,280 proteins available for quantitative evaluation, 472 showed high relative occupancy (>0.6) in the cytoplasmic fraction, of which six proteins were annotated as present in the nucleus (Figure 4D). They were IDR-containing proteins, such as PEST proteolytic signal-containing nuclear protein, Pcnp [39]. Based on further curation using the integrated cellular localization prediction software BUSCA [40] and experimental evidence from human ortholog data [41], these proteins may localize to the cytoplasm (Supplementary Table S3).

Intracellular Behavior of Bcnt/Cfdp1 in Digitonin-Mediated Release

To further investigate the cytoplasmic distribution of Bcnt/Cfdp1 strictly, C2C12 and HEK293T cell cultures on dishes were directly treated with digitonin to obtain the cytoplasmic soluble proteins. Digitonin is a unique detergent that selectively binds to the cholesterol in membranes, creating pores in cholesterol-rich membranes that allow macromolecules to pass through [42]. In addition, it has been observed that digitonin does not flip across membranes with low cholesterol content, such as the nuclear envelope [43]. Therefore, digitonin is currently one of the most appropriate methods for isolating the cytosolic proteins of mammalian cells while exerting minimal effects on the intracellular behavior of other proteins, provided that the treatment conditions are appropriately determined.

A) In an isotonic solution

C2C12 myoblasts on the dish were exposed to isotonic solutions containing different concentrations of digitonin. The released eluate and corresponding residual cell contents were collected and subjected to Western blotting analysis. A prominent band of ~43 kDa was observed in the eluate at a digitonin concentration of 25 μg/mL, showing a similar pattern to that of the cytoplasmic soluble proteins Rasa1 and Gapdh (Figure 5A). The ~43 kDa band was excised from the gel and subjected to mass spectrometry analysis (Figure 5B). The data showed that Bcnt/Cfdp1 was ranked high in the Sum Pep Score of 124 identified proteins annotated as cytoplasmic proteins except for nuclear protein Trip13 (Supplementary Table S4 and Figure 5C). However, the human ortholog TRIP13 has been reported to be distributed diffusely in the cytoplasm, in addition to the nucleus [44]. Thus, these results support the cytoplasmic distribution of Bcnt/Cfdp1 in C2C12 myoblasts.

Furthermore, we developed a three-step strategy to enhance the collection of cytoplasmic components, as shown in Figure 6. The supernatant released from the digitonin-treated cells was initially collected, centrifuged, and labeled Cyt#1. Subsequently, the cell debris on the dish was gently scraped off, and the supernatant was collected by centrifugation (Cyt#2). Next, the supernatant was obtained by washing the pellet with an isotonic solution (Cyt#3). Following the

concentration of each fraction using trichloroacetic acid (TCA), the proteins were dissolved in sodium dodecyl sulfate (SDS)-containing buffer. Equal relative volumes of each fraction were subjected to Western blotting analysis, as shown in Figure 3. The elution pattern of Bcnt/Cfdp1 was similar to that of the cytoplasmic soluble marker proteins, such as Rasa1 and Gapdh (Figure 5D).

B) In Ficoll/sucrose solution

Macromolecular crowding has been shown to affect protein properties in living cells [2, 45], nuclear structure and function in preparation [46], and subcellular distribution in subcellular fractionation [47]. Therefore, digitonin treatment was performed in the HEPES-buffered Ficoll/sucrose solution (HFS) instead of an isotonic solution to prepare the cytoplasmic components from the cells appropriately. Cyt#1–3 fractions were prepared using the above three-step process (Figure 6) and subjected to Western blotting analysis. Notably, the elution pattern of Bcnt/Cfdp1 differed from that of the cytoplasmic soluble marker proteins (Figure 7A). A significant quantity of Bcnt/Cfdp1 was detected within the Cyt#3 fraction, whereas only a tiny amount was observed within the Cyt#1 and #2 fractions at lower digitonin concentrations. A comparative preparation was conducted using direct digitonin-mediated release in the HFS solution on HEK293T cells grown on polylysine-coated dishes (Figure 7B). The distribution pattern of BCNT/CFDP1 in HEK293T cells was similar to that observed in C2C12 cells. A significantly higher concentration of BCNT/CFDP1 was also observed in the Cyt#3 fraction than in the Cyt#1 and #2 fractions (Figure 7A). These results indicated that macromolecular crowding affects the mobility of Bcnt/Cfdp1 in C2C12 and HEK293T cells during the three-step preparation, suggesting that Bcnt/Cfdp1 exhibits flexible behavior in the process.

Mobility of Bcnt/Cfdp1 in the Cytoplasm

We expected that the differential elution pattern of digitonin-mediated release in the HFS solution would reflect the mobility or integrity of each protein in the above three-step preparation. To examine the perspective of the property, we prepared Cyt#1 and #2* similarly to Figure 7, except that the pellet of Cyt#2 was washed with HFS solution instead of isotonic solution, and the washed supernatant was combined and designated as Cyt#2* (Figure 8A). These two fractions were subjected to quantitative LC-MS/MS analysis, and 2472 proteins were identified, of which 2442 were qualified for ratio evaluation (Supplementary Table S5). Among them, 51 glycolysis/glucogenesis- and TCA-related enzymes were listed, and their Cyt#1 to Cyt#2* ratios (Cyt#1/2*) were plotted as LOG2 (Figure 8 B and C, and Supplementary Table S6). The ratios of glycolysis-related proteins were above 1 (>0 in LOG2 unit), but TCA cycle-

278   or membrane-associated proteins had ratios significantly below 1 (< 0 in LOG2), while the

279   Bcnt/Cfdp1 ratio was 0.17 (-2.55 in LOG2) (Supplementary Table S5). Remarkably, the four

280   isoenzymes of cytoplasmic and mitochondrial types revealed clear property trends.

281   Table 1. Ct#1/2* ratios of four representative glycolysis/TCA-related isomers

Protein Names	Gene Symbol (Alias)	Intracellular localization	Mw (kDa)	Cyt#1/2* (Ratios)	Cyt#1/2*(LOG2)
Bucentaur/Craniofacial development protein1	Bcnt/Cfdp1 (cp27)		33.4	0.17	-2.56
Aconitase	Aco1	cytoplasmic	99	2.27	1.18
	Aco2	mitochondrial	85.4	0.072	-3.80
Enolase	Eno1	cytoplasmic	47.1	2.36	1.24
	Eno2	mitochondrial	37.2	0.055	-4.18
	Eno3	cytoplasmic	56.1	2.33	1.22
Aspartate aminotransferase	Got1	cytoplasmic	46.2	2.41	1.27
	Got2	mitochondrial	47.4	0.03	-5.06
Malate dehydrogenase	Mdh1	cytoplasmic	40	1.835	0.88
	Mdh2	mitochondrial	35.6	0.038	-4.72

Cyt#1 and 2* were prepared from the digitonin-mediated cytoplasmic fractions of C2C12 cells using

a three-step process, as shown in Figure 6. Each fraction was analyzed by quantitative mass spectroscopy.

The Cyt#1/2* ratios of three representative isomers were listed and compared.

Furthermore, the ratios Cyt#1/2* of 321 RNA-binding proteins selected based on published data

were significantly smaller than that of the entire population (p-value <0.01) (Figure 8C and

Supplementary Table S7). These results support the differential elution patterns of each protein

and reflect the mobility or integrity of each protein during the three-step preparation, which is

affected by macromolecular crowding.

DISCUSSION

Although the lack of high-quality anti-Bcnt/Cfdp1 antibodies made the analysis difficult in this

study, reliable data were obtained by quantitative mass spectrometry. Nuclear fractions and

extracts of differentiated myotubes of C2C12 cells showed several off-target bands of Western

blots using A305-624A-M as the anti-CFDP1 antibody [33], which is currently considered the

most reliable antibody in our Western blot analysis. These difficulties forced us to focus on the

cytoplasmic soluble proteins, which did not allow us to take advantage of the characteristics of

C2C12 cells. In this study, cytochemical approaches showed that Bcnt/Cfdp1 was mainly

observed in the nucleus, whereas biochemical fractionation detected the molecule at high levels

in the cytoplasmic fraction. Here, we discuss these contradictory results from the viewpoints of

nuclear export trafficking and macromolecular crowding and also address the potential

functionality of Bcnt/Cfdp1 in the Srcap-C chromatin frame.

Bcnt/Cfdp1 Subcellular Distribution Determined by Cytochemical Approaches

As reported in a previous study, we showed that tag-free mouse Bcnt/Cfdp1 is predominantly

localized in the nuclei of human HEK293T cells using antibodies against mouse Bcnt/Cfdp1

peptides (Figure 1A, modified from Nakashima *et al*. [37]). This strategy was excellent, but

these antibodies react with off-targets in Western blotting in HEK293T and C2C12 cells.

Therefore, we used FLAG- and fluorescent protein-tagged Bcnt/Cfdp1 to assess its subcellular

localization. The results showed that the signal was predominantly observed in the nuclei of

C2C12 myoblasts and differentiated myotubes (Figure 1B-1 and 1C). Furthermore, using

higher-resolution imaging in myoblast, Bcnt/Cfdp1 was detected in a segregated pattern, the

regions with the lower DAPI staining (Figure 1B-2). The result is consistent with the

observation that the *Drosophila* homolog YETI has been mapped to multiple sites on the

euchromatic region on salivary gland polytene chromosomes [11].

When SRCAP-C is prepared at a higher-salt extraction buffer such as 300 mM KCl,

BCNT/CFDP1 is scarcely captured as its component among other subunits [13, 48], which is, in

part, why Bcnt/Cfdp1 has often been neglected [11]. However, a recent study revealed that

BCNT/CFDP1 was significantly recovered when a lower-salt extraction buffer (150 mM NaCl)

was used for isolation from the reconstituted system composed of recombinant proteins

(personal communication, Dr. Shinya Watanabe, University of Massachusetts Medical School).

This finding suggests that either weak ionic interactions are critical for BCNT/CFDP1 to be a

component of the SRCAP-C or that high salt concentrations disrupt the specific environment,

such as a condensate, in which Bcnt/Cfdp1 is incorporated into SRCAP-C. The N-terminal

peptide of SWC5 was shown to have a potential for condensate formation through molecular

dynamics simulations [49]. Although the amino acid sequence of the N-terminal regions is quite

different between SWC5 and BCNT/CFDP1, both are IDRs rich in negatively and positively

charged amino acid residues. Thus, we explored the possibility of condensate formation of

BCNT/CFDP1 in the nucleus. The IDR of BCNT/CFD1 (the N-terminal 273 of 299 whole

amino acids) was inserted into an optoDroplet assay vector containing a nuclear localization

signal, and condensate formation was monitored [50] (Supplementary Figure S1). However, no

positive results were obtained.

Considering the findings of the higher-resolution imaging in which Bcnt/Cfdp1 is detected in

the lower chromatin density regions and the fact that the interchromatin is a porous medium [6],

Bcnt/Cfdp1 may dynamically behave in the nucleus for coupling with Srcap-C. Further insights

into the physical properties of Bcnt/Cfdp1 are expected in the context of weak ionic interactions

in Srcap-C formation.

Nonetheless, considerable concerns remain with respect to the cytochemical results. First, we

could not correctly evaluate the weak fluorescent signals in the cytoplasm, which were observed

at the level of fluorescence microscopy. However, it was challenging to visualize after

paraformaldehyde (PFA) fixation and subsequent anti-FLAG antibody detection, which are used

to minimize artifact signals. Therefore, to accurately evaluate these weak signals, considering

the side effects of exogenous protein expression, improvements in fixation and visualization

methods, including expression systems that can control gene copy numbers, are necessary.

Second, the fluorescent protein tag, corresponding to a mass of approximately 27 kDa, may

inhibit passive nuclear export [8], resulting in overestimating Bcnt/Cfdp1 nuclear localization.

Thus, it remains essential to prepare higher-quality anti-Bcnt/Cfdp1 antibodies.

Effectiveness and Limitations of Western Blotting Monitoring

We employed A305-624A-M as an anti-Bcnt/Cfdp1 antibody to effectively monitor the ~43

kDa band in Western blotting. The presence of Bcnt/Cfdp1 molecules in the ~43 kDa region

was confirmed by enrichment followed by mass spectrometry (Supplementary Figure S2 and

Supplementary Table S1). However, in contrast to the cytoplasmic soluble proteins, the

particulates and the extract of differentiated C2C12 myotubes showed several off-target bands

in the wide range of ~ 50 kDa (Figures 2C, 4, and 8). We reported similar bands in the

precipitate fraction of T-Rex cell extracts solubilized in Radio-Immunoprecipitation Assay

(RIPA) buffer (Supplementary Figure S4 in Iwashita *et al.* [33]). During myotube

differentiation, another signal of ~ 50 kDa gradually increased, whereas Bcnt/Cfdp1 levels

decreased in the later stages (Figure 2E and Supplementary Figure S3). The latter was consistent

with mRNA expression levels [51] (courtesy of Dr. Yan Fei Gao). Concerning the ~43 kDa

doublets, which are caused by phosphorylation in T-Rex cells (a HEK293 cell derivative) at the

250th serine residue of BCNT/CFDP1 [10] or the 245th serine residue of mouse Bcnt/Cfdp1

[33], its intensity in C2C12 cells was too weak to be accurately assessed. Taking these results

together, we focused on their cytosolic extracts without mentioning the doublets.

Subcellular Fractionation Using the Available Fractionation Kit

We first performed subcellular fractionation using a Thermo Scientific kit containing multiple

non-ionic detergents in each step [38]. Using the process, significant bands of 43–45 kDa were

observed in the cytoplasmic fractions of both C2C12 and HEK293T cells, and other bands of

~50 kDa were detected in the respective nuclear fractions (Figure 3). The presence of

Bcnt/Cfdp1 in the cytoplasmic fraction was confirmed via mass spectrometry (Figure 5C from

Supplementary Table S2). Non-ionic detergents are often combined with a hypotonic solution to

achieve optimal results in sequential subcellular fractionation and have been reported to be
particularly effective in separating the cytoplasmic and nuclear compartments [52]. However,
the possibility that non-ionic detergents, even at low concentrations, affect lipid-lipid and lipid-
protein interactions, affecting the membrane-associated population of the targets and
compromising membrane integrity cannot be ignored.
The nucleus is protected by a fibrous layer called the lamina, which lies beneath the nuclear
envelope [53]. Therefore, the apparent intactness of the nucleus after various treatments does
not necessarily guarantee the integrity of the nuclear envelope. The nuclear membrane is much
more elastic than previously thought. Its integrity, even its rupture, is influenced by a
physiological response [54] and through signaling pathways, such as the activation of Akt
kinase [55]. Regarding the distribution of Bcnt/Cfdp1, it is interesting that HMGB1 was
detected in the cytoplasm using the non-ionic detergent IGEPAL (Nonidet P-40), but in the
nucleus using the Dounce homogenization method in lung cancer cell lines [56]. The
discrepancy may be attributed to the properties of HMGB1 that include nuclear localization
signals in the N-terminal and IDRs in the C-terminal. Notably, blocking the nuclear localization
signals of HMGB1 in leucocytes redirects it to the cytoplasm, eventually leading to secretion
[57]. In our mass analysis, Hmgb1 was mainly detected in the cytoplasmic fraction in the
subcellular fractionation of C2C12 and annotated in most compartments, including the non-
structural extracellular matrix (Supplementary Table S2). HMGB1, found initially as a nuclear
protein, may have properties similar to those of Bcnt/Cfdp1 in flexible intracellular behavior.
Regarding the possibility of Bcnt/Cfdp1 secretion, it was reported as an unconventionally
secreted protein in the conditioned medium of C2C12 myotubes [58]. Furthermore, a secretome
prediction tool supported the result: OutCyte, http://www.outcyte.com/(courtesy of Dr. Gereon
Poschmann) [59]. Although we tried to confirm the possibility, obtaining the C2C12 cell culture
medium supernatant was difficult even in myoblast cultures, as dead cells constituted at least
1% of the total. Thus, it was impossible to reach a definitive conclusion, and further research is
needed.
401
Digitonin-Mediated Extraction in Isotonic or Ficoll/Sucrose Solution
To examine Bcnt/Cfdp1 in the cytoplasm more strictly, we directly prepared the cytoplasmic
soluble proteins from cell cultures on the dishes by digitonin treatment. The method was
expected to allow the extraction of cytosolic proteins with minimal interference from other
proteins. In the experiment, Bcnt/Cfdp1 was detected along with the cytoplasmic soluble
proteins Rasa1 and Gapdh in isotonic buffer at a concentration of 25 μg/mL (0.0025%) (Figure
5A and Supplementary Table S4). Notably, a nuclear protein, Trip13 (thyroid hormone receptor
interactor 13), was released with the digitonin treatment and showed similar mobility to

Bcnt/Cfdp1 (Supplementary Table S2 and S4). Indeed, the human ortholog TRIP13 has been

reported to be significant trafficking in the cytoplasm and the nucleus [44]. These results

support the cytoplasmic distribution of Bcnt/Cfdp1 and the specific proteins that exhibit similar

molecular mobility. On the other hand, it was difficult to obtain reproducible release patterns at

lower digitonin concentrations because of the tailing of cytoplasmic soluble marker proteins

under the direct elution method, especially in the HFS solution.

Since macromolecular crowding is critical for molecular behavior [2, 45], we referred to the

following reports to examine the subcellular behavior and distribution of Bcnt/Cfdp1. First,

solutions without macromolecular crowding may not fully preserve the nuclear structure and

function during nuclear isolation [46], and second, the intracellular trafficking of SIRT1 may

depend on macromolecular crowding [47]. Therefore, according to the report, we used the HFS

solution instead of an isotonic solution in the digitonin treatment [47].

Notably, Bcnt/Cfdp1 was distinctly extracted compared to the cytoplasmic soluble proteins,

especially Rasa1, appearing in the Cyt#3 fraction rather than in the Cyt #1 and Cyt #2 fractions

(Figure 7). Cyt#3 was the supernatant fraction obtained by scraping the digitonin-treated cell

debris and washing the pellet with an isotonic solution, followed by centrifugation (Figure 6). A

comparable extraction pattern was also observed at a higher digitonin concentration of 250

μg/mL in C2C12 cells and 50 μg/mL in HEK293T cells.

We prepared the fractions of Cyt#1 and Cyt#2*, the latter obtained by washing the pellet of

Cyt#2 with HFS solution and combining the washed supernatant (Figure 8). The mass analysis

of Cyt#1 to #2* of ratios reflected the mobility of each protein in the cytoplasm. The ratio of

Bcnt/Cfdp1 was 0.17 (-2.55 in LOG2), suggesting that Bcnt/Cfdp1 exists in a complex in the

cytoplasm, probably associated with RNA or RNA-binding proteins [28]. This result reflects an

essentially similar phenomenon, as reported; the subcellular distribution of SIRT1 differs

depending on whether the fractionation solution is composed of macromolecular densities [47].

Although the rate at which the nuclear membrane is partially damaged, resulting in leakage out

of the nucleus during the three-step preparation, should not be underestimated, we propose that

Bcnt/Cfdp1 and some specific proteins have physical properties that allow them to be readily

mobile from the nucleus to the cytoplasm, which is affected by macromolecular crowding.

These findings have practical implications for future research and applications in protein

behavior.

Beyond the Limitations of Digitonin Treatment -Conflicting Results for the Subcellular

Localization of Bcnt/Cfdp1

The discrepancy in the subcellular distribution of Bcnt/Cfdp1 described above is a complex and

vital issue that presents significant challenges. As mentioned earlier, several factors could lead

to discrepancy. Two situations lead to overestimating nuclear localization: the cytochemical

approach could not capture the weak signals in the cytoplasm, and the fluorescent protein tag

could inhibit passive nuclear export. On the other hand, mobility due to nuclear membrane

damage in the digitonin-mediated preparation could lead to the overestimation of cytoplasmic

localization.

It was challenging to completely recover the cytoplasmic soluble proteins released by direct

digitonin treatment, especially in an HFS solution. Subsequently, we developed a three-step

strategy for the preparation of digitonin-mediated extracts. The Cyt#3 fraction obtained by

washing the digitonin-mediated cell debris pellet contains the proteins of the residual

cytoplasmic population, which may include the attached one to various organelle membranes. In

addition, proteins present in the nucleus may leak out of the nucleus due to damage to the

nuclear membrane during the three-step procedure. However, given the relatively selective

activity of digitonin in membrane poration, which depends on the cell type and physiological

state, it is inherently challenging to avoid partial membrane damage during digitonin treatment

to harvest the cytoplasmic components completely. For example, although digitonin treatment

can be mild, i.e., using 0.001% in phosphate-buffered saline (PBS) for 2 min on ice, it still

caused partial damage to the nuclear envelope, as evidenced by staining with an anti-lamin B1

Ab [60]. In addition, digitonin is capable of disordering the alkyl chains of phospholipid

monolayers, and its effect is more potent on phosphatidylethanolamine and phosphatidylserine

than phosphatidylcholine [61]. Orre *et al.* performed a large-scale analysis of subcellular

localization in five cancer cell lines using a uniform fractionation method that involved the

digitonin treatment of cell cultures on dishes, followed by homogenization in a hypotonic buffer

[41]. Whereas the data showed 80% agreement between the cell lines, significant discrepancies

were attributed to the differences in the localization of proteins, including BCNT/CFDP1,

between the cytosol and nucleus (https://lehtio-lab.se/subcellbarcode/). This discrepancy might

be due to the different susceptibility of the nuclear envelope in some cells than in others during

the fractionation or due to other factors, such as differences in the levels of post-translational

phosphorylation in each cell. Furthermore, the unique intracellular mobility that we observed on

Bcnt/Cfdp1 and some proteins may also contribute similarly.

Beyond these difficulties of evaluation in digitonin-mediated preparation, our data suggest that

Bcnt/Cfdp1 and some proteins are readily mobile from the nucleus. Recent studies have

revealed unique features for the nuclear membrane compared to other cell membranes; nuclear

lipid membranes are much more elastic than other membranes [62], and their lipid composition

is closely related to membrane integrity [63]. Additionally, NPC, the sole transport gateway

between the cytoplasm and nucleus, comprises approximately 300 IDPs, rendering the pore a

flexible rather than a rigid structure. Interestingly, while folded protein transport has revealed a

size-dependent exclusion, where larger proteins are restricted, IDPs can diffuse through the

NPC without energy or specific transport receptors due to their lack of a fixed structure,

allowing them to navigate the pores more efficiently [9]. These results support the possibility

that Bcnt/Cfdp1 and some specific proteins may be able to diffuse rapidly through the NPC into

the cytoplasm during the subcellular fractionation. However, leakage due to partial damage to

the nuclear envelope cannot be underestimated in the three-step processes.

Biological Functionality of Bcnt/Cfdp1

Bcnt/Cfdp1 plays a pivotal role in developmental processes, as evidenced by its mutants in

Drosophila [20], zebrafish [21, 22], and mice [23, https://www.jax.org/strain/033834]. These

biological variations include alterations to the cell cycle, cell migration, and apoptosis, which

subsequently influence cancer prognosis and susceptibility [28, 29].

The zebrafish mutant *gazami*, lacking the seventh exon of Bcnt/Cfdp1 (encoding the C-terminal

31 amino acid residues), exhibited abnormal neuronal differentiation in brain granule cells. The

elevated expression levels of cyclin B1 (*Ccnb1*) have been shown to contribute to inappropriate

differentiation [21]. However, its upregulation might be one of the gene expression variations

induced by deficient *cfdp1*, similar to the *Bcnt/Cfdp1* knocked down in the Cfdp1-K1 cells [33].

Interestingly, many genes related to the Wnt or Akt pathways and apoptosis/autophagy, such as

Wls, *Wnt4*, *Anxa2*, *Bcl2*, and *Akt2,* are included, up- or downregulated in Cfdp1-K1 cells

(Supplementary Table S8). They are involved in cell cycle control, cell migration, and

apoptosis. Indeed, the Wnt signaling pathway is suppressed specifically in extraluminal

mesenchymal cells of the valves of zebrafish Bcnt/Cfdp1-deficient embryos [22].

The broad effects of deficient *Bcnt/Cfdp1* may be comparable to natural genetic variations in the

fission yeast *swc5*. The mutant exhibited abnormal mRNA, antisense, and noncoding RNA

expression, possibly caused by improper histone exchange reactions [34]. Srcap-C has been

demonstrated to be involved in differentiation processes in intestinal epithelial cells controlled

by H2A.Z in association with Wnt signaling [64]. Assuming that Bcnt/Cfdp1, similarly to

SWC5 in SWR1-C, is a key factor for Srcap-C activity [14-16], a deficiency in **Bcnt/Cfdp1** may

induce inappropriate histone exchange reaction activity, resulting in changes to the

transcriptional landscape by Srcap-C. The phenotype of the mutant *gazami* described above may

be a clear example.

Notably, some reports of the biological responses of perturbed *BCNT/CFDP1* mentioned earlier

are partially associated with *BCAR1***,** a proximal neighboring gene of *BCNT/CFDP1* located on

chromosome 16q23.1 and has multiple functions [65]. The *BCAR1–CFDP1–TMEM170A* locus,

which is highly conserved across vertebrates, contains several single nucleotide polymorphisms

(SNPs) associated with diseases, such as vascular diseases [66] and Dupuytren's disease [67].

Although these diseases may be highly related to variations in the Wnt/Notch/Akt signaling

pathways, the causal relationship between these SNPs and diseases remains unclear.

CONCLUSION

Metazoan Bcnt/Cfdp1 is predicted to have a similar function to yeast SWC5 as a component of

Srcap-C. Consistent with this prediction, both untagged mouse Bcnt/Cfdp1 transfected into

HEK293T cells and fluorescent protein-tagged mouse Bcnt/Cfdp1 in C2C12 cells were detected

primarily in the nucleus by cytochemical approaches. However, by detergent-based biochemical

fractionation of C2C12 and HEK293T cells, endogenous Bcnt/Cfdp1 was detected mainly in the

cytoplasmic fraction. Furthermore, when cell cultures on the dish were treated with digitonin,

Bcnt/Cfdp1 was released and detected as a cytoplasmic protein, while different elution patterns

were observed between isotonic and Ficoll/sucrose solutions. Although our results on the

intracellular behavior of Bcnt/Cfdp1 reveal its mobility in the presence of detergent rather than

the export trafficking, we argue that Bcnt/Cfdp1 and some specific proteins have physical

properties that allow them to be readily mobile from the nucleus to the cytoplasm, which is

affected by macromolecular crowding. The unique property of Bcnt/Cfdp1 intracellular mobility

remains unclear, but its elastic properties at the cytoplasm/nucleoplasm interface may be

relevant. These issues are the most critical aspects of this study and are expected to become

more evident as our understanding of the nuclear membrane expands beyond classical cell

membrane studies [68].

Data Availability Statement

540　All data reported in this paper, including the proteomics mass spec dataset, are available without

541　restriction.

MATERIALS AND METHODS

544　Supplementary Table S9 provides detailed information on the material resources used in the

545　study, except for some experiments mentioned therein.

Cell Culture

C2C12 cells were obtained from the European Collection of Authenticated Cell Cultures through KAC Co., Ltd. (Kyoto), expanded, and aliquots were stored at –80°C using Bambanker. HEK293T cells were provided by Dr. Rie Komori (Tokushima Bunri University) and stored in liquid nitrogen using Cellbanker. Both types of cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM)–10% FCS and subcultured using Accutase, as described previously [33]. Briefly, after incubation for 5 min at room temperature, the cells were gently homogenized using a 1000 μL tip, mixed with medium, centrifuged, and resuspended in the medium unless otherwise specified. The cell numbers were counted using a hemocytometer (a disposable cell counting plate of the Improved Neubauer Type), and dead cells were estimated via the trypan blue exclusion assay. Before seeding, the dishes were incubated with the medium in a CO_2 incubator for at least 30 min.

C2C12 Differentiation

C2C12 cells (5×10^4 per 35 mm dish) were cultured for 2 days, and then the medium was replaced with DMEM–2% horse serum (designated as day 0). On the corresponding day, the cultures were washed with chilled HEPES buffered saline (HBS) twice and homogenized in 100 μL Lysis buffer (LS:1% SDS, 1 mM EDTA in 10 mM HEPES–NaOH, pH 7.5) supplemented with inhibitors of proteases and phosphatases using a silicon policeman. Each extract was collected in a 1.5 ml tube, boiled immediately for 3 min, and then sonicated for 2.5 min. A constant volume of each extract was subjected to 12.5% SDS-polyacrylamide gel electrophoresis (SDS/PAGE), followed by Western blot analysis.

Coating Culture Dishes with Polylysine

According to the protocol, 60 mm dishes were coated with polylysine with some modifications (https://assets.thermofisher.com/TFS-Assets/BID/Flyers/gibco-poly-d-lysine-flyer.pdf). Each dish was shaken with 1 mL of 50 μg/mL poly-d-lysine solution on a seesaw shaker for 1 h at room temperature. To examine fluorescent protein-tagged mBcnt/Cfdp1, a commercially available 4-well polylysine-coated chamber (Matsunami, Osaka) was used.

Construction of Plasmids

All Polymerase Chain Reaction (PCR) primers are listed in Supplementary Table S10. Tag-free and FLAG-tagged Bcnt/Cfdp1 were generated as previously described [37]. The same primers were used to generate mCherry- and EGFP-tagged MCS or mouse Bcnt/Cfdp1 using their respective expression vectors (NCBI accession numbers LC311025, LC311020, LC311024, and LC311019).

Transfection of Plasmids

Transfections were carried out using Lipofectamine 3000 according to the manufacturer's protocol with minor modifications. Prior to transfection, each half of the medium was saved once, and the plasmids were transfected. After 4 h, the saved medium was returned, and the cells were cultured for 20-44 h. For the fluorescent protein observation, C2C12 cells were seeded at a density of 1×10^4 cells per well in 0.75 mL medium per well of polylysine-coated chamber, which was pre-incubated with culture medium in a CO_2 incubator. After 20 h culturing, the medium was replaced with 0.5 mL of antibiotic-free medium, and cells were transfected with 0.25 μg of each plasmid using Lipofectamine 3000 at a DNA-to-Lipofectamine ratio of 1:2. Following 4 h of culture, 0.5 mL of medium containing 20% FCS was added to each well. The cells were then cultured for an additional 20 h.

Fluorescent Proteins by Immunocytochemistry

The transfected and further cultured cells were washed twice with cold HBS and fixed with a final concentration of 4% PFA by dilution in 0.1 M phosphate buffer (pH 7.2). The cells were washed with 10 mM PBS to remove PFA. All the blocking serum, antibodies, and nuclear staining dye were used at the indicated dilutions with 10 mM PBS containing 0.2% Triton X-100. The cells were blocked with 5% normal donkey serum for 30 min at room temperature, then incubated with the anti-DYKDDDDK tag monoclonal Ab (1:1000) at 4°C overnight. The immunoreactivity was visualized with Cy3-conjugated donkey anti-mouse IgG (1:200) or CF™488A-conjugated donkey anti-mouse IgG (1:200). DAPI (1 μg/mL) was used for nuclear staining. The immunostained cells were examined by phase contrast and fluorescence microscopy (TE2000-U, Nikon).

Enrichment of ~43 kDa Band by Western Blot Monitoring

A puromycin-resistant cell line of AB2.2 mouse embryonic stem cells was gifted from Dr. Tsutomu Kumazaki (Tokushima Bunri University), who obtained the original AB2.2 cells from the Sanger Institute (Cambridge, UK) and generated the cell line by transfection with pMSCVpuro plasmid using PolyFect. The cells were expanded under serum- and feeder-layer-free culture conditions as described in [33] without G418 and puromycin. The cells were harvested using Accutase and rinsed with chilled HBS. The cell suspension aliquots were snap-frozen in liquid nitrogen and stored at −80°C until use. For the experiment, the cell suspension (equivalent to 3×10^7 cells) was thawed and resuspended in 300 μL of Solution 1 (a cytoplasmic extraction buffer from Atto, Tokyo). The suspension was homogenized in a Biomasher II, rotating the pestle 10 times, sonicated in an ice-water bath (10 s, three times), and then centrifuged at $25,000 \times g$ for 30 min at 4°C. Subsequently, CHAPSO ($C_{32}H_{58}N_2O_8S$) was added

to the supernatant at a final concentration of 1% (w/v). The extracts were slowly loaded onto

heparin-agarose (200 μL bed volume in a spin column) by cycling three times after the resin had

been equilibrated with Solution 1. Nonidet-P40 was included in all buffers at a final

concentration of 0.05% v/v, which was utilized for the subsequent chromatographic procedures.

After washing the resin with 20 mM Tris-NaOH, pH 7.5, the bound proteins were eluted twice

with 200 μL of Tris buffer containing different concentrations of NaCl. Each elution was

allowed 5 min to settle at each elution step. The eluates from the heparin column

chromatography with 0.05–0.1 M NaCl were combined and loaded onto Phos-tag resin (150-μL

volume in a spin column) by three cycles. The resin was washed thrice with 150 μL of wash

buffer, followed by a buffer containing 1M NaCl. The bound proteins were released using 4 ×

100 μL of phosphate-containing elution buffer without Nonidet-P40 following the

manufacturer's protocol (https://labchem-

wako.fujifilm.com/jp/category/docs/01072_doc01.pdf). The eluates were concentrated by

deoxycholate-TCA precipitation, as described in a standard protocol

(https://www.sigmaaldrich.cn/deepweb/assets/sigmaaldrich/product/documents/132/021/protprb

ul.pdf). One aliquot of preparation was stored in a buffer designed for two-dimensional gel

electrophoresis (Solution 2 from Atto Corp.). Following SDS/PAGE separation, the gel was

stained in silver staining, and a gel piece in the range of ~43 kDa was excised for mass

spectroscopy analysis.

Subcellular Fractionation using Available Fractionation Kit

C2C12 or HEK293T cells were harvested using Accutase, washed with DMEM-1% polyvinyl

alcohol (PVA), and then washed with chilled PBS. Cells were fractionated according to the

manufacturer's protocol, with modifications to minimize the contamination for the next step

(Thermo Fisher Scientific, https://assets.thermofisher.com/TFS-

Assets/LSG/manuals/MAN0011667_Subcellular_Protein_Fraction_CulturedCells_UG.pdf).

The samples were centrifuged, and the resulting supernatant was combined with the original

solution. The nuclear extraction buffer without $CaCl_2$ and nuclease was used for washing for

chromatin-bound nuclear fractionation. Each fraction was concentrated by TCA precipitation, as

described below. The resulting pellets were then dissolved in LS buffer and added to

concentrated SDS/PAGE sample buffer at a final concentration of 1 x boiled for 3 min. Finally,

equal relative volumes of each fraction were subjected to Western blotting analysis. To evaluate

the resulting five fractions, Rasa1 and Gapdh were used as marker proteins for the cytoplasmic

fraction, Vdac1 for the membrane fraction, NonO for the soluble nuclear fraction, and Histone

H3 for the chromatin-bound fraction. For quantitative mass spectrometry analysis, C2C12 cells

were seeded at 1.7×10^6 per 100 mm dish x 4 and cultured for 23 h. Duplicate samples were

prepared independently in subsequent steps. The cells were rinsed twice with 10 mL of warm

HBS and treated with 2 mL of Accutase per 100 mm dish. The incubation was carried out at

room temperature for 10 min with gentle rocking on a seesaw shaker while rotating the cell dish

at 90 degrees per 2 min. The suspended cells were gently homogenized with a 1000 μL tip to

prepare a single-cell suspension, which was then collected in a 15 mL tube. The cell

suspensions were combined by washing the dishes twice with 2 mL of chilled DMEM-1% PVA

and centrifuged for 1 min at 4°C. The percentages of trypan-blue-positive cells in the two

groups were 1.8 and 2.2%, respectively. The cells were subjected to a subcellular fractionation

at 10^6 cells per 100 μL of cytoplasmic solution from the kit. Each subcellular fractionated

sample was concentrated with TCA, as described above.

Preparation of Digitonin Solution

659 A stock solution of digitonin was prepared by mixing 25 mg per mL of H_2O at ~80°C. The

660 solution was then aliquoted and stored at –30°C. Before usage, the thawed aliquot was briefly

661 sonicated, heated to ~60°C, cooled to room temperature, and diluted as needed.

Digitonin-Mediated Release in Isotonic Solution from C2C12 Cells

C2C12 cells (4 x 10^5 cells) were seeded in a 6-well dish and cultured in 2 mL DMEM–10%

FCS for 26 h. After two rinses with 2 mL of chilled HBS, the plates were transferred to a cold

room, and sample preparation was performed below 4°C. The residual buffer was wholly

removed with filter paper, and the cells were immersed in 0.5 mL of isotonic solution (HEPES–

KOH, pH 7.5, 150 mM KCl, 2 mM $MgCl_2$, 1 mM EGTA, 1 mM β-mercaptoethanol). The plate

was gently rocking on a seesaw shaker for 10 min. Each fluid was collected into a 1.5 mL tube

and centrifuged at $10,000 \times g$ for 10 min at 4 °C. Concurrently, using a silicone policeman, each

corresponding remaining cell in a 6-well dish was lysed in 100 μL of LS buffer, which included

a cocktail of inhibitors against proteases and phosphatases. Extracts were collected into 1.5 mL

tubes using a 200 μL bore-type tip. After vortexing and boiling for 5 min, the samples were

sonicated for 2.5 min. Each sample was concentrated with TCA and dissolved in LS buffer

using a mixer for over 1 h (Tomy Digital Biology, Tokyo). Final samples were prepared by

mixing 2× SDS/PAGE sample buffer to adjust to 1 x.

Subcellular Fractionation with Digitonin in a Three-Step Process

C2C12 cells (3 x 10^5) or HEK293T cells (4 x 10^5 cells in the polylysine-coated dish) were

cultured in a 60 mm dish for 42–44 h. Washing was performed in the same way as described

earlier. The cells were immersed in 1 mL of digitonin-containing isotonic solution or HFS

solution (6.8% Ficoll, 0.27 M sucrose, 3 mM $CaCl_2$, 2 mM $MgCl_2$ in 20 mM HEPES–KOH,

pH7.5) and gently rocked for 10 min while rotating the cell dish at 90 degrees per min on a

seesaw shaker to ensure even distribution of the solution. The dishes were placed on ice, and the

supernatant fluid was collected in a 1.5 mL tube. The tube was centrifuged at $1000 \times \textbf{\textit{g}}$, 10 min,

4°C, and the resulting supernatant was designated Cyt#1, while the pellets were examined as

debris. The remaining cell layer was gently scraped off into a 0.5 mL isotonic or HFS solution

without digitonin using a cell scraper, and the extract was collected into a 1.5 mL tube using a

bore-type 1000 μL tip. The sample was mixed by vortexing for 5 s at dial 8, left to settle for 15

min on ice, and then centrifuged at $1000 \times g$ for 10 min at 4°C; the supernatant was saved as

Cyt#2. The pellet was then homogenized in 500 μL isotonic solution. The homogenates were

mixed by vortexing and gentle pipetting with a bore-type 1000 μL tip and left for 5 min on ice

before centrifugation at $3000 \times \textbf{\textit{g}}$ for 10 min at 4°C. The supernatant was saved as Cyt#3. The

pellets were resuspended in 150 μL of RIPA buffer (20 mM Tris-HCl, pH 7.5, 150 mM NaCl,

0.5% sodium deoxycholate, 1% Nonidet-P40, 1 mM EDTA) containing proteinase and

phosphatase inhibitors. In some experiments, the pellets were treated with 0.2% Nonidet-P40

and analyzed as an NP40 fraction before being dissolved in RIPA buffer. Subsequently, they

were sonicated in an ice-water bath for 2.5 min (10 s intervals, 15 times) and centrifuged at

$25,000 \times g$ for 30 min at 4°C. The supernatant was designated RIPA-Sup, while the pellets were

dissolved in 50 μL LS. They were vortexed, immediately boiled for 3 min, and analyzed as the

RIPA–Ppt fraction.

Treatment with esiRNA

C2C12 cells were seeded in a 24-well plate at 2×10^4 cells per well. After 20 h culture, the

medium was replaced with 300 μL of either regular medium (A group) or FCS-free DMEM (B

group). Four hours later, transfection was performed with 0, 50, or 100 nM of esiRNA targeting

mouse *Cfdp1* (Sigma-Aldrich, dissolved in 1 mM sodium citrate, pH 6.4) using Lipofectamine

3000. After 4 h, 300 μL of regular medium (to A group) or DMEM containing 20% FCS (to B

group) was added, and the cells were cultured for an additional 44 h. The cells were rinsed twice

with chilled HBS and scraped off using a silicone policeman in 50 μL of LS buffer

supplemented with inhibitors against proteases and phosphatases. The cell lysates were

harvested in a 1.5 mL tube and then boiled immediately. After sonication (10-s intervals x 10),

the samples were centrifuged at $25,000 \times g$ for 30 min at 22°C. Constant amounts of protein (20

μg) were subjected to Western blot analysis. For examination of siRNA effects on subcellular

fractionation, C2C12 cells were seeded at a density of 5×10^4 or 1×10^5/35 mm dish and

cultured for 19 h, treated with esiRNA (0, 50, or 100 nM) using Lipofectamine for 44 h. The

digitonin-mediated fractionation was carried out in an HFS solution similar to the above three-

step process with the following modification. After obtaining the supernatant of Cyt#1 using 0.4

mL of an HFS solution containing 250 μg/ml digitonin, the residual cell layers were scraped off

in 0.2 mL of an HFS solution and centrifuged without vortexing. The pellets were suspended in
0.2 mL HFS solution with brief vortexing and standby for 15 min on ice, then centrifuged, and
the supernatant was combined as Cyt#2*. After washing the second pellets with HBS, the
pellets were resolved in LS buffer, and the fraction was designated as Particulate. Cyt#1 and
#2* were precipitated by TCA treatment, and the precipitates were resolved in 80 μL LS buffer.
Equal amounts of proteins in each fraction of Cyt#1 (15 μg), Cyt#2*(7 μg), and Particulate (25
μg) were subjected to Western blot analysis.

Examination of Potential LLPS of BCNT/CFDP1 in the Nuclei
Plasmid construction and optoDroplet assays were performed in the same way as previously
reported [50]. The IDR of BCNT/CFDP1 was estimated using the PONDR VSL2 algorithm
(http://www.pondr.com/). The IDR fragment of BCNT/CFDP1 (residues 1–273) was amplified
by PCR from a human cDNA library (generated from total RNA extracted from HEK293T
cells. Primers were described in Supplementary Table S10). An optoDroplet assay vector was
constructed from the Cry2 vector and mCherry using the NEBuilder HiFi DNA Assembly Kit to
incorporate the BCNT/CFDP1 fragment. HEK293T cells were transfected with the plasmids
using TurboFect, and potential LLPS activity was measured after 48 h by image monitoring
(Nikon Ti2 inverted microscope with the A1R confocal laser scanning system). After acquiring
background mCherry images, cells were imaged using blue light stimulation (488 nm
stimulation/3 s, interval/17 s, mCherry imaging/10 s x 3), followed by mCherry imaging
(imaging/10 s and interval/50 s x 10).

Isolation of Transiently Expressed FLAG-tagged Bcnt/Cfdp1
C2C12 cells (5 x 10^5 cells/100 mm dish) were cultured for 21 h, transfected with the F-
Bcnt/Cfdp1 plasmid (10 μg), and cultured for an additional 42 h until they were nearly
confluent. The cells were rinsed twice with chilled HBS on ice and scraped off using a cell
scraper in 0.5 ml RIPA buffer containing inhibitors against proteinases and phosphatases. The
lysates were collected in a 1.5 mL tube, sonicated for 2.5 min (15 × 10-s pulses at 10-s
intervals), and centrifuged at 25,000 × g for 30 min at 4°C. The resulting supernatant was snap-
frozen in liquid nitrogen and stored at −80°C until use. The frozen extract was thawed,
sonicated in an ice-water bath (3 × 10-s pulses at 10-s intervals), and centrifuged at 10,000 × g
for 1 min at 4°C. The isolation of F-Bcnt/Cfdp1 was essentially the same as described
previously [33] except for using RIPA buffer to equilibrate the agarose before mixing the cell
extract.

TCA Precipitation

TCA precipitation was carried out according to a protocol with some modifications

(https://www.thermofisher.com/document-connect/document-

connect.html?url=https://assets.thermofisher.com/TFS-

Assets%2FLSG%2Fmanuals%2FMAN0011667_Subcellular_Protein_Fraction_CulturedCells_

UG.pdf). The pellets were washed once with acetone and twice with a 4:1 acetone/water

mixture. The samples were sonicated in an ice-water bath (10-s x 2) and centrifuged during each

washing step. Finally, the sample tubes were heated in a bath at ~95°C for 2 min to evaporate.

Immunoblotting and Densitometric Analysis

Immunoblotting was performed using SDS/PAGE on a 15% gel, followed by electroblotting

onto a polyvinylidene fluoride (PVDF) membrane, as previously described [33, 37]. The Abs

used in the experiments are listed in Supplementary Table S9. Immunoreactivity was detected

using horseradish peroxidase (HRP)-conjugated secondary Abs via a chemical fluorescent

method as previously described [33, 37]. For the subcellular fractionation assay, after

SDS/PAGE and transferred to a PVDF membrane, the membrane was briefly rinsed with water,

followed by Tris-buffered saline containing 0.1% Tween-20 (TBT), then wrapped in plastic

wrap and cut into three parts, higher than 75 kDa (upper), 75–25 kDa (middle), and below 25

kDa, using a cutter based on the position of the prestained SDS/PAGE standard molecular

weight markers. It was sometimes cut into four parts: higher than 75 kDa, 75–40 kDa, 40–20

kDa, and lower than 20 kDa. Each filter was blocked with 0.1 % skim milk in TBT for over 2 h

at room temperature and then incubated at 4°C overnight with Abs against Rasa 1 (upper),

CFDP1 (middle), and HstH3 (lower). The upper and middle membrane parts were subjected to

stripping, followed by blocking, and processed to access by Abs against nestin, NonO for a 75–

40 Da range membrane, and Vdac1 for a 40–25 or 40–20 kDa range membrane. Finally, the

middle or 40–20 kDa range membrane was stripped and then incubated with an HRP-

conjugated anti-Gapdh Ab after blocking. Multi-box strays were routinely used for all

incubation procedures of blocking, reaction with Abs, and washing (Ina Optika MWB-06 or

MWB-4). All images were captured at 5.51M Pixel via a GeneGNone-5 (Syngene Bio Imaging)

using GeneSnaps software and saved as tiff files. The densitometric evaluation of Western blot

signals was conducted using the Band/Peak Quantification plugin in Fiji software [69].

Mass Spectroscopy Analysis

The Coomassie brilliant blue (CBB)-stained bands or Western blot-positive bands in the ~43-

kDa range in the SDS/PAGE gel were excised and destained, and the samples were processed in

duplicate as previously described [10, 33, 37]. They were subjected to trypsin digestion and

subsequently analyzed using nano LC-MS/MS with a Q Exactive mass spectrometer. Each

sample was run in duplicate or triplicate for mass spectrometry analysis. In the study of the

subcellular fractionated samples, SDS/PAGE was performed, and electrophoresis was stopped

upon the entry of the samples into the separation gel to treat each as a single band. Each band

was excised and subjected to reductive acrylamidation. The samples were digested with trypsin,

and equal amounts of each were analyzed in triplicate using the Q Exactive HF-X. The data

were analyzed using Proteome Discoverer 3.0 and MASCOT server 2.8 for protein

identification and quantification (Supplementary Table S2). For examination of Cyt#1/#2*

ratios, each fraction precipitated with TCA was also subjected to brief SDS/PAGE as described

above. Each protein concentration was determined based on amino acid composition and

analyzed in triplicate. Data were acquired using the Proteome Discoverer 2.4 and MASCOT

server 2.7 (Supplementary Table S5). The data of Tables S2 and S5 are available via

ProteomeXchange with the identifier PXD063720

http://proteomecentral.proteomexchange.org/. The distribution of Cyt#1 to Cyt#2* ratios was

presented as violin plots, and their statistical analysis was performed using EZR [70]

(https://cran.r-project.org/web/packages/RcmdrPlugin.EZR/RcmdrPlugin.EZR.pdf).

Figure Legends

Figure 1

Detection of Bcnt/Cfdp1 primarily in the nucleus by a cytochemical approach

A cytochemical approach was employed to detect the localization of Bcnt/Cfdp1, primarily in

the nucleus.

A. Use of tag-free mouse Bcnt/Cfdp1 in HEK293T cells

HEK293T cells were cultured in dishes coated with polylysine, transfected with plasmids

encoding tag-free mouse Bcnt/Cfdp1 or unrelated proteins, and immunostained with an anti-

Bcnt-N Ab (a, b) or anti-Bcnt-Cter Ab (c, d). The image was visualized using Alexa488-

conjugated secondary Ab. The two Abs were generated against two peptides, respectively. The

N-terminal amino acid sequence of Bcnt/Cfdp1 was aligned with its human counterpart, with

red letters indicating different amino acids. The original images are published by Nakashima et

al. [36].

B. Use of fluorescent protein-tagged mouse Bcnt/Cfdp1 in C2C12 myoblasts

B-1. C2C12 cells were cultured in dishes coated with polylysine and transfected with plasmids

encoding EGFP-MCS (a, b) or EGFP-Bcnt/Cfdp1 (c, d). Following a 24 h incubation, the cells

were fixed with 4% PFA on ice. The samples were immunostained with anti-FLAG Ab

followed by CF488A conjugated secondary Ab. The nuclei were stained with DAPI (b, d). The

merged images of EGFP and DAPI on phase contrast pictures are shown (b, d). The scale bars

indicate 25 μm.

B-2. C2C12 myoblasts transfected with Flag-mCherry-Bcnt/Cfdp1, similarly cultured and

stained as Figure 1B-1, were observed at higher resolution using Leica Mica microscopy.

Bcnt/Cfdp1 was stained with anti-Flag Ab followed by Cy3-conjugated secondary Ab (b); the

nuclei were stained with DAPI (c), and the merged image of mCherry and DAPI is shown (a).

The scale bar indicates 10 μm.

C. In differentiated C2C12 myotubes

After 24 h of culturing the EGFP-MCS (a, b), EGFP-Bcnt/Cfdp1 (e, f, i, j), mCherry-MCS (c, d)

or mCherry-Bcnt/Cfdp1 (g, h, k, l) transfectants, the medium was replaced with differentiation

medium (DMEM containing 2% horse serum (DM)). The cells were cultured for an additional 4

days with daily DM changes, after which they were fixed with 4% PFA. The samples were

immunostained with an anti-FLAG Ab followed by CF488A- or Cy3-conjugated secondary Ab,

and the nuclei were stained with DAPI (b, d, f, h, j, l). The merged images of EGFP, mCherry,

and DAPI on phase contrast pictures are shown (b, d, f, h, j, l). Scale bars indicate 50 μm (a–h)

and 25 μm (i–l).

Figure 2

Validity of Western blotting monitoring in C2C12 cell subcellular fractionation

A. Western blotting signals of Bcnt/Cfdp1 in C2C12 cells

Extracts from whole C2C12 cells (equivalent to 7.5×10^4) or mouse embryo brains (15 μg) were

subjected to Western blotting analysis using A305-624A-M as an anti-Bcnt/Cfdp1 Ab.

B. Effects of esiRNA knockdown of Bcnt/Cfdp1 in C2C12 cells.

C2C12 cells were treated with 0, 50, or 100 nM esiRNA for 24 h. Each extract was prepared by

scraping off the cells in LS buffer and boiling. After centrifugation, equal amounts of protein

(20 μg) were subjected to Western blotting analysis. The upper and middle sections of the image

show Western blotting signals using Abs against Rasa1 and Bcnt/Cfdp1, respectively. The red

and blue arrows correspond to the target and upper band. At the bottom, densitometric analysis

of blots of Bcnt/Cfdp1 (red bar), "upper band(s)" (dark blue), and Rasa1 (light blue) are shown

as bar graphics with the mean ± SEM of duplicate experiments.

C. Effects of esiRNA on the cytosolic and particulate proteins

C2C12 cells were cultured in 35 mm dishes and treated with esiRNA (0 or 50 nM). Cytosolic

and particulate proteins were obtained using an HFS solution containing 250 μg/ml digitonin, as

detailed later in Figures 6-8. Briefly, digitonin-mediated released fluid was the cytosolic

fraction, and the particulates were obtained by scrapping off the remaining cell layer in HFS

solution followed by centrifugation. The cytosolic fraction was concentrated with TCA

precipitation followed by resuspending in LS buffer, while the particulate fraction was directly

resolved in LS buffer. Equal relative amounts of each fraction (approximately 15 or 25 μg of

protein, respectively) were subjected to Western blotting analysis to assess the degree of

separation using anti-histone H3 Ab in addition to that in Fig. 3B. The red and blue arrows

correspond to the target and "upper band," respectively. Asterisks ($*$) indicate off-target bands.

D.   The presence of the "upper band" in the FLAG-tagged Bcnt/Cfdp1

C2C12 cells were transfected with F-MCS or F-Bcnt/Cfdp1 (Tf: transfection). After 24 h of

culture, one sample from each group was replaced with serum-free medium (St: serum-starved)

and cultured for 6 h. All cell cultures were harvested and subjected to Western blotting analysis

using anti-Bcnt/Cfdp1 (upper panel) or anti-FLAG (lower panel) Abs. The blue arrows indicate

the "upper band" of F-Bcnt/Cfdp1.

E.   The appearance of a cross-reacted 50 kDa band during C2C12 differentiation

C2C12 cell cultures were switched from regular medium to DMEM containing 2% horse serum,

designated as day 0. Cell extracts were collected and subjected to Western blotting analysis the

following day. The red and blue arrows indicate Bcnt/Cfdp1 and an unrelated 50 kDa band. At

the bottom, densitometric analysis of blots of Bcnt/Cfdp1 (red bar) and 50 kDa band (dark blue)

are shown as bar graphics with the mean ± SEM of duplicate experiments.

Figure 3

The presence of Bcnt/Cfdp1 in the cytoplasmic fraction determined using a widely used kit

For subcellular fractionation of C2C12 or HEK293T cells, each cell suspension (1×10^6 cells)

was subjected to fractionation according to the manufacturer's protocol, which was slightly

modified with additional washes to minimize contamination at each step. Each fraction was

concentrated by TCA precipitation and resolved in LS buffer. To show the relative occupancy

of all fractions except WCE, equal volumes of each were subjected to Western blotting analysis.

The following proteins were accessed: Rasa1 (Ras GTPase activating protein 1), Gapdh

(glyceraldehyde-3-phosphate dehydrogenase), Vdac1 (voltage-gated anion channel 1), non-POU

domain-containing octamer-binding protein (NonO), HstH3 (histone H3), and Bcnt/Cfdp1 using

A305-624A-M. The asterisks in the signals detected with the anti-Bcnt/Cfdp1 Ab indicate the

off-target band. Abbreviations at the top of the figure. WCE: whole cell extract, Cyt:

cytoplasmic extract, Mem: membrane extract, s-N: soluble nuclear extract, c-N: chromatin-

bound nuclear extract, and Ppt: pellet extract.

Figure 4

Quantitative analysis of the subcellular fractionated samples accessed by Western blotting

or by mass spectrometry

A. Distribution of six marker proteins in each fraction accessed by Western blotting

Densitometric analysis of Western blots of six biomarker proteins (Bcnt/Cfdp1, Rasa 1,

Gapdh, Vdac1, NonO, and HstH3) in C2C12 subcellular fractions of Figure 3 is shown. Data

represent the relative occupancy of each marker protein in the five fractions, in which total

amounts are set as 1.0, and the bars represent the mean ± SEM of results obtained from five

independent experiments. Cyt, cytoplasmic fraction; Mem, membrane fraction; s-N, soluble

nuclear fraction; c-N, chromatin-bound fraction; Ppt, precipitates.

B–D

Two independent C2C12 myoblast cultures were fractionated in a manner analogous to that

described in Figure 4. Each fraction was subjected to quantitative mass spectrometry analysis. A

total of 4,489 proteins were identified, and the average abundance of each component was

calculated and displayed graphically. Cyt, cytoplasmic fraction; Mem, membrane fraction; s-N,

soluble nuclear fraction; c-N, chromatin-bound fraction; CSK, cytoskeleton.

B. Distribution of six marker proteins in each fraction accessed by mass analysis

Data represent the relative amounts of each marker protein per cell extract in the five fractions,

in which total amounts of Bcnt/Cfdp1 are set at 1.0, and bars represent the mean ± SEM of

results obtained from two independent experiments.

C. The intracellular distributions of three paraspeckle components and 18 heterogeneous

nuclear ribonucleoproteins

The data are shown in relative quantitative amounts, with the total Bcnt/Cfdp1 set at 1.0. The

data represent the means of two experiments.

D. The intracellular distribution of six proteins

The intracellular distribution of proteins screened as those detected in the cytoplasmic fraction
despite being annotated localized in the nucleus is shown. The relative distribution of each
protein is shown from the mean value of the two experiments.

Figure 5
Release of Bcnt/Cfdp1 by digitonin treatment in an isotonic solution

A. Cell cultures in 12-well dishes were exposed to a range of digitonin concentrations in an
isotonic solution for 10 min on a rocking shaker. After the supernatant of each eluate was
collected, the corresponding extracts of the residual cellular debris were scraped in an SDS-
containing buffer and boiled. Equal volumes of each sample were subjected to Western
blotting using Abs against Rasa1 (upper box), Bcnt/Cfdp1 (middle box), and Gapdh (lower
box). After Western blotting analysis, the membrane was stained with CBB to show the 37–
75 kDa region (at the bottom).
B. The eluate supernatant was subjected to SDS/PAGE, and after staining the gel with CBB, the
~43 kDa region was excised for mass spectroscopic analysis.
C. After destaining the gel pieces, they were subjecting tryptic digests to liquid
chromatography-tandem mass spectrometry analysis. Of the 124 proteins identified
(Supplementary Table S4), data are shown for Bcnt/Cfdp1, with 41% recovered by the
analyzed fragments.
D. Using the three-step preparation of the digitonin-mediated eluate, as illustrated in Figure 7,
Cyt #1–3 fractions were obtained using digitonin at 125 µg/mL in an isotonic solution.
Following TCA precipitation, each pellet was dissolved in LS buffer (containing 1% SDS),
and relatively equal volumes of each fraction were subjected to Western blotting analysis, as
described in the legend of Figure 3. The dark blue arrow and asterisks correspond to the
"upper band" of Bcnt/Cfdp1 and probable off-target band(s). Densitometric analysis of the
Western blotting is shown at the bottom. The data represent the relative occupancy of each
of the six marker proteins in each fraction (Cyt#1–RIPA–Ppt), with the total amounts of
each marker protein set at 1.0.

Figure 6
Schematic diagram of the three-step preparation of the digitonin-mediated eluates

Cultured cells were immersed in an isotonic or HEPES/Ficoll/sucrose (HFS) solution containing

digitonin while the dish was rotated. The eluate supernatant was referred to as Cyt#1 after

centrifugation. The remaining cell debris was scraped off using a cell scraper in a digitonin-free

isotonic or HFS solution, and the supernatant was stored as Cyt#2 after centrifugation. The

pellet was washed by suspension in an isotonic solution, and the supernatant was stored as

Cyt#3 following centrifugation. The pellet was resuspended in RIPA buffer, sonicated, and

centrifuged. The supernatant was used as the RIPA fraction, the pellet was dissolved in SDS-

containing buffer, and the supernatant was analyzed as the Ppt fraction. In the case of isotonic

solution, the pellets were solved in 0.2% Nonidet-40. The details of the procedure are described

in the Materials and Methods section.

Figure 7

Differential release of Bcnt/Cfdp1 from cytosolic proteins by digitonin treatment in HFS

The Cyt#1–3 fractions were obtained using digitonin in HFS solution at 50 or 250 μg/mL in

C2C12 cells or 50 μg/mL in HEK293T cells, as illustrated in Figure 6. Following TCA

precipitation, each pellet was dissolved in LS buffer, and relatively equal volumes of each

fraction were subjected to Western blotting analysis, as described in the legend in Figure 3.

Densitometric analysis of the Western blotting is shown at the bottom. The data represent the

relative occupancy of each of the six marker proteins in each fraction (Cyt#1–RIPA–Ppt), with

the total amounts of each marker protein set at 1.0.

Figure 8

Mobility or integrity of digitonin-mediated released proteins in the Cytoplasm -Evaluation

from the ratios of two fractions-

A. C2C12 cells treated with esiRNA (0, 50, or 100 nM) were subjected to digitonin-mediated

fractionation in HFS solution at 250 μg/mL digitonin, as shown in Figure 6. But, after

obtaining Cyt#2, its pellet was washed with HFS solution but not with isotonic solution, and

the washed supernatant was combined to give Cyt#2*. Furthermore, the precipitate was

dissolved in LS buffer after washing with HBS as a Particulate fraction. Equal amounts of

protein from Cyt#1 (15 μg), Cyt#2* (7 mg), and Particulate (25 mg) fractions were

subjected to Western blot analysis. These different protein amounts were adjusted so that

the relative occupancy volumes of the three fractions were approximately equal. Each Cyt#1

and #2* fraction was subjected to quantitative mass spectrometry analysis (Supplementary

Table S5).

B. Fifty-one glycolysis/glucogenesis/TCA cycle/organelle-associated enzymes were listed

from the 2442 qualified proteins by quantitative mass analysis (Supplementary S6). Each

Cyt#1/2* ratio value is plotted on the vertical axis as LOG2, while the horizontal axis shows

its molecular weight (Mw). Glycolysis-related cytosolic proteins (magenta circles),

organelle-related proteins (Diverse, orange circles), and TCA-related mitochondrial proteins

(dark blue circles) are shown.

C. The distribution of Cyt#1/2* ratios of 51-enzymes shown in Figure 8B (magenta, orange,

dark blue), 2442 qualified total proteins (purple), and 321 RNA-binding proteins (green),

which are selected based on published data were presented (Supplementary Table S7) as

violin plots, respectively, using EZR software. The rectangles within the violin plots

indicate the first and third quartiles, with central + as mean values. Results of statistical

analysis are shown. ** indicates a significant difference with $P<0.01$

Acknowledgments

We thank Dr. Tsutomu Kumazaki for the puromycin-resistant AB2.2 cell line. We appreciate

Drs. Shinya Watanabe and Dimitris Beis for information on the properties of Bcnt/cfdp1. We

acknowledge Drs. Yan Fei Gao, Gereon Poschmann, Lauri A. Aaltonen, and Justin Cotney for

sharing unpublished data. We also thank the following scientists for their valuable insights

during this research: Drs. Satoshi Yoshitome, Yasuhiro Kashino, Atsushi Iwai, Masahiro

Nishibori, Naoyuki Kataoka, Richard Schulz, Heiko Heerklotz, Song-Tao Liu, Takuto Shimizu,

and Mitsuru Okuwaki. We were encouraged by Dr. Stephane Twigg for his comments regarding

the nomenclature of the CFDP1 gene. We are willing to thank Editage (www. editing. com) for

precision editing the manuscript. We were indebted to Professors Si-Young Song, Hiromi

Nochi, and Takashi Tominaga for their support and encouragement.

References

1. Antifeeva, I.A., Fonin, A.V., Fefilova, A.S., Stepanenko, O.V., Povarova, O.I., Silonov, S.A., et al. (2022) Liquid-liquid phase separation as an organizing principle of intracellular space: overview of the evolution of the cell compartmentalization concept. Cell Mol Life Sci. 79, 251. 10.1007/s00018-022-04276-4

2. Jalihal, A.P., Schmidt, A., Gao, G., Saffron, R., Little, S.R., Pitchiaya, S., et al. (2021) Hyperosmotic phase separation: Condensates beyond inclusions, granules and organelles. J Biol Chem. 296, 100044. 10.1074/jbc.REV120.010899

3. Watson, J.L., Seinkmane, E., Styles, C.T., Mihut, A., Krüger, L.K., McNally, K.E., et al. (2023) Macromolecular condensation buffers intracellular water potential. Nature 623, 842-852. 10.1038/s41586-023-06626-z

4. Zhao, Y.G., Zhang, H. (2020) Phase separation in membrane biology: The interplay between membrane-bound organelles and membraneless condensates. Dev Cell 55, 30-44. 10.1016/j.devcel.2020.06.033

5. Kim, T., Jaeyoon Yoo, J., Do, S., Hwang, D.S., YongKeun Park, Y.K. (2023) RNA-mediated demixing transition of low-density condensates. Nat Commun. 14, 2425. 10.1038/s41467-023-38118-z

6. Hertzog, M., Erdel, F. (2023) The material properties of the cell nucleus: A matter of scale. Cells 12, 1958. 10.3390/cells12151958

7. Hampoelz, B., Andres-Pons, A., Kastritis, P., Beck, M. (2019) Structure and assembly of the nuclear pore complex. Annu Rev Biophys. 48, 515-536. 10.1016/j.devcel.2020.06.033

8. Timney, B.L., Raveh, B., Mironska, R., Trivedi, J.M., Kim, S.J, Russel, D., et al. (2016) Simple rules for passive diffusion through the nuclear pore complex. J Cell Biol. 215, 57-76. 10.1016/j.devcel.2020.06.033

9. Junod, S.L., Kelich, J.M., Ma, J., Yang, W. (2020) Nucleocytoplasmic transport of intrinsically disordered proteins studied by high-speed super-resolution microscopy. Protein Sci. 29, 1459-1472. 10.1073/pnas.1914313117

10. Iwashita, S., Suzuki, T., Yasuda, T., Nakashima, K., Sakamoto, T., Kohno, T., et al. (2015) Mammalian Bcnt/Cfdp1, a potential epigenetic factor characterized by an acidic stretch

in the disordered N-terminal and Ser250 phosphorylation in the conserved C-terminal regions. Biosci Rep. 35, e00228. 10.1042/BSR20150111

11. Messina, G., Celauro, E., Atterrato, M.T, Giordano, E., Iwashita, S., Dimitri, P. (2015) The Bucentaur (BCNT) protein family: a long-neglected class of essential proteins required for chromatin/chromosome organization and function. Chromosoma 124, 153-162. 10.1007/s00412-014-0503-8

12. Yu, J., Sui, F., Gu, F., Li, W., Yu, Z., Wang, Q., et al. (2024) Structural insights into histone exchange by human SRCAP complex. Cell Discov. 10, 15. 10.1038/s41421-023-00640-1

13. Cai, Y., Jin, J., Florens, L., Swanson, S.K., Kusch, T., Li, B., et al. (2005) The mammalian YL1 protein is a shared subunit of the TRRAP/TIP60 histone acetyltransferase and SRCAP complexes. J Biol Chem. 280, 13665-13670. 10.1074/jbc.M500001200

14. Huang, Y., Sun, L., Pierrakeas, L., Dai, L., Pana, L., Luk, E., et al. (2020) Role of a DEF/Y motif in histone H2A-H2B recognition and nucleosome editing. Proc Natl Acad Sci U S A. 117, 3543-3550. 10.1073/pnas.1914313117

15. Jalal, A.S.B., Girvan, P., Chua, E. Y. D., Liu L., Wang, S., McCormack, E. A., et al. (2024) Stabilization of the hexasome intermediate during histone exchange by yeast SWR1 complex. Mol Cell 84, 3871-3884.e9. 10.1016/j.molcel.2024.08.015

16. Sun, L., Luk, E. (2017) Dual function of Swc5 in SWR remodeling ATPase activation and histone H2A eviction. Nucleic Acids Res. 45, 9931-9946. 10.1093/nar/gkx589

17. Ishida, H., Kono, H. (2022) Free energy landscape of H2A-H2B displacement from nucleosome. J Mol Biol. 434, 167707. 10.1016/j.jmb.2022.167707

18. Saccharomyces Genome Database. https://www.yeastgenome.org/

19. Morillo-Huesca, M., Clemente-Ruiz, M., Andújar, E., Prado, F. (2010) The SWR1 histone replacement complex causes genetic instability and genome-wide transcription misregulation in the absence of H2A.Z. PLoS One 5, e12143. 10.1371/journal.pone.0012143

20. Messina, G., Damia, E., Fanti, L., Atterrato, M.T., Celauro, E., Mariotti, F.R, et al. (2014) *Yeti*, an essential *Drosophila melanogaster* gene, encodes a protein required for chromatin organization. J Cell Sci. 127, 2577-2588. 10.1242/jcs.150243

21. Itoh, T., Inoue, S., Sun, X., Kusuda, R., Hibi, M., Shimizu, T. (2021) Cfdp1 controls the cell cycle and neural differentiation in the zebrafish cerebellum and retina. Dev Dyn. 250, 1618-1633. 10.1002/dvdy.371

22. Giardoglou, P., Deloukas, P., Dedoussis, G., Beis, D. (2023) *Cfdp1* Is Essential for Cardiac Development and Function. Cells 12, 1994. 10.3390/cells12151994

23. Ito, Y. (2012) CP27: Expression, Regulation, and Function of a Chromatin Complex Member (Doctoral dissertation, University of Illinois at Chicago)

24. Doolan, P., Barron, N., Kinsella, P., Clarke, C., Meleady, P., O'Sullivan, F., et al. (2012) Microarray expression profiling identifies genes regulating sustained cell specific productivity (S-Qp) in CHO K1 production cell lines. Biotechnol J. 7, 516-526. 10.1002/biot.201100255

25. Xu, L., He, Y., Ding, Y., Liu, G.E., Zhang, H., Cheng, H.H., et al. (2018) Genetic assessment of inbred chicken lines indicates genomic signatures of resistance to Marek's disease. J Anim Sci Biotechnol. 9, 65. 10.1186/s40104-018-0281-x

26. Messina, G., Atterrato, M.T., Prozzillo, Y., Piacentini, L., Losada, A., Dimitri, P. (2017) The human Cranio Facial Development Protein 1 (Cfdp1) gene encodes a protein required for the maintenance of higher-order chromatin organization. Sci Rep. 7, 45022. 10.1038/srep45022

27. Moustakas, A., Heldin, C.H. (2007) Signaling networks guiding epithelial-mesenchymal transitions during embryogenesis and cancer progression. Cancer Sci. 98, 1512-1520. 10.1111/j.1349-7006.2007.00550.x

28. Ming, R., Li, X., Wang, E., Wei, J., Liu, B., Zhou, P., et al. (2022) The prognostic signature of head and neck squamous cell carcinoma constructed by immune-related RNA-binding proteins. Front Oncol. 12, 795781. 10.3389/fonc.2022.795781

29. Zhou, Y., Qiu, J., Liu, S., Wang, P., Ma, D., Zhang, G., et al. (2023) CFDP1 promotes hepatocellular carcinoma progression through activating NEDD4/PTEN/PI3K/AKT signaling pathway. Cancer Med. 12, 425-444. 10.1002/cam4.4919

30. Cuadrado, A., Corrado, N., Perdiguero, E., Lafarga, V., Muñoz-Canoves, P., Nebreda, A.R. (2010) Essential role of p18Hamlet/SRCAP-mediated histone H2A.Z chromatin incorporation in muscle differentiation. EMBO J. 29, 2014-2025. 10.1038/emboj.2010.85

31. Dong, S., Han, J., Chen, J., Liu, T., Huen, M.Y.S., Yang, Y. (2014) The human SRCAP chromatin remodeling complex promotes DNA-end resection. Curr Biol. 24, 2097-2110. 10.1016/j.cub.2014.07.081

32. Välimäki, N., Jokinen, V., Cajuso, T., Kuisma, H., Taira, A., Dagnaud, O., et al. (2023) Inherited mutations affecting the SRCAP complex are central in moderate-penetrance predisposition to uterine leiomyomas. Am J Hum Genet. 110, 460-474. 10.1016/j.ajhg.2023.01.009

33. Iwashita, S., Suzuki, T., Kiriyama, Y., Dohmae, N., Ohoka, Y., Song, S.Y., et al. (2020) Overcoming off-targets: assessing Western blot signals for Bcnt/Cfdp1, a tentative component of the chromatin remodeling complex. Biosci Rep. 40, BSR20194012. 10.1042/BSR20194012

34. Clement-Ziza, M., Marsellach, F., Codlin, S., Papadakis, M.A., Reinhardt, S., Rodríguez-López, M., et al. (2014) Natural genetic variation impacts expression levels of coding, non-coding, and antisense transcripts in fission yeast. Mol Syst Biol. 10, 764. 10.15252/msb.20145123

35. Hagting, A., Jackman, M., Simpson, K., Pines, J. (1999) Translocation of cyclin B1 to the nucleus at prophase requires a phosphorylation-dependent nuclear import signal, Curr Biol **9**,680-9. 10.1016/s0960-9822(99)80308-x.

36. Pinarbasi, E.S., Tolga Cağatay, T., Fung, H.Y.J., Li, Y.C., Chook, Y.M., Thomas, P.J. (2018) Active nuclear import and passive nuclear export are the primary determinants of TDP-43 localization. Sci Rep. 8, 7083. 10.1038/s41598-018-25008-4

37. Nakashima, K., Iwashita, S., Suzuki, T., Kato, C., Kohno, T., Kamei, Y., et al. (2019) A spatial similarity of stereochemical environments formed by amino acid residues defines a common epitope of two non-homologous proteins. Sci Rep. 9, 14818. 10.1038/s41598-019-51350-2

38. Ramsby, M.L., Makowski, G.S. (2005) Differential detergent fractionation of eukaryotic cells. In: Walker, J.M. (ed) The Proteomics Protocols Handbook. Springer Protocols Handbooks. Humana Press. 10.1385/1-59259-890-0:037

39. Afzal, A., Sarfraz, M., Li, G.L., Ji, S., Duan, S.F., Khan, N.H., et al. (2019) Taking a holistic view of PEST-containing nuclear protein (PCNP) in cancer biology. Cancer Med. 8, 6335-6343. 10.1002/cam4.2465

40. Savojardo, C., Martelli, P.L., Fariselli, P., Profiti, G., Casadio, R. (2018) BUSCA: an integrative web server to predict subcellular localization of proteins. Nucleic Acids Res. 46, W459–W466. 10.1093/nar/gky320

41. Orre, L.M., Vesterlund, M., Pan, Y., Arslan, T., Zhu, Y., Woodbridge, A.F., et al. (2019) SubCellBarCode: Proteome-wide mapping of protein localization and relocalization. Mol Cell 73, 166-182.e7. 10.1016/j.molcel.2018.11.035

42. Lorent, J.H., Quetin-Leclercq, J., Mingeot-Leclercq, M.P. (2014) The amphiphilic nature of saponins and their effects on artificial and biological membranes and potential consequences for red blood and cancer cells. Org Biomol Chem. 12, 8803-8822. 10.1039/C4OB01652A

43. Fan, H.Y., Heerklotz, H. (2017) Digitonin does not flip across cholesterol-poor membranes. J Colloid Interface Sci. 504, 283-293. 10.1016/j.jcis.2017.05.034

44. Wang, K., Sturt-Gillespie, B., Hittle, J.C., Macdonald, D., Chan, G.K., Yen, T.J., et al. (2014) Thyroid hormone receptor interacting protein 13 (TRIP13) AAA-ATPase is a novel mitotic checkpoint-silencing protein. J Biol Chem. 289, 23928-23937. 10.1074/jbc.M114.585315

45. Minton, A.P. (2006) How can biochemical reactions within cells differ from those in test tubes? J Cell Sci. 119, 2863-2869. 10.1242/jcs.03063

46. Hancock, R., Hadj-Sahraoui, Y. (2009) Isolation of cell nuclei using inert macromolecules to mimic the crowded cytoplasm. PLoS One 4, e7560. 10.1080/15384101.2016.1211214

47. Sun, L., Fang, J. (2016) Macromolecular crowding effect is critical for maintaining SIRT1's nuclear localization in cancer cells. Cell Cycle 15, 2647-2655. 10.1080/15384101.2016.1211214

48. Yu, J., Sui, F., Gu, F., Li, W., Yu, Z., Wang, Q., et al. (2024) Structural insights into histone exchange by human SRCAP complex. Cell Discov. 10, 15. 10.1038/s41421-023-00640-1

49. Chu, W.T., Wang, J. (2021) Thermodynamic and sequential characteristics of phase separation and droplet formation for an intrinsically disordered region/protein ensemble. PLoS Comput Biol. 17, e1008672. 10.1371/journal.pcbi.1008672

50. Shin, Y., Berry, J., Pannucci, N., Haataja, M.P., Toettcher, J.E., Brangwynne, C.P. (2017) Spatiotemporal control of intracellular phase transitions using light-activated optoDroplets. Cell 168, 159-171. 10.1016/j.cell.2016.11.054

51. Li, H.K., Zhou, Y., Ding, J., Xiong, L., Shi, Y.X., He, Y.J., et al. (2020) LRTM1 promotes the differentiation of myoblast cells by negatively regulating the FGFR1 signaling pathway. Exp Cell Res. 396, 112237. 10.1016/j.yexcr.2020.112237

52. Senichkin, V.V., Prokhorova, E.A., Zhivotovsky, B., Kopeina, G.S. (2021) Simple and efficient protocol for subcellular fractionation of normal and apoptotic cells. Cells 10, 852. 10.3390/cells10040852

53. Houben, F., Ramaekers, F.C.S., Snoeckx, L.H.E.H., Broers, J.L.V. (2007) Role of nuclear lamina-cytoskeleton interactions in the maintenance of cellular strength. Biochim Biophys Acta 1773, 675-686. 10.1016/j.bbamcr.2006.09.018

54. Gunn, A.L., Yashchenko, A.I., Dubrulle, J., Johnson, J., Hatch, E.M. (2024) A high-content screen reveals new regulators of nuclear membrane stability. Sci Rep. 14, 6013. 10.1038/s41598-024-56613-1

55. Fan, J.R., Chang, S.N., Chu, C.T., Chen, H.C. (2023) AKT2-mediated nuclear deformation leads to genome instability during epithelial-mesenchymal transition. iScience 26, 106992. 10.1016/j.isci.2023.106992

56. Herrmann, C., Avgousti, D.C., Weitzman, M.D. (2017) Differential salt fractionation of nuclei to analyze chromatin-associated proteins from cultured mammalian cells. Bio Protoc. 7, e2175. 10.21769/BioProtoc.2175

57. Youn, J.H., Shin, J.S. (2006) Nucleocytoplasmic shuttling of HMGB1 is regulated by phosphorylation that redirects it toward secretion[1]. **J Immunol.** 177, 7889-7897. 10.4049/jimmunol.177.11.7889

58. Cox, J., Jensen, L.J., Meissner, F., Mann, M. (2015) Secretome analysis of lipid-induced insulin resistance in skeletal muscle cells by a combined experimental and bioinformatics workflow. J Proteome Res. 14, 4885-4895. 10.1021/acs.jproteome.5b00720

59. Zhao, L., Poschmann, G., Waldera-Lupa, D., Rafiee, N., Kollmann, M., Stühler, K. (2019) OutCyte: a novel tool for predicting unconventional protein secretion. Sci Rep. 9, 19448. 10.1038/s41598-019-55351-z

60. Chien, H.M., He, R.Y., Lee, C.C., Huang, Y.A., Hung, I.J., Hou, K.T., et al. (2021) Nanoscopic investigation of C9orf72 poly-GA oligomers on nuclear membrane disruption by a photoinducible platform. Commun Chem. 4, 111. 10.1038/s42004-021-00547-6

61. Orczyk, M., Wojciechowski, K., Brezesinski, G. (2017) Disordering effects of digitonin on phospholipid monolayers. Langmuir 33, 3871-3881. 10.1021/acs.langmuir.6b04613

62. Dazzoni, R., Grélard, A., Morvan, E., Bouter, A., Applebee, C.J., Loquet, A. (2020) The unprecedented membrane deformation of the human nuclear envelope. in a magnetic field, indicates formation of nuclear membrane invaginations. *Sci Rep* **10**,5147. 10.1038/s41598-020-61746-0

63. Romanauska, A., Köhler A (2023) Lipid saturation controls nuclear envelope function. *Nat Cell Biol* **25**,1290-1302. 10.1038/s41556-023-01207-8

64. Rispal, J., Escaffit, F., Trouche, D. (2020) Chromatin dynamics in intestinal epithelial homeostasis: A paradigm of cell fate determination versus cell plasticity. Stem Cell Rev Rep. 16, 1062-1080. 10.1007/s12015-020-10055-0

65. Leal, M.P.C., Sciortino, M., Tornillo, G., Colombo, S., Defilippi, P., Cabodi, S. (2015) p130Cas/BCAR1 scaffold protein in tissue homeostasis and pathogenesis. Gene 562, 1-7. 10.1016/j.gene.2015.02.027

66. Nurnberg, S.T., Guerraty, M.A., Wirka, R.C., Rao, H.S., Pjanic, M., Norton, S., et al. (2020) Genomic profiling of human vascular cells identifies TWIST1 as a causal gene for common vascular diseases. PLoS Genet. 16, e1008538. 10.1371/journal.pgen.1008538

67. Riesmeijer, S.A., Kamali, Z., Ng, M., Drichel, D., Piersma, B., Becker, K., et al. (2024) A genome-wide association meta-analysis implicates Hedgehog and Notch signaling in Dupuytren's disease. Nat Commun. 15, 199. 10.1038/s41467-023-44451-0

68. Anand, D., Chaudhuri, A. (2023) Grease in the nucleus: Insights into the dynamic life of nuclear membranes. J Membr Biol. 256, 137-145. 10.1007/s00232-022-00272-8

69. Ohgane, K., Yoshioka, H. (2019) Quantification of gel bands by an Image J macro, band/peak quantification tool. Protocols.io. 10.17504/protocols.io.7vghn3w

70. Kanda, Y. (2013) Investigation of the freely available easy-to-use software 'EZR' for medical statistics. Bone Marrow Transplant 48, 452-458. 10.1038/bmt.2012.244

Footnote references

‡The official name of *Bcnt/Cfdp1* is *Cfdp1*. However, genetic studies of disease (1) or transcriptome analysis (2) do not provide solid evidence for its involvement in craniofacial development, including tooth development (3). Therefore, the nomenclature of *Cfdp1* is misleading and may need to be more accurate. To maintain consistency with previous studies, the name *Bcnt/Cfdp1* is used instead of *Cfdp1* in this article.

1. Twigg, S.R.F., Wilkie, A.O.M. (2015) New insights into craniofacial malformations. Hum Mol Genet. 24, R50-R59. 10.1093/hmg/ddv228

2. Yankee, T.N., Oh, S., Winchester, E.W., Wilderman, A., Robinson, K., Gordon, T., et al. (2023) Integrative analysis of transcriptome dynamics during human craniofacial development identifies candidate disease genes. Nat Commun. 14, 4623. 10.1038/s41467-023-40363-1

3. Hallikas, O., Roy, R.D., Christensen, M.M., Renvoisé, E., Sulic, A.M., Jernvall, J. (2021) System-level analyses of keystone genes required for mammalian tooth development. J Exp Zool B Mol Dev Evol. 336, 7, 10.1002/jez.b.23009

Detection of Bcnt/Cfdp1 primarily in the nucleus by a cytochemical approach

A   **Use of tag-free mouse Bcnt/Cfdp1 in HEK293T cells**

B   **Use of fluorescent protein-tagged mouse Bcnt/Cfdp1 in C2C12 myoblasts**

C **In differentiated C2C12 myotubes**

Figure 2

Validity of Western blotting monitoring in C2C12 cell subcellular fractionation

Presence of Bcnt/Cfdp1 in the cytoplasmic fraction determined using a widely used kit

Quantitative analysis of subcellular fractionated samples

A Marker proteins by Western blotting analysis

B Marker proteins by Mass analysis

C Paraspeckle components and Hnrnps

D *Six "nuclear proteins" detected in the cytoplasmic fraction*

Figure 5

Release of Bcnt/Cfdp1 by digitonin treatment in an isotonic solution

Figure 7

Differential release of Bcnt/Cfdp1 from cytosolic proteins by digitonin treatment in HFS

Mobility or integrity of digitonin-mediated released proteins in the Cytoplasm
-Evaluation from the ratios of two fractions-