Preparation duration shapes the goal-directed tuning of stretch reflex responses

Robin Rohlén[1,2*], Frida Torell[1], Michael Dimitriou[1]

[1] Department of Medical and Translational Biology, Umeå University, Umeå, Sweden

[2] Department of Diagnostics and Intervention, Umeå University, Umeå, Sweden

* Correspondence to Robin Rohlén (robin.rohlen@umu.se).

ORCID: Robin Rohlén (0000-0003-4328-5467), Frida Torell (0000-0002-6294-7844), Michael Dimitriou (0000-0001-9890-2974)

Acknowledgements: We thank Carola Hjältén and Anders Bäckström for their technical support. This work was supported by grants awarded to M.D. by Hjärnfonden (FO2024-0425-HK-88) and R.R. by the Swedish Research Council for Sport Science (P2025-0173).

Statements and Declarations: The authors have no competing interests to declare that are relevant to the content of this study.

Author contributions: All authors contributed to the study's conception and design. RR performed data curation and analysis. RR wrote the first draft of the manuscript. All authors reviewed and edited the manuscript. All authors read and approved the final manuscript.

Abstract

Stretch reflex responses allow for effective resistance against postural perturbations, but reflex gains can also be modulated to accommodate the execution of voluntary movements. Recent findings suggest that movement preparation includes independent and goal-directed tuning of muscle spindles, facilitating a congruent modulation of reflex gains. In the context of delayed-reach, goal-directed tuning of stretch reflex gains is present or enhanced when the homonymous muscle is unloaded (i.e., antagonist load) and the preparatory delay is sufficiently long (>250 ms and <750 ms). This study examined how multiple preparatory delays impact the goal-directed modulation of short- and long-latency stretch reflex (SLR and LLR) gains of the pectoralis, anterior and posterior deltoid muscles in a delayed-reach task. We used electromyographic signals to quantify stretch reflex responses of healthy subjects that experienced haptic perturbations induced by a robotic manipulandum in a delayed-reach task. We found that a preparatory delay of 300 ms is sufficient for goal-directed tuning of SLR responses in the pectoralis and posterior deltoid muscles, and 400 ms allows for even stronger goal-directed tuning of stretch reflex responses. In contrast, there was no goal-directed modulation in the SLR of the anterior deltoid. Tuning of LLR gains was robustly goal-dependent throughout, with negligible modulation across preparatory delays. Our results clarify the minimum preparatory time required for goal-directed tuning of stretch reflex gains and characterise the relationship between reflex gains and preparation duration. This relationship is not strictly linear, likely reflecting the interplay of multiple feedback mechanisms functioning at different time frames.

Keywords: preparatory delay, reaching task, stretch reflex, perturbation, electromyography

Introduction

Voluntary movement normally involves a preparation period where neural activity is adjusted before movement initiation (Kutas and Donchin, 1974; Wise, 1985; Ghez *et al.*, 1997). This preparatory neural activity correlates with movement direction (Tanji and Evarts, 1976), movement speed (Churchland, Santhanam and Shenoy, 2006), reach distance (Messier and Kalaska, 2000), reach trajectory (Hocherman and Wise, 1991), and visual target location (Batista *et al.*, 2007). Moreover, reaching movements involve the formation of feedforward motor commands and a feedback policy that modulates long-latency stretch reflex responses (LLRs) before movement onset (Todorov and Jordan, 2002; Shadmehr and Krakauer, 2008; Wagner and Smith, 2008; Ahmadi-Pajouh, Towhidkhah and Shadmehr, 2012; Yeo, Franklin and Wolpert, 2016).

Unexpected stretching of muscles in the upper extremities can produce a short-latency reflex response (SLR) beginning ~25 ms after stretch onset and an LLR beginning ~50 ms after stimulus onset (Hammond, 1956; Marsden, Merton and Morton, 1972). Both the SLR and the LLR are considered involuntary, with voluntary control of the upper extremities beginning ~100 ms after the onset of the triggering sensory event (Yang *et al.*, 2011). It has long been known that stretch reflex responses can allow for effective resistance against unwanted postural perturbations (Nichols and Houk, 1976). Moreover, reflex responses can also be modulated subconsciously to accommodate the execution of voluntary movements. For example, in tasks involving active reaching, significant modulation of reflex responses has been observed according to target shape (Nashed, Crevecoeur and Scott, 2012), obstacles (Nashed, Crevecoeur and Scott, 2014), static and moving targets (Cluff and Scott, 2015), and target cue direction (Pruszynski, Kurtzer and Scott, 2008; Scott, 2016).

In the upper extremities, the SLR is often characterised by a pre-load sensitivity referred to as

automatic gain scaling, whereas the LLR is found to be robustly task-dependent (Pruszynski *et*

al., 2009, 2011). More recently, the early LLR has been shown to contain a stabilising

component modulated independently of the voluntary action planned for a queued target, while

the late LLR displays task dependency (Lee and Perreault, 2019). Such studies reinforce the

idea that at least certain components of the LLR are highly adaptable and not strictly reflex in

nature (Shemmell, Krutky and Perreault, 2010). As mentioned above, goal-directed feedback

controllers affecting LLRs are thought to be already loaded during the reach preparation phase

(Ahmadi-Pajouh, Towhidkhah and Shadmehr, 2012). Although it is still unclear what specific

mechanism produces this tuned reflex modulation before movement initiation and exactly how

cortical preparation manifests as improved motor performance, it is currently believed that

cortical preparatory activity serves a dynamic system that defines the initial conditions for the

progression of goal-directed movement (Churchland *et al.*, 2010, 2012).

Recent findings propose that movement preparation includes the independent and goal-directed

tuning of muscle spindles, leading to a congruent modulation of reflex gains (Papaioannou and

Dimitriou, 2021). These findings suggest that two output mechanisms are involved in the

preparation of voluntary reaching, i.e., one targeting alpha motor neurons and another

implicating independent control of gamma motor neurons (Dimitriou, 2021, 2022). By

modulating the sensitivity of muscle spindles and stretch reflexes during preparation via gamma

control, the nervous system may adjust sensory feedback and reflex motor responses (i.e.,

muscle stiffness during movement) independently of any coinciding muscle force during

preparation. For instance, reflex gains are tuned according to the target direction in a way that

facilitates the upcoming voluntary reach (Torell *et al.*, 2023). Specifically, this goal-directed

tuning of stretch reflex gains, including at the SLR epoch, was present or enhanced when the

homonymous muscle was unloaded (i.e., antagonist load) and the preparatory delay was

sufficiently long (>250 ms and <750 ms). However, in this behavioural paradigm, the minimal

preparatory delay required for goal-directed tuning of stretch reflex gains was not quantified,

and the temporal evolution of such tuning is currently unclear.

In this study, we examine how a series of multiple preparatory delays impact the goal-directed

modulation of stretch reflex gains of the pectoralis, anterior and posterior deltoid muscles. To

achieve this, we used electromyographic (EMG) signals to quantify stretch reflex responses of

healthy subjects that experienced perturbations of their right upper limb introduced by a robotic

manipulandum during a delayed-reach task. Our findings clarify the minimal preparatory time

needed for goal-directed tuning of stretch reflex gains and reveal how these gains are modulated

as a function of preparation duration.

Methods

Subjects

A total of 21 right-handed and neurologically healthy subjects (10 males of mean age $27.1 \pm$

5.3 years and 11 females of mean age 24.1 ± 2.1 years) participated in the study. Data from four

subjects were not included in statistical analyses, one due to poor adaptation to the task and

three due to low-quality EMG signals. The remaining 17 subjects were eight males of mean age

27.6 ± 5.9 years and nine females of mean age 24.4 ± 2.0 years. All 21 subjects were financially

compensated for their participation and gave informed, written consent before participating in

the study. The subjects gave written informed consent, and the study was performed in line with

the principles of the Declaration of Helsinki. It was approved by the Regional Ethical Review

Board in Umeå.

Experimental setup

Subjects sat upright in a customised adjustable chair in front of the Kinarm robotic platform

(Kinarm End-Point Robot, BKIN Technologies). The subjects used their right hand to grasp the

robotic manipulandum (Fig. 1a). The right forearm was placed inside a customised foam

structure, resting on an airsled that allowed frictionless arm movement in a 2D plane. To ensure

a secure mechanical connection, the forearm, foam-cushioned airsled, the Kinarm handle, and

the hand were secured using a leather fabric with Velcro attachments. This attachment also

fixed the wrist in straight alignment with the forearm throughout the experiment. The robotic

platform measured kinematic data regarding the hand's position, and sensors inside the robotic

handle recorded the forces exerted by the subjects' right hand (six-axis force transducer;

Mini40-R, ATI Industrial Automation). Position and force data were sampled at 1 kHz.

Bipolar surface EMG electrodes (Bagnoli DE-2.1, Delsys Inc., Natick, MA, USA) with

10.0x1.0 mm silver contacts and 10 mm inter-electrode distance were placed on the pectoralis

major, posterior deltoid, and anterior deltoid muscles. Before placing the electrodes, the skin

was cleaned with alcohol, and a double-sided adhesive tape was attached to the electrodes and

coated with conductive gel. The electrodes were secured using surgical tape to ensure good

contact throughout the session. A ground electrode (Dermatrode HE-R Reference Electrode

type 00200–3400, 5 cm diameter, American Imex) was placed on the processus spinosus of the

C7 region. The EMG signals were sampled at 1 kHz, bandpass filtered at 20-500 Hz, and A/D

converted with 16-bit resolution using the Bagnoli EMG amplifier (Delsys Inc., Natick, MA,

USA).

Fig. 1 *The robotic manipulandum and experimental timeline. (**a**) Participants were seated upright in an adjustable chair with their right (dominant) hand secured to a robotic handle. A visual cursor representing the position of the hand, two visual targets and the origin were projected onto a one-way mirror, which also prevented a direct view of the hand (for details, see Methods). (**b**) Each trial began when the cursor was moved to the origin. A 4N load (800 ms rise-time and 1200 ms hold) was then applied in the front-and-left and right-and-back direction (135 and 315 degrees, respectively). During this phase, participants had to resist the load and maintain their hand at the origin. A visual target was then cued (through a red filled circle) for 250, 300, 350, 400, 450 or 500 ms. A position-controlled perturbation of the hand is then applied in the 135- or 315-degree direction (3.5 cm over 150 ms). The position of the visual cursor was frozen at the origin during the haptic perturbation. Upon completion of the perturbation, the cued target turned into a green-filled circle (the "Go" signal), and the participants were required to move the cursor to this target promptly*

Experimental design

The experimental design of the delayed-reach task involving haptic perturbations is summarised in Fig. 1 and is similar to what has been employed elsewhere, e.g., (Torell *et al.*, 2023). Specifically, a one-way mirror prevented a view of the hand and robotic handle, and visual

stimuli were projected in the plane of movement. Visual stimuli included a moving white cursor

of 1 cm in diameter, representing the hand's position. In addition, two targets (1.25 cm radius)

located 9 cm from the centre point ('origin'; 0.75 cm radius) were displayed as orange circle

outlines unless cued. The targets were placed in a front-and-left and right-and-back direction,

corresponding to 135 and 315 degrees.

Each trial started by moving the cursor to the origin. For the trial to progress, the cursor had to

remain immobile inside the origin for a random wait period between 1 and 1.5 s, after which a

4N load (800 ms rise time and 1200 ms hold) was applied in the 135 or 315 degree direction

for 2 s. One of the targets was then cued by suddenly turning from an orange outline to a filled

red circle. After a pre-defined preparatory delay (250, 300, 350, 400, 450 or 500 ms), a position-

controlled perturbation of the hand was applied in one of the two directions (3.5 cm with 150

ms rise-time and no hold). The cursor position was frozen during the perturbation. At the onset

of the perturbation, the filled red circle also turned green (the "Go" signal). At the end of the

perturbation, the subjects had to reach and remain within the target for 300 ms. After this, the

subjects received visual feedback on their performance, and a correct trial was defined as

reaching the target between 400 and 1400 ms after perturbation onset. Moving the cursor to the

origin began the subsequent trial.

This study represents a full factorial design without centre points, i.e., each block contained 48

unique trials in a randomised order comprising two targets to be cued, two load directions, two

perturbation directions, and six preparatory delays (2x2x2x6=48 trials). For the first five

subjects, there was a total of 13 blocks, while for the latter subjects, there were 18 blocks,

resulting in 864 trials. The subjects were instructed to take one-minute breaks after

approximately consecutive blocks. However, the subjects were free to rest after completing any

trial by pulling the cursor to the side of the workspace.

Data processing

EMG data were high-pass filtered using a third-order, zero-phase-lag Butterworth filter with a

20 Hz cutoff and rectified. For each trial, the time between issuing the perturbation and

movement onset (defined as 5% of peak hand velocity) was virtually identical across trials and

subjects, given the nature of the hand perturbation (i.e., 18 ms). Therefore, all data were aligned

on perturbation onset as determined by the KINARM and then shifted by a fixed delay of 18

ms. The raw EMG data were normalised by z-scoring, as described elsewhere (Dimitriou, 2014,

2016), to allow for inter-subject analysis. Briefly, this normalisation procedure involves

concatenating all EMG signals from all blocks and calculating each muscle's sample mean and

standard deviation separately. We also removed the offset of the baseline around the

perturbation (-25 to 25 ms) to ensure it was zero before averaging.

The first four blocks of trials were viewed as familiarisation trials and were not included in the

subsequent analyses (see Results section). We removed trials (on average 7% across all

participants) when the EMG or kinematics had a wiggly baseline when expected to be stable,

e.g., due to the subject not being fully relaxed. For the EMG, a wiggly baseline within 200 and

0 ms before the perturbation was defined empirically as above 0.1 in mean absolute deviation.

For the kinematics, the hand positions (both x- and y-axis) were defined empirically as outliers

if they were 15 cm from the centre point within the first 15 ms after the perturbation.

Since this study focused on stretch reflex responses, we only analysed data from stretching

muscles by analysing specific combinations of muscle and perturbation direction. To simplify

analyses of individual muscles, the median EMG signal of each muscle was generated for each

subject and trial type (i.e., for each load, perturbation, preparatory delay, and target direction),

where the EMG signals were smoothed with a moving mean of five ms). Data directly

representing goal-directed differences were generated by contrasting (subtracting) EMG values

in trials where reaching the cued target required stretching the homonymous muscle vs. values

from corresponding epochs in trials where reaching the target required muscle shortening.

Statistical analysis

We analysed the goal-directed differences in stretch reflex responses for the different muscles

and preparation delays using the Wilcoxon signed rank test with the False Discovery Rate

(FDR) method to correct for multiple comparisons. We also tested the differences in the stretch

reflex modulation between preparatory delays using the Kruskal-Wallis test with Fisher's least

significant difference (LSD) post-hoc tests. We used a significance level of 0.05. All data

processing and statistics were performed using MATLAB (Version 2023b; MathWorks).

Results

Task performance

The subjects' performance in the delayed-reach task reached a plateau relatively quickly, with

the average performance in terms of % correct trials being high and stable after the first four

blocks (Fig. 2a). Therefore, the first four blocks of trials were deemed familiarisation blocks

and were not included in subsequent statistical analyses. Although the subjects generally

performed well at the task, each subject occasionally initiated movements before receiving the

'Go' cue (2.6 ± 2.3 % of all trials), a phenomenon mainly observed during the longest

preparatory delays, i.e., 450-500 ms (Fig. 2b). Although infrequent overall, these "false starts"

persisted despite repeated instructions to wait for the 'Go' cue. That is, the experimenter

provided relevant instructions before data collection began and repeatedly after false starts, but

interestingly, a relatively small number of such trials often persisted throughout the experiment.

Fig. 2 *Reach performance and premature movement initiation. (**a**) The subjects' performance*

at the task, in terms of % correct trials in each block of trials. A correct trial involved reaching

the cued target within a certain time interval (see main text for more details). Performance

reached a high and rather stable plateau after the first four blocks, which were deemed

*familiarisation blocks and not included in subsequent analyses. (**b**) Each subject occasionally*

initiated movements prematurely, i.e., before the 'Go' cue (2.6 ± 2.3 % of all trials), mainly

during the longest preparatory delays (450-500 ms). These "false starts" occurred even though

the subjects were repeatedly instructed to wait for the 'Go' cue

Goal-directed modulation of the stretch reflex responses – Pectoralis muscle

250 A representative example of hand positions, forces, and z-scored pectoralis EMG activity

251 averaged across subjects for trials with a 400 ms preparatory delay is illustrated in Fig. 3a

252 (unloaded pectoralis) and another in Fig. 3b (loaded pectoralis). As indicated in these figures,

a preparatory delay of 400 ms is sufficient for inducing goal-directed tuning of SLR and LLR

responses, at least when the homonymous muscle is unloaded (Fig. 3a). Also, note that the

averaged hand positions and forces (along the x- and y-axis) are virtually identical during the

haptic perturbation, regardless of the load direction (Fig. 3).

Fig. 3 *Averaged responses across subjects for trials with a preparatory delay of 400 ms.*

Averaged hand positions, haptic forces, and z-scored EMG activity across subjects for trials

*with a 400 ms preparatory delay when the pectoralis muscle was unloaded (**a**) and loaded (**b**).*

The vertical dashed lines indicate the epochs of the short- and long-latency stretch reflexes

(i.e., SLR, early LLR, and late LLR) and the voluntary activity epoch ('VOL'). Curve shading

denotes ± 1 SE. In addition to z-transformation, the baseline offset of the EMG signals (pre-

reflex activity) was removed at the level of single trials (see main text for more details)

Our analyses indicated that goal-directed pectoralis EMG activity across subjects varied as a

function of preparatory delay (Fig. 4a). Analysis of individual EMG signals of the unloaded

pectoralis (i.e., individual subject averages across trials) indicated a significant goal-directed

difference in SLR ($p < 0.05$) for the 300-500 ms preparation delays (Fig. 4b). In contrast, we

found no significant goal-directed difference in SLR when the preparatory delay was 250 ms.

When the pectoralis was loaded, analyses indicated no goal-directed differences in SLR,

regardless of the preparatory delay duration (Fig. 4c). Regarding the early and late LLR of the

pectoralis muscle, analyses showed a significant goal-directed difference ($p < 0.01$) at all

preparatory delays and load conditions (Fig. 4d-g).

Pairwise comparison of differences in SLR EMG in the unloaded pectoralis muscle as a

function of preparation duration (Fig. 4b) indicated significantly weaker goal-directed tuning

of SLR following the shortest delay (250 ms) vs. all other delays ($p < 0.05$), except for the 450

ms preparation delay ($p = 0.057$). For the early LLR (Fig. 4d), analyses revealed a significant

difference in EMG activity only between the 250 and 400 ms delays ($p < 0.05$). For the late

LLR (Fig. 4f), we found significantly weaker goal-directed tuning of EMG activity in the

shortest preparatory delay vs. all other delays ($p < 0.05$), except for the 300 ms one ($p = 0.31$).

Analyses indicated no significant difference in goal-directed tuning of voluntary EMG activity

between the preparatory delays (not shown).

Fig. 4 *Goal-directed difference of pectoralis EMG activity. (**a**) Goal-directed difference of*

*pectoralis EMG as a function of preparatory delay and load condition. (**b**) There was a goal-*

directed difference in SLR of the unloaded pectoralis following 300-500 ms preparatory delays.

There was a significant difference in SLR EMG between the shortest preparatory delay (250

*ms) and all other delays (p < 0.05), except for the 450 ms one (p = 0.057). (**c**) There were no*

goal-directed differences in SLR responses of the loaded pectoralis, regardless of preparatory

*delay duration. (**d-e**) There was a significant goal-directed difference in early LLR across all*

preparatory delays (p < 0.01) for both the unloaded and loaded muscles. However, there was

only a significant difference in early LLR between the 250 and 400 ms delays (p < 0.05) when

*the muscle was unloaded. (**f-g**) There was a significant goal-directed difference in late LLR*

across all preparatory delays (p < 0.01) for both the unloaded and loaded muscles. There was

a significant difference in late LLR between the shortest delay (250 ms) and all other delays (p

< 0.05), except for the 300 ms (p = 0.31). Dots represent single-subject EMG values averaged

across the relevant trials (see Methods), grey background rectangles represent upper and lower

quartiles, and thin vertical lines represent 95% data range. * p < 0.05, ** p < 0.01, *** p <

0.001

Goal-directed modulation of the stretch reflex responses – Posterior and anterior deltoid

muscles

There was a significant goal-directed difference in SLR EMG of the unloaded posterior deltoid

muscle (i.e., pectoralis antagonist) following preparatory delays of 300-400 ms ($p < 0.05$; Fig.

5a). A pairwise comparison of differences in SLR EMG showed no significant difference

between 300 and 400 ms preparatory delays ($p > 0.05$; Fig. 5a). In contrast, there was a

significant difference between 350/400 ms and {250, 450, 500} ms ($p < 0.05$). For the early

and late LLR, there was a significant goal-directed difference across all preparatory delays (p

< 0.001; Fig. 5b-c). A pairwise comparison of differences in early LLR EMG showed a

significant difference between the shortest delay (250 ms) and all other delays ($p < 0.05$), except

for the 300 ms delay ($p = 0.43$). Analyses indicated no significant impact of preparation duration

on the voluntary muscle activity (not shown).

In contrast to the pectoralis and posterior deltoid muscles, there was no goal-directed difference

in the SLR of the anterior deltoid (Fig. 5d). However, there was a goal-directed difference

across all preparatory delays in the early and late LLR ($p < 0.05$; Fig. 5e-f), except for the 250

ms delay in early LLR (Fig. 5f). A pairwise comparison of differences in SLR EMG showed

no significant difference between the preparatory delays ($p > 0.05$; Fig. 5d). For the early LLR

(Fig. 5e), we found a significant difference between the shortest delay (250 ms) and all other

delays ($p < 0.05$), except for 300 ms and 450 ms. For the late LLR (Fig. 5f), the longer

preparatory delays were associated with a larger difference in EMG activity than the shorter

delays. Finally, analyses indicated no significant impact of preparation duration on the

voluntary muscle activity (not shown).

Fig. 5 *Goal-directed difference of the unloaded posterior and anterior deltoid muscles. (**a**)*

There was a goal-directed difference in SLR EMG of the posterior deltoid at preparatory delays

of 300-400 ms (p < 0.05), and there was no significant difference in SLR EMG among those

delays. In contrast, there was a significant difference between 350/400 ms and {250, 450, 500}

*ms (p < 0.05). (**b-c**) There was a goal-directed difference in early and late LLR EMG of the*

posterior deltoid across all preparatory delays (p < 0.001). For the early LLR activity, there

was a significant difference between the shortest delay (250 ms) and all other delays (p < 0.05),

*except for the 300 ms delay (p = 0.43). (**d**) There was no goal-directed difference in the SLR of*

*the anterior deltoid. (**e-f**) There was a goal-directed difference in early and late LLR EMG of*

the anterior deltoid across all preparatory delays (p < 0.001), except for the 250 ms delay in

early LLR. For the early LLR (e), we found a significant difference between the shortest delay

(250 ms) and all other delays (p < 0.05), except for 300 ms and 450 ms. For the late LLR (f),

the longer preparatory delays were associated with a larger difference in EMG activity than

*the shorter delays. * p < 0.05, ** p < 0.01, *** p < 0.001*

Discussion

This study aimed to identify the minimum preparatory time required for goal-directed tuning

of SLR and LLR responses and to characterise the incremental evolution of such tuning as a

function of preparation duration. The two main findings of this study are that 1) a preparatory

delay of 300 ms is sufficient for goal-directed tuning of SLR responses in the pectoralis and

posterior deltoid muscles (e.g., Figs. 4-5) and 2) that an additional 100 ms (i.e., preparation of

400 ms) allows for even stronger goal-directed modulation of stretch reflex responses.

Our results confirm previous findings that preparatory delays of 250 ms are not associated with

goal-directed SLR responses (Torell *et al.*, 2023). While pertinent underlying mechanisms may

already be at work during preparatory delays of 250 ms or shorter (Papaioannou and Dimitriou,

2021), longer preparatory delays are required for the systematic manifestation of goal-directed

tuning of SLR responses, as assessed by surface EMG. Also, in line with previous findings, we

show that unloading of the homonymous muscle facilitates the goal-directed tuning of SLRs.

In contrast, muscle loading leads to 'automatic' gain-scaling of the SLR, regardless of goal and

LLRs responses are robustly goal-dependent, regardless of background muscle loading

(Pruszynski *et al.*, 2009; Papaioannou and Dimitriou, 2021; Torell *et al.*, 2023).

The optimal preparatory delay of 400 ms, found here, is approximately 200 ms longer than the

goal-directed modulation of reflexes compared to what Yang and colleagues found (Yang *et*

al., 2011). This difference is likely related to methodological differences and our participants

preparing to perform a multi-joint task, not a single-joint task. Single-joint movements allow

for a more controlled study of the preparatory processes. However, multi-joint movements

provide a more naturalistic context for studying how the sensorimotor system coordinates

complex actions, such as reach-to-grasp. In other words, it appears that the nervous system

needs an additional ~200 ms to sufficiently incorporate the goal-directed preparation of multi-

joint reaching vs. single-joint movement.

Interestingly, for preparatory delays longer than 400 ms, the goal-directed tuning of the SLR

(at least) seems to drop, and such longer preparatory delays are associated with more frequent

"false starts", i.e., trials with premature movement initiation (i.e., before receiving a "Go"

signal). False starts occurred even though the subjects were instructed to wait for the "Go"

signal before and after these occurred, suggesting the engagement of specific cognitive

mechanisms when the preparatory delay is long enough. The false start trials could occur for a

myriad of reasons. When a series of alternative preparatory durations are involved in a task,

given sufficient preparatory time the participants might explicitly or implicitly try to predict

when the "Go" cue will occur, and this process biasing their decision to initiate movement i.e.,

anticipating the Go cue to come sooner than it does for the longer preparatory delays. Some

participants might tend to act more impulsively and, therefore, be more likely to act in

anticipation of a "Go" signal. When trying to wait, the CNS (and PNS) prepares for the

upcoming movement. The cognitive preparation can sometimes "leak" into premature muscle

activation, resulting in a false start (Forgaard *et al.*, 2019). Our findings suggest ≥ 450 ms

preparatory delay is sufficient time for this cognitive motor leakage to occur in delayed reach.

In Figs. 4-5, we observed a decrease in goal-directed tuning of stretch reflex responses (Δ EMG

activity) and voluntary activity, around 450 ms, for different preparation delays in the unloaded

pectoralis muscle and posterior deltoid. The reason for this decrease is not directly apparent and

requires further investigation. One can hypothesise that around 450 ms, there is a transient shift

in the dominant preparatory processes. For instance, there might be an initial phase of more

reflexive motor activity followed by a more specific phase of inhibition or refinement of the

motor control policy involving higher cognitive functions. This could temporarily reduce the

overall excitability of the reflex pathways. Preparatory delays longer than 450 ms could be

sufficient for the higher cortical preparatory processes to dominate, and including EEG in future

studies could shed light on this interesting phenomenon. On the other hand, having many

different delays may engage a different mechanism than a task involving a discrete dichotomy

in preparatory delays, i.e., long vs. short delay, with one being too short and one being

sufficient. Indeed, this could explain why 250 ms was too little while 750 ms sufficed in a

previous study (Torell *et al.*, 2023), also characterised by a far smaller number of false-start

trials.

In contrast to the case of the pectoralis and posterior deltoid muscle, there was no goal-directed

tuning of SLR responses in the anterior deltoid muscle. In our task, the anterior deltoid

functioned as a pectoralis synergist. This, in turn, may imply that, at least after task

familiarisation, the sensorimotor system does not apply independent fusimotor control to all

synergists potentially engaged in a task but rather to a subset of these. In other words,

independent fusimotor control appears highly selective across the muscles that could be

similarly engaged as agonists and antagonists in a motor task. Congruent with the idea that

feedback controllers are loaded before movement initiation (Ahmadi-Pajouh, Towhidkhah and

Shadmehr, 2012), we identify the minimum amount of time required by these controllers for

manifesting goal-directed modulation of reflex responses, highlighting that feedback

controllers likely also engage top-down fusimotor control (Dimitriou, 2022).

In the current study, reflex responses appear strongly inhibited or even absent under certain

experimental conditions. This can be attributed to the context-dependent modulation of the

stretch reflex. In some situations, the CNS might actively suppress or gate the stretch reflex to

allow for controlled movement (Shemmell, An and Perreault, 2009). If muscle stretch occurs

during such higher-order suppression or gating, the resulting contraction might be weaker than

expected, potentially being perceived as absent or a minimal response. In the current study, the

task strives to be as naturalistic as possible, allowing the subjects to perform the task without

being too constrained, which may result in a different recruitment of motor units (Hodson-Tole

and Wakeling, 2009). Since we used bipolar surface EMG electrodes to quantify the stretch

reflex responses in this naturalistic task, this may partly explain the variability in reflex

responses across participants. To overcome this limitation, using multi-channel surface EMG

to cover a larger part of the muscle would lead to higher signal-to-noise estimates of the stretch

reflex responses.

In conclusion, we examined how multiple preparatory delays impact the goal-directed

modulation of stretch reflex gains of the pectoralis, anterior and posterior deltoid muscles in a

multi-joint reaching task. We found that a preparatory delay of 300 ms is sufficient for goal-

directed tuning of SLR responses in the pectoralis and posterior deltoid muscles, and 400 ms

allows for even stronger goal-directed modulation of stretch reflex responses. Our results clarify

the minimum preparatory time required for goal-directed tuning of stretch reflex gains at the

level of the dominant upper limb and characterise the relationship between reflex gains and

preparation duration. We show that this relationship is not strictly linear, likely reflecting the

interplay of multiple feedback mechanisms functioning at different time frames.

Declarations

Consent to participate and publish

The subjects gave written informed consent to participate in this study. The authors affirm that human research participants provided informed consent to submit the manuscript to a scientific journal.

Funding

This work was supported by grants awarded to M.D. by Hjärnfonden (FO2024-0425-HK-88) and R.R. by the Swedish Research Council for Sport Science (P2025-0173).

Competing interests

The authors have no competing interests to declare that are relevant to the content of this study.

Ethics approval

The subjects gave written informed consent, and the study was performed in line with the principles of the Declaration of Helsinki. It was approved by the Regional Ethical Review Board in Umeå.

Data and code availability

Data and source code are available upon request.

Authors' contribution statements

All authors contributed to the study's conception and design. RR performed data curation and analysis. RR wrote the first draft of the manuscript. All authors reviewed and edited the manuscript. All authors read and approved the final manuscript.

References

Ahmadi-Pajouh, M.A., Towhidkhah, F. and Shadmehr, R. (2012) 'Preparing to Reach: Selecting an Adaptive Long-Latency Feedback Controller', *Journal of Neuroscience*, 32(28), pp. 9537–9545. Available at: https://doi.org/10.1523/JNEUROSCI.4275-11.2012.

Batista, A.P. *et al.* (2007) 'Reference Frames for Reach Planning in Macaque Dorsal Premotor Cortex', *Journal of Neurophysiology*, 98(2), pp. 966–983. Available at: https://doi.org/10.1152/jn.00421.2006.

Churchland, M.M. *et al.* (2010) 'Cortical Preparatory Activity: Representation of Movement or First Cog in a Dynamical Machine?', *Neuron*, 68(3), pp. 387–400. Available at: https://doi.org/10.1016/j.neuron.2010.09.015.

Churchland, M.M. *et al.* (2012) 'Neural population dynamics during reaching', *Nature*, 487(7405), pp. 51–56. Available at: https://doi.org/10.1038/nature11129.

Churchland, M.M., Santhanam, G. and Shenoy, K.V. (2006) 'Preparatory Activity in Premotor and Motor Cortex Reflects the Speed of the Upcoming Reach', *Journal of Neurophysiology*, 96(6), pp. 3130–3146. Available at: https://doi.org/10.1152/jn.00307.2006.

Cluff, T. and Scott, S.H. (2015) 'Apparent and Actual Trajectory Control Depend on the Behavioral Context in Upper Limb Motor Tasks', *Journal of Neuroscience*, 35(36), pp. 12465–12476. Available at: https://doi.org/10.1523/JNEUROSCI.0902-15.2015.

Dimitriou, M. (2014) 'Human Muscle Spindle Sensitivity Reflects the Balance of Activity between Antagonistic Muscles', *Journal of Neuroscience*, 34(41), pp. 13644–13655. Available at: https://doi.org/10.1523/JNEUROSCI.2611-14.2014.

Dimitriou, M. (2016) 'Enhanced Muscle Afferent Signals during Motor Learning in Humans', *Current Biology*, 26(8), pp. 1062–1068. Available at: https://doi.org/10.1016/j.cub.2016.02.030.

Dimitriou, M. (2021) 'Crosstalk proposal: There is much to gain from the independent control of human muscle spindles', *The Journal of Physiology*, 599(10), pp. 2501–2504. Available at: https://doi.org/10.1113/JP281338.

Dimitriou, M. (2022) 'Human muscle spindles are wired to function as controllable signal-processing devices', *eLife*. Edited by E. Perreault and T.R. Makin, 11, p. e78091. Available at: https://doi.org/10.7554/eLife.78091.

Forgaard, C.J. *et al.* (2019) 'Influence of kinesthetic motor imagery and effector specificity on the long-latency stretch response', *Journal of Neurophysiology*, 122(5), pp. 2187–2200. Available at: https://doi.org/10.1152/jn.00159.2019.

Ghez, C. *et al.* (1997) 'Discrete and continuous planning of hand movements and isometric force trajectories', *Experimental Brain Research*, 115(2), pp. 217–233. Available at: https://doi.org/10.1007/PL00005692.

Hammond, P. (1956) 'The influence of prior instruction to the subject on an apparently involuntary neuromuscular response', *J. Physiol.*, 132, pp. 17–19.

Hocherman, S. and Wise, S.P. (1991) 'Effects of hand movement path on motor cortical activity in awake, behaving rhesus monkeys', *Experimental Brain Research*, 83(2), pp. 285–302. Available at: https://doi.org/10.1007/BF00231153.

Hodson-Tole, E.F. and Wakeling, J.M. (2009) 'Motor unit recruitment for dynamic tasks: current understanding and future directions', *Journal of Comparative Physiology B*, 179(1), pp. 57–66. Available at: https://doi.org/10.1007/s00360-008-0289-1.

Kutas, M. and Donchin, E. (1974) 'Studies of Squeezing: Handedness, Responding Hand, Response Force, and Asymmetry of Readiness Potential', *Science*, 186(4163), pp. 545–548. Available at: https://doi.org/10.1126/science.186.4163.545.

Lee, H. and Perreault, E.J. (2019) 'Stabilizing stretch reflexes are modulated independently from the rapid release of perturbation-triggered motor plans', *Scientific Reports*, 9(1), p. 13926.

Available at: https://doi.org/10.1038/s41598-019-50460-1.

Marsden, C.D., Merton, P.A. and Morton, H.B. (1972) 'Servo Action in Human Voluntary Movement', *Nature*, 238(5360), pp. 140–143. Available at: https://doi.org/10.1038/238140a0.

Messier, J. and Kalaska, J.F. (2000) 'Covariation of Primate Dorsal Premotor Cell Activity With Direction and Amplitude During a Memorized-Delay Reaching Task', *Journal of Neurophysiology*, 84(1), pp. 152–165. Available at: https://doi.org/10.1152/jn.2000.84.1.152.

Nashed, J.Y., Crevecoeur, F. and Scott, S.H. (2012) 'Influence of the behavioral goal and environmental obstacles on rapid feedback responses', *Journal of Neurophysiology*, 108(4), pp. 999–1009. Available at: https://doi.org/10.1152/jn.01089.2011.

Nashed, J.Y., Crevecoeur, F. and Scott, S.H. (2014) 'Rapid Online Selection between Multiple Motor Plans', *Journal of Neuroscience*, 34(5), pp. 1769–1780. Available at: https://doi.org/10.1523/JNEUROSCI.3063-13.2014.

Nichols, T.R. and Houk, J.C. (1976) 'Improvement in linearity and regulation of stiffness that results from actions of stretch reflex', *Journal of Neurophysiology*, 39(1), pp. 119–142. Available at: https://doi.org/10.1152/jn.1976.39.1.119.

Papaioannou, S. and Dimitriou, M. (2021) 'Goal-dependent tuning of muscle spindle receptors during movement preparation', *Science Advances*, 7(9), p. eabe0401. Available at: https://doi.org/10.1126/sciadv.abe0401.

Pruszynski, J.A. *et al.* (2009) 'Temporal Evolution of "Automatic Gain-Scaling"', *Journal of Neurophysiology*, 102(2), pp. 992–1003. Available at: https://doi.org/10.1152/jn.00085.2009.

Pruszynski, J.A. *et al.* (2011) 'Primary motor cortex underlies multi-joint integration for fast feedback control', *Nature*, 478(7369), pp. 387–390. Available at: https://doi.org/10.1038/nature10436.

Pruszynski, J.A., Kurtzer, I. and Scott, S.H. (2008) 'Rapid Motor Responses Are Appropriately Tuned to the Metrics of a Visuospatial Task', *Journal of Neurophysiology*, 100(1), pp. 224–238. Available at: https://doi.org/10.1152/jn.90262.2008.

Scott, S.H. (2016) 'A Functional Taxonomy of Bottom-Up Sensory Feedback Processing for Motor Actions', *Trends in Neurosciences*, 39(8), pp. 512–526. Available at: https://doi.org/10.1016/j.tins.2016.06.001.

Shadmehr, R. and Krakauer, J.W. (2008) 'A computational neuroanatomy for motor control', *Experimental Brain Research*, 185(3), pp. 359–381. Available at: https://doi.org/10.1007/s00221-008-1280-5.

Shemmell, J., An, J.H. and Perreault, E.J. (2009) 'The Differential Role of Motor Cortex in Stretch Reflex Modulation Induced by Changes in Environmental Mechanics and Verbal Instruction', *Journal of Neuroscience*, 29(42), pp. 13255–13263. Available at: https://doi.org/10.1523/JNEUROSCI.0892-09.2009.

Shemmell, J., Krutky, M.A. and Perreault, E.J. (2010) 'Stretch sensitive reflexes as an adaptive mechanism for maintaining limb stability', *Clinical Neurophysiology*, 121(10), pp. 1680–1689. Available at: https://doi.org/10.1016/j.clinph.2010.02.166.

Tanji, J. and Evarts, E.V. (1976) 'Anticipatory activity of motor cortex neurons in relation to direction of an intended movement', *Journal of Neurophysiology*, 39(5), pp. 1062–1068. Available at: https://doi.org/10.1152/jn.1976.39.5.1062.

Todorov, E. and Jordan, M.I. (2002) 'Optimal feedback control as a theory of motor coordination', *Nature Neuroscience*, 5(11), pp. 1226–1235. Available at: https://doi.org/10.1038/nn963.

Torell, F. *et al.* (2023) 'Assistive Loading Promotes Goal-Directed Tuning of Stretch Reflex Gains', *eNeuro*, 10(2). Available at: https://doi.org/10.1523/ENEURO.0438-22.2023.

Wagner, M.J. and Smith, M.A. (2008) 'Shared Internal Models for Feedforward and Feedback Control', *Journal of Neuroscience*, 28(42), pp. 10663–10673. Available at: https://doi.org/10.1523/JNEUROSCI.5479-07.2008.

Wise, S.P. (1985) 'The primate premotor cortex: past, present, and preparatory', *Annual review of neuroscience*, 8, pp. 1–19. Available at: https://doi.org/10.1146/annurev.ne.08.030185.000245.

Yang, L. *et al.* (2011) 'Rapid motor responses quickly integrate visuospatial task constraints', *Experimental Brain Research*, 211(2), pp. 231–242. Available at: https://doi.org/10.1007/s00221-011-2674-3.

Yeo, S.-H., Franklin, D.W. and Wolpert, D.M. (2016) 'When Optimal Feedback Control Is Not Enough: Feedforward Strategies Are Required for Optimal Control with Active Sensing', *PLOS Computational Biology*, 12(12), p. e1005190. Available at: https://doi.org/10.1371/journal.pcbi.1005190.