Title: Long-term P fertilization influences microbial carbon use efficiency and soil organic

matter decomposition in non-allophanic Andosols.

Authors:

Wako KOIZUMI[1], Timothy J CLOUGH[2], Soichi KOJIMA[1], Tomoyuki MAKINO[1], Soh

SUGIHARA[3], Yoshitaka UCHIDA[4], Toru HAMAMOTO[1]*

Address:

[1]Graduate School of Agricultural Science, Tohoku University, 468-1 Aramaki Aoba Aoba-ku

Sendai Miyagi 9808572, Japan

[2]Faculty of Agriculture and Life Sciences, Lincoln University, Lincoln, New Zealand

[3]Graduate School of Agricultural Science, Tokyo University of Agriculture and Technology,

3-5-8 Saiwai-cho, Fuchu-shi, Tokyo 1838509, Japan

[4]Research Faculty of Agriculture, Hokkaido University, Kita 9 Nishi 9 Kita-Ku, Sapporo,

Hokkaido, 060-8589, Japan

*Correspondence: T Hamamoto, toru.hamamoto.d1@tohoku.ac.jp

Number of Figures: three

Number of Tables: two

Keywords: Microbial carbon use efficiency; Microbial respiration; Soil organic matter priming;

Phosphorus-mining; Non-allophanic Andosol

Abstract

Phosphorus (P) availability affects soil carbon (C) cycling such as microbial C use efficiency

(CUE) and priming effects (PEs). While non-allophanic Andosols are characterized by high

organic C content and strong P retention, the effects of different P fertilization regime on C

dynamics in these soils remain poorly understood. In this study, we conducted a 20-day

incubation experiment using ^{13}C-enriched glucose to investigate how different soil P levels

(Truog-P: 157 mg P kg^{-1} and 12 mg P kg^{-1}) impacted microbial C dynamics in non-allophanic

Andosols from contrasting field management practices. Our results showed that soil organic

matter (SOM) priming is associated with P fertilization management, with total primed CO_2-C

emissions remaining low in these soils. In the high-P soils, glucose and nitrogen (N) addition

resulted in negative PEs, whereas in the low-P soils, the same treatment stimulated microbial

SOM mining, resulting in positive PEs. Additionally, higher glucose-derived CUE was found in

the high-P soils than in low-P soils after 20 days of incubation. These findings suggest that

long-term P fertilization influences both substrate-induced microbial assimilation and SOM

decomposition, with P limitation potentially promoting SOM mining which, along with

concurrent soil acidity and exchangeable Al toxicity, modulates CUE. This study provides

insights for improving C sequestration in non-allophanic Andosols through soil fertility

management.

1. Introduction

43  Soil carbon (C) plays a central role in the global C cycle with the potential to mitigate the

44 elevated atmospheric C dioxide (CO_2) concentration (Jackson et al. 2017). Soil microbes can

45 degrade soil organic C (SOC) emitting it as CO_2 (Wieder et al. 2013). Increases in the stock of

46 SOC may occur as a result of increased microbial biomass, in response to inputs of C to the soil

47 (Kallenbach et al. 2016; Wang et al. 2021).

49  Microbial C use efficiency (CUE), the ratio of C allocated to microbial biomass growth to

50 that respired as CO_2, is positively correlated with SOC levels (Tao et al. 2023). The magnitude

51 of CUE has also been shown to depend on microbial community structure (Maynard et al. 2017),

52 and various abiotic factors, such as soil pH and exchangeable Al, which can alter microbial

53 metabolism and C allocation (Jones et al. 2019; Schroeder et al. 2024), clay content (Islam et al.

54 2023), nutrient availability (Manzoni et al. 2012; Spohn 2016), and substrate type (Jones et al.

55 2018). Limited nutrient availability and unbalanced substrate additions may lower CUE by

56 promoting excessive enzyme allocation and respiration, described as overflow metabolism (e.g.,

57 Tempest and Neijssel 1992; Russell and Cook 1995; Schimel and Weimtraub 2003; Keiblinger

58 et al. 2010). According to Manzoni et al. (2012), CUE is also influenced by nitrogen (N) and

59 phosphorus (P) availability, since microbes require these nutrients to balance anabolic and

catabolic processes. P is an essential nutrient for soil microbes, as it is a key component of DNA,

RNA, and has a critical role in phospholipid synthesis, and energy transfer. Depletion of P

resources constrains microbial investment in biomass production, shifting the C allocation

toward enzyme production, which can lower CUE (Zhai et al. 2022). In addition, limited P

availability represents a significant agronomic challenge, as it directly constrains plant

productivity and undermines soil health. Microbes compete with plants for P, thereby limiting

plant activity by reducing the plant-available P in P-limited soil (Jiang et al. 2024).

An imbalance of available nutrients leads to microbial SOM decomposition and nutrient

acquisition, observed as positive priming effects (PEs) (Blagodatskaya and Kuzyakov 2008;

Nottingham et al. 2015; Bernard et al. 2022). For example, the addition of a high C:N substrate

induces microbial N acquisition from SOM, a process known as 'N mining', and positive PEs

are found when compared to the addition of low C:N substrates, (Craine et al. 2007; Xu et al.

2024). Immobilization and nutrient mining also applies to P, with high C:P substrates inducing

positive PEs to occur under P-limited soil conditions. In some cases, this may also involve the

mobilization of P compounds from mineral phases through the mechanisms such as the action of

competing ligands, protons, or electron-donating compounds (Spohn 2024; Lieberman et al.

2025). Additionally, P addition to soils may contribute to SOM destabilization through abiotic

processes. Inputs of P promotes the desorption of the organic C adsorbed to mineral surface by

ligand exchange or the dissolution of the organic matter formed complexes with metal such as

aluminum (Al) and iron (Fe) (Kaiser and Zech 1999; Li et al. 2020; Spohn et al. 2022; Mori

2023). Previous studies have indicated that P addition increases the bioavailable C pool and C

mineralization, contributing to the SOM decomposition in various soils, including Andosols

(Miyazawa et al. 2013; Scott et al. 2015; Spohn and Schleuss 2019; Li et al. 2020).

Considerable research interest has recently focused on P-induced PEs and CUE, and an

increasing number of studies examining the effects of P availability on PEs have been

conducted using direct P substrate addition, with contrasting results: some showing higher

positive PEs but higher CUE (Wang et al. 2025), others with negative PEs (Wang et al. 2014;

Sawada et al. 2023) or negative PEs with high CUE (Zeng et al. 2025), or with higher positive

PEs but no effects on CUE (Mehnaz et al. 2019). However, short-term additions may not fully

capture the sustained microbial and abiotic responses observed under long-term fertilization.

Long-term P fertilization in the field provides a more realistic framework than short-term

addition for assessing PE responses to exogenous C inputs (Qin et al. 2024). Limited studies

using this approach compared to studies using direct P addition have also reported contrasting

results. Some observed increased CUE and no significant changes in SOC levels with sustained

P input (Li et al. 2025a), while others found enhanced SOM mineralization and positive PEs (Li

et al. 2023; Qin et al. 2024). Moreover, Widdig et al. (2020) reported that P fertilization had no

effect on CUE. These varied findings emphasize that microbial responses to P availability are

highly context-dependent (i.e., microbial nutrient mining hypothesis and microbial

stoichiometric decomposition hypothesis, Hessen et al. 2004; Craine et al. 2007). There remains

a critical knowledge gap in understanding how P availability influences microbial CUE and PEs

across different soil types and the quality or amount of substrate, particularly in soils with

unique properties such as Andosols.

Derived from volcanic ash, Andosols are mainly located on the Pacific coast (e.g., Japan,

Indonesia, New Zealand, and Chile), but are also present in parts of Europe (e.g., Italy; Iamarino

and Terribile 2008; Takahashi 2020; Matus et al. 2024). They are important for soil C

sequestration due to their abundance of Al-organic matter complexes which contributes to high

SOM stability (Nanzyo 2002; Takahashi and Dahlgren 2016). Andosols are classified according

to their significant P retention and are mainly divided into two groups: allophanic and

non-allophanic, which are distinguished by the active Al composition (Takahashi and Dahlgren

2016; IUSS Working Group WRB 2022). Non-allophanic Andosols, dominated by Al-organic

matter complexes, account for about 30% of Japanese arable Andosols (Saigusa and Matsuyama

1998). Non-allophanic Andosols are characterized by high acidity and high P sorption due to

Al-organic matter complexes (Shoji et al. 1987). Thus, non-allophanic variants display greater

Al phytotoxicity, which regulates the availability of P, consequently restricting crop growth (Ito

et al. 2011). Given the dependence of soil microbial activity on P availability, it is highly

expected that microbial CUE is affected by P application. Moreover, long-term P fertilization in

the field provides insights into the impacts of sustained P input on C decomposition, which

cannot be replicated in short-term laboratory experiments (Qin et al. 2024). However, it remains

unknown how CUE responds to long-term P addition in non-allophanic Andosols, and how

P-induced SOM changes interact microbial C cycling in non-allophanic Andosols.

In this study, we conducted an incubation experiment to investigate the effect of long-term P

fertilization on microbial C utilization in non-allophanic Andosols. We used ^{13}C glucose to

evaluate the impact of P availability on CUE and soil ^{13}C recovery. We hypothesized that: (i) in

unfertilized soils, substrate additions would promote P-mining, accelerating the positive PEs;

and (ii) higher CUE would be observed in P-fertilized soils, as the nutrient balance would better

match microbial requirements compared to unfertilized soils.

2. Materials and methods

2.1. Soil sampling

Soil samples were collected at 0-10 cm depth in April 2023 from two contrasting fertilization

fields at Field Science Center, Graduate School of Agricultural Science, Tohoku University,

Japan (38°74'N, 140°75 'E, 190 m above mean sea-level). The soils were classified as

non-allophanic Andosols according to the Soil Classification System of Japan (Obara et al.

2011). Two contrasting soils with different P content were sampled (Table 1). The HP soil,

which had been P-fertilized since 1987, exhibited a high Truog-P content (157.31 mg P kg^{-1}).

The amount of P input from the triple superphosphate and the fused magnesium phosphate

varied depending on the cultivation years, ranging from 31 to 361 kg P ha^{-1} yr^{-1}. Additionally,

cattle manure was applied in certain years. In contrast, the low-P soil (LP) has been cultivated

since 2009 without P fertilization, had a significantly lower Truog-P content (12.10 mg P kg^{-1}).

Detailed fertilization and cultivation histories are described in Table S1. The HP soil (pH 5.29)

had a significantly higher pH than the LP soil (pH 4.49). Correspondingly, the LP soil showed a

significantly higher amount of exchangeable Al (21.30 meq kg^{-1}) compared to the HP soil (0

meq kg^{-1}). The collected soils were sieved through a 2 mm sieve, and all visible roots were

removed, then soils were stored at 4□ until further use.

Table 1. Chemical and biological properties in HP and LP soils (mean (SD)). For each property, significant differences between HP and LP soils

were determined by independent samples t-tests.

	$pH_{(CaCl2)}$	Total C	Total N	Total P	C:N	MBC	MBP	log(MBC:MBP)	Resin P	Truog P	HCl-P
		(g C kg^{-1})	(g N kg^{-1})	(g P kg^{-1})		(mg C kg^{-1})	(mg P kg^{-1})	molar basis	(mg P kg^{-1})	(mg P kg^{-1})	(g P kg^{-1})
HP	5.29 (0.03)	88.07 (3.38)	5.23 (0.23)	2.35 (0.27)	16.83	1368.71 (56.55)	75.06 (18.31)	1.68	122.39 (6.31)	157.31 (5.07)	1.41 (0.029)
LP	4.49 (0.06)	91.24 (1.99)	5.36 (0.10)	1.37 (0.14)	17.01	1343.65 (142.09)	21.31 (11.63)	2.31	29.13 (1.33)	12.10 (0.31)	0.54 (0.063)
P values	<0.0001	n.s	n.s	<0.01	<0.05	n.s	<0.01	<0.05	<0.001	<0.0001	<0.0001

150 MBC, Microbial biomass C: MBP, microbial biomass P: MBC: MBP, their molar-based ratio: Resin P, Resin-extractable P. Truog P and HCl-P

151 represent Truog-extractable P and 1M HCl-extractable P, respectively.

2.2. Experimental design

Prior to the experiment, the sieved soils were pre-incubated in the dark at 25□ for 14 days

and maintained at a gravimetric moisture content of 45% (w/w). Pre-incubated soil was placed

in plastic bags (each equivalent to 100 g dry weight). Soil samples received a substrate solution

containing either: (1) water ("Control"), (2) ^{13}C-enriched glucose ("Glc"), or (3) ^{13}C-enriched

glucose and $(NH_4)_2SO_4$ ("Glc+N"). There were 24 samples in total (two soils × three glucose

levels × four replicates). The rate of added ^{13}C-enriched glucose (4 atom%, D-GLUCOSE

(U–$^{13}C_6$, 99%), Cambridge, 450 mg C kg^{-1} soil) was set to be 25 to 50% of the MBC in order to

increase microbial activity while avoiding exponential microbial growth and shifting microbial

community composition (Blagodatskaya and Kuzyakov 2008; Sawada et al. 2023). The

concentration of added mineral N (61 mg N kg^{-1} soil) was determined according to mean

microbial C:N:P ratios, 60:7:1 (Cleveland and Liptzin 2007). The experiment was then started

by wetting to 55% gravimetric soil moisture content with each substrate solution.

2.3. CO_2 emission

For measurement of soil respiration, each treatment was placed in a 450 mL airtight glass jar

with a CO_2 alkali trap (5 mL of 1 M NaOH) and incubated at 25□ for 20 days. Ten mL of 0.01

M HCl solution was also placed in the glass jar to maintain the soil moisture. The collected

NaOH solutions received 8 mL of 1 M $SrCl_2$ to form $SrCO_3$ precipitate, and the resulting

mixture of precipitate and solution was titrated using a 1 M HCl solution (Zibilske 1994; Qiao

et al. 2014). The collected NaOH solutions received 8 mL of 1 M $SrCl_2$ to form $SrCO_3$

precipitate and this was titrated using a 1 M HCl solution. After titrating, the precipitates were

washed three times with 30mL water and then dried at 105□. The dried $SrCO_3$ was packed in tin

capsules to analyze for ^{13}C enrichment on DELTA V plus isotope ratio mass spectrometer

(IRMS, Thermo Fisher Scientific, Bremen, Germany). The calculation of glucose-derived

$CO_2\text{-}C_{glc}$ was adapted from Mehnaz et al. (2019) and Brant et al. (2006):

$CO_2\text{-}C_{glc} = CO_2\text{-}C \times (^{13}C_c - {}^{13}C_{none}) / (^{13}C_{label} - {}^{13}C_{none})$ (1)

Where $CO_2\text{-}C$ is the cumulative $CO_2\text{-}C$ respired from the soil with glucose addition; $^{13}C_c$,

$^{13}C_{none}$ and ^{13}C labels are the ^{13}C atom% values of the $CO_2\text{-}C$ measured in the NaOH solutions

with and without glucose addition and the added glucose substrate, respectively. Soil-derived

$CO_2\text{-}C$ ($CO_2\text{-}C_{soil}$) in glc and glcN treatments was calculated as the difference between the total

($CO_2\text{-}C$) and glucose-derived $CO_2\text{-}C_{glc}$.

    2.4. Soil microbial biomass C and P

Soil microbial biomass C (MBC) was determined with a modified chloroform fumigated

method (Beck et al. 1997). Briefly, two samples of 5 g dry soil of each sample with and without

24 h fumigation using chloroform were extracted with 25 mL of 0.05 M K_2SO_4 and analyzed for

total organic C using a TOC-TN analyzer (TOC-V CPH/CPN, Shimadzu, Japan). The MBC was

calculated from the difference in total organic C concentrations in samples with and without

fumigation divided by an extraction efficiency of 0.45 (Beck et al. 1997). The ^{13}C atom% in

each extract was measured on an IRMS (Thermo Fisher Scientific, USA). The ^{13}C atom% of the

microbial biomass ($^{13}C_{mic}$) and the glucose-derived C in the microbial biomass ($MBC_{glucose}$) were

also calculated by the following equation (Mehnaz et al. 2019):

$^{13}C_{mic} = (C_f \times {}^{13}C_f \ - \ C_{nf} \times {}^{13}C_{nf}) / (C_f - C_{nf})$ (2)

$MBC_{glucose} = MBC \times \ ({}^{13}C_{mic,c} - {}^{13}C_{mic,none}) \ / ({}^{13}C_{label} - {}^{13}C_{mic,none})$ (3)

where C_f and $^{13}C_f$ are the amount of C and ^{13}C atom% in the fumigated extracts, and C_{nf} and

$^{13}C_{nf}$ are those in the non fumigated, respectively.

Organic C within K_2SO_4 extracts from non-fumigated soil was denoted as Dissolved organic C

(DOC). The terms $^{13}C_{mic,c}$ and $^{13}C_{mic,none}$ are the $^{13}C_{mic}$ values with and without glc addition.

Soil microbial biomass P (MBP) was measured using an anion exchange membrane method

according to Kouno et al. (1995). Briefly, soil samples (1 g dry mass equivalent) were shaken

and extracted in a 20 ml mixture of deionized water and resin for 16 h. For fumigated samples

only, 2 ml of hexanol was added instead of chloroform used in the method of Kouno et al.

(1995). The resin was then removed and the phosphate was recovered by shaking for 1 h in 20

ml of 0.5 M HCl. We added 25 and 40 mg P kg^{-1} depending on P contents as spike considering

the P adsorption in LP and HP soils, respectively (Brookes et al. 1982). The extracted P was

determined with the malachite green colorimetric method (Ohno and Zibilske 1991; Jeannotte et

al. 2004; Song et al. 2019). The extracts were diluted appropriately to avoid interference with

HCl (D'Angelo et al. 2001) and analyzed by absorbance measurement at 630 nm using a

microplate reader (iMark, Bio-Rad Laboratories, USA). The MBP was calculated from the

difference in extracted P concentrations between samples with and without fumigation divided

by 0.40 (Brookes et al. 1982).

        2.5. Soil P measurements

Phosphate within non-applied hexanol extracts for analysis of MBP was denoted as Resin

extractable P (Resin P). Resin P represent a part of labile P pools (Erinle et al. 2020) and avoid

fixation of P by soil colloid during the extraction in Andosols (Kouno et al. 1995). However, we

note that accurately assessing P availability in Andosols with high P sorption capacity using a

single chemical extraction method is difficult (Fixen and Grove 1990; Sugito and Shinano 2013).

1M HCl extractable P described by DeLuca et al. (2015) was also measured. It primarily

represents P mobilized by protons released by microbes or plants mainly from apatite and other

recalcitrant inorganic P forms, and also partially extracts Al/Fe-bound P and phosphate which is

weakly adsorbed to clay particles or present in inorganic precipitates (Kuo 1995; Pingree et al.

2017; Wuenscher et al. 2015; Li et al. 2019). This fraction was denoted as HCl-P. It was

measured by extracting 0.5 g of dry mass equivalent soil with 10 ml of 1 M HCl for 3 h at 200

rev min^{-1} at room temperature. After filtering, the inorganic P was determined by the malachite

green method at 630 nm using a microplate reader (iMark, Bio-Rad Laboratories, USA) as for

the MBP analysis.

2.6. Inorganic N measurements

Inorganic N (IN) was determined as the sum of NH_4-N and NO_3-N. Soil samples (5 g

dry mass equivalent) from the 2- and 20-day incubation periods were extracted with 25 ml of 1

M KCl by shaking for 30 minutes. The concentrations of NH_4-N and NO_3-N in the

filtrates were analyzed using a flow injection analyzer (AQLA-700, Aqualab Co., Ltd., Japan),

with NH_4^+-N measured by absorbance at 630 nm and NO_3^--N by absorbance at 540 nm as

described by Hamamoto and Uchida (2015).

    2.7. Substrate-derived C use efficiency and Priming effects

The calculations of glucose-derived C use efficiency (CUE) and priming effects (PEs) were

adapted from Mehnaz et al. (2019). While the CUE measured in this study strictly represents

glucose use efficiency, we used term "glucose-derived CUE" to maintain consistency with

similar previous studies (e.g., Cui et al. 2022; Karhu et al. 2022, Kalu et al. 2024, Xiao et al.

2021). Microbial glucose-derived CUE ($CUE_{glucose}$) of the added glucose was calculated using

the following equation:

$$CUE_{glucose} = MBC_{glucose} / (MBC_{glucose} + CO_2\text{-}C_{glucose}) \qquad (4)$$

The PEs were calculated as the increase in native soil organic C released after substrate

application also using the following equation (Mehnaz et al. 2019):

$$PEs = CO_2\text{-}C_{soil} - CO_2\text{-}C_{ctr} \qquad (5)$$

Where CO_2-C_{soil} and CO_2-C_{ctr} are the soil-derived CO_2-C respired from the soil with glucose

addition and the cumulative CO_2-C respired from the soil without glucose addition, respectively.

2.8. Soil extracellular enzyme activities

The potential activity of acid phosphomonoesterase (EC 3.1.3.2), P-acquiring extracellular

enzyme activity (EEA), was measured fluorometrically with methylumbelliferone (Bell et al.

2013). Briefly, soil samples (1.5 g dry mass equivalent) with 45 mL of deionized water were

shaken for 30 minutes (Deng et al. 2011; Li et al. 2025b). 800 µL of mixed soil slurry was taken

into a deep-well plate, then 200 µL of 1 mM substrate (4-Methylumbelliferyl phosphate) was

added and incubated at 35 °C for 1.5 hours. After incubation samples were centrifuged for 3

minutes at 2500 xg and moved to the 96-well black microplate. Fluorescence intensity was

measured using a Varioskan ALF multi-mode microplate reader (Thermo Scientific, USA) at

345 nm excitation and 450 nm emission. EEA was expressed as nmol $g^{-1} h^{-1}$.

2.9. Statistics

All statistical analyses were performed with R (version 4.2.2; R Core Team. 2023). We

conducted a two-way analysis of variance (ANOVA) including the interaction between soils and

substrates to assess their effects on the total CO_2-C, glucose-derived and soil-derived CO_2-C

using linear model (Bates et al. 2015). We also conducted three-way ANOVA including the

interaction between soils and substrates, soils and time, and substrates and time to assess their

effects total MBC, glucose-derived and soil-derived MBC, DOC, MBP, HCl-P, Resin P, and

EEA using linear model. Prior to the analysis, the datasets were rank-transformed to meet the

assumptions of ANOVA. Model was validated by examining residuals using the DHARMa

package (Hartig 2022). The data were further analyzed using Tukey's HSD test to find pairwise

differences between soil type and substrate combinations at $\alpha = 0.05$ using multcomp package

(Hothorn et al. 2008). The effects of soil P fertilization on each result, N addition on PEs was

determined by independent samples t-tests.

3. Results

3.1. CO_2 emission

The HP soil had a higher cumulative CO_2 emission (339.3 mg C kg^{-1} soil) than the LP soil

(308.6 mg C kg^{-1} soil) and there were significant differences when averaged across substrate

treatments during the incubation period (Fig. 1 and S1). Substrate additions significantly

stimulated the CO_2 emissions from both soils at day 1 immediately after the incubation (Fig. S1

and Table S3). This initial increase was primally due to the glucose-derived CO_2 emission,

which peaked at 161.2 mg C kg^{-1} in the HP soil and 154.5 mg C kg^{-1} in the LP soil across

substarte treatments (Fig. S1). Among substarte treatments, the glc treatments had significantly

higher cumulative CO_2 emissions than the glcN treatments in each soil (Fig. 1 and Table S3).

The cumulative SOM-derived CO_2 emission was higher in the HP soil than the LP soil

averaging across the substrate treatments (p<0.05, Fig. 1). Substrate additions increased

SOM-derived CO_2 emission in each soil type 6 days after incubation (p<0.001, Table S3). In the

HP soil, the glc treatment increased the SOM-derived CO_2 emissions, while in the LP soil, glc

and glcN treatments significantly increased the SOM-derived CO_2 emissions.

In contrast to the total and SOM-derived CO_2 emissions, the cumulative glucose-derived CO_2

emission showed no significant differences among soils and substrate treatments throughout the

incubation period (Fig. S2).

Fig. 1. Cumulative CO_2 emissions (mg C kg^{-1} soil) over 20-days incubation. Treatments include

soil without any amendment (Ctr), soil with ^{13}C-enriched glucose (glc), and soil with

^{13}C-enriched glucose and N substrates (glcN). Different color bars show cumulative CO_2

emissions (mean ± SD, n=4, but n=3 in LP Ctr due to an operational error during the

experiment) derived from added glucose (light green) and soil organic matter (grey). Uppercase

and lowercase letters indicate significant differences ($p < 0.05$) in total and SOM-derived CO_2-C

among treatments, respectively.

The PEs during incubation were lower in the HP soil than the LP across the treatments ($p < 0.05$

at 1, 6, 8 days of incubation, $p < 0.01$ and $p < 0.001$ at 2 days and 12, 20 days of incubation,

respectively) (Fig. 2, Tables S3 and S4). The glc addition had significantly higher PEs compared

to the glcN addition in both soils. The addition of glc and glcN in the LP soil showed positive

PEs. However, the effect of substrate treatments on PEs in the HP soil contrasted with the LP

soil; the glc addition showed a positive PEs, but glcN addition showed negative PEs. The total

primed CO_2-C as a % of added glucose C ranged from –0.71 to 12.88 % (Table S4).

Fig. 2. The priming effects (PEs) in HP and LP soils over 20 days. Treatments include soil with

^{13}C-enriched glucose (glc) and soil with ^{13}C-enriched glucose and N substrates (glcN). Green

and blue lines show the priming effect (mean ± SD, n = 4) in glc and glcN, respectively, for

each soil. * (p <0.05), ** (p <0.01) and blank (not significant) indicate significance level of N

addition effect for each soil separately.

3.2. Microbial biomass, soil chemical properties, and soil enzyme activity

The HP soil had significantly higher total MBC (MBC_{Total}) than the LP soil at 20 days of

incubation, whereas no significant differences were observed after 2 days of incubation (Fig. 3,

Table S5 and S6). Substrate additions did not affect MBC_{Total} in either soil at either 2 or 20 days

of incubation (Table S5 and S6). For the SOM-derived microbial biomass C (MBC_{SOM}),

substrate additions significantly decreased MBC_{SOM} in the HP soil compared to the control at 2

days of incubation (Fig. 3, Table S5 and S6). In contrast, the LP soil showed no differences in

MBC_{SOM} among substrate treatments at 2 days of incubation. Glucose-derived microbial

biomass C (MBC_{Glc}) showed no significant differences among soils and substrate treatments at 2

days of incubation (Fig. 3, Table S5 and S6). However, at 20 days of incubation, MBC_{Glc} was

higher in the HP soil than in the LP soil. In both soils, MBC_{Glc} was significantly lower at 20

days of incubation compared to 2 days. The P-acquiring enzyme activity was higher in HP soils

than in LP soils (Table S5 and S6). The addition of GlcN significantly increased the P acquire

enzyme activity in LP soils at 2 and 20 days of incubation.

Fig. 3. Microbial biomass C (MBC) (mg C kg^{-1} soil) in 2 and 20-days incubation. Treatments

include soil without any amendment (Control), soil with ^{13}C-enriched glucose (glc), and soil

with ^{13}C-enriched glucose and N substrates (glcN). Different color bars show cumulative MBC

(mean ± SD, n =4, but n =3 in LP Ctr due to an operational error during the experiment) derived

from added glucose (light green) and soil organic matter (grey). Uppercase and lowercase letters

indicate significant differences (p <0.05) in glucose-derived MBC and SOM-derived MBC

among substrate treatments and soils at each sampling time, respectively.

The HP soil had significantly higher MBP and a lower log ratio of MBC:MBP compared to

the LP soil at 2 and 20 days of incubation (Table S5 and S6). However, substrate additions had

no significant effect on MBP in the both soils during the incubation period.

Soil pH, Resin P, and HCl-P were consistently higher in the HP soils than in the LP soil during

the incubation period (Table S6 and S7). Resin P increased from 0 to 2 days of incubation in HP

soil ($p <0.001$), while there was no significant difference in LP soil ($p =0.065$). Then Resin P

decreased from 2 to 20 days in LP soil ($p <0.001$), whereas no significant difference was found

in HP soil ($p =0.41$).

        3.3. Microbial C use efficiency

Glucose-derived microbial C use efficiency (CUE) showed no significant differences between

the two soils or substrate additions at 2 days after incubation (Table 2). However, at 20 days

after incubation, glucose-derived CUE in the HP soil was higher than in the LP soil ($p <0.05$).

The HP soil with glc showed the highest glucose use efficiency, while the LP soil with glc had

the lowest glucose use efficiency at 20 days after incubation.

Table 2. C use efficiency (CUE) in the HP and LP soils at 2 and 20 days after incubation (mean (SD)). Different letters indicate significant

differences (p <0.05) among treatments for each soil type.

		CUE			
Soil	Substrate	2-day		20-day	
HP	glc	0.65 (0.025)		0.44 (0.015)	a
	glcN	0.61 (0.038)		0.42 (0.009)	ab
LP	glc	0.62 (0.035)		0.39 (0.012)	b
	glcN	0.63 (0.012)		0.43 (0.025)	a
ANOVA P values					
Soil		n.s		<0.05	
Substrate		n.s		n.s	
Soil × Sub		n.s		<0.01	

4. Discussion

Our results show that contrasting long-term P fertilization affected on the microbial

respirations. The high-P soils had a significantly higher cumulative CO_2 emissions in the

control compared to the low-P soils, and this trend also remained consistent regardless of the

glc or glcN addition. This finding suggests that P input potentially enhanced the microbial

activities due to increasing available P and could also promote the desorption of the organic

C, thereby increasing available C pools (Kaiser and Zech 1999; Miyazawa et al. 2013; Scott

et al. 2015, Spohn and Schleuss 2019, Li et al. 2020; Spohn et al. 2022).

PEs were also influenced by P fertilization (Fig. 2). Notably, N-mining was suggested in

both soils by the higher soil-derived CO emissions with glucose-only addition compared to

glucose+N addition (Li et al. 2017). However, the overall response of PEs to substrate

addition clearly contrasted with soil P status. Specifically, when N was not limited (i.e., in

the glucose+N treatment), negative PEs were found in the high-P soil, whereas positive PEs

were observed in the low-P soil. Combined with the significantly increased phosphatase

activity in the low-P soil (Table S5), this result indicates that the positive PEs are associated

with SOM degradation driven by P-mining, supporting our first hypothesis. In addition, the

low-P soils had consistently lower Resin P and higher MBC:P ratios compared to the high-P

soils (Table 1, S6, and S7). Under nutrient-imbalanced conditions, microbes invest abundant

elements into decomposing SOM to obtain limited nutrients, thereby maximizing their

productivity (Allison and Vitousek 2005; Mooshammer et al. 2014). Moreover, MBP levels

in the low-P soil remained stable throughout the incubation period, possibly because positive

priming induced P-mining and maintained microbial P (Spohn 2016).

In agreement with our second hypothesis, the P fertilization was also associated with CUE.

After 20 days of incubation, the high-P soils showed significantly higher glucose-derived

CUE than the low-P soils (Figs. 1, 3 and Table 2). These long-term CUE measurements

(>48h) suggest the efficiency of C stabilization through the recycling of the microbial

necromass and exudates (Geyer et al. 2016; Kalu et al. 2024), whereas the short-term

incubation CUE measurements (<48 h) may reflect the substrate utilization for gross

production (Brant et al. 2006; Geyer et al. 2016). After 20 days of incubation, in the low-P

soil, microbes allocated more energy to catabolic pathways to acquire P, which in turn

reduced C available for growth and resulted lower glucose-derived CUE than in the high-P

soil (Bernard et al. 2022). These findings suggest that differences in P fertilization between

the two soils are a key factor explaining the variation in CUE and PEs.

The two soils used in this study significantly differed in pH and exchangeable acidity due

to long-term P fertilization, both of which are important factors influencing microbial activity.

The lower pH throughout the incubation period and higher exchangeable Al in the low-P

soils compared to the high-P soils likely suppressed the microbial respirations and activities

(Table1 and S7). In general, higher PEs occur in soils with neutral pH compared to the soils

with acidic pH because the microbial activity, the community structure, and the enzyme

synthesis are more pronounced (Blagodatskaya and kuzyakov 2008; Blagodatskaya and

Anderson 1998). However, previous study suggested that greater positive PEs observed in

strongly acidic soils are associated with the size of labile C pools and net increase in MBC

(Aye et al. 2018). In this study, although MBC showed no significant differences among soils,

higher DOC concentrations were shown in the low-P soils across the treatments, which

corresponded with the greater primed CO_2 emissions observed (Fig. 3 and Table S5). These

findings indicated a potential role of low pH in affecting DOC pools and microbial resource

accessibility. In contrast, in the low-P soils, MBP showed significant lower than the high-P

soils, and as discussed above, the higher MBC:P ratio likely reflects SOM degradation

induced by P-mining.

In addition to PEs, soil pH is also important for understanding the CUE results; however,

after 2 days of incubation, glucose-derived CUE showed no significant differences among

soils (Table 2). It is possible that the effect of P fertilization on microbial assimilation of

substrates (i.e., short-term CUE) after 2 days of incubation was offset by: i) availability of

organic C; ii) the soil pH, both specific to non-allopnanic Andosols. Soil C content is one of

the regulating factors of microbial assimilation (Kamble and Bååth 2014); Andosols have a

large amount of organic C content due to Al-soil organic matter complexes (Takahashi and

Dahlgren 2016). Moreover, it has been suggested that non-allophanic Andosols have higher

organic C contents than other Andosols (Nanzyo et al. 1993). Previous studies showed that

microbial growth and CUE are strongly related to C availability rather than N and P

availability (Wang et al. 2021; Morris et al. 2021; Jiang et al. 2025). In this study, we found

that the high-P and the low-P soils contain approximately 10% TC, this high C content may

have influenced glucose-derived CUE after 2 days of incubation. Moreover, soil pH can alter

the composition of microbial communities, and consequently affect CUE (Manzoni et al.

2012; Zhen et al. 2019). Rousk et al. (2011) reported that fungal growth is suppressed at

$pH_{(H2O)} > 5.5$, contrasted with bacterial growth limited at $pH_{(H2O)} < 5.5$. Keiblinger et al.

(2010) reported that fungi induced higher CUE than bacteria, and fungal CUE was affected

by C availability and bacterial CUE limited by N and P availability, respectively. Thus, it is

possible that the microbial community composition was adapted to low pH conditions and

the high C content of the soil and these factors may have contributed to the marginal effect

of P fertilization on glucose-derived CUE in the non-allophanic Andosol.

Jones et al. (2019) reported that low pH and associated exchangeable Al decrease CUE

through metabolic shifts and greater C diversion into maintenance and catabolic processes,

rather than through a reduction in the ability to utilize labile C. In our study, the amount of

exchangeable Al was 0 meq kg^{-1} in the high-P soil and 21.30 meq kg^{-1} in the low-P soil,

respectively. It is possible that the lower glucose-derived CUE in the low-P soils after 20

days of incubation, when applied C substrate was largely consumed, was due to the toxicity

of exchangeable Al, as reflected in the differing amounts of Al^{3+}, in addition to low soil P

level. Although Jones et al. (2019) focused on short-term effects (2-72 h), our study

suggested that these factors possibly affect CUE over longer time scales than several hours.

The absence of a difference in CUE between the high-P and the low-P soils on 2 days of

incubation, however, may have been due to the influence of organic C availability, as

mentioned in the previous paragraph. Therefore, the low CUE observed in the low-P soils is

likely attributable to C investment for P acquisition and the response to acid and Al stresses.

Our results showed lower primed CO_2-C of added glucose C in non-allophanic Andosols

compared to previous studies. In the current, the values ranged from –1 to 13 % in the

non-allophanic Andosols over 20 days incubation compared to values of 28 to 420 % for

other soil types (Table S4) (Shahbaz et al. 2018; Mehnaz et al. 2019; Sawada et al. 2023).

This comparison is valid as the values were calculated from data between 18 to 42 days of

the entire incubation period. In contrast, the size of the MBC pool was comparable or even

larger than the size in these studies (100-1190 mg C kg^{-1} soil), suggesting a higher total

microbial biomass including potentially active microbes. This lower PEs in the

non-allophanic Andosols were possibly due to the abundance of Al-organic matter

complexes. The stability of SOM depends on Al-organic matter complexes and soil acidity

(Miyazawa et al. 2013). These complexes promote the soil aggregates formation and are also

suggested to physically suppress the microbial SOM decomposition (Matsui et al. 2016).

Thus, it is suggested that the low PEs reflect SOM stability in non-allophanic Andosols.

Our findings indicate that maintaining C stocks in non-allophanic Andosols depends not

only on adequate long-term P fertilization but also on appropriate regulation of soil acidity.

However, we acknowledge that our experiment did not directly represent the temporal Al

toxicity effects on CUE, and we could not statistically separate the individual contributions

of these overlapping factors to the observed responses. Future studies need to demonstrate

these causal relationships by employing controlled manipulations of soil pH and investigate

the relationships between Al bioavailability and C dynamics at different soil P levels.

5. Conclusion

This study demonstrated that substrate-induced SOM decomposition in non-allophanic

Andosols is strongly influenced by the long-term P fertilization. The addition of glucose with

N substrate promoted positive PEs under low P conditions, while negative PEs occurred

under high P conditions. Throughout the incubation, PEs in the low-P soils were greater than

in the high-P soils. Additionally, in the low-P soils, substrate additions did not affect MBP

and the ratio of MBC:P throughout the incubation period. These results suggest that a lack of

P affected the impact on SOM decomposition due to the maintaining microbial P. Greater

CUE was observed in the high-P soils than in the low-P soils after 20 days of incubation,

with no difference after 2 days of incubation. These results indicate that under P-limited

conditions combined with soil acidity and exchangeable Al, the microbial allocation of C for

P acquisition regulates substrate-induced microbial assimilation and SOM decomposition.

Future research needs to elucidate the interactive effects of P availability, soil pH, and

exchangeable Al on microbial C dynamics in non-allophanic Andosols.

Author's Contributions

W.K., T.M., and T.H. planned and designed the research. W.K. and T.H. conducted the

incubation experiments. W.K. and T.H. analyzed the data, wrote the main manuscript, and

prepared the figures and tables. All authors reviewed the manuscript.

Acknowledgments

This work was financially supported by the JSPS Grant-in-Aid for JSPS Fellows

(23K14056), and the Yamaguchi Educational and Scholarship Foundation. This study was

supported by Support system for young researchers to use research equipment in Tohoku

University.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

Allison, S. D., Vitousek, P. M. (2005). Responses of extracellular enzymes to simple and

complex nutrient inputs. Soil Biol Biochem, 37(5), 937–944.

Aye, N. S., Butterly, C. R., Sale, P. W., & Tang, C. (2018). Interactive effects of initial pH

and nitrogen status on soil organic carbon priming by glucose and lignocellulose. Soil Biol

Biochem, 123, 33-44.

Bates, D., Mächler, M., Bolker, B. M., Walker, S. C. (2015). Fitting linear mixed-effects

models using lme4. J Stat Softw, 67(1), 1–48.

Beck, T., Joergensen, R. G., Kandeler, E., Makeschin, F., Nuss, E., Oberholzer, H. R., Scheu,

S. (1997). An inter-laboratory comparison of ten different ways of measuring soil microbial

biomass C. Soil Biol Biochem, 29(7), 1023–1032.

Bell, C. W., Fricks, B. E., Rocca, J. D., Steinweg, J. M., McMahon, S. K., Wallenstein, M. D.

(2013). High-throughput fluorometric measurement of potential soil extracellular enzyme

activities. JoVE, 81, e50961.

Bernard, L., Basile-Doelsch, I., Derrien, D., Fanin, N., Fontaine, S., Guenet, B., Karimi, B.,

Marsden, C., Maron, P. A. (2022). Advancing the mechanistic understanding of the priming

effect on soil organic matter mineralisation. Funct Ecol, 36(6), 1355–1377.

Blagodatskaya, E. V., and Anderson, T. H. (1998). Interactive effects of pH and substrate

quality on the fungal-to-bacterial ratio and qCO2 of microbial communities in forest soils.

Soil Biol Biochem, 30(10-11), 1269-1274.

Blagodatskaya, E., Kuzyakov, Y. (2008). Mechanisms of real and apparent priming effects

and their dependence on soil microbial biomass and community structure: Critical review.

Biol Fertil Soil, 45(2), 115–131.

Brant, J. B., Sulzman, E. W., Myrold, D. D. (2006). Microbial community utilization of

added carbon substrates in response to long-term carbon input manipulation. Soil Biol

Biochem, 38(8), 2219–2232.

Brookes, P. C., Powlson, D. S., Jenkinson, D. S. (1982). Measurement of microbial biomass

phosphorus in soil. Soil Biol Biochem, 14(4), 319–330.

Cleveland, C. C., Liptzin, D. (2007). C:N:P stoichiometry in soil: Is there a "Redfield ratio"

for the microbial biomass? Biogeochemistry, 85(3), 235–252.

Craine, J. M., Morrow, C., Fierer, N. (2007). Microbial nitrogen limitation increases

decomposition. Ecology, 88(8), 2105–2113.

Cui, Y., Moorhead, D. L., Wang, X., Xu, M., Wang, X., Wei, X., Zhu, Z., Ge, T., Peng, S.,

Zhu, B., Zhang, X., Fang, L. (2022). Decreasing microbial phosphorus limitation increases

soil carbon release. Geoderma, 419, 115868.

D'Angelo, E., Crutchfield, J., Vandiviere, M. (2001). Rapid, sensitive, microscale

determination of phosphate in water and soil. J Environ Qual, 30, 2206–2209.

DeLuca, T. H., Glanville, H. C., Harris, M., Emmett, B. A., Pingree, M. R. A., de Sosa, L. L.,

Morenà, C., Jones, D. L. (2015). A novel biologically-based approach to evaluating soil

phosphorus availability across complex landscapes. Soil Biol Biochem, 88, 110–119.

Deng, S., Kang, H., Freeman, C. (2011). Microplate fluorimetric assay of soil

enzymes. Methods of soil enzymology, 9, 311-318.

Erinle, K. O., Doolette, A., Marschner, P. (2020). Changes in phosphorus pools in the

detritusphere induced by removal of P or switch of residues with low and high C/P ratio. Biol

Fertil Soils, 56(1), 1–10.

Fixen, P. E., Grove, J. H. (1990). Testing soils for phosphorus. Soil testing and plant analysis,

3, 141-180.

Geyer, K. M., Kyker-Snowman, E., Grandy, A. S., Frey, S. D. (2016). Microbial carbon use

efficiency: accounting for population, community, and ecosystem-scale controls over the fate

of metabolized organic matter. Biogeochemistry, 127(2–3), 173–188.

Hamamoto, T., Uchida, Y. (2015). Sodium contents in dairy cow urine and soil aggregate

sizes influence the amount of nitrogen lost from soil. Appl Environ Soil Sci, 2015(1),

275985.

Hartig, F. (2016). DHARMa: residual diagnostics for hierarchical (multi-level/mixed)

regression models. CRAN: Contributed Packages.

Hessen, D. O., Ågren, G. I., Anderson, T. R., Elser, J. J., De Ruiter, P. C. (2004). Carbon

sequestration in ecosystems: the role of stoichiometry. Ecology, 85(5), 1179-1192.

Hothorn, T., Hornik, K., Van De Wiel, M. A. Zeileis, A. (2008). Implementing a class of

permutation tests: the coin package. J Stat Softw, 28, 1-23.

Iamarino, M., Terribile, F. (2008). The importance of andic soils in mountain ecosystems: a

pedological investigation in Italy. Eur J Soil Sci, 59(6), 1284-1292.

Islam, M. R., Singh, B., Dijkstra, F. A. (2023). Microbial carbon use efficiency of glucose

varies with soil clay content: A meta-analysis. Appl Soil Ecol, 181, 104636.

Ito, T., Kikawa, N., Saigusa, M. (2011). Phosphorus sorption and bioavailability in

allophanic and non-allophanic Andosols. Pedologist, 55(2), 84–88 (in Japanese).

IUSS Working Group WRB. (2022). World reference base for soil resources. International

soil classification system for naming soils and creating legends for soil maps (4th ed.). IUSS.

Jackson, R. B., Lajtha, K., Crow, S. E., Hugelius, G., Kramer, M. G., Piñeiro, G. (2017). The

ecology of soil carbon: pools, vulnerabilities, and biotic and abiotic controls. Annu Rev Ecol

Evol Syst, 48, 419–445.

Jeannotte, R., Sommerville, D. W., Hamel, C., Whalen, J. K. (2004). A microplate assay to

measure soil microbial biomass phosphorus. Biol Fertil Soil, 40(3), 201–205.

Jiang, M., Crous, K. Y., Carrillo, Y., Macdonald, C. A., Anderson, I. C., Boer, M. M., Farrell,

M., Gherlenda, A. N., Castañeda-Gómez, L., Hasegawa, S., et al. (2024). Microbial

competition for phosphorus limits the CO_2 response of a mature forest. Nature, 630(8017),

660–665.

Jiang, Q., Yan, L., Wang, H., Dai, X., Meng, S., Fu, X., Yang, F., Ma, Z., Shi, W., Lv, W.,

Wang, Y., Gao, D. (2025). Nutrient additions have minor impact on microbial carbon use

efficiency. Agric ecosyst environ., 385, 109582.

Jones, D. L., Cooledge, E. C., Hoyle, F. C., Griffiths, R. I., Murphy, D. v. (2019). pH and

exchangeable aluminum are major regulators of microbial energy flow and carbon use

efficiency in soil microbial communities. Soil Biol Biochem, 138, 107584.

Jones, D. L., Hill, P. W., Smith, A. R., Farrell, M., Ge, T., Banning, N. C., Murphy, D. v.

(2018). Role of substrate supply on microbial carbon use efficiency and its role in

interpreting soil microbial community-level physiological profiles (CLPP). Soil Biol

Biochem, 123, 1–6.

Kaiser, K., Zech, W. (1999). Release of natural organic matter sorbed to oxides and a subsoil.

Soil Sci Soc Am J, 63(5), 1157–1166.

Kallenbach, C. M., Frey, S. D., Grandy, A. S. (2016). Direct evidence for microbial-derived

soil organic matter formation and its ecophysiological controls. Nat Commun, 7(1), 1–10.

Kalu, S., Seppänen, A., Mganga, K. Z., Sietiö, O. M., Glaser, B., Karhu, K. (2024). Biochar

reduced the mineralization of native and added soil organic carbon: evidence of negative

priming and enhanced microbial carbon use efficiency. Biochar, 6(1), 1–14.

Kamble, P. N., Bååth, E. (2014). Induced N-limitation of bacterial growth in soil: Effect of

carbon loading and N status in soil. Soil Biol Biochem, 74, 11–20.

Karhu, K., Alaei, S., Li, J., Merilä, P., Ostonen, I., Bengtson, P. (2022). Microbial carbon use

efficiency and priming of soil organic matter mineralization by glucose additions in boreal

forest soils with different C: N ratios. Soil Biol Biochem, 167, 108615.

Keiblinger, K. M., Hall, E. K., Wanek, W., Szukics, U., Hämmerle, I., Ellersdorfer, G., Böck,

S., Strauss, J., Sterflinger, K., Richter, A., Zechmeister-Boltenstern, S. (2010). The effect of

resource quantity and resource stoichiometry on microbial carbon-use-efficiency. FEMS

Microbiol Ecol, 73(3), 430–440.

Kouno, K., Tuchiya, Y., Ando, T. (1995). Measurement of soil microbial biomass phosphorus

by an anion exchange membrane method. Soil Biol Biochem, 27(10), 1353–1357.

Kuo, S. (2018). Phosphorus. Methods of Soil Analysis, Part 3: Chemical Methods, 869–919.

Li, C., Wade, J., Vollbracht, K., Hooper, D. G., Wills, S. A., Margenot, A. J. (2025b). Do

chromogenic assays of soil enzyme activities need buffers? More disadvantages than

advantages of modified universal buffer in the para-nitrophenyl-based assay of

phosphomonoesterase and β-glucosidase activities. Soil Biol Biochem, 202, 109704.

Li, J., Hoang, K. T. K., Hassan, N., Marschner, P. (2019). Vermicompost influences soil P

pools and available N—effect of placement and combination with inorganic fertiliser. Soil

Sci Plant Nutr, 19(4), 900-905.

Li, J., Liu, Z. F., Jin, M. K., Zhang, W., Lambers, H., Hui, D., Liang, C., Zhang, J., Wu, D.,

Sardans, J., Peñuelas, J., Petticord, D. F., Frey, D. W., Zhu, Y. (2023). Microbial controls

over soil priming effects under chronic nitrogen and phosphorus additions in subtropical

forests. ISME J, 17(12), 2160–2168.

Li, Q., Tian, Y., Zhang, X., Xu, X., Wang, H., Kuzyakov, Y. (2017). Labile carbon and

nitrogen additions affect soil organic matter decomposition more strongly than temperature.

Appl Soil Ecol, 114, 152–160.

Li, S., Cui, Y., Moorhead, D. L., Dijkstra, F. A., Sun, L., Xia, Z., Gao, Y., Ma, Q., Yu, W.

(2025a). Phosphorus limitation regulates the responses of microbial carbon metabolism to

long-term combined additions of nitrogen and phosphorus in a cropland. Soil Biol Biochem,

200, 109614.

Li, Y., Wang, T., Camps-Arbestain, M., Suárez-Abelenda, M., Whitby, C. P. (2020). Lime

and/or phosphate application affects the stability of soil organic carbon: evidence from

changes in quantity and chemistry of the soil water-extractable organic matter. Environ Sci

Technol, 54(21), 13908–13916.

Lieberman, H. P., von Sperber, C., Kallenbach, C. M. (2025). Soil phosphorus dynamics are

an overlooked but dominant control on mineral‐associated organic matter. Glob Chang

Biol, 31(7), e70307.

Manzoni, S., Taylor, P., Richter, A., Porporato, A., Ågren, G. I. (2012). Environmental and

stoichiometric controls on microbial carbon-use efficiency in soils. New Phytol, 196(1),

79–91.

Matsui, T., Takahashi, T., Kanno, H., Nanzyo, M. (2016). Inhibitory effect of amorphous

aluminum hydroxide gels on organic carbon decomposition in non-allophanic Andosols.

Pedologist, 60(1), 4-13.

Matus, F., Salazar, O., Aburto, F., Zamorano, D., Nájera, F., Jovanović, R., Guerra, C.,

Reyes-Rojas, L., Seguel, O., Pfeiffer, M., Dörner, J., Valle, S., Radic-Schilling, S., Duarte, E.

(2024). Perspective of soil carbon sequestration in Chilean volcanic soils. Npj Mater Sustain,

2(1), 1–7.

Maynard, D. S., Crowther, T. W., Bradford, M. A. (2017). Fungal interactions reduce carbon

use efficiency. Ecol Lett, 20(8), 1034–1042.

Mehnaz, K. R., Corneo, P. E., Keitel, C., Dijkstra, F. A. (2019). Carbon and phosphorus

addition effects on microbial carbon use efficiency, soil organic matter priming, gross

nitrogen mineralization and nitrous oxide emission from soil. Soil Biol Biochem, 134,

175–186.

Miyazawa, M., Takahashi, T., Sato, T., Kanno, H., Nanzyo, M. (2013). Factors controlling

accumulation and decomposition of organic carbon in humus horizons of Andosols: A case

study for distinctive non-allophanic Andosols in northeastern Japan. Biol Fertil Soil, 49(7),

929–938.

Mooshammer, M., Wanek, W., Zechmeister-Boltenstern, S., Richter, A. (2014).

Stoichiometric imbalances between terrestrial decomposer communities and their resources:

mechanisms and implications of microbial adaptations to their resources. Front Microbiol, 5,

22.

Mori, T. (2023). P fertilization experiments require reinterpretation: Abiotic elevation of

available C and N could influence microbial processes in soil. Appl Soil Ecol, 189, 104899.

Morris, K. A., Richter, A., Migliavacca, M., Schrumpf, M. (2022). Growth of soil microbes

is not limited by the availability of nitrogen and phosphorus in a Mediterranean oak-savanna.

Soil Biol Biochem, 169, 108680.

Nanzyo, M. (2002). Unique properties of volcanic ash soils. Glob Environ Res, 6(2), 99-112.

Nanzyo, M., Dahlgren, R., Shoji, S. (1993). Chemical characteristics of volcanic ash soils.

Developments in soil science, 21, 145-187.

Nottingham, A. T., Turner, B. L., Stott, A. W., Tanner, E. V. J. (2015). Nitrogen and

phosphorus constrain labile and stable carbon turnover in lowland tropical forest soils. Soil

Biol Biochem, 80, 26–33.

Obara, H., Ohkura, T., Takata, Y., Kohyama, K., Maejima, Y., Hamazaki, T. (2011).

Comprehensive soil classification system of Japan first approximation. Nogyo Kankyo

Gijutsu Kenkyusho Hokoku Bull Natl Inst Agro-Environ Sci, 29, 1–73.

Ohno, T., Zibilske, L. M. (1991). Determination of low concentrations of phosphorus in soil

extracts using malachite green. Soil Sci Soc Am J, 55(3), 892–895.

Pingree, M. R., Makoto, K., DeLuca, T. H. (2017). Interactive effects of charcoal and

earthworm activity increase bioavailable phosphorus in sub-boreal forest soils. Biol Fertil

Soil, 53(8), 873-884.

Qiao, N. A., Schaefer, D., Blagodatskaya, E., Zou, X., Xu, X., Kuzyakov, Y. (2014). Labile

carbon retention compensates for CO_2 released by priming in forest soils. Global Change

Biol, 20(6), 1943-1954.

Qin, W., Feng, J., Zhang, Q., Yuan, X., Ren, F., Zhou, H., Zhu, B. (2024). Different

responses of priming effects in long-term nitrogen-and phosphorus-fertilized soils to

exogenous carbon inputs. Plant and Soil, 500(1), 647–663.

R Core Team. (2023). R: A Language and Environment for Statistical Computing. Vienna,

Austria: R Foundation for Statistical Computing.

Rousk, J., Brookes, P. C., Bååth, E. (2011). Fungal and bacterial growth responses to N

fertilization and pH in the 150-year 'Park Grass' UK grassland experiment. FEMS Microbiol

Ecol, 76(1), 89–99.

Russell, J. B., Cook, G. M. (1995). Energetics of bacterial growth: balance of anabolic and

catabolic reactions. Microbiol Rev, 59(1), 48–62.

Saigusa, M., Matsuyama, N. (1998). Distribution of allophanic Andosols and non-allophanic

Andosols in Japan. Tohoku J. Agr. Res., 48, 75–83.

Sawada, K., Kunito, T., Watanabe, T., Kitagawa, N., Lyu, H., Nguyen, H. L., Toyota, K.,

Funakawa, S. (2023). Positive priming effects through microbial P-mining in tropical forest

soils under N2-fixing trees. Soil Sci. Plant Nutr, 69(4), 209–214.

Schimel, J. P., Weintraub, M. N. (2003). The implications of exoenzyme activity on

microbial carbon and nitrogen limitation in soil: a theoretical model. Soil Biol Biochem,

35(4), 549–563.

Schroeder, J., Dămătîrcă, C., Bölscher, T., Chenu, C., Elsgaard, L., Tebbe, C. C., Skadell, L.,

Poeplau, C. (2024). Liming effects on microbial carbon use efficiency and its potential

consequences for soil organic carbon stocks. Soil Biol Biochem, 191, 109342.

Scott, J. T., Lambie, S. M., Stevenson, B. A., Schipper, L. A., Parfitt, R. L., McGill, A. C.

(2015). Carbon and nitrogen leaching under high and low phosphate fertility pasture with

increasing nitrogen inputs. Agric ecosyst environ, 202, 139–147.

Shahbaz, M., Kumar, A., Kuzyakov, Y., Börjesson, G., Blagodatskaya, E. (2018). Priming

effects induced by glucose and decaying plant residues on SOM decomposition: a

three-source 13C/14C partitioning study. Soil Biol Biochem, 121, 138–146.

Shoji, S., Suzuki, Y., Saigusa, M. (1987). Clay Mineralogical and Chemical Properties of

Nonallophanic Andepts (Andisols) from Oregon, USA. Soil Sci Soc Am J, 51(4), 986–990.

Song, S. hui, Wen, Y. jie, Zhang, J. yao, Wang, H. (2019). Rapid spectrophotometric

measurement with a microplate reader for determining phosphorus in NaHCO3 soil extracts.

Microchem J, 146, 210–213.

Spohn, M. (2016). Element cycling as driven by stoichiometric homeostasis of soil

microorganisms. Basic Appl Ecol, 17(6), 471–478.

Spohn, M. (2024). Preferential adsorption of nitrogen-and phosphorus-containing organic

compounds to minerals in soils: A review. Soil Biol Biochem, 194, 109428.

Spohn, M., Diáková, K., Aburto, F., Doetterl, S. Borovec, J. (2022). Sorption and desorption

of organic matter in soils as affected by phosphate. Geoderma, 405, 115377.

Spohn, M., Schleuss, P. M. (2019). Addition of inorganic phosphorus to soil leads to

desorption of organic compounds and thus to increased soil respiration. Soil Biol Biochem,

130, 220–226.

Sugito, T., Shinano, T. (2013). Estimating phosphorus availability in andosols using soil

biomass phosphorus as an indicator. JARQ, 47(4), 347–351.

Takahashi, T. (2020). The diversity of volcanic soils: focusing on the function of

aluminum–humus complexes. Soil Sci Plant Nutr, 66(5), 666–672.

Takahashi, T., Dahlgren, R. A. (2016). Nature, properties and function of aluminum–humus

complexes in volcanic soils. Geoderma, 263, 110–121.

Tao, F., Huang, Y., Hungate, B. A., Manzoni, S., Frey, S. D., Schmidt, M. W. I., Reichstein,

M., Carvalhais, N., Ciais, P., Jiang, L., Lehmann, J., Wang, Y., Houlton, B. Z., Ahrens, B.,

Mishra, U., Hugelius, G., Hocking, T. D., Lu, X., Shi, Z., Viatkin, K., Vargas, R., Yigini, Y.,

Omuto, C., Malik, A. A., Peralta, G., Cuevas-Corona, R., Di Paolo, L. E., Luotto, I., Liao, C.,

Liang, Y., Saynes, V. S., Huang, X., Luo, Y. (2023). Microbial carbon use efficiency

promotes global soil carbon storage. Nature, 618, 981–985.

Tempest, D. W., Neijssel, O. M. (1992). Physiological and energetic aspects of bacterial

metabolite overproduction. FEMS Microbiol Lett, 100(1-3), 169–76.

Wang, C., Qu, L., Yang, L., Liu, D., Morrissey, E., Miao, R., Liu, Z., Wang, Q., Fang, Y., Bai,

E. (2021). Large-scale importance of microbial carbon use efficiency and necromass to soil

organic carbon. Global Change Biol, 27(10), 2039–2048.

Wang, H., Ye, W., He, W., Guo, Z., Hu, G., Lou, Y., Yang, Q., Yang, Z., Sun, Y., Pan, H.,

Zhuge, Y. (2025). Phosphorus addition increases soil organic matter priming in a coastal

saline soil. Soil Biol Biochem, 208, 109862.

Wang, Q., Wang, S., He, T., Liu, L., Wu, J. (2014). Response of organic carbon

mineralization and microbial community to leaf litter and nutrient additions in subtropical

forest soils. Soil Biol Biochem, 71, 13–20.

Widdig, M., Schleuss, P. M., Biederman, L. A., Borer, E. T., Crawley, M. J., Kirkman, K. P.,

Seabloom, E. W., Wragg, P. D., Spohn, M. (2020). Microbial carbon use efficiency in

grassland soils subjected to nitrogen and phosphorus additions. Soil Biol Biochem, 146,

107815.

Wieder, W. R., Bonan, G. B., Allison, S. D. (2013). Global soil carbon projections are

improved by modelling microbial processes. Nat Clim Change, 3(10), 909–912.

Wuenscher, R., Unterfrauner, H., Peticzka, R., Zehetner, F. (2015). A comparison of 14 soil

phosphorus extraction methods applied to 50 agricultural soils from Central Europe, Plant

Soil Environ, 61, 86–96.

Xiao, Q., Huang, Y., Wu, L., Tian, Y., Wang, Q., Wang, B., Xu, M., Zhang, W. (2021).

Long-term manuring increases microbial carbon use efficiency and mitigates priming effect

via alleviated soil acidification and resource limitation. Biol Fertil Soils, 57(7), 925–934.

Xu, S., Delgado-Baquerizo, M., Kuzyakov, Y., Wu, Y., Liu, L., Yang, Y., Li, Y., Yu, Y., Zhu,

B., Yao, H. (2024). Positive soil priming effects are the rule at a global scale. Global Change

Biol, 30(9).

Zeng, Q., Zhang, Q., Mei, K., Feng, J., Yuan, X., Liu, Y., Xu, M., Sun, H., Zhu, B., Chen, Y.

(2025). Increasing phosphorus availability reduces priming effect by facilitating microbial

carbon use efficiency in a subtropical forest soil. Biol Fertil Soil, 61(5), 909–923.

Zhai, Z., Luo, M., Yang, Y., Liu, Y., Chen, X., Zhang, C., Huang, J., Chen, J. (2022).

Trade-off between microbial carbon use efficiency and microbial phosphorus limitation

under salinization in a tidal wetland. CATENA, 209, 105809.

Zheng, Q., Hu, Y., Zhang, S., Noll, L., Böckle, T., Richter, A., Wanek, W. (2019). Growth

explains microbial carbon use efficiency across soils differing in land use and geology. Soil

Biol Biochem, 128, 45–55.

Zibilske, L. M. (1994). Carbon mineralization. Methods of soil analysis: Part 2

microbiological and biochemical properties, 5, 835–863.