The absence of the *ompA* and type VI secretion system in the

successful lineage of *Acinetobacter baumannii* ST19

Authors: Adam Valcek[1, 2*], Diana Isabela Costescu Strachinaru[3], Kristina

Nesporova[4], Patrick Soentjens[3,5] Anke Stoefs[6,7], Charles Van der Henst[1, 2*]

Affiliations:

1. Microbial Resistance and Drug Discovery, VIB-VUB Center for Structural Biology,

VIB, Flanders Institute for Biotechnology, Brussels, Belgium

2. Structural Biology Brussels, Vrije Universiteit Brussel (VUB), Brussels, Belgium

3. Center for Infectious Diseases, Queen Astrid Military Hospital, Brussels, Belgium

4. Microbial Systems Cell Biology Group, Department of Biology, KU Leuven,

Leuven, Belgium

5. Department of Clinical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

6. Department of Microbiology, Queen Astrid Military Hospital, Brussels, Belgium

7. Department of Microbiology, Universitair Ziekenhuis Brussels, Brussels Belgium

* These authors contributed equally to this work and share the last authorship.

Corresponding authors: Adam Valcek (*adam.valcek@vub.be),* Charles Van der

Henst (*charles.vanderhenst@vub.vib.be*)

Abstract

Acinetobacter baumannii is a critical, opportunistic pathogen due to its multi- to

pandrug-resistance including resistance to last-resort antibiotics such as

carbapenems. While *A. baumannii* can acquire environmental DNA through multiple

mechanisms including natural competence, one of the most common is acquisition

via plasmids by conjugation. One of the challenges for conjugation is an active Type

VI Secretion System (T6SS), which kills nonkin bacteria and therefore inhibits the

conjugation. The acquisition of resistance genes has helped the spread of resistant

clones of *A. baumannii.* The most prominently disseminated clones belong to

specific sequence types [(ST), Pasteur scheme], namely ST1, ST2, ST10, ST15,

ST25, ST79 and ST85 with the ST2 being the most detected ST. While ST1 and ST2

were thoroughly studied, there are new emerging clones whose genetic background

facilitating the rise is still unknown or poorly understood. One of the rising clones is

A. baumannii ST19, recently reported to be frequently occurring in Georgia and

Ukraine. In this study, we have whole-genome sequenced three isolates of *A.*

baumannii ST19 which were then assessed for biofilm production and virulence in

the *Galleria mellonella in vivo* infection model. Further 156 *A. baumannii* ST19

genomes were obtained from public repositories. In total, 157/159 genomes revealed

loss of *ompA* and T6SS locus, which was in 56/157 genomes (35,7%) replaced by

ΔTn*9* carrying chloramphenicol and ΔTn*10* carrying tetracycline resistance genes

and formaldehyde and chlorite resistance genes. Hindered T6SS-loss-mediated

contact-killing might promote the acquisition of genetic material supporting the

success of the *A. baumannii* ST19 lineage.

Importance

Acinetobacter baumannii is a nosocomial pathogen with immense potential to

become multidrug- to pan-drug-resistant, driven in part to its capacity to acquire DNA

from the environment through conjugation and natural competence. Conjugation

might however be hindered by an active type VI secretion system (T6SS), which kills

nonkin bacteria. In *A. baumannii* ST19, the absence of the T6SS enables the co-

existence and conjugation of nonkin bacteria. This, together with the reduced

metabolic burden of lacking T6SS and further acquisition of antibiotic and non-

antibiotic resistance traits, might enable the emergence of a successful, resistant,

and potentially high-risk *A. baumannii* lineage ST19.

Introduction

Acinetobacter baumannii is an emerging opportunistic nosocomial pathogen that

causes a variety of infections including urinary tract infections, wound infections,

meningitis, ventilator-associated pneumonia, blood-stream infection, and

endocarditis (1). *A. baumannii* is a member of ESKAPE pathogens (*Enterococcus*

faecium, *Staphylococcus   aureus*, *Klebsiella   pneumoniae*, *Acinetobacter*

baumannii, *Pseudomonas aeruginosa* and *Enterobacter* spp.) (2), and due to its

frequent resistance to last-line antibiotics such as carbapenems, it has been

designated as a critical priority in the list of pathogens of importance for studying and

developing new antibiotics by the World Health Organization's list (3, 4).

Besides conventional antibiotics, *A. baumannii* can withstand challenging

environment conditions such as oxidative stress, prolonged periods of desiccation,

human serum but also disinfectants (5).

The ability of *A. baumannii* to acquire exogenous DNA via natural competence (6)

and conjugation (7) contributes to high plasticity of its genome (6). The horizontal

gene transfer plays significant role in the evolution of *A. baumannii* lineages,

resulting in number of highly successful clonal lineages such as ST1, ST2, ST10,

ST15, ST25, ST79 and ST85 (Pasteur scheme) (8, 9). While ST1 and ST2 are

responsible for the majority of infections caused by extensively-drug resistant *A.*

baumannii, some clones are specific to certain geographical locations (9) , such as

ST25 (Europe, United Arab Emirates, South America, Asia) (10) , ST32 (South

Korea, Europe, the USA, Canada and the Middle East) (11) or ST78 (Italy, Ukraine,

USA) (12).

While the horizontal gene transfer, especially plasmid transfer via conjugation, can

shape entire lineages and their success (13) an active Type VI secretion system

(T6SS) poses a unique challenge to conjugative plasmids, as *Acinetobacter* plasmid

donors and recipients may kill each other (14). The T6SS machinery, including outer

membrane protein *ompA*, is usually encoded in a single chromosomal locus (15).

The T6SS of *A. baumannii* has been shown to kill *E. coli*, *K. pneumoniae* and *P.*

aeruginosa, while different strains of *A. baumannii* are also able to kill each other

(15, 16). While the role of *ompA* was demonstrated to influence the virulence and

biofilm formation of *A. baumannii* (17) and was as well proposed as a possible

therapeutic and vaccine target (18) the role of T6SS in *Galleria mellonella* model of

virulence is believed to be indirect and strain dependent (19). Similarly, the presence

T6SS in *A. baumannii* was also linked to higher biofilm-forming activity and better

survival in the presence of normal human serum (20). Notably, the OmpA is rather

required for the formation of thick biofilms on polystyrene, than essential for biofilm

development, (21) and implicated in inflammation, tissue destruction in a murine

model of infection (22).

In this study, we have characterized a rising lineage of *A. baumannii* ST19, of which

occurrence is increasing regionally (Ukraine, Georgia) (23) in its ability to form

biofilm on abiotic surfaces and its virulence potential in *G. mellonella in vivo* infection

model. Furthermore, we have performed comparative genomic analysis of the global

population of *A. baumannii* ST19 in order to decipher genomic traits that make this

lineage emergingly successful and to assess its threat in becoming world-wide

disseminated clone.

110 *Results*

111 *A. baumannii* ST19 strains and genomes

A concerning report on the increasing occurrence of *Acinetobacter baumannii* ST19

encoding carbapenemases was recently published (23), raising questions about

factors contributing the success of this lineage. Three isolates of *A. baumannii* ST19

were obtained from two different patients (patient 1 - wound, rectal swab; patient 2 -

rectal swab) and long-read whole-genome sequenced and *de novo* assembled.

A further 156 genomes (including 97 from Luo et al. (23)) of *A. baumannii* ST19 were

retrieved from public repositories (GenBank and BIGSdb). *A. baumannii* ST19 has

been occurring with increased frequency (Figure 1), suggesting it is successfully

adapted to the hospital environment, the host or both.

Figure 1: Temporal and spatial occurrence of *A. baumannii* ST19

The examined genomes carried a variety of virulence and resistance genes (Figure

2), including resistance to last-resort antibiotics such as carbapenems. One-third

(n=53) out of 159 genomes described in this study encoded at least one

carbapenemase gene (bla_{OXA-23}, bla_{OXA-72}, bla_{NDM-1} or bla_{NDM-5}) with 6 encoding two

and 9 encoding three carbapenemases. Furthermore, a broad variety of plasmid

replicons was detected (Figure 2), suggesting easy acquisition of genetic material.

<mark>Figure 2: BOTTOM</mark>

Figure 2: Phylogenetic analysis of 3 isolates and further 156 of *A. baumannii* ST19

genomes, depicting year of isolation, country, capsule locus type (KL), outer core

lipooligosaccharide type (OCL), and presence of GI*sul2* genetic island, chlorite

dismutase, resistance, T6SS, formaldehyde, plasmid replicons, resistance, and

virulence genes, respectively.

Genomic analysis suggests that the possible underlying reason for its success to be

the acquisition of multiple partial transposons, including ΔTn*9* (24) carrying

chloramphenicol resistance gene and ΔTn*10* carrying the tetracycline resistance

gene. Interestingly, 61/159 (38,4 %) genomes of *A. baumannii* ST19 carried chlorite

dismutase and formaldehyde resistance (*frmRAB*) genes, with 98,4% AA identity to

Massilia glaciei and 96,7% to *Ideonella dechloratans*, where it combats oxidative

stress (25). To our knowledge, this is the first time chlorite dismutase has been

reported in the genome of *A. baumannii*.

Furthermore, these transposons disrupted and replaced the locus encoding *ompA*

and the T6SS, impairing persistence as the examined isolates were weak biofilm

producers (Figure 3). The entire region of T6SS and *ompA* deletion exhibits

extensive genetic rearrangements, as ΔTn*9* and ΔTn*10* were acquired and disrupted

by IS*26* and IS*4321,* and IS*26*, respectively.

Figure 3: Crystal violet quantification of biofilm production in three ST19 strains of *A.*

baumannii, AB5075 and non-inoculated LB medium as a control.

However, the virulence in the *G. mellonella* infection model varied (Figure 4),

supporting the observation that T6SS is not directly involved in virulence in *G.*

mellonella (19). The absence of T6SS, enabling conjugal acquisition of genetic

material (14) such as antibiotic, formaldehyde and chlorite resistance, could explain

their successful rise and frequent occurrence.

Figure 4: *In vivo* virulence in *G. mellonella* of three ST19 *A. baumannii* strains and AB5075 showing lower virulence potential of *A. baumannii* ST19 compared to AB5075 at infection dose of 2×10^7 CFU/mL. The x axis shows percentage of survival and the y axis the elapsed time in hours.

Genetic variability of the *A. baumannii* ST19 lineage

167
168

The whole genome sequences were subjected to genotyping [sequence type, capsule locus (KL) type, outer core lipooligosaccharide locus (OCL) type, antimicrobial resistance, and virulence genes]. A phylogenetic analysis was performed on these isolates, including genomes of the same ST from public repositories (GenBank and BIGSdb) (Figure 2).

The vast majority (157/159) of *A. baumannii* ST19 harbored GI*sul2,* a chromosomally carried Genomic Island encoding sulphonamide resistance determinant (*sul2*) along with putative arsenic resistance genes (26). We have detected a range of 0 to 6 plasmid replicons (27) in the genomes of *A. baumannii*

ST19, with rP-T1 replicon being the most prevalent (n=122/159), followed by r3-T23

(n=46/159), r3-T12 and r1-T8 (n=44/159 both).

Interestingly, 18 isolates did not carry any detectable plasmid replicon raising a

question whether there is another mechanism limiting acquisition of plasmids or

whether they have not encountered conjugative plasmid yet. However, the

Acinetobacter typing schemes are still being developed (27), possibly undetected (or

below the detection threshold) plasmids must be considered as well.

While various ST (Pasteur scheme) such as ST2 or ST1 often belong to various KL

types (6), the ST19 lineage exhibited limited variability with 69/159 genomes being

KL91, 66/159 KL15, followed by KL1 (13/159), KL147 (8/159), KL17 (2/159) and

KL40 (1/159). Interestingly, 53/55 OCL1 encoded KL15 (2/55 OCL1 were KL17), the

most predominant OCL type, OCL5 (n=87) was variable and encompassed KL91,

KL1, KL147 and KL15. Furthermore, all of the genomes of KL15/OCL1 formed one

large, genetically related cluster (Figure 2).

The capsule abundance was quantified in a gradient of silica. While the relative

abundance was same for the tested *A. baumannii* ST19 isolates, it was lower

amount of capsular polysaccharide than in AB5075 (Figure 5). While the capsulation

level was same for all three isolates, the virulence phenotype was not identical

(Figure 4), supporting previous observations of high capsulation not correlating with

higher virulence potential (6).

Figure 5: The quantification of the capsular polysaccharide production in the gradient

of colloidal silica (Ludox LS) showing relatively high capsulation of all three tested *A.*

baumannii ST19 strains. The AB5075 strain was tested as a reference.

Evaluating the T6SS locus in the ST19 lineage of *A. baumannii* we identified

variability in genetic origin of the deletion of these genes. Of 61/159 genomes

carrying chlorite dismutase and formaldehyde resistance genes in the locus of T6SS

and *ompA*, 55/61 harbored ΔTn*10* carrying the tetracycline and ΔTn*9* carrying

chloramphenicol resistance genes (Figure 6). Five formaldehyde resistance and

chlorite dismutase-encoding genomes (5/61) did not harbor ΔTn*10* carrying the

tetracycline resistance and 1/61 did not harbor ΔTn*9* carrying chloramphenicol

resistance genes. However, mapping on the reference genome (0604 -

SAMN46706488) confirmed, that in all these genomes (61/159) these genetic

rearrangements displaced locus encoding T6SS and the *ompA* gene.

Figure 6: Former T6SS and *ompA* locus of *A. baumannii* ST19 (0788 - SAMN46706486) as a representative genome showing deletion mediated by ΔTn*10*, ΔTn*9*, and formaldehyde resistance locus.

Interestingly, 2/159 genomes of *A. baumannii* ST19 (GCA_042643835.1 and GCA_040868255.1) still encoded T6SS and *ompA* locus. However, compared to AB5075-VUB (accession nr. CP070362), a number of SNPs (99% pairwise identity over 19417 bp) was detected in these two genomes, while the broader genetic neighbourhood only sporadically exhibited SNPs. The genomes GCA_042643835.1 and GCA_040868255.1 belonged to KL40/OCL2 and KL17/OCL1, respectively, which are rare for this lineage (a sole KL40 and one of two KL17, respectively). While none of the SNPs were non-sense, the influence of missense or synonymous (e.g., via tRNA) on the functionality of T6SS remains to be explored.

Discussion

Acinetobacter has the core components of the T6SS conserved and found in a single chromosomal locus (15). The *Acinetobacter* strains use type VI secretion system (T6SS) to kill nonkin bacteria such as *E. coli*, *K. pneumoniae* and *P. aeruginosa* (16, 28). The active T6SS is also a challenge to conjugation and horizontal gene transmission (15). The absence of T6SS could therefore allow co-existence of nonkin bacterial species and strains. We hypothesize that this might and explain original

acquisition of genetic material from these species and possibly evolution of the *A.*

baumannii ST19 lineage.

While the Tn*10* (29) was described, the best match of its nucleotide sequence was

detected with *S. enterica* serovar Typhi and *S. flexneri*. The ΔTn*10* observed in this

study has the highest similarity to regions from *Enterobacter hormaechei* IncHI2

palsmids (>99% nucleotide identity and 97% query coverage). Furthermore, the

ΔTn*9* (30) examined in this study was BLASTn-detected (with >99% nucleotide

identity and 100% query coverage) in *Klebsiella pneumoniae* IncF plasmids.

Although *adhC* genes facilitating resistance to formaldehyde which are native to

Acinetobacter have been reported (31), we have identified locus, likely originating

from *Enterobacter hormaechei* IncHI2 plasmid, or *Klebsiella pneumoniae* (or

quasipneumoniae) IncF plasmid (95% nucleotide identity and 100% coverage).

However, the origin of the chlorite dismutase gene remains undeciphered, while the

BLASTn of here observed chlorite dismutase results in identical hits (>96%

nucleotide identity and 98% query coverage) in various species, such as

Pseudomonas aeruginosa (chromosome), *S. enterica* (IncHI2 plasmid), *K.*

pneumoniae (IncF plasmid), *E. hormaechei* (chromosome), but also *Vibrio*

rumoiensis (plasmid) *Raoultella ornithinolytica* (IncF plasmid) *Leclercia*

adecarboxylata (IncHI2 plasmid).

Interestingly, the virulence in *G. mellonella* in this study varied, supporting previous

observations that T6SS is not directly implicated in pathogenicity, and the virulence

in *G. mellonella* might be indirect (19). As isolates with T6SS disrupted by IS*Aba1*

were observed (32) it has been suggested that T6SS is not essential to cause the

disease.

The OmpA is an outer membrane protein (33), encoded in the T6SS locus and is

known to be involved in the formation of a translocation pore (34). The OmpA has

been extensively studied for its role in *Acinetobacter* virulence (35). Furthermore, the

OmpA plays a role in the development of robust biofilms on plastic, and A549 human

alveolar epithelial cells (18, 21). This supports the observation of low biofilm forming

ability of strains examined in this study. Furthermore, the loss of the *ompA* gene in

the *A. baumannii* ST19 may have further consequences as it might render one of the

most prominent vaccines targets ineffective (18, 36).

Conclusion

A. baumannii ST19 has been occurring with increased frequency, suggesting it has

successfully adapted to the hospital environment. The loss of T6SS in the (nearly)

entire lineage of *A. baumannii* ST19 points towards a beneficial evolutionary step, a

fitness trade-off towards favouring environments where antibiotic resistance is key

for survival. The increasing prevalence of the *A. baumannii* ST19 lineage and its

antibiotic and non-antibiotic (environmental stresses and disinfectants) resistance is

highly concerning and warrants greater attention in surveillance efforts.

Material and Methods

Bacterial isolates, genomes and metadata

Three isolates of *A. baumannii* ST19 were obtained from two different patients

(patient 1 - wound, rectal swab; patient 2 - rectal swab). A further 156 genomes

(including 97 from Luo et al.(23)) of *A. baumannii* ST19 were retrieved from public

repositories (GenBank and BIGSdb) as of March 2025.

DNA extraction and whole genome sequencing

The DNA was extracted using DNeasy Blood & Tissue Kit (QIAGEN). A rapid

barcoding kit (V14) (ONT) was then used for library preparation which was then

subjected to long-read whole-genome sequencing using the R10.4.1 flowcell with the

PromethION (P2 Solo) platform by Nanopore.

Genome assembly and annotation

The long-read sequences were basecalled using the SUP model and adaptor and

quality (Q ≤ 15) trimmed using Porechop v0.2.2 (https://github.com/rrwick/Porechop)

and NanoFilt v2.8.0 (37), respectively. The reads were then assembled using

Flye2.9.3 (38).

Phylogenetic analysis

The maximum-likelihood tree was constructed by Prokka (39) generated gff files

used for the core-genome alignment created by Roary (40). RAxML (41) was used

for the calculation of the phylogenetic tree under GTRGAMMA model of rate

heterogeneity method supported by 500 bootstraps. The tree was visualized in iTOL

(42).

Identification of resistance determinants

The resistance and virulence genes were detected using ABRicate

(https://github.com/tseemann/abricate) employing ResFinder (43) and VFDB 2025

(44) databases, respectively, with a 90% threshold for both gene identity and

coverage. The typing of capsule-encoding loci (KL) and lipooligosaccharide outer

core loci (OCL) was determined using Kaptive (45). The ABRicate

(https://github.com/tseemann/abricate) was used to detect the presence of plasmid

replicons using AcinetobacrPlasmidTyping database

(https://github.com/MehradHamidian/AcinetobacterPlasmidTyping) (27), while the

genes encoding GIs*ul2*, chlorite dismutase, formaldehyde resistance genes, *ompA*

and T6SS-encoding genes were detected using a custom database.

Capsule production

The production of capsular polysaccharide was assessed as described before (6).

Briefly, 1mL of overnight culture was centrifuged (2 minutes, 7000g), supernatant

was removed a resuspended 1mL of PBS. Then, 875μL of the suspension was

mixed with 125μL of Ludox LS colloidal silica [30% (wt/wt) suspension in H_2O] and

centrifuged (30 minutes, 12000g) and the distance of the center of the band from the

bottom of the microtube was measured. The measurements were taken in biological

triplicate.

Biofilm quantification assay

The relative amount of biofilm was measured using a biofilm assay in a 96-well flat-

bottomed polystyrene plate. The overnight cultures were adjusted to an OD_{600} of

0.05 in Luria–Bertoni medium and 200 μL of bacterial suspension were then

inoculated into 12 wells per isolate and incubated statically, overnight at 37°C. The

wells were then washed twice with 200 μL of PBS, air-dried, and stained with crystal

violet 0.1% v/v for 15 min. Then, the crystal violet was pipetted out and wells were

again washed twice with 200 μL of PBS and air-dried. The stain was then solubilised

with ethanol 95% v/v for 5 min and transferred to a new 96-well plate and measured

in a plate reader (Cytation, Agilent BioTek) at OD_{570}. Each experiment was

performed in biological triplicate.

***Galleria mellonella in vivo* infection model of virulence**

The *G. mellonella* infections assay was performed as previously described (46).

Briefly, an overnight culture was washed with physiological saline (PS) (0.9% NaCl)

and diluted to known Colony Forming Unites (CFU) at $OD_{600}1$. Then approximately 2

$\times 10^7$ CFU/mL of *A. baumannii* in 10 µL of PS was injected in the last left proleg

using a 0.3 mL insulin syringe (BD MicroFine). Each strain was injected into 10

larvae, and 10 larvae were injected with PS as a negative control. The experiments

were carried out in duplicates, and the survival of larvae was visualized in Kaplan-

Meier curve in a time span of 72 hours.

Data availability

The whole genome sequencing data were deposited in GenBank under following

accession numbers; strain 0788 - SAMN46706486, strain 0482 -

SAMN48114201and strain 0604 - SAMN46706488.

Ethical statement

The clinical isolates sequenced in this study were obtained within retrospective study

conducted under the ethical approval granted by the C. H. U. Brugmann CE

2023/51.

Acknowledgements

AV is a recipient of a junior postdoctoral fellowship of the Research Foundation – Flanders (FWO; file number 1287223N). CVDH is supported by the Flanders Institute for Biotechnology (VIB). Sequencing was performed by VIB Nucleomics Core (www.nucleomics.be). The Vande Velde Lab kindly provided *G. mellonella* larvae.

References

1. Brek TM, Muhajir AA, Alkuwaity KK, Haddad MA, Alattas EM, Eisa ZM, Al-Thaqafy MS, Albarraq AM, Al-Zahrani IA. Genomic insights of predominant international high-risk clone ST2 Acinetobacter baumannii isolates in Saudi Arabia. J Glob Antimicrob Resist. 2025 Mar 28;42:243-252. doi: 10.1016/j.jgar.2025.03.012. Epub ahead of print. PMID: 40158887.

2. Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens. Expert Rev Anti Infect Ther. 2013 Mar;11(3):297-308. doi: 10.1586/eri.13.12. PMID: 23458769.

3. WHO Bacterial Priority Pathogens List, 2024: bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance. Geneva: World Health Organization; 2024. Licence: CC BY-NC-SA 3.0 IGO.

4. Antochevis LC, Wilhelm CM, Arns B, Sganzerla D, Sudbrack LO, Nogueira TCRL, Guzman RD, Martins AS, Cappa DS, Dos Santos ÂC, Pascual JC, Perugini VH, Vespero EC, Rigatto MHP, Pereira DC, Lutz L, Leão RS, Marques EA, Henrique DM, Coelho AAM, Frutuoso LL, de A Sousa EE, Abreu Guimarães LF, Ferreira ALP, Castiñeiras AC, Alves MD, Telles JP, Yamada CH, de Almeida FP, Girão ES, de Sousa PCP, de Melo AGND, Mendes ET, Rocha VFD, Neves EDS, Ribeiro MT, Starling

CEF, Oliveira MS, Sampaio JLM, Martins AF, Barth AL, Zavascki AP; ASCENSION Study Group. World Health Organization priority antimicrobial resistance in *Enterobacterales, Acinetobacter baumannii, Pseudomonas aeruginosa, Staphylococcus aureus* and *Enterococcus faecium* healthcare-associated bloodstream infections in Brazil (ASCENSION): a prospective, multicentre, observational study. Lancet Reg Health Am. 2025 Jan 30;43:101004. doi: 10.1016/j.lana.2025.101004. PMID: 39957800; PMCID: PMC11830303.

5. Hooppaw AJ, McGuffey JC, Di Venanzio G, Ortiz-Marquez JC, Weber BS, Lightly TJ, van Opijnen T, Scott NE, Cardona ST, Feldman MF. The Phenylacetic Acid Catabolic Pathway Regulates Antibiotic and Oxidative Stress Responses in Acinetobacter. mBio. 2022 Jun 28;13(3):e0186321. doi: 10.1128/mbio.01863-21. Epub 2022 Apr 25. PMID: 35467424; PMCID: PMC9239106.

6. Valcek A, Philippe C, Whiteway C, Robino E, Nesporova K, Bové M, Coenye T, De Pooter T, De Coster W, Strazisar M, Van der Henst C. Phenotypic Characterization and Heterogeneity among Modern Clinical Isolates of Acinetobacter baumannii. Microbiol Spectr. 2023 Feb 14;11(1):e0306122. doi: 10.1128/spectrum.03061-22. Epub 2022 Dec 8. PMID: 36475894; PMCID: PMC9927488.

7. Brovedan MA, Cameranesi MM, Limansky AS, Morán-Barrio J, Marchiaro P, Repizo GD. What do we know about plasmids carried by members of the Acinetobacter genus? World J Microbiol Biotechnol. 2020 Jul 13;36(8):109. doi: 10.1007/s11274-020-02890-7. PMID: 32656745.

8. Cain AK, Hamidian M. Portrait of a killer: Uncovering resistance mechanisms and global spread of Acinetobacter baumannii. PLoS Pathog. 2023 Aug 10;19(8):e1011520. doi: 10.1371/journal.ppat.1011520. PMID: 37561719; PMCID: PMC10414682.

9. Hamidian M, Nigro SJ. Emergence, molecular mechanisms and global spread of

carbapenem-resistant *Acinetobacter baumannii*. Microb Genom. 2019

Oct;5(10):e000306. doi: 10.1099/mgen.0.000306. PMID: 31599224; PMCID:

PMC6861865.

10. Sahl JW, Del Franco M, Pournaras S, Colman RE, Karah N, Dijkshoorn L, Zarrilli R.

Phylogenetic and genomic diversity in isolates from the globally distributed

Acinetobacter baumannii ST25 lineage. Sci Rep. 2015 Oct 14;5:15188. doi:

10.1038/srep15188. PMID: 26462752; PMCID: PMC4604477.

11. Tobin LA, Abu Sabah E, Lebreton F, Myers GSA, McGann PT, Hamidian M.

Genomic analysis of early ST32 Acinetobacter baumannii strains recovered in US

military treatment facilities reveals distinct lineages and links to the origins of the

Tn6168 ampC transposon. J Antimicrob Chemother. 2025 Mar 3;80(3):666-675. doi:

10.1093/jac/dkae454. PMID: 39680383; PMCID: PMC11879209.

12. Harmer CJ, Cahill SM, Kenyon JJ, Hall RM. Aminoglycoside resistance genes in

early members of the Acinetobacter baumannii ST78A (SMAL, Italian clone) reside

in an IS26-bounded island in the chromosome. J Antimicrob Chemother. 2024 May

2;79(5):1014-1018. doi: 10.1093/jac/dkae064. PMID: 38530861; PMCID:

PMC11062947.

13. Gao Y, Li H, Chen H, Zhang J, Wang R, Wang Z, Wang H. Origin, Phylogeny, and

Transmission of the Epidemic Clone ST208 of Carbapenem-Resistant Acinetobacter

baumannii on a Global Scale. Microbiol Spectr. 2022 Jun 29;10(3):e0260421. doi:

10.1128/spectrum.02604-21. Epub 2022 May 31. PMID: 35638783; PMCID:

PMC9241911.

14. Di Venanzio G, Moon KH, Weber BS, Lopez J, Ly PM, Potter RF, Dantas G, Feldman

MF. Multidrug-resistant plasmids repress chromosomally encoded T6SS to enable

their dissemination. Proc Natl Acad Sci U S A. 2019 Jan 22;116(4):1378-1383. doi:

10.1073/pnas.1812557116. Epub 2019 Jan 9. PMID: 30626645; PMCID:

PMC6347727.

15. Weber BS, Kinsella RL, Harding CM, Feldman MF. The Secrets of Acinetobacter

Secretion. Trends Microbiol. 2017 Jul;25(7):532-545. doi: 10.1016/j.tim.2017.01.005.

Epub 2017 Feb 16. PMID: 28216293; PMCID: PMC5474180.

16. Repizo GD, Gagné S, Foucault-Grunenwald ML, Borges V, Charpentier X, Limansky

AS, Gomes JP, Viale AM, Salcedo SP. Differential Role of the T6SS in Acinetobacter

baumannii Virulence. PLoS One. 2015 Sep 24;10(9):e0138265. doi:

10.1371/journal.pone.0138265. PMID: 26401654; PMCID: PMC4581634.

17. Cerqueira GM, Peleg AY. Insights into Acinetobacter baumannii pathogenicity.

IUBMB Life. 2011 Dec;63(12):1055-60. doi: 10.1002/iub.533. Epub 2011 Oct 12.

PMID: 21989983.

18. Nie D, Hu Y, Chen Z, Li M, Hou Z, Luo X, Mao X, Xue X. Outer membrane protein

A (OmpA) as a potential therapeutic target for Acinetobacter baumannii infection. J

Biomed Sci. 2020 Jan 18;27(1):26. doi: 10.1186/s12929-020-0617-7. PMID:

31954394; PMCID: PMC6969976.

19. Flaugnatti N, Bader L, Croisier-Coeytaux M, Blokesch M. Capsular polysaccharide

restrains type VI secretion in *Acinetobacter baumannii*. Elife. 2025 Jan 3;14:e101032.

doi: 10.7554/eLife.101032. PMID: 39749675; PMCID: PMC11731876.

20. Kim J, Lee JY, Lee H, Choi JY, Kim DH, Wi YM, Peck KR, Ko KS. Microbiological

features and clinical impact of the type VI secretion system (T6SS) in Acinetobacter

baumannii isolates causing bacteremia. Virulence. 2017 Oct 3;8(7):1378-1389. doi:

10.1080/21505594.2017.1323164. Epub 2017 Apr 27. PMID: 28448786; PMCID:

PMC5711433.

21. Gaddy JA, Tomaras AP, Actis LA. The Acinetobacter baumannii 19606 OmpA protein plays a role in biofilm formation on abiotic surfaces and in the interaction of this pathogen with eukaryotic cells. Infect Immun. 2009 Aug;77(8):3150-60. doi: 10.1128/IAI.00096-09. Epub 2009 May 26. PMID: 19470746; PMCID: PMC2715673..

22. Choi CH, Lee JS, Lee YC, Park TI, Lee JC. Acinetobacter baumannii invades epithelial cells and outer membrane protein A mediates interactions with epithelial cells. BMC Microbiol. 2008 Dec 10;8:216. doi: 10.1186/1471-2180-8-216. PMID: 19068136; PMCID: PMC2615016.

23. Luo TL, Martin MJ, Kovalchuk V, Kondratiuk V, Trapaidze N, Metreveli M, Hulseberg CE, Dao HD, Kwak YI, Maybank R, Musich TA, Scherer MR, Bennett JW, Mc Gann PT, Lebreton F. Detection of carbapenemase producing *Acinetobacter baumannii* ST19 from Georgia and Ukraine carrying bla_{OXA-23}, bla_{OXA-72}, and/or bla_{NDM-5}, December 2019 to June 2023. Euro Surveill. 2024 Jun;29(24):2400259. doi: 10.2807/1560-7917.ES.2024.29.24.2400259. PMID: 38873796; PMCID: PMC11177571.

24. Boyd D, Taylor G, Fuller J, Bryce E, Embree J, Gravel D, Katz K, Kibsey P, Kuhn M, Langley J, Mataseje L, Mitchell R, Roscoe D, Simor A, Thomas E, Turgeon N, Mulvey M; Canadian Nosocomial Infection Surveillance Program. Complete Sequence of Four Multidrug-Resistant MOBQ1 Plasmids Harboring blaGES-5 Isolated from Escherichia coli and Serratia marcescens Persisting in a Hospital in Canada. Microb Drug Resist. 2015 Jun;21(3):253-60. doi: 10.1089/mdr.2014.0205. Epub 2014 Dec 29. PMID: 25545311.

25. Shaffer JMC, Giddings LA, Samples RM, Mikucki JA. Genomic and phenotypic characterization of a red-pigmented strain of *Massilia frigida* isolated from an

Antarctic microbial mat. Front Microbiol. 2023 May 12;14:1156033. doi: 10.3389/fmicb.2023.1156033. PMID: 37250028; PMCID: PMC10213415.

26. Hamidian M, Hall RM. Acinetobacter baumannii ATCC 19606 Carries GIsul2 in a Genomic Island Located in the Chromosome. Antimicrob Agents Chemother. 2016 Dec 27;61(1):e01991-16. doi: 10.1128/AAC.01991-16. PMID: 27795382; PMCID: PMC5192138.

27. Lam MMC, Koong J, Holt KE, Hall RM, Hamidian M. Detection and Typing of Plasmids in Acinetobacter baumannii Using *rep* Genes Encoding Replication Initiation Proteins. Microbiol Spectr. 2023 Feb 14;11(1):e0247822. doi: 10.1128/spectrum.02478-22. Epub 2022 Dec 6. PMID: 36472426; PMCID: PMC9927589.

28. Weber BS, Ly PM, Irwin JN, Pukatzki S, Feldman MF. A multidrug resistance plasmid contains the molecular switch for type VI secretion in Acinetobacter baumannii. Proc Natl Acad Sci U S A. 2015 Jul 28;112(30):9442-7. doi: 10.1073/pnas.1502966112. Epub 2015 Jul 13. PMID: 26170289; PMCID: PMC4522760.

29. Chalmers R, Sewitz S, Lipkow K, Crellin P. Complete nucleotide sequence of Tn10. J Bacteriol. 2000 May;182(10):2970-2. doi: 10.1128/JB.182.10.2970-2972.2000. PMID: 10781570; PMCID: PMC102010.

30. Alton NK, Vapnek D. Nucleotide sequence analysis of the chloramphenicol resistance transposon Tn9. Nature. 1979 Dec 20-27;282(5741):864-9. doi: 10.1038/282864a0. PMID: 390403.

31. Echenique JR, Dorsey CW, Patrito LC, Petroni A, Tolmasky ME, Actis LA. Acinetobacter baumannii has two genes encoding glutathione-dependent formaldehyde dehydrogenase: evidence for differential regulation in response to iron.

Microbiology (Reading). 2001 Oct;147(Pt 10):2805-2815. doi: 10.1099/00221287-147-10-2805. PMID: 11577159.

32. Wright MS, Haft DH, Harkins DM, Perez F, Hujer KM, Bajaksouzian S, Benard MF, Jacobs MR, Bonomo RA, Adams MD. New insights into dissemination and variation of the health care-associated pathogen Acinetobacter baumannii from genomic analysis. mBio. 2014 Jan 21;5(1):e00963-13. doi: 10.1128/mBio.00963-13. Erratum in: mBio. 2015 Jun 04;6(3):e00585. doi: 10.1128/mBio.00585-15. PMID: 24449752; PMCID: PMC3903280.

33. Shrivastava S, Mande SS. Identification and functional characterization of gene components of Type VI Secretion system in bacterial genomes. PLoS One. 2008 Aug 13;3(8):e2955. doi: 10.1371/journal.pone.0002955. PMID: 18698408; PMCID: PMC2492809.

34. Wang Y. The function of OmpA in Escherichia coli. Biochem Biophys Res Commun. 2002 Mar 29;292(2):396-401. doi: 10.1006/bbrc.2002.6657. PMID: 11906175.

35. Shadan A, Pathak A, Ma Y, Pathania R, Singh RP. Deciphering the virulence factors, regulation, and immune response to *Acinetobacter baumannii* infection. Front Cell Infect Microbiol. 2023 Feb 23;13:1053968. doi: 10.3389/fcimb.2023.1053968. PMID: 36968113; PMCID: PMC10038080.

36. Lau YT, Tan HS. *Acinetobacter baumannii* subunit vaccines: recent progress and challenges. Crit Rev Microbiol. 2024 Aug;50(4):434-449. doi: 10.1080/1040841X.2023.2215303. Epub 2023 May 21. PMID: 37211625.

37. De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack: visualizing and processing long-read sequencing data. Bioinformatics. 2018 Aug 1;34(15):2666-2669. doi: 10.1093/bioinformatics/bty149. PMID: 29547981; PMCID: PMC6061794.

38. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads

using repeat graphs. Nat Biotechnol. 2019 May;37(5):540-546. doi: 10.1038/s41587-

019-0072-8. Epub 2019 Apr 1. PMID: 30936562.

39. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014 Jul

15;30(14):2068-9. doi: 10.1093/bioinformatics/btu153. Epub 2014 Mar 18. PMID:

24642063.

40. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush

D, Keane JA, Parkhill J. Roary: rapid large-scale prokaryote pan genome analysis.

Bioinformatics. 2015 Nov 15;31(22):3691-3. doi: 10.1093/bioinformatics/btv421.

Epub 2015 Jul 20. PMID: 26198102; PMCID: PMC4817141.

41. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large   phylogenies.   Bioinformatics.   2014   May   1;30(9):1312-3.   doi:

10.1093/bioinformatics/btu033.   Epub   2014   Jan   21.   PMID: 24451623;   PMCID:

PMC3998144.

42. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic

tree display and annotation. Nucleic Acids Res. 2021 Jul 2;49(W1):W293-W296. doi:

10.1093/nar/gkab301. PMID: 33885785; PMCID: PMC8265157.

43. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A,

Allesoe RL, Rebelo AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B,

Werner G, Bender JK, Stingl K, Nguyen M, Coppens J, Xavier BB, Malhotra-Kumar

S, Westh H, Pinholt M, Anjum MF, Duggett NA, Kempf I, Nykäsenoja S, Olkkola S,

Wieczorek K, Amaro A, Clemente L, Mossong J, Losch S, Ragimbeau C, Lund O,

Aarestrup FM. ResFinder 4.0 for predictions of phenotypes from genotypes. J

Antimicrob Chemother. 2020 Dec 1;75(12):3491-3500. doi: 10.1093/jac/dkaa345.

PMID: 32780112; PMCID: PMC7662176.

44. Zhou S, Liu B, Zheng D, Chen L, Yang J. VFDB 2025: an integrated resource for

exploring anti-virulence compounds. Nucleic Acids Res. 2025 Jan 6;53(D1):D871-

D877. doi: 10.1093/nar/gkae968. PMID: 39470738; PMCID: PMC11701737.

45. Wyres KL, Cahill SM, Holt KE, Hall RM, Kenyon JJ. Identification of *Acinetobacter*

baumannii loci for capsular polysaccharide (KL) and lipooligosaccharide outer core

(OCL) synthesis in genome assemblies using curated reference databases compatible

with *Kaptive*. Microb Genom. 2020 Mar;6(3):e000339. doi: 10.1099/mgen.0.000339.

PMID: 32118530; PMCID: PMC7200062.

46. Whiteway C, Valcek A, Philippe C, Strazisar M, De Pooter T, Mateus I, Breine A, Van

der Henst C. Scarless excision of an insertion sequence restores capsule production

and virulence in Acinetobacter baumannii. ISME J. 2022 May;16(5):1473-1477. doi:

10.1038/s41396-021-01179-3. Epub 2021 Dec 23. PMID: 34949784; PMCID:

PMC9038732.

Figure 2: Phylogenetic analysis of 3 isolates and further 156 of *A. baumannii* ST19 genomes, depicting year of isolation, country,

capsule locus type (KL), outer core lipooligosaccharide type (OCL), and presence of GI*sul2* genetic island, chlorite dismutase,

resistance, T6SS, formaldehyde, plasmid replicons, resistance, and virulence genes, respectively.

Tree scale: 0.001

Galleria mellonella infection assay

Tn*10* (AP000342) tetracycline resistance

Tn*9* (CP010373) chloramphenicol resistance

chlorite resistance

Δ T6SS
A. baumannii ST19

formaldehyde resistance
(LC589173)