Title: The absence of the Type VI Secretion System in the successful

lineage of *Acinetobacter baumannii* ST19

Running title: Loss of Type VI Secretion System in *A. baumannii* ST19

Authors: Adam Valcek[1, 2*], Diana Isabela Costescu Strachinaru[3], Kristina Nesporova[4],

Patrick Soentjens[3,5] Anke Stoefs[6,7], Charles Van der Henst[1, 2*]

Affiliations:

1. Microbial Resistance and Drug Discovery, VIB-VUB Center for Structural Biology, VIB,

Flanders Institute for Biotechnology, Brussels, Belgium

2. Structural Biology Brussels, Vrije Universiteit Brussel (VUB), Brussels, Belgium

3. Center for Infectious Diseases, Queen Astrid Military Hospital, Brussels, Belgium

4. Microbial Systems Cell Biology Group, Department of Biology, KU Leuven, Leuven,

Belgium

5. Department of Clinical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

6. Department of Microbiology, Queen Astrid Military Hospital, Brussels, Belgium

7. Department of Microbiology, Universitair Ziekenhuis Brussels, Brussels Belgium

* These authors contributed equally to this work and share the last authorship.

Corresponding authors: Adam Valcek (*adam.valcek@vub.be),* Charles Van der Henst

(*charles.vanderhenst@vub.vib.be*)

Abstract

Acinetobacter baumannii is an opportunistic pathogen, often multi- to pandrug-resistant,

including to last-resort antibiotics such as carbapenems. *A. baumannii* can acquire DNA

through multiple mechanisms including natural competence and conjugation. An active Type

VI Secretion System (T6SS), which kills nonkin bacteria and challenges and inhibits the

conjugation. The acquisition of resistance genes has helped the spread of resistant clones of

A. baumannii, such as those of Sequence Types (ST^Pas), ST1, ST2, ST10, ST15, ST25, ST79

and ST85. While ST1 and ST2 were thoroughly studied, there are new emerging clones

whose genetic background facilitating the rise is still unknown or poorly understood.

The emergence of *Acinetobacter baumannii* ST19 first peaked in 2003, coinciding with the

Iraq conflict and increased detection among U.S. military personnel. Its prevalence later

surged in Georgia and Ukraine, around the onset of the conflict in Ukraine in 2022,

associating it with regions affected by military activity.

In this study, we have whole-genome sequenced three isolates of *A. baumannii* ST19 which

and further 156 *A. baumannii* ST19 genomes were obtained from public repositories. In total,

157/159 genomes revealed loss of T6SS locus, which was in 56/157 genomes (35,7%)

replaced by ΔTn*9* carrying chloramphenicol and ΔTn*10* carrying tetracycline resistance

genes, and formaldehyde and chlorite resistance genes. Surprisingly, the antibiotic resistance-

encoding transposons likely originated from Enterobacteriaceae plasmids. The loss of a

functional T6SS in *A. baumannii* ST19 may potentially facilitating horizontal gene transfer

and promoting a cooperative or less competitive lifestyle providing a selective advantage at

the population level.

Introduction

Acinetobacter baumannii is an emerging opportunistic nosocomial pathogen that causes a variety of infections including urinary tract infections, wound infections, meningitis, ventilator-associated pneumonia, blood-stream infection, and endocarditis (1). *A. baumannii* is a member of ESKAPE pathogens (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacter* spp.) (2), and due to its frequent resistance to last-line antibiotics such as carbapenems, it has been designated as a critical priority in the list of pathogens of importance for studying and developing new antibiotics by the World Health Organization's list (3, 4).

Besides conventional antibiotics, *A. baumannii* can withstand challenging environment conditions such as oxidative stress, prolonged periods of desiccation, human serum but also disinfectants (5).

The ability of *A. baumannii* to acquire exogenous DNA via natural competence (6) and conjugation (7) contributes to high plasticity of its genome (6). The horizontal gene transfer plays significant role in the evolution of *A. baumannii* lineages, resulting in number of highly successful clonal lineages such as ST1, ST2, ST10, ST15, ST25, ST79 and ST85 (Pasteur scheme) (8, 9). While ST1 and ST2 are responsible for the majority of infections caused by extensively-drug resistant *A. baumannii*, some clones are specific to certain geographical locations (9) , such as ST25 (Europe, United Arab Emirates, South America, Asia) (10) , ST32 (South Korea, Europe, the USA, Canada and the Middle East) (11) or ST78 (Italy, Ukraine, USA) (12).

While the horizontal gene transfer, especially plasmid transfer via conjugation, can shape entire lineages and their success (13), an active Type VI secretion system (T6SS) poses a

unique challenge to conjugative plasmids, as *Acinetobacter* plasmid donors and recipients

may kill each other (14). The T6SS machinery is usually encoded in a single chromosomal

locus (15). The T6SS of *A. baumannii* has been shown to mediate the killing of *E. coli*, *K.*

pneumoniae and *P. aeruginosa*, while different strains of *A. baumannii* are also able to kill

each other (15, 16).

In this study, we have assessed genetic determinants disrupting the T6SS-encoding locus in a

rising lineage of *A. baumannii* ST19, of which occurrence is increasing regionally (Ukraine,

Georgia) (17). Furthermore, we have performed comparative genomic analysis of the global

population of *A. baumannii* ST19 in order to decipher genomic traits that make this lineage

emergingly successful and to assess its threat in becoming world-wide disseminated clone.

Results

A. baumannii ST19 strains and genomes

A concerning report on the increasing occurrence of *Acinetobacter baumannii* ST19 encoding

carbapenemases was recently published (17), raising questions about factors contributing the

success of this lineage. Three isolates of *A. baumannii* ST19 were obtained from two

different patients (patient 1 - wound, rectal swab; patient 2 - rectal swab) and long-read

whole-genome sequenced and *de novo* assembled.

A further 156 genomes (including 97 from Luo et al. (17)) of *A. baumannii* ST19 were

retrieved from public repositories (GenBank and BIGSdb). The occurrence of *A. baumannii*

ST19 appears to spike during specific time periods that correlate with military conflicts

(Figure 1), such as the Iraq War (2003) and the war in Ukraine (2022).

Figure 1: Temporal and spatial occurrence of *A. baumannii* ST19 based on genomes

deposited in public repositories and available metadata.

The examined genomes carried a variety of virulence and resistance genes (Figure 2),

including resistance to last-resort antibiotics such as carbapenems. One-third (n=53) out of

159 genomes described in this study encoded at least one carbapenemase gene (bla_{OXA-23},

bla_{OXA-72}, bla_{NDM-1} or bla_{NDM-5}) with 6 encoding two and 9 encoding three carbapenemases.

Furthermore, a broad variety of plasmid replicons was detected (Figure 2), suggesting an easy

acquisition of genetic material by the horizontal gene transfer.

Figure 2: BOTTOM

Figure 2: Phylogenetic analysis of 3 isolates and further 156 of *A. baumannii* ST19 genomes,

depicting year of isolation, country, capsule locus type (KL), outer core lipooligosaccharide

type (OCL), and presence of GI*sul2* genetic island, chlorite dismutase, resistance, T6SS,

formaldehyde, plasmid replicons, resistance, and virulence genes, respectively.

Genomic analysis suggests that the possible underlying reason for its success to be the acquisition of multiple partial transposons, including ΔTn*9* (18) carrying chloramphenicol resistance gene and ΔTn*10* (19) carrying the tetracycline resistance gene. Interestingly, 61/159 (38,4 %) genomes of *A. baumannii* ST19 carried chlorite dismutase and formaldehyde resistance (*frmRAB*) genes, with 98,4% amino acid identity to *Massilia glaciei* and 96,7% to *Ideonella dechloratans*, where it combats oxidative stress (20). To our knowledge, this is the first time chlorite dismutase has been reported in the genome of *A. baumannii*.

Furthermore, these transposons disrupted and replaced the locus encoding the T6SS. The entire region of T6SS deletion exhibits extensive genetic rearrangements, as ΔTn*9* and ΔTn*10* were acquired and disrupted by IS*26* and IS*4321,* and IS*26*, respectively.

The absence of a functional T6SS, likely enabling conjugal acquisition of genetic material (14) such as antibiotic, formaldehyde and chlorite resistance, could represent one of the reasons behind their successful rise.

Genetic variability of the *A. baumannii* ST19 lineage

The whole genome sequences were subjected to genotyping [sequence type, capsule locus (KL) type, outer core lipooligosaccharide locus (OCL) type, antimicrobial resistance, and virulence genes]. A phylogenetic analysis was performed on these isolates, including genomes of the same ST from public repositories (GenBank and BIGSdb) (Figure 2).

The vast majority (157/159) of *A. baumannii* ST19 harbored GI*sul2,* a chromosomally carried Genomic Island encoding sulphonamide resistance determinant (*sul2*) along with putative arsenic resistance genes (21). We have detected a range of 0 to 6 plasmid replicons (22) in the genomes of *A. baumannii* ST19, with rP-T1 replicon being the most prevalent (n=122/159), followed by r3-T23 (n=46/159), r3-T12 and r1-T8 (n=44/159 both).

Interestingly, 18 isolates did not carry any detectable plasmid replicon raising a question

whether there is another mechanism limiting acquisition of plasmids or whether they have not

encountered conjugative plasmid yet. However, the *Acinetobacter* typing schemes are still

being developed (22), possibly undetected (or below the detection threshold) plasmids must

be considered as well.

While various ST (Pasteur scheme) such as ST2 or ST1 often belong to various KL types (6),

the ST19 lineage exhibited limited variability with 69/159 genomes being KL91, 66/159

KL15, followed by KL1 (13/159), KL147 (8/159), KL17 (2/159) and KL40 (1/159).

Interestingly, 53/55 OCL1 encoded KL15 (2/55 OCL1 were KL17), the most predominant

OCL type, OCL5 (n=87) was variable and encompassed KL91, KL1, KL147 and KL15.

Furthermore, all of the genomes of KL15/OCL1 formed one large, genetically related cluster

(Figure 2).

By evaluating the T6SS locus in the ST19 lineage of *A. baumannii*, we identified variability

in genetic origin of the deletion of these genes. Of 61/159 genomes carrying chlorite

dismutase and formaldehyde resistance genes in the locus of T6SS, 55/61 harbored ΔTn*10*

carrying the tetracycline and ΔTn*9* carrying chloramphenicol resistance genes (Figure 3).

Five formaldehyde resistance and chlorite dismutase-encoding genomes (5/61) did not harbor

ΔTn*10* carrying the tetracycline resistance and 1/61 did not harbor ΔTn*9* carrying

chloramphenicol resistance genes. However, mapping on the reference genome (0604 -

SAMN46706488) confirmed, that in all these genomes (61/159) these genetic rearrangements

displaced locus encoding T6SS.

Figure 3: Depiction of the T6SS locus disruption of *A. baumannii* ST19 (0482 -

SAMN48114201) as a representative genome, compared to AB5075-VUB (CP070362)

showing deletion mediated by ΔTn*9*, chlorite dismutase, formaldehyde resistance locus

frmRAB, and ΔTn*10*. The alignments are based on amino acid identity.

Interestingly, 2/159 genomes of *A. baumannii* ST19 (GCA_042643835.1 and

GCA_040868255.1) still encoded a likely functional T6SS. However, compared to AB5075-

VUB (accession nr. CP070362), a number of SNPs (99% pairwise identity over 19417 bp)

was detected in these two genomes, while the broader genetic neighbourhood only

sporadically exhibited SNPs. The genomes GCA_042643835.1 and GCA_040868255.1

belonged to KL40/OCL2 and KL17/OCL1, respectively, which are rare for this lineage (a

sole KL40 and one of two KL17, respectively). While none of the SNPs were non-sense, the

influence of missense or synonymous (e.g., via tRNA) on the functionality of T6SS remains

to be explored.

Discussion

Acinetobacter has the core components of the T6SS conserved and found in a single

chromosomal locus (15). The *A. baumannii* strains use Type VI Secretion System (T6SS) to

kill nonkin *A. baumannii* as well as other bacterial species such as *E. coli*, *K. pneumoniae* and

P. aeruginosa, (16, 23). The active T6SS can be a challenge to conjugation and horizontal

gene transmission (15). The absence of T6SS could therefore allow co-existence of nonkin

bacterial species and strains. We hypothesize that this might and explain original acquisition

of genetic material from these species and possibly evolution of the *A. baumannii* ST19

lineage.

The Tn*10* was originally identified and described in *Shigella flexneri* (19). This corresponds

to the best match of the ΔTn*10* nucleotide sequence in this study was detected for *S. enterica*

serovar Typhi and *S. flexneri*. The ΔTn*10* observed in this study has the highest similarity to

regions from *Enterobacter hormaechei* IncHI2 palsmids (>99% nucleotide identity and 97%

query coverage). Furthermore, the ΔTn*9* (24) examined in this study was BLASTn-detected

(with >99% nucleotide identity and 100% query coverage) in *Klebsiella pneumoniae* IncF

plasmids.

Although *adhC* genes facilitating resistance to formaldehyde which are native to

Acinetobacter have been reported (25), we have identified locus, likely originating from

Enterobacter hormaechei IncHI2 plasmid, or *Klebsiella pneumoniae* (or *quasipneumoniae*)

IncF plasmid (95% nucleotide identity and 100% coverage). However, the origin of the

chlorite dismutase gene remains undeciphered, while the BLASTn of here observed chlorite

dismutase results in identical hits (>96% nucleotide identity and 98% query coverage) in

various species, such as *P. aeruginosa* (chromosome), *S. enterica* (IncHI2 plasmid), *K.*

pneumoniae (IncF plasmid), *E. hormaechei* (chromosome), but also *Vibrio rumoiensis*

(plasmid) *Raoultella ornithinolytica* (IncF plasmid) *Leclercia adecarboxylata* (IncHI2

plasmid).

The total of 61/159 of genomes were identified to encode the same genetic cargo (ΔTn*9,*

ΔTn*10*, chlorite dismutase and the formaldehyde resistance cluster) as the one present in 3

strains of ours (0788, 0482, and 0604) in the region where the T6SS locus was expected. For

the rest of genomes (98/157; as 2 genomes carried the T6SS locus) not encoding T6SS

machinery, we could not identify the underlying mechanisms of loss due to draft-genomes

assemblies and possible crucial information missing. However, the absence of genetic cargo

expected to replace the T6SS locus could suggest either different way of T6SS locus

displacement or further genomic rearrangement, possibly leading to formation of further

subclades of *A. baumannii* ST19 lineage. Alternatively, the subclade with no detectable T6SS

locus displacement perhaps never encoded the T6SS in the first place.

Conclusion

A. baumannii ST19 has been in occurring in multiple spikes throughout last two decades,

especially correlating with war conflicts such as the conflict in Iraq and in Ukraine. The

genetic background of this lineage suggests its successful adaptation to the hospital

environment. The loss of T6SS in the (nearly) entire lineage of *A. baumannii* ST19 points

towards a beneficial evolutionary step, a fitness trade-off towards favouring environments

where antibiotic resistance is key for survival. The increasing prevalence of the *A. baumannii*

ST19 lineage and its antibiotic and non-antibiotic (environmental stresses and disinfectants)

resistance is highly concerning and warrants greater attention in surveillance efforts.

Material and Methods

Data Summary

The whole genome sequencing data were deposited in GenBank under following accession

numbers; strain 0788 - SAMN46706486, strain 0482 - SAMN48114201 and strain 0604 -

SAMN46706488.

Bacterial isolates, genomes and metadata

Three isolates of *A. baumannii* ST19 were obtained from two different patients (patient 1 -

wound, rectal swab; patient 2 - rectal swab). A further 156 genomes (including 97 from Luo

et al. (17)) of *A. baumannii* ST19 were retrieved from public repositories (GenBank and

BIGSdb) as of March 2025.

DNA extraction and whole genome sequencing

The DNA was extracted using DNeasy Blood & Tissue Kit (QIAGEN). A rapid barcoding kit

(V14) (ONT) was then used for library preparation which was then subjected to long-read

whole-genome sequencing using the R10.4.1 flowcell with the PromethION (P2 Solo)

platform by Nanopore.

Genome assembly and annotation

The long-read sequences were basecalled using the SUP model and adaptor and quality

($Q \leq 15$) trimmed using Porechop v0.2.2 (https://github.com/rrwick/Porechop) and

NanoFilt v2.8.0 (26), respectively. The reads were then assembled using Flye2.9.3 (27).

Phylogenetic analysis

The maximum-likelihood tree was constructed by Prokka (28) generated gff files used for the

core-genome alignment created by Roary (29). RAxML (30) was used for the calculation of

the phylogenetic tree under GTRGAMMA model of rate heterogeneity method supported by

500 bootstraps. The tree was visualized in iTOL (31).

Identification of resistance determinants

The resistance and virulence genes were detected using ABRicate

(https://github.com/tseemann/abricate) employing ResFinder 4.0 (32) and VFDB 2025 (33)

databases, respectively, with a 90% threshold for both gene identity and coverage. The typing

of capsule-encoding loci (KL) and lipooligosaccharide outer core loci (OCL) was determined

using Kaptive (34). The ABRicate (https://github.com/tseemann/abricate) was used to detect

the presence of plasmid replicons using AcinetobacrPlasmidTyping database

(https://github.com/MehradHamidian/AcinetobacterPlasmidTyping) (21), while the genes

encoding GIs*ul2*, chlorite dismutase, formaldehyde resistance genes, and T6SS-encoding

genes were detected using a custom database.

Conflict of interest

The authors declare that there are no conflicts of interest.

Ethical statement

The clinical isolates sequenced in this study were obtained within retrospective study

conducted under the ethical approval granted by the C. H. U. Brugmann CE 2023/51.

Acknowledgements

AV is a recipient of a junior postdoctoral fellowship of the Research Foundation – Flanders

(file number 1287223N). CVDH is supported by the Flanders Institute for Biotechnology

(VIB). KN was funded by the European Union under Horizon Europe: MSCA Actions

(Project 101105027) and by a junior postdoctoral fellowship from the Research Foundation -

Flanders (file number 1251624N). Sequencing was performed by VIB Nucleomics Core

(www.nucleomics.be).

References

1. Brek TM, Muhajir AA, Alkuwaity KK, Haddad MA, Alattas EM, Eisa ZM, Al-

Thaqafy MS, Albarraq AM, Al-Zahrani IA. Genomic insights of predominant

international high-risk clone ST2 *Acinetobacter baumannii* isolates in Saudi Arabia. J

Glob Antimicrob Resist. 2025 Mar 28;42:243-252. doi: 10.1016/j.jgar.2025.03.012.

Epub ahead of print. PMID: 40158887.

2. Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens.

Expert Rev Anti Infect Ther. 2013 Mar;11(3):297-308. doi: 10.1586/eri.13.12. PMID:

23458769.

3. WHO Bacterial Priority Pathogens List, 2024: bacterial pathogens of public health

importance to guide research, development and strategies to prevent and control

antimicrobial resistance. Geneva: World Health Organization; 2024. Licence: CC BY-

NC-SA 3.0 IGO.

4. Antochevis LC, Wilhelm CM, Arns B, Sganzerla D, Sudbrack LO, Nogueira TCRL,

Guzman RD, Martins AS, Cappa DS, Dos Santos ÂC, Pascual JC, Perugini VH,

Vespero EC, Rigatto MHP, Pereira DC, Lutz L, Leão RS, Marques EA, Henrique DM,

Coelho AAM, Frutuoso LL, de A Sousa EE, Abreu Guimarães LF, Ferreira ALP,

Castiñeiras AC, Alves MD, Telles JP, Yamada CH, de Almeida FP, Girão ES, de Sousa

PCP, de Melo AGND, Mendes ET, Rocha VFD, Neves EDS, Ribeiro MT, Starling

CEF, Oliveira MS, Sampaio JLM, Martins AF, Barth AL, Zavascki AP; ASCENSION

Study Group. World Health Organization priority antimicrobial resistance

in *Enterobacterales, Acinetobacter baumannii, Pseudomonas aeruginosa, Staphylococcus aureus* and *Enterococcus faecium* healthcare-associated bloodstream infections in Brazil (ASCENSION): a prospective, multicentre, observational study. Lancet Reg Health Am. 2025 Jan 30;43:101004. doi: 10.1016/j.lana.2025.101004. PMID: 39957800; PMCID: PMC11830303.

5. Hooppaw AJ, McGuffey JC, Di Venanzio G, Ortiz-Marquez JC, Weber BS, Lightly TJ, van Opijnen T, Scott NE, Cardona ST, Feldman MF. The Phenylacetic Acid Catabolic Pathway Regulates Antibiotic and Oxidative Stress Responses in *Acinetobacter*. mBio. 2022 Jun 28;13(3):e0186321. doi: 10.1128/mbio.01863-21. Epub 2022 Apr 25. PMID: 35467424; PMCID: PMC9239106.

6. Valcek A, Philippe C, Whiteway C, Robino E, Nesporova K, Bové M, Coenye T, De Pooter T, De Coster W, Strazisar M, Van der Henst C. Phenotypic Characterization and Heterogeneity among Modern Clinical Isolates of *Acinetobacter baumannii*. Microbiol Spectr. 2023 Feb 14;11(1):e0306122. doi: 10.1128/spectrum.03061-22. Epub 2022 Dec 8. PMID: 36475894; PMCID: PMC9927488.

7. Brovedan MA, Cameranesi MM, Limansky AS, Morán-Barrio J, Marchiaro P, Repizo GD. What do we know about plasmids carried by members of the *Acinetobacter* genus? World J Microbiol Biotechnol. 2020 Jul 13;36(8):109. doi: 10.1007/s11274-020-02890-7. PMID: 32656745.

8. Cain AK, Hamidian M. Portrait of a killer: Uncovering resistance mechanisms and global spread of *Acinetobacter baumannii*. PLoS Pathog. 2023 Aug 10;19(8):e1011520. doi: 10.1371/journal.ppat.1011520. PMID: 37561719; PMCID: PMC10414682.

9. Hamidian M, Nigro SJ. Emergence, molecular mechanisms and global spread of carbapenem-resistant *Acinetobacter baumannii*. Microb Genom. 2019

Oct;5(10):e000306. doi: 10.1099/mgen.0.000306. PMID: 31599224; PMCID: PMC6861865.

10. Sahl JW, Del Franco M, Pournaras S, Colman RE, Karah N, Dijkshoorn L, Zarrilli R. Phylogenetic and genomic diversity in isolates from the globally distributed *Acinetobacter baumannii* ST25 lineage. Sci Rep. 2015 Oct 14;5:15188. doi: 10.1038/srep15188. PMID: 26462752; PMCID: PMC4604477.

11. Tobin LA, Abu Sabah E, Lebreton F, Myers GSA, McGann PT, Hamidian M. Genomic analysis of early ST32 *Acinetobacter baumannii* strains recovered in US military treatment facilities reveals distinct lineages and links to the origins of the Tn*6168 ampC* transposon. J Antimicrob Chemother. 2025 Mar 3;80(3):666-675. doi: 10.1093/jac/dkae454. PMID: 39680383; PMCID: PMC11879209.

12. Harmer CJ, Cahill SM, Kenyon JJ, Hall RM. Aminoglycoside resistance genes in early members of the *Acinetobacter baumannii* ST78A (SMAL, Italian clone) reside in an IS26-bounded island in the chromosome. J Antimicrob Chemother. 2024 May 2;79(5):1014-1018. doi: 10.1093/jac/dkae064. PMID: 38530861; PMCID: PMC11062947.

13. Gao Y, Li H, Chen H, Zhang J, Wang R, Wang Z, Wang H. Origin, Phylogeny, and Transmission of the Epidemic Clone ST208 of Carbapenem-Resistant *Acinetobacter baumannii* on a Global Scale. Microbiol Spectr. 2022 Jun 29;10(3):e0260421. doi: 10.1128/spectrum.02604-21. Epub 2022 May 31. PMID: 35638783; PMCID: PMC9241911.

14. Di Venanzio G, Moon KH, Weber BS, Lopez J, Ly PM, Potter RF, Dantas G, Feldman MF. Multidrug-resistant plasmids repress chromosomally encoded T6SS to enable their dissemination. Proc Natl Acad Sci U S A. 2019 Jan 22;116(4):1378-1383. doi:

10.1073/pnas.1812557116. Epub 2019 Jan 9. PMID: 30626645; PMCID: PMC6347727.

15. Weber BS, Kinsella RL, Harding CM, Feldman MF. The Secrets of *Acinetobacter* Secretion. Trends Microbiol. 2017 Jul;25(7):532-545. doi: 10.1016/j.tim.2017.01.005. Epub 2017 Feb 16. PMID: 28216293; PMCID: PMC5474180.

16. Repizo GD, Gagné S, Foucault-Grunenwald ML, Borges V, Charpentier X, Limansky AS, Gomes JP, Viale AM, Salcedo SP. Differential Role of the T6SS in *Acinetobacter baumannii* Virulence. PLoS One. 2015 Sep 24;10(9):e0138265. doi: 10.1371/journal.pone.0138265. PMID: 26401654; PMCID: PMC4581634.

17. Luo TL, Martin MJ, Kovalchuk V, Kondratiuk V, Trapaidze N, Metreveli M, Hulseberg CE, Dao HD, Kwak YI, Maybank R, Musich TA, Scherer MR, Bennett JW, Mc Gann PT, Lebreton F. Detection of carbapenemase producing *Acinetobacter baumannii* ST19 from Georgia and Ukraine carrying bla_{OXA-23}, bla_{OXA-72}, and/or bla_{NDM-5}, December 2019 to June 2023. Euro Surveill. 2024 Jun;29(24):2400259. doi: 10.2807/1560-7917.ES.2024.29.24.2400259. PMID: 38873796; PMCID: PMC11177571.

18. Boyd D, Taylor G, Fuller J, Bryce E, Embree J, Gravel D, Katz K, Kibsey P, Kuhn M, Langley J, Mataseje L, Mitchell R, Roscoe D, Simor A, Thomas E, Turgeon N, Mulvey M; Canadian Nosocomial Infection Surveillance Program. Complete Sequence of Four Multidrug-Resistant MOBQ1 Plasmids Harboring bla_{GES-5} Isolated from *Escherichia coli* and *Serratia marcescens* Persisting in a Hospital in Canada. Microb Drug Resist. 2015 Jun;21(3):253-60. doi: 10.1089/mdr.2014.0205. Epub 2014 Dec 29. PMID: 25545311.

19. Chalmers R, Sewitz S, Lipkow K, Crellin P. Complete nucleotide sequence of Tn*10*. J

Bacteriol. 2000 May;182(10):2970-2. doi: 10.1128/JB.182.10.2970-2972.2000.

PMID: 10781570; PMCID: PMC102010.

20. Shaffer JMC, Giddings LA, Samples RM, Mikucki JA. Genomic and phenotypic

characterization of a red-pigmented strain of *Massilia frigida* isolated from an

Antarctic microbial mat. Front Microbiol. 2023 May 12;14:1156033. doi:

10.3389/fmicb.2023.1156033. PMID: 37250028; PMCID: PMC10213415.

21. Hamidian M, Hall RM. *Acinetobacter baumannii* ATCC 19606 Carries GI*sul2* in a

Genomic Island Located in the Chromosome. Antimicrob Agents Chemother. 2016

Dec 27;61(1):e01991-16. doi: 10.1128/AAC.01991-16. PMID: 27795382; PMCID:

PMC5192138.

22. Lam MMC, Koong J, Holt KE, Hall RM, Hamidian M. Detection and Typing of

Plasmids in Acinetobacter baumannii Using *rep* Genes Encoding Replication

Initiation Proteins. Microbiol Spectr. 2023 Feb 14;11(1):e0247822. doi:

10.1128/spectrum.02478-22. Epub 2022 Dec 6. PMID: 36472426; PMCID:

PMC9927589.

23. Weber BS, Ly PM, Irwin JN, Pukatzki S, Feldman MF. A multidrug resistance

plasmid contains the molecular switch for type VI secretion in *Acinetobacter*

baumannii. Proc Natl Acad Sci U S A. 2015 Jul 28;112(30):9442-7. doi:

10.1073/pnas.1502966112. Epub 2015 Jul 13. PMID: 26170289; PMCID:

PMC4522760.

24. Alton NK, Vapnek D. Nucleotide sequence analysis of the chloramphenicol resistance

transposon Tn*9*. Nature. 1979 Dec 20-27;282(5741):864-9. doi: 10.1038/282864a0.

PMID: 390403.

25. Echenique JR, Dorsey CW, Patrito LC, Petroni A, Tolmasky ME, Actis LA. *Acinetobacter baumannii* has two genes encoding glutathione-dependent formaldehyde dehydrogenase: evidence for differential regulation in response to iron. Microbiology (Reading). 2001 Oct;147(Pt 10):2805-2815. doi: 10.1099/00221287-147-10-2805. PMID: 11577159.

26. De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack: visualizing and processing long-read sequencing data. Bioinformatics. 2018 Aug 1;34(15):2666-2669. doi: 10.1093/bioinformatics/bty149. PMID: 29547981; PMCID: PMC6061794.

27. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat graphs. Nat Biotechnol. 2019 May;37(5):540-546. doi: 10.1038/s41587-019-0072-8. Epub 2019 Apr 1. PMID: 30936562.

28. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014 Jul 15;30(14):2068-9. doi: 10.1093/bioinformatics/btu153. Epub 2014 Mar 18. PMID: 24642063.

29. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D, Keane JA, Parkhill J. Roary: rapid large-scale prokaryote pan genome analysis. Bioinformatics. 2015 Nov 15;31(22):3691-3. doi: 10.1093/bioinformatics/btv421. Epub 2015 Jul 20. PMID: 26198102; PMCID: PMC4817141.

30. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics. 2014 May 1;30(9):1312-3. doi: 10.1093/bioinformatics/btu033. Epub 2014 Jan 21. PMID: 24451623; PMCID: PMC3998144.

31. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic

tree display and annotation. Nucleic Acids Res. 2021 Jul 2;49(W1):W293-W296. doi:

10.1093/nar/gkab301. PMID: 33885785; PMCID: PMC8265157.

32. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A,

Allesoe RL, Rebelo AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B,

Werner G, Bender JK, Stingl K, Nguyen M, Coppens J, Xavier BB, Malhotra-Kumar

S, Westh H, Pinholt M, Anjum MF, Duggett NA, Kempf I, Nykäsenoja S, Olkkola S,

Wieczorek K, Amaro A, Clemente L, Mossong J, Losch S, Ragimbeau C, Lund O,

Aarestrup FM. ResFinder 4.0 for predictions of phenotypes from genotypes. J

Antimicrob Chemother. 2020 Dec 1;75(12):3491-3500. doi: 10.1093/jac/dkaa345.

PMID: 32780112; PMCID: PMC7662176.

33. Zhou S, Liu B, Zheng D, Chen L, Yang J. VFDB 2025: an integrated resource for

exploring anti-virulence compounds. Nucleic Acids Res. 2025 Jan 6;53(D1):D871-

D877. doi: 10.1093/nar/gkae968. PMID: 39470738; PMCID: PMC11701737.

34. Wyres KL, Cahill SM, Holt KE, Hall RM, Kenyon JJ. Identification of *Acinetobacter*

baumannii loci for capsular polysaccharide (KL) and lipooligosaccharide outer core

(OCL) synthesis in genome assemblies using curated reference databases compatible

with *Kaptive*. Microb Genom. 2020 Mar;6(3):e000339. doi: 10.1099/mgen.0.000339.

PMID: 32118530; PMCID: PMC7200062.

Tree scale: 0.001

443   Figure 2: Phylogenetic analysis of 3 isolates and further 156 of *A. baumannii* ST19 genomes, depicting year of isolation, country, capsule locus

444   type (KL), outer core lipooligosaccharide type (OCL), and presence of GI*sul2* genetic island, chlorite dismutase, T6SS, *frmRAB* operon, plasmid

445   replicons, resistance, and virulence genes, respectively.

446