Title: The absence of the Type VI Secretion System in the successful

lineage of *Acinetobacter baumannii* ST19

Running title: Loss of Type VI Secretion System in *A. baumannii* ST19

Authors: Adam Valcek[1, 2*], Diana Isabela Costescu Strachinaru[3], Kristina Nesporova[4],

Patrick Soentjens[3,5] Anke Stoefs[6,7], Charles Van der Henst[1, 2*]

Affiliations:

1. Microbial Resistance and Drug Discovery, VIB-VUB Center for Structural Biology, VIB,

Flanders Institute for Biotechnology, Brussels, Belgium

2. Structural Biology Brussels, Vrije Universiteit Brussel (VUB), Brussels, Belgium

3. Center for Infectious Diseases, Queen Astrid Military Hospital, Brussels, Belgium

4. Microbial Systems Cell Biology Group, Department of Biology, KU Leuven, Leuven,

Belgium

5. Department of Clinical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

6. Department of Microbiology, Queen Astrid Military Hospital, Brussels, Belgium

7. Department of Microbiology, Universitair Ziekenhuis Brussels, Brussels Belgium

* These authors contributed equally to this work and share the last authorship.

Corresponding authors: Adam Valcek (*adam.valcek@vub.be),* Charles Van der Henst

(*charles.vanderhenst@vub.vib.be*)

Abstract

Acinetobacter baumannii is an opportunistic pathogen, often multi- to pandrug-resistant, including to last-resort antibiotics such as carbapenems. *A. baumannii* can acquire DNA through multiple mechanisms including natural competence and conjugation. An active Type VI Secretion System (T6SS), which kills nonkin bacteria and challenges and inhibits the conjugation. The acquisition of resistance genes has helped the spread of resistant clones of *A. baumannii*, such as those of Sequence Types (ST^Pas), ST1, ST2, ST10, ST15, ST25, ST79 and ST85. While ST1 and ST2 were thoroughly studied, there are new emerging clones whose genetic background facilitating the rise is still unknown or poorly understood.

The emergence of *Acinetobacter baumannii* ST19 first peaked in 2003, coinciding with the Iraq conflict and increased detection among U.S. military personnel. Its prevalence later surged in Georgia and Ukraine, around the onset of the conflict in Ukraine in 2022, associating it with regions affected by military activity.

In this study, we have sequenced whole-genome three isolates of *A. baumannii* ST19 and compared them with further 156 *A. baumannii* ST19 genomes were obtained from public repositories. In total, 157/159 genomes revealed loss of T6SS locus, which was replaced in 56/157 genomes (35,7%) by ΔTn*9* carrying chloramphenicol and ΔTn*10* carrying tetracycline resistance genes, and formaldehyde and chlorite resistance genes. Surprisingly, the antibiotic resistance-encoding transposons likely originated from Enterobacteriaceae plasmids. The loss of a functional T6SS in *A. baumannii* ST19 may potentially facilitate horizontal gene transfer and promoting a cooperative or less competitive lifestyle providing a selective advantage at the population level.

Introduction

Acinetobacter baumannii is an emerging opportunistic nosocomial pathogen that causes a variety of infections including urinary tract infections, wound infections, meningitis, ventilator-associated pneumonia, blood-stream infection, and endocarditis (1). *A. baumannii* is a member of ESKAPE pathogens (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacter* spp.) (2), and due to its frequent resistance to last-line antibiotics such as carbapenems, it has been designated as a critical priority in the list of pathogens of importance for studying and developing new antibiotics by the World Health Organization's list (3, 4).

Besides conventional antibiotics, *A. baumannii* can withstand challenging environmental conditions such as oxidative stress, prolonged periods of desiccation, human serum but also disinfectants (5).

The ability of *A. baumannii* to acquire exogenous DNA via natural competence (6) and conjugation (7) contributes to high plasticity of its genome (6). The horizontal gene transfer plays significant role in the evolution of *A. baumannii* lineages, resulting in number of highly successful clonal lineages such as ST1, ST2, ST10, ST15, ST25, ST79 and ST85 (Pasteur scheme) (8, 9). While ST1 and ST2 are responsible for the majority of infections caused by extensively-drug resistant *A. baumannii*, some clones are specific to certain geographical locations (9) , such as ST25 (Europe, United Arab Emirates, South America, Asia) (10) , ST32 (South Korea, Europe, the USA, Canada and the Middle East) (11) or ST78 (Italy, Ukraine, USA) (12).

While the horizontal gene transfer, especially plasmid transfer via conjugation, can shape

entire lineages and their success (13), an active Type VI secretion system (T6SS) poses a

unique challenge to conjugative plasmids, as *Acinetobacter* plasmid donors and recipients

may kill each other (14). The T6SS machinery is usually encoded in a single chromosomal

locus (15). The T6SS of *A. baumannii* has been shown to mediate the killing of *E. coli*, *K.*

pneumoniae and *P. aeruginosa*, while different strains of *A. baumannii* are also able to kill

each other (15, 16).

In this study, we have assessed genetic determinants disrupting the T6SS-encoding locus in a

rising lineage of *A. baumannii* ST19, of which occurrence is increasing regionally (Ukraine,

Georgia) (17). Furthermore, we have performed comparative genomic analysis of the global

population of *A. baumannii* ST19 in order to decipher genomic traits that make this lineage

emergingly successful and to assess its threat in becoming a world-wide disseminated clone.

Results

A. baumannii ST19 strains and genomes

A concerning report on the increasing occurrence of *Acinetobacter baumannii* ST19 encoding

carbapenemases was recently published (17), raising questions about factors contributing the

success of this lineage. Three isolates of *A. baumannii* ST19 were obtained from two

different patients (patient 1 - wound, rectal swab; patient 2 - rectal swab) and long-read

whole-genome sequenced and *de novo* assembled.

A further 156 genomes (including 97 from Luo et al. (17)) of *A. baumannii* ST19 were

retrieved from public repositories (GenBank and BIGSdb). The occurrence of *A. baumannii*

ST19 appears to spike during specific time periods that correlate with military conflicts

(Figure 1), such as the Iraq War (2003) and the war in Ukraine (2022).

Figure 1: Temporal and spatial occurrence of *A. baumannii* ST19 based on genomes
deposited in public repositories and available metadata.

The examined genomes carried a variety of virulence and resistance genes (Supplementary
Figure 1), including resistance to last-resort antibiotics such as carbapenems. One-third
(n=53) out of 159 genomes described in this study encoded at least one carbapenemase gene
(bla_{OXA-23}, bla_{OXA-72}, bla_{NDM-1} or bla_{NDM-5}) with 6 encoding two and 9 encoding three
carbapenemases. Furthermore, a broad variety of plasmid replicons was detected
(Supplementary Figure 1), suggesting an easy acquisition of genetic material by the
horizontal gene transfer.

Genomic analysis suggests that the possible underlying reason for its success to be the
acquisition of multiple partial transposons, including ΔTn*9* (18) carrying chloramphenicol
resistance gene and ΔTn*10* (19) carrying the tetracycline resistance gene. Interestingly,

61/159 (38,4 %) genomes of *A. baumannii* ST19 carried chlorite dismutase and formaldehyde

resistance (*frmRAB*) genes, with 98,4% amino acid identity to *Massilia glaciei* and 96,7% to

Ideonella dechloratans, where it combats oxidative stress (20). To our knowledge, this is the

first time chlorite dismutase has been reported in the genome of *A. baumannii*.

Furthermore, these transposons disrupted and replaced the locus encoding the T6SS. The

entire region of T6SS deletion exhibits extensive genetic rearrangements, as ΔTn*9* and

ΔTn*10* were acquired and disrupted by IS*26* and IS*4321,* and IS*26*, respectively.

The absence of a functional T6SS, likely enabling conjugal acquisition of genetic material

(14) such as antibiotic, formaldehyde and chlorite resistance, could represent one of the

reasons behind their successful rise.

Genetic variability of the *A. baumannii* ST19 lineage

The whole genome sequences were subjected to genotyping [sequence type, capsule locus

(KL) type, outer core lipooligosaccharide locus (OCL) type, antimicrobial resistance, and

virulence genes]. A phylogenetic analysis was performed on these isolates, including

genomes of the same ST from public repositories (GenBank and BIGSdb) (Figure 2,

Supplementary Figure 1).

Figure 2: Phylogenetic analysis of 3 isolates and further 156 of *A. baumannii* ST19 genomes,

depicting year of isolation, country, capsule locus type (KL), outer core lipooligosaccharide

type (OCL), and presence of GI*sul2* genetic island, chlorite dismutase, T6SS, *frmRAB* genes,

respectively.

While various MLST (Multilocus Sequence Type) ST (Pasteur scheme) such as ST2 or ST1

often belong to various KL types (6), the ST19 lineage exhibited limited variability with

69/159 genomes being KL91, 66/159 KL15, followed by KL1 (13/159), KL147 (8/159),

KL17 (2/159) and KL40 (1/159). Interestingly, 53/55 OCL1 encoded KL15 (2/55 OCL1 were

KL17), the most predominant OCL type, OCL5 (n=87) was variable and encompassed KL91,

KL1, KL147 and KL15. Furthermore, all of the genomes of KL15/OCL1 formed one large,

genetically related cluster (Figure 2).

The vast majority (157/159) of *A. baumannii* ST19 harbored GI*sul2,* a chromosomally carried

Genomic Island encoding sulphonamide resistance determinant (*sul2*) along with putative

arsenic resistance genes (21). We have detected a range of 0 to 6 plasmid replicons (22) in the

genomes of *A. baumannii* ST19, with rP-T1 replicon being the most prevalent (n=122/159),

followed by r3-T23 (n=46/159), r3-T12 and r1-T8 (n=44/159 both).

Interestingly, 18 isolates did not carry any detectable plasmid replicon raising a question

whether there is another mechanism limiting acquisition of plasmids or whether they have not

encountered a conjugative plasmid yet. However, the *Acinetobacter* typing schemes are still

being developed (22), possibly undetected (or below the detection threshold) plasmids must

be considered as well.

By evaluating the T6SS locus in the ST19 lineage of *A. baumannii*, we identified variability

in genetic origin of the deletion of these genes. Of 61/159 genomes carrying chlorite

dismutase and formaldehyde resistance genes in the locus of T6SS, 55/61 harbored ΔTn*10*

carrying the tetracycline and ΔTn*9* carrying chloramphenicol resistance genes (Figure 3).

Five formaldehyde resistance and chlorite dismutase-encoding genomes (5/61) did not harbor

ΔTn*10* carrying the tetracycline resistance and 1/61 did not harbor ΔTn*9* carrying

chloramphenicol resistance genes. However, mapping on the reference genome (0604 -

SAMN46706488) confirmed, that in all these genomes (61/159) these genetic rearrangements

displaced locus encoding T6SS.

Figure 3: Depiction of the T6SS locus disruption of *A. baumannii* ST19 (0482 -
SAMN48114201) as a representative genome, compared to AB5075-VUB (CP070362)

showing deletion mediated by ΔTn*9*, chlorite dismutase, formaldehyde resistance locus

frmRAB, and ΔTn*10*. The alignments are based on amino acid identity.

Interestingly, 2/159 genomes of *A. baumannii* ST19 (GCA_042643835.1 and

GCA_040868255.1) still encoded a likely functional T6SS. However, compared to AB5075-

VUB (accession nr. CP070362), a number of SNPs (99% pairwise identity over 19417 bp)

was detected in these two genomes, while the broader genetic neighbourhood only

sporadically exhibited SNPs. The genomes GCA_042643835.1 and GCA_040868255.1

belonged to KL40/OCL2 and KL17/OCL1, respectively, which are rare for this lineage (a

sole KL40 and one of two KL17, respectively). While none of the SNPs were non-sense, the

influence of missense or synonymous (e.g., via tRNA) on the functionality of T6SS remains

to be explored.

The loss of the T6SS in non-ST19 lineages of *A. baumannii*

The presence of the T6SS locus was assessed in 8218 genomes of *A. baumannii* available in

BIGSdb. The lack of the T6SS was detected in 1287 genomes (including 159 genomes of *A.*

baumannii ST19). The most dominant lineage of *A. baumannii* lacking the T6SS in this

comparison was ST2 (n=435; 33,7%) followed by ST19 (n=159; 12,3%) and ST94 (n=82;

6,4%) (Supplementary Figure 3AB).

The phylogenetic comparison of all 1287 (Supplementary Figure 2) revealed clustering of all

A. baumannii ST19 and 103 non-ST19 genomes together. The non-ST19 genomes in the

ST19 cluster consisted of 93 single-locus variant (SLV) sequence types including one novel

ST, and 10 double-locus variant (DLV) sequence types including one novel ST

(Supplementary Figure 3C).

The presence of *frmRAB*, and chlorite dismutase-encoding genes was detected predominantly

in *A. baumannii* ST19, but also in two *A. baumannii* ST315 genomes within the same cluster.

A single genome (ST2525) outside of the ST19 cluster was found to carry the *frmRAB* genes.

Discussion

Acinetobacter has the core components of the T6SS conserved and found in a single

chromosomal locus (15). The *A. baumannii* strains use Type VI Secretion System (T6SS) to

kill nonkin *A. baumannii* as well as other bacterial species such as *E. coli*, *K. pneumoniae* and

P. aeruginosa, (16, 23). The active T6SS can be a challenge to conjugation and horizontal

gene transmission (15). The absence of T6SS could therefore allow co-existence of nonkin

bacterial species and strains. We hypothesize that this might and explain original acquisition

of genetic material from these species and possibly evolution of the *A. baumannii* ST19

lineage.

The Tn*10* was originally identified and described in *Shigella flexneri* (19). This corresponds

to the best match of the ΔTn*10* nucleotide sequence in this study was detected for

Salmonella. enterica serovar Typhi and *S. flexneri*. The ΔTn*10* observed in this study has the

highest similarity to regions from *Enterobacter hormaechei* IncHI2 plasmids (>99%

nucleotide identity and 97% query coverage). Furthermore, the ΔTn*9* (24) examined in this

study was BLASTn-detected (with >99% nucleotide identity and 100% query coverage) in

Klebsiella pneumoniae IncF plasmids.

Although *adhC* genes facilitating resistance to formaldehyde which are native to

Acinetobacter have been reported (25), we have identified locus, likely originating from an

Enterobacter hormaechei IncHI2 plasmid, or a *Klebsiella pneumoniae* (or *quasipneumoniae*)

IncF plasmid (95% nucleotide identity and 100% coverage). However, the origin of the

chlorite dismutase gene remains undeciphered, while the BLASTn of the chlorite dismutase

gene observed in this study results in identical hits (>96% nucleotide identity and 98% query

coverage) in various species, such as *P. aeruginosa* (chromosome), *S. enterica* (IncHI2

plasmid), *K. pneumoniae* (IncF plasmid), *E. hormaechei* (chromosome), but also *Vibrio*

rumoiensis (plasmid) *Raoultella ornithinolytica* (IncF plasmid), and *Leclercia*

adecarboxylata (IncHI2 plasmid).

The total of 61/159 of genomes were identified to encode the same genetic cargo (ΔTn*9,*

ΔTn*10*, chlorite dismutase and the formaldehyde resistance cluster) as the one present in 3

strains of ours (0788, 0482, and 0604) in the region where the T6SS locus was expected. For

the rest of genomes (98/157; as 2 genomes carried the T6SS locus) not encoding T6SS

machinery, we could not identify the underlying mechanisms of loss due to draft-genome

assemblies and possible crucial information missing. However, the absence of genetic cargo

expected to replace the T6SS locus could suggest either a different way of T6SS locus

displacement or further genomic rearrangement, possibly leading to formation of further

subclades of the *A. baumannii* ST19 lineage. Alternatively, the subclade with no detectable

T6SS locus displacement perhaps never encoded the T6SS in the first place.

From the total of 1287 genomes, 179 different sequence types of *A. baumannii* lacking the

T6SS were detected. Furthermore, 35 genomes were typed as novel ST, with at least one

novel allele, and therefore some ST might be repetitive. While additional non-ST19 *A.*

baumannii lineages were detected that lack the T6SS, the specific displacement and

acquisition of the ΔTn*9*, ΔTn*10*, *frmRAB*, and chlorite dismutase gene combination was

observed only in *A. baumannii* ST19. The lineages *A. baumannii* ST19 (except for two

genomes) and ST94 (SLV of ST19) globally lacked the T6SS. While the most genomes

lacking the T6SS were detected from *A. baumannii* ST2 lineage, this is skewed by the

overrepresentation of this lineage in the dataset from BIGSdb (n=6402) and the T6SS might

have been replaced or lost, as observed in other cases of clinical isolates lacking some or all

T6SS genes (35, 36).

The *A. baumannii* lineage ST94 (SLV of ST19) completely lacked the T6SS, however no

frmRAB or chlorite dismutase genes were detected in this lineage. Unfortunately, the lack of

complete chromosomal sequences of *A. baumannii* ST94 lineage from this dataset hinders

resolution of the nature of the T6SS loss.

The loss of the T6SS in entire lineages of *A. baumannii* could also be used as an

epidemiological marker. While there were reports of sporadic loss of part or the entire T6SS

locus in *A. baumannii* clinical isolates (37, 38), the lineage-wide loss could be an indicator

for the detection of a difficult-to-treat lineage, widely resistant to disinfectants and

antibiotics, with high persistence potential.

Conclusion

A. baumannii ST19 has been in occurring in multiple spikes in prevalence throughout last

two decades, especially correlating with war conflicts such as the conflict in Iraq and in

Ukraine. The genetic background of this lineage suggests its successful adaptation to the

hospital environment. The loss of T6SS in the (nearly) entire lineage of *A. baumannii* ST19

points towards a beneficial evolutionary step, a fitness trade-off towards favouring

environments where antibiotic resistance is key for survival. The increasing prevalence of the

A. baumannii ST19 lineage and its antibiotic and non-antibiotic (environmental stresses and

disinfectants) resistance is highly concerning and warrants greater attention in surveillance

efforts.

Material and Methods

Data Summary

The whole genome sequencing data were deposited in GenBank under following accession

numbers; strain 0788 - SAMN46706486, strain 0482 - SAMN48114201 and strain 0604 -

SAMN46706488.

Bacterial isolates, genomes and metadata

Three isolates of *A. baumannii* ST19 were obtained from two different patients (patient 1 -

wound, rectal swab; patient 2 - rectal swab). A further 156 genomes (including 97 from Luo

et al. (17)) of *A. baumannii* ST19 were retrieved from public repositories (GenBank and

BIGSdb) as of November 2025. In order to analyse other lineages of *A. baumnannii* for T6SS

locus deletion, all available genomes o *A. baumannii* in BIGSdb (n=8218) were downloaded

and screened.

DNA extraction and whole genome sequencing

The DNA was extracted using DNeasy Blood & Tissue Kit (QIAGEN). A rapid barcoding kit

(V14) (ONT) was then used for library preparation which was then subjected to long-read

whole-genome sequencing using the R10.4.1 flowcell with the PromethION (P2 Solo)

platform by Nanopore.

Genome assembly and annotation

The long-read sequences were basecalled using the SUP model and adaptor and quality

($Q \leq 15$) trimmed using Porechop v0.2.2 (https://github.com/rrwick/Porechop) and

NanoFilt v2.8.0 (26), respectively. The reads were then assembled using Flye2.9.3 (27).

Phylogenetic analysis

The maximum-likelihood trees were constructed by Prokka (28) generated gff files used for

the core-genome alignment created by Roary (29). RAxML (30) was used for the calculation

of the phylogenetic tree under GTRGAMMA model of rate heterogeneity method supported

by 500 bootstraps. The tree was visualized in iTOL (31).

Identification of resistance determinants

The resistance and virulence genes were detected using ABRicate

(https://github.com/tseemann/abricate) employing ResFinder 4.0 (32) and VFDB 2025 (33)

databases, respectively, with a 90% threshold for both gene identity and coverage. The typing

of capsule-encoding loci (KL) and lipooligosaccharide outer core loci (OCL) was determined

using Kaptive (34). The ABRicate (https://github.com/tseemann/abricate) was used to detect

the presence of plasmid replicons using AcinetobacrPlasmidTyping database

(https://github.com/MehradHamidian/AcinetobacterPlasmidTyping) (21), while the genes

encoding GIs*ul2*, chlorite dismutase, formaldehyde resistance genes, and T6SS-encoding

genes were detected using a custom database.

Conflict of interest

The authors declare that there are no conflicts of interest.

Ethical statement

The clinical isolates sequenced in this study were obtained within retrospective study

conducted under the ethical approval granted by the C. H. U. Brugmann CE 2023/51.

Acknowledgements

AV is a recipient of a junior postdoctoral fellowship of the Research Foundation – Flanders

(file number 1287223N). CVDH is supported by the Flanders Institute for Biotechnology

(VIB). KN was funded by the European Union under Horizon Europe: MSCA Actions

(Project 101105027) and by a junior postdoctoral fellowship from the Research Foundation -

Flanders (file number 1251624N). Sequencing was performed by VIB Nucleomics Core

(www.nucleomics.be).

322 **References**

1. Brek TM, Muhajir AA, Alkuwaity KK, Haddad MA, Alattas EM, Eisa ZM, Al-

Thaqafy MS, Albarraq AM, Al-Zahrani IA. Genomic insights of predominant

international high-risk clone ST2 *Acinetobacter baumannii* isolates in Saudi Arabia. J

Glob Antimicrob Resist. 2025 Mar 28;42:243-252. doi: 10.1016/j.jgar.2025.03.012.

Epub ahead of print. PMID: 40158887.

2. Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens.

Expert Rev Anti Infect Ther. 2013 Mar;11(3):297-308. doi: 10.1586/eri.13.12. PMID:

23458769.

3. WHO Bacterial Priority Pathogens List, 2024: bacterial pathogens of public health

importance to guide research, development and strategies to prevent and control

antimicrobial resistance. Geneva: World Health Organization; 2024. Licence: CC BY-

NC-SA 3.0 IGO.

4. Antochevis LC, Wilhelm CM, Arns B, Sganzerla D, Sudbrack LO, Nogueira TCRL,

Guzman RD, Martins AS, Cappa DS, Dos Santos ÂC, Pascual JC, Perugini VH,

Vespero EC, Rigatto MHP, Pereira DC, Lutz L, Leão RS, Marques EA, Henrique DM,

Coelho AAM, Frutuoso LL, de A Sousa EE, Abreu Guimarães LF, Ferreira ALP,

Castiñeiras AC, Alves MD, Telles JP, Yamada CH, de Almeida FP, Girão ES, de Sousa

PCP, de Melo AGND, Mendes ET, Rocha VFD, Neves EDS, Ribeiro MT, Starling CEF, Oliveira MS, Sampaio JLM, Martins AF, Barth AL, Zavascki AP; ASCENSION Study Group. World Health Organization priority antimicrobial resistance in *Enterobacterales, Acinetobacter baumannii, Pseudomonas aeruginosa, Staphylococcus aureus* and *Enterococcus faecium* healthcare-associated bloodstream infections in Brazil (ASCENSION): a prospective, multicentre, observational study. Lancet Reg Health Am. 2025 Jan 30;43:101004. doi: 10.1016/j.lana.2025.101004. PMID: 39957800; PMCID: PMC11830303.

5. Hooppaw AJ, McGuffey JC, Di Venanzio G, Ortiz-Marquez JC, Weber BS, Lightly TJ, van Opijnen T, Scott NE, Cardona ST, Feldman MF. The Phenylacetic Acid Catabolic Pathway Regulates Antibiotic and Oxidative Stress Responses in *Acinetobacter*. mBio. 2022 Jun 28;13(3):e0186321. doi: 10.1128/mbio.01863-21. Epub 2022 Apr 25. PMID: 35467424; PMCID: PMC9239106.

6. Valcek A, Philippe C, Whiteway C, Robino E, Nesporova K, Bové M, Coenye T, De Pooter T, De Coster W, Strazisar M, Van der Henst C. Phenotypic Characterization and Heterogeneity among Modern Clinical Isolates of *Acinetobacter baumannii*. Microbiol Spectr. 2023 Feb 14;11(1):e0306122. doi: 10.1128/spectrum.03061-22. Epub 2022 Dec 8. PMID: 36475894; PMCID: PMC9927488.

7. Brovedan MA, Cameranesi MM, Limansky AS, Morán-Barrio J, Marchiaro P, Repizo GD. What do we know about plasmids carried by members of the *Acinetobacter* genus? World J Microbiol Biotechnol. 2020 Jul 13;36(8):109. doi: 10.1007/s11274-020-02890-7. PMID: 32656745.

8. Cain AK, Hamidian M. Portrait of a killer: Uncovering resistance mechanisms and global spread of *Acinetobacter baumannii*. PLoS Pathog. 2023 Aug

10;19(8):e1011520. doi: 10.1371/journal.ppat.1011520. PMID: 37561719; PMCID:

PMC10414682.

9. Hamidian M, Nigro SJ. Emergence, molecular mechanisms and global spread of

carbapenem-resistant *Acinetobacter baumannii*. Microb Genom. 2019

Oct;5(10):e000306. doi: 10.1099/mgen.0.000306. PMID: 31599224; PMCID:

PMC6861865.

10. Sahl JW, Del Franco M, Pournaras S, Colman RE, Karah N, Dijkshoorn L, Zarrilli R.

Phylogenetic and genomic diversity in isolates from the globally distributed

Acinetobacter baumannii ST25 lineage. Sci Rep. 2015 Oct 14;5:15188. doi:

10.1038/srep15188. PMID: 26462752; PMCID: PMC4604477.

11. Tobin LA, Abu Sabah E, Lebreton F, Myers GSA, McGann PT, Hamidian M.

Genomic analysis of early ST32 *Acinetobacter baumannii* strains recovered in US

military treatment facilities reveals distinct lineages and links to the origins of the

Tn*6168 ampC* transposon. J Antimicrob Chemother. 2025 Mar 3;80(3):666-675. doi:

10.1093/jac/dkae454. PMID: 39680383; PMCID: PMC11879209.

12. Harmer CJ, Cahill SM, Kenyon JJ, Hall RM. Aminoglycoside resistance genes in

early members of the *Acinetobacter baumannii* ST78A (SMAL, Italian clone) reside

in an IS26-bounded island in the chromosome. J Antimicrob Chemother. 2024 May

2;79(5):1014-1018. doi: 10.1093/jac/dkae064. PMID: 38530861; PMCID:

PMC11062947.

13. Gao Y, Li H, Chen H, Zhang J, Wang R, Wang Z, Wang H. Origin, Phylogeny, and

Transmission of the Epidemic Clone ST208 of Carbapenem-Resistant *Acinetobacter*

baumannii on a Global Scale. Microbiol Spectr. 2022 Jun 29;10(3):e0260421. doi:

10.1128/spectrum.02604-21. Epub 2022 May 31. PMID: 35638783; PMCID:

PMC9241911.

14. Di Venanzio G, Moon KH, Weber BS, Lopez J, Ly PM, Potter RF, Dantas G, Feldman

MF. Multidrug-resistant plasmids repress chromosomally encoded T6SS to enable

their dissemination. Proc Natl Acad Sci U S A. 2019 Jan 22;116(4):1378-1383. doi:

10.1073/pnas.1812557116. Epub 2019 Jan 9. PMID: 30626645; PMCID:

PMC6347727.

15. Weber BS, Kinsella RL, Harding CM, Feldman MF. The Secrets of *Acinetobacter*

Secretion. Trends Microbiol. 2017 Jul;25(7):532-545. doi: 10.1016/j.tim.2017.01.005.

Epub 2017 Feb 16. PMID: 28216293; PMCID: PMC5474180.

16. Repizo GD, Gagné S, Foucault-Grunenwald ML, Borges V, Charpentier X, Limansky

AS, Gomes JP, Viale AM, Salcedo SP. Differential Role of the T6SS in *Acinetobacter*

baumannii Virulence. PLoS One. 2015 Sep 24;10(9):e0138265. doi:

10.1371/journal.pone.0138265. PMID: 26401654; PMCID: PMC4581634.

17. Luo TL, Martin MJ, Kovalchuk V, Kondratiuk V, Trapaidze N, Metreveli M,

Hulseberg CE, Dao HD, Kwak YI, Maybank R, Musich TA, Scherer MR, Bennett JW,

Mc Gann PT, Lebreton F. Detection of carbapenemase producing *Acinetobacter*

baumannii ST19 from Georgia and Ukraine carrying bla_{OXA-23}, bla_{OXA-72},

and/or bla_{NDM-5}, December 2019 to June 2023. Euro Surveill. 2024

Jun;29(24):2400259. doi: 10.2807/1560-7917.ES.2024.29.24.2400259. PMID:

38873796; PMCID: PMC11177571.

18. Boyd D, Taylor G, Fuller J, Bryce E, Embree J, Gravel D, Katz K, Kibsey P, Kuhn M,

Langley J, Mataseje L, Mitchell R, Roscoe D, Simor A, Thomas E, Turgeon N,

Mulvey M; Canadian Nosocomial Infection Surveillance Program. Complete

Sequence of Four Multidrug-Resistant MOBQ1 Plasmids Harboring bla_{GES-5} Isolated

from *Escherichia coli* and *Serratia marcescens* Persisting in a Hospital in Canada.

Microb Drug Resist. 2015 Jun;21(3):253-60. doi: 10.1089/mdr.2014.0205. Epub 2014 Dec 29. PMID: 25545311.

19. Chalmers R, Sewitz S, Lipkow K, Crellin P. Complete nucleotide sequence of Tn*10*. J Bacteriol. 2000 May;182(10):2970-2. doi: 10.1128/JB.182.10.2970-2972.2000. PMID: 10781570; PMCID: PMC102010.

20. Shaffer JMC, Giddings LA, Samples RM, Mikucki JA. Genomic and phenotypic characterization of a red-pigmented strain of *Massilia frigida* isolated from an Antarctic microbial mat. Front Microbiol. 2023 May 12;14:1156033. doi: 10.3389/fmicb.2023.1156033. PMID: 37250028; PMCID: PMC10213415.

21. Hamidian M, Hall RM. *Acinetobacter baumannii* ATCC 19606 Carries GI*sul2* in a Genomic Island Located in the Chromosome. Antimicrob Agents Chemother. 2016 Dec 27;61(1):e01991-16. doi: 10.1128/AAC.01991-16. PMID: 27795382; PMCID: PMC5192138.

22. Lam MMC, Koong J, Holt KE, Hall RM, Hamidian M. Detection and Typing of Plasmids in Acinetobacter baumannii Using *rep* Genes Encoding Replication Initiation Proteins. Microbiol Spectr. 2023 Feb 14;11(1):e0247822. doi: 10.1128/spectrum.02478-22. Epub 2022 Dec 6. PMID: 36472426; PMCID: PMC9927589.

23. Weber BS, Ly PM, Irwin JN, Pukatzki S, Feldman MF. A multidrug resistance plasmid contains the molecular switch for type VI secretion in *Acinetobacter baumannii*. Proc Natl Acad Sci U S A. 2015 Jul 28;112(30):9442-7. doi: 10.1073/pnas.1502966112. Epub 2015 Jul 13. PMID: 26170289; PMCID: PMC4522760.

24. Alton NK, Vapnek D. Nucleotide sequence analysis of the chloramphenicol resistance
transposon Tn*9*. Nature. 1979 Dec 20-27;282(5741):864-9. doi: 10.1038/282864a0.
PMID: 390403.

25. Echenique JR, Dorsey CW, Patrito LC, Petroni A, Tolmasky ME, Actis LA.
Acinetobacter baumannii has two genes encoding glutathione-dependent
formaldehyde dehydrogenase: evidence for differential regulation in response to iron.
Microbiology (Reading). 2001 Oct;147(Pt 10):2805-2815. doi: 10.1099/00221287-
147-10-2805. PMID: 11577159.

26. De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack:
visualizing and processing long-read sequencing data. Bioinformatics. 2018 Aug
1;34(15):2666-2669. doi: 10.1093/bioinformatics/bty149. PMID: 29547981; PMCID:
PMC6061794.

27. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads
using repeat graphs. Nat Biotechnol. 2019 May;37(5):540-546. doi: 10.1038/s41587-
019-0072-8. Epub 2019 Apr 1. PMID: 30936562.

28. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014 Jul
15;30(14):2068-9. doi: 10.1093/bioinformatics/btu153. Epub 2014 Mar 18. PMID:
24642063.

29. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush
D, Keane JA, Parkhill J. Roary: rapid large-scale prokaryote pan genome analysis.
Bioinformatics. 2015 Nov 15;31(22):3691-3. doi: 10.1093/bioinformatics/btv421.
Epub 2015 Jul 20. PMID: 26198102; PMCID: PMC4817141.

30. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics. 2014 May 1;30(9):1312-3. doi:

10.1093/bioinformatics/btu033. Epub 2014 Jan 21. PMID: 24451623; PMCID:

PMC3998144.

31. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic

tree display and annotation. Nucleic Acids Res. 2021 Jul 2;49(W1):W293-W296. doi:

10.1093/nar/gkab301. PMID: 33885785; PMCID: PMC8265157.

32. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A,

Allesoe RL, Rebelo AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B,

Werner G, Bender JK, Stingl K, Nguyen M, Coppens J, Xavier BB, Malhotra-Kumar

S, Westh H, Pinholt M, Anjum MF, Duggett NA, Kempf I, Nykäsenoja S, Olkkola S,

Wieczorek K, Amaro A, Clemente L, Mossong J, Losch S, Ragimbeau C, Lund O,

Aarestrup FM. ResFinder 4.0 for predictions of phenotypes from genotypes. J

Antimicrob Chemother. 2020 Dec 1;75(12):3491-3500. doi: 10.1093/jac/dkaa345.

PMID: 32780112; PMCID: PMC7662176.

33. Zhou S, Liu B, Zheng D, Chen L, Yang J. VFDB 2025: an integrated resource for

exploring anti-virulence compounds. Nucleic Acids Res. 2025 Jan 6;53(D1):D871-

D877. doi: 10.1093/nar/gkae968. PMID: 39470738; PMCID: PMC11701737.

34. Wyres KL, Cahill SM, Holt KE, Hall RM, Kenyon JJ. Identification of *Acinetobacter*

baumannii loci for capsular polysaccharide (KL) and lipooligosaccharide outer core

(OCL) synthesis in genome assemblies using curated reference databases compatible

with *Kaptive*. Microb Genom. 2020 Mar;6(3):e000339. doi: 10.1099/mgen.0.000339.

PMID: 32118530; PMCID: PMC7200062.

35. Dong JF, Liu CW, Wang P, Li L, Zou QH. The type VI secretion system in

Acinetobacter baumannii clinical isolates and its roles in antimicrobial resistance

acquisition.   Microb   Pathog.   2022   Aug;169:105668.   doi:

10.1016/j.micpath.2022.105668.   Epub   2022   Jul   8.   PMID: 35811021.

36. Meumann EM, Anstey NM, Currie BJ, Piera KA, Kenyon JJ, Hall RM, Davis JS,

Sarovich DS. Genomic epidemiology of severe community-onset *Acinetobacter*

baumannii infection. Microb Genom. 2019 Mar;5(3):e000258. doi:

10.1099/mgen.0.000258. Epub 2019 Feb 26. PMID: 30806611; PMCID:

PMC6487312.

37. Wright MS, Haft DH, Harkins DM, Perez F, Hujer KM, Bajaksouzian S, Benard MF,

Jacobs MR, Bonomo RA, Adams MD. New insights into dissemination and variation

of the health care-associated pathogen *Acinetobacter baumannii* from genomic

analysis. mBio. 2014 Jan 21;5(1):e00963-13. doi: 10.1128/mBio.00963-13. Erratum

in: mBio. 2015 Jun 04;6(3):e00585. doi: 10.1128/mBio.00585-15. PMID: 24449752;

PMCID: PMC3903280.

38. Repizo GD, Espariz M, Seravalle JL, Salcedo SP. Bioinformatic Analysis of the Type

VI Secretion System and Its Potential Toxins in the *Acinetobacter* Genus. Front

Microbiol. 2019 Nov 1;10:2519. doi: 10.3389/fmicb.2019.02519. PMID: 31736933;

PMCID: PMC6838775.

Supplementary Figure 1: Phylogenetic tree of all *A. baumannii* genomes available in BIGSdb that do not encode the T6SS. The cluster

containing the ST19 lineage is highlighted; turquoise indicates ST19, and gray indicates non-ST19.

Supplementary Figure 2: Phylogenetic analysis of three isolates and an additional 156 *A. baumannii* ST19 genomes, depicting the year of

isolation, country, capsule locus type (KL), outer core lipooligosaccharide type (OCL), and the presence of the GI*sul2* genomic island, chlorite

dismutase, T6SS, the *frmRAB* operon, plasmid replicons, and resistance and virulence genes, respectively.

Supplementary Figure 3: A, Quantification of the 20 most prevalent sequence types (STs) of *A. baumannii* genomes from BIGSdb lacking the

T6SS. B, Pie chart showing the percentage distribution of the 10 most prevalent STs of *A. baumannii* genomes from BIGSdb lacking the T6SS.

C, Numeric prevalence of STs within the cluster containing the ST19 lineage shown in Supplementary Figure 1.

512