The Impact of Visual Perturbations on Balance Control during Walking

Yaqi Li[1], Eugenie Lambrecht[2], Sjoerd M. Bruijn[1], Jaap H. van Dieën[1]*

1   Department of Human Movement Science, Vrije Universiteit Amsterdam, Amsterdam Movement Sciences, Amsterdam, Netherlands
2   Department of Rehabilitation Sciences and Physiotherapy, University of Antwerp, Antwerp, Belgium

Abstract

Visual perturbations may lead to a perception of self-motion and affect balance control. We studied effects of different visual perturbations in 16 healthy young participants walking on a treadmill, by assessing foot placement and center of mass (CoM) movement. Three different visual perturbations were applied: fixating a stationary target at eye level while the background moved to the right (MB), tracking a target moving rightward over a stationary background with head rotation (MT-HR), and tracking the moving target with eye movement only (MT-EM). Linear models were fit to the kinematics data to predict foot placement from CoM state at mid-swing. Over the whole trial, MT-HR and MT-EM caused an increase in step width variability, CoM position variability and foot placement residual error. During visual perturbation epochs specifically, in MB, a left deviation of foot and CoM trajectories was observed from the start of background movement. In MT-HR and MT-EM, a right deviation of foot and CoM trajectories was observed only after the target had stopped moving. Contrary to our expectations, foot placement errors did not coincide with subsequent CoM deviations in the opposite direction. An obvious change in frontal plane trunk orientation was found only in MT-HR. While all visual perturbations affected control of the CoM trajectory in the frontal plane, these effects appeared caused by effects on heading as well as balance control. Head rotation appears to additionally disturb balance through a coupling with trunk orientation.

Keywords: visual perturbations, foot placement, stability, locomotion

Introduction

Balance control is a dynamic process that continuously stabilizes the trajectory of the body's center of mass (CoM) against the destabilizing effects of gravity encountered by humans in upright postures (Bruijn and van Dieen 2018). The main mechanism to maintain balance during walking is the selection of the appropriate location for foot placement in relation to the ongoing movement of the CoM. This control has been identified by a linear relation between the CoM state during the swing phase and the location of foot placement at the end of the swing phase, and is thought to be achieved at least in part actively (van Leeuwen, Bruijn et al. 2022). For this control, the state of the CoM is estimated from sensory information provided by the visual, vestibular, and proprioceptive systems (Perry, Santos et al. 2001, Betker, Moussavi et al. 2008, Magnani, van Dieën et al. 2023). This information is integrated by the central nervous system (CNS), which then sends signals to alfa-motor neurons to execute muscular responses to achieve the appropriate foot placement.

The visual system is not only used to maintain balance but also to gather information about the environment by fixating on or tracking visual targets. Fixating on a stationary target while the background is moving can perturb balance control. Previous studies have shown that background movement can lead to a perception of self-motion in the direction opposite to the background movement (Nakamura 1996, Nakamura, Palmisano et al. 2016). Such illusions lead to balance responses that move the CoM in the direction of the background movement. In walking, it has been found that subjects step in the direction of the illusory self-movement and displace their CoM in the opposite direction (Reimann, Fettrow et al. 2018).

Another potential visual perturbation during walking may be caused by visual tracking of moving targets. Tracking a moving target causes relative movement of the visual background on the retina, which may induce a similar illusion as described above. Tracking a moving target is usually completed by coordinated eye and head rotation movements (Gadotti, Elbaum et al. 2016, Srivastava, Ahmad et al. 2018, Yamazoe, Mitsugami et al. 2019). Thus, besides affecting visual information, visual tracking will also have effects on the vestibular and neck proprioceptive systems and possibly mechanical effects on balance. Previous literature has shown changes in the gait pattern due to visual tracking and related head movements. Magnani et al (2020) demonstrated that yaw head motion made gait less stable. Vallis and Patla (2004) showed that head rotation during walking coincided with a mediolateral deviation of the CoM trajectory.

As stated above, the CoM trajectory during walking is mainly stabilized by foot placement (Bruijn and van Dieen 2018, van Leeuwen, Bruijn et al. 2022). However, it is not yet clear how fixating on a stationary target over a moving background and tracking a moving target affect control of the CoM by foot placement. For the latter condition, it has not been studied how rotating the head and re-directing the gaze contribute to balance perturbations. In the current study, we investigated the

relationship between the CoM state and foot placement under the aforementioned visual perturbations during walking. We hypothesized that (1) visual perturbation will cause errors in foot placement error and specifically that (2) people will step in the direction opposite to the background movement and in the direction of the target movement during these visual perturbations, which (3) will subsequently cause displacement of the CoM in the opposite direction.

Methods

Participants

Sixteen healthy individuals (age: 23.4±3.9 years, height: 1.70±0.09 m, weight: 62.2±11.7 kg) volunteered for this study. All participants reported themselves to be free from any neurological or musculoskeletal disorders that could negatively affect their balance or walking performance. All participants walked without assistive devices. Ethical approval was obtained from the ethics committee of the Faculty of Behavioral and Movement Sciences at Vrije Universiteit Amsterdam (VCWE-2023-170). All procedures conformed to the Declaration of Helsinki and participants signed informed consent before participation.

Instruments

All measurements were conducted on a system that integrated a motion capture system sampling at 50 samples/sec (Optotrak, Northern Digital, Waterloo ON, Canada) with an instrumented treadmill (Motek-Forcelink, Amsterdam Netherlands), and a data projector. Clusters of three markers were affixed to the participant's bodies at the following locations: the posterior surfaces of both heels, the sacrum at the level of S2, the trunk at the level of T6, and the back of head. A screen (2.50 m* 1.56 m) was placed 1.60 m in front of the participants with the projector behind it to present the visual perturbations. **Figure 1A** shows the full set-up.

Visual perturbations

A background of 12 uniformly distributed black-and-white vertical stripes (0.205 m * 1.560 m) was presented on the screen. A red target dot measuring 12 cm in diameter was positioned at the center of the screen. The height of the red target was adjusted to be at the participant's eye level. Participants walked under four different conditions: (a) Normal walking (NW), in which the whole scene was stationary. Participants were asked to walk normally and weren't instructed to look at a specific point. (b) Moving background (MB), during this perturbation, the red target dot was fixed in the middle, while the black-and-white background moved horizontally from the middle to the right side corresponding to a 45 degrees movement in 4s, stayed at the side for 8s, and then moved back to the middle. In this condition, participants were asked to keep looking at the stationary red target dot in the middle. (c) Moving target with head rotation (MT-HR), during this perturbation, the black-and-white background was

stationary, while the red target dot moved horizontally over 45 degrees to the right side in 4s and stayed at the side for 8s then moved back to the middle. Participants were asked to track the moving target. (d) Moving target with eye movement only (MT-EM), during this perturbation, the scene was the same as during MT-HR, but the participants were asked to track the horizontally moving target rolling their eyes only while keeping their head stationary. All conditions are illustrated in **Figure1B.**

Visual perturbations were triggered at right heel strikes. Approximately 14 perturbations in total were provided in each walking trial. The first perturbation was triggered at the 20th right heel strike. Six to eight right heel strikes occurred randomly after every perturbation before a new perturbation was provided.

Procedures

The treadmill speed was set to 3.6km/h. To enhance immersion, the lights in the lab were turned off, and the curtains were closed. Before the actual measurement, there was a 10-minute period during which participants were familiarized with the walking speed and all perturbations. Subsequently, a normal walking trial was conducted, followed by the three perturbation trials in a randomized order. Each trial lasted 5 minutes. A 5-minute seated rest was given between trials.

Figure 1 Illustration of the experimental setup (A) and visual perturbation conditions (B).

Data analysis

The mediolateral CoM position was approximated as the average position of the pelvis marker cluster. Mediolateral foot positions were represented by both heel clusters. Trunk orientation in the frontal plane was calculated based on the trunk cluster relative to the laboratory coordinate system.

Gait characteristics reflecting balance were calculated over the whole walking trial, including step width, step width variability, step frequency, CoM position variability, trunk orientation variability, and foot placement residual error. Step width was defined as the mediolateral distance between the heel positions of both feet at heel strike. Step frequency was defined as the number of steps per minute. The foot placement residual error was defined as the RMS of the foot placement error based on the model described below.

We used the complete time series of each trial to determine a linear model predicting foot placement by the preceding CoM state. The model linked the variance in mediolateral foot placement at heel strike to the variance in mediolateral CoM position and velocity during the preceding swing phase (Wang and Srinivasan 2014, Arvin, Hoozemans et al. 2018, van Leeuwen, van Dieen et al. 2020) as:

$$FP_i = \beta_1 * CoM_{i, mid-swing} + \beta_2 * vCoM_{i, mid-swing} + \varepsilon_i$$

With foot placement of the leading leg (FP) at step i and position of CoM at mid-swing ($CoM_{i, mid-swing}$) defined relative to the stance foot of the trailing leg, $vCoM_{i, mid-swing}$ referring to the velocity of the CoM at mid-swing. β_1 and β_2 are regression coefficients and ε_i is the error for every step. The RMS of ε over all steps was defined as the residual error in foot placement.

We analysed the time series of foot position, foot placement error, CoM position, and trunk orientation in detail from the start of the visual perturbation to its end, and divided these episodes into two phases: moving and stationary phases (Figure 1B). The continuous variables were low-pass filtered at 0.01Hz (2nd order bidirectional Butterworth filter) to eliminate the fluctuations related to the stride cycle. From all these variables we subtracted the value at the first sample of the visual perturbation epoch. Peak position during each phase was selected as the maximum absolute value. We determined the average values of these variables among subjects over perturbations. Foot placement errors in both phases of the visual perturbation were extracted from the foot placement model described above.

Statistics

All statistical analyses were performed in Matlab (R2023a, The Mathworks Inc., Natick, MA). For all statistical tests, the assumption of normality was checked using a Shapiro-Wilks test. To assess the overall effects of visual perturbations on gait characteristics, repeated measures analyses of variance with Visual perturbation as a factor was used. For variables that did not follow a normal distribution (i.e. CoM variability and foot placement residual error) the Friedman test was used. For the perturbation epochs, one-sample *t*-tests were used to assess if averaged peak foot positions, peak CoM position, peak trunk orientation, and foot placement errors per step were significantly different from zero. As foot placement error didn't follow a normal distribution, Wilcoxon signed-rank tests were used. Repeated measures

analyses of variance with two factors (Visual perturbation and Phase) were used for peak foot positions, peak CoM position, peak trunk orientation in the perturbation epochs (MB, MT-HR and MT-EM). When we found main effects or interaction effects to be significant, we performed Bonferroni corrected pairwise post-hoc comparisons to test for differences among visual perturbations for each phase (moving and stationary phases). For all repeated measures analyses, if the assumption of sphericity was violated, the Greenhouse-Geisser correction was be applied to adjust the degrees of freedom. For all analyses, the significance level (p-value) was set at 0.05.

Results

Destabilizing effects of visual perturbations over the whole trial

There were no significant effects of Condition on step width and step frequency. Step width variability was significantly affected by Visual perturbation ($F_{visual\ perturbation}$ (df = 3,45) = 10.744, p < 0.01) (**Figure 2**). Step width variability in the NW condition was significantly smaller than in the MT-HR and MT-EM conditions (p < 0.01; p < 0.01, respectively). There was no significant difference between MB and the other conditions.

CoM position variability was significantly affected by Visual perturbation ($\chi^2_{visual\ perturbation}$ (df = 3,45) = 30.22, p < 0.05). CoM position variability in the NW condition was significantly smaller than in the MT-HR and MT-EM conditions (p < 0.01; p < 0.01, respectively), which were not significantly different from each other. There was no significant difference between MB and the other conditions.

Trunk orientation variability was significantly affected by Visual perturbation ($F_{visual\ perturbation}$ (df = 3,45) = 31.534, p < 0.05). During NW, it was significantly smaller than during MT-HR (p < 0.01) and MT-EM (p < 0.01) conditions. MT-HR and MT-EM conditions also had significantly greater trunk orientation variability than MB (p < 0.01; p < 0.01). Trunk orientation variability in the MB condition was not significantly different from that in NW.

Foot placement residual error was significantly affected by Visual perturbation ($\chi^2_{visual\ perturbation}$ (df = 3,45) = 29.77, p < 0.01). It was significantly smaller in NW than in MT-HR (p < 0.01) and MT-EM (p < 0.01). Foot placement residual error in MT-HR was significantly greater than in MB (p < 0.01), while MB was not significantly different from NW.

Figure 2 Effects of Visual perturbation conditions on step width, step width variability, step frequency, CoM position variability, and trunk orientation variability. Variability of the kinematic variables was quantified as the standard deviation over steps. Significant effects are indicated by an asterisk. NW: normal walking, MB: moving background, MT-HR: moving target with head rotation, MT-EM: moving target with eye movement.

Destabilizing effects during the visual perturbation

Qualitatively, foot trajectories changed during the visual perturbations (**Figure 3**). In the MB condition, participants placed their feet in the direction opposite to the background movement from the start of the moving phase. A peak was reached at the beginning of the stationary phase and then the foot trajectories tended to gradually return to the original location. In the MT-HR condition, participants placed their feet more in the movement direction of the target during the moving phase and continued deviating throughout the stationary phase. In the MT-EM condition, participants placed their feet more in the direction of the target from the end of the moving phase and continued deviating throughout the stationary phase.

Figure 3 Deviations in foot and CoM positions during visual perturbation epochs. All the data were referenced to the first sample of the visual perturbation epoch. From left to right: (A) MB: moving background. (B) MT-HR: moving target with head rotation, (C) MT-EM: moving target with eye movement. The vertical dashed lines separate the different phases: the line at 0s marks the start of the moving phase, the line at 4s marks the start of the stationary phase, and the line at 12s marks the end of the stationary phase. The shaded area represents the between-participants standard deviations.

Figure 4 shows the peak foot positions averaged over perturbations during the moving and stationary phases relative to the start of the visual perturbation for all perturbation conditions. In MB, foot positions deviated significantly leftward in the moving phase ($p < 0.01$) and remained deviated to the left in the stationary phase ($p < 0.01$). In both MT-HR and MT-EM, foot positions were not significantly different from zero in the moving phase (MT-HR: $p = 0.27$; MT-EM: $p = 0.83$). However, they significantly deviated to the right in the stationary phase (MT-HR: $p < 0.05$; MT-EM: $p < 0.05$). We note that this effect was not very consistent between participants.

A two-way repeated measures ANOVA showed significant effects of Visual perturbation on averaged foot peak positions ($F_{visual\ perturbation}$ (df = 2,30) = 17.006, $p < 0.01$). Post-hoc results showed that MB was significantly different from MT-HR ($p < 0.01$), and MT-EM ($p < 0.01$). Phase also had a significant effect on averaged foot peak positions (F_{phase} (df = 1,15) = 9.4062, $p < 0.01$). The interaction between Visual perturbation and Phase ($F_{visual\ perturbation*phase}$ (df = 2,30) = 8.8686, $p < 0.01$) indicated that during the stationary phase the differences in averaged foot peak positions between visual perturbations were larger than during the moving phase.

Figure 4 Averaged peak foot positions relative to the first sample of the visual perturbation epoch in the MP (moving phase) and SP (stationary phase), respectively. From left to right: MB: moving background; MT-HR: moving target with head rotation; MT-EM: moving target with eye movement. Significant differences from zero are indicated by an asterisk.

The first ten foot placement errors after the visual perturbation of each condition can be seen in **Figure 5**. In MB, leftward (negative) foot placement errors were significant at the second ($p < 0.01$), third ($p < 0.01$), fourth ($p < 0.01$), and sixth steps ($p < 0.05$), and a rightward foot placement error was significant at the tenth step ($p < 0.05$). In MT-HR, rightward foot placement errors were significant at the fourth ($p < 0.01$), seventh ($p < 0.01$), eighth ($p < 0.05$), ninth ($p < 0.01$), and tenth ($p < 0.05$) steps. In MT-EM, a leftward foot placement error was significant in the second step ($p < 0.01$) and a rightward foot placement error in the tenth step ($p < 0.05$).

Figure 5 Foot placement error of the first ten steps after the onset of the visual perturbation. From left to right: MB: moving background, MT-HR: moving target with head rotation, MT-EM: moving target with eye movement. Steps with significant differences from zero are indicated by an asterisk.

Qualitatively, in all conditions, the CoM appeared to follow the positions of the feet (**Figure 6**). In MB, significant leftward deviations occurred in the moving phase ($p < 0.01$), which remained significant in the stationary phase ($p < 0.01$). In both MT-HR and MT-EM, peak CoM positions were not significantly different from zero in the moving phase (MT-HR: $p = 0.40$; MT-EM: $p = 0.98$). However, they deviated significantly rightward in the stationary phase (MT-HR: $p < 0.05$; MT-EM: $p < 0.05$). Generally, the peak CoM positions deviated leftward during MB, whereas in MT-HR and MT-ER, they deviated to the right.

Peak CoM positions in MB were significantly ($F_{visual\ perturbation}(df = 2,30) = 14.918$, $p < 0.01$) smaller than in MT-HR ($p < 0.01$) and MT-EM ($p < 0.01$). While there was no significant difference between MT-HR and MT-EM. A significant effect of Phase was also observed ($F_{phase}(df = 1,15) = 8.2733$, $p < 0.05$), showing that the CoM deviation in the stationary phase was larger than in the moving phase ($p < 0.05$). The interaction between Visual perturbation and Phase ($F_{visual\ perturbation*phase} (df = 2,30) = 9.0611$, $p < 0.01$) indicated that during the stationary phase the differences in peak CoM positions between visual perturbations were larger than during the moving phase.

Figure 6 Peak CoM positions relative to the first sample of the visual perturbation epoch in the moving phase (left) and stationary phase (right). MB: moving background; MT-HR: moving target with head rotation; MT-EM: moving target with eye movement. Significant differences from zero are indicated by an asterisk.

Peak frontal plane trunk orientations during the moving and stationary phases are shown in **Figure 7**. In MB, trunk orientation did not significantly differ from zero in either phase (moving phase: p = 0.38; stationary phase: p = 0.76). In MT-HR, the trunk was significantly oriented leftward during both moving (p < 0.01) and stationary phases (p < 0.01). In MT-EM, trunk orientation was not significantly different from zero in the moving phase (p = 0.08), while it deviated leftward in the stationary phase (p < 0.01).

Peak trunk orientation depended on the visual perturbation condition ($F_{visual\ perturbation}$(df = 2,30) = 24.581, p < 0.01). In MT-HR, it was significantly smaller (more leftward) than in both MT-EM (p < 0.01) and MB (p < 0.01), but there was no significant difference between MB and MT-EM (p = 0.17). A significant effect of Phase was observed as well (F_{phase}(df = 1,15) = 16.271, p < 0.01), indicating the leftward tilt became larger during the stationary phase. The interaction between Visual perturbation and Phase ($F_{visual\ perturbation*phase}$ (df = 2,30) = 5.9438, p < 0.01) indicated that during the stationary phase, the differences in trunk orientation between visual perturbations were larger than in the moving phase.

Figure 7 Frontal plane trunk orientation during visual perturbation epochs. From left to right: MB: moving background. MT-HR: moving target with head rotation. MT-EM: moving target with eye movement. All the data were referenced to the first sample of the visual perturbation epoch. The panels in the first row show the deviations of trunk orientation. The vertical dashed lines in the figures at the first row separate the different phases: the line at 0s marks the start of the moving phase, the line at 4s marks the start of the stationary phase, and the line at 12s samples marks the end of the stationary phase. The shaded area represents the between-participants standard deviations. The panels in second row show the peak frontal plane trunk orientation in the moving phase (MP) and stationary phase (SP), respectively. Significant differences from zero are indicated by an asterisk.

Discussion

The visual system can be used to gather information from the environment to control balance. However, this information can in some situations be considered as a perturbation (Logan, Kiemel et al. 2010, Mohebbi, Amiri et al. 2022). For example, on a stationary train, one may perceive to be moving when a train slowly passes by. In this study, we applied different types of visual perturbations to study if and how these affect balance during walking through eliciting changes in foot placement. Our results

indicated that the variability of trunk sway and foot placement increased over the entire walking trial only when tracking a moving target, either with head rotation (MT-HR) or with eye movement only (MT-EM). During the visual perturbation epochs, fixating a stationary target while the background moved caused a deviation of foot and CoM trajectories opposite to the background movement. Tracking a moving target with head rotation or with eye movement only did not cause obvious deviations of foot and CoM trajectories during the moving phase, but did cause deviations in the target direction during the stationary phase.

Over the whole walking trial, a negative effect of visual perturbations on gait characteristics was only found in MT-HR and MT-EM. In these two conditions, step width variability, CoM position variability, trunk orientation variability, and foot placement residual error were significantly higher than in NW. This indicates that visually tracking a moving target destabilizes walking even in healthy young participants. Unexpectedly, MB did not induce similar increases in gait characteristics. Since foot and CoM trajectories in MB deviated but returned towards normal already during the stationary phase, the effect in MB may have been diluted by the unperturbed steps which may have masked the impact of the perturbation on the trial as a whole.

As expected, the deviation in foot placement during the moving phase varied between conditions. MB caused a deviation opposite to the background movement. However, in MT-HR and MT-EM, no obvious deviations were observed in this phase, in line with findings from a study by Vallis and Patla (2004). This may be explained by correction based on other information. For one, in the target tracking tasks, the CNS likely interpreted the relative backround movement as a predictable outcome of self-motion, attenuating its perceptual salience and reducing the need for behavioral adjustments (Spering, Schutz et al. 2011). Other sources of information that might help to distinguish self-motion and external motion may be: (1) the stationary backgrond sensed by pheripheral vision: the fixed black-and-white vertical stripes background provided a reference, (2) proprioception: proprioceptive information from the neck and eye extraocular muscles would indicate self-motion (Wang and Pan 2013, Blumer, Carrero-Rojas et al. 2024), (3) the presence of an efference copy: predictions based on an efference copy of the motor commands related to target tracking would also indicate self-motion (Wolpert and Ghahramani 2000). In addition, since the onset of the target movements was unpredictable and fast, saccades may have been used during tracking instead of smooth pursuit movements (Leigh and Kennard 2004). While smooth pursuit eye movements appear to disturb balance during walking, saccades were not found to do so (Thomas, Dewhurst et al. 2017). Saccadic eye movements may induce suppression of visual information outlasting the duration of the saccade itself (Ibbotson and Krekelberg 2011), which may have reduced sensitivity to the visual perturbation.

The deviation in foot placement trajectories during the stationary phase in MB became smaller, while it became larger in MT-HR and MT-EM. Obviously,

accumulation of small errors over the moving phase may have occurred, leading to significant deviations only in the stationary phase. This finding may also reflect that heading direction was aligned with gaze direction in this phase in all conditions. It has been proposed that head orientation or gaze direction provides the CNS with a frame of reference, and that a change of this reference releases a steering synergy which affects heading direction (Hollands, Patla et al. 2002, Vallis and Patla 2004, Bernardin, Kadone et al. 2012, Belmonti, Cioni et al. 2013). Finally, walking with a large head or eye rotation angle may be considered as a dual task, decreasing the control over heading, but it is not obvious why this would lead to deviations in a specific direction.

We expected foot placement errors in the direction of illusory self-motion. In MB perturbations, where the background moved to the right, participants indeed stepped left of the predicted foot placement from the second through the fourth step throughout almost the whole moving phase, which would align with an illusion of a deviation in the CoM state (postion or velocity) in this phase. In MT-HR perturbations, with a right target movement, as expected rightward foot placement errors appeared at the 4[th] step, around the transition between the moving and stationary phases, and at the 7[th], 9[th] and 10[th] steps during stationary phase. In MT-EM, this happened only at the 10[th] step. In this condition, the second step (of the left foot) was unexpectedly placed leftwards.

Generally, a lateral step is associated with a force accelerating the CoM in the opposite direction (Reimann, Fettrow et al. 2018). However, none of the foot placement errors were associated with a subsequent movement of the CoM to the oppposite side. Instead, in contrast with our expectation, the foot placement deviations coincided with a CoM displacement in the same direction. Possibly, the CoM trajectory was adjusted immediately after foot placement through ankle moments bringing the CoM in the same direction (van Leeuwen, van Dieen et al. 2020). Our predictions were based on the expected role of visual information in sensing the state of the CoM and the role of foot placement in stabilizng the CoM trajectory, both in the frontal plane. The conflicting findings may be attributable to the fact, already alluded to above, that the visual tasks may affect heading estimates and that foot placement also plays a role in control of heading.

A significant frontal trunk roll was found in MT-HR, which is consistent with previous literature (Patla, Adkin et al. 1999, Vallis and Patla 2004, van Andel, Schmidt et al. 2023). Van Andel et al. (2023) illustrated that head rotation was associated with frontal plane rotation of the trunk. Previous studies (Patla, Adkin et al. 1999, Vallis and Patla 2004) proposed that this frontal trunk roll is used as a strategy to push the CoM in the desired direction when making a turn and this would match with our participants walking in the direction of the target. However, in our study, trunk roll only occurred in MT-HR but not in MT-EM, where a similar deviation in gait direction was observed. Before making a turn, people usually rotate their head in the direction of the turn as well (Grasso, Prevost et al. 1998, Hicheur, Vieilledent et al. 2005,

Dollack, Perusquia-Hernandez et al. 2019). Therefore, we assume that the trunk roll may be a result of a mechanical coupling between head rotation and trunk roll. In a post hoc test on whether such trunk roll is also present when turning the head while standing (see **Figure 8,** data from the same subjects as in the main results), we found this to be the case, supporting that this could result from a mechanical coupling.

Figure 8 Frontal plane trunk orientation deviations during visual perturbation epoch in standing. All the participants performed the visual perturbation conditions in standing. For each condition, five repeated standing trials were conducted. During recording, participants were instructed to fixate the target or track the target either by head or eyes and stand still for 40s. During each trial one one visual perturbation was imposed . The visual perturbation began around the 10th second and lasted for 16s. The setting of visual perturbation was the same as in walking. All the data were referenced to the first sample of the visual perturbation epoch. The vertical dashed lines in the figures separate the different phases: the line at 0s marks the start of the moving phase, the line at 4s marks the start of the stationary phase, and the line at 12s marks the end of the stationary phase. The shaded area represents the between-participants standard deviations. From left to right column: MB: moving background. MT-HR: moving target with head rotation. MT-EM: moving target with eye movement.

This study has several limitations. First, we only recruited healthy young adults, and results may be different for example in older adults, who have a high risk of falling (Lutz, Sanderson et al. 2008). Secondly, the experiment was conducted on a treadmill. This partially constrains walking speed and walking direction, which may have had an impact on participant's responses to the visual perturbations. Thirdly, we did not assess foot orientation. Foot orientation may have played a role in

redirecting the CoM trajectories. Further investigations may also consider the role of ankle moments for keeping balance during visual perturbations.

Conclusion

Our results showed that visual perturbations challenge frontal plane balance, or in other words the control of the CoM trajectories in this plane during walking. This was in line with expected effects on sensory estimates of the CoM state used for the control of foot placement in stabilizing the frontal plane CoM trajectory. Mechanical effects of head rotation, associated with visual tracking, had an additional disturbing effect on balance. However, effects of the gaze direction on heading appeared to affect gait simultaneously and cannot easily be distinguished from responses related to balance control.

Acknowledgment

This study was supported by a scholarship (No. 202308310074) from the China Scholarship Council (CSC). The authors appreciate all participants for their voluntary participation. The authors thank Richard Casius, Leon Schutte and Paul Pesman for the technical assistance. The authors also thank Raven Huiberts for his help in data collection.

Reference

Arvin, M., M. J. M. Hoozemans, M. Pijnappels, J. Duysens, S. M. Verschueren and J. H. van Dieen (2018). "Where to Step? Contributions of Stance Leg Muscle Spindle Afference to Planning of Mediolateral Foot Placement for Balance Control in Young and Old Adults." Front Physiol **9**: 1134.

Belmonti, V., G. Cioni and A. Berthoz (2013). "Development of anticipatory orienting strategies and trajectory formation in goal-oriented locomotion." Exp Brain Res **227**(1): 131-147.

Bernardin, D., H. Kadone, D. Bennequin, T. Sugar, M. Zaoui and A. Berthoz (2012). "Gaze anticipation during human locomotion." Exp Brain Res **223**(1): 65-78.

Betker, A. L., Z. M. K. Moussavi and T. Szturm (2008). "Ambulatory Center of Mass Prediction Using Body Accelerations and Center of Foot Pressure." IEEE Trans Biomed Eng **55**(11): 2491-2498.

Blumer, R., G. Carrero-Rojas, P. M. Calvo, J. Streicher, R. R. de la Cruz and A. M. Pastor (2024). "Proprioceptors in extraocular muscles." Exp Physiol **109**(1): 17-26.

Bruijn, S. M. and J. H. van Dieen (2018). "Control of human gait stability through foot placement." J R Soc Interface **15**(143).

Dollack, F., M. Perusquia-Hernandez, H. Kadone and K. Suzuki (2019). "Head Anticipation During Locomotion With Auditory Instruction in the Presence and Absence of Visual Input." Front Hum Neurosci 13: 293.

Gadotti, I. C., L. Elbaum, Y. Jung, V. Garbalosa, S. Kornbluth, B. Da Costa, K. Maitra and D. Brunt (2016). "Evaluation of eye, head and trunk coordination during target tracking tasks." Ergonomics 59(11): 1420-1427.

Grasso, R., P. Prevost, Y. P. Ivanenko and A. Berthoz (1998). "Eye-head coordination for the steering of locomotion in humans: an anticipatory synergy." Neurosci Lett 253(2): 115-118.

Hicheur, H., S. Vieilledent and A. Berthoz (2005). "Head motion in humans alternating between straight and curved walking path: combination of stabilizing and anticipatory orienting mechanisms." Neurosci Lett 383(1-2): 87-92.

Hollands, M. A., A. E. Patla and J. N. Vickers (2002). ""Look where you're going!": gaze behaviour associated with maintaining and changing the direction of locomotion." Exp Brain Res 143(2): 221-230.

Ibbotson, M. and B. Krekelberg (2011). "Visual perception and saccadic eye movements." Curr Opin Neurobiol 21(4): 553-558.

Leigh, R. J. and C. Kennard (2004). "Using saccades as a research tool in the clinical neurosciences." Brain 127(Pt 3): 460-477.

Logan, D., T. Kiemel, N. Dominici, G. Cappellini, Y. Ivanenko, F. Lacquaniti and J. J. Jeka (2010). "The many roles of vision during walking." Exp Brain Res 206(3): 337-350.

Lutz, W., W. Sanderson and S. Scherbov (2008). "The coming acceleration of global population ageing." Nature 451(7179): 716-719.

Magnani, R. M., S. M. Bruijn, J. H. van Dieen and M. F. Vieira (2020). "Head orientation and gait stability in young adults, dancers and older adults." Gait Posture 80: 68-73.

Magnani, R. M., J. H. van Dieën and S. M. Bruijn (2023). "Effects of vestibular stimulation on gait stability when walking at different step widths." Exp Brain Res 241(1): 49-58.

Mohebbi, A., P. Amiri and R. E. Kearney (2022). "Identification of human balance control responses to visual inputs using virtual reality." J Neurophysiol 127(4): 1159-1170.

Nakamura, S. (1996). "Effects of background stimulation upon eye-movement information." Percept Mot Skills 82(2): 627-635.

Nakamura, S., S. Palmisano and J. Kim (2016). "Relative Visual Oscillation Can Facilitate Visually Induced Self-Motion Perception." Iperception **7**(4): 2041669516661903.

Patla, A. E., A. Adkin and T. Ballard (1999). "Online steering: coordination and control of body center of mass, head and body reorientation." Exp Brain Res **129**(4): 629-634.

Perry, S. D., L. C. Santos and A. E. Patla (2001). "Contribution of vision and cutaneous sensation to the control of centre of mass (COM) during gait termination." Brain Res **913**(1): 27-34.

Reimann, H., T. Fettrow and J. J. Jeka (2018). "Strategies for the control of balance during locomotion." Kinesiology Review **7**(1): 18-25.

Reimann, H., T. Fettrow, E. D. Thompson and J. J. Jeka (2018). "Neural Control of Balance During Walking." Front Physiol **9**: 1271.

Spering, M., A. C. Schutz, D. I. Braun and K. R. Gegenfurtner (2011). "Keep your eyes on the ball: smooth pursuit eye movements enhance prediction of visual motion." J Neurophysiol **105**(4): 1756-1767.

Srivastava, A., O. F. Ahmad, C. P. Pacia, M. Hallett and C. Lungu (2018). "The Relationship between Saccades and Locomotion." J Mov Disord **11**(3): 93-106.

Thomas, N. M., S. Dewhurst, T. M. Bampouras, T. Donovan, A. Macaluso and G. Vannozzi (2017). "Smooth pursuits decrease balance control during locomotion in young and older healthy females." Exp Brain Res **235**(9): 2661-2668.

Vallis, L. A. and A. E. Patla (2004). "Expected and unexpected head yaw movements result in different modifications of gait and whole body coordination strategies." Exp Brain Res **157**(1): 94-110.

van Andel, S., A. R. Schmidt and P. A. Federolf (2023). "Distinct coordination patterns integrate exploratory head movements with whole-body movement patterns during walking." Sci Rep **13**(1): 1235.

van Leeuwen, A. M., S. M. Bruijn and J. H. van Dieen (2022). "Mechanisms that stabilize human walking." Braz J Mot Behav **16**(5): 326-351.

van Leeuwen, A. M., J. H. van Dieen, A. Daffertshofer and S. M. Bruijn (2020). "Active foot placement control ensures stable gait: Effect of constraints on foot placement and ankle moments." PLoS One **15**(12): e0242215.

Wang, J. and Y. Pan (2013). "Eye proprioception may provide real time eye position information." Neurol Sci **34**(3): 281-286.

Wang, Y. and M. Srinivasan (2014). "Stepping in the direction of the fall: the next foot placement can be predicted from current upper body state in steady-state walking." Biol Lett **10**(9).

Wolpert, D. M. and Z. Ghahramani (2000). "Computational principles of movement neuroscience." <u>Nat Neurosci</u> **3 Suppl**: 1212-1217.

Yamazoe, H., I. Mitsugami, T. Okada and Y. Yagi (2019). "Analysis of head and chest movements that correspond to gaze directions during walking." <u>Exp Brain Res</u> **237**(11): 3047-3058.