BENCHMARKING UNIVERSAL MACHINE LEARNING FORCE FIELDS WITH HYDROGEN-BONDING COOPERATIVITY

Xinping Feng[1,2,3], You Xu[*1,2,3], Jing Huang[*1,2,3]

[1]Key Laboratory of Structural Biology of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, Zhejiang 310024, China

[2]Westlake AI Therapeutics Laboratory, Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, Zhejiang 310024, China

[3]Institute of Biology, Westlake Institute for Advanced Study, Hangzhou, Zhejiang 310024, China

ABSTRACT

Machine learning force fields (MLFFs) offer a promising balance between quantum mechanical (QM) accuracy and molecular mechanics efficiency. While MLFFs have shown strong performance in modeling short-range interactions and reproducing potential energy surfaces, their ability to capture long-range cooperative effects remains underexplored. In this study, we assess the ability of three MLFF models — ANI, MACE-OFF, and Orb — to reproduce cooperative interactions arising from environmental induction and dispersion, which are essential for many biomolecular processes. Using a recently proposed framework, we quantify hydrogen bond (H-bond) cooperativity in N-methylacetamide polymers. Our results show that all MLFFs capture cooperativity to some extent, with MACE-OFF yielding the closest agreement with QM data. These findings highlight the importance of evaluating many-body effects in MLFFs and suggest that H-bond cooperativity can serve as a useful benchmark for improving their physical fidelity.

Keywords Machine learning force field · cooperative effects · self-assembly · neural network potential · hydrogen bond

1 Introduction

Machine learning potentials (MLPs), sometimes also referred to as machine learning force fields (MLFFs), have emerged as a significant advancement in computational chemistry, offering a balance between the high accuracy of quantum mechanics (QM) and the computational efficiency of molecular mechanics (MM) [1, 2]. By leveraging machine learning algorithms, MLPs learn the statistical relationship between molecular structures and their potential energies from large datasets. Various methodologies, including kernel-based approaches and neural network (NN) models, have demonstrated notable success in simulating and predicting the properties of complex chemical systems.

NN potentials are particularly promising because deep NNs excel at fitting high dimensional data distributions, enabling them to capture intricate intra- and intermolecular interactions [3, 4, 5, 6, 7, 8]. The seminal work of Behler and Parrinello introduced a scheme in which the total potential energy is decomposed into atomic contributions, each predicted by an NN that takes atom-centered environment descriptors as input [9]. The field has progressed rapidly since then, with more advanced network architectures designed to better preserve physical symmetries and improve training efficiency [10, 11, 12, 13, 14]. Whereas early MLPs were usually trained bespoke on data generated for the particular system studied, recent efforts have shifted toward pre-training models for broad, off-the-shelf use——much like traditional force fields (FFs) that are carefully parametrized once and then distributed to end-users. Resonating

---

with the wider ML move toward "foundation models", an increasing number of universal MLFFs covering substantial regions of chemical space are now becoming available.

One of the most popular universal MLFFs is the ANI series developed by Roitberg and co-workers. The ANI-2x model, employs transfer learning: a network first trained on a large DFT dataset is fine-tuned with a Ser, high-level CCSD(T) dataset, achieving accuracy across a broad chemical space that includes C, H, N, O, S, F, and Cl [15]. Another notable model, MACE, extends message-passing neural networks (MPNNs) by incorporating higher-order equivariant messages, boosting both efficiency and accuracy [13]. The MACE open force fields (MACE-OFFs) were trained on diverse datasets including organic molecules, water clusters, and small peptides, and have demonstrated state-of-the-art performance not only in reproducing potential energy surfaces (PES) and atomic forces, but also in predicting condensed-phase properties such as liquid densities and solvation free energies [16, 17]. Meanwhile, the Orb model, designed for large-scale materials simulations, employs a scalable graph NN architecture that preserves rotational invariance and allows efficient modeling of long-range dispersion interactions via diffusion pre-training and D3 correction [14].

In classical FFs, non-bonded interactions are evaluated throughout the space. In particular, electrostatics are typically handled with the particle meshed Ewald (PME) method, and more recently the LJ-PME method has been adopted to fully account for the van der Waals interactions[18, 19]. This contrasts with MLFFs, in which atomic energies are computed from the information contained within a direct cutoff (typically 4.0 Å to 6.0 Å). Interactions beyond this range are presumed to be captured indirectly, for example through multiple passes of message passing, yet the extent to which MLFFs reproduce full long range interaction remains to be systematically benchmarked [20]. Long-range interactions are essential in chemical and biological systems dominated by non-covalent forces, for example proteins. A prime example is the cooperative effects of hydrogen bonds (H-bonds) in stabilizing protein structures. Once a few H-bonds form between residues, subsequent H-bond formation becomes energetically more favorable, facilitating protein folding, assembly, and aggregation [21, 22, 23].

H-bond cooperativity arises mainly from electronic induction, reflecting the molecular polarizability in response to the surrounding environment. QM methods such as CCSD(T), MP2, and DFT can accurately capture these effects, as can polarizable MM force fields, which explicitly account for charge redistribution [24, 25, 26, 27]. In contrast, additive force fields fail to reproduce such cooperative effects. The quantification of cooperative energy has historically varied due to differences in computational methods, model systems, and definitions of cooperativity. QM calculations of various H-bonded polymers, including N-methylacetamide (NMA) [28], water [29], N-methylformamide (NMF) [30], formamide [31], and alanine peptides [32, 33], have yielded cooperative energy estimates ranging from 3 to 26 kcal/mol, depending on the specific formulation used.

Recent methodological advancements have enabled more rigorous quantification of cooperative effects and improved benchmark for classical MM force fields [34]. In this recent study, we introduced a general framework that defines cooperativity as the difference between the interaction energy of an isolated dimer (V_{A-B}^{AB}) and its interaction energy in the presence of a third molecule (V_{A-B}^{ABC}) (Figure 1a). This cooperative energy is computed using the internal energies of optimized geometries (Equation 1).

$$\Delta V_{A-B} = -(U_{ABC}^{ABC} - U_{BC}^{ABC} - U_{AC}^{ABC} + U_{C}^{ABC} - U_{AB}^{AB} + U_{A}^{AB} + U_{B}^{AB}) \tag{1}$$

As the superscripts indicate, the first four terms (internal energies of ABC, BC, AC, and C) are computed using the geometry and basis set of complex ABC, while the remaining three terms (internal energies of AB, A, and B) are obtained using the geometry and basis set of complex AB. Here, A, B, and C represent any number of molecules. The framework accounts for geometric and inductive effects of molecule C on the interaction between A and B while excluding contributions from C's internal interactions. This formulation thus enables a systematic analysis of many-body effects in various molecular environments.

Here, we apply this theoretical framework to evaluate the ability of MLFFs to capture H-bond cooperatives in MLFFs. We used the same model systems and QM references, where NMA chains were optimized at the ωB97XD/cc-pVTZ level and energies computed at the RI-MP2/aug-cc-pVTZ level. Our findings reveal that all MLFFs considered, including the ANI, MACE-OFF, and Orb models, capture cooperativity to some extent but with considerable variance in accuracy. Despite being trained on high-quality QM data covering chemical spaces, these MLFFs differ substantially in how they represent environment-dependent interactions. Our work provides a critical benchmark for MLFFs and sheds light on how these models capture complex cooperative effects.

2 Method

Homogeneous NMA polymers identical to the previous study were used [34], where NMAs oriented in parallel configuration were arranged to form hydrogen bonds (H-bonds) in a head-to-tail way. As the fundamental H-bonded block, an NMA dimer was used, in which a hydrogen bond was formed between the carbonyl oxygen of molecule A (O_A) and the amide nitrogen of molecule B (N_B). The dimer was constrained to remain planar.

The NMA dimer adopted two distinct conformations. In the syn conformation, atoms N_A and N_B are positioned on the same side of the C_A-O_A axis, corresponding to dihedral ϕ_1 (N_A-C_A-O_A-N_B) of $0°$. In contrast, in the anti conformation, N_A and N_B lay on opposite sides of the C_A-O_A axis, with ϕ_1 at $180°$ (Figure 1b). Two well-ordered NMA polymer patterns were examined: one in which identical dimers formed an arc pattern and another where alternating syn and anti dimers resulted in a linear pattern. All polymer structures were built starting from a syn dimer and extended up to 12 NMAs. The head, middle, and tail H-bonds were calculated within each polymer to assess the cooperative effects of chain elongation on different sides (Figure 1c, d).

The cooperativity of H-bond interactions was evaluated for three families of MLFFs (Table 1), ANI models including 1ccx [35] and 2x [15], Orb models [14] including d3-v2, v2, and v3-conservative-20-omat [36], and MACE-OFF models including 23-S, 23-M, 23b-M [16], 23-SC [37], and 24-M [17]. The energy terms of equation 1 were calculated from the corresponding geometries that were energetically minimized. The initial conformations were taken from QM-optimized geometries. During minimization, torsions ϕ_1 and ϕ_2 (C_A-O_A-N_B-C_B) was constrained to maintain planarity across all NMA molecules. For Orb models, structure optimization using the L-BFGS algorithm and single-point potential energy calculations were performed via the ASE calculator interface [38]. For MACE-OFF and ANI models, structures were minimized using the L-BFGS algorithm, and potential energy calculations were carried out in OpenMM [39], employing a customized API (openmm-ml) designed for ML models in simulations (https://github.com/jharrymoore/openmm-ml/tree/main).

3 Results

3.1 NMA dimer interaction and conformation

NMA serves as a model system for protein backbone interactions in the parametrization of classical force fields. The H-bond interaction between the amide and the carbonyl group stabilizes NMA dimer in a planar conformation (Figure 1b, c). For longer NMA polymers, however, the global energy minimum usually corresponds to irregular aggregated structures with different potential energy models. To systematically evaluate the cooperative effects of each elongating NMA that mimic the formation of secondary structures, planarity restraints were applied to all polymers during optimization, including the reference dimer. In our previous QM calculations, both syn and anti conformations were optimized using density functional theory (DFT) at the ωB97XD/cc-pVTZ level, with interaction energies evaluated at the SCS-MP2/aug-cc-pVTZ level. These calculations indicated an energy difference of approximately 0.6 kcal/mol between the two conformations. Among MLFFs, all MACE-OFF models provided consistent interaction energy estimates, whereas ANI and Orb models exhibited greater variation across their respective variants (Figure 2). Most MLFF models overestimated the interaction energies, though the deviations for all models except ANI-2x and Orb-v2 remained within 2 kcal/mol, which is comparable to those observed in MM force fields.

Without restraints, geometric optimization using MACE-OFF and ANI still recovered the correct dimer conformations (Figure 1b, c), indicating that both force fields provide an adequate description of NMA interaction. In contrast, Orb-v2 models failed to reproduce the H-bonding conformation, instead favoring stacked structures (Figure S1a). This issue persisted for longer chains, where Orb-v2 consistently failed to achieve H-bonded conformations, even under planarity restraints (Figure S1b). To enforce a regular polymer pattern capable of capturing H-bond cooperativity, a much looser convergence tolerance (4.6 kcal/mol/Å) had to be applied on Orb-v2 compared to the 0.00024 kcal/mol/Å used for MACE-OFF and ANI. This limitation was partially addressed in the recently released Orb-v3-conservative-20-omat, which showed improved recovery of the H-bonded dimer and better polymer geometry optimization under reasonable convergence criteria. Nonetheless, Orb-v3 models continued to exhibit challenges when applied to long polymers in the presence of water. As a result, the cooperative energy for Orb models were calculated using QM geometries rather than Orb optimized polymer structures (Figure S1c).

3.2 Cooperative energy of H-bonds

Although the interaction energies of isolated dimers differ slightly between syn and anti conformations, the overall profile of cooperativity for arc and linear polymer patterns remains similar, as shown in the previous study. Such trend was also consistently observed across all MLFFs evaluated in this work. For clarity, the main text focuses on arc

polymers composed of repeating syn dimers, while the corresponding results for linear polymers are provided in the Supporting Information (SI).

Cooperativity was present across all three MLFFs, confirming their reproduction in polarizability. However, the addition of NMAs to either side of the dimer produced markedly different cooperative effects among these models (Figure 3, Figure S2 for linear polymers). The MACE-OFF models closely reproduced the MP2 reference, whereas ANI models substantially underestimated and Orb models significantly overestimated cooperative energies. Consistent with the QM reference, the maximum cooperative energies on both terminal H-bonds were equivalent, measuring 1.4-1.6 kcal/mol in MACE variants, 0.3-0.6 kcal/mol in ANI, 7.5-10 kcal/mol in Orb-v2 models, and 3.5-4 kcal/mol in Orb-v3 model. The maximum cooperative energy for middle hydrogen bonds, as calculated by MP2, was 4.2 kcal/mol. This value was closely reproduced by MACE (2.8-3.2 kcal/mol), underestimated by ANI (0.4-1.0 kcal/mol), significantly overestimated by Orb-v2 (18-20 kcal/mol), and partially corrected in Orb-v3 (7-8 kcal/mol).

In ANI and MACE-OFF models, cooperativity reached a plateau immediately after adding a single NMA on either side, with no further increase upon polymer elongation. MACE-OFF models therefore correctly reproduced the cooperative effect of adding one NMA, while underestimating the effect of further elongation. All the MACE-OFF variants performed equivalently in cooperative energy, except for 23-SC, which exhibited a observable drop, likely due to its modified repulsive atomic interactions that are specifically introduced for free energy perturbation calculations [37]. Different from ANI and MACE-OFF, cooperativity in Orb models increased progressively and converged after the addition of five or more NMAs, closely resembling the QM profile.

3.3 Cooperativity in presence of water molecules

To further assess the impact of solvent, all polymer chain termini were capped with water molecules (Figure 4, Figure S3 for linear polymers). One water molecule formed a single H-bond with the amide group, while three additional water molecules interacted with the carbonyl group via two H-bonds and formed two additional H-bonds among themselves (Figure 1d, e). In SCS-MP2 reference of NMA dimer H-bond, the cooperative contribution of this four-water flanking arrangement (2.5 kcal/mol, Figure 4a) was comparable to that of two NMA molecules (2.8 kcal/mol, Figure 3b). The MACE-OFF models closely reproduced the contribution, yielding values in the range of 2.2-2.9 kcal/mol. ANI models continued to underestimate the cooperativity, suggesting a limited sensitivity to solvent-induced effects (Figure 4d). In contrast, the Orb models showed smaller deviations in cooperative energy for the water-flanked dimers than for the NMA-flanked ones: Orb_d3-v2 predicted a value of 1.5 kcal/mol, while Orb-v2 yielded 2.8 kcal/mol, both in reasonable agreement with the QM reference (Figure 4g).

Chain extension was performed by inserting additional NMA between the terminal NMA and the adjacent water molecule(s). Due to the contribution of terminal water molecules, the maximum cooperativity observed in SCS-MP2 calculations increased to 3.1 kcal/mol at the head H-bond and 3.9 kcal/mol at the tail H-bond (Figure 4a, c), compared to 2.2 kcal/mol in the NMA-only system. Among the MLFFs, MACE-OFF models closely reproduced the QM reference values, predicting cooperativities of 2.2-2.4 kcal/mol at the head and 2.6-3.5 kcal/mol at the tail. These results suggest that the polarizability of both water and NMA in MACE-OFF aligns well with QM behavior. ANI models still consistently underestimated cooperativity upon elongation, indicating limited response to environmental effects. Orb models continued to overestimate cooperativity, and terminal values exceeded those of the NMA-only system by about 2 kcal/mol, suggesting the Orb water model is reasonably accurate, while the primary source of deviation lies in its NMA representation. Moreover, the trend of increasing cooperativity with NMA extension under the Orb models was qualitatively consistent with QM, which indicates Orb may effectively capture interaction effects of longer range than ANI and MACE-OFF.

In addition to geometric failures observed in the Orb models, outliers in energetics were also identified for MACE-OFF and ANI due to imperfect geometry optimization. Compared to MM force fields, geometry optimization using MLFFs proved more challenging, especially for larger systems. This is consistent with the common assumption that the potential energy surfaces of MLFFs may be less smooth than those of classical FFs. As chain length increased, it became increasingly difficult to locate an energy minimum while maintaining the planarity required for analyzing H-bond cooperativity.

3.4 H-bond distance in NMA polymers

The H-bond distance serves as an indicator of interaction strength, with QM-optimized geometries showing progressively shorter bond lengths as cooperativity increases. Among the MLFFs, both ANI and MACE-OFF reproduced dimer H-bond distances in good agreement with QM, indicating their reliability in modeling direct interactions between two NMA molecules (Figure 5, Figure S4 for linear polymers). On the other hand, since Orb models failed to generate

H-bonded polymer geometries under standard energy minimization, they were excluded for evaluating H-bond distance trends.

Upon the addition of one NMA molecule on each side of a dimer, the H-bond distances predicted by MACE-OFF models, including 23-SC, correlated well with their corresponding cooperative energies (Figure 5a, b, c). Distance shortening was evident for both termini, with further reductions in the middle H-bonds of the polymers, consistent with the QM trend. These results also indicate that M-sized MACE-OFF models offer a more accurate description of H-bond distances compared to the S counterpart. Furthermore, as with the energy profiles, the magnitude of distance reduction in MACE-OFF variant was no longer pronounced for further NMA chain extension.

The cooperative effects on H-bond distance in ANI models were much weaker than MACE-OFF and notable deficiencies of optimization in longer polymer were also observed (Figure 5d, e, f). While ANI-1ccx captured a slight distance shortening consistent with cooperative effects, ANI-2x failed to reflect any significant trend. This discrepancy contrasts with their energy profiles, where ANI-2x exhibited more pronounced responses than ANI-1ccx.

In summary, MACE-OFF variants provided the best agreement with QM data for both cooperative energy and H-bond distances. Notably, the polarizability of both NMA and water was correctly captured in MACE-OFF models, reproducing the expected QM trends. Among MACE-OFF variants, M-sized models outperformed S and 23-SC models, but no significant differences were observed between MACE-OFF23 and MACE-OFF24. Although ANI models correctly reproduced NMA dimer H-bonding geometries, they failed to capture the cooperative effects on both energy and H-bond distance. Orb-v2 models were unable to reproduce the correct dimer geometry and drastically overestimated the cooperative energy for NMA, but the polarizability of water was better depicted. Additionally, while MACE-OFF and ANI captured cooperative effects only upon the addition of one NMA, Orb models more closely resembled the QM trend, where cooperative effects continued to accumulate with further NMA chain extension.

4 Discussion

Quantifying cooperativity provides a sensitive metric for evaluating a force field's ability to generalize across electrostatic environments and intermolecular interactions. In our previous study,[34] we benchmarked classical MM force fields using the same NMA polymer systems. Additive force fields, by construction, yield strictly zero cooperativity due to the absence of polarization in their potential energy functions. In contrast, polarizable models such as Drude and AMOEBA explicitly incorporate electronic induction and showed clear improvements over additive FFs in reproducing cooperative effects. However, even these advanced FF models underestimated cooperative energy by approximately 20% and 40%, respectively, when compared to QM references.

Leveraging the growing availability of high-level QM data and computational resources, MLFFs offer the potential to model complex systems with improved generality and accuracy. In this study, we assessed three representative MLFFs (ANI, MACE-OFF, and Orb) using the NMA benchmark system designed to probe electrostatic induction and dispersion contributions across increasing chain lengths. Despite differences in architectures and training data, all three models were trained on QM-calculated energies and forces with the goal of achieving broad chemical transferability. A key aspect of comparison lies in their capacity to capture polarizability and long-range interactions – factors central to H-bond cooperativity. As such, the cooperativity trend in NMA polymers provides a stringent test of how well each MLFF models many-body electrostatics across molecular chains.

None of the MLFFs fully reproduced the cooperative energy across all chain lengths with quantitative fidelity. Both ANI and MACE-OFF failed to capture the growth of cooperative effects as the number of NMA units exceeded two, highlighting limitations in modeling long-range polarization. ANI, while successful in modeling small molecules, consistently underestimated cooperative energy, reflecting its lack of explicit electrostatic induction. MACE-OFF, by contrast, achieved better agreement at short to medium ranges, likely due to its rotationally equivariant architecture and exposure to peptide-like fragments and solvated molecular complexes during the training. Nonetheless, its sensitivity plateaued beyond trimer systems, suggesting a finite non-bonded perception range.

Orb demonstrated the clearest monotonic rise in cooperativity with chain length, indicating a strong response to long-range correlations. This performance likely arises from its attention-based architecture that incorporate receptive field. However, Orb significantly overestimated the magnitude of cooperative energy, implying an over-polarizable potential. This may result from its pre-training strategies that prioritize data diversity and generalization but do not sufficiently sample bio-organic environments. While Orb captures the qualitative trend of cooperativity, its quantitative predictions deviates substantially from QM references.

These findings highlight a broader challenge in MLFF development: while modern architectures can encode complex spatial correlations, accurately capturing cooperative effects requires more than architectural sophistication or transfer

learning. The large discrepancies between ML models reflect a combination of architectural biases, training data gaps, and the difficulty of learning subtle many-body polarization phenomena from general-purpose datasets.

Our results point to the need for better benchmarks and descriptors of cooperativity. Current MLFF evaluations often focus on local accuracy metrics (e.g., forces, energies, torsion scans), which may overlook systematic deficiencies in long-range interactions. As cooperative H-bonding is ubiquitous in biomolecular systems, we propose that quantifying cooperativity in controlled model systems should become a standard component of MLFF validation. Adopting this approach can help pinpoint force field limitations and inform the development of next-generation neural network potentials that incorporate inductive and many-body effects in a physically meaningful way.

5   Conclusion

This study evaluated the capability of MLFFs to reproduce cooperative H-bonding interactions in NMA polymers, a representative system for biomolecular noncovalent interactions. By quantifying the cooperative energy and structural changes, we assessed three state-of-the-art MLFFs — ANI, MACE-OFF, and Orb — against high-level QM calculations. Significant differences were observed among the models, demonstrating that H-bond cooperativity can serve as a critical benchmark for evaluating the accuracy and transferability of MLFFs.

Universal MLFFs have rapidly evolved with the general aim of simulating complex chemical and biological systems. The benchmark results presented in this work underscore the importance of evaluating MLFFs not only in terms of pairwise accuracy but also in their treatment of collective many-body phenomena. H-bond cooperativity provides a rigorous and physically interpretable benchmark for assessing long-range induction and dispersion effects, which are crucial for accurately modeling biomolecules. Our findings suggest that while current MLFFs show promising accuracy for isolated interactions, further development is needed to better account for cooperative behavior across extended systems, particularly in biological and condensed-phase applications.

Acknowledgements

This work was supported by the "Pioneer" and "Leading Goose" R&D Program of Zhejiang (2023C03109, 2024SSYS0036), the National Natural Science Foundation of China (32171247, 21803057). We thank the Westlake University Supercomputer Center for computational resources and related assistance.

Conflict of interest

No conflict of interest

References

[1] Oliver T. Unke, Stefan Chmiela, Huziel E. Sauceda, Michael Gastegger, Igor Poltavsky, Kristof T. Schütt, Alexandre Tkatchenko, and Klaus-Robert Müller. Machine Learning Force Fields. *Chemical Reviews*, 121(16):10142–10186, August 2021.

[2] Yuanqing Wang, Kenichiro Takaba, Michael S. Chen, Marcus Wieder, Yuzhi Xu, Tong Zhu, John Z. H. Zhang, Arnav Nagle, Kuang Yu, Xinyan Wang, Daniel J. Cole, Joshua A. Rackers, Kyunghyun Cho, Joe G. Greener, Peter Eastman, Stefano Martiniani, and Mark E. Tuckerman. On the design space between molecular mechanics and machine learning force fields, September 2024.

[3] Ye Ding and Jing Huang. Implementation and Validation of an OpenMM Plugin for the Deep Potential Representation of Potential Energy. *International Journal of Molecular Sciences*, 25(3):1448, January 2024.

[4] Ye Ding and Jing Huang. DP/MM: A Hybrid Model for Zinc–Protein Interactions in Molecular Dynamics. *The Journal of Physical Chemistry Letters*, 15(2):616–627, January 2024.

[5] Ye Ding, Kuang Yu, and Jing Huang. Data science techniques in biomolecular force field development. *Current Opinion in Structural Biology*, 78:102502, February 2023.

[6] Wenyi Zhang, Kaiyue Zhang, and Jing Huang. A Simple Way to Incorporate Target Structural Information in Molecular Generative Models. *Journal of Chemical Information and Modeling*, 63(12):3719–3730, June 2023.

[7] Huayang Liu, Zilin Song, Yu Zhang, Bihan Wu, Dinghao Chen, Ziao Zhou, Hongyue Zhang, Sangshuang Li, Xinping Feng, Jing Huang, and Huaimin Wang. De novo design of self-assembling peptides with antimicrobial activity guided by deep learning. *Nature Materials*, pages 1–12, March 2025.

[8] David Rosenberger, Justin S. Smith, and Angel E. Garcia. Modeling of Peptides with Classical and Novel Machine Learning Force Fields: A Comparison. *The Journal of Physical Chemistry B*, 125(14):3598–3612, April 2021.

[9] Jörg Behler and Michele Parrinello. Generalized Neural-Network Representation of High-Dimensional Potential-Energy Surfaces. *Physical Review Letters*, 98(14):146401, April 2007.

[10] J. S. Smith, O. Isayev, and A. E. Roitberg. ANI-1: An extensible neural network potential with DFT accuracy at force field computational cost. *Chemical Science*, 8(4):3192–3203, March 2017.

[11] Linfeng Zhang, Jiequn Han, Han Wang, Roberto Car, and Weinan E. Deep Potential Molecular Dynamics: A Scalable Model with the Accuracy of Quantum Mechanics. *Physical Review Letters*, 120(14):143001, April 2018.

[12] Simon Batzner, Albert Musaelian, Lixin Sun, Mario Geiger, Jonathan P. Mailoa, Mordechai Kornbluth, Nicola Molinari, Tess E. Smidt, and Boris Kozinsky. E(3)-equivariant graph neural networks for data-efficient and accurate interatomic potentials. *Nature Communications*, 13(1):2453, May 2022.

[13] Ilyes Batatia, Dávid Péter Kovács, Gregor N. C. Simm, Christoph Ortner, and Gábor Csányi. MACE: Higher Order Equivariant Message Passing Neural Networks for Fast and Accurate Force Fields, January 2023.

[14] Mark Neumann, James Gin, Benjamin Rhodes, Steven Bennett, Zhiyi Li, Hitarth Choubisa, Arthur Hussey, and Jonathan Godwin. Orb: A Fast, Scalable Neural Network Potential, October 2024.

[15] Christian Devereux, Justin S. Smith, Kate K. Huddleston, Kipton Barros, Roman Zubatyuk, Olexandr Isayev, and Adrian E. Roitberg. Extending the Applicability of the ANI Deep Learning Molecular Potential to Sulfur and Halogens. *Journal of Chemical Theory and Computation*, 16(7):4192–4202, July 2020.

[16] Dávid Péter Kovács, J. Harry Moore, Nicholas J. Browning, Ilyes Batatia, Joshua T. Horton, Venkat Kapil, William C. Witt, Ioan-Bogdan Magdău, Daniel J. Cole, and Gábor Csányi. MACE-OFF23: Transferable Machine Learning Force Fields for Organic Molecules, December 2023.

[17] Dávid Péter Kovács, J. Harry Moore, Nicholas J. Browning, Ilyes Batatia, Joshua T. Horton, Yixuan Pu, Venkat Kapil, William C. Witt, Ioan-Bogdan Magdău, Daniel J. Cole, and Gábor Csányi. MACE-OFF: Transferable Short Range Machine Learning Force Fields for Organic Molecules, January 2025.

[18] Alison N. Leonard, Andrew C. Simmonett, Frank C. IV Pickard, Jing Huang, Richard M. Venable, Jeffery B. Klauda, Bernard R. Brooks, and Richard W. Pastor. Comparison of Additive and Polarizable Models with Explicit Treatment of Long-Range Lennard-Jones Interactions Using Alkane Simulations. *Journal of Chemical Theory and Computation*, 14(2):948–958, February 2018.

[19] You Xu and Jing Huang. Validating the CHARMM36m protein force field with LJ-PME reveals altered hydrogen bonding dynamics under elevated pressures. *Communications Chemistry*, 4(1):1–11, June 2021.

[20] Dylan M. Anstine and Olexandr Isayev. Machine Learning Interatomic Potentials and Long-Range Physics. *The Journal of Physical Chemistry A*, 127(11):2417–2431, March 2023.

[21] A. Subha Mahadevi and G. Narahari Sastry. Cooperativity in Noncovalent Interactions. *Chemical Reviews*, 116(5):2775–2825, March 2016.

[22] Yang Li, Yiguang Wang, Gang Huang, and Jinming Gao. Cooperativity Principles in Self-Assembled Nanomedicine. *Chemical Reviews*, 118(11):5359–5391, June 2018.

[23] Christopher A. Hunter and Harry L. Anderson. What is Cooperativity? *Angewandte Chemie International Edition*, 48(41):7488–7499, 2009.

[24] Pedro E. M. Lopes, Jing Huang, Jihyun Shim, Yun Luo, Hui Li, Benoît Roux, and Alexander D. MacKerell. Polarizable Force Field for Peptides and Proteins Based on the Classical Drude Oscillator. *Journal of Chemical Theory and Computation*, 9(12):5430–5449, December 2013.

[25] Fang-Yu Lin, Jing Huang, Poonam Pandey, Chetan Rupakheti, Jing Li, Benoı T. Roux, and Alexander D. MacKerell. Further Optimization and Validation of the Classical Drude Polarizable Protein Force Field. *Journal of Chemical Theory and Computation*, 16(5):3221–3239, May 2020.

[26] Pengyu Ren and Jay W. Ponder. Polarizable Atomic Multipole Water Model for Molecular Mechanics Simulation. *The Journal of Physical Chemistry B*, 107(24):5933–5947, June 2003.

[27] Jay W. Ponder, Chuanjie Wu, Pengyu Ren, Vijay S. Pande, John D. Chodera, Michael J. Schnieders, Imran Haque, David L. Mobley, Daniel S. Lambrecht, Robert A. Jr. DiStasio, Martin Head-Gordon, Gary N. I. Clark, Margaret E. Johnson, and Teresa Head-Gordon. Current Status of the AMOEBA Polarizable Force Field. *The Journal of Physical Chemistry B*, 114(8):2549–2564, March 2010.

[28] Hong Guo and Martin Karplus. Ab initio studies of hydrogen bonding of N-methylacetamide: Structure, cooperativity, and internal rotational barriers. *The Journal of Physical Chemistry*, 96(18):7273–7287, September 1992.

[29] Lars Ojamaee and Kersti Hermansson. Ab Initio Study of Cooperativity in Water Chains: Binding Energies and Anharmonic Frequencies. *The Journal of Physical Chemistry*, 98(16):4271–4282, April 1994.

[30] Hong Guo, Nohad Gresh, Bernard P. Roques, and Dennis R. Salahub. Many-Body Effects in Systems of Peptide Hydrogen-Bonded Networks and Their Contributions to Ligand Binding: A Comparison of the Performances of DFT and Polarizable Molecular Mechanics. *The Journal of Physical Chemistry B*, 104(41):9746–9754, October 2000.

[31] Nadya Kobko and J. J. Dannenberg. Cooperativity in Amide Hydrogen Bonding Chains. Relation between Energy, Position, and H-Bond Chain Length in Peptide and Protein Folding Models. *The Journal of Physical Chemistry A*, 107(48):10389–10395, December 2003.

[32] Yun-Dong Wu and Yi-Lei Zhao. A Theoretical Study on the Origin of Cooperativity in the Formation of 310- and α-Helices. *Journal of the American Chemical Society*, 123(22):5313–5319, June 2001.

[33] Alexandre V. Morozov, Kiril Tsemekhman, and David Baker. Electron Density Redistribution Accounts for Half the Cooperativity of α Helix Formation. *The Journal of Physical Chemistry B*, 110(10):4503–4505, March 2006.

[34] You Xu and Jing Huang. Quantifying the cooperativity of backbone hydrogen bonding. *bioRxiv : the preprint server for biology*, 2025.

[35] Justin S. Smith, Benjamin T. Nebgen, Roman Zubatyuk, Nicholas Lubbers, Christian Devereux, Kipton Barros, Sergei Tretiak, Olexandr Isayev, and Adrian Roitberg. Outsmarting Quantum Chemistry Through Transfer Learning, July 2018.

[36] Benjamin Rhodes, Sander Vandenhaute, Vaidotas Šimkus, James Gin, Jonathan Godwin, Tim Duignan, and Mark Neumann. Orb-v3: Atomistic simulation at scale, April 2025.

[37] J. Harry Moore, Daniel J. Cole, and Gabor Csanyi. Computing hydration free energies of small molecules with first principles accuracy, May 2024.

[38] Ask Hjorth Larsen, Jens Jørgen Mortensen, Jakob Blomqvist, Ivano E Castelli, Rune Christensen, Marcin Dułak, Jesper Friis, Michael N Groves, Bjørk Hammer, Cory Hargus, Eric D Hermes, Paul C Jennings, Peter Bjerre Jensen, James Kermode, John R Kitchin, Esben Leonhard Kolsbjerg, Joseph Kubal, Kristen Kaasbjerg, Steen Lysgaard, Jón Bergmann Maronsson, Tristan Maxson, Thomas Olsen, Lars Pastewka, Andrew Peterson, Carsten Rostgaard, Jakob Schiøtz, Ole Schütt, Mikkel Strange, Kristian S Thygesen, Tejs Vegge, Lasse Vilhelmsen, Michael Walter, Zhenhua Zeng, and Karsten W Jacobsen. The atomic simulation environment—a Python library for working with atoms. *Journal of Physics: Condensed Matter*, 29(27):273002, June 2017.

[39] Peter Eastman, Raimondas Galvelis, Raúl P. Peláez, Charlles R. A. Abreu, Stephen E. Farr, Emilio Gallicchio, Anton Gorenko, Michael M. Henry, Frank Hu, Jing Huang, Andreas Krämer, Julien Michel, Joshua A. Mitchell, Vijay S. Pande, João PGLM Rodrigues, Jaime Rodriguez-Guerra, Andrew C. Simmonett, Sukrit Singh, Jason Swails, Philip Turner, Yuanqing Wang, Ivy Zhang, John D. Chodera, Gianni De Fabritiis, and Thomas E. Markland. OpenMM 8: Molecular Dynamics Simulation with Machine Learning Potentials. *The Journal of Physical Chemistry B*, 128(1):109–116, January 2024.

Figure 1: **Scheme of cooperativity calculation and models.** (a) Cooperativity is the difference of dimer interaction between two-body and three-body systems. (b) Two conformations of NMA dimer optimized at the ωB97XD/cc-pVTZ level of QM. (c) An arc decamer with exclusively syn conformation of dimer blocks. (d) A linear decamer with alternating syn and anti blocks. In (c) and (d), the water capping sites on termini are illustrated; and the cooperative energies calculated for the first, middle, and last H-bonds to evaluate the effect of NMAs extending on B-side, both sides and A-side, respectively, are highlighted with transparent boxes.

Figure 2: **Dimer interaction of each model.** (a) syn conformation; (b) anti conformation. The x-axis indicates the group of force fields and legend of variants is shown on the right.

Figure 3: **Comparison of H-bond cooperative energies in arc-shaped NMA polymers calculated using QM and MLFFs.** The x-axis represents the number of NMA units in the polymer, while the y-axis shows cooperative energy in kcal/mol. Black dots and lines represent the QM reference, and colored dots and lines correspond to various MLFF predictions as indicated in the legends. (a-c) Cooperative energies for head, middle, and tail H-bonds using MACE-OFF models. Legend labels SC, S, and M denote the soft-core, S, and M model variants, respectively. (d-f) Cooperative energies for head, middle, and tail H-bonds using ANI models. (g-i) Cooperative energies for head, middle, and tail H-bonds using Orb models.

Figure 4: **Cooperative energy in arc-shaped NMA polymers in the presence of water molecules.** Data organization and representation are identical to those in Figure 3.

Figure 5: **Comparison of H-bond distances in arc-shaped NMA polymers calculated using QM and MLFFs.** The x-axis represents the number of NMA units in the polymer, while the y-axis shows the distance between O_A and N_B. Other data organization and representation are identical to those in Figure 3.

		training dataset			architecture		long-range effect	
		elements	level of theory	dataset	model	input	cutoff (Å)	electrostatics
MACE-OFF	23-SC	H, C, N, O, F, P, S, Cl, Br, I	ωB97M-D3(BJ)/ def2-TZVPPD	SPICE v1 (neutral charge) + QMugs + water cluster (≤ 50)	higher-order equivariant message passing (2-layer) GNN	node with spherical harmonics	5	not explicitly included
	23-SMALL						4.5	
	23-MEDIUM						5	
	23b-MEDIUM						6	
	24-MEDIUM			SPICE v2 (updated PubChem molecules and amino acid -ligand pairs) + water cluster (≤ 90)			6	
ANI	ANI-1ccx	H, C, N, O	CCSD(T)*/CBS†	enhanced ANI-1x dataset (with active learning)	Feed-forward NN with modified BP symmetry	atomic environment vectors	~5	not explicitly included
	ANI-2x	H, C, N, O, F, S, Cl	ωB97x/631G*	ANI-2x dataset				
Orb	Orb-v2	89 elements	DFT with PBE	MPtraj + Alexandria datasets	GNN with deep layer and diffusion pretraining	node with radial basis functions	adjustable to 10Å (≤ 20 neighbors)	receptive field
	Orb-d3-v2		DFT with PBE + D3					
	orb-v3		AIMD	OMat24 dataset				

† computationally efficient approximation of CCSD(T)/CBS

Table 1: **Feature comparison of the MLFFs evaluated in this study.**