Perceptual Prediction Error Supports Implicit Process in Motor Learning

Xiaoyue Zhang[1], Wencheng Wu[1], Kunlin Wei[1,2,3,4] *

[1]School of Psychological and Cognitive Sciences, Peking University, Beijing, China
[2]Beijing Key Laboratory of Behavior and Mental Health, Beijing, China
[3]Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, China
[4]National Key Laboratory of General Artificial Intelligence, Beijing, China

***Correspondence to: Kunlin Wei
School of Psychological and Cognitive Sciences
Peking University
Yiheyuan Road
Beijing, 100871, China
wei.kunlin@pku.edu.cn

Abstract

Error-based learning underlies motor learning, but what specific motor error drives implicit learning, the procedural component of motor skill, is unclear. A typical action consists of a movement and a performance outcome, e.g., grabbing a coffee cup involves a reaching movement and its actual landing of the hand relative to the target cup. While performance error is fundamental for the cognitive component of motor learning, what error, either performance or movement prediction error, underlies implicit motor learning has not been resolved. These two errors are hard to disentangle as the performance outcome is an integral part of the movement. Here we used the classical visuomotor adaptation paradigm, in which people learn to counter visual perturbations by deliberately aiming off the target, to dissociate the performance error from the prediction error. Using a series of behavioral experiments and model comparisons, we revealed that movement prediction error, but not performance error, can parsimoniously explain diverse learning effects. Importantly, despite the perturbation is visual, the movement prediction error is not specified in visual terms, but determined by a perceptual estimate of the hand kinematics. In other words, contrary to the widely-held concept of sensory prediction error, a perceptual prediction error drives implicit motor learning.

Introduction

Learning from errors is a fundamental aspect of human cognition, particularly in the realms of perception and action [1-3]. A classic example is the basketball player's effort to improve their jump shot; when a shot lands left of the basket, the player consciously adjusts their aim to the right in subsequent attempts. This explicit learning is complemented by an implicit learning process, which involves subconscious fine-tuning of bodily movements without conscious correction or even awareness of these fine-tunes. While performance error is recognized as a key driver of explicit learning across various motor tasks [4-6], the specific errors that facilitate implicit learning—a defining feature of motor skill acquisition—are still a matter of intense debate.

The debate on motor errors centers around two primary categories: performance error or movement prediction error [7]. Performance error is typically defined as the discrepancy

between movement outcome and the intended goal [8-13], while movement prediction
error refers to the discrepancy between the executed movement and its predicted sensory
consequences [14-18]. The complexity of dissociating these two errors arises from their
simultaneous occurrence during action execution. For instance, when reaching for a
coffee cup, the desired trajectory of the hand and the desired final position must both be
considered [19]. The trajectory prescribes a movement prediction error, and the final
position prescribes a performance error, both theoretically leading to implicit learning for
subsequent movements [2, 14, 18, 20].

Prominent computational models of motor learning [8-13] and foundational textbooks [21]
have primarily focused on performance error (PE) to explain implicit learning behaviors.
In contrast, a growing body of behavioral and neurophysiological research highlights the
role of sensory prediction error (SPE) —the difference between the actual movement
trajectory and the intended movement trajectory—as a critical factor in driving implicit
learning [14, 15, 18, 22-26]. Resolving this theoretical inconsistency in the understanding
of implicit motor learning is not only essential for cognitive science [7] but also promises
to translate algorithmic insights from human procedural learning into applied fields such
as robotic control and artificial intelligence [27].

In this study, we harness the classical visuomotor rotation (VMR) adaptation paradigm —
an established framework for investigating error processing in motor learning—to
elucidate the competing error signals that contribute to implicit learning [18, 20]. During
the VMR task, participants encounter a hand cursor that moves in a skewed direction
relative to their actual hand motion, prompting them to re-aim their movements away from
the intended target. This dissociation between movement prediction (re-aiming) and goal
attainment (cursor motion) allows us to separate PE and SPE within a single action
(Figure 1A): PE is the deviation of the final cursor position from the target, while SPE is
characterized by the deviation of the visual cursor motion from the re-aiming movement.
Importantly, SPE is based solely on visual feedback, with the implicit assumption that the
motor system adapts itself based on the visual consequence of actions, i.e., the skewed
cursor motion. However, spatially locating one's moving body effector is the foundation

of motor control [28-30]. We thus propose a novel third category of error: perceptual

prediction error (PPE), which represent the discrepancy between the *perceived* hand

movement from the predicted re-aiming movement. Previously, the discrepancy between

the perceived hand movement and the intended movement has been proposed to drive

error-clamp adaptation [31, 32], a special case of VMR where explicit learning is absent.

However, in the error-clamp adaptation, the predicted direction and the target direction

are spatially aligned, and thus PE and PPE are indissociable. In VMR adaptation with

explicit learning, the re-aiming direction naturally dissociate these two errors, enabling an

opportunity to critically examine their roles.

Here we designed a series of six experiments that systematically manipulated VMR

perturbations, revealing distinct characteristics of implicit motor learning. Through

comprehensive behavioral analyses and modeling, we present compelling evidence that

perceptual prediction error, rather than performance error or sensory prediction error,

serves as the primary driver of implicit motor learning. This finding offers a parsimonious

explanation for our new results while reconciling them with established literature in the

field.

Results

Errors and models

In the VMR paradigm, participants need to move a directionally rotated cursor toward the

target, and the directional deviation of the cursor (x_v) from the target defines the PE

(Figure 1A). PE drives the participants to adapt their movements in the subsequent trial

by consciously re-aiming away from the target in the opposite direction of the rotation

perturbation [33]. This re-aiming reflects the participant's prediction of where their hand

will move, generating a sensory prediction of the reaching action (x_u). The deviation

between the actual cursor motion and this prediction is regarded as a sensory prediction

error, given its purely visual nature [7, 18]. However, we propose that since the predicted

variable pertains to the intended *hand* direction, the relevant feedback should also

concern the hand rather than the cursor [31]. Thus, the perceived hand $(\hat{x}_{hand})$, rather

than the cursor, is compared to the predicted hand (x_u), with their discrepancy constitutes

a perceptual prediction error. The perceived hand ($\hat{x}_{hand}$) is derived from a combination

of multiple sensory cues, including the visual cue (x_v) from the cursor, the proprioceptive

cue (x_p) from the actual hand, and the sensory prediction (x_u). These cues are integrated

in accordance with Bayesian principles, with each cue weighted by its relative reliability.

The three types of errors—PE, SPE, and PPE—exhibit distinct temporal profiles

throughout the learning process, such as during a 30° counterclockwise VMR (Figure 1B

& C). When the perturbation is introduced, the cursor initially deviates 30° from the target,

and through trial-by-trial gradual learning, the cursor home back toward the target. This

results in an exponential decrease in PE, driven by the gradual correction of hand

movement direction. In contrast, re-aiming behavior—which reflects explicit learning—

initially rises sharply and then exhibits a small decline over the course of learning [34].

The re-aiming behavior dictates the dynamics of the prediction errors (Figure 1C): SPE

remains large throughout the learning process because the re-aiming direction stays

consistently offset from the cursor. In contrast, PPE remains relatively small and stable

during learning, as the re-aiming direction is close to the perceived hand direction, which

is derived from a combination of sensory cues (x_u, x_v, and x_p; Figure 1A).

To determine which error best explains the diverse learning phenomena observed in

motor adaptation, we implemented a unified model framework. State-space models,

which have been successfully used to model error-based learning in visuomotor

adaptation [8, 10, 35, 36], were applied to test whether implicit learning is driven by PE,

SPE, or PPE (see Methods for model details).

The perceptual estimation of hand location is contingent on the relative reliability of the

individual sensory cues. Importantly, visual uncertainty for increases with larger SPEs

[31], as participants tend to fixate not on the target but closer to their re-aiming direction

during the movement [37, 38] (Figure S1). This results in the cursor deviating from the

fixation point by an angular separation equivalent to the magnitude of the SPE. Our

previous work demonstrated that visual uncertainty regarding cursor motion increases

linearly with this eccentricity [31]. We also confirmed that nonlinear, instead of linear,
eccentricity effect on visual uncertainty would not affect model comparison results (Figure
S4, Table S7). While large visual perturbations induce both large SPE and PE (at least
during early learning; Figure 1C), its effect on PPE is dampened due to the high visual
uncertainty associated with large perturbations. Visual uncertainty reduces the influence
of the visual feedback on the perceived hand direction ($\hat{x}_{hand}$) according to Bayesian
principles of perception [39]. By incorporating the perceptual estimate of hand direction
into the prediction error, the dynamics of PPE diverge significantly from those of PE and
SPE, thus predicting implicit learning differently.

Figure 1. Illustration of sensorimotor cues and error dynamics in visuomotor rotation
(VMR) learning. **(A)** Schematic representation of the key sensory cues and errors during
VMR learning. The rotated visual cursor, actual hand position, and re-aiming direction
correspond to the visual cue (x_v), proprioceptive cue (x_p), and sensory prediction (x_u),
respectively. Performance error (PE) is the discrepancy between the cursor and the target,
sensory prediction error (SPE) is the discrepancy between the cursor and the sensory
prediction, and perceptual prediction error (PPE) is the discrepancy between the
prediction and the perceived hand direction ($\hat{x}_{hand}$) combination, computed from a

Bayesian integration of x_v, x_p, and x_u. **(B)** Simulation of a 30° VMR learning using the

PPE model. The time-series of the variables from (A) are shown, with solid lines

representing variables that are directly observable, and the dashed line represents the

perceived hand direction ($\hat{x}_{hand}$), which is inferred but not directly observable. **(C)** Error

dynamics during learning under a 30° VMR perturbation. Simulated PE and SPE match

well with the observed data [34] represented by the dots. Distinctly, simulated PPE

remains small and stable, but quantitatively match our previous estimation of this error

[31].

Experiment 1: Perturbation size-dependency of implicit learning reflects the

dynamics of PPE, not PE or SPE

Manipulating the size of visual perturbations is a straightforward way to probe error-based

learning and to differentiate between distinct error signals. In VMR, larger perturbations

increase overall and explicit learning, but implicit learning remains stable across a broad

range of perturbation sizes [16, 34, 40, 41]. This classical finding has been interpreted as

a hard limitation of implicit learning [20, 32, 34, 40] or a result of competition of explicit

and implicit learning, where the explicit learning "siphons" away performance error [10].

The three error types—PE, SPE, and PPE—behave differently during early learning.

Both PE and SPE increase with larger perturbations, but PPE remains small (Figure 1B).

Previous studies have often overlooked early learning, focusing on late learning [10]. We

examined the entire learning process using various perturbation sizes (15°, 30°, 60°, and

90° VMR) by analyzing a prior dataset first [34] (Figure 2A). Since the participants

reported their aiming direction for each trial, we could calculate the PE and SPE

throughout learning (Figure 2B). As expected, larger perturbations led to an increase in

both PE and SPE during early learning, but the scaling effect maintains during late

learning only for SPE, not for PE (Figure 2B, middle and right; Figure 2D). Remarkably,

implicit adaptation did not scale with perturbation size. Instead, it remained unchanged or

even slightly decreased, both in early and late learning (Figure 2E; one-way ANOVA,

early: $F_{(3,36)} = 0.2$, $p = 0.897$; late: $F_{(3,36)} = 1.44$, $p = 0.247$). Neither the PE nor SPE

model captured this pattern, as both models predicted a linear increase in early implicit

learning with perturbation size. The SPE model also incorrectly predicted a linear increase in late implicit learning (Figure 2E). In contrast, the PPE model successfully explained across perturbation sizes for both early and late stages (Figure 2C & 2E), as PPE remains largely invariant across perturbation sizes (Figure 2B left & 2D). Model comparison showed that the PPE model significantly outperformed both the PE and SPE models (R^2 = 0.708, 0.177, and -0.730, BIC = 182.04, 260.69, and 528.63 for PPE, PE, and SPE models, respectively; Table S1).

The fact that PPE does not exhibit a linear increase with perturbation size, unlike PE and SPE, stems from the nature of perceptual estimation. PPE is based on the perceived hand direction, which integrates visual and proprioceptive cues, each weighted by its uncertainty. Prior models based on PE and SPE have overlooked the fact that visual uncertainty increases with perturbation size [31]. During VMR learning, participants tend to fixate near their re-aiming direction, not on the target [37, 38]. We confirmed this through a control experiment where we measured eye fixations during VMR learning (Figure S1). Larger perturbations had larger explicit learning with fixation further away from the target. This greater angular separation between the cursor and the fixation point increases visual uncertainty, reducing the influence of visual perturbation on hand localization ($\hat{x}_{hand}$) according to Bayesian principles [31]. By incorporating visual uncertainty into the PPE model, we captured the classical finding of invariant implicit adaptation over perturbation size. This result suggests the apparent limit on implicit learning is not a reflection of a fundamental constraint in the learning system, nor is it due to competition for performance error.

Figure 2. Behavioral and model fitting results from *Bond & Taylor, 2015*. **(A)** Total, explicit, and implicit learning during VMR adaptation with varying perturbation sizes. Explicit learning was estimated from participants' reported aiming directions. Solid lines represent the mean, and shaded areas depict the standard error (SEM) for each perturbation condition. **(B)** Changes in perceptual prediction error (PPE), performance error (PE), and sensory prediction error (SPE) during learning. PE and SPE were computed from data,

while PPE was estimated through model fitting. The three errors show distinct temporal profiles, particularly during early learning. **(C)** Model fits for implicit learning across perturbation sizes, with explicit learning treated as known data. The PPE model accurately replicated the implicit learning patterns. In contrast, the PE model performed poorly for large perturbations (60° and 90°), and the SPE model struggled with small perturbations (15° and 30°). **(D)** The absolute error magnitudes of PPE, PE, and SPE as a function of perturbation size during early (solid lines) and late (dashed lines) learning. While PE and SPE scale with perturbation sizes, PPE remain relatively stable across perturbations. **(E)** Data and model predictions for early (left) and late (right) implicit learning. Both early and late learning show minimal scaling across perturbation sizes, in contrast to the predictions of the PE and SPE models. The model predictions for implicit learning reflect the error dynamics illustrated in (D). Error bars represent SEMs for the data and bootstrap-derived standard deviations for the model.

The trial-by-trial aiming reports in *Bond & Taylor., 2015* enabled quantification of PE and SPE throughout learning. However, frequent aiming reports can alter the composition of explicit and implicit learning [33, 34, 42, 43]. To minimize this potential confound, we conducted Experiment 1, which largely replicated Bond et al.'s study but removed the trial-by-trial aiming reports. Instead, we inserted a small number of exclusion trials during late learning to specifically measure implicit learning [43-45]. Additionally, the PPE model predicts a biased perceived hand localization after each movement, which cannot be directly measured. In Experiment 1, we measured this bias indirectly by requiring the participants judge the location of their passively moved hand immediately after an adaptation trial [31] (Figure 3A; People's judgement should be biased in the direction of the visual perturbation, an effect termed as proprioceptive bias [45-50]. Importantly, we hypothesized that the magnitude of proprioceptive biases should correlate with the PPE and the implicit learning estimated by the model.

Experiment 1 recruited four independent groups of participants (n = 15 per group) to adapt to 15°, 30°, 60°, or 90° VMR (Figure 3A). The overall learning patterns were consistent with previous studies [34, 41, 51]. Implicit learning, measured through exclusion trials

during late adaptation, exhibited a slightly nonlinear dependency on perturbation size (Figure 3D). A one-way ANOVA revealed a significant main effect of perturbation size in implicit learning ($F(3,56) = 7.15$, $p = 0.0004$). Post-hoc tests showed that implicit learning was significantly smaller for the 90° perturbation compared to both 15° ($p = 0.046$) and 30° ($p < .0001$), suggesting a concave size-dependency when continuous aiming reports were not required.

Despite reasonable fits to total and explicit learning (Figure 3C), the PE and SPE models incorrectly predicted a scaling effect of implicit learning with perturbation size. This discrepancy arose PE and SPE scaled with the perturbation size (Figure 3B), as observed in Bond & Taylor's study (Figure 2B). In contrast, the PPE model accurately predicted implicit learning, showing that PPE first increased and then decreased with perturbation size, leading to a concave pattern that matches the empirical data (Figure 3B & C, $R^2 = 0.969$; Table S2). The model's parameters aligned with previous estimates, with implicit and explicit learning rates ($B_i = 0.151$ and $B_e = 0.513$) comparable to previous studies [8, 9]. The slope of visual uncertainty with perturbation size ($k = 0.204$) was slightly lower than that in the error-clamp paradigm ($k = 0.31$) [31], likely because participants in the VMR paradigm focus more on the cursor to ensure accurate performance. Eye-tracking data from the control experiment confirmed that participants tended to fixate back to the cursor after completing their movements (Figure S1B). This contrasts with the error-clamp paradigm, where participants are instructed to ignore the cursor and fixate on the target, resulting in higher visual uncertainty.

Figure 3. Results of Experiment 1. **(A)** Learning curves for VMR across different perturbation sizes. Learning starts after baseline, gradually reaches a steady-state, and then decreases significantly during the no-feedback washout phase. Colored lines represent the mean, and shaded areas represent the SEM for each condition. Dark and light gray shading regions indicate the timing of proprioception test trials ("Prop-test") and exclusion trials ("Ex-test"). The group average and SEM for implicit learning and proprioceptive bias are shown. Proprioceptive judgements are biased in the direction of the visual perturbation, which is opposite to the direction of learning. **(B)** Time course of the three error types, with data shown as solid lines and model estimates as dash lines. Shading denotes SEM. Similar to *Bond & Taylor, 2015*, perturbation size scales the early PE and overall SPE, but not PPE. **(C)** Model fitting results for total, explicit, and implicit learning. All three models reproduced the scaling effect of perturbation size for total and explicit learning. However, for implicit learning, only the PPE model accurately predicted the concave relationship observed in the data, whereas both PE and SPE models erroneously predicted a linear scaling effect. Error bars represent SEM for data and bootstrapped standard deviations for the model. The black dots represent individual data. **(D)** Implicit learning and proprioceptive bias as a function of perturbation size. Both exhibited a similar nonlinear relationship (* denote $p < .05$ and *** denote $p < .001$). Note that the proprioceptive bias has been sign-flipped here to facilitate comparison with

implicit learning. **(E)** Implicit learning was significantly correlated with proprioceptive bias on the individual level, consistent with the predictions of the PPE model. Each dot denotes the two variables measured in each condition.

The proprioceptive biases significantly correlated with implicit learning, as predicted by the PPE model. During late adaptation, participants' right hands, the hand used for reaching adaptation, were passively moved to a random position near the target, and they were asked to indicate the direction of the right hand using their left hand. This passive localization test revealed a proprioceptive bias towards the visual perturbation, a phenomenon known as proprioceptive recalibration [31, 45, 46, 48, 52]. Importantly, the bias followed a concave trend across perturbation size, mirroring the pattern of implicit learning (Figure 3D). A one-way ANOVA revealed a significant main effect of perturbation size ($F(3,56) = 3.7$, $p = 0.017$), with post-hoc test revealing a higher bias for 30° than for 90° perturbations ($p = 0.011$). A correlation analysis across participants showed a significant positive correlation between proprioceptive bias and implicit adaptation (Figure 3E; $R = 0.41$, $p = .001$). This high correlation aligns with the PPE model, as both measures are driven by the misperceived hand direction under visuomotor rotation ($\hat{x}_{hand}$, Figure 1). In the proprioception test trial, the proprioceptive judgment is biased by the misperception of the hand location experienced after the preceding adaptation trial ends [31], resulting in a small but consistent perceptual bias in the direction of visual perturbation.

In summary, implicit learning exhibited either a concave size-dependency [51] (Experiment 1) or invariance [34] across perturbation sizes, depending on the task specifics, such as the presence of frequent aiming reports. Neither the PE nor SPE model could account for these size-dependencies, as their error sources did not follow a similar size-dependency (Figure 2D & 3B). Interestingly, although PE converges a small value across different perturbation sizes during late learning, enabling the PE model to capture late learning size dependencies [10], it fails to account for early learning, where it erroneously predicts a size-scaling effect. Only the PPE model provides a parsimonious

explanation the size dependencies of implicit learning across both early and late learning
phases (Figures 2F & 3C).

Experiment 2: Stepwise perturbations enhance implicit learning due to increased
PPE

The differences between the three competing errors are most pronounced at the time
when a perturbation is initially introduced (Figure 1C, 2B & 3B). Therefore, if the
perturbation size changes *during* learning, we expect these errors to have diverging
effects. A previous study has shown that, compared to a single large perturbation (e.g.,
a 60° VMR, referred to as one-step perturbation herein), incrementally increasing
perturbations (e.g., a 15→30→45→60°, referred to as stepwise perturbation) boosts
implicit learning while reducing explicit learning [53]. The reduction in explicit adaptation
is expected due to smaller PE in stepwise conditions. However, the increase of implicit
learning challenges the conventional view that implicit learning is stereotypical and
unaffected by various manipulations [16, 34, 40, 41], and its mechanism remains unclear.
Importantly, stepwise adaptation offers a critical test of the three error models. For both
PE and SPE, one-step perturbation generates larger errors than stepwise perturbation
during early learning, predicting greater implicit learning for the one-step condition (Figure
4B, middle and right). In contrast, the PPE model predicts the opposite: smaller PPE for
one-step perturbation due to the discounting effect of visual uncertainty, and thus, less
implicit learning (Figure 4B, left).

To test these predictions in Experiment 2, we recruited two groups of participants: one
group experienced a one-step 60° VMR perturbation (n = 15), while the other group
experienced a stepwise 15°→30°→45°→60° VMR perturbation (n = 15; Figure 4A).
Implicit learning was assessed using exclusion trials during late learning. Both groups
achieved similar steady-state performance by the end of learning (two-sample *t*-test, $t(28)$
= -1.796, p = 0.083), but their implicit learning patterns diverged (Figure 4C). The stepwise
group showed increasing implicit learning with each perturbation step ($F(3,21)$ = 11.639,
p < 0.001), while the one-step group showed no significant change in implicit adaptation
over time ($F(3,21)$ = 1.938, p = 0.154). Notably, from the second exclusion trial block

onward, the stepwise group exhibited significantly greater implicit learning than the one-
step group (*p* = 0.011). By the final exclusion trial block (at 60° VMR), implicit learning in
the stepwise group was nearly double that of the one-step group (26.33° ± 2.74° vs. 12.74°
± 1.45°; *p* = .001). In contrast, explicit learning was consistently larger for the one-step
group for each exclusion trial block (*p* < 0.001).

Figure 4. Results of Experiment 2. **(A)** Learning of the one-step (yellow) and stepwise
(brown) group. The dark grey shading indicates the timing range for the proprioception
test trials, and the light grey shading indicates the timing range for the exclusion trials.
Squares with error bars represent implicit learning measured via exclusion trials ("Ex-
test"), while diamonds with error bars represent proprioceptive bias measured by
proprioception test trials ("Prop-test"). **(B)** Time courses of the errors. The solid lines and
dots represent data and the dashed lines represent model estimates. PE and SPE were
larger for one-step than for stepwise perturbations during early learning, while PPE

showed the opposite condition difference. **(C)** Time courses of explicit and implicit

learning. Both explicit and implicit learning increased over time for stepwise adaptation,

but remained unchanged for one-step adaptation. Note the difference between conditions:

while explicit learning was larger for one-step adaptation, implicit learning was larger for

stepwise adaptation especially during late learning (* denote $p < 0.05$, ** denote $p < 0.01$,

and *** denote $p < 0.001$). Error bars are SEMs for the data. The black dots represent

individual data. **(D)** Model fittings for implicit learning. Error bars are bootstrap-derived

standard deviations for the model (resample size = 5000). Only the PPE model replicated

the actual learning patterns, while both PE and SPE model predicted one-step adaptation

was larger than stepwise adaptation in most cases. **(E)** Time course of proprioceptive

bias in the two conditions. The sign of proprioceptive bias was flipped to aid the

comparison. With larger PPE, stepwise adaptation elicited larger proprioceptive bias than

one-step adaptation. **(F)** Implicit learning and proprioceptive bias plotted as a function of

perturbation size during stepwise adaptation. **(G)** Implicit learning was significantly

correlated with proprioceptive bias on the individual level. Each dot denotes the two

variables measured in each perturbation size during stepwise adaptation.

All three models predict the gradual increase in implicit learning with stepwise

perturbations, as all errors increase incrementally (Figure 4B & 4D). However, only the

PPE model correctly captures the relatively smaller implicit learning for the one-step

perturbation compared to stepwise adaptation (Figure 4D). The PE and SPE models

erroneously predict the opposite. Model comparison confirmed the superior performance

of the PPE model (R^2 = 0.960, 0.933, and 0.925; BIC = 264.81, 309.59, and 317.67 for

PPE, PE, and SPE models, respectively; more model comparisons, see Table S3). The

superior performance of the PPE model was obtained even though we used the visual

uncertainty parameter (k) estimated from Experiment 1, which also involved a one-step

perturbation.

As in Experiment 1, proprioceptive bias in this experiment correlated with implicit learning,

further supporting the PPE model. Proprioceptive bias increased with stepwise

perturbation (Figure 4E, $F(3,25) = 8.874$, $p < 0.001$), but remained unchanged during the

one-step perturbation ($F(3,25) = 0.538$, $p = 0.661$). These proprioceptive bias patterns

mirrored those of PPE and implicit learning (Figure 4F). To analyze the individual

difference, we used a mixed linear model to isolate the between-subject variability and

revealed a significant positive correlation between proprioceptive bias and implicit

learning ($\beta = 0.91$, $p < 0.001$; Figure 4G). This result was consistent with previous studies

on proprioception and motor adaptation [31, 32, 46, 48], supporting that implicit

adaptation and proprioceptive bias are driven by the same error source, i.e., PPE.

One might argue that the one-step perturbation amplifies explicit learning compared to

the stepwise perturbation, while the model-based analyses above assume fixed explicit

learning parameters across both conditions. Indeed, when faced with a large, salient

perturbation, individuals may rely more heavily on explicit learning strategies [10, 53]. To

address this, we allowed the explicit learning parameters to vary between the two

conditions and re-evaluated the models (Figure S2; Table S3). As expected, the explicit

learning rate was higher in the one-step condition across all models. However, even with

varying explicit learning parameters, both the PE and SPE models still failed to account

for the observed differences between conditions (Figure S2B). While the PE model

slightly improved its fit compared to the fixed-parameter case (Figure 4), it could not

capture the implicit learning patterns for small stepwise perturbations (15° and 30°, Figure

S2B). Furthermore, this improvement came at the cost of unrealistic model parameters

(Table S3). For example, the explicit learning rate in the stepwise condition was estimated

to be as low as 0.218, even smaller than the implicit learning rate (0.223), which

contradicts the typical finding that explicit learning rates are several times larger than

implicit learning rates [8, 9, 54]. The SPE model showed minimal improvement, continuing

to underestimate implicit learning in the stepwise condition and overestimate it in the one-

step condition (Figure S2B). In contrast, the PPE model continued to provide excellent

fits across both conditions, yielding the best overall fit ($R^2 = 0.962$, 0.958, and 0.926, BIC

= 268.76, 280.84, and 325.39 for PPE, PE, and SPE models, respectively).

We further validated the models using cross-validation, fitting each model with data from

the stepwise group and testing it on the one-step group. Although the PE and SPE models

fit the stepwise group reasonably well, their predictions for the one-step group were highly inaccurate (Figure S2C). Both models overestimated implicit learning and underestimated explicit learning in the one-step condition. In contrast, the PPE model provided a near-perfect match in the cross-validation test (Figure S2C; Table S3), further reinforcing its robustness in capturing the dynamics of both implicit and explicit learning across different perturbation schedules.

Experiment 3: Suppression of explicit learning uncovers nonlinear implicit learning compatible only with PPE

When a visual perturbation is first introduced, PE and SPE equal the perturbation size, as the participant has not yet developed a re-aiming strategy. In contrast, PPE is less than the visual perturbation because it derives multisensory cue combination for hand localization (Figure 1A). Thus, the learning to the very first perturbation can highlight the key differences between errors: both PE and SPE predict a linear learning function relative to perturbation size, while PPE predicts a nonlinear function, as larger perturbation increase visual uncertainty, reducing its own influence on hand localization [31, 55].

In Experiment 3 (n = 18), we used trial-by-trial random perturbations to suppress explicit learning and isolate the single-trial implicit learning. Unlike blocked perturbations as in Experiment 1 and 2, random perturbations prevent the accumulation of implicit learning. The participants performed single-target reaches, but each perturbation trial was perturbed by a visuomotor rotation with a pseudo-random size and direction (0, ±4°, ±8°, ±16°, ±32°, and ±64°), and was flanked by two no-feedback null trials (Figure 5A). Participants were instructed to aim directly at the target in the null trials, thus discouraging explicit learning. Implicit learning was measured by comparing hand movement directions between the pre- and post-perturbation null trials [31]. The perturbation sequence, though appeared random, was identical for all participants. The perturbation sequence had a zero mean, allowing any residual explicit learning to average out.

The participants displayed adaptive changes immediately after the perturbation sequence was introduced, as evidenced by increased variability in hand directions (Figure 5B). We fit the learning data with the three competing models. Since PE and SPE are equal to the perturbation size in single-trial learning, we compared only the PPE model and PE model. The PPE model outperformed the PE model in explaining the learning response to random perturbations (Figure 5B, R^2 = 0.471 vs -0.683; BIC = 46.56 vs 82.00). As predicted by the PPE model, implicit learning follows a nonlinear pattern: it increased with perturbation size until a point, then perturbation size grew larger (Figure 5C; repeated-measure ANOVA: $F(3.2, 53.8)$ = 45.897, $p < .0001$; 4° vs 8°: p = 0.005; 8° vs 32°: p = 0.001; 32° vs 64°: $p < .0001$). This nonlinear pattern was consistent with previous findings from various motor adaptation paradigms [31, 32, 51, 55]. In contrast, the PE (SPE) model predicted a linear pattern (linear regression: , with = -0.007 and = -0.064, R^2 = 0.908, F = 89.1, $p < .0001$), that drastically deviated from the observed data (Figure 5C; $R^2 < 0$, $RMSE$ = 2.449°), while the PPE model captured the nonlinear pattern well (R^2 = 0.954, $RMSE$ = 0.684°). Despite the randomness of the perturbations, the slope of visual uncertainty (k) estimated in this experiment was consistent with that from the block-design experiments (k = 0.195 from Experiment 2 and k = 0.204 from Experiment 1 and 2), further supporting the robustness of the PPE model across different experimental designs.

Figure 5. Results of Experiment 3. **(A)** Experimental design. In feedback (FB) trials, participants experienced visuomotor rotations with pseudo-random perturbation sizes and directions (0, ±4°, ±8°, ±16°, ±32°, and ±64°). Each feedback trial was flanked by two no-feedback (NFB) null trials, where participants were instructed to move directly to the target. Single-trial implicit learning was measured by the directional change between pre- and post-perturbation null trials. **(B)** Time course of hand movement directions across trials, along with model fits. The grey line (with shaded SEM) represents the data, with stars denoting feedback trials and circles representing no-feedback trials. The PPE model (red) provided a better fit to the data than the PE/SPE model (dark blue). **(C)** Single-trial learning as a function of perturbation size, derived from both trial-by-trial data and model fits. The PE/SPE model predicted a linear relationship between perturbation size and learning, while the PPE model predicted a nonlinear relationship. Color shades represent SEMs for the data and bootstrapped standard deviations for the model. The light grey lines represent individual participants' data.

Despite the task instructions, participants might have strategically re-aimed away from the visual target in the null trials, potentially "contaminated" the estimates of implicit

learning with explicit learning. To assess this, we employed a model-based analysis to
include an explicit learning component. The results from the PPE model revealed that the
estimated explicit learning parameters were quite small (learning rate B_e = 0.047 and
retention A_e = 0.037), indicating a negligible contribution from explicit learning.
Additionally, incorporating explicit learning into the model only marginally increase the R^2
but result in a higher BIC for the PPE model (R^2: 0.471 vs. 0.500; BIC = 46.56 vs. 52.82).
When we constrained the lower limit of A_e to typical values (> 0.5), the inclusion of explicit
learning actually worsened the model fit (R^2 = 0.471 vs. 0.453, BIC = 46.56 vs. 57.26).
Similarly, the fit of the PE/SPE models deteriorated when explicit learning was added
(Table S4). Notably, the PPE model still substantially outperformed the PE and SPE
models even with the inclusion of explicit learning (Table S4). Overall, we conclude that
random perturbations primarily lead to implicit learning with a nonlinear size dependency,
which can only be explained by the PPE model.

Experiment 4: Implicit learning elicited by instructed re-aiming is driven by PPE
One of the seminal studies in motor neuroscience supporting the role of SPE employed
the instructed re-aiming paradigm in VMR [15]. In this paradigm, participants are
instructed to re-aim to a neighboring target whose angular deviation from the original
target matches the imposed visual perturbation (e.g., 45°; Figure 6A). This re-aiming
nullifies performance error (PE) immediately. Intriguingly, despite this, participants begin
to drift away from the re-aiming target involuntarily over repeated trials, increasing PE.
This drift has been interpreted as evidence that SPE drives implicit learning [15]. Recently,
Albert et al. (2022) proposed an alternative explanation, suggesting that the drift could be
driven by PE relative to the re-aiming target (PE_re-aiming)—a form of PE that arises between
the cursor and the re-aiming target rather than the original target. Notably, this new PE
formulation is essentially equivalent to SPE (Figure 6B). However, our PPE could also
explain this drift: the perceived hand position is "pulled" toward the visual cursor, deviating
from the re-aiming target, thus generating a PPE and driving implicit learning. Our model
simulations indicate that all three error models (PE, SPE, and PPE) can explain the
original findings by Mazzoni & Krakauer (Figure 6C). However, the models make distinct
predictions when the re-aiming offset size changes. If the angular separation between the

re-aiming and original targets increases (e.g., from 45° to 90°), both SPE and PE_{re-aiming}

would predict a larger implicit drift, as the errors increase with the offset. In contrast, the

PPE model predicts a reduction in implicit drift because larger visual perturbations reduce

the influence of PPE due to increased visual uncertainty.

Figure 6. Data and model fitting for the instructed re-aiming paradigm (15) and

Experiment 4. **(A)** Instructed re-aiming paradigm illustration. Participants initially moved

a cursor toward a target without perturbation during baseline trials, followed by a single

rotation trial with a 45° visuomotor rotation. During subsequent instructed re-aiming trials,

they were instructed to move their hand toward a neighboring "re-aiming target" 45° away,

allowing the cursor to perfectly hit the original target. **(B)** Motor errors during instructed

re-aiming. Despite re-aiming to the new target, participants' hands unknowingly "drift"

away from the re-aiming target. In addition to the three errors shown in Figure 1A, a new

error termed re-aiming performance error (PE_{re-aiming})—the deviation of the visual cursor

from the re-aiming target—is introduced [10, 26]. This error is equivalent to SPE. **(C)**

Performance relative to the original target before and after instructed re-aiming. Data from

Mazzoni & Krakauer (2006). After the initial 45° rotation trial, participants performed

instructed re-aiming trials, with their hands drifting away from the re-aiming target. This

implicit drift can be explained equally well by the PPE, PE, and SPE models. **(D)** Results

of Experiment 4. Participants exhibited an implicit drift away from the intended direction

after instructed re-aiming, with the drift being significantly larger for the 45° re-aiming

offset compared to the 90° re-aiming offset. Colored lines with shaded areas represent

means and SEMs for the two conditions. **(E)** Extent of implicit learning observed and

predicted by the three competing models. The PPE model accurately predicts smaller

implicit learning for the 90° condition compared to the 45° condition, while both the PE

and SPE models erroneously predict the opposite. Error bars represent SEMs for the data

and bootstrap-derived standard deviations for the model. The black dots represent

individual data.

To test these diverging predictions, Experiment 4 recruited two groups of participants who

adapted to either a large (90°, n = 15) or small (45°, n = 15) re-aiming offset. Upon re-

aiming, participants' movements began to drift in the opposite direction of the perturbation.

Critically, the drift for the 90° group was significantly smaller than that for the 45° group

(Figure 6E). The final drift in the 90° condition was considerably smaller than in the 45°

condition (13.17° ± 1.29° for 45°, 8.44° ± 1.22° for 90°; two-sample t-test: $t(28) = 2.58$, p

= 0.016). The PPE model readily explains this condition difference, as it predicts smaller

implicit drift for larger re-aiming offsets due to the reduced influence of visual

perturbations on hand localization. In contrast, both the PE and SPE models incorrectly

predict that the drift should increase in the 90° condition (Figure 6E; Table S5). The visual

uncertainty parameter (k) estimated from this experiment (0.211) was similar to those in

previous experiments, further supporting the robustness of the PPE model.

Experiment 5: Performance error based on spontaneous re-aiming fails to induce

implicit learning

In Experiment 4, participants were instructed to re-aim, preventing the spontaneous

formation of explicit learning. However, one might argue PE and SPE could still account

for implicit drift if re-aiming occurred spontaneously rather than being explicitly instructed.

To address this, Experiment 5 introduced a re-aiming target after participants had fully

developed their own explicit learning. In this experiment, participants (n = 20) adapted to

a 45° VMR, with explicit learning intermittently measured using aiming reports during adaptation (Figure 7A). After reaching a steady-state learning phase (spontaneous re-aiming in Figure 7B), we presented a re-aiming target that aligned with each participant's last reported aiming direction. Participants were instructed to aim for this re-aiming target for 30 additional trials to assess potential implicit drift (instructed re-aiming in Figure 7B). Since this re-aiming target was consistent with the participants' spontaneous re-aiming direction, both SPE and PPE should remain unchanged (Figure 7C), and both models would predict no further implicit learning. In contrast, the PE model predicts that the introduction of the re-aiming target would create a new performance error ($PE_{re\text{-}aiming}$), leading to further implicit learning beyond the learning asymptote (Figure 7C).

Participants reported an average explicit learning of 29.53° ± 1.63° in their last aiming report, consistent with previous studies [8, 34]. However, no further increase in implicit learning occurred after the re-aiming target was introduced. The hand deviation from the original target with the re-aiming target (43.01° ± 0.50°) was statistically indistinguishable from the level before introducing the re-aiming target (43.25° ± 0.79°, $t(19) = 0.327$, $p = 0.747$, $BF = 0.24$). Moreover, implicit learning, as measured by exclusion trials, did not change when comparing the last exclusion trial cycle before re-aiming target introduction to the first trial cycle of the washout phase (9.04° ± 0.84° vs. 7.44° ± 0.66°, $t(19) = 1.888$, $p = 0.075$; Figure 7D). These findings show that introducing a re-aiming target after explicit learning has been formed does not induce further implicit learning, contradicting the PE model's predictions. Instead, the null result aligns with the prediction of the PPE model (Figure 7D), which also provided a better fit to the data ($R^2 = 0.880$ vs. 0.337 for the PE model, Table S6).

Figure 7. Results of Experiment 5. **(A)** Illustration of the re-aiming paradigm. Participants

reported their aiming direction using a numeric ring. After reaching asymptotic learning

for a 45° VMR, the last reported aiming direction was used to designate a re-aiming target

for each participant. **(B)** Learning in Experiment 5. Participants adapted to a 45° VMR

with explicit learning measured through verbal aiming reports and implicit learning

assessed using exclusion trials. After reaching the learning asymptote, participants

continued to aim at their chosen direction, now marked by a re-aiming target. The

introduction of the re-aiming target divides the learning into two phases, one for

spontaneous re-aiming and the other for instructed re-aiming. **(C)** Time courses of PPE,

PE, SPE, PE_re-aiming before and after the introduction of the re-aiming target. PE, SPE,

and PE_re-aiming were calculated from the data, while PPE was estimated by model fitting.

Note that the introduction of the re-aiming target results in a new PE_re-aiming, while PPE

and SPE remain unchanged. **(D)** Implicit learning observed and predicted by the models.
Implicit learning was measured using exclusion trials. The models were fit to the data
before the re-aiming target was introduced and used to predict learning afterwards. The
PE model failed to predict the observed re-aiming performance, while the PPE model
accurately predicted the absence of further implicit learning. Error bars represent SEMs
for the data and bootstrapped standard deviations for the model predictions. The black
dots represent individual data.

Experiment 6: Direct measures of PPE reveals its correlation to implicit learning
Thus far, PPE has been implicated through model analysis or indirectly measured by
proprioceptive bias in Experiments 1 and 2. In Experiment 6, we directly measured PPE
by asking participants to locate both their aiming and produced movement. Our model
simulation indicates that PPE should remain small throughout the learning process
(Figure 1C), and should exhibited a similar perturbation size-dependency as
proprioceptive bias and implicit learning (Figure 2D and 3D). To test these predictions,
we randomly assigned 30 participants to three conditions with 15°, 30°, and 90° VMR,
respectively (n = 10 each). For a probe trial, a reaching movement was preceded by a
pre-movement aiming report and followed by a post-movement report of perceived hand
direction (Figure 8A). The direction difference between these two reports reflects the
perceived error. Indeed, when the visual perturbation was turned on, the perceived hand
direction deviated away from the re-aiming direction and towards the visual perturbation
(Figure 8B). Thus, the PPE was significantly negative (15°: -1.99° ± 0.84°, $t(9)$ = -2.36, p
= 0.042; 30°: -4.39° ± 0.47°, $t(9)$ = -9.42, $p < 0.001$; 90°: -1.86° ± 0.65°, $t(9)$ = -2.87, p =
0.018), in the opposite direction of implicit learning (Figure 8C). Notably, the size of PPE
was rather small and remained small throughout learning, consistent with our initial model
simulations (Figure 1C). Importantly, PPE followed a nonlinear concave function of
perturbation size ($F(2,27)$ = 4.52, p = 0.020, post-hoc test: 30° vs. 90°: p = 0.038),
resembling the model-estimated PPE based on data from Bond 2015 and Experiment 1
(Figure 2D and 3D) and the proprioceptive bias measured in Experiment 1 (Figure 3D).
Furthermore, the concave PPE pattern revealed here was accompanied by a similar
concave pattern in implicit learning (Figure 8D; $F(2,25)$ = 5.04, p = 0.015, post-hoc test:

30° vs. 90°: *p* = 0.012). Across participants, PPE also positively correlated with implicit
learning (Figure 8E; *R* = 0.41, *p* = 0.031). Hence, the direct measures of PPE provide
further support for the driving role of PPE in implicit learning.
665

666
Figure 8. Results of Experiment 6. **(A)** Illustration of the probe trial that directly measures
PPE. Participants reported their aiming direction and perceived hand direction before and
after a perturbed reach, by placing a visual marker with their left hand via the KINARM

robot. The just-experienced visual perturbation biases the perceived hand direction away
from the aiming direction. **(B)** Learning performance and reported aiming/hand directions
from the three perturbation-size conditions are separately shown. Movement direction
relative to the target shows typical adaptation changes. Aiming direction and perceived
hand direction changes along with learning. Hand direction is more rotated than the
aiming direction, manifesting implicit learning; the aiming direction is more rotated than
the perceived hand direction, manifesting a perceptual error relative to the intended hand
direction (i.e., PPE). Shades and error bars denote SEM. **(C)** PPE as a function of probe
trials. PPE becomes negative with perturbation, in line with its causal role in driving
adaptation in the positive direction. **(D)** The magnitudes of both PPE and implicit learning
show a concave dependency on perturbation size. * denotes $p < 0.05$ for post-hoc
comparison. **(E)** Implicit learning significantly correlates with PPE as revealed by
generalized linear regression.
683
Discussion
Using a combination of systematic behavioral experiments and computational modeling,
our study provides converging evidence that explicit and implicit processes in
sensorimotor learning are driven by distinct error signals. Explicit learning is driven by
performance error, while implicit learning is governed by perceptual prediction error. By
leveraging the classical visuomotor rotation paradigm, we disentangled three potential
error signals for implicit learning—performance error, sensory prediction error, and
perceptual prediction error—across five experiments. Our findings demonstrate that
implicit learning exhibits a perturbation size-dependency that can only be explained by
PPE, not by PE or SPE [34] (Experiment 1 and 2). The perceptual prediction error can be
indirectly measured by proprioception tests, which show people's perceived hand is
biased and this bias in correlated with implicit learning (Experiment 1 and 2). When explicit
learning is suppressed using random perturbations, implicit learning follows a nonlinear
size-dependency, again consistent only with PPE (Experiment 3). When explicit learning
is constrained by instructed re-aiming (Experiment 4), implicit learning develops in a
pattern that directly contradicts the predictions of SPE-based learning [15]. When a re-
aiming strategy is introduced after explicit learning has already been established, no

further implicit learning occurs, refuting the predictions of PE-based learning (Experiment
5). Directly measured by probe trials, PPE is found to remain negative with a small yet
significant offset from the aiming direction (Experiment 6), a pattern drastically different
from PE and SPE yet matching the model predictions (Figure 1B and 1C). Individual
analysis reveals a positive correlation between PPE and implicit learning. Similar to
implicit learning and proprioceptive bias, the measured PPE shows a concave
dependency on perturbation size, further supporting its role in driving learning and biasing
proprioceptive judgment. Overall, these results challenge the prevalent theories that
attribute implicit motor learning to either performance or sensory prediction errors. Instead,
they highlight the critical role of perceptual prediction error, defined as the deviation of the
perceptual estimate of hand motion from the predicted (desired) motion, in driving implicit
motor learning.

Performance error does not drive implicit motor learning
Performance error is a well-established signal for learning in various domains, including
perceptual learning [56], reinforcement learning [57, 58], decision-making [59, 60], and
the explicit components of motor learning [8-10, 12, 18]. However, whether performance
error also underlies the implicit component of motor learning—traditionally associated
with procedural learning—remains a topic of heated debate. While the dominant view
posits that sensory prediction error drives implicit learning [7, 14, 15, 18, 22, 23], all the
influential state-space models endorse performance error as the key driver [8-10, 12]. A
major challenge in this debate is the difficulty of dissociating performance error from
movement prediction error, as performance error relates to goal attainment and is often
tightly coupled with the movement itself. Previous studies have attempted to separate
these errors by introducing "target jumps" during movement [23, 61-63], thus creating a
performance error that is less correlated with the movement. However, such
manipulations tend to distract participants, affecting both explicit and implicit learning [23,
61, 62]. Furthermore, the target jump is likely to be interpreted as an environmental
change, which leads to minimal learning within the motor system [55, 64]. Indeed, the
target jump has been shown to be ineffective in eliciting implicit learning [23, 65]. In
contrast, our study employed the classical VMR paradigm, which does not require

additional manipulations of the perturbation feedback. The presence of explicit re-
aiming—whether spontaneously learned or instructed—naturally separates the predicted
movement from the goal-seeking cursor, allowing us to dissociate movement prediction
error from performance error (Figure 1A). Our findings strongly suggest that performance
error cannot account for the various patterns of implicit motor learning observed, despite
its clear role in explicit learning.
738
It is worth noting that Albert and colleagues (2022) also employed a similar modeling
approach using extensive datasets from VMR paradigms, but they reached the
contrasting conclusion that both implicit and explicit learning are driven by the same
performance error. However, their analysis focused exclusively on late learning (final
learning extent) without considering the entire adaptation process. In late learning,
performance error and perceptual prediction error tend to converge in size (Figure 2B, 3B
& 7C), potentially leading to misleading model results. In our study, we demonstrated that
early learning—where PE and PPE differ significantly—reveals the limitations of the PE
model (Figure 2E). Additionally, Albert et al. did not compare stepwise adaptation with
one-step adaptation, a critical contrast that highlights the erroneous predictions of both
PE and SPE models (Figure 4D). In summary, by analyzing the full learning process, as
opposed to selective learning segments, our study refutes the notion that performance
error drives implicit motor learning.
752

Motor prediction error from multisensory integration, not a single modality
Our findings call for a shift in how prediction error is conceptualized in the context of motor
control and learning, emphasizing its perceptual nature rather than a strictly sensory one.
Traditional models of motor prediction error have largely focused on visual prediction
errors derived from the discrepancy between predicted and actual visual feedback during
movements [18, 20]. This approach has been applied not only to visual perturbations such
as visuomotor rotation and saccadic adaptation [66, 67], but also to non-visual
perturbations such as force [11, 68-70] and inertial perturbations [71, 72]. However,
accumulating evidence suggests that learning of force and inertial perturbations can occur
without visual feedback [73-76], contradicting the idea that visual errors alone drive motor

adaptation in these contexts. Defining prediction error solely in terms of visual metrics is

increasingly seen as inadequate, as several recent studies have noted [31, 32]. Despite

this, the field continues to label prediction errors in motor control as sensory prediction

errors, reinforcing their unimodal, primarily visual nature [18, 20]. Our study provides

strong evidence that visual prediction error cannot adequately capture the dynamics of

implicit learning, even when the perturbation itself is visual. For instance, SPE scales

dramatically with perturbation size (Figure 2B & 3B), leading to exaggerated predictions

regarding implicit learning, which often exhibits more stable or mild changes. In contrast,

perceptual prediction error (PPE), which is derived from multisensory integration, remains

more resilient to changes in visual perturbation size and can parsimoniously explain not

only our data but also key findings in the literature [15, 34, 53].

Our findings suggest that perceptual prediction error drives implicit learning beyond

upper-limb movements, as supported by emerging evidence from other motor learning

paradigms. One such example is saccadic adaptation, which explores how the

oculomotor system adapts to spatial perturbations of gaze targets [67, 77, 78]. In this

domain, performance error (i.e., the retinal distance between the post-saccadic target and

the fovea) does not seem to support implicit learning [67, 79]. Similar to reaching studies,

saccadic adaptation research has traditionally focused on sensory prediction error (i.e.,

the deviation between the post-saccadic visual error and an internally predicted visual

error [67, 79, 80]. However, recent modeling work suggests that both motoric and

perceptual changes in saccadic adaptation are better explained by a postdiction-based

error [81]. This postdiction error refers to the discrepancy between the eye-muscle motor

command and the perceived target location, the latter arising from a combination of the

actual saccadic error and the predicted saccadic displacement. This concept mirrors our

perceptual estimate of hand localization in reaching adaptation, wherein multiple

sensorimotor cues are integrated to inform the perceptual error driving adaptation.

Notably, saccadic adaptation assumes a stable target location across saccades,

reflecting the oculomotor system's assumption of spatial stability [82-84]. Similarly, in

reaching tasks, the manual control system assumes that the hand should consistently

and accurately reach the intended target. In this light, implicit learning functions to

recalibrate the motor system to ensure the maintenance of stability and accuracy, aided
by perceptual prediction error.
796
Perceptual prediction error may also underlie implicit learning in speech adaptation,
where the articulatory system adapts to auditory perturbations [85, 86]. Most studies in
speech adaptation emphasized auditory prediction error, a modality-specific form of
sensory prediction error [87-92]. However, recent modeling work has demonstrated that
a state estimate of the task, rather than sensory prediction error, best explains the
observed learning patterns [93, 94]. This state estimate is derived from a combination of
the previous articulatory state, an efference copy of the motor command, and auditory
feedback, creating a multimodal perceptual estimate that updates the articulatory-to-task
transformation—essentially, an internal model. Thus, even in the context of auditory
perturbations, a multimodal perceptual estimate, rather than a unimodal auditory
prediction error, appears to drive speech adaptation.
808
Theoretical advance of PPE model relative to previous work
Tsay et al. has recently proposed that implicit adaptation should be understood as a
process of proprioceptive re-alignment of the hand towards the target (PreMo model) [32].
They challenged the traditional visual-centric view and emphasized the critical role of
localizing one's effector in motor learning. However, as shown in our previous work [31],
the assumption of the PreMo model, i.e., the effect of visual perturbation on proprioceptive
feedback follows a ramp function of perturbation size, is not supported by empirical
findings [31, 46]. The PreMo model also predicts that the perceived hand direction should
be unbiased in late learning (i.e., re-aligned), contradicting to our findings in Experiment
6 which revealed a persistent perceptual error in late learning. Most importantly, the
PreMo model specifically aims to account for the error-clamp adaptation, a special form
of visuomotor rotation that invokes implicit process only [16]. Our PPE framework, instead,
aims to provide a unified account for both explicit and implicit learning processes, which
ubiquitously co-exist in various motor learning tasks [18].
823
Dual learning reflects the dual imperatives of motor control

The dual learning based on the two distinct error signals could reflect the dual imperatives
of motor control: achieving action goals and resolving sensory conflicts. Researchers
have shown that unconscious movements tend to resolve multisensory conflicts, as
opposed to achieving explicit goals [95-98]. For instance, in the rubber hand illusion
paradigm, participants exhibit subtle, unconscious movements to align visual and
proprioceptive cues, not to achieve a specific goal but to reduce sensory conflict [95, 96].
This suggests that the motor system, apart from controlling explicit actions, implicitly
adjusts movements to minimize sensory mismatches [99, 100]. Similarly, in the VMR
learning task, sensory feedback from different modalities (e.g., visual and proprioceptive)
creates a perceptual error relative to the predicted hand movement, leading the motor
system to implicitly recalibrate movements to resolve the conflict. Here explicit learning
corresponds to the goal-directed imperative of motor control, driven by PE. In contrast,
implicit learning aligns with the imperative of minimizing sensory conflict, driven by PPE.
838

The relationship between explicit and implicit learning
Our findings on PPE provide new insights into the relationship between explicit and
implicit learning, a central question in motor learning. Previous studies have suggested
that these two processes either complement each other [12, 33] or compete [10, 41, 101,
102]. However, our PPE model suggests that explicit and implicit learning are functionally
independent, driven by distinct error signals, and their relationship is task-specific. In
tasks where PE and PPE are aligned, explicit and implicit learning can work
synergistically, as seen in typical adaptation tasks (Experiments 1 and 2). However, when
these errors diverge, the two learning processes can conflict. For example, in Mazzoni &
Krakauer (2006) and Experiment 4, when participants were instructed to re-aim explicitly,
implicit learning caused a drift away from the re-aiming target, paradoxically worsening
performance (Figure 6). This conflict reveals that implicit learning, driven by PPE, can
sometimes undermine explicit learning, especially when their errors differ. Only when
performance error grows large enough and sustains for an extended period of time do
individuals adjust their explicit strategy to correct for performance error [26]. Similar
conflict or decoupling between explicit and implicit learning also occurs in the mirror-
reversal paradigm, in which implicit learning initially worsens performance [13, 103, 104],

even after a few days of training [103, 105, 106]. Only after sufficient exposure of this
conflict, people slowly develop explicit re-aiming strategies to override implicit learning
and thus improve the overall performance [103, 104]. Interestingly, in cases of conflict,
implicit learning—driven by the minimization of PPE—tends to be prioritized over
performance error correction. This is reminiscent of the competition between Type I
(unconscious, fast) and Type II (conscious, slow) cognitive processes in human decision-
making [11, 107].

Past research has reported that better explicit learners tend to exhibit less implicit learning,
implying a competitive relationship between the two processes [10, 41, 101, 102].
However, this conclusion is often based on inter-dependent learning measures, as implicit
learning is typically measured using exclusion trials, while explicit learning is inferred by
subtracting implicit learning from total learning [10, 41, 43, 51]. Given that most
participants reach a learning ceiling, this subtraction method risks producing a false
negative correlation. When explicit and implicit learning are measured independently—
e.g., using aiming reports for explicit learning and exclusion trials for implicit learning—
their relationship does not show a consistent negative correlation but varies depending
on the context. In some cases, there is even a positive correlation between the two [108]
(Figure S3). Additionally, during early learning, when learning is not yet complete with a
relatively large PE, the negative correlation between explicit and implicit learning
disappears (Experiment 5; Figure S3), challenging the notion that these processes
compete for a shared performance error signal. In summary, our findings suggest that
explicit and implicit learning are independent processes that can interact in a task-specific
way. Their relationship is not inherently competitive but rather context-dependent, driven
by the alignment or misalignment of the underlying error signals.

Proprioceptive bias resulted from perceptual bias related to perturbed action
Proprioceptive bias observed in motor adaptation highlights how motor learning can
modify perception. Our findings suggest that both motoric and perceptual changes during
motor learning might follow the principles of Bayesian cue integration. In implicit motor
learning, PPE arises from a misperception of the hand's position ($\hat{x}_{hand}$). Similarly, in

proprioception tests, biased judgments result from the same biased cue (provided by
previous adaptation trials. Consistent with this model, our results show that both
proprioceptive bias and implicit learning are modulated by ongoing adaptation (Figures 3
& 4). The model further explains the varied effects of perturbation size on proprioceptive
bias, which can be either linear [46, 47] or constant [52], an inconsistency left unexplained.
A linear dependency emerges with stepwise perturbations, where each step-up of
perturbation incrementally increases the perceived hand position [46, 47] (Experiment 2).
In contrast, a constant dependency is associated with one-step perturbations, leading to
similar across perturbation sizes [52] (Experiment 1). The significant correlation between
implicit learning and proprioceptive bias further supports the idea that changes in
perceived hand position underlie these perceptual shifts (Experiments 1 and 2). The
concept based on aligns with the view that proprioceptive bias results from multisensory
integration [45, 48-50]. A few studies have theorized that internal model formed by motor
adaptation underlies proprioceptive bias [50, 109, 110]. However, internal model typically
develops slowly, but proprioceptive bias develops rapidly with perturbation [45, 49, 111,
112] and diminishes rapidly without it [31]. This makes internal model an unlikely
explanation. Thus, Bayesian cue combination provides a unified explanation for both
implicit motor learning and accompanied perceptual changes.
905

Implications for neurophysiological studies
The compositional nature of motor learning—wherein explicit and implicit learning are
driven by distinct error signals—offers a new framework for exploring their neural
underpinnings. Traditionally, neurophysiological studies have focused on the neural
correlates of total learning, implicating both sensorimotor regions (e.g., cerebellum and
sensorimotor cortex [18, 113-117]; and regions associated with cognitive control and
decision-making (e.g., prefrontal cortex and basal ganglia [118-120]). By decomposing
total learning into explicit and implicit components, future research can better delineate
the neural circuits underlying these distinct processes [121], with particular focus on the
cerebellum and posterior parietal cortex (PPC) involved in learning acquisition [22, 117,
120, 122, 123]. Furthermore, most previous studies have used either sensory prediction
errors [114, 124-126] or performance errors [127-130] to identify neural correlates of

motor learning. Our finding that perceptual prediction error underlies implicit learning, with a distinct temporal profile from other errors, suggests more tailored neural assays could be developed [131]. Behavioral paradigms that effectively modulate perceptual prediction error, like the stepwise perturbations in Experiment 2, could be particularly useful for future investigations. Additionally, our results emphasize the importance of localizing one's effector in motor control and learning, requiring integration of motor prediction, visual, and proprioceptive cues on a trial-by-trial basis (Figure 1A). The neural substrates for encoding and integrating these cues overlap with key networks involved in visuomotor learning, including the premotor cortex for intended cursor movement [132, 133], PPC for learning-related visual representations [120] and visuomotor transformation [134], the cerebellum for error processing and acquisition of internal models [124-126], and the somatosensory cortex that contributing to the acquisition and retention of motor learning [135-137]. The interplay between motor learning and minimizing position errors of body effectors warrants further examination with neurophysiological approaches. By incorporating insights from our findings, future studies can better understand the neural mechanisms that support the distinct yet interconnected processes of explicit and implicit motor learning.

Methods

Participants

We recruited 204 college students from Peking University (118 females, 22.76 ± 2.53 years, mean ± SD). All participants were right-handed and had normal or corrected-to-normal vision. Participants were naïve to the purpose of the experiment.

Ethics Statement

The study was approved by the Ethical Committee of the School of Psychological and Cognitive Sciences at Peking University. All participants provided written informed consent to participate in the study and received monetary compensation upon completion of the experiment.

 Apparatus

In Experiments 1 and 2, we used the KINARM planar robotic manipulandum with a virtual-

reality system (BKIN Technologies Ltd., Kingston, Canada). Participants sat in a chair

and held the two robot handles with their left and right hands, respectively. The right hand

performed the reaching task, and the left hand indicated the perceived direction of the

right hand if required. A semi-silvered mirror positioned below eye level displayed visual

stimuli and obscured the participants' view of their hands and the robotic manipulandum.

The sampling rate of the KINARM was 1,000 Hz.

In Experiments 3, 4, and 5, we used a horizontally placed digitizing tablet (48.8 x 30.5 cm,

Intuos 4 PTK-1240, Wacom, Saitama, Japan) to measure people's hand movements.

Participants sat in front of a vertically-placed LCD screen (29.6 x 52.7 cm, Dell, Round

Rock, TX, US) with a customized wooden blinder placed over the tablet to occlude the

vision of the hand. Using a stylus with their right hand, participants performed the reaching

task on the tablet. The sampling rate of the tablet was 200Hz. Procedures and data

acquisition were customized coded (Psychtoolbox-3-3.0.19.2 in MATLAB).

The control experiment used a similar setup to Experiments 3, 4, and 5, with the addition

of eye tracking. A slightly smaller digitizing tablet (31.1 x 21.6 cm, Intuos M, Wacom,

Saitama, Japan) was used. The vertically-placed LCD screen was positioned

approximately 50cm in front of a chin rest, with a bracket occluding the participants' view

of the tablet and their hands. Movement trajectories were sampled at 200Hz by the

digitizing tablet. Eye movements were continuously tracked at 500Hz using a video-based

eye tracker (Eyelink 1000, SR Research), which was placed beneath the LCD screen.

Experiment 1

Sixty participants were randomly assigned to one of four groups (n = 15 each). Each

group adapted to visuomotor rotation of one of four possible sizes (15°, 30°, 60°, or 90°).

Participants made reaching movements from a central start position towards an up-right

(defined as 45°) target located 10 cm away. The start position, target, and cursor were

displayed as filled circles, with a 5-mm radius purple circle for the start position, a 5-mm

radius blue circle for the target, and a 3-mm radius white circle for the cursor. Each trial began with participants holding their right hand at the start position for 800ms. The target then appeared and signaled participants to make a quick, straight sliding movement through it. The trial ended when the distance between the hand and the start position exceeded 10 cm. If the movement duration from leaving the start position to reaching 10 cm exceeded 300 ms, a warning message, "too slow", would appear on the screen.

The experiment included three phases: a 35-trial baseline phase with veridical feedback, a 75-trial adaptation phase with VMR feedback, and a 20-trial washout phase without cursor feedback. The direction of the VMR perturbation was counterbalanced across participants. To assess implicit learning at the asymptote of learning, we inserted exclusion trials at the 50, 55, 60, and 65th trials during the adaptation phase. In these exclusion trials, participants were instructed to move their hand straight to the target without any strategic re-aiming. The washout phase used the same exclusion instruction, which the experimenter emphasized during data collection.

To measure proprioceptive bias, we inserted proprioception test trials at the 15, 20, 25, 30, and 35th trials during the adaptation phase (Figure 3A). Additional proprioception test trials were conducted at the 5, 10, 15, and 20th trials during the baseline phase to assess baseline proprioceptive bias. In the proprioception test trial, participants were instructed to relax their right arm, which was then passively moved to one of four locations (± 5° and ± 15° away from the reaching target). The passive movement lasted for 1000 ms and followed a straight path with a minimum-jerk speed profile. During this movement, a white ring, centered at the start position, expanded in sync with the hand's center-out displacement to indicate the movement distance. After the right hand reached the target location, participants held their right hand steady and used the left robot handle to indicate the perceived position of their right hand. Moving the left handle controlled the rotation of a visual "dial" displayed along an arc that passed through the proprioceptive targets (31). Participants held the dial at their perceived location for 1,000 ms, which was recorded as their perceptual judgement.

Experiment 2

Experiment 2 used the same task setup as in Experiment 1. We recruited two groups of

participants: one group experienced stepwise perturbations (n = 15), while the other group

experienced a one-step perturbation (n = 15). The stepwise group experienced a stepwise

15°→30°→45°→60° VMR perturbation, with each perturbation presented for 40 trials.

The one-step group experienced a 60° perturbation for 160 consecutive trials (Figure 4A).

Within each 40-trial perturbation block, we inserted exclusion trials at the 30 and 35th

trials to measure implicit learning, and proprioception test trials at the 5, 10, 15, and 20th

trials to measure proprioceptive bias. Before the adaptation phase, all participants

completed a 35-trial baseline phase with veridical feedback. After the adaptation phase,

they went through a washout phase without visual feedback, following the same exclusion

instruction.

1023

Experiment 3

Experiment 3 investigated the size-dependency of implicit learning when explicit learning

was suppressed by the randomness of perturbations. To prevent cumulative learning over

trials, three targets (25°, 45°, and 65° at an 8 cm distance) were used. Participants (n =

18) began with a baseline phase consisting of 15 veridical-feedback trials and 15

interleaved no-feedback null trials. In subsequent adaptation phase, trials were organized

into 6-trial mini blocks, where each target was presented once randomly and each

perturbation trial was flanked by two no-feedback null trials (Figure 5A). During

perturbation trials, participants were instructed to control the cursor to hit the target, while

null trials were served as exclusion trials. Visuomotor rotation of 0°, 4°, 8°, 16°, 32°, and

64°, either clockwise or counter-clockwise was randomly used in each perturbation trial.

These 11 perturbations were repeated nine times, leading to a total of 99 perturbation

trials and 99 null trials in the adaptation phase. The wide range of rotation sizes enabled

us to uncover the linear or nonlinear size-dependency of implicit adaptation [55, 138]. The

randomness of perturbation size and direction would prevent participants forming an

adaptive strategy. Single-trial learning was quantified as the difference in movement

direction between the two adjacent null trials before and after a perturbation trial.

1041

Experiment 4

Experiment 4 examined VMR adaptation with instructed re-aiming, as in the seminar
study by *Mazzoni & Krakauer, 2006*. However, two different sizes of VMR were used for
two separate groups of participants, i.e., 45° (n = 15) or 90° VMR (n = 15). The target,
presented as a 5-mm radius white circle, would appear at one of eight possible locations
(0°, 45°, 90°, 135°, 180°, 225°, 270°, and 315°) 8 cm away, randomly in an eight-trial
cycle. In the baseline phase, participants were instructed to control the cursor to hit the
target with veridical feedback. In the adaptation phase, a new re-aiming target was
introduced starting from the first perturbation trial. The re-aiming target, which was either
45° or 90° but in the opposite rotation direction of the imposed VMR, was presented as a
5-mm radius green circle, distinct from the original target. For a consecutive of 120 trials,
participants were required to move their hand to the re-aiming target to make the cursor
"hit" the original target. Half of the participants in each group experienced clockwise
perturbations and the other half counterclockwise perturbations. Typically, people could
maintain their aim at the re-aiming target despite their hand implicitly drifted away from it,
across the span of 120 trials [26].

1058

Experiment 5

Experiment 5 examined the instructed re-aiming introduced at the end of VMR adaptation,
as opposed to at its beginning (as in Experiment 4). Four targets were used, located at
45°, 135°, 225°, and 315°, 8 cm away, with trials organized into 4-trial cycles. The
baseline phase consisted of 3 trial cycles with veridical feedback trial and 1 trial cycle
without feedback. The subsequent adaptation phase consisted of 60 cycles with a 45°
VMR. Within the adaptation phase, we intermittently measured the explicit adaptation
through aiming reports at the 5, 19, 33, and 47th cycles. In these trials, the target was
surrounded by a ring of 63 numbered visual landmarks spaced 5.625° apart, with 0°
centered on the target. Before the reach, participants verbally reported the landmark
number they planned to reach. Their reported aiming directions were recorded by the
experimenter. To measure implicit adaptation, we inserted exclusion trials with no visual
feedback at the 6, 20, 34, 48th cycles. In the ensuing re-aiming phase, we presented a
re-aiming target at the last reported aiming direction, tailored to each individual participant.

Participants were required to move their hand straight to this re-aiming targe for 40 cycles.

Note that the re-aiming target would still make the cursor reached the original target since

it was aligned with the participant's spontaneous explicit learning asymptote. We inserted

exclusion trials at the 1st, 11, and 21th cycles in the re-aiming phase. In the final washout

phase, participants completed an additional 15 cycles of exclusion trials.

1078

Experiment 6

Experiment 6 aimed to directly measure PPE, defined as the difference between the

intended movement direction (aiming direction) and the perceived hand direction. Using

similar setup and procedures as Experiment 1 and 2, Experiment 6 recruited thirty

participants and randomly assigned them to learn one of three VMR perturbations (15°,

30°, and 90° VMR; n = 10 each). Participants went through three learning phases, i.e.,

baseline (20 trials), adaptation (44 trials), and washout (20 trials). Every four trials they

performed a probe trial for measuring PPE (Figure 8A). The first eight adaptation trials

were not probed to avoid affecting participants' formation of strategic re-aiming during this

critical initial learning phase [45, 139]. For each probe trial, participants held their right

hand at the start position for 800 ms, and then a target (a 5-mm radius blue circle), a

white arc, and a purple visual marker were displayed. Participants were asked to report

their intended aiming direction ("Where will your right hand go?") by moving the left

KINARM handle to dial the visual marker, just as in Experiment 2. Participants

subsequently performed the reaching movement, with or without the cursor perturbation,

by their right hand. After the reach, their right hand was passively returned to the starting

position by the robotic manipulandum. The arc and visual marker reappeared to prompt

participants to report the perceived direction of their actual hand movement ("Where was

your right hand?"), again using the left KINARM handle to dial the visual marker. PPE

was computed as the angular difference between these two reported directions.

1099

Control experiment with eye-tracking

The control experiment examined whether people fixate in the re-aiming direction during

reaching movements in typical VMR adaptation. Participants (n = 16) learned 45° and 90°

VMR perturbations in two consecutive sessions, with the order counterbalanced. Each

session included a baseline, adaptation, and washout phase, as in Experiment 1, with 40,

160, and 20 trials, respectively. Each trial began when the participant moved the cursor

(a 3-mm radius white circle) into the starting position (a 5-mm radius purple circle) using

a stylus. After holding the position for 500ms, a target (a 5-mm blue circle) was presented

10 cm away at a random direction between 0° and 360°. Participants started to reach the

target with the cursor once the target disappeared. The target presentation duration

followed a uniform distribution from 800 to 1200 ms, making the movement trigger less

predictable. If the movement was initiated before the target disappearance, the trial was

aborted and a text message "reacting too fast" would appear on the screen. Veridical or

perturbed cursor feedback was shown during the movement and the target re-appeared

once the movement ended and lasted for 1000 ms. The disappearance of the target

prevented some participants keeping their fixation at the starting position without change

to their re-aiming direction. To avoid slow movements, a text message "moving too slow"

would appear if the movement time exceeded 400ms.

1118

The three error models

Perceptual prediction error (PPE) model: The PPE model hinges on the perceptual

estimation of effector position that is dynamically updated through Bayesian cue

combination. For reaching, the perceptual estimate of the hand movement direction

$(\hat{x}_{hand})$ results from the integration of three sensory cues: the visual cue (x_v) from the

cursor, the proprioceptive cue (x_p) from the hand, and the sensory prediction of the

reaching action (x_u). Following Bayesian principles [39], the perceptual estimate is

$$1126 \qquad \hat{x}_{hand} = \sum_i W_i x_i, \ with \quad W_i = \frac{1/\sigma_i^2}{\sum_j 1/\sigma_j^2}, \ i, j = u, p, v \qquad (1)$$

where the weight of each cue is determined by their uncertainty $(\boldsymbol{\sigma})$ relative to each other.

Notably, in the VMR paradigm, the predicted reaching direction is the aiming direction.

The rotated cursor (x_v) biases the hand estimate $(\hat{x}_{hand})$ toward the cursor's direction.

The deviation from the predicted (re-aiming) direction constitutes the perceptual

prediction error, which drives implicit adaptation on the subsequent trial (Eq. 2). We

assume that learning proceeds on a trial-by-trial basis thus a state-space framework is

used. Explicit learning is driven by performance error as in existing models (Eq. 3) [8, 9,

10, 36, 140]. For each trial, the learning is updated by

1135
$$x_{t+1}^i = A^i x_t^i - B^i \left(\hat{x}_{Hand,t} - x_{u,t} \right) \tag{2}$$

1136
$$x_{t+1}^e = A^e x_t^e - B^e e_t^{PE} \tag{3}$$

1137
$$x_{t+1} = x_{t+1}^e + x_{t+1}^i \tag{4}$$

1138 Where total learning (x) is the sum of explicit learning (x^e) and implicit learning (x^i). *A* is

1139 the retention rate capturing the inter-trial forgetting and *B* is the learning rate capturing

1140 the proportion of error correction.

1141

1142 As individuals fixate near the re-aiming direction during the movement [37, 38], the visual

1143 cursor directionally differs from the fixation by an amount of $(x_v - x_u)$. This eccentricity

1144 linearly increases visual uncertainty about cursor motion direction, as shown both in visual

1145 perception studies and in our previous work [31]. Thus, the visual uncertainty (σ_v) follows

1146 a linear function:

1147
$$\sigma_v = k * |x_v - x_u| + b \tag{5}$$

1148 where *k* and *b* are the slope and intercept of the visual uncertainty, respectively.

1149
1150 *Performance error (PE) model:* The performance error (PE) model, also called the

1151 competition model [10] or the two-state model [11], has been widely used to explain motor

1152 adaptation [8-13]. The Eq. 2 in the PPE model is replaced by

1153
$$x_{t+1}^i = A^i x_t^i - B^i e_t^{PE} \tag{6}$$

1154

1155 *Sensory-prediction error (SPE) model:* Implicit motor learning has been suggested to be

1156 driven by sensory prediction error [15, 22, 29]. In the VMR paradigm, SPE is defined as

1157 the deviation of the perturbing cursor from the aiming direction [20]. Therefore, the SPE

1158 model replaces Eq.2 in the PPE model by

1159
$$x_{t+1}^i = A^i x_t^i - B^i \left(r - x_t^i \right), \tag{7}$$

1160   where $r - x_t^i$ is SPE and r is the VMR size.

1161

1162

1163

Data analysis

Hand kinematics: In Experiments 1 and 2, hand positions and velocities were directly

acquired from the KINARM robot at a fixed sampling rate of 1,000 Hz. The raw kinematic

data were smoothed using a fifth-order Savitzky-Golay filter with a window length of 50

ms. In Experiments 3, 4, and 5, hand positions were recorded at a sampling rate of 125

Hz. The movement direction was determined by computing the direction of the vector

spanning from the starting position to the hand position at the peak outward velocity. We

defined the direction to counter the perturbation as the positive direction.

1172

Gaze: The gaze position obtained in the control experiment was first preprocessed to

remove blinks. Consistent with previous studies [37], we used the last fixation before

movement onset to quantify the participants' aiming direction. Trials were excluded if the

fixation before movement onset was missing (due to a blink), or near the start position (<

40% of target distance), or beyond the target (>150% of target distance). These

exclusions amounted to 1.67% of the trials in the 45°-VMR condition and 3.61% in the

90°-VMR condition. Other direction measures, such as the gaze position at movement

onset, yielded similar results.

1181

Statistical analysis: For the dataset from *Bond & Taylor 2015,* we quantified early and late

learning by using the data from the first 5 cycles and the last 5 cycles in the adaptation

phase, respectively. Total learning was defined as the hand's directional deviation from

the target. Explicit learning was quantified as the direction deviation of the aiming report

from the target. Implicit learning was calculated by subtracting explicit learning from total

learning. Group differences were examined using one-way independent-measure

ANOVAs. In Experiments 1 and 2, implicit learning was quantified as the average hand

deviation from the target in the four exclusion trials. Total learning was calculated as the

average hand deviation from the target in the last cycle during the adaptation phase.

Explicit learning was calculated by subtracting implicit learning from total learning.

Proprioceptive bias was quantified as the average angular difference between the perceived (reported) and actual hand directions over the six proprioception test trials. Group differences were examined using one-way independent-measure ANOVAs, with pairwise post-hoc comparisons conducted using Bonferroni correction. For Experiment 2, two-way mixed ANOVAs were performed, with group as the between-group factor and learning time as the within-group factor. For the correlation analysis of Experiment 2, we used a mixed linear model: , where y is implicit learning, is proprioceptive bias, is the random factor of subject, and is the residual error. In Experiment 3, single-trial learning was defined as the angular difference in movement direction between two consecutive no-feedback null trials before and after a perturbation trial. Given the symmetry between CW and CCW perturbations, we pool the data to quantify the size dependency on the absolute perturbation size. Repeated-measures ANOVAs were used to examine the effect of perturbation size on single-trial learning. In Experiment 4, independent t-tests were used to compare implicit learning between 45°-VMR and 90°-VMR conditions. In Experiment 5, we used paired t-tests to compare implicit learning before and after introducing the re-aiming target. Implicit learning was quantified by both subtraction and exclusion methods. In Experiment 6, we used one-sample t-tests to examined whether PPE was significantly negative, and used one-way ANOVAs to examine group differences in PPE and implicit learning. Post-hoc comparisons were conducted with Bonferroni correction. In the eye-tracking experiment, we used paired t tests condition differences (90° vs. 45°) in re-aiming direction. We used a mixed linear model for analyzing the correlation between explicit learning and eye-fixation direction: , where is explicit learning, is eye-fixation direction, is the random factor of subject, and is the residual error. For all experiments, we excluded the trials with movement time exceeding 400 ms or below 150 ms, as well as the trials with systematic movements to the incorrect direction (i.e., movement direction less than -20° or exceeded 120°). These criteria resulted in the removal of less than 1% of trials on average for each experiment.

Model analysis

1221   Both PE and SPE models have four parameters $\Theta = [A^e, B^e, A^i, B^i]$. The PPE model has

1222   an additional four parameters $\Theta = [\sigma_u, \sigma_p, k, b]$. Not all the experiments need the full

1223   models (as detailed below).

1224

Bond & Taylor, 2015. We fitted the trial-by-trial average learning data from all the four

perturbation conditions simultaneously (Figure 2C). No fitting was needed for explicit

learning since it was measured throughout. The parameters related to sensory

uncertainty (σ_u, σ_p, k, b) had been estimated from our previous work with visuomotor

rotation perturbations [31]. Three of them were fixed, including b = 1.853°, σ_u = 5.05°, σ_p

= 11.12°. Note, the parameter k, the linear slope of visual uncertainty relative to

perturbation size, was left as a free parameter. This is because the previous work used

the error-clamp paradigm, which reinforced fixation on the reach target rather than on the

cursor. In contrast, participants in the VMR paradigm here, look at the cursor at the end

of the reaching movement, which presumably reduces the effect of eccentricity on visual

uncertainty. Therefore, we left k as a free parameter for all the experiments examined

here. Hence, the PPE model had three free parameters $\Theta = [k, A^i, B^i]$, while the PE and

SPE models each had two free parameters $\Theta = [A^i, B^i]$. Bootstrapping was used to

estimate the standard deviations of the model fits. Participants in each group were

resampled with replacement 5000 times, each time producing a new average learning

sequence.

1241

Experiment 1 and 2. The trial-by-trial learning data include total learning and implicit

learning (intermittently measured during the adaptation phase) from all the four conditions

(Experiment 1; Figure 3C) or the two conditions (Experiment 2; Figure 4D). The PPE

model had five free parameters $\Theta = [k, A^i, B^i, A^e, B^e]$. Since the one-step condition in

Experiment 2 was identical to the conditions in Experiment 1 (only differing in the duration

of the adaptation phase), we applied the k value estimated from Experiment 1 to the one-

step condition in Experiment 2. The stepwise condition, however, retained k as a free

parameter (k_{step} in Table S3) since the participants would be attracted by the sudden

stepwise change in cursor rotation, which would reduce visual uncertainty. Through fitting

the data from both two conditions (Figure 4D), we obtained the value of k_{step}. We used

the values of k and k_{step} for cross-validation (Figure S2B). The SPE and PE models had

four free parameters $\Theta = [A^e, B^e, A^i, B^i]$.

1254

Experiment 3, 4 and 5. We first assumed that the single-trial learning in Experiment 3 was

implicit. Thus, the PPE model had three free parameters $\Theta = [k, A^i, B^i]$ to fit the average

trial-by-trial changes (Figure 5B). The averaging was feasible since the pseudo-random

perturbation sequence was identical for each participant. For fitting, we set the lower limit

of at 0.7 for all the models. Based on fitting the trial-by-trial learning, we calculated the

model-predicted single-trial learning as a function of perturbation size (Figure 5D). To

examine whether there was explicit learning at work, we then included the explicit learning

component and evaluated whether it improved model fitting. In Experiment 4, there was

only implicit learning since participants were unaware of their hand drifting away from the

instructed re-aiming target. Thus, only implicit learning was fitted for the 45°- and 90°-

VMR conditions at once. The PPE model had three free parameters $\Theta = [k, A^i, B^i]$, while

the SPE and PE models had two free parameters $\Theta = [A^i, B^i]$. For Experiment 5, the data

used for model fitting included the trial-by-trial changes in total learning as well as the

intermittently-measured explicit learning. Models were fitted to the data before introducing

the re-aiming target and then used to predict the learning afterwards (Figure 7D). The

PPE model thus had five free parameters $\Theta = [k, A^i, B^i, A^e, B^e]$, while the other two models

had four free parameters $\Theta = [A^i, B^i, A^e, B^e]$.

1272

All model fittings were conducted using *fmincon* function in MATLAB (2022b, MathWorks,

Natick, MA, US). To ensure the model adhered to established assumptions, we

constrained the parameters with $A^e < A^i$ and $B^e > B^i$ as commonly assumed for explicit

and implicit learning components [8, 9, 11]. All the obtained parameter values, along with

quality measures of the model fit, including R^2, root mean squared error (*RMSE*), Akaike

information criterion (AIC), and Bayesian information criterion (BIC), were provided in

Table S1 to S6.

1280

1281   **Data availability**

1282   Data and code are available at: https://osf.io/tjc52/

1283

Acknowledgements: We would like to thank Dr. Jordan Taylor for providing the raw data

from his study [34]. **Funding**: This work was supported by STI2030-Major Projects

(2021ZD0202600) and the National Natural Science Foundation of China (32071047,

32471099, 6206113600, 31871102), awarded to KW. **Authors contributions:**

Conceptualization: KW, XZ, Data curation: XZ, WW; Formal analysis: XZ; Funding

acquisition: KW; Methodology: KW, XZ, WW; Software: XZ, WW; Supervision: KW;

Visualization: XZ; Writing: KW, XZ.

1291

Supplementary materials

 Figure S1. Control experiment examining eye fixation during visuomotor rotation
 adaptation. **(A)** Illustration of the experimental procedure. In each trial, participants first
 held their hand at the start position for 500 ms, awaiting the appearance of a target
 randomly positioned between 0° and 360°. They were instructed not to move until the

target disappeared (~1000 ms after appearance). Visual feedback of the hand cursor was
withheld until the trial ended, when it re-appeared together with the target as performance
feedback. Eye tracking measurements were taken during reaching. **(B)** Time course of
the gaze direction. Different colors represent different timings of a trial, including target
preview, reaction time (RT), movement duration, and terminal feedback. Two
representative trials from a participant are shown for the 45°-VMR (upper panel) and 90°-
VMR (lower panel) conditions, respectively. The participant fixated on the starting position
during the target preview and RT, then shifted fixation toward the re-aiming direction
during the movement, maintaining this fixation until the terminal feedback was provided
(purple line). The re-aiming direction was less rotated than the imposed VMR. After the
movement ended and visual feedback was provided, participants tended to fixate back to
the cursor (yellow line). We calculated the probability of fixations in the target area (± 22.5°
around the target direction) after movement: 64.28 ± 4.91% in the 45°-VMR condition and
43.23 ± 6.80% in the 90°-VMR condition. **(C)** Average hand movement and eye fixation
directions during adaptation to the 45° and 90° VMR. The fixation direction was defined
as the last fixation before movement onset [37]. The shaded areas denote the standard
errors across participants. From early learning, participants started to fixated away from
the target direction, closer to their explicit re-aiming direction, and this fixation "strategy"
persisted throughout the adaptation phase. Note that the fixation deviated further from
the target in the 90°-VMR condition compared to the 45°-VMR condition, consistent with
the condition difference in re-aiming and explicit learning. In the washout phase, their
fixation returned to the target as instructed. **(D)** Participants' fixation direction in the 90°-
VMR condition was significantly higher than that in the 45°-VMR condition, during both
early (45°: 19.00°± 2.23°; 90°: 44.49°± 4.43°; $t_{(15)}$ = -6.38, p < .0001) and late learning
(45°: 21.63°± 2.56°; 90°: 57.88°± 4.18°; $t_{(15)}$ = -9.21, p < .0001). The larger the fixation
direction, the greater the re-aiming deviates away from the target. Early and late learning
refer to the first 5 and last 5 trial cycles during adaptation. **(E)** Participants' eye-fixation
direction was positively correlated with their explicit learning ($β$ = 0.80, p = 0.014), the
latter estimated from the first exclusion trial in the washout phase.
1328
1329

1330

Figure S2. Model comparison results from Experiment 2. **(A)** Implicit learning data in
Experiment 2, a reproduction of Figure 4C right panel. **(B)** Model fitting with varying
explicit parameters across the two conditions. Compared to the model fitting in the main
text (Figure 4D), here we allowed explicit parameters to vary across the two conditions.
The PE and SPE models still failed to reproduce the data, especially the relative size of
implicit learning between the two perturbation conditions. The PPE model remained
capable of explaining the data well with variable explicit learning parameters. These
results suggest that the failure of PE and SPE to explain implicit learning across
conditions was not due to an unconsidered potential parametric change in explicit learning.
(C) Cross-validation tests for the three models. Each model was fit to the data from the
stepwise condition and then used to predict implicit learning in the one-step condition.
Neither PE or SPE model captures the learning pattern, with mismatches similar to their
original model fitting in Figure 4D. In contrast, the PPE model's predictions match well
with the empirical data across conditions. In **(B)** and **(C)**, error bars represent
bootstrapped standard deviations (resample size = 5000).

1346

1347

Figure S3. Correlation between explicit and implicit learning varies widely when the two learning components are independently measured. Data from three separate studies are presented, with explicit learning measured by aiming reports and implicit learning

measured by no-feedback exclusion trials. **(A)** Correlations between explicit and implicit learning in different perturbation size conditions, including 15°, 30°, 60°, and 90° VMR (n = 10 for each). Due to the small sample size and variability across condition, support for a negative correlation was insufficient. Data from *Bond & Taylor, 2015*. **(B)** Correlations across perturbation size conditions of 15°, 30°, 45°, and 60° VMR (n = 33, 54, 28, 61, respectively) from *D'Amario et al., 2024*. With decent sample sizes, positive, negative or null correlation was observed in these different conditions. **(C)** Correlations were inconsistent when the format of visual perturbation varied but with a fixed perturbation size of 45° VMR. Data from *D'Amario et al., 2024*. Continuous cursor feedback (n = 51) is the conventional feedback format, terminal feedback promotes explicit learning and reduce implicit learning (n = 35), delayed feedback reduces implicit learning (n = 39), and cursor jump feedback usually promotes explicit learning and reduces implicit learning (n = 32). Only the continuous-feedback condition showed a negative correlation, no correlation was detected for the other three feedback formats. **(D)** Early learning did not show a negative correlation. In our Experiment 5 (n = 20), explicit learning and implicit learning were independently measured four times during the 60 trial-cycle learning period. The two early instances (the 5th and 19th cycles) did not show significant negative correlations, the third instance (the 33th cycle) showed a marginal effect, and the fourth instance (the 47th cycle) showed a significant negative correlation. Thus, though performance error is most prominent during early learning, there is no evidence that the two learning processes compete for this error and then lead to a negative correlation. Each dot represents an individual's data.

Figure S4. The performance of the PPE model was compared by assuming a linear and an exponential visual uncertainty function. The observed and model-predicted implicit learning for Bond & Taylor 2015 (A), Experiment 1 (B), and Experiment 2 (C). The two versions of the PPE model similarly explain the data. (D) The visual uncertainty functions are estimated from the three datasets. Note the estimates are similar for the linear and the exponential functions, and they are also close to the visual uncertainty measured in our previous work [31].

1401 **Table S1**. Model comparisons and parameters for the dataset from *Bond & Taylor, 2015*

	R^2	$RMSE(°)$	AIC	BIC	B_i	A_i	k
PPE Model	**0.708**	**1.684**	**172.81**	**182.04**	0.158	0.931	0.209
PE Model	0.177	2.188	254.54	260.69	0.072	0.999	/
SPE Model	-0.730	5.054	522.48	528.63	0.015	0.916	/

1402

1403 **Table S2**. Model comparisons and parameters for the data from Experiment 1

	R^2	$RMSE(°)$	AIC	BIC	B_i	A_i	B_e	A_e	k
PPE Model	**0.969**	**4.807**	**361.67**	**375.27**	0.151	0.981	0.513	0.981	0.204
PE Model	0.957	5.870	404.43	415.31	0.069	0.991	0.450	0.991	/
SPE Model	0.954	6.044	411.00	421.87	0.014	0.979	0.505	0.979	/

1404

1405 **Table S3**. Model comparisons and parameters for the data from Experiment 2

1406 **Table S3a.** Fitting the one-step and stepwise groups together with a single set of
1407 parameters

	R^2	$RMSE(°)$	AIC	BIC	B_i	A_i	B_e	A_e	k	k_{step}
PPE Model	**0.960**	**3.965**	**252.43**	**264.81**	0.245	0.927	0.889	0.927	0.204	0.139
PE Model	0.933	5.245	299.68	309.59	0.263	0.958	0.647	0.958	/	/
SPE Model	0.925	5.491	307.76	317.67	0.046	0.906	0.908	0.906	/	/

1408

1409 **Table S3b.** Assuming explicit learning changes across conditions, fitting the one-step and
1410 stepwise groups with separate explicit learning parameters

							Stepwise			One-step		
	R^2	$RMSE(°)$	AIC	BIC	B_i	A_i	B_e	A_e	k_{step}	B_e	A_e	k
PPE Model	**0.962**	**3.854**	**251.42**	**268.76**	0.237	0.930	0.636	0.924	0.136	0.940	0.930	0.204
PE Model	0.958	4.234	265.97	280.84	0.223	0.970	0.218	0.990	/	0.792	0.970	/
SPE Model	0.926	5.453	310.53	325.39	0.047	0.902	0.672	0.945	/	0.939	0.902	/

1411

1412 **Table S3c.** Cross-validation: first fit the model by using the data from the stepwise group
1413 only, and then predict the data for the one-step group

	R^2	$RMSE(°)$	AIC	BIC	B_i	A_i	B_e	A_e	k	k_{step}
PPE Model	**0.913**	**5.868**	**165.72**	**174.64**	0.304	0.911	0.660	0.911	0.204	0.139
PE Model	0.661	9.031	201.66	208.80	0.382	0.946	0.454	0.946	/	/
SPE Model	0.583	9.638	207.38	214.52	0.132	0.859	0.774	0.859	/	/

1414

Table S4. Model comparisons and parameters for the data from Experiment 3

Table S4a. For the trial-by-trial learning sequence (Figure 5B)

		R^2	RMSE(°)	AIC	BIC	B_i	A_i	B_e	A_e	k
PPE Model	**Imp**	**0.471**	**1.294**	**39.988**	**46.557**	0.418	0.700	/	/	0.193
	Imp+Exp	0.453	1.317	46.307	57.256	0.424	0.700	0.010	0.500	0.201
PE Model	Imp	-0.682	1.747	77.622	82.001	0.054	0.700	/	/	/
	Imp+Exp	-0.682	1.747	81.622	90.381	0.054	0.700	0.496	0.542	/
SPE Model	Imp	-0.682	1.747	77.622	82.001	0.054	0.700	/	/	/
	Imp+Exp	-0.698	1.750	81.848	90.607	0.054	0.700	0.010	0.500	/

Table S4b. For the single-trial learning function (Figure 5C)

	R^2	RMSE (°)
PPE Model	**0.954**	**0.684**
PE/SPE Model	-0.321	2.449

Table S5. Model comparisons and parameters for the data from Experiment 4

	R^2	RMSE (°)	AIC	BIC	B_i	A_i	k
PPE Model	**0.972**	**0.524**	**-32.81**	**-28.61**	0.396	0.840	0.211
PE Model	-0.866	4.496	94.19	97.00	0.034	0.796	/
SPE Model	-1.150	4.457	93.67	96.48	0.025	0.833	/

Table S6. Model comparisons and parameters for the data from Experiment 5: Cross-validation (Fitting before-re-aiming phase, predicting after-re-aiming phase).

	R^2	RMSE (°)	AIC	BIC	B_i	A_i	B_e	A_e	k
PPE Model	**0.880**	**5.641**	**155.33**	**164.01**	0.082	0.966	0.304	0.966	0.201
PE Model	0.337	13.825	228.62	235.57	0.082	0.966	0.304	0.966	/
SPE Model	0.859	5.843	156.59	163.54	0.016	0.954	0.359	0.954	/

1432 **Table S7:** Model comparisons of the PPE models, which assume linear, exponential, or

1433 power-law visual uncertainty function.

1434 PPE^{exp} model: assuming $\sigma_v = k'e^{\alpha\theta} + (b - k')$

1435 PPE^{power} model: assuming $\sigma_v = k'\theta^{\alpha} + b$

1436

1437 **Table S7a.** Model comparisons for *Bond & Taylor, 2015. Note that the parameters of*

1438 *power and exponential functions indicate that these functions are close to linear ones.*

	R^2	$RMSE(°)$	AIC	BIC	B_i	A_i		
PPE Model	0.708	1.684	172.81	182.04	0.158	0.931	/	/
PPE^{power} Model	0.978	1.173	59.05	71.35	0.531	0.977	1.49	0.674
PPE^{exp} Model	0.680	1.777	191.93	204.24	0.143	0.930	99.93	0.002
PE Model	0.177	2.188	254.54	260.69	0.072	0.999	/	/
SPE Model	-0.730	5.054	522.48	528.63	0.015	0.916	/	/

1439

1440 **Table S7b.** Model comparisons for Experiment 1. *Note the parameters of power and*

1441 *exponential functions indicate these functions are close to a linear one.*

	R^2	$RMSE(°)$	AIC	BIC	B_i	A_i	B_e	A_e		
PPE Model	0.969	4.807	359.67	370.55	0.151	0.981	0.513	0.981	/	/
PPE^{power} Model	0.971	4.612	350.44	361.31	0.505	0.982	0.505	0.982	1.28	0.681
PPE^{exp} Model	0.968	4.832	360.84	371.71	0.014	0.979	0.505	0.979	99.99	0.002
PE Model	0.957	5.870	404.43	415.31	0.069	0.991	0.450	0.991	/	/
SPE Model	0.954	6.044	411.00	421.87	0.014	0.979	0.505	0.979	/	/

1442

1443 **Table S7c.** Model comparisons for Experiment 2. *Note the parameters of power and*

1444 *exponential functions indicate these functions are close to a linear one.*

	R^2	$RMSE(°)$	AIC	BIC	B_i	A_i	B_e	A_e		
PPE Model	0.960	3.965	252.43	264.81	0.245	0.927	0.889	0.927	/	/
PPE^{power} Model	0.962	3.897	253.39	270.73	0.409	0.941	0.870	0.941	0.43	0.889
PPE^{exp} Model	0.962	3.900	253.55	270.89	0.360	0.940	0.874	0.940	99.73	0.003
PE Model	0.933	5.245	299.68	309.59	0.263	0.958	0.647	0.958	/	/
SPE Model	0.925	5.491	307.76	317.67	0.046	0.906	0.908	0.906	/	/

1445

1446

1447

1448

Reference

1. Clark A. Whatever next? Predictive brains, situated agents, and the future of cognitive science. Behavioral and brain sciences. 2013;36(3):181-204.
2. Wolpert DM, Diedrichsen J, Flanagan JR. Principles of sensorimotor learning. Nature Reviews Neuroscience. 2011;12(12):739.
3. Dosher B, Lu Z-L. Visual perceptual learning and models. Annual review of vision science. 2017;3(1):343-63.
4. Ellis R. Implicit and explicit learning, knowledge and instruction. Implicit and explicit knowledge in second language learning, testing and teaching. 2009;42:3-25.
5. Green TD, Flowers JH. Implicit versus explicit learning processes in a probabilistic, continuous fine-motor catching task. Journal of motor behavior. 1991;23(4):293-300.
6. Batterink LJ, Reber PJ, Neville HJ, Paller KA. Implicit and explicit contributions to statistical learning. Journal of memory and language. 2015;83:62-78.
7. Jordan MI, Rumelhart DE. Forward models: Supervised learning with a distal teacher. Cognitive science. 1992;16(3):307-54.
8. McDougle SD, Bond KM, Taylor JA. Explicit and implicit processes constitute the fast and slow processes of sensorimotor learning. Journal of Neuroscience. 2015;35(26):9568-79.
9. McDougle SD, Bond KM, Taylor JA. Implications of plan-based generalization in sensorimotor adaptation. Journal of neurophysiology. 2017;118(1):383-93.
10. Albert ST, Jang J, Modchalingam S, Marius't Hart B, Henriques D, Lerner G, et al. Competition between parallel sensorimotor learning systems. elife. 2022;11:e65361.
11. Smith MA, Ghazizadeh A, Shadmehr R. Interacting adaptive processes with different timescales underlie short-term motor learning. PLoS Biol. 2006;4(6):e179.
12. Miyamoto YR, Wang S, Smith MA. Implicit adaptation compensates for erratic explicit strategy in human motor learning. Nature neuroscience. 2020;23(3):443-55.
13. Hadjiosif AM, Krakauer JW, Haith AM. Did we get sensorimotor adaptation wrong? Implicit adaptation as direct policy updating rather than forward-model-based learning. Journal of Neuroscience. 2021;41(12):2747-61.
14. Shadmehr R, Smith MA, Krakauer JW. Error correction, sensory prediction, and adaptation in motor control. Annual review of neuroscience. 2010;33:89-108.
15. Mazzoni P, Krakauer JW. An implicit plan overrides an explicit strategy during visuomotor adaptation. J Neurosci. 2006;26(14):3642-5.
16. Morehead JR, Taylor JA, Parvin DE, Ivry RB. Characteristics of implicit sensorimotor adaptation revealed by task-irrelevant clamped feedback. Journal of cognitive neuroscience. 2017;29(6):1061-74.
17. Wolpert D, Kawato M. Multiple paired forward and inverse models for motor control. Neural Networks. 1998;11(7-8):1317-29.
18. Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM. Motor learning. Compr Physiol. 2019;9(2):613-63.
19. Harris CM, Wolpert DM. Signal-dependent noise determines motor planning. Nature. 1998;394(6695):780-4.
20. Morehead JR, Orban de Xivry JJ. A synthesis of the many errors and learning processes of visuomotor adaptation. BioRxiv. 2021 Mar 15:2021-03.

21. Schmidt RA, Lee TD, Winstein C, Wulf G, Zelaznik HN. Motor control and learning:
A behavioral emphasis: Human kinetics; 2018.
22. Tseng Y, Diedrichsen J, Krakauer J, Shadmehr R, Bastian A. Sensory prediction
errors drive cerebellum-dependent adaptation of reaching. Journal of
Neurophysiology. 2007;98(1):54.
23. Tsay JS, Haith AM, Ivry RB, Kim HE. Interactions between sensory prediction error
and task error during implicit motor learning. PLoS computational biology.
2022;18(3):e1010005.
24. Schween R, Taube W, Gollhofer A, Leukel C. Online and post-trial feedback
differentially affect implicit adaptation to a visuomotor rotation. Experimental brain
research. 2014;232:3007-13.
25. Lee K, Oh Y, Izawa J, Schweighofer N. Sensory prediction errors, not performance
errors, update memories in visuomotor adaptation. Scientific reports.
2018;8(1):16483.
26. Taylor JA, Ivry RB. Flexible cognitive strategies during motor learning. PLoS
computational biology. 2011;7(3):e1001096.
27. Kayacan E, Chowdhary G. Tracking error learning control for precise mobile robot
path tracking in outdoor environment. Journal of Intelligent & Robotic Systems.
2019;95:975-86.
28. Sober SJ, Sabes PN. Flexible strategies for sensory integration during motor
planning. Nature neuroscience. 2005;8(4):490-7.
29. Kawato M. Internal models for motor control and trajectory planning. Current Opinion
in Neurobiology. 1999;9(6):718-27.
30. Hogan N, Flash T. Moving gracefully: quantitative theories of motor coordination.
Trends in Neurosciences. 1987;10(4):170-4.
31. Zhang Z, Wang H, Zhang T, Nie Z, Wei K. Perceptual error based on Bayesian cue
combination drives implicit motor adaptation. eLife. 2024;13:RP94608.
32. Tsay JS, Kim H, Haith AM, Ivry RB. Understanding implicit sensorimotor adaptation
as a process of proprioceptive re-alignment. Elife. 2022;11:e76639.
33. Taylor JA, Krakauer JW, Ivry RB. Explicit and implicit contributions to learning in a
sensorimotor adaptation task. Journal of Neuroscience. 2014;34(8):3023-32.
34. Bond KM, Taylor JA. Flexible explicit but rigid implicit learning in a visuomotor
adaptation task. Journal of neurophysiology. 2015;113(10):3836-49.
35. Cheng S, Sabes PN. Modeling sensorimotor learning with linear dynamical systems.
Neural Comput. 2006;18(4):760-93.
36. Herzfeld DJ, Vaswani PA, Marko MK, Shadmehr R. A memory of errors in
sensorimotor learning. Science. 2014;345(6202):1349-53.
37. Bromberg Z, Donchin O, Haar S. Eye movements during visuomotor adaptation
represent only part of the explicit learning. Eneuro. 2019;6(6).
38. de Brouwer AJ, Albaghdadi M, Flanagan JR, Gallivan JP. Using gaze behavior to
parcellate the explicit and implicit contributions to visuomotor learning. Journal of
neurophysiology. 2018;120(4):1602-15.
39. Ernst M, Banks M. Humans integrate visual and haptic information in a statistically
optimal fashion. Nature 2002;415(6870):429-33.

40. Kim HE, Morehead JR, Parvin DE, Moazzezi R, Ivry RB. Invariant errors reveal limitations in motor correction rather than constraints on error sensitivity. Communications biology. 2018;1(1):19.

41. Neville K-M, Cressman EK. The influence of awareness on explicit and implicit contributions to visuomotor adaptation over time. Experimental brain research. 2018;236:2047-59.

42. Maresch J, Mudrik L, Donchin O. Measures of explicit and implicit in motor learning: what we know and what we don't. Neuroscience & Biobehavioral Reviews. 2021;128:558-68.

43. Maresch J, Werner S, Donchin O. Methods matter: Your measures of explicit and implicit processes in visuomotor adaptation affect your results. European Journal of Neuroscience. 2021;53(2):504-18.

44. Werner S, Van Aken BC, Hulst T, Frens MA, Van Der Geest JN, Strüder HK, et al. Awareness of sensorimotor adaptation to visual rotations of different size. PLoS One. 2015;10(4):e0123321.

45. Ruttle JE, 't Hart BM, Henriques DY. Implicit motor learning within three trials. Scientific Reports. 2021;11(1):1627.

46. Salomonczyk D, Cressman EK, Henriques DY. Proprioceptive recalibration following prolonged training and increasing distortions in visuomotor adaptation. Neuropsychologia. 2011;49(11):3053-62.

47. Henriques DY, Cressman EK. Visuomotor adaptation and proprioceptive recalibration. Journal of motor behavior. 2012;44(6):435-44.

48. Salomonczyk D, Cressman EK, Henriques DY. The role of the cross-sensory error signal in visuomotor adaptation. Experimental Brain Research. 2013;228:313-25.

49. Ruttle JE, 't Hart BM, Henriques DY. The fast contribution of visual-proprioceptive discrepancy to reach aftereffects and proprioceptive recalibration. PLoS One. 2018;13(7):e0200621.

50. Rossi C, Bastian AJ, Therrien AS. Mechanisms of proprioceptive realignment in human motor learning. Current Opinion in Physiology. 2021;20:186-97.

51. Tsay JS, Lee A, Ivry RB, Avraham G. Moving outside the lab: The viability of conducting sensorimotor learning studies online. arXiv preprint arXiv:210713408. 2021.

52. Modchalingam S, Vachon CM, 't Hart BM, Henriques DY. The effects of awareness of the perturbation during motor adaptation on hand localization. PLoS One. 2019;14(8):e0220884.

53. Modchalingam S, Ciccone M, D'Amario S, 't Hart BM, Henriques DY. Adapting to visuomotor rotations in stepped increments increases implicit motor learning. Scientific Reports. 2023;13(1):5022.

54. Coltman SK, Cashaback JG, Gribble PL. Both fast and slow learning processes contribute to savings following sensorimotor adaptation. Journal of neurophysiology. 2019;121(4):1575-83.

55. Wei K, Kording K. Relevance of error: what drives motor adaptation? Journal of Neurophysiology. 2009;101(2):655-64.

56. Lu Z-L, Dosher BA. Current directions in visual perceptual learning. Nature reviews psychology. 2022;1(11):654-68.

57. Holroyd CB, Coles MG. The neural basis of human error processing: reinforcement
learning, dopamine, and the error-related negativity. Psychological review.
2002;109(4):679.
58. Niv Y. Reinforcement learning in the brain. Journal of Mathematical Psychology.
2009;53(3):139-54.
59. Johnson DD, Blumstein DT, Fowler JH, Haselton MG. The evolution of error: Error
management, cognitive constraints, and adaptive decision-making biases. Trends in
ecology & evolution. 2013;28(8):474-81.
60. Law C-T, Gold JI. Reinforcement learning can account for associative and perceptual
learning on a visual-decision task. Nature neuroscience. 2009;12(5):655-63.
61. Leow LA, Marinovic W, de Rugy A, Carroll TJ. Task errors contribute to implicit
aftereffects in sensorimotor adaptation. European Journal of Neuroscience.
2018;48(11):3397-409.
62. Leow L-A, Marinovic W, de Rugy A, Carroll TJ. Task errors drive memories that
improve sensorimotor adaptation. Journal of neuroscience. 2020;40(15):3075-88.
63. Al-Fawakhiri N, Ma A, Taylor JA, Kim OA. Exploring the role of task success in implicit
motor adaptation. Journal of neurophysiology. 2023;130(2):332-44.
64. Berniker M, Körding K. Estimating the sources of motor errors for adaptation and
generalization. Nature Neuroscience. 2008;11(12):1454-61.
65. Oza A, Kumar A, Sharma A, Mutha PK. Limb-related sensory prediction errors and
task-related performance errors facilitate human sensorimotor learning through
separate mechanisms. PLoS Biology. 2024;22(7):e3002703.
66. Pelisson D, Alahyane N, Panouilleres M, Tilikete C. Sensorimotor adaptation of
saccadic eye movements. Neuroscience & Biobehavioral Reviews. 2010;34(8):1103-
20.
67. Wong AL, Shelhamer M. Saccade adaptation improves in response to a gradually
introduced stimulus perturbation. Neuroscience letters. 2011;500(3):207-11.
68. Gonzalez Castro LN, Monsen CB, Smith MA. The binding of learning to action in
motor adaptation. PLoS Computational Biology. 2011;7(6):e1002052.
69. Shadmehr R, Mussa-Ivaldi FA. Adaptive representation of dynamics during learning
of a motor task. The Journal of Neuroscience. 1994;14(5):3208-24.
70. Thoroughman KA, Shadmehr R. Learning of action through adaptive combination of
motor primitives. Nature. 2000;407(6805):742-7.
71. Brown LE, Rosenbaum DA, Sainburg RL. Limb position drift: implications for control
of posture and movement. Journal of neurophysiology. 2003;90(5):3105-18.
72. Duff SV, Sainburg RL. Lateralization of motor adaptation reveals independence in
control of trajectory and steady-state position. Experimental brain research.
2007;179:551-61.
73. DiZio P, Lackner JR. Congenitally blind individuals rapidly adapt to Coriolis force
perturbations of their reaching movements. Journal of neurophysiology.
2000;84(4):2175-80.
74. Franklin DW, So U, Burdet E, Kawato M. Visual feedback is not necessary for the
learning of novel dynamics. PloS one. 2007;2(12):e1336.
75. Marko MK, Haith AM, Harran MD, Shadmehr R. Sensitivity to prediction error in reach
adaptation. Journal of neurophysiology. 2012;108(6):1752-63.

76. Striemer CL, Enns JT, Whitwell RL. Visuomotor adaptation in the absence of input
from early visual cortex. Cortex. 2019;115:201-15.
77. Deubel H. Adaptivity of gain and direction in oblique saccades. Eye movements from
physiology to cognition: Elsevier; 1987. p. 181-90.
78. McLaughlin SC. Parametric adjustment in saccadic eye movements. Perception &
Psychophysics. 1967;2(8):359-62.
79. Collins T, Wallman J. The relative importance of retinal error and prediction in
saccadic adaptation. Journal of Neurophysiology. 2012;107(12):3342-8.
80. Bahcall DO, Kowler E. Illusory shifts in visual direction accompany adaptation of
saccadic eye movements. Nature. 1999;400(6747):864-6.
81. Masselink J, Lappe M. Visuomotor learning from postdictive motor error. Elife.
2021;10:e64278.
82. Deubel H, Schneider WX, Bridgeman B. Postsaccadic target blanking prevents
saccadic suppression of image displacement. Vision research. 1996;36(7):985-96.
83. Wurtz RH. Neuronal mechanisms of visual stability. Vision research.
2008;48(20):2070-89.
84. Burr DC, Morrone MC. Constructing stable spatial maps of the word. Perception.
2012;41(11):1355-72.
85. Nasir SM, Ostry DJ. Auditory plasticity and speech motor learning. Proceedings of
the National Academy of Sciences. 2009;106(48):20470-5.
86. Houde JF, Jordan MI. Sensorimotor adaptation in speech production. Science.
1998;279(5354):1213-6.
87. Daliri A, Dittman J. Successful auditory motor adaptation requires task-relevant
auditory errors. Journal of Neurophysiology. 2019;122(2):552-62.
88. Parrell B, Agnew Z, Nagarajan S, Houde J, Ivry RB. Impaired feedforward control
and enhanced feedback control of speech in patients with cerebellar degeneration.
Journal of Neuroscience. 2017;37(38):9249-58.
89. Parrell B, Ivry RB, Nagarajan SS, Houde JF. Intact correction for self-produced vowel
formant variability in individuals with cerebellar ataxia regardless of auditory
feedback availability. Journal of Speech, Language, and Hearing Research.
2021;64(6S):2234-47.
90. Kim KS, Max L. Speech auditory-motor adaptation to formant-shifted feedback lacks
an explicit component: Reduced adaptation in adults who stutter reflects limitations
in implicit sensorimotor learning. European Journal of Neuroscience.
2021;53(9):3093-108.
91. Kim KS, Hinkley LB, Dale CL, Nagarajan SS, Houde JF. Neurophysiological
evidence of sensory prediction errors driving speech sensorimotor adaptation.
BioRxiv. 2023.
92. Kim KS, Gaines JL, Parrell B, Ramanarayanan V, Nagarajan SS, Houde JF.
Mechanisms of sensorimotor adaptation in a hierarchical state feedback control
model of speech. PLoS Computational Biology. 2023;19(7):e1011244.
93. Parrell B, Ramanarayanan V, Nagarajan S, Houde J. The FACTS model of speech
motor control: Fusing state estimation and task-based control. PLoS computational
biology. 2019;15(9):e1007321.

94. Patri J-F, Perrier P, Schwartz J-L, Diard J. What drives the perceptual change resulting from speech motor adaptation? Evaluation of hypotheses in a Bayesian modeling framework. PLoS computational biology. 2018;14(1):e1005942.

95. Botvinick M, Cohen J. Rubber hands 'feel' touch that eyes see. Nature. 1998;391(6669):756-.

96. Sanchez-Vives MV, Spanlang B, Frisoli A, Bergamasco M, Slater M. Virtual hand illusion induced by visuomotor correlations. PloS one. 2010;5(4):e10381.

97. Asai T. Illusory body-ownership entails automatic compensative movement: for the unified representation between body and action. Experimental brain research. 2015;233:777-85.

98. Lanillos P, Franklin S, Maselli A, Franklin DW. Active strategies for multisensory conflict suppression in the virtual hand illusion. Scientific Reports. 2021;11(1):22844.

99. Maselli A, Lanillos P, Pezzulo G. Active inference unifies intentional and conflict-resolution imperatives of motor control. PLoS computational biology. 2022;18(6):e1010095.

100. Rengarajan S, Cannon J, Baron B, Mohan N, Chukoskie L, editors. Measuring the Interaction of Conflict-Minimizing and Goal-Seeking Motor Imperatives in Autism Spectrum Disorder. International Simulation and Gaming Association Conference; 2022: Springer.

101. Benson BL, Anguera JA, Seidler RD. A spatial explicit strategy reduces error but interferes with sensorimotor adaptation. Journal of neurophysiology. 2011;105(6):2843-51.

102. Flegal KE, Anderson MC. Overthinking skilled motor performance: Or why those who teach can't do. Psychonomic Bulletin & Review. 2008;15:927-32.

103. Wilterson SA, Taylor JA. Implicit visuomotor adaptation remains limited after several days of training. eneuro. 2021;8(4).

104. Gastrock RQ, 't Hart BM, Henriques DY. Distinct learning, retention, and generalization in de novo learning. bioRxiv. 2023:2023.10. 02.560506.

105. Telgen S, Parvin D, Diedrichsen J. Mirror reversal and visual rotation are learned and consolidated via separate mechanisms: recalibrating or learning de novo? Journal of Neuroscience. 2014;34(41):13768-79.

106. Kasuga S, Kurata M, Liu M, Ushiba J. Alteration of a motor learning rule under mirror-reversal transformation does not depend on the amplitude of visual error. Neuroscience research. 2015;94:62-9.

107. Kahneman D. Thinking, fast and slow. Farrar, Straus and Giroux. 2011.

108. D'Amario S, Ruttle JE, 't Hart BM, Henriques DY. Implicit Adaptation is Fast, Robust and Independent from Explicit Adaptation. bioRxiv. 2024:2024.04. 10.588930.

109. Mostafa AA, 't Hart BM, Henriques DY. Motor learning without moving: Proprioceptive and predictive hand localization after passive visuoproprioceptive discrepancy training. PLoS One. 2019;14(8):e0221861.

110. 't Hart BM, Henriques DY. Separating predicted and perceived sensory consequences of motor learning. PLoS One. 2016;11(9):e0163556.

111. Ruttle JE, Cressman EK, 't Hart BM, Henriques DY. Time course of reach adaptation and proprioceptive recalibration during visuomotor learning. PLoS One. 2016;11(10):e0163695.

112. Ruttle JE, 't Hart BM, Henriques DY. Reduced feedback barely slows down proprioceptive recalibration. Journal of Neurophysiology. 2022;128(6):1625-33.

113. Tzvi E, Koeth F, Karabanov AN, Siebner HR, Krämer UM. Cerebellar–premotor cortex interactions underlying visuomotor adaptation. Neuroimage. 2020;220:117142.

114. Tzvi E, Gajiyeva L, Bindel L, Hartwigsen G, Classen J. Coherent theta oscillations in the cerebellum and supplementary motor area mediate visuomotor adaptation. NeuroImage. 2022;251:118985.

115. Donchin O, Rabe K, Diedrichsen J, Lally N, Schoch B, Gizewski ER, et al. Cerebellar regions involved in adaptation to force field and visuomotor perturbation. Journal of neurophysiology. 2012;107(1):134-47.

116. Krakauer JW, Ghilardi M-F, Mentis M, Barnes A, Veytsman M, Eidelberg D, et al. Differential cortical and subcortical activations in learning rotations and gains for reaching: a PET study. Journal of neurophysiology. 2004;91(2):924-33.

117. Izawa J, Criscimagna-Hemminger SE, Shadmehr R. Cerebellar contributions to reach adaptation and learning sensory consequences of action. Journal of Neuroscience. 2012;32(12):4230-9.

118. Seidler R, Noll D, Chintalapati P. Bilateral basal ganglia activation associated with sensorimotor adaptation. Experimental brain research. 2006;175:544-55.

119. Anguera JA, Russell CA, Noll DC, Seidler RD. Neural correlates associated with intermanual transfer of sensorimotor adaptation. Brain research. 2007;1185:136-51.

120. Haar S, Donchin O, Dinstein I. Dissociating visual and motor directional selectivity using visuomotor adaptation. Journal of Neuroscience. 2015;35(17):6813-21.

121. Griffa G, Deleglise A, Solano A, Jacobacci F, De Pino G, Della-Maggiore V. The human hippocampus is involved in implicit motor learning. bioRxiv. 2024:2024.08.30.610548.

122. Della-Maggiore V, McIntosh AR. Time course of changes in brain activity and functional connectivity associated with long-term adaptation to a rotational transformation. Journal of neurophysiology. 2005;93(4):2254-62.

123. Galea JM, Vazquez A, Pasricha N, Orban de Xivry J-J, Celnik P. Dissociating the roles of the cerebellum and motor cortex during adaptive learning: the motor cortex retains what the cerebellum learns. Cerebral cortex. 2011;21(8):1761-70.

124. Imamizu H, Miyauchi S, Tamada T, Sasaki Y, Takino R, PuÈtz B, et al. Human cerebellar activity reflecting an acquired internal model of a new tool. Nature. 2000;403(6766):192-5.

125. Schlerf J, Ivry RB, Diedrichsen J. Encoding of sensory prediction errors in the human cerebellum. Journal of Neuroscience. 2012;32(14):4913-22.

126. Wolpert DM, Miall RC, Kawato M. Internal models in the cerebellum. Trends in cognitive sciences. 1998;2(9):338-47.

127. Mista C, Arguissain F, Ranieri A, Nielsen J, Andersen H, Manresa JB, et al. Spatio-temporal modulation of cortical activity during motor deadaptation depends on the feedback of task-related error. Behavioural Brain Research. 2024;468:115024.

128. Popa LS, Hewitt AL, Ebner TJ. Predictive and feedback performance errors are signaled in the simple spike discharge of individual Purkinje cells. Journal of Neuroscience. 2012;32(44):15345-58.

129. Popa LS, Streng ML, Ebner TJ. Long-term predictive and feedback encoding of
motor signals in the simple spike discharge of Purkinje cells. Eneuro. 2017;4(2).
130. Popa LS, Ebner TJ. Cerebellum, predictions and errors. Frontiers in cellular
neuroscience. 2019;12:524.
131. Torrecillos F, Alayrangues J, Kilavik BE, Malfait N. Distinct modulations in
sensorimotor postmovement and foreperiod β-band activities related to error
salience processing and sensorimotor adaptation. Journal of Neuroscience.
2015;35(37):12753-65.
132. Haar S, Donchin O. A revised computational neuroanatomy for motor control. Journal
of Cognitive Neuroscience. 2020;32(10):1823-36.
133. Hardwick RM, Lesage E, Eickhoff CR, Clos M, Fox P, Eickhoff SB. Multimodal
connectivity of motor learning-related dorsal premotor cortex. Neuroimage.
2015;123:114-28.
134. Tanaka H, Sejnowski TJ, Krakauer JW. Adaptation to visuomotor rotation through
interaction between posterior parietal and motor cortical areas. Journal of
neurophysiology. 2009;102(5):2921-32.
135. Darainy M, Manning TF, Ostry DJ. Disruption of somatosensory cortex impairs motor
learning and retention. Journal of Neurophysiology. 2023;130(6):1521-8.
136. Ebrahimi S vdVB, Ostry DJ. The Consolidation of Newly Learned Movements
Depends upon the Somatosensory Cortex in Humans. Journal of Neuroscience.
2024;44(32).
137. Ebrahimi S, Ostry DJ. The human somatosensory cortex contributes to the encoding
of newly learned movements. Proceedings of the National Academy of Sciences.
2024 Feb 6;121(6):e2316294121.
138. Hutter SA, Taylor JA. Relative sensitivity of explicit reaiming and implicit motor
adaptation. Journal of neurophysiology. 2018;120(5):2640-8.
139. McDougle SD, Taylor JA. Dissociable cognitive strategies for sensorimotor learning.
Nat Commun. 2019;10(1):40.
140. Cheng S, Sabes PN. Modeling sensorimotor learning with linear dynamical systems.
Neural computation. 2006;18(4):760-93.
1792