**Perceptual Prediction Error Supports Implicit Process in Motor Learning**

Xiaoyue Zhang[1], Wencheng Wu[1], Kunlin Wei[1,2,3,4] *

[1]School of Psychological and Cognitive Sciences, Peking University, Beijing, China
[2]Beijing Key Laboratory of Behavior and Mental Health, Beijing, China
[3]Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, China
[4]National Key Laboratory of General Artificial Intelligence, Beijing, China

*Correspondence to: Kunlin Wei
        School of Psychological and Cognitive Sciences
        Peking University
        Yiheyuan Road
        Beijing, 100871, China
        wei.kunlin@pku.edu.cn

**Abstract**

Error-based learning underlies motor learning, but what specific motor error drives implicit learning, the procedural component of motor skill, is unclear. A typical action consists of a movement and a performance outcome, e.g., grabbing a coffee cup involves a reaching movement and its actual landing of the hand relative to the target cup. While performance error is fundamental for the cognitive component of motor learning, what error, either performance or movement prediction error, underlies implicit motor learning has not been resolved. These two errors are hard to disentangle as the performance outcome is an integral part of the movement. Here we used the classical visuomotor adaptation paradigm, in which people learn to counter visual perturbations by deliberately aiming off the target, to dissociate the performance error from the prediction error. Using a series of behavioral experiments and model comparisons, we revealed that movement prediction error, but not performance error, can parsimoniously explain diverse learning effects. Importantly, despite the perturbation is visual, the movement prediction error is not specified in visual terms, but determined by a perceptual estimate of the hand kinematics. In other words, contrary to the widely-held concept of sensory prediction error, a perceptual prediction error drives implicit motor learning.

**Introduction**

Learning from errors is a fundamental aspect of human cognition, particularly in the realms of perception and action (Clark, 2013; Dosher & Lu, 2017; Wolpert et al., 2011). A classic example is the basketball player's effort to improve their jump shot; when a shot lands left of the basket, the player consciously adjusts their aim to the right in subsequent attempts. This explicit learning is complemented by an implicit learning process, which involves subconscious fine-tuning of bodily movements without conscious correction or even awareness of these fine-tunes. While performance error is recognized as a key driver of explicit learning across various motor tasks (Batterink et al., 2015; Ellis, 2009; Green & Flowers, 1991), the specific errors that facilitate implicit learning—a defining feature of motor skill acquisition—are still a matter of intense debate.

The debate on motor errors centers around two primary categories: performance error or
movement prediction error (Jordan & Rumelhart, 1992). Performance error is typically
defined as the discrepancy between movement outcome and the intended goal (Albert et
al., 2022; Hadjiosif et al., 2021; Miyamoto et al., 2020; Smith et al., 2006), while movement
prediction error refers to the discrepancy between the executed movement and its
predicted sensory consequences (Krakauer et al., 2019; Mazzoni & Krakauer, 2006;
Morehead et al., 2017; Shadmehr et al., 2010; Wolpert & Kawato, 1998). The complexity
of dissociating these two errors arises from their simultaneous occurrence during action
execution. For instance, when reaching for a coffee cup, the desired trajectory of the hand
and the desired final position must both be considered (Harris & Wolpert, 1998). The
trajectory prescribes a movement prediction error, and the final position prescribes a
performance error, both theoretically leading to implicit learning for subsequent
movements (Krakauer et al., 2019; Morehead & Xivry, 2021; Shadmehr et al., 2010;
Wolpert et al., 2011).

Prominent computational models of motor learning (Albert et al., 2022; Hadjiosif et al.,
2021; McDougle et al., 2015, 2017; Miyamoto et al., 2020; Smith et al., 2006) and
foundational textbooks (e.g., Schmidt et al., 2018) have primarily focused on performance
error (PE) to explain implicit learning behaviors. In contrast, a growing body of behavioral
and neurophysiological research highlights the role of sensory prediction error (SPE) —
the difference between the actual movement trajectory and the intended movement
trajectory—as a critical factor in driving implicit learning (Krakauer et al., 2019; Lee et al.,
2018; Mazzoni & Krakauer, 2006; Schween et al., 2014; Shadmehr et al., 2010; Taylor &
Ivry, 2011; Tsay et al., 2022a; Tseng et al., 2007). Resolving this theoretical inconsistency
in the understanding of implicit motor learning is not only essential for cognitive science
(Jordan & Rumelhart, 1992) but also promises to translate algorithmic insights from
human procedural learning into applied fields such as robotic control and artificial
intelligence (Kayacan & Chowdhary, 2019).

In this study, we harness the classical visuomotor rotation (VMR) adaptation paradigm —
an established framework for investigating error processing in motor learning—to

elucidate the competing error signals that contribute to implicit learning (Krakauer et al.,

2019; Morehead & Xivry, 2021). During the VMR task, participants encounter a hand

cursor that moves in a skewed direction relative to their actual hand motion, prompting

them to re-aim their movements away from the intended target. This dissociation between

movement prediction (re-aiming) and goal attainment (cursor motion) allows us to

separate PE and SPE within a single action (Figure 1A): PE is the deviation of the final

cursor position from the target, while SPE is characterized by the deviation of the visual

cursor motion from the re-aiming movement. Importantly, SPE is based solely on visual

feedback, with the implicit assumption that the motor system adapts itself based on the

visual consequence of actions, i.e., the skewed cursor motion. However, spatially locating

one's moving body effector is the foundation of motor control (Hogan & Flash, 1987;

Kawato, 1999; Sober & Sabes, 2005). We thus propose a novel third category of error:

perceptual prediction error (PPE), which represent the discrepancy between the

*perceived* hand movement from the predicted re-aiming movement. The perceived hand

is based on sensory integration of multiple spatial cues (Zhang et al., 2024), contrasting

with SPE, which relies exclusively on visual information.

To investigate these concepts, we designed a series of five experiments that

systematically manipulated VMR perturbations, revealing distinct characteristics of

implicit motor learning. Through comprehensive behavioral analyses and modeling, we

present compelling evidence that perceptual prediction error, rather than performance

error or sensory prediction error, serves as the primary driver of implicit motor learning.

This finding offers a parsimonious explanation for our new results while reconciling them

with established literature in the field.

**Results**

***Errors and models***

In the VMR paradigm, participants need to move a directionally rotated cursor toward the

target, and the directional deviation of the cursor ($x_v$) from the target defines the PE

(Figure 1A). PE drives the participants to adapt their movements in the subsequent trial

by consciously  re-aiming away from the target in the opposite direction of the rotation

perturbation (Taylor et al., 2014). This re-aiming reflects the participant's prediction of
where their hand will move, generating a sensory prediction of the reaching action ($x_u$).
The deviation between the actual cursor motion and this prediction is regarded as a
sensory prediction error, given its purely visual nature (Jordan & Rumelhart, 1992;
Krakauer et al., 2019). However, we propose that since the predicted variable pertains to
the intended *hand* direction, the relevant feedback should also concern the hand rather
than the cursor (Zhang et al., 2024). Thus, the perceived hand ($\hat{x}_{hand}$), rather than the
cursor, is compared to the predicted hand ($x_u$), with their discrepancy constitutes a
perceptual prediction error. The perceived hand ($\hat{x}_{hand}$) is derived from a combination of
multiple sensory cues, including the visual cue ($x_v$) from the cursor, the proprioceptive
cue ($x_p$) from the actual hand, and the sensory prediction ($x_u$). These cues are integrated
in accordance with Bayesian principles, with each cue weighted by its relative reliability.
136
The three types of errors—PE, SPE, and PPE—exhibit distinct temporal profiles
throughout the learning process, such as during a 30° counterclockwise VMR (Figure 1B
& C). When the perturbation is introduced, the cursor initially deviates 30° from the target,
and through trial-by-trial gradual learning, the cursor home back toward the target. This
results in an exponential decrease in PE, driven by the gradual correction of hand
movement direction. In contrast, re-aiming behavior—which reflects explicit learning—
initially rises sharply and then exhibits a small decline over the course of learning (Bond
& Taylor, 2015). The re-aiming behavior dictates the dynamics of the prediction errors
(Figure 1C): SPE remains large throughout the learning process because the re-aiming
direction stays consistently offset from the cursor. In contrast, PPE remains relatively
small and stable during learning, as the re-aiming direction is close to the perceived hand
direction, which is derived from a combination of sensory cues ($x_u$, $x_v$, and $x_p$; Figure 1A).
149
To determine which error best explains the diverse learning phenomena observed in
motor adaptation, we implemented a unified model framework. State-space models,
which have been successfully used to model error-based learning in visuomotor
adaptation (Albert et al., 2022; Cheng & Sabes, 2006; Herzfeld et al., 2014; McDougle et

al., 2015, 2017), were applied to test whether implicit learning is driven by PE, SPE, or

PPE (see Methods for model details).

The perceptual estimation of hand location is contingent on the relative reliability of the

individual sensory cues. Importantly, visual uncertainty for $x_v$ increases with larger SPEs

(Zhang et al., 2024), as participants tend to fixate not on the target but closer to their re-

aiming direction during the movement (Figure S1; Bromberg et al., 2019; de Brouwer et

al., 2018). This results in the cursor deviating from the fixation point by an angular

separation equivalent to the magnitude of the SPE. Our previous work demonstrated that

visual uncertainty regarding cursor motion increases linearly with this eccentricity (Zhang

et al., 2024). We also confirmed that nonlinear, instead of linear, eccentricity effect on

visual uncertainty would not affect model comparison results (Figure S4, Table S7). While

large visual perturbations induce both large SPE and PE (at least during early learning;

Figure 1C), its effect on PPE is dampened due to the high visual uncertainty associated

with large perturbations. Visual uncertainty reduces the influence of the visual feedback

on the perceived hand direction ($\hat{x}_{hand}$) according to Bayesian principles of perception

(Ernst & Banks, 2002). By incorporating the perceptual estimate of hand direction into the

prediction error, the dynamics of PPE diverge significantly from those of PE and SPE,

thus predicting implicit learning differently.

**Figure 1**. Illustration of sensorimotor cues and error dynamics in visuomotor rotation (VMR) learning. **(A)** Schematic representation of the key sensory cues and errors during VMR learning. The rotated visual cursor, actual hand position, and re-aiming direction correspond to the visual cue ($x_v$), proprioceptive cue ($x_p$), and sensory prediction ($x_u$), respectively. Performance error (PE) is the discrepancy between the cursor and the target, sensory prediction error (SPE) is the discrepancy between the cursor and the sensory prediction, and perceptual prediction error (PPE) is the discrepancy between the prediction and the perceived hand direction ($\hat{x}_{hand}$) combination, computed from a Bayesian integration of $x_v$, $x_p$, and $x_u$. **(B)** Simulation of a 30° VMR learning using the PPE model. The time-series of the variables from (A) are shown, with solid lines representing variables that are directly observable, and the dashed line represents the perceived hand direction ($\hat{x}_{hand}$), which is inferred but not directly observable. **(C)** Error dynamics during learning under a 30° VMR perturbation. Simulated PE and SPE match well with the observed data (dots; Bond & Taylor, 2015). Distinctly, simulated PPE remains small and stable, but quantitatively match our previous estimation of this error (Zhang et al., 2024).

### *Experiment 1: Perturbation size-dependency of implicit learning reflects the dynamics of PPE, not PE or SPE*

Manipulating the size of visual perturbations is a straightforward way to probe error-based learning and to differentiate between distinct error signals. In VMR, larger perturbations increase overall and explicit learning, but implicit learning remains stable across a broad range of perturbation sizes (Bond & Taylor, 2015; Kim et al., 2018; Morehead et al., 2017; Neville & Cressman, 2018). This classical finding has been interpreted as a hard limitation of implicit learning (Bond & Taylor, 2015; Kim et al., 2018; Morehead & Xivry, 2021; Tsay et al., 2021, 2022b) or a result of competition of explicit and implicit learning, where the explicit learning "siphons" away performance error (Albert et al., 2022).

The three error types—PE, SPE, and PPE—behave differently during early learning. Both PE and SPE increase with larger perturbations, but PPE remains small (Figure 1B). Previous studies have often overlooked early learning, focusing on late learning (e.g., Albert et al., 2022). We examined the entire learning process using various perturbation sizes (15°, 30°, 60°, and 90° VMR) by analyzing a prior dataset first (Figure 2A; Bond & Taylor, 2015). Since the participants reported their aiming direction for each trial, we could calculate the PE and SPE throughout learning (Figure 2B). As expected, larger perturbations led to an increase in both PE and SPE during early learning, but the scaling effect maintains during late learning only for SPE, not for PE (Figure 2B, middle and right; Figure 2D). Remarkably, implicit adaptation did not scale with perturbation size. Instead, it remained unchanged or even slightly decreased, both in early and late learning (Figure 2E; one-way ANOVA, early: $F(3,36) = 0.2$, $p = 0.897$; late: $F(3,36) = 1.44$, $p = 0.247$). Neither the PE nor SPE model captured this pattern, as both models predicted a linear increase in early implicit learning with perturbation size. The SPE model also incorrectly predicted a linear increase in late implicit learning (Figure 2E). In contrast, the PPE model successfully explained across perturbation sizes for both early and late stages (Figure 2C & 2E), as PPE remains largely invariant across perturbation sizes (Figure 2B left & 2D). Model comparison showed that the PPE model significantly outperformed both the PE and SPE models ($R^2$ = 0.708, 0.177, and -0.730, BIC = 182.04, 260.69, and 528.63 for PPE, PE, and SPE models, respectively; Table S1).

The fact that PPE does not exhibit a linear increase with perturbation size, unlike PE and SPE, stems from the nature of perceptual estimation. PPE is based on the perceived hand direction, which integrates visual and proprioceptive cues, each weighted by its uncertainty. Prior models based on PE and SPE have overlooked the fact that visual uncertainty increases with perturbation size (Zhang et al., 2024). During VMR learning, participants tend to fixate near their re-aiming direction, not on the target (Bromberg et al., 2019; de Brouwer et al., 2018). We confirmed this through a control experiment where we measured eye fixations during VMR learning (Figure S1). Larger perturbations had larger explicit learning with fixation further away from the target. This greater angular separation between the cursor and the fixation point increases visual uncertainty, reducing the influence of visual perturbation on hand localization ($\hat{x}_{hand}$) according to Bayesian principles (Zhang et al., 2024). By incorporating visual uncertainty into the PPE model, we captured the classical finding of invariant implicit adaptation over perturbation size. This result suggests the apparent limit on implicit learning is not a reflection of a fundamental constraint in the learning system, nor is it due to competition for performance error.

**Figure 2**. Behavioral and model fitting results from *Bond & Taylor, 2015*. **(A)** Total, explicit, and implicit learning during VMR adaptation with varying perturbation sizes. Explicit learning was estimated from participants' reported aiming directions. Solid lines represent the mean, and shaded areas depict the standard error (SEM) for each perturbation condition. **(B)** Changes in perceptual prediction error (PPE), performance error (PE), and sensory prediction error (SPE) during learning. PE and SPE were computed from data,

while PPE was estimated through model fitting. The three errors show distinct temporal
profiles, particularly during early learning. **(C)** Model fits for implicit learning across
perturbation sizes, with explicit learning treated as known data. The PPE model
accurately replicated the implicit learning patterns. In contrast, the PE model performed
poorly for large perturbations (60° and 90°), and the SPE model struggled with small
perturbations (15° and 30°). **(D)** The absolute error magnitudes of PPE, PE, and SPE as
a function of perturbation size during early (solid lines) and late (dashed lines) learning.
While PE and SPE scale with perturbation sizes, PPE remain relatively stable across
perturbations. **(E)** Data and model predictions for early (left) and late (right) implicit
learning. Both early and late learning show minimal scaling across perturbation sizes, in
contrast to the predictions of the PE and SPE models. The model predictions for implicit
learning reflect the error dynamics illustrated in (D). Error bars represent SEMs for the
data and bootstrap-derived standard deviations for the model.

The trial-by-trial aiming reports in *Bond & Taylor., 2015* enabled quantification of PE and
SPE throughout learning. However, frequent aiming reports can alter the composition of
explicit and implicit learning (Bond & Taylor, 2015; Maresch et al., 2021a, 2021b; Taylor
et al., 2014). To minimize this potential confound, we conducted Experiment 1, which
largely replicated Bond et al.'s study but removed the trial-by-trial aiming reports. Instead,
we inserted a small number of exclusion trials during late learning to specifically measure
implicit learning (Maresch et al., 2021b; Ruttle et al., 2021; Werner et al., 2015).
Additionally, the PPE model predicts a biased perceived hand localization after each
movement, which cannot be directly measured.  In Experiment 1, we measured this bias
indirectly by requiring the participants judge the location of their passively moved hand
immediately after an adaptation trial (Figure 3A; Zhang et al., 2024). People's judgement
should be biased in the direction of the visual perturbation, an effect termed as
proprioceptive bias (Henriques & Cressman, 2012; Salomonczyk et al., 2011, 2013; Rossi
et al., 2021; Ruttle et al., 2016, 2018, 2021, 2022). Importantly, we hypothesized that the
magnitude of proprioceptive biases should correlate with the PPE and the implicit learning
estimated by the model.

Experiment 1 recruited four independent groups of participants (n = 15 per group) to adapt
to 15°, 30°, 60°, or 90° VMR (Figure 3A). The overall learning patterns were consistent
with previous studies (Bond & Taylor, 2015; Neville & Cressman, 2018; Tsay et al., 2021).
Implicit learning, measured through exclusion trials during late adaptation, exhibited a
slightly nonlinear dependency on perturbation size (Figure 3D). A one-way ANOVA
revealed a significant main effect of perturbation size in implicit learning ($F$(3,56) = 7.15,
$p$ = 0.0004). Post-hoc tests showed that implicit learning was significantly smaller for the
90° perturbation compared to both 15° ($p$ = 0.046) and 30° ($p$ < .0001), suggesting a
concave size-dependency when continuous aiming reports were not required.

Despite reasonable fits to total and explicit learning (Figure 3C), the PE and SPE models
incorrectly predicted a scaling effect of implicit learning with perturbation size. This
discrepancy arose PE and SPE scaled with the perturbation size (Figure 3B), as observed
in Bond & Taylor's study (Figure 2B). In contrast, the PPE model accurately predicted
implicit learning, showing that PPE first increased and then decreased with perturbation
size, leading to a concave pattern that matches the empirical data (Figure 3B & C, $R^2$ =
0.969; Table S2). The model's parameters aligned with previous estimates, with implicit
and explicit learning rates ($B_i$ = 0.151 and $B_e$ = 0.513) comparable to previous studies
(McDougle et al., 2015, 2017). The slope of visual uncertainty with perturbation size ($k$ =
0.204) was slightly lower than that in the error-clamp paradigm ($k$ = 0.31; Zhang et al.,
2024), likely because participants in the VMR paradigm focus more on the cursor to
ensure accurate performance. Eye-tracking data from the control experiment confirmed
that participants tended to fixate back to the cursor after completing their movements
(Figure S1B). This contrasts with the error-clamp paradigm, where participants are
instructed to ignore the cursor and fixate on the target, resulting in higher visual
uncertainty.

**Figure 3**. Results of Experiment 1. **(A)** Learning curves for VMR across different perturbation sizes. Learning starts after baseline, gradually reaches a steady-state, and then decreases significantly during the no-feedback washout phase. Colored lines represent the mean, and shaded areas represent the SEM for each condition. Dark and light gray shading regions indicate the timing of proprioception test trials ("Prop-test") and exclusion trials ("Ex-test"). The group average and SEM for implicit learning and proprioceptive bias are shown. Proprioceptive judgements are biased in the direction of the visual perturbation, which is opposite to the direction of learning. **(B)** Time course of the three error types, with data shown as solid lines and model estimates as dash lines. Shading denotes SEM. Similar to *Bond & Taylor, 2015*, perturbation size scales the early PE and overall SPE, but not PPE. **(C)** Model fitting results for total, explicit, and implicit learning. All three models reproduced the scaling effect of perturbation size for total and explicit learning. However, for implicit learning, only the PPE model accurately predicted the concave relationship observed in the data, whereas both PE and SPE models erroneously predicted a linear scaling effect. Error bars represent SEM for data and bootstrapped standard deviations for the model. The black dots represent individual data. **(D)** Implicit learning and proprioceptive bias as a function of perturbation size. Both exhibited a similar nonlinear relationship (* denote $p < .05$ and *** denote $p < .001$). Note that the proprioceptive bias has been sign-flipped here to facilitate comparison with

implicit learning. **(E)** Implicit learning was significantly correlated with proprioceptive bias

on the individual level, consistent with the predictions of the PPE model. Each dot denotes

the two variables measured in each condition.

The proprioceptive biases significantly correlated with implicit learning, as predicted by

the PPE model. During late adaptation, participants' right hands, the hand used for

reaching adaptation, were passively moved to a random position near the target, and they

were asked to indicate the direction of the right hand using their left hand. This passive

localization test revealed a proprioceptive bias towards the visual perturbation, a

phenomenon known as proprioceptive recalibration (Modchalingam et al., 2019; Ruttle et

al., 2021; Salomonczyk et al., 2011, 2013; Zhang et al., 2024). Importantly, the bias

followed a concave trend across perturbation size, mirroring the pattern of implicit learning

(Figure 3D). A one-way ANOVA revealed a significant main effect of perturbation size

($F(3,56)$ = 3.7, $p$ = 0.017), with post-hoc test revealing a higher bias for 30° than for 90°

perturbations ($p$ = 0.011). A correlation analysis across participants showed a significant

positive correlation between proprioceptive bias and implicit adaptation (Figure 3E; $R$ =

0.41, $p$ = .001). This high correlation aligns with the PPE model, as both measures are

driven by the misperceived hand direction under visuomotor rotation ($\hat{x}_{hand}$, Figure 1). In

the proprioception test trial, the proprioceptive judgment is biased by the misperception

of the hand location experienced after the preceding adaptation trial ends (Zhang et al.,

2024), resulting in a small but consistent perceptual bias in the direction of visual

perturbation.

In summary, implicit learning exhibited either a concave size-dependency (Experiment 1;

Tsay et al., 2021) or invariance (Bond & Taylor, 2015) across perturbation sizes,

depending on the task specifics, such as the presence of frequent aiming reports. Neither

the PE nor SPE model could account for these size-dependencies, as their error sources

did not follow a similar size-dependency (Figure 2D & 3B). Interestingly, although PE

converges a small value across different perturbation sizes during late learning, enabling

the PE model to capture late learning size dependencies (Albert et al., 2022), it fails to

account for early learning, where it erroneously predicts a size-scaling effect. Only the

PPE model provides a parsimonious explanation the size dependencies of implicit learning across both early and late learning phases (Figures 2F & 3C).

### ***Experiment 2: Stepwise perturbations enhance implicit learning due to increased PPE***

The differences between the three competing errors are most pronounced at the time when a perturbation is initially introduced (Figure 1C, 2B & 3B). Therefore, if the perturbation size changes *during* learning, we expect these errors to have diverging effects. A previous study has shown that, compared to a single large perturbation (e.g., a 60° VMR, referred to as one-step perturbation herein), incrementally increasing perturbations (e.g., a 15→30→45→60°, referred to as stepwise perturbation) boosts implicit learning while reducing explicit learning (Modchalingam et al., 2023). The reduction in explicit adaptation is expected due to smaller PE in stepwise conditions. However, the increase of implicit learning challenges the conventional view that implicit learning is stereotypical and unaffected by various manipulations (Bond & Taylor, 2015; Kim et al., 2018; Morehead et al., 2017; Neville & Cressman, 2018), and its mechanism remains unclear. Importantly, stepwise adaptation offers a critical test of the three error models. For both PE and SPE, one-step perturbation generates larger errors than stepwise perturbation during early learning, predicting greater implicit learning for the one-step condition (Figure 4B, middle and right). In contrast, the PPE model predicts the opposite: smaller PPE for one-step perturbation due to the discounting effect of visual uncertainty, and thus, less implicit learning (Figure 4B, left).

To test these predictions in Experiment 2, we recruited two groups of participants: one group experienced a one-step 60° VMR perturbation (n = 15), while the other group experienced a stepwise 15°→30°→45°→60° VMR perturbation (n = 15; Figure 4A). Implicit learning was assessed using exclusion trials during late learning. Both groups achieved similar steady-state performance by the end of learning (two-sample *t*-test, $t(28)$ = -1.796, $p$ = 0.083), but their implicit learning patterns diverged (Figure 4C). The stepwise group showed increasing implicit learning with each perturbation step ($F(3,21)$ = 11.639, $p < 0.001$), while the one-step group showed no significant change in implicit adaptation

over time ($F(3,21) = 1.938$, $p = 0.154$). Notably, from the second exclusion trial block onward, the stepwise group exhibited significantly greater implicit learning than the one-step group ($p = 0.011$). By the final exclusion trial block (at 60° VMR), implicit learning in the stepwise group was nearly double that of the one-step group (26.33° ± 2.74° vs. 12.74° ± 1.45°; $p = .001$). In contrast, explicit learning was consistently larger for the one-step group for each exclusion trial block ($p < 0.001$).

**Figure 4**. Results of Experiment 2. **(A)** Learning of the one-step (yellow) and stepwise (brown) group. The dark grey shading indicates the timing range for the proprioception test trials, and the light grey shading indicates the timing range for the exclusion trials. Squares with error bars represent implicit learning measured via exclusion trials ("Ex-test"), while diamonds with error bars represent proprioceptive bias measured by proprioception test trials ("Prop-test"). **(B)** Time courses of the errors. The solid lines and dots represent data and the dashed lines represent model estimates. PE and SPE were

larger for one-step than for stepwise perturbations during early learning, while PPE showed the opposite condition difference. **(C)** Time courses of explicit and implicit learning. Both explicit and implicit learning increased over time for stepwise adaptation, but remained unchanged for one-step adaptation. Note the difference between conditions: while explicit learning was larger for one-step adaptation, implicit learning was larger for stepwise adaptation especially during late learning (* denote $p < 0.05$, ** denote $p < 0.01$, and *** denote $p < 0.001$). Error bars are SEMs for the data. The black dots represent individual data. **(D)** Model fittings for implicit learning. Error bars are bootstrap-derived standard deviations for the model (resample size = 5000). Only the PPE model replicated the actual learning patterns, while both PE and SPE model predicted one-step adaptation was larger than stepwise adaptation in most cases. **(E)** Time course of proprioceptive bias in the two conditions. The sign of proprioceptive bias was flipped to aid the comparison. With larger PPE, stepwise adaptation elicited larger proprioceptive bias than one-step adaptation. **(F)** Implicit learning and proprioceptive bias plotted as a function of perturbation size during stepwise adaptation. **(G)** Implicit learning was significantly correlated with proprioceptive bias on the individual level. Each dot denotes the two variables measured in each perturbation size during stepwise adaptation.

All three models predict the gradual increase in implicit learning with stepwise perturbations, as all errors increase incrementally (Figure 4B & 4D). However, only the PPE model correctly captures the relatively smaller implicit learning for the one-step perturbation compared to stepwise adaptation (Figure 4D). The PE and SPE models erroneously predict the opposite. Model comparison confirmed the superior performance of the PPE model ($R^2$ = 0.960, 0.933, and 0.925; BIC = 264.81, 309.59, and 317.67 for PPE, PE, and SPE models, respectively; more model comparisons, see Table S3). The superior performance of the PPE model was obtained even though we used the visual uncertainty parameter ($k$) estimated from Experiment 1, which also involved a one-step perturbation.

As in Experiment 1, proprioceptive bias in this experiment correlated with implicit learning, further supporting the PPE model. Proprioceptive bias increased with stepwise

perturbation (Figure 4E, $F(3,25) = 8.874$, $p < 0.001$), but remained unchanged during the

one-step perturbation ($F(3,25) = 0.538$, $p = 0.661$). These proprioceptive bias patterns

mirrored those of PPE and implicit learning (Figure 4F). To analyze the individual

difference, we used a mixed linear model to isolate the between-subject variability and

revealed a significant positive correlation between proprioceptive bias and implicit

learning ($\beta = 0.91$, $p < 0.001$; Figure 4G). This result was consistent with previous studies

on proprioception and motor adaptation (Salomonczyk et al., 2011, 2013; Tsay et al.,

2022b; Zhang et al., 2024), supporting that implicit adaptation and proprioceptive bias are

driven by the same error source, i.e., PPE.

One might argue that the one-step perturbation amplifies explicit learning compared to

the stepwise perturbation, while the model-based analyses above assume fixed explicit

learning parameters across both conditions. Indeed, when faced with a large, salient

perturbation, individuals may rely more heavily on explicit learning strategies (Albert et

al., 2022; Modchalingam et al., 2023). To address this, we allowed the explicit learning

parameters to vary between the two conditions and re-evaluated the models (Figure S2;

Table S3). As expected, the explicit learning rate was higher in the one-step condition

across all models. However, even with varying explicit learning parameters, both the PE

and SPE models still failed to account for the observed differences between conditions

(Figure S2B). While the PE model slightly improved its fit compared to the fixed-parameter

case (Figure 4), it could not capture the implicit learning patterns for small stepwise

perturbations (15° and 30°, Figure S2B). Furthermore, this improvement came at the cost

of unrealistic model parameters (Table S3). For example, the explicit learning rate in the

stepwise condition was estimated to be as low as 0.218, even smaller than the implicit

learning rate (0.223), which contradicts the typical finding that explicit learning rates are

several times larger than implicit learning rates (Coltman et al., 2019; McDougle et al.,

2015, 2017). The SPE model showed minimal improvement, continuing to underestimate

implicit learning in the stepwise condition and overestimate it in the one-step condition

(Figure S2B). In contrast, the PPE model continued to provide excellent fits across both

conditions, yielding the best overall fit ($R^2 = 0.962$, 0.958, and 0.926, BIC = 268.76, 280.84,

and 325.39 for PPE, PE, and SPE models, respectively).

We further validated the models using cross-validation, fitting each model with data from the stepwise group and testing it on the one-step group. Although the PE and SPE models fit the stepwise group reasonably well, their predictions for the one-step group were highly inaccurate (Figure S2C). Both models overestimated implicit learning and underestimated explicit learning in the one-step condition. In contrast, the PPE model provided a near-perfect match in the cross-validation test (Figure S2C; Table S3), further reinforcing its robustness in capturing the dynamics of both implicit and explicit learning across different perturbation schedules.

***Experiment 3: Suppression of explicit learning uncovers nonlinear implicit learning compatible only with PPE***

When a visual perturbation is first introduced, PE and SPE equal the perturbation size, as the participant has not yet developed a re-aiming strategy. In contrast, PPE is less than the visual perturbation because it derives multisensory cue combination for hand localization (Figure 1A). Thus, the learning to the very first perturbation can highlight the key differences between errors: both PE and SPE predict a linear learning function relative to perturbation size, while PPE predicts a nonlinear function, as larger perturbation increase visual uncertainty, reducing its own influence on hand localization (Wei & Kording, 2009; Zhang et al., 2024).

In Experiment 3 (n = 18), we used trial-by-trial random perturbations to suppress explicit learning and isolate the single-trial implicit learning. Unlike blocked perturbations as in Experiment 1 and 2, random perturbations prevent the accumulation of implicit learning. The participants performed single-target reaches, but each perturbation trial was perturbed by a visuomotor rotation with a pseudo-random size and direction (0, ±4°, ±8°, ±16°, ±32°, and ±64°), and was flanked by two no-feedback null trials (Figure 5A). Participants were instructed to aim directly at the target in the null trials, thus discouraging explicit learning. Implicit learning was measured by comparing hand movement directions between the pre- and post-perturbation null trials (Zhang et al., 2024). The perturbation

sequence, though appeared random, was identical for all participants. The perturbation

sequence had a zero mean, allowing any residual explicit learning to average out.

The participants displayed adaptive changes immediately after the perturbation sequence

was introduced, as evidenced by increased variability in hand directions (Figure 5B). We

fit the learning data with the three competing models. Since PE and SPE are equal to the

perturbation size in single-trial learning, we compared only the PPE model and PE model.

The PPE model outperformed the PE model in explaining the learning response to

random perturbations (Figure 5B, $R^2$ = 0.471 vs -0.683; BIC = 46.56 vs 82.00). As

predicted by the PPE model, implicit learning follows a nonlinear pattern: it increased with

perturbation size until a point, then perturbation size grew larger (Figure 5C; repeated-

measure ANOVA: $F(3.2,53.8)$ = 45.897, $p$ < .0001; 4° vs 8°: $p$ = 0.005; 8° vs 32°: $p$ =

0.001; 32° vs 64°: $p$ < .0001). This nonlinear pattern was consistent with previous findings

from various motor adaptation paradigms (Tsay et al., 2021; Wei & Kording, 2009; Zhang

et al., 2024). In contrast, the PE (SPE) model predicted a linear pattern (linear regression:

$y = a + b\theta$, with $a$ = -0.007 and $b$ = -0.064, $R^2$ = 0.908, $F$ = 89.1, $p$ < .0001), that

drastically deviated from the observed data (Figure 5C; $R^2$ < 0, $RMSE$ = 2.449°), while

the PPE model captured the nonlinear pattern well ($R^2$ = 0.954, $RMSE$ = 0.684°). Despite

the randomness of the perturbations, the slope of visual uncertainty ($k$) estimated in this

experiment was consistent with that from the block-design experiments ($k$ = 0.195 from

Experiment 2 and $k$ = 0.204 from Experiment 1 and 2), further supporting the robustness

of the PPE model across different experimental designs.

**Figure 5**. Results of Experiment 3. **(A)** Experimental design. In feedback (FB) trials, participants experienced visuomotor rotations with pseudo-random perturbation sizes and directions (0, ±4°, ±8°, ±16°, ±32°, and ±64°). Each feedback trial was flanked by two no-feedback (NFB) null trials, where participants were instructed to move directly to the target. Single-trial implicit learning was measured by the directional change between pre- and post-perturbation null trials. **(B)** Time course of hand movement directions across trials, along with model fits. The grey line (with shaded SEM) represents the data, with stars denoting feedback trials and circles representing no-feedback trials. The PPE model (red) provided a better fit to the data than the PE/SPE model (dark blue). **(C)** Single-trial learning as a function of perturbation size, derived from both trial-by-trial data and model fits. The PE/SPE model predicted a linear relationship between perturbation size and learning, while the PPE model predicted a nonlinear relationship. Color shades represent SEMs for the data and bootstrapped standard deviations for the model. The light grey lines represent individual participants' data.

Despite the task instructions, participants might have strategically re-aimed away from the visual target in the null trials, potentially "contaminated" the estimates of implicit

learning with explicit learning. To assess this, we employed a model-based analysis to
include an explicit learning component. The results from the PPE model revealed that the
estimated explicit learning parameters were quite small (learning rate $B_e$ = 0.047 and
retention $A_e$ = 0.037), indicating a negligible contribution from explicit learning.
Additionally, incorporating explicit learning into the model only marginally increase the $R^2$
but result in a higher BIC for the PPE model ($R^2$: 0.471 vs. 0.500; BIC = 46.56 vs. 52.82).
When we constrained the lower limit of $A_e$ to typical values (> 0.5), the inclusion of explicit
learning actually worsened the model fit ($R^2$ = 0.471 vs. 0.453, BIC = 46.56 vs. 57.26).
Similarly, the fit of the PE/SPE models deteriorated when explicit learning was added
(Table S4). Notably, the PPE model still substantially outperformed the PE and SPE
models even with the inclusion of explicit learning (Table S4). Overall, we conclude that
random perturbations primarily lead to implicit learning with a nonlinear size dependency,
which can only be explained by the PPE model.
546
***Experiment 4: Implicit learning elicited by instructed re-aiming is driven by PPE***
One of the seminal studies in motor neuroscience supporting the role of SPE employed
the instructed re-aiming paradigm in VMR (Mazzoni & Krakauer, 2006). In this paradigm,
participants are instructed to re-aim to a neighboring target whose angular deviation from
the original target matches the imposed visual perturbation (e.g., 45°; Figure 6A). This re-
aiming nullifies performance error (PE) immediately. Intriguingly, despite this, participants
begin to drift away from the re-aiming target involuntarily over repeated trials, increasing
PE. This drift has been interpreted as evidence that SPE drives implicit learning (Mazzoni
& Krakauer, 2006). Recently, Albert et al. (2022) proposed an alternative explanation,
suggesting that the drift could be driven by PE relative to the re-aiming target (PE_{re-}
_{aiming})—a form of PE that arises between the cursor and the re-aiming target rather than
the original target. Notably, this new PE formulation is essentially equivalent to SPE
(Figure 6B). However, our PPE could also explain this drift: the perceived hand position
is "pulled" toward the visual cursor, deviating from the re-aiming target, thus generating a
PPE and driving implicit learning. Our model simulations indicate that all three error
models (PE, SPE, and PPE) can explain the original findings by Mazzoni & Krakauer
(Figure 6C). However, the models make distinct predictions when the re-aiming offset

size changes. If the angular separation between the re-aiming and original targets
increases (e.g., from 45° to 90°), both SPE and $PE_{re-aiming}$ would predict a larger implicit
drift, as the errors increase with the offset. In contrast, the PPE model predicts a reduction
in implicit drift because larger visual perturbations reduce the influence of PPE due to
increased visual uncertainty.

**Figure 6.** Data and model fitting for the instructed re-aiming paradigm (Mazzoni &
Krakauer, 2006) and Experiment 4. **(A)** Instructed re-aiming paradigm illustration.
Participants initially moved a cursor toward a target without perturbation during baseline
trials, followed by a single rotation trial with a 45° visuomotor rotation. During subsequent
instructed re-aiming trials, they were instructed to move their hand toward a neighboring
"re-aiming target" 45° away, allowing the cursor to perfectly hit the original target. **(B)**
Motor errors during instructed re-aiming. Despite re-aiming to the new target, participants'
hands unknowingly "drift" away from the re-aiming target. In addition to the three errors
shown in Figure 1A, a new error termed re-aiming performance error ($PE_{re-aiming}$)—the
deviation of the visual cursor from the re-aiming target—is introduced (Albert et al., 2022;
Taylor & Ivry, 2011). This error is equivalent to SPE. **(C)** Performance relative to the
original target before and after instructed re-aiming. Data from Mazzoni & Krakauer

(2006). After the initial 45° rotation trial, participants performed instructed re-aiming trials,

with their hands drifting away from the re-aiming target. This implicit drift can be explained

equally well by the PPE, PE, and SPE models. **(D)** Results of Experiment 4. Participants

exhibited an implicit drift away from the intended direction after instructed re-aiming, with

the drift being significantly larger for the 45° re-aiming offset compared to the 90° re-

aiming offset. Colored lines with shaded areas represent means and SEMs for the two

conditions. **(E)** Extent of implicit learning observed and predicted by the three competing

models. The PPE model accurately predicts smaller implicit learning for the 90° condition

compared to the 45° condition, while both the PE and SPE models erroneously predict

the opposite. Error bars represent SEMs for the data and bootstrap-derived standard

deviations for the model. The black dots represent individual data.

To test these diverging predictions, Experiment 4 recruited two groups of participants who

adapted to either a large (90°, n = 15) or small (45°, n = 15) re-aiming offset. Upon re-

aiming, participants' movements began to drift in the opposite direction of the perturbation.

Critically, the drift for the 90° group was significantly smaller than that for the 45° group

(Figure 6E). The final drift in the 90° condition was considerably smaller than in the 45°

condition (13.17° ± 1.29° for 45°, 8.44° ± 1.22° for 90°; two-sample *t*-test: $t(28) = 2.58$, *p*

= 0.016). The PPE model readily explains this condition difference, as it predicts smaller

implicit drift for larger re-aiming offsets due to the reduced influence of visual

perturbations on hand localization. In contrast, both the PE and SPE models incorrectly

predict that the drift should increase in the 90° condition (Figure 6E; Table S5). The visual

uncertainty parameter (*k*) estimated from this experiment (0.211) was similar to those in

previous experiments, further supporting the robustness of the PPE model.

***Experiment 5: Performance error based on spontaneous re-aiming fails to induce***

***implicit learning***

In Experiment 4, participants were instructed to re-aim, preventing the spontaneous

formation of explicit learning. However, one might argue PE and SPE could still account

for implicit drift if re-aiming occurred spontaneously rather than being explicitly instructed.

To address this, Experiment 5 introduced a re-aiming target after participants had fully

developed their own explicit learning. In this experiment, participants (n = 20) adapted to a 45° VMR, with explicit learning intermittently measured using aiming reports during adaptation (Figure 7A). After reaching a steady-state learning phase (spontaneous re-aiming in Figure 7B), we presented a re-aiming target that aligned with each participant's last reported aiming direction. Participants were instructed to aim for this re-aiming target for 30 additional trials to assess potential implicit drift (instructed re-aiming in Figure 7B). Since this re-aiming target was consistent with the participants' spontaneous re-aiming direction, both SPE and PPE should remain unchanged (Figure 7C), and both models would predict no further implicit learning. In contrast, the PE model predicts that the introduction of the re-aiming target would create a new performance error (PE_re-aiming), leading to further implicit learning beyond the learning asymptote (Figure 7C).

Participants reported an average explicit learning of 29.53° ± 1.63° in their last aiming report, consistent with previous studies (Bond & Taylor, 2015; McDougle et al., 2015). However, no further increase in implicit learning occurred after the re-aiming target was introduced. The hand deviation from the original target with the re-aiming target (43.01° ± 0.50°) was statistically indistinguishable from the level before introducing the re-aiming target (43.25° ± 0.79°, $t(19) = 0.327$, $p = 0.747$, $BF = 0.24$). Moreover, implicit learning, as measured by exclusion trials, did not change when comparing the last exclusion trial cycle before re-aiming target introduction to the first trial cycle of the washout phase (9.04° ± 0.84° vs. 7.44° ± 0.66°, $t(19) = 1.888$, $p = 0.075$; Figure 7D). These findings show that introducing a re-aiming target after explicit learning has been formed does not induce further implicit learning, contradicting the PE model's predictions. Instead, the null result aligns with the prediction of the PPE model (Figure 7D), which also provided a better fit to the data ($R^2 = 0.880$ vs. 0.337 for the PE model, Table S6).

**Figure 7**. Results of Experiment 5. **(A)** Illustration of the re-aiming paradigm. Participants

reported their aiming direction using a numeric ring. After reaching asymptotic learning

for a 45° VMR, the last reported aiming direction was used to designate a re-aiming target

for each participant. **(B)** Learning in Experiment 5. Participants adapted to a 45° VMR

with explicit learning measured through verbal aiming reports and implicit learning

assessed using exclusion trials. After reaching the learning asymptote, participants

continued to aim at their chosen direction, now marked by a re-aiming target. The

introduction of the re-aiming target divides the learning into two phases, one for

spontaneous re-aiming and the other for instructed re-aiming. **(C)** Time courses of PPE,

PE, SPE, PE_re-aiming before and after the introduction of the re-aiming target. PE, SPE,

and PE_re-aiming were calculated from the data, while PPE was estimated by model fitting.

Note that the introduction of the re-aiming target results in a new PE_re-aiming, while PPE

and SPE remain unchanged. **(D)** Implicit learning observed and predicted by the models.

Implicit learning was measured using exclusion trials. The models were fit to the data

before the re-aiming target was introduced and used to predict learning afterwards. The

PE model failed to predict the observed re-aiming performance, while the PPE model

accurately predicted the absence of further implicit learning. Error bars represent SEMs

for the data and bootstrapped standard deviations for the model predictions. The black

dots represent individual data.

***Experiment 6: Direct measures of PPE reveals its correlation to implicit learning***

Thus far, PPE has been implicated through model analysis or indirectly measured by

proprioceptive bias in Experiments 1 and 2. In Experiment 6, we directly measured PPE

by asking participants to locate both their aiming and produced movement. Our model

simulation indicates that PPE should remain small throughout the learning process

(Figure 1C), and should exhibited a similar perturbation size-dependency as

proprioceptive bias and implicit learning (Figure 2D and 3D). To test these predictions,

we randomly assigned 30 participants to three conditions with 15°, 30°, and 90° VMR,

respectively (n = 10 each). For a probe trial, a reaching movement was preceded by a

pre-movement aiming report and followed by a post-movement report of perceived hand

direction (Figure 8A). The direction difference between these two reports reflects the

perceived error. Indeed, when the visual perturbation was turned on, the perceived hand

direction deviated away from the re-aiming direction and towards the visual perturbation

(Figure 8B). Thus, the PPE was significantly negative (15°: -1.99° ± 0.84°, $t(9)$ = -2.36, $p$

= 0.042; 30°: -4.39° ± 0.47°, $t(9)$ = -9.42, $p$ < 0.001; 90°: -1.86° ± 0.65°, $t(9)$ = -2.87, $p$ =

0.018), in the opposite direction of implicit learning (Figure 8C). Notably, the size of PPE

was rather small and remained small throughout learning, consistent with our initial model

simulations (Figure 1C). Importantly, PPE followed a nonlinear concave function of

perturbation size ($F(2,27)$ = 4.52, $p$ = 0.020, post-hoc test: 30° vs. 90°: $p$ = 0.038),

resembling the model-estimated PPE based on data from Bond 2015 and Experiment 1

(Figure 2D and 3D) and the proprioceptive bias measured in Experiment 1 (Figure 3D).

Furthermore, the concave PPE pattern revealed here was accompanied by a similar

concave pattern in implicit learning ($F(2,25)$ = 5.04, $p$ = 0.015, post-hoc test: 30° vs. 90°:

*p* = 0.012; Figure 8D). Across participants, PPE also positively correlated with implicit

learning (Figure 8E; *R* = 0.41, *p* = 0.031). Hence, the direct measures of PPE provide

further support for the driving role of PPE in implicit learning.

**Figure 8.** Results of Experiment 6. **(A)** Illustration of the probe trial that directly measures

PPE. Participants reported their aiming direction and perceived hand direction before and

after a perturbed reach, by placing a visual marker with their left hand via the KINARM

robot. The just-experienced visual perturbation biases the perceived hand direction away

from the aiming direction. **(B)** Learning performance and reported aiming/hand directions

from the three perturbation-size conditions are separately shown. Movement direction

relative to the target shows typical adaptation changes. Aiming direction and perceived

hand direction changes along with learning. Hand direction is more rotated than the

aiming direction, manifesting implicit learning; the aiming direction is more rotated than

the perceived hand direction, manifesting a perceptual error relative to the intended hand

direction (i.e., PPE). Shades and error bars denote SEM. **(C)** PPE as a function of probe

trials. PPE becomes negative with perturbation, in line with its causal role in driving

adaptation in the positive direction. **(D)** The magnitudes of both PPE and implicit learning

show a concave dependency on perturbation size. * denotes $p < 0.05$ for post-hoc

comparison. **(E)** Implicit learning significantly correlates with PPE as revealed by

generalized linear regression.

## Discussion

Using a combination of systematic behavioral experiments and computational modeling,

our study provides converging evidence that explicit and implicit processes in

sensorimotor learning are driven by distinct error signals. Explicit learning is driven by

performance error, while implicit learning is governed by perceptual prediction error. By

leveraging the classical visuomotor rotation paradigm, we disentangled three potential

error signals for implicit learning—performance error, sensory prediction error, and

perceptual prediction error—across five experiments. Our findings demonstrate that

implicit learning exhibits a perturbation size-dependency that can only be explained by

PPE, not by PE or SPE (Experiment 1 and 2, and Bond & Taylor, 2015). The perceptual

prediction error can be indirectly measured by proprioception tests, which show people's

perceived hand is biased and this bias in correlated with implicit learning (Experiment 1

and 2). When explicit learning is suppressed using random perturbations, implicit learning

follows a nonlinear size-dependency, again consistent only with PPE (Experiment 3).

When explicit learning is constrained by instructed re-aiming (Experiment 4), implicit

learning develops in a pattern that directly contradicts the predictions of SPE-based

learning (Mazzoni & Krakauer, 2006). When a re-aiming strategy is introduced after

explicit learning has already been established, no further implicit learning occurs, refuting the predictions of PE-based learning (Experiment 5). Directly measured by probe trials, PPE is found to remain negative with a small yet significant offset from the aiming direction (Experiment 6), a pattern drastically different from PE and SPE yet matching the model predictions (Figure 1B and 1C). Individual analysis reveals a positive correlation between PPE and implicit learning. Similar to implicit learning and proprioceptive bias, the measured PPE shows a concave dependency on perturbation size, further supporting its role in driving learning and biasing proprioceptive judgment. Overall, these results challenge the prevalent theories that attribute implicit motor learning to either performance or sensory prediction errors. Instead, they highlight the critical role of perceptual prediction error, defined as the deviation of the perceptual estimate of hand motion from the predicted (desired) motion, in driving implicit motor learning.

### *Performance error does not drive implicit motor learning*

Performance error is a well-established signal for learning in various domains, including perceptual learning (Lu & Dosher, 2022), reinforcement learning (Holroyd & Coles, 2002; Niv, 2009), decision-making (Johnson et al., 2013; Law & Gold, 2009), and the explicit components of motor learning (Albert et al., 2022; Krakauer et al., 2019; McDougle et al., 2015, 2017; Miyamoto et al., 2020). However, whether performance error also underlies the implicit component of motor learning—traditionally associated with procedural learning—remains a topic of heated debate. While the dominant view posits that sensory prediction error drives implicit learning (Jordan & Rumelhart, 1992; Krakauer et al., 2019; Mazzoni & Krakauer, 2006; Shadmehr et al., 2010; Tsay et al., 2022a; Tseng et al., 2007), all the influential state-space models endorse performance error as the key driver (Albert et al., 2022; McDougle et al., 2015, 2017; Miyamoto et al., 2020). A major challenge in this debate is the difficulty of dissociating performance error from movement prediction error, as performance error relates to goal attainment and is often tightly coupled with the movement itself. Previous studies have attempted to separate these errors by introducing "target jumps" during movement (Al-Fawakhiri et al., 2023; Leow et al., 2018, 2020; Tsay et al., 2022a), thus creating a performance error that is less correlated with the movement. However, such manipulations tend to distract participants, affecting both explicit and

implicit learning (Leow et al., 2018, 2020; Tsay et al., 2022a). Furthermore, the target
jump is likely to be interpreted as an environmental change, which leads to minimal
learning within the motor system (Berniker & Körding, 2008; Wei & Kording, 2009). Indeed,
the target jump has been shown to be ineffective in eliciting implicit learning (Oza et al.,
2024; Tsay et al., 2022a). In contrast, our study employed the classical VMR paradigm,
which does not require additional manipulations of the perturbation feedback. The
presence of explicit re-aiming—whether spontaneously learned or instructed—naturally
separates the predicted movement from the goal-seeking cursor, allowing us to dissociate
movement prediction error from performance error (Figure 1A). Our findings strongly
suggest that performance error cannot account for the various patterns of implicit motor
learning observed, despite its clear role in explicit learning.

It is worth noting that Albert and colleagues (2022) also employed a similar modeling
approach using extensive datasets from VMR paradigms, but they reached the
contrasting conclusion that both implicit and explicit learning are driven by the same
performance error. However, their analysis focused exclusively on late learning (final
learning extent) without considering the entire adaptation process. In late learning,
performance error and perceptual prediction error tend to converge in size (Figure 2B, 3B
& 7C), potentially leading to misleading model results. In our study, we demonstrated that
early learning—where PE and PPE differ significantly—reveals the limitations of the PE
model (Figure 2E). Additionally, Albert et al. did not compare stepwise adaptation with
one-step adaptation, a critical contrast that highlights the erroneous predictions of both
PE and SPE models (Figure 4D). In summary, by analyzing the full learning process, as
opposed to selective learning segments, our study refutes the notion that performance
error drives implicit motor learning.

***Motor prediction error from multisensory integration, not a single modality***
Our findings call for a shift in how prediction error is conceptualized in the context of motor
control and learning, emphasizing its perceptual nature rather than a strictly sensory one.
Traditional models of motor prediction error have largely focused on visual prediction
errors derived from the discrepancy between predicted and actual visual feedback during

movements (Krakauer et al., 2019; Morehead & Xivry, 2021). This approach has been
applied not only to visual perturbations such as visuomotor rotation and saccadic
adaptation (Pelisson et al., 2010; Wong & Shelhamer, 2011), but also to non-visual
perturbations such as force (Gonzalez Castro et al., 2011; Shadmehr & Mussa-Ivaldi,
1994; Smith et al., 2006; Thoroughman & Shadmehr, 2000) and inertial perturbations
(Brown et al., 2003; Duff & Sainburg, 2007). However, accumulating evidence suggests
that learning of force and inertial perturbations can occur without visual feedback (DiZio
& Lackner, 2000; Franklin et al., 2007; Marko et al., 2012; Striemer et al., 2019),
contradicting the idea that visual errors alone drive motor adaptation in these contexts.
Defining prediction error solely in terms of visual metrics is increasingly seen as
inadequate, as several recent studies have noted (Tsay et al., 2022; Zhang et al., 2024).
Despite this, the field continues to label prediction errors in motor control as sensory
prediction errors, reinforcing their unimodal, primarily visual nature (Krakauer et al., 2019;
Morehead & Xivry, 2021). Our study provides strong evidence that visual prediction error
cannot adequately capture the dynamics of implicit learning, even when the perturbation
itself is visual. For instance, SPE scales dramatically with perturbation size (Figure 2B &
3B), leading to exaggerated predictions regarding implicit learning, which often exhibits
more stable or mild changes. In contrast, perceptual prediction error (PPE), which is
derived from multisensory integration, remains more resilient to changes in visual
perturbation size and can parsimoniously explain not only our data but also key findings
in the literature (e.g., Bond & Taylor, 2015; Mazzoni & Krakauer, 2006; Modchalingam et
al., 2023).

Our findings suggest that perceptual prediction error drives implicit learning beyond
upper-limb movements, as supported by emerging evidence from other motor learning
paradigms. One such example is saccadic adaptation, which explores how the
oculomotor system adapts to spatial perturbations of gaze targets (Deubel, 1987;
McLaughlin, 1967; Wong & Shelhamer, 2011). In this domain, performance error (i.e., the
retinal distance between the post-saccadic target and the fovea) does not seem to
support implicit learning (Collins & Wallman, 2012; Wong & Shelhamer, 2011). Similar to
reaching studies, saccadic adaptation research has traditionally focused on sensory

prediction error (i.e., the deviation between the post-saccadic visual error and an

internally predicted visual error; Bahcall & Kowler, 1999; Collins & Wallman, 2012; Wong

& Shelhamer, 2011). However, recent modeling work suggests that both motoric and

perceptual changes in saccadic adaptation are better explained by a postdiction-based

error (Masselink & Lappe, 2021). This postdiction error refers to the discrepancy between

the eye-muscle motor command and the perceived target location, the latter arising from

a combination of the actual saccadic error and the predicted saccadic displacement. This

concept mirrors our perceptual estimate of hand localization in reaching adaptation,

wherein multiple sensorimotor cues are integrated to inform the perceptual error driving

adaptation. Notably, saccadic adaptation assumes a stable target location across

saccades, reflecting the oculomotor system's assumption of spatial stability (Burr &

Morrone, 2012; Deubel et al., 1996; Wurtz, 2008). Similarly, in reaching tasks, the manual

control system assumes that the hand should consistently and accurately reach the

intended target. In this light, implicit learning functions to recalibrate the motor system to

ensure the maintenance of stability and accuracy, aided by perceptual prediction error.

Perceptual prediction error may also underlie implicit learning in speech adaptation,

where the articulatory system adapts to auditory perturbations (Houde & Jordan, 1998;

Nasir & Ostry, 2009). Most studies in speech adaptation emphasized auditory prediction

error, a modality-specific form of sensory prediction error (Daliri & Dittman, 2019; Kim et

al., 2023a, 2023b; Kim & Max, 2021; Parrell et al., 2017, 2021). However, recent modeling

work has demonstrated that a state estimate of the task, rather than sensory prediction

error, best explains the observed learning patterns (Parrell et al., 2019; Patri et al., 2018)..

This state estimate is derived from a combination of the previous articulatory state, an

efference copy of the motor command, and auditory feedback, creating a multimodal

perceptual estimate that updates the articulatory-to-task transformation—essentially, an

internal model. Thus, even in the context of auditory perturbations, a multimodal

perceptual estimate, rather than a unimodal auditory prediction error, appears to drive

speech adaptation.

***Theoretical advance of PPE model relative to previous work***

Tsay et al. has recently proposed that implicit adaptation should be understood as a
process of proprioceptive re-alignment of the hand towards the target (PreMo model,
Tsay et al., 2022b). However, as shown in our previous work (Zhang et al., 2024), the
assumption of the PreMo model, i.e., the effect of visual perturbation on proprioceptive
feedback follows a ramp function of perturbation size, is not supported by empirical
findings (Zhang et al., 2024; Salomonczyk et al., 2011). The PreMo model also predicts
that the perceived hand direction should be unbiased in late learning (i.e., re-aligned),
contradicting to our findings in Experiment 6 which revealed a persistent perceptual error
in late learning. Most importantly, the PreMo model specifically aims to account for the
error-clamp adaptation, a special form of visuomotor rotation that invokes implicit process
only (Morehead et al., 2017). Our PPE framework, instead, aims to provide a unified
account for both explicit and implicit learning processes, which ubiquitously co-exist in
various motor learning tasks (Krakauer et al., 2019).

***Dual learning reflects the dual imperatives of motor control***
The dual learning based on the two distinct error signals could reflect the dual imperatives
of motor control: achieving action goals and resolving sensory conflicts. Researchers
have shown that unconscious movements tend to resolve multisensory conflicts, as
opposed to achieving explicit goals (Asai, 2015; Botvinick & Cohen, 1998; Lanillos et al.,
2021; Sanchez-Vives et al., 2010). For instance, in the rubber hand illusion paradigm,
participants exhibit subtle, unconscious movements to align visual and proprioceptive
cues, not to achieve a specific goal but to reduce sensory conflict (Botvinick & Cohen,
1998; Sanchez-Vives et al., 2010). This suggests that the motor system, apart from
controlling explicit actions, implicitly adjusts movements to minimize sensory mismatches
(Maselli et al., 2022; Rengarajan et al., 2022). Similarly, in the VMR learning task, sensory
feedback from different modalities (e.g., visual and proprioceptive) creates a perceptual
error relative to the predicted hand movement, leading the motor system to implicitly
recalibrate movements to resolve the conflict.  Here explicit learning corresponds to the
goal-directed imperative of motor control, driven by PE. In contrast, implicit learning aligns
with the imperative of minimizing sensory conflict, driven by PPE.

### The relationship between explicit and implicit learning

Our findings on PPE provide new insights into the relationship between explicit and implicit learning, a central question in motor learning. Previous studies have suggested that these two processes either complement each other (Miyamoto et al., 2020; Taylor et al., 2014) or compete (Albert et al., 2022; Benson et al., 2011; Flegal & Anderson, 2008; Neville & Cressman, 2018). However, our PPE model suggests that explicit and implicit learning are functionally independent, driven by distinct error signals, and their relationship is task-specific. In tasks where PE and PPE are aligned, explicit and implicit learning can work synergistically, as seen in typical adaptation tasks (Experiments 1 and 2). However, when these errors diverge, the two learning processes can conflict. For example, in Mazzoni & Krakauer (2006) and Experiment 4, when participants were instructed to re-aim explicitly, implicit learning caused a drift away from the re-aiming target, paradoxically worsening performance (Figure 6). This conflict reveals that implicit learning, driven by PPE, can sometimes undermine explicit learning, especially when their errors differ. Only when performance error grows large enough and sustains for an extended period of time do individuals adjust their explicit strategy to correct for performance error (Taylor & Ivry, 2011). Similar conflict or decoupling between explicit and implicit learning also occurs in the mirror-reversal paradigm, in which implicit learning initially worsens performance (Gastrock et al., 2023; Hadjiosif et al., 2021; Wilterson & Taylor, 2021), even after a few days of training (Kasuga et al., 2015; Telgen et al., 2014; Wilterson & Taylor, 2021). Only after sufficient exposure of this conflict, people slowly develop explicit re-aiming strategies to override implicit learning and thus improve the overall performance (Gastrock et al., 2023; Wilterson & Taylor, 2021). Interestingly, in cases of conflict, implicit learning—driven by the minimization of PPE—tends to be prioritized over performance error correction. This is reminiscent of the competition between Type I (unconscious, fast) and Type II (conscious, slow) cognitive processes in human decision-making (Kahneman, 2011).

Past research has reported that better explicit learners tend to exhibit less implicit learning, implying a competitive relationship between the two processes (Albert et al., 2022; Benson et al., 2011; Flegal & Anderson, 2008; Neville & Cressman, 2018). However, this

conclusion is often based on inter-dependent learning measures, as implicit learning is
typically measured using exclusion trials, while explicit learning is inferred by subtracting
implicit learning from total learning (Albert et al., 2022; Maresch, Werner, et al., 2021;
Neville & Cressman, 2018; Tsay et al., 2021). Given that most participants reach a
learning ceiling, this subtraction method risks producing a false negative correlation.
When explicit and implicit learning are measured independently—e.g., using aiming
reports for explicit learning and exclusion trials for implicit learning—their relationship
does not show a consistent negative correlation but varies depending on the context. In
some cases, there is even a positive correlation between the two (Figure S3; D'Amario et
al., 2024). Additionally, during early learning, when learning is not yet complete with a
relatively large PE, the negative correlation between explicit and implicit learning
disappears (Experiment 5; Figure S3), challenging the notion that these processes
compete for a shared performance error signal. In summary, our findings suggest that
explicit and implicit learning are independent processes that can interact in a task-specific
way. Their relationship is not inherently competitive but rather context-dependent, driven
by the alignment or misalignment of the underlying error signals.

***Proprioceptive bias resulted from perceptual bias related to perturbed action***
Proprioceptive bias observed in motor adaptation highlights how motor learning can
modify perception. Our findings suggest that both motoric and perceptual changes during
motor learning might follow the principles of Bayesian cue integration. In implicit motor
learning, PPE arises from a misperception of the hand's position ($\hat{x}_{hand}$). Similarly, in
proprioception tests, biased judgments result from the same biased cue ($\hat{x}_{hand}$) provided
by previous adaptation trials. Consistent with this model, our results show that both
proprioceptive bias and implicit learning are modulated by ongoing adaptation (Figures 3
& 4). The model further explains the varied effects of perturbation size on proprioceptive
bias, which can be either linear (Henriques & Cressman, 2012; Salomonczyk et al., 2011)
or constant (Modchalingam et al., 2019), an inconsistency left unexplained. A linear
dependency emerges with stepwise perturbations, where each step-up of perturbation
incrementally increases the perceived hand position $\hat{x}_{hand}$ (Salomonczyk et al., 2011;
also see Experiment 2). In contrast, a constant dependency is associated with one-step

perturbations, leading to similar $\hat{x}_{hand}$ across perturbation sizes (Modchalingam et al.,
2019; Experiment 1). The significant correlation between implicit learning and
proprioceptive bias further supports the idea that changes in perceived hand position
underlie these perceptual shifts (Experiments 1 and 2). The concept based on $\hat{x}_{hand}$
aligns with the view that proprioceptive bias results from multisensory integration (Rossi
et al., 2021; Ruttle et al., 2018, 2021; Salomonczyk et al., 2013). A few studies have
theorized that internal model formed by motor adaptation underlies proprioceptive bias('t
Hart & Henriques, 2016; Mostafa et al., 2019; Rossi et al., 2021). However, internal model
typically develops slowly, but proprioceptive bias develops rapidly with perturbation
(Ruttle et al., 2016, 2018, 2021, 2022) and diminishes rapidly without it (Zhang et al.,
2024). This makes internal model an unlikely explanation. Thus, Bayesian cue
combination provides a unified explanation for both implicit motor learning and
accompanied perceptual changes.

***Implications for neurophysiological studies***
The compositional nature of motor learning—wherein explicit and implicit learning are
driven by distinct error signals—offers a new framework for exploring their neural
underpinnings. Traditionally, neurophysiological studies have focused on the neural
correlates of total learning, implicating both sensorimotor regions (e.g., cerebellum and
sensorimotor cortex; Donchin et al., 2012; Izawa et al., 2012; Krakauer et al., 2004, 2019;
Tzvi et al., 2020, 2022) and regions associated with cognitive control and decision-making
(e.g., prefrontal cortex and basal ganglia; Anguera et al., 2007; Haar et al., 2015; Seidler
et al., 2006). By decomposing total learning into explicit and implicit components, future
research can better delineate the neural circuits underlying these distinct processes
(Griffa et al., 2024), with particular focus on the cerebellum and posterior parietal cortex
(PPC) involved in learning acquisition (Della-Maggiore & McIntosh, 2005; Galea et al.,
2011; Haar et al., 2015; Izawa et al., 2012; Tseng et al., 2007). Furthermore, most
previous studies have used either sensory prediction errors (Imamizu et al., 2000; Schlerf
et al., 2012; Tzvi et al., 2022; Wolpert et al., 1998) or performance errors (Mista et al.,
2024; Popa & Ebner, 2019; Popa et al., 2012, 2017) to identify neural correlates of motor
learning. Our finding that perceptual prediction error underlies implicit learning, with a

distinct temporal profile from other errors, suggests more tailored neural assays could be developed (Torrecillos et al., 2015). Behavioral paradigms that effectively modulate perceptual prediction error, like the stepwise perturbations in Experiment 2, could be particularly useful for future investigations. Additionally, our results emphasize the importance of localizing one's effector in motor control and learning, requiring integration of motor prediction, visual, and proprioceptive cues on a trial-by-trial basis (Figure 1A). The neural substrates for encoding and integrating these cues overlap with key networks involved in visuomotor learning, including the premotor cortex for intended cursor movement (Haar & Donchin, 2020; Hardwick et al., 2015), PPC for learning-related visual representations (Haar et al., 2015) and visuomotor transformation (Tanaka et al., 2009), the cerebellum for error processing and acquisition of internal models (Imamizu et al., 2000; Schlerf et al., 2012; Wolpert et al., 1998), and the somatosensory cortex that contributing to the acquisition and retention of motor learning (Darainy et al., 2023; Ebrahimi & Ostry, 2024; Ebrahimi et al., 2024). The interplay between motor learning and minimizing position errors of body effectors warrants further examination with neurophysiological approaches. By incorporating insights from our findings, future studies can better understand the neural mechanisms that support the distinct yet interconnected processes of explicit and implicit motor learning.

**Methods**

***Participants***

We recruited 174 college students from Peking University (101 females, 73 males, 22.76 ± 2.53 years, mean ± SD). All participants were right-handed and had normal or corrected-to-normal vision. Participants were naïve to the purpose of the experiment and received monetary compensation upon completion of the experiment.

***Apparatus***

In Experiments 1 and 2, we used the KINARM planar robotic manipulandum with a virtual-reality system (BKIN Technologies Ltd., Kingston, Canada). Participants sat in a chair and held the two robot handles with their left and right hands, respectively. The right hand

1002    performed the reaching task, and the left hand indicated the perceived direction of the

1003    right hand if required. A semi-silvered mirror positioned below eye level displayed visual

1004    stimuli and obscured the participants' view of their hands and the robotic manipulandum.

1005    The sampling rate of the KINARM was 1,000 Hz.

1006

In Experiments 3, 4, and 5, we used a horizontally placed digitizing tablet (48.8 x 30.5 cm,

Intuos 4 PTK-1240, Wacom, Saitama, Japan) to measure people's hand movements.

Participants sat in front of a vertically-placed LCD screen (29.6 x 52.7 cm, Dell, Round

Rock, TX, US) with a customized wooden blinder placed over the tablet to occlude the

vision of the hand. Using a stylus with their right hand, participants performed the reaching

task on the tablet. The sampling rate of the tablet was 200Hz. Procedures and data

acquisition were customized coded (Psychtoolbox-3-3.0.19.2 in MATLAB).

1014

The control experiment used a similar setup to Experiments 3, 4, and 5, with the addition

of eye tracking. A slightly smaller digitizing tablet (31.1 x 21.6 cm, Intuos M, Wacom,

Saitama, Japan) was used. The vertically-placed LCD screen was positioned

approximately 50cm in front of a chin rest, with a bracket occluding the participants' view

of the tablet and their hands. Movement trajectories were sampled at 200Hz by the

digitizing tablet. Eye movements were continuously tracked at 500Hz using a video-based

eye tracker (Eyelink 1000, SR Research), which was placed beneath the LCD screen.

1022

***Experiment 1***

Sixty participants were randomly assigned to one of four groups (n = 15 each). Each

group adapted to visuomotor rotation of one of four possible sizes (15°, 30°, 60°, or 90°).

Participants made reaching movements from a central start position towards an up-right

(defined as 45°) target located 10 cm away. The start position, target, and cursor were

displayed as filled circles,  with a 5-mm radius purple circle for the start position, a 5-mm

radius blue circle for the target, and a 3-mm radius white circle for the cursor. Each trial

began with participants holding their right hand at the start position for 800ms. The target

then appeared and signaled participants to make a quick, straight sliding movement

through it. The trial ended when the distance between the hand and the start position

exceeded 10 cm. If the movement duration from leaving the start position to reaching 10

cm exceeded 300 ms, a warning message, "too slow", would appear on the screen.

1035

The experiment included three phases: a 35-trial baseline phase with veridical feedback,

a 75-trial adaptation phase with VMR feedback, and a 20-trial washout phase without

cursor feedback. The direction of the VMR perturbation was counterbalanced across

participants. To assess implicit learning at the asymptote of learning, we inserted

exclusion trials at the 50, 55, 60, and 65th trials during the adaptation phase. In these

exclusion trials, participants were instructed to move their hand straight to the target

without any strategic re-aiming. The washout phase used the same exclusion instruction,

which the experimenter emphasized during data collection.

1044

To measure proprioceptive bias, we inserted proprioception test trials at the 15, 20, 25,

30, and 35th trials during the adaptation phase (Figure 3A). Additional proprioception test

trials were conducted at the 5, 10, 15, and 20th trials during the baseline phase to assess

baseline proprioceptive bias. In the proprioception test trial, participants were instructed

to relax their right arm, which was then passively moved to one of four locations (± 5° and

± 15° away from the reaching target). The passive movement lasted for 1000 ms and

followed a straight path with a minimum-jerk speed profile. During this movement, a white

ring, centered at the start position, expanded in sync with the hand's center-out

displacement to indicate the movement distance. After the right hand reached the target

location, participants held their right hand steady and used the left robot handle to indicate

the perceived position of their right hand. Moving the left handle controlled the rotation of

a visual "dial" displayed along an arc that passed through the proprioceptive targets

(Zhang et al., 2024). Participants held the dial at their perceived location for 1,000 ms,

which was recorded as their perceptual judgement.

1059

1060    ***Experiment 2***

1061    Experiment 2 used the same task setup as in Experiment 1. We recruited two groups of

1062    participants: one group experienced stepwise perturbations (n = 15), while the other group

1063    experienced a one-step perturbation (n = 15). The stepwise group experienced a stepwise

15°→30°→45°→60° VMR perturbation, with each perturbation presented for 40 trials. The one-step group experienced a 60° perturbation for 160 consecutive trials (Figure 4A). Within each 40-trial perturbation block, we inserted exclusion trials at the 30 and 35th trials to measure implicit learning, and proprioception test trials at the 5, 10, 15, and 20th trials to measure proprioceptive bias. Before the adaptation phase, all participants completed a 35-trial baseline phase with veridical feedback. After the adaptation phase, they went through a washout phase without visual feedback, following the same exclusion instruction.

### Experiment 3

Experiment 3 investigated the size-dependency of implicit learning when explicit learning was suppressed by the randomness of perturbations. To prevent cumulative learning over trials, three targets (25°, 45°, and 65° at an 8 cm distance) were used. Participants (n = 18) began with a baseline phase consisting of 15 veridical-feedback trials and 15 interleaved no-feedback null trials. In subsequent adaptation phase, trials were organized into 6-trial mini blocks, where each target was presented once randomly and each perturbation trial was flanked by two no-feedback null trials (Figure 5A). During perturbation trials, participants were instructed to control the cursor to hit the target, while null trials were served as exclusion trials. Visuomotor rotation of 0°, 4°, 8°, 16°, 32°, and 64°, either clockwise or counter-clockwise was randomly used in each perturbation trial. These 11 perturbations were repeated nine times, leading to a total of 99 perturbation trials and 99 null trials in the adaptation phase. The wide range of rotation sizes enabled us to uncover the linear or nonlinear size-dependency of implicit adaptation (Hutter & Taylor, 2018; Wei & Kording, 2009). The randomness of perturbation size and direction would prevent participants forming an adaptive strategy. Single-trial learning was quantified as the difference in movement direction between the two adjacent null trials before and after a perturbation trial.

### Experiment 4

Experiment 4 examined VMR adaptation with instructed re-aiming, as in the seminar study by *Mazzoni & Krakauer, 2006*. However, two different sizes of VMR were used for

two separate groups of participants, i.e., 45° (n = 15) or 90° VMR (n = 15). The target, presented as a 5-mm radius white circle, would appear at one of eight possible locations (0°, 45°, 90°, 135°, 180°, 225°, 270°, and 315°) 8 cm away, randomly in an eight-trial cycle. In the baseline phase, participants were instructed to control the cursor to hit the target with veridical feedback. In the adaptation phase, a new re-aiming target was introduced starting from the first perturbation trial. The re-aiming target, which was either 45° or 90° but in the opposite rotation direction of the imposed VMR, was presented as a 5-mm radius green circle, distinct from the original target. For a consecutive of 120 trials, participants were required to move their hand to the re-aiming target to make the cursor "hit" the original target. Half of the participants in each group experienced clockwise perturbations and the other half counterclockwise perturbations. Typically, people could maintain their aim at the re-aiming target despite their hand implicitly drifted away from it, across the span of 120 trials (Taylor & Ivry, 2011).

### *Experiment 5*

Experiment 5 examined the instructed re-aiming introduced at the end of VMR adaptation, as opposed to at its beginning (as in Experiment 4). Four targets were used, located at 45°, 135°, 225°, and 315°, 8 cm away, with trials organized into 4-trial cycles. The baseline phase consisted of 3 trial cycles with veridical feedback trial and 1 trial cycle without feedback. The subsequent adaptation phase consisted of 60 cycles with a 45° VMR. Within the adaptation phase, we intermittently measured the explicit adaptation through aiming reports at the 5, 19, 33, and 47th cycles. In these trials, the target was surrounded by a ring of 63 numbered visual landmarks spaced 5.625° apart, with 0° centered on the target. Before the reach, participants verbally reported the landmark number they planned to reach. Their reported aiming directions were recorded by the experimenter. To measure implicit adaptation, we inserted exclusion trials with no visual feedback at the 6, 20, 34, 48th cycles. In the ensuing re-aiming phase, we presented a re-aiming target at the last reported aiming direction, tailored to each individual participant. Participants were required to move their hand straight to this re-aiming targe for 40 cycles. Note that the re-aiming target would still make the cursor reached the original target since it was aligned with the participant's spontaneous explicit learning asymptote. We inserted

1126    exclusion trials at the 1st, 11, and 21th cycles in the re-aiming phase. In the final washout

1127    phase, participants completed an additional 15 cycles of exclusion trials.

1128

***Experiment 6***

Experiment 6 aimed to directly measure PPE, defined as the difference between the

intended movement direction (aiming direction) and the perceived hand direction. Using

similar setup and procedures as Experiment 1 and 2, Experiment 6 recruited thirty

participants and randomly assigned them to learn one of three VMR perturbations (15°,

30°, and 90° VMR; n = 10 each). Participants went through three learning phases, i.e.,

baseline (20 trials), adaptation (44 trials), and washout (20 trials).  Every four trials they

performed a probe trial for measuring PPE (Figure 8A). The first eight adaptation trials

were not probed to avoid affecting participants' formation of strategic re-aiming during this

critical initial learning phase (McDougle & Taylor, 2019; Ruttle et al., 2021). For each

probe trial, participants held their right hand at the start position for 800 ms, and then a

target (a 5-mm radius blue circle), a white arc, and a purple visual marker were displayed.

Participants were asked to report their intended aiming direction ("Where will your right

hand go?") by moving the left KINARM handle to dial the visual marker, just as in

Experiment 2. Participants subsequently performed the reaching movement, with or

without the cursor perturbation, by their right hand. After the reach, their right hand was

passively returned to the starting position by the robotic manipulandum. The arc and

visual marker reappeared to prompt participants to report the perceived direction of their

actual hand movement ("Where was your right hand?"), again using the left KINARM

handle to dial the visual marker. PPE was computed as the angular difference between

these two reported directions.

1150

***Control experiment with eye-tracking***

The control experiment examined whether people fixate in the re-aiming direction during

reaching movements in typical VMR adaptation. Participants (n = 16) learned 45° and 90°

VMR perturbations in two consecutive sessions, with the order counterbalanced. Each

session included a baseline, adaptation, and washout phase, as in Experiment 1, with 40,

160, and 20 trials, respectively. Each trial began when the participant moved the cursor

(a 3-mm radius white circle) into the starting position (a 5-mm radius purple circle) using

a stylus. After holding the position for 500ms, a target (a 5-mm blue circle) was presented

10 cm away at a random direction between 0° and 360°. Participants started to reach the

target with the cursor once the target disappeared. The target presentation duration

followed a uniform distribution from 800 to 1200 ms, making the movement trigger less

predictable. If the movement was initiated before the target disappearance, the trial was

aborted and a text message "reacting too fast" would appear on the screen. Veridical or

perturbed cursor feedback was shown during the movement and the target re-appeared

once the movement ended and lasted for 1000 ms. The disappearance of the target

prevented some participants keeping their fixation at the starting position without change

to their re-aiming direction. To avoid slow movements, a text message "moving too slow"

would appear if the movement time exceeded 400ms.

1169

***The three error models***

*Perceptual prediction error (PPE) model:* The PPE model hinges on the perceptual

estimation of effector position that is dynamically updated through Bayesian cue

combination. For reaching, the perceptual estimate of the hand movement direction

$(\hat{x}_{hand})$ results from the integration of three sensory cues: the visual cue $(x_v)$ from the

cursor, the proprioceptive cue $(x_p)$ from the hand, and the sensory prediction of the

reaching action $(x_u)$. Following Bayesian principles (Ernst & Banks, 2002), the perceptual

estimate is

1178
$$\hat{x}_{hand} = \sum_i W_i x_i, \text{ with } W_i = \frac{1/\sigma_i^2}{\sum_j 1/\sigma_j^2}, \; i,j = u, p, v \tag{1}$$

where the weight of each cue is determined by their uncertainty $(\sigma)$ relative to each other.

Notably, in the VMR paradigm, the predicted reaching direction is the aiming direction.

The rotated cursor $(x_v)$ biases the hand estimate $(\hat{x}_{hand})$ toward the cursor's direction.

The deviation from the predicted (re-aiming) direction constitutes the perceptual

prediction error, which drives implicit adaptation on the subsequent trial (Eq. 2). We

assume that learning proceeds on a trial-by-trial basis thus a state-space framework is

used. Explicit learning is driven by performance error $e_t^{PE}$, as in existing models (Eq. 3;

Albert et al., 2022; Cheng & Sabes, 2006; Herzfeld et al., 2014; McDougle et al., 2015, 2017). For each trial, the learning is updated by

$$x_{t+1}^i = A^i x_t^i - B^i \left( \hat{x}_{Hand,t} - x_{u,t} \right) \tag{2}$$

$$x_{t+1}^e = A^e x_t^e - B^e e_t^{PE} \tag{3}$$

$$x_{t+1} = x_{t+1}^e + x_{t+1}^i \tag{4}$$

Where total learning ($x$) is the sum of explicit learning ($x^e$) and implicit learning ($x^i$). *A* is the retention rate capturing the inter-trial forgetting and *B* is the learning rate capturing the proportion of error correction.

As individuals fixate near the re-aiming direction during the movement (Bromberg et al., 2019; de Brouwer et al., 2018), the visual cursor directionally differs from the fixation by an amount of $(x_v - x_u)$. This eccentricity linearly increases visual uncertainty about cursor motion direction, as shown both in visual perception studies and in our previous work (Zhang et al., 2024). Thus, the visual uncertainty ($\sigma_v$) follows a linear function:

$$\sigma_v = k * |x_v - x_u| + b \tag{5}$$

where *k* and *b* are the slope and intercept of the visual uncertainty, respectively.

*Performance error (PE) model:* The performance error (PE) model, also called the competition model (Albert et al., 2022) or the two-state model (Smith et al., 2006), has been widely used to explain motor adaptation (Albert et al., 2022; Hadjiosif et al., 2021; McDougle et al., 2015, 2017; Miyamoto et al., 2020; Smith et al., 2006). The Eq. 2 in the PPE model is replaced by

$$x_{t+1}^i = A^i x_t^i - B^i e_t^{PE} \tag{6}$$

*Sensory-prediction error (SPE) model:* Implicit motor learning has been suggested to be driven by sensory prediction error (SPE; Kawato, 1999; Mazzoni & Krakauer, 2006; Tseng et al., 2007). In the VMR paradigm, SPE is defined as the deviation of the perturbing

1213 cursor from the aiming direction (Morehead & Xivry, 2021). Therefore, the SPE model

1214 replaces Eq.2 in the PPE model by

$$x_{t+1}^i = A^i x_t^i - B^i (r - x_t^i),\tag{7}$$

1216 where $r - x_t^i$ is SPE and $r$ is the VMR size.

1217
1218

### Data analysis

*Hand kinematics:* In Experiments 1 and 2, hand positions and velocities were directly

acquired from the KINARM robot at a fixed sampling rate of 1,000 Hz. The raw kinematic

data were smoothed using a fifth-order Savitzky-Golay filter with a window length of 50

ms. In Experiments 3, 4, and 5, hand positions were recorded at a sampling rate of 125

Hz. The movement direction was determined by computing the direction of the vector

spanning from the starting position to the hand position at the peak outward velocity. We

defined the direction to counter the perturbation as the positive direction.

1227

*Gaze:* The gaze position obtained in the control experiment was first preprocessed to

remove blinks. Consistent with previous studies (e.g., Bromberg et al., 2019), we used

the last fixation before movement onset to quantify the participants' aiming direction.

Trials were excluded if the fixation before movement onset was missing (due to a blink),

or near the start position (< 40% of target distance), or beyond the target (>150% of target

distance). These exclusions amounted to 1.67% of the trials in the 45°-VMR condition

and 3.61% in the 90°-VMR condition. Other direction measures, such as the gaze position

at movement onset, yielded similar results.

1236

*Statistical analysis:* For the dataset from *Bond & Taylor 2015,* we quantified early and late

learning by using the data from the first 5 cycles and the last 5 cycles in the adaptation

phase, respectively. Total learning was defined as the hand's directional deviation from

the target. Explicit learning was quantified as the direction deviation of the aiming report

from the target. Implicit learning was calculated by subtracting explicit learning from total

learning. Group differences were examined using one-way independent-measure

ANOVAs. In Experiments 1 and 2, implicit learning was quantified as the average hand

deviation from the target in the four exclusion trials. Total learning was calculated as the

average hand deviation from the target in the last cycle during the adaptation phase.

Explicit learning was calculated by subtracting implicit learning from total learning.

Proprioceptive bias was quantified as the average angular difference between the

perceived (reported) and actual hand directions over the six proprioception test trials.

Group differences were examined using one-way independent-measure ANOVAs, with

pairwise post-hoc comparisons conducted using Bonferroni correction. For Experiment 2,

two-way mixed ANOVAs were performed, with group as the between-group factor and

learning time as the within-group factor. For the correlation analysis of Experiment 2, we

used a mixed linear model: $y = \beta_0 + \beta_1 * x_{prop} + \beta_2 * x_{sub|prop} + \epsilon$, where y is implicit

learning, $x_{prop}$ is proprioceptive bias, $x_{sub|prop}$ is the random factor of subject, and $\epsilon$ is

the residual error. In Experiment 3, single-trial learning was defined as the angular

difference in movement direction between two consecutive no-feedback null trials before

and after a perturbation trial. Given the symmetry between CW and CCW perturbations,

we pool the data to quantify the size dependency on the absolute perturbation size.

Repeated-measures ANOVAs were used to examine the effect of perturbation size on

single-trial learning. In Experiment 4, independent t-tests were used to compare implicit

learning between 45°-VMR and 90°-VMR conditions. In Experiment 5, we used paired t-

tests to compare implicit learning before and after introducing the re-aiming target. Implicit

learning was quantified by both subtraction and exclusion methods. In Experiment 6, we

used one-sample t-tests to examined whether PPE was significantly negative, and used

one-way ANOVAs to examine group differences in PPE and implicit learning. Post-hoc

comparisons were conducted with Bonferroni correction. In the eye-tracking experiment,

we used paired t tests condition differences (90° vs. 45°) in re-aiming direction. We used

a mixed linear model for analyzing the correlation between explicit learning and eye-

fixation direction: $y = \beta_0 + \beta_1 * x_{aim} + \beta_2 * x_{condition} + \beta_3 * x_{sub|(aim*condition)} + \epsilon$, where

$y$ is explicit learning, $x_{aim}$ is eye-fixation direction, $x_{sub|(aim*condition)}$ is the random factor

of subject, and $\epsilon$ is the residual error. For all experiments, we excluded the trials with

movement time exceeding 400 ms or below 150 ms, as well as the trials with systematic

movements to the incorrect direction (i.e., movement direction less than -20° or exceeded

120°). These criteria resulted in the removal of less than 1% of trials on average for each

experiment.

1276

1277    ***Model analysis***

1278    Both PE and SPE models have four parameters $\Theta = [A^e, B^e, A^i, B^i]$. The PPE model has

1279    an additional four parameters $\Theta = [\sigma_u, \sigma_p, k, b]$. Not all the experiments need the full

1280    models (as detailed below).

1281

*Bond & Taylor, 2015.* We fitted the trial-by-trial average learning data from all the four

perturbation conditions simultaneously (Figure 2C). No fitting was needed for explicit

learning since it was measured throughout. The parameters related to sensory

uncertainty ($\sigma_u, \sigma_p, k, b$) had been estimated from our previous work with visuomotor

rotation perturbations (Zhang et al., 2024). Three of them were fixed, including $b$ = 1.853°,

$\sigma_u$ = 5.05°, $\sigma_p$ = 11.12°. Note, the parameter $k$, the linear slope of visual uncertainty

relative to perturbation size, was left as a free parameter. This is because the previous

work used the error-clamp paradigm, which reinforced fixation on the reach target rather

than on the cursor. In contrast, participants in the VMR paradigm here, look at the cursor

at the end of the reaching movement, which presumably reduces the effect of eccentricity

on visual uncertainty. Therefore, we left $k$ as a free parameter for all the experiments

examined here. Hence, the PPE model had three free parameters $\Theta = [k, A^i, B^i]$, while

the PE and SPE models each had two free parameters $\Theta = [A^i, B^i]$. Bootstrapping was

used to estimate the standard deviations of the model fits. Participants in each group were

resampled with replacement 5000 times, each time producing a new average learning

sequence.

1298

*Experiment 1 and 2.* The trial-by-trial learning data include total learning and implicit

learning (intermittently measured during the adaptation phase) from all the four conditions

(Experiment 1; Figure 3C) or the two conditions (Experiment 2; Figure 4D). The PPE

model had five free parameters $\Theta = [k, A^i, B^i, A^e, B^e]$. Since the one-step condition in

Experiment 2 was identical to the conditions in Experiment 1 (only differing in the duration

of the adaptation phase), we applied the $k$ value estimated from Experiment 1 to the one-

step condition in Experiment 2. The stepwise condition, however, retained $k$ as a free

parameter ($k_{step}$ in Table S3) since the participants would be attracted by the sudden

stepwise change in cursor rotation, which would reduce visual uncertainty. Through fitting

the data from both two conditions (Figure 4D), we obtained the value of $k_{step}$. We used

the values of $k$ and $k_{step}$ for cross-validation (Figure S2B). The SPE and PE models had

four free parameters $\Theta = [A^e, B^e, A^i, B^i]$.

1311

*Experiment 3, 4 and 5.* We first assumed that the single-trial learning in Experiment 3 was

implicit. Thus, the PPE model had three free parameters $\Theta = [k, A^i, B^i]$ to fit the average

trial-by-trial changes (Figure 5B). The averaging was feasible since the pseudo-random

perturbation sequence was identical for each participant. For fitting, we set the lower limit

of $A^i$ at 0.7 for all the models. Based on fitting the trial-by-trial learning, we calculated the

model-predicted single-trial learning as a function of perturbation size (Figure 5D). To

examine whether there was explicit learning at work, we then included the explicit learning

component and evaluated whether it improved model fitting. In Experiment 4, there was

only implicit learning since participants were unaware of their hand drifting away from the

instructed re-aiming target. Thus, only implicit learning was fitted for the 45°- and 90°-

VMR conditions at once. The PPE model had three free parameters $\Theta = [k, A^i, B^i]$, while

the SPE and PE models had two free parameters $\Theta = [A^i, B^i]$. For Experiment 5, the data

used for model fitting included the trial-by-trial changes in total learning as well as the

intermittently-measured explicit learning. Models were fitted to the data before introducing

the re-aiming target and then used to predict the learning afterwards (Figure 7D). The

PPE model thus had five free parameters $\Theta = [k, A^i, B^i, A^e, B^e]$, while the other two models

had four free parameters $\Theta = [A^i, B^i, A^e, B^e]$.

1329

All model fittings were conducted using *fmincon* function in MATLAB (2022b, MathWorks,

Natick, MA, US). To ensure the model adhered to established assumptions, we

constrained the parameters with $A^e < A^i$ and $B^e > B^i$ as commonly assumed for explicit

and implicit learning components (McDougle et al., 2015, 2017; Smith et al., 2006). All

the obtained parameter values, along with quality measures of the model fit, including $R^2$,

root mean squared error (*RMSE*), Akaike information criterion (AIC), and Bayesian
information criterion (BIC), were provided in Table S1 to S6.

1337

1338 **Data availability**

1339 Data and code are available at: https://osf.io/tjc52/

1340

**Acknowledgements:** We would like to thank Dr. Jordan Taylor for providing the raw data
from his study (Bond & Taylor, 2015). **Funding**: This work was supported by STI2030-
Major Projects (2021ZD0202600) and the National Natural Science Foundation of China
(32071047, 32471099, 6206113600, 31871102), awarded to KW. **Authors**
**contributions:** Conceptualization: KW, XZ, Data curation: XZ, WW; Formal analysis: XZ;
Funding acquisition: KW; Methodology: KW, XZ, WW; Software: XZ, WW; Supervision:
KW; Visualization: XZ; Writing: KW, XZ.

1348

**Supplementary materials**

 **Figure S1**. Control experiment examining eye fixation during visuomotor rotation

 adaptation. **(A)** Illustration of the experimental procedure. In each trial, participants first

 held their hand at the start position for 500 ms, awaiting the appearance of a target

 randomly positioned between 0° and 360°. They were instructed not to move until the

target disappeared (~1000 ms after appearance). Visual feedback of the hand cursor was withheld until the trial ended, when it re-appeared together with the target as performance feedback. Eye tracking measurements were taken during reaching. **(B)** Time course of the gaze direction. Different colors represent different timings of a trial, including target preview, reaction time (RT), movement duration, and terminal feedback. Two representative trials from a participant are shown for the 45°-VMR (upper panel) and 90°-VMR (lower panel) conditions, respectively. The participant fixated on the starting position during the target preview and RT, then shifted fixation toward the re-aiming direction during the movement, maintaining this fixation until the terminal feedback was provided (purple line). The re-aiming direction was less rotated than the imposed VMR. After the movement ended and visual feedback was provided, participants tended to fixate back to the cursor (yellow line). We calculated the probability of fixations in the target area (± 22.5° around the target direction) after movement: 64.28 ± 4.91% in the 45°-VMR condition and 43.23 ± 6.80% in the 90°-VMR condition. **(C)** Average hand movement and eye fixation directions during adaptation to the 45° and 90° VMR. The fixation direction was defined as the last fixation before movement onset (Bromberg et al., 2019). The shaded areas denote the standard errors across participants. From early learning, participants started to fixated away from the target direction, closer to their explicit re-aiming direction, and this fixation "strategy" persisted throughout the adaptation phase. Note that the fixation deviated further from the target in the 90°-VMR condition compared to the 45°-VMR condition, consistent with the condition difference in re-aiming and explicit learning. In the washout phase, their fixation returned to the target as instructed. **(D)** Participants' fixation direction in the 90°-VMR condition was significantly higher than that in the 45°-VMR condition, during both early (45°: 19.00°± 2.23°; 90°: 44.49°± 4.43°; $t_{(15)}$ = -6.38, $p$ < .0001) and late learning (45°: 21.63°± 2.56°; 90°: 57.88°± 4.18°; $t_{(15)}$ = -9.21, $p$ < .0001). The larger the fixation direction, the greater the re-aiming deviates away from the target. Early and late learning refer to the first 5 and last 5 trial cycles during adaptation. **(E)** Participants' eye-fixation direction was positively correlated with their explicit learning ($\beta$ = 0.80, $p$ = 0.014), the latter estimated from the first exclusion trial in the washout phase.

1387

**Figure S2**. Model comparison results from Experiment 2. **(A)** Implicit learning data in Experiment 2, a reproduction of Figure 4C right panel. **(B)** Model fitting with varying explicit parameters across the two conditions. Compared to the model fitting in the main text (Figure 4D), here we allowed explicit parameters to vary across the two conditions. The PE and SPE models still failed to reproduce the data, especially the relative size of implicit learning between the two perturbation conditions. The PPE model remained capable of explaining the data well with variable explicit learning parameters. These results suggest that the failure of PE and SPE to explain implicit learning across conditions was not due to an unconsidered potential parametric change in explicit learning. **(C)** Cross-validation tests for the three models. Each model was fit to the data from the stepwise condition and then used to predict implicit learning in the one-step condition. Neither PE or SPE model captures the learning pattern, with mismatches similar to their original model fitting in Figure 4D. In contrast, the PPE model's predictions match well with the empirical data across conditions. In **(B)** and **(C)**, error bars represent bootstrapped standard deviations (resample size = 5000).

1403

1404

**Figure S3**. Correlation between explicit and implicit learning varies widely when the two learning components are independently measured. Data from three separate studies are presented, with explicit learning measured by aiming reports and implicit learning

measured by no-feedback exclusion trials. **(A)** Correlations between explicit and implicit learning in different perturbation size conditions, including 15°, 30°, 60°, and 90° VMR (n = 10 for each). Due to the small sample size and variability across condition, support for a negative correlation was insufficient. Data from *Bond & Taylor, 2015*. **(B)** Correlations across perturbation size conditions of 15°, 30°, 45°, and 60° VMR (n = 33, 54, 28, 61, respectively) from *D'Amario et al., 2024*. With decent sample sizes, positive, negative or null correlation was observed in these different conditions. **(C)** Correlations were inconsistent when the format of visual perturbation varied but with a fixed perturbation size of 45° VMR. Data from *D'Amario et al., 2024*. Continuous cursor feedback (n = 51) is the conventional feedback format, terminal feedback promotes explicit learning and reduce implicit learning (n = 35), delayed feedback reduces implicit learning (n = 39), and cursor jump feedback usually promotes explicit learning and reduces implicit learning (n = 32). Only the continuous-feedback condition showed a negative correlation, no correlation was detected for the other three feedback formats. **(D)** Early learning did not show a negative correlation. In our Experiment 5 (n = 20), explicit learning and implicit learning were independently measured four times during the 60 trial-cycle learning period. The two early instances (the 5th and 19th cycles) did not show significant negative correlations, the third instance (the 33th cycle) showed a marginal effect, and the fourth instance (the 47th cycle) showed a significant negative correlation. Thus, though performance error is most prominent during early learning, there is no evidence that the two learning processes compete for this error and then lead to a negative correlation. Each dot represents an individual's data.

1440

**Figure S4.** The performance of the PPE model was compared by assuming a linear and an exponential visual uncertainty function. The observed and model-predicted implicit learning for Bond & Taylor 2015 (A), Experiment 1 (B), and Experiment 2 (C). The two versions of the PPE model similarly explain the data. (D) The visual uncertainty functions are estimated from the three datasets. Note the estimates are similar for the linear and the exponential functions, and they are also close to the visual uncertainty measured in our previous work (Zhang et al., 2024).

1448

1449

1450

1451

1452

1453

1454

1455

1456

1457

1458 **Table S1**. Model comparisons and parameters for the dataset from *Bond & Taylor, 2015*

|  | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $k$ |
|---|---|---|---|---|---|---|---|
| **PPE Model** | **0.708** | **1.684** | **172.81** | **182.04** | 0.158 | 0.931 | 0.209 |
| PE Model | 0.177 | 2.188 | 254.54 | 260.69 | 0.072 | 0.999 | / |
| SPE Model | -0.730 | 5.054 | 522.48 | 528.63 | 0.015 | 0.916 | / |

1459

1460 **Table S2**. Model comparisons and parameters for the data from Experiment 1

|  | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $B_e$ | $A_e$ | $k$ |
|---|---|---|---|---|---|---|---|---|---|
| **PPE Model** | **0.969** | **4.807** | **361.67** | **375.27** | 0.151 | 0.981 | 0.513 | 0.981 | 0.204 |
| PE Model | 0.957 | 5.870 | 404.43 | 415.31 | 0.069 | 0.991 | 0.450 | 0.991 | / |
| SPE Model | 0.954 | 6.044 | 411.00 | 421.87 | 0.014 | 0.979 | 0.505 | 0.979 | / |

1461

1462 **Table S3**. Model comparisons and parameters for the data from Experiment 2

1463 **Table S3a.** Fitting the one-step and stepwise groups together with a single set of

1464 parameters

|  | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $B_e$ | $A_e$ | $k$ | $k_{step}$ |
|---|---|---|---|---|---|---|---|---|---|---|
| **PPE Model** | **0.960** | **3.965** | **252.43** | **264.81** | 0.245 | 0.927 | 0.889 | 0.927 | 0.204 | 0.139 |
| PE Model | 0.933 | 5.245 | 299.68 | 309.59 | 0.263 | 0.958 | 0.647 | 0.958 | / | / |
| SPE Model | 0.925 | 5.491 | 307.76 | 317.67 | 0.046 | 0.906 | 0.908 | 0.906 | / | / |

1465

1466 **Table S3b.** Assuming explicit learning changes across conditions, fitting the one-step and

1467 stepwise groups with separate explicit learning parameters

|  |  |  |  |  |  |  | Stepwise |  |  | One-step |  |  |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
|  | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $B_e$ | $A_e$ | $k_{step}$ | $B_e$ | $A_e$ | $k$ |
| **PPE Model** | **0.962** | **3.854** | **251.42** | **268.76** | 0.237 | 0.930 | 0.636 | 0.924 | 0.136 | 0.940 | 0.930 | 0.204 |
| PE Model | 0.958 | 4.234 | 265.97 | 280.84 | 0.223 | 0.970 | 0.218 | 0.990 | / | 0.792 | 0.970 | / |
| SPE Model | 0.926 | 5.453 | 310.53 | 325.39 | 0.047 | 0.902 | 0.672 | 0.945 | / | 0.939 | 0.902 | / |

1468

1469 **Table S3c.** Cross-validation: first fit the model by using the data from the stepwise group

1470 only, and then predict the data for the one-step group

|  | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $B_e$ | $A_e$ | $k$ | $k_{step}$ |
|---|---|---|---|---|---|---|---|---|---|---|
| **PPE Model** | **0.913** | **5.868** | **165.72** | **174.64** | 0.304 | 0.911 | 0.660 | 0.911 | 0.204 | 0.139 |
| PE Model | 0.661 | 9.031 | 201.66 | 208.80 | 0.382 | 0.946 | 0.454 | 0.946 | / | / |
| SPE Model | 0.583 | 9.638 | 207.38 | 214.52 | 0.132 | 0.859 | 0.774 | 0.859 | / | / |

1471

1472  **Table S4**. Model comparisons and parameters for the data from Experiment 3

1473  **Table S4a.** For the trial-by-trial learning sequence (Figure 5B)

| | | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $B_e$ | $A_e$ | $k$ |
|---|---|---|---|---|---|---|---|---|---|---|
| **PPE Model** | **Imp** | **0.471** | **1.294** | **39.988** | **46.557** | 0.418 | 0.700 | / | / | 0.193 |
| | Imp+Exp | 0.453 | 1.317 | 46.307 | 57.256 | 0.424 | 0.700 | 0.010 | 0.500 | 0.201 |
| PE Model | Imp | -0.682 | 1.747 | 77.622 | 82.001 | 0.054 | 0.700 | / | / | / |
| | Imp+Exp | -0.682 | 1.747 | 81.622 | 90.381 | 0.054 | 0.700 | 0.496 | 0.542 | / |
| SPE Model | Imp | -0.682 | 1.747 | 77.622 | 82.001 | 0.054 | 0.700 | / | / | / |
| | Imp+Exp | -0.698 | 1.750 | 81.848 | 90.607 | 0.054 | 0.700 | 0.010 | 0.500 | / |

1474

1475  **Table S4b.** For the single-trial learning function (Figure 5C)

| | $R^2$ | $RMSE (°)$ |
|---|---|---|
| **PPE Model** | **0.954** | **0.684** |
| PE/SPE Model | -0.321 | 2.449 |

1476

1477

1478  **Table S5**. Model comparisons and parameters for the data from Experiment 4

| | $R^2$ | $RMSE (°)$ | AIC | BIC | $B_i$ | $A_i$ | $k$ |
|---|---|---|---|---|---|---|---|
| **PPE Model** | **0.972** | **0.524** | **-32.81** | **-28.61** | 0.396 | 0.840 | 0.211 |
| PE Model | -0.866 | 4.496 | 94.19 | 97.00 | 0.034 | 0.796 | / |
| SPE Model | -1.150 | 4.457 | 93.67 | 96.48 | 0.025 | 0.833 | / |

1479

1480

1481  **Table S6**. Model comparisons and parameters for the data from Experiment 5: Cross-
1482  validation (Fitting before-re-aiming phase, predicting after-re-aiming phase).

| | $R^2$ | $RMSE (°)$ | AIC | BIC | $B_i$ | $A_i$ | $B_e$ | $A_e$ | $k$ |
|---|---|---|---|---|---|---|---|---|---|
| **PPE Model** | **0.880** | **5.641** | **155.33** | **164.01** | 0.082 | 0.966 | 0.304 | 0.966 | 0.201 |
| PE Model | 0.337 | 13.825 | 228.62 | 235.57 | 0.082 | 0.966 | 0.304 | 0.966 | / |
| SPE Model | 0.859 | 5.843 | 156.59 | 163.54 | 0.016 | 0.954 | 0.359 | 0.954 | / |

1483

1484

1485

1486

1487

1488

1489 **Table S7:** Model comparisons of the PPE models, which assume linear, exponential, or

1490 power-law visual uncertainty function.

1491 PPE$^{exp}$ model: assuming $\sigma_v = k'e^{\alpha\theta} + (b - k')$

1492 PPE$^{power}$ model: assuming $\sigma_v = k'\theta^\alpha + b$

1493

1494 **Table S7a.** Model comparisons for *Bond & Taylor, 2015. Note that the parameters of*

1495 *power and exponential functions indicate that these functions are close to linear ones.*

| | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $k'$ | $\alpha$ |
|---|---|---|---|---|---|---|---|---|
| PPE Model | 0.708 | 1.684 | 172.81 | 182.04 | 0.158 | 0.931 | / | / |
| PPE$^{power}$ Model | 0.978 | 1.173 | 59.05 | 71.35 | 0.531 | 0.977 | 1.49 | 0.674 |
| PPE$^{exp}$ Model | 0.680 | 1.777 | 191.93 | 204.24 | 0.143 | 0.930 | 99.93 | 0.002 |
| PE Model | 0.177 | 2.188 | 254.54 | 260.69 | 0.072 | 0.999 | / | / |
| SPE Model | -0.730 | 5.054 | 522.48 | 528.63 | 0.015 | 0.916 | / | / |

1496

1497 **Table S7b.** Model comparisons for Experiment 1. *Note the parameters of power and*

1498 *exponential functions indicate these functions are close to a linear one.*

| | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $B_e$ | $A_e$ | $k'$ | $\alpha$ |
|---|---|---|---|---|---|---|---|---|---|---|
| PPE Model | 0.969 | 4.807 | 359.67 | 370.55 | 0.151 | 0.981 | 0.513 | 0.981 | / | / |
| PPE$^{power}$ Model | 0.971 | 4.612 | 350.44 | 361.31 | 0.505 | 0.982 | 0.505 | 0.982 | 1.28 | 0.681 |
| PPE$^{exp}$ Model | 0.968 | 4.832 | 360.84 | 371.71 | 0.014 | 0.979 | 0.505 | 0.979 | 99.99 | 0.002 |
| PE Model | 0.957 | 5.870 | 404.43 | 415.31 | 0.069 | 0.991 | 0.450 | 0.991 | / | / |
| SPE Model | 0.954 | 6.044 | 411.00 | 421.87 | 0.014 | 0.979 | 0.505 | 0.979 | / | / |

1499

1500 **Table S7c.** Model comparisons for Experiment 2. *Note the parameters of power and*

1501 *exponential functions indicate these functions are close to a linear one.*

| | $R^2$ | $RMSE(°)$ | AIC | BIC | $B_i$ | $A_i$ | $B_e$ | $A_e$ | $k'$ | $\alpha$ |
|---|---|---|---|---|---|---|---|---|---|---|
| PPE Model | 0.960 | 3.965 | 252.43 | 264.81 | 0.245 | 0.927 | 0.889 | 0.927 | / | / |
| PPE$^{power}$ Model | 0.962 | 3.897 | 253.39 | 270.73 | 0.409 | 0.941 | 0.870 | 0.941 | 0.43 | 0.889 |
| PPE$^{exp}$ Model | 0.962 | 3.900 | 253.55 | 270.89 | 0.360 | 0.940 | 0.874 | 0.940 | 99.73 | 0.003 |
| PE Model | 0.933 | 5.245 | 299.68 | 309.59 | 0.263 | 0.958 | 0.647 | 0.958 | / | / |
| SPE Model | 0.925 | 5.491 | 307.76 | 317.67 | 0.046 | 0.906 | 0.908 | 0.906 | / | / |

1502

1503

1504

1505

## Reference

Al-Fawakhiri, N., Ma, A., Taylor, J. A., & Kim, O. A. (2023). Exploring the role of task success in implicit motor adaptation. *Journal of neurophysiology, 130*(2), 332-344.

Albert, S. T., Jang, J., Modchalingam, S., Marius't Hart, B., Henriques, D., Lerner, G., Della-Maggiore, V., Haith, A. M., Krakauer, J. W., & Shadmehr, R. (2022). Competition between parallel sensorimotor learning systems. *Elife, 11*, e65361.

Anguera, J. A., Russell, C. A., Noll, D. C., & Seidler, R. D. (2007). Neural correlates associated with intermanual transfer of sensorimotor adaptation. *Brain Research, 1185*, 136-151.

Asai, T. (2015). Illusory body-ownership entails automatic compensative movement: for the unified representation between body and action. *Experimental Brain Research, 233*, 777-785.

Bahcall, D. O., & Kowler, E. (1999). Illusory shifts in visual direction accompany adaptation of saccadic eye movements. *Nature, 400*(6747), 864-866.

Batterink, L. J., Reber, P. J., Neville, H. J., & Paller, K. A. (2015). Implicit and explicit contributions to statistical learning. *Journal of memory and language, 83*, 62-78.

Benson, B. L., Anguera, J. A., & Seidler, R. D. (2011). A spatial explicit strategy reduces error but interferes with sensorimotor adaptation. *Journal of neurophysiology, 105*(6), 2843-2851.

Berniker, M., & Körding, K. (2008). Estimating the sources of motor errors for adaptation and generalization. *Nature neuroscience, 11*(12), 1454-1461.

Bond, K. M., & Taylor, J. A. (2015). Flexible explicit but rigid implicit learning in a visuomotor adaptation task. *Journal of neurophysiology, 113*(10), 3836-3849.

Botvinick, M., & Cohen, J. (1998). Rubber hands 'feel'touch that eyes see. *Nature, 391*(6669), 756-756.

Bromberg, Z., Donchin, O., & Haar, S. (2019). Eye movements during visuomotor adaptation represent only part of the explicit learning. *Eneuro, 6*(6).

Brown, L. E., Rosenbaum, D. A., & Sainburg, R. L. (2003). Limb position drift: implications for control of posture and movement. *Journal of neurophysiology, 90*(5), 3105-3118.

Burr, D. C., & Morrone, M. C. (2012). Constructing stable spatial maps of the word. *Perception, 41*(11), 1355-1372.

Cheng, S., & Sabes, P. N. (2006). Modeling sensorimotor learning with linear dynamical systems. *Neural Comput, 18*(4), 760-793.

Cheng, S., & Sabes, P. N. (2006). Modeling sensorimotor learning with linear dynamical systems. *Neural computation, 18*(4), 760-793.

Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future of cognitive science. *Behavioral and brain sciences, 36*(3), 181-204.

Collins, T., & Wallman, J. (2012). The relative importance of retinal error and prediction in saccadic adaptation. *Journal of neurophysiology, 107*(12), 3342-3348.

Coltman, S. K., Cashaback, J. G., & Gribble, P. L. (2019). Both fast and slow learning processes contribute to savings following sensorimotor adaptation. *Journal of neurophysiology, 121*(4), 1575-1583.

D'Amario, S., Ruttle, J. E., 't Hart, B. M., & Henriques, D. Y. (2024). Implicit Adaptation is Fast, Robust and Independent from Explicit Adaptation. *bioRxiv*, 2024.2004. 2010.588930.

Daliri, A., & Dittman, J. (2019). Successful auditory motor adaptation requires task-relevant auditory errors. *Journal of neurophysiology, 122*(2), 552-562.

Darainy, M., Manning, T. F., & Ostry, D. J. (2023). Disruption of somatosensory cortex impairs motor learning and retention. *Journal of Neurophysiology, 130*(6), 1521-1528.

de Brouwer, A. J., Albaghdadi, M., Flanagan, J. R., & Gallivan, J. P. (2018). Using gaze behavior to parcellate the explicit and implicit contributions to visuomotor learning. *Journal of neurophysiology, 120*(4), 1602-1615.

Della-Maggiore, V., & McIntosh, A. R. (2005). Time course of changes in brain activity and functional connectivity associated with long-term adaptation to a rotational transformation. *Journal of neurophysiology, 93*(4), 2254-2262.

Deubel, H. (1987). Adaptivity of gain and direction in oblique saccades. In *Eye movements from physiology to cognition* (pp. 181-190). Elsevier.

Deubel, H., Schneider, W. X., & Bridgeman, B. (1996). Postsaccadic target blanking prevents saccadic suppression of image displacement. *Vision research, 36*(7), 985-996.

DiZio, P., & Lackner, J. R. (2000). Congenitally blind individuals rapidly adapt to Coriolis force perturbations of their reaching movements. *Journal of neurophysiology, 84*(4), 2175-2180.

Donchin, O., Rabe, K., Diedrichsen, J., Lally, N., Schoch, B., Gizewski, E. R., & Timmann, D. (2012). Cerebellar regions involved in adaptation to force field and visuomotor perturbation. *Journal of neurophysiology, 107*(1), 134-147.

Dosher, B., & Lu, Z.-L. (2017). Visual perceptual learning and models. *Annual review of vision science, 3*(1), 343-363.

Duff, S. V., & Sainburg, R. L. (2007). Lateralization of motor adaptation reveals independence in control of trajectory and steady-state position. *Experimental Brain Research, 179*, 551-561.

Ebrahimi, S., van der Voort, B., & Ostry, D. J. (2024). The Consolidation of Newly Learned Movements Depends upon the Somatosensory Cortex in Humans. *Journal of Neuroscience, 44*(32).

Ebrahimi, S., & Ostry, D. J. (2024). The human somatosensory cortex contributes to the encoding of newly learned movements. *Proceedings of the National Academy of Sciences, 121*(6), e2316294121.

Ellis, R. (2009). Implicit and explicit learning, knowledge and instruction. *Implicit and explicit knowledge in second language learning, testing and teaching, 42*, 3-25.

Ernst, M., & Banks, M. (2002). Humans integrate visual and haptic information in a statistically optimal fashion. *Nature 415*(6870), 429-433.

Flegal, K. E., & Anderson, M. C. (2008). Overthinking skilled motor performance: Or why those who teach can't do. *Psychonomic Bulletin & Review, 15*, 927-932.

Franklin, D. W., So, U., Burdet, E., & Kawato, M. (2007). Visual feedback is not necessary for the learning of novel dynamics. *PloS one, 2*(12), e1336.

Galea, J. M., Vazquez, A., Pasricha, N., Orban de Xivry, J.-J., & Celnik, P. (2011). Dissociating the roles of the cerebellum and motor cortex during adaptive learning:

the motor cortex retains what the cerebellum learns. *Cerebral cortex, 21*(8), 1761-
1770.
Gastrock, R. Q., 't Hart, B. M., & Henriques, D. Y. (2023). Distinct learning, retention, and
generalization in de novo learning. *bioRxiv*, 2023.2010. 2002.560506.
Gonzalez Castro, L. N., Monsen, C. B., & Smith, M. A. (2011). The binding of learning to
action in motor adaptation. *PLoS computational biology, 7*(6), e1002052.
Green, T. D., & Flowers, J. H. (1991). Implicit versus explicit learning processes in a
probabilistic, continuous fine-motor catching task. *Journal of motor behavior, 23*(4),
293-300.
Griffa, G., Deleglise, A., Solano, A., Jacobacci, F., De Pino, G., & Della-Maggiore, V.
(2024). The human hippocampus is involved in implicit motor learning. *bioRxiv*,
2024.2008. 2030.610548.
Haar, S., & Donchin, O. (2020). A revised computational neuroanatomy for motor control.
*Journal of Cognitive Neuroscience, 32*(10), 1823-1836.
Haar, S., Donchin, O., & Dinstein, I. (2015). Dissociating visual and motor directional
selectivity using visuomotor adaptation. *Journal of Neuroscience, 35*(17), 6813-
6821.
Hadjiosif, A. M., Krakauer, J. W., & Haith, A. M. (2021). Did we get sensorimotor
adaptation wrong? Implicit adaptation as direct policy updating rather than forward-
model-based learning. *Journal of Neuroscience, 41*(12), 2747-2761.
Hardwick, R. M., Lesage, E., Eickhoff, C. R., Clos, M., Fox, P., & Eickhoff, S. B. (2015).
Multimodal connectivity of motor learning-related dorsal premotor cortex.
*Neuroimage, 123*, 114-128.
Harris, C. M., & Wolpert, D. M. (1998). Signal-dependent noise determines motor
planning. *Nature, 394*(6695), 780-784.
Henriques, D. Y., & Cressman, E. K. (2012). Visuomotor adaptation and proprioceptive
recalibration. *Journal of motor behavior, 44*(6), 435-444.
Herzfeld, D. J., Vaswani, P. A., Marko, M. K., & Shadmehr, R. (2014). A memory of errors
in sensorimotor learning. *science, 345*(6202), 1349-1353.
Hogan, N., & Flash, T. (1987). Moving gracefully: quantitative theories of motor
coordination. *Trends in Neurosciences, 10*(4), 170-174.
Holroyd, C. B., & Coles, M. G. (2002). The neural basis of human error processing:
reinforcement learning, dopamine, and the error-related negativity. *Psychological*
*review, 109*(4), 679.
Houde, J. F., & Jordan, M. I. (1998). Sensorimotor adaptation in speech production.
*science, 279*(5354), 1213-1216.
Hutter, S. A., & Taylor, J. A. (2018). Relative sensitivity of explicit reaiming and implicit
motor adaptation. *Journal of neurophysiology, 120*(5), 2640-2648.
Imamizu, H., Miyauchi, S., Tamada, T., Sasaki, Y., Takino, R., PuÈtz, B., Yoshioka, T., &
Kawato, M. (2000). Human cerebellar activity reflecting an acquired internal model
of a new tool. *Nature, 403*(6766), 192-195.
Izawa, J., Criscimagna-Hemminger, S. E., & Shadmehr, R. (2012). Cerebellar
contributions to reach adaptation and learning sensory consequences of action.
*Journal of Neuroscience, 32*(12), 4230-4239.

Johnson, D. D., Blumstein, D. T., Fowler, J. H., & Haselton, M. G. (2013). The evolution
of error: Error management, cognitive constraints, and adaptive decision-making
biases. *Trends in ecology & evolution, 28*(8), 474-481.
Jordan, M. I., & Rumelhart, D. E. (1992). Forward models: Supervised learning with a
distal teacher. *Cognitive science, 16*(3), 307-354.
Kahneman, D. (2011). Thinking, fast and slow. *Farrar, Straus and Giroux*.
Kasuga, S., Kurata, M., Liu, M., & Ushiba, J. (2015). Alteration of a motor learning rule
under mirror-reversal transformation does not depend on the amplitude of visual
error. *Neuroscience research, 94*, 62-69.
Kawato, M. (1999). Internal models for motor control and trajectory planning. *Current
opinion in neurobiology, 9*(6), 718-727.
Kayacan, E., & Chowdhary, G. (2019). Tracking error learning control for precise mobile
robot path tracking in outdoor environment. *Journal of Intelligent & Robotic
Systems, 95*, 975-986.
Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R., & Ivry, R. B. (2018). Invariant
errors reveal limitations in motor correction rather than constraints on error
sensitivity. *Communications biology, 1*(1), 19.
Kim, K. S., Gaines, J. L., Parrell, B., Ramanarayanan, V., Nagarajan, S. S., & Houde, J.
F. (2023a). Mechanisms of sensorimotor adaptation in a hierarchical state
feedback control model of speech. *PLoS computational biology, 19*(7), e1011244.
Kim, K. S., Hinkley, L. B., Dale, C. L., Nagarajan, S. S., & Houde, J. F. (2023b).
Neurophysiological evidence of sensory prediction errors driving speech
sensorimotor adaptation. *bioRxiv*.
Kim, K. S., & Max, L. (2021). Speech auditory-motor adaptation to formant-shifted
feedback lacks an explicit component: Reduced adaptation in adults who stutter
reflects limitations in implicit sensorimotor learning. *European Journal of
Neuroscience, 53*(9), 3093-3108.
Krakauer, J. W., Ghilardi, M.-F., Mentis, M., Barnes, A., Veytsman, M., Eidelberg, D., &
Ghez, C. (2004). Differential cortical and subcortical activations in learning
rotations and gains for reaching: a PET study. *Journal of neurophysiology, 91*(2),
924-933.
Krakauer, J. W., Hadjiosif, A. M., Xu, J., Wong, A. L., & Haith, A. M. (2019). Motor
Learning. *Compr Physiol, 9*(2), 613-663.
Lanillos, P., Franklin, S., Maselli, A., & Franklin, D. W. (2021). Active strategies for
multisensory conflict suppression in the virtual hand illusion. *Scientific Reports,
11*(1), 22844.
Law, C.-T., & Gold, J. I. (2009). Reinforcement learning can account for associative and
perceptual learning on a visual-decision task. *Nature neuroscience, 12*(5), 655-
663.
Lee, K., Oh, Y., Izawa, J., & Schweighofer, N. (2018). Sensory prediction errors, not
performance errors, update memories in visuomotor adaptation. *Scientific Reports,
8*(1), 16483.
Leow, L.-A., Marinovic, W., de Rugy, A., & Carroll, T. J. (2020). Task errors drive
memories that improve sensorimotor adaptation. *Journal of Neuroscience, 40*(15),
3075-3088.

Leow, L. A., Marinovic, W., de Rugy, A., & Carroll, T. J. (2018). Task errors contribute to
implicit aftereffects in sensorimotor adaptation. *European Journal of Neuroscience,*
*48*(11), 3397-3409.
Lu, Z.-L., & Dosher, B. A. (2022). Current directions in visual perceptual learning. *Nature*
*reviews psychology, 1*(11), 654-668.
Maresch, J., Mudrik, L., & Donchin, O. (2021). Measures of explicit and implicit in motor
learning: what we know and what we don't. *Neuroscience & Biobehavioral Reviews,*
*128*, 558-568.
Maresch, J., Werner, S., & Donchin, O. (2021). Methods matter: Your measures of explicit
and implicit processes in visuomotor adaptation affect your results. *European*
*Journal of Neuroscience, 53*(2), 504-518.
Marko, M. K., Haith, A. M., Harran, M. D., & Shadmehr, R. (2012). Sensitivity to prediction
error in reach adaptation. *Journal of neurophysiology, 108*(6), 1752-1763.
Maselli, A., Lanillos, P., & Pezzulo, G. (2022). Active inference unifies intentional and
conflict-resolution imperatives of motor control. *PLoS computational biology, 18*(6),
e1010095.
Masselink, J., & Lappe, M. (2021). Visuomotor learning from postdictive motor error. *Elife,*
*10*, e64278.
Mazzoni, P., & Krakauer, J. W. (2006). An implicit plan overrides an explicit strategy
during visuomotor adaptation. *J Neurosci, 26*(14), 3642-3645.
McDougle, S. D., Bond, K. M., & Taylor, J. A. (2015). Explicit and implicit processes
constitute the fast and slow processes of sensorimotor learning. *Journal of*
*Neuroscience, 35*(26), 9568-9579.
McDougle, S. D., Bond, K. M., & Taylor, J. A. (2017). Implications of plan-based
generalization in sensorimotor adaptation. *Journal of neurophysiology, 118*(1),
383-393.
McDougle, S. D., & Taylor, J. A. (2019). Dissociable cognitive strategies for sensorimotor
learning. *Nature communications, 10*(1), 40.
McLaughlin, S. C. (1967). Parametric adjustment in saccadic eye movements. *Perception*
*& Psychophysics, 2*(8), 359-362.
Mista, C., Arguissain, F., Ranieri, A., Nielsen, J., Andersen, H., Manresa, J. B., &
Andersen, O. (2024). Spatio-temporal modulation of cortical activity during motor
deadaptation depends on the feedback of task-related error. *Behavioural Brain*
*Research, 468*, 115024.
Miyamoto, Y. R., Wang, S., & Smith, M. A. (2020). Implicit adaptation compensates for
erratic explicit strategy in human motor learning. *Nature neuroscience, 23*(3), 443-
455.
Modchalingam, S., Ciccone, M., D'Amario, S., 't Hart, B. M., & Henriques, D. Y. (2023).
Adapting to visuomotor rotations in stepped increments increases implicit motor
learning. *Scientific Reports, 13*(1), 5022.
Modchalingam, S., Vachon, C. M., 't Hart, B. M., & Henriques, D. Y. (2019). The effects
of awareness of the perturbation during motor adaptation on hand localization.
*PloS one, 14*(8), e0220884.
Morehead, J. R., Taylor, J. A., Parvin, D. E., & Ivry, R. B. (2017). Characteristics of implicit
sensorimotor adaptation revealed by task-irrelevant clamped feedback. *Journal of*
*Cognitive Neuroscience, 29*(6), 1061-1074.

Morehead, J. R., & Xivry, J. J. O. D. (2021). A Synthesis of the Many Errors and Learning Processes of Visuomotor Adaptation.

Mostafa, A. A., 't Hart, B. M., & Henriques, D. Y. (2019). Motor learning without moving: Proprioceptive and predictive hand localization after passive visuoproprioceptive discrepancy training. *PloS one, 14*(8), e0221861.

Nasir, S. M., & Ostry, D. J. (2009). Auditory plasticity and speech motor learning. *Proceedings of the National Academy of Sciences, 106*(48), 20470-20475.

Neville, K.-M., & Cressman, E. K. (2018). The influence of awareness on explicit and implicit contributions to visuomotor adaptation over time. *Experimental Brain Research, 236*, 2047-2059.

Niv, Y. (2009). Reinforcement learning in the brain. *Journal of Mathematical Psychology, 53*(3), 139-154.

Oza, A., Kumar, A., Sharma, A., & Mutha, P. K. (2024). Limb-related sensory prediction errors and task-related performance errors facilitate human sensorimotor learning through separate mechanisms. *PLoS biology, 22*(7), e3002703.

Parrell, B., Agnew, Z., Nagarajan, S., Houde, J., & Ivry, R. B. (2017). Impaired feedforward control and enhanced feedback control of speech in patients with cerebellar degeneration. *Journal of Neuroscience, 37*(38), 9249-9258.

Parrell, B., Ivry, R. B., Nagarajan, S. S., & Houde, J. F. (2021). Intact correction for self-produced vowel formant variability in individuals with cerebellar ataxia regardless of auditory feedback availability. *Journal of Speech, Language, and Hearing Research, 64*(6S), 2234-2247.

Parrell, B., Ramanarayanan, V., Nagarajan, S., & Houde, J. (2019). The FACTS model of speech motor control: Fusing state estimation and task-based control. *PLoS computational biology, 15*(9), e1007321.

Patri, J.-F., Perrier, P., Schwartz, J.-L., & Diard, J. (2018). What drives the perceptual change resulting from speech motor adaptation? Evaluation of hypotheses in a Bayesian modeling framework. *PLoS computational biology, 14*(1), e1005942.

Pelisson, D., Alahyane, N., Panouilleres, M., & Tilikete, C. (2010). Sensorimotor adaptation of saccadic eye movements. *Neuroscience & Biobehavioral Reviews, 34*(8), 1103-1120.

Popa, L. S., & Ebner, T. J. (2019). Cerebellum, predictions and errors. *Frontiers in cellular neuroscience, 12*, 524.

Popa, L. S., Hewitt, A. L., & Ebner, T. J. (2012). Predictive and feedback performance errors are signaled in the simple spike discharge of individual Purkinje cells. *Journal of Neuroscience, 32*(44), 15345-15358.

Popa, L. S., Streng, M. L., & Ebner, T. J. (2017). Long-term predictive and feedback encoding of motor signals in the simple spike discharge of Purkinje cells. *Eneuro, 4*(2).

Rengarajan, S., Cannon, J., Baron, B., Mohan, N., & Chukoskie, L. (2022). Measuring the Interaction of Conflict-Minimizing and Goal-Seeking Motor Imperatives in Autism Spectrum Disorder. International Simulation and Gaming Association Conference,

Rossi, C., Bastian, A. J., & Therrien, A. S. (2021). Mechanisms of proprioceptive realignment in human motor learning. *Current Opinion in Physiology, 20*, 186-197.

Ruttle, J. E., 't Hart, B. M., & Henriques, D. Y. (2018). The fast contribution of visual-proprioceptive discrepancy to reach aftereffects and proprioceptive recalibration. *PloS one, 13*(7), e0200621.

Ruttle, J. E., 't Hart, B. M., & Henriques, D. Y. (2021). Implicit motor learning within three trials. *Scientific Reports, 11*(1), 1627.

Ruttle, J. E., 't Hart, B. M., & Henriques, D. Y. (2022). Reduced feedback barely slows down proprioceptive recalibration. *Journal of neurophysiology, 128*(6), 1625-1633.

Ruttle, J. E., Cressman, E. K., 't Hart, B. M., & Henriques, D. Y. (2016). Time course of reach adaptation and proprioceptive recalibration during visuomotor learning. *PloS one, 11*(10), e0163695.

Salomonczyk, D., Cressman, E. K., & Henriques, D. Y. (2011). Proprioceptive recalibration following prolonged training and increasing distortions in visuomotor adaptation. *Neuropsychologia, 49*(11), 3053-3062.

Salomonczyk, D., Cressman, E. K., & Henriques, D. Y. (2013). The role of the cross-sensory error signal in visuomotor adaptation. *Experimental Brain Research, 228*, 313-325.

Sanchez-Vives, M. V., Spanlang, B., Frisoli, A., Bergamasco, M., & Slater, M. (2010). Virtual hand illusion induced by visuomotor correlations. *PloS one, 5*(4), e10381.

Schlerf, J., Ivry, R. B., & Diedrichsen, J. (2012). Encoding of sensory prediction errors in the human cerebellum. *Journal of Neuroscience, 32*(14), 4913-4922.

Schmidt, R. A., Lee, T. D., Winstein, C., Wulf, G., & Zelaznik, H. N. (2018). *Motor control and learning: A behavioral emphasis*. Human kinetics.

Schween, R., Taube, W., Gollhofer, A., & Leukel, C. (2014). Online and post-trial feedback differentially affect implicit adaptation to a visuomotor rotation. *Experimental Brain Research, 232*, 3007-3013.

Seidler, R., Noll, D., & Chintalapati, P. (2006). Bilateral basal ganglia activation associated with sensorimotor adaptation. *Experimental Brain Research, 175*, 544-555.

Shadmehr, R., & Mussa-Ivaldi, F. A. (1994). Adaptive representation of dynamics during learning of a motor task. *The Journal of Neuroscience, 14*(5), 3208-3224.

Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction, and adaptation in motor control. *Annual review of neuroscience, 33*, 89-108.

Smith, M. A., Ghazizadeh, A., & Shadmehr, R. (2006). Interacting adaptive processes with different timescales underlie short-term motor learning. *PLoS Biol, 4*(6), e179.

Sober, S. J., & Sabes, P. N. (2005). Flexible strategies for sensory integration during motor planning. *Nature neuroscience, 8*(4), 490-497.

Striemer, C. L., Enns, J. T., & Whitwell, R. L. (2019). Visuomotor adaptation in the absence of input from early visual cortex. *Cortex, 115*, 201-215.

't Hart, B. M., & Henriques, D. Y. (2016). Separating predicted and perceived sensory consequences of motor learning. *PloS one, 11*(9), e0163556.

Tanaka, H., Sejnowski, T. J., & Krakauer, J. W. (2009). Adaptation to visuomotor rotation through interaction between posterior parietal and motor cortical areas. *Journal of neurophysiology, 102*(5), 2921-2932.

Taylor, J. A., & Ivry, R. B. (2011). Flexible cognitive strategies during motor learning. *PLoS computational biology, 7*(3), e1001096.

Taylor, J. A., Krakauer, J. W., & Ivry, R. B. (2014). Explicit and implicit contributions to learning in a sensorimotor adaptation task. *Journal of Neuroscience, 34*(8), 3023-3032.

Telgen, S., Parvin, D., & Diedrichsen, J. (2014). Mirror reversal and visual rotation are learned and consolidated via separate mechanisms: recalibrating or learning de novo? *Journal of Neuroscience, 34*(41), 13768-13779.

Thoroughman, K. A., & Shadmehr, R. (2000). Learning of action through adaptive combination of motor primitives. *Nature, 407*(6805), 742-747.

Torrecillos, F., Alayrangues, J., Kilavik, B. E., & Malfait, N. (2015). Distinct modulations in sensorimotor postmovement and foreperiod β-band activities related to error salience processing and sensorimotor adaptation. *Journal of Neuroscience, 35*(37), 12753-12765.

Tsay, J. S., Haith, A. M., Ivry, R. B., & Kim, H. E. (2022a). Interactions between sensory prediction error and task error during implicit motor learning. *PLoS computational biology, 18*(3), e1010005.

Tsay, J. S., Kim, H., Haith, A. M., & Ivry, R. B. (2022b). Understanding implicit sensorimotor adaptation as a process of proprioceptive re-alignment. *Elife, 11*, e76639.

Tsay, J. S., Lee, A., Ivry, R. B., & Avraham, G. (2021). Moving outside the lab: The viability of conducting sensorimotor learning studies online. *arXiv preprint arXiv:2107.13408*.

Tsay, J. S., Chandy, A. M., Chua, R., Miall, R. C., Cole, J., Farnè, A., ... & Sarlegna, F. R. (2024). Minimal impact of chronic proprioceptive loss on implicit sensorimotor adaptation and perceived movement outcome. *Journal of Neurophysiology, 132*(3), 770-780.

Tsay, J. S., Tan, S., Chu, M. A., Ivry, R. B., & Cooper, E. A. (2023). Low vision impairs implicit sensorimotor adaptation in response to small errors, but not large errors. *Journal of cognitive neuroscience, 35*(4), 736-748.

Tseng, Y., Diedrichsen, J., Krakauer, J., Shadmehr, R., & Bastian, A. (2007). Sensory prediction errors drive cerebellum-dependent adaptation of reaching. *Journal of neurophysiology, 98*(1), 54.

Tzvi, E., Gajiyeva, L., Bindel, L., Hartwigsen, G., & Classen, J. (2022). Coherent theta oscillations in the cerebellum and supplementary motor area mediate visuomotor adaptation. *Neuroimage, 251*, 118985.

Tzvi, E., Koeth, F., Karabanov, A. N., Siebner, H. R., & Krämer, U. M. (2020). Cerebellar–premotor cortex interactions underlying visuomotor adaptation. *Neuroimage, 220*, 117142.

Wei, K., & Kording, K. (2009). Relevance of error: what drives motor adaptation? *Journal of neurophysiology, 101*(2), 655-664.

Werner, S., Van Aken, B. C., Hulst, T., Frens, M. A., Van Der Geest, J. N., Strüder, H. K., & Donchin, O. (2015). Awareness of sensorimotor adaptation to visual rotations of different size. *PloS one, 10*(4), e0123321.

Wilterson, S. A., & Taylor, J. A. (2021). Implicit visuomotor adaptation remains limited after several days of training. *Eneuro, 8*(4).

Wolpert, D., & Kawato, M. (1998). Multiple paired forward and inverse models for motor control. *Neural Networks, 11*(7-8), 1317-1329.

Wolpert, D. M., Diedrichsen, J., & Flanagan, J. R. (2011). Principles of sensorimotor learning. *Nature reviews neuroscience, 12*(12), 739.

Wolpert, D. M., Miall, R. C., & Kawato, M. (1998). Internal models in the cerebellum. *Trends in cognitive sciences, 2*(9), 338-347.

Wong, A. L., & Shelhamer, M. (2011). Saccade adaptation improves in response to a gradually introduced stimulus perturbation. *Neuroscience letters, 500*(3), 207-211.

Wurtz, R. H. (2008). Neuronal mechanisms of visual stability. *Vision research, 48*(20), 2070-2089.

Zhang, Z., Wang, H., Zhang, T., Nie, Z., & Wei, K. (2024). Perceptual error based on Bayesian cue combination drives implicit motor adaptation. *Elife, 13*, RP94608.