Housing environment bilaterally alters transcriptomic profile in the rat hippocampal CA1 region

Short title: Transcriptomic profiles of the left and right hippocampal CA1 regions

Azusa Kubota[1], Kentaro Kojima[2], Shinnosuke Koketsu[3], Takayuki Kannon[2,4], Takehiro Sato[2,5], Kazuyoshi Hosomichi[2,6], Yoshiaki Shinohara[7*], Atsushi Tajima[1,2*]

[1] Department of Bioinformatics and Genomics, Graduate School of Advanced Preventive Medical Sciences, Kanazawa University, Kanazawa, Japan

[2] Department of Bioinformatics and Genomics, Graduate School of Medical Sciences, Kanazawa University, Kanazawa, Japan

[3] Department of Physical Therapy, Faculty of Medical Science, Nagoya Women's University, Nagoya, Japan

[4] Department of Biomedical Data Science, Fujita Health University School of Medicine, Toyoake, Japan

[5] Department of Human Biology and Anatomy, Graduate School of Medicine, University of the Ryukyu, Nishihara, Japan

[6] Laboratory of Computational Genomics, School of Life Science, Tokyo University of

Pharmacy and Life Sciences, Tokyo, Japan

[7] Department of Anatomy and Systems Biology, Faculty of Medicine, University of Yamanashi,

Yamanashi, Japan

*Corresponding authors

Email: atajima@med.kanazawa-u.ac.jp (AT); yoshinohara@yamanashi.ac.jp (YS)

Abstract

Brain asymmetry is a fundamental feature of neural organization. However, the molecular basis of hippocampal lateralization in response to environmental stimuli remains poorly understood. Here, we examined the transcriptomic profiles of the left and right hippocampal CA1 regions in rats reared under isolated or enriched housing conditions to elucidate hemisphere-specific responses and shared molecular adaptations. RNA-sequencing analysis revealed lateralized differences in the number and identity of differentially expressed genes, accompanied by distinct biological themes, as indicated by overrepresentation and gene set enrichment analysis. The left CA1 region was prominently engaged in pathways related to synaptic organization and mitochondrial function, whereas the right CA1 region exhibited enrichment in transcriptional regulation and RNA metabolic processes. Despite these asymmetries, co-expression and protein–protein interaction network analyses revealed shared molecular architectures. Immediate early genes formed consistent central hubs across both hemispheres, and a common Mecp2–Grin2b–Cdkl5–Tet3 protein interaction cluster was identified as a potential integrative regulatory module. Additional enrichment analysis of differentially expressed genes shared between hemispheres further highlighted conserved responses, particularly in synaptic plasticity and cell–cell communication. Together, these findings demonstrate that the left and right CA1 regions employ distinct yet partially convergent transcriptional programs to adapt to environmental stimuli. This coordinated

45   molecular asymmetry provides novel insights into hippocampal lateralization and its role in

46   experience-dependent brain plasticity.

48   Introduction

Numerous animals and humans have left–right asymmetric brains; in humans and

primates, the acquisition of higher brain functions is closely related to brain asymmetry [1,2].

Electroencephalography and functional magnetic resonance imaging are used to characterize

time- and space-dependent functional differences between the left and right cerebrum of the

human brain [3]. These functional differences are thought to be due to various responses to

environmental stimuli, including task performance and stress exposure. Although lateral

specialization of the brain has been extensively studied, the molecular factors and intricate

mechanisms remain largely unknown.

Studies on hippocampal asymmetry suggest that the genetic mechanisms of visceral

and brain lateralization are different. Chiral movement of the cilia induces a leftward nodal

flow during embryonic development and consistently produces predictable asymmetry in the

arrangement of the thoracic and abdominal organs in adulthood [4]. Normal ciliary function in

wild-type mice is associated with an uneven hippocampus, and the disruption of ciliary

motility leads to the randomization of organ laterality. In mice with arbitrarily inhibited ciliary

movements, bilateral right isomerism occurs in the hippocampal circuitry regardless of the

original direction of organ laterality [5]. Notably, specific gene expression patterns

occasionally drive molecular variations in the brain. For example, the *iv* mouse, which has a

mutation in the *Lrd* (left-right dynein) gene, exhibits disrupted organ lateralization and loss of

hippocampal lateralization, ultimately resulting in spatial reference and working memory

deficits [6].

In rodents, the size of the synapses of the CA1 pyramidal cells varies depending on

whether they receive input from the left or right CA3 pyramidal cells [7,8]. This intrinsic

asymmetry is closely associated with the differential expression patterns of glutamate receptor

subunits, which, in turn, correlate with synaptic size and plasticity potential. Individual

environmental experiences can further modulate this inherent asymmetry. For instance, rats

reared in enriched environments characterized by running wheels, tunnels, and toys exhibit

increased synaptic density, specifically in the right CA1 region, along with enhanced

interhemispheric gamma coherence [9]. Environmental enrichment also leads to greater

dendritic complexity, maintenance of spine density, and increased branching of hippocampal

CA1 pyramidal neurons in the hippocampus [10]. These findings suggest that environmental

stimulation actively facilitates functional lateralization within the hippocampal circuitry

through activity-dependent synaptic and dendritic reorganization. Likewise, environmental

enrichment induces profound neuroanatomical and molecular changes in other hippocampal

subregions [11]. In the dentate gyrus, enrichment robustly stimulates adult neurogenesis,

synaptogenesis, and survival of new neurons, while also increasing levels of neurotrophic

factors and enhancing synaptic transmission. CA3, while less frequently studied, also exhibits

increased synaptic plasticity and network connectivity, supporting improved pattern

completion and associative memory. Furthermore, environmentally induced lateralization and

plasticity are not unique to rodents. They have been observed across diverse species, including

zebrafish, birds, and humans, indicating that experience-dependent brain asymmetry is a

highly conserved and functionally important phenomenon [12,13]. Therefore, to fully

understand the intricate mechanisms underlying neural asymmetry formation, it is essential to

consider not only the structural and electrophysiological aspects, but also the dynamic,

context-dependent gene expression responses of hippocampal regions to environmental

stimuli.

Differential mRNA profiles between the whole left and right rat hippocampi have

been observed using microarrays [14], and interhemispheric differences in miRNA profiles

have been found in the rat hippocampal CA3 region [15]. Moreover, previous reports revealed

that housing environments alter gene expression in the rodent brain, including the

hippocampus [16-18]. However, the genes responsible for neurological laterality in the rodent

hippocampus and the environmental influence on asymmetry remain unknown. Therefore, in

this study, we aimed to elucidate the transcriptomic profiles of the left and right CA1 regions

and to assess their respective sensitivity to an enriched environment using RNA-sequencing

(RNA-seq). Through systematic analyses of differentially expressed genes (DEGs) across hemispheric and environmental contrasts, including counts, functional enrichment, and co-expression/protein–protein interaction networks, we identified both distinct and shared molecular responses of the bilateral CA1 regions to environmental stimuli.

Materials and methods

Animals and microdissection

Male Long Evans rats were reared under either isolated (ISO) or enriched (ENR) conditions. Rats in the ISO group were caged individually after weaning at 21 days of age and raised in standard cages (length, 32 cm; width, 22 cm; height, 13.5 cm). In the ENR group, five to six male littermates were housed together in a larger cage (length, 44 cm; width, 27 cm; height, 18.7 cm) with a ladder, running wheels, tunnels, and toys, the location of which was changed every 3–4 days. Hippocampal slices were prepared, as described previously, 4 weeks after rearing under the respective environmental conditions [7,8,15]. Briefly, the rats were euthanized with isoflurane, and their brains were immediately washed with ice-cold artificial cerebrospinal fluid (ACSF). After cooling in ACSF for 15 min, hippocampal slices were cut into 400-μm-thick sections using a McIlwain tissue chopper. The bilateral rat CA1 regions were manually dissected under a stereoscopic microscope (Olympus, Tokyo, Japan) using a handmade microblade. Three biological replicates were prepared for sequencing from the left

121   and right CA1 regions. To minimize the effects of changes in the number of animals in the cage,

122   ENR brain slices were prepared on the same day. The Animal Experimental Committee of

123   Nagoya City University approved the study design (approval number: H29M-44). The animal

124   study protocol complied with the ARRIVE guidelines 2.0 (https://arriveguidelines.org).

Library preparation and RNA-seq

Total RNA was extracted from rat CA1 region tissues using the RNeasy® Mini Kit

(Qiagen, Germany), according to the manufacturer's protocol. All RNA samples exhibited an

appropriate range of RNA integrity number values, as measured using a bioanalyzer (Agilent,

USA). The cDNA libraries for RNA-seq were prepared from 500 ng of extracted RNA using

the TruSeq® Stranded mRNA Sample Preparation Kit (Illumina, USA), according to the

manufacturer's instructions. The libraries were sequenced on an Illumina HiSeq2000 to obtain

101-bp paired-end reads for each sample.

Computational quantification of gene expression

The sequenced reads were processed using fastp [19] (version 0.23.4) to eliminate

low-quality reads and trim adapter sequences. Transcript quantification was performed using

Salmon [20] (version 1.10.3) with Ensembl release 113 of the mRatBN7.2 cDNA assembly.

Salmon output files were converted into read-count data and summarized from transcript-level

to gene-level quantification using tximport [21] (version 1.34.0). The raw gene count data for

downstream analyses are provided in the S1 Data. The converted raw gene count data were

normalized using tximport with the transcripts per million (TPM) method for multivariate

analysis, as described below. Hierarchical clustering was conducted to summarize sample

similarities based on Pearson's correlation, and multidimensional scaling was used to

visualize the relationships among the samples. TPM-normalized gene count data are provided

in S2 Data.

Differentially expressed genes (DEGs) analysis

Statistical analyses of DEGs were performed using edgeR [22-24] (version 4.4.2),

which applies an empirical Bayes framework to stabilize dispersion estimates and is widely

recognized as an appropriate method for RNA-seq datasets with limited biological replicates.

After filtering out genes with very low expression using the edgeR::filterByexpr() function, the

quasi-likelihood F-test was used to evaluate the significance of gene expression differences.

An appropriate design matrix was formulated for the paired left–right sample quasi-likelihood

F-test analysis. Nominal p-values were adjusted using false discovery rate (FDR) correction. In

this step, the left group was treated as the control in the left–right comparison, and the ISO

group was treated as the control in the ISO-ENR comparison. DEGs were detected using

arbitrary thresholds of FDR = 0.1 and $\log_2$(fold-change) ($\log_2$FC) = $\pm\log_2 1.5$ (= ± 0.58).

Overlapping ISO-ENR DEGs between the left and right CA1 regions were visualized as Venn

diagrams using the jvenn web application (http://bioinfo.genotoul.fr/jvenn) [25]. The primary

assembly FASTA and genomic annotation GTF files of mRatBN7.2 were downloaded from

Ensembl release 113.

Correlation and multivariate analysis

161   The $\log_2FC$ values of the 14,368 common Ensembl gene IDs from housing condition

162 comparisons of the left and right CA1 regions were used for correlation analysis. To ensure the

163 robustness of multivariate analysis, gene-wise expression variability was quantified using the

164 mean absolute $\log_2FC$, and the top 500 genes were selected based on this metric for heatmap

165 generation.

Reverse transcription quantitative polymerase chain reaction (RT-qPCR)

For validation purposes, two representative DEGs identified by RNA-seq were

selected for RT-qPCR analysis. The same RNA samples used for sequencing were also

utilized for qPCR validation, with the analysis aimed at confirming the consistency of

expression trends rather than providing independent replication. The number of biological

replicates was identical to that used in the RNA-seq experiment (n = 3 per condition).

Reverse transcription was performed using ReverTra Ace® qPCR RT Master Mix (TOYOBO,

Japan). The PCR mixture was prepared with 10.0 μL of THUNDERBIRD® Next SYBR

qPCR Mix (TOYOBO, Japan), 0.3 μM of forward and reverse primer, 1.0 μL of cDNA, and

7.8 μL of sterilized water. QuantStudio 6 Pro (Applied Biosystems, USA) was used for qPCR

analysis. A relative quantification method [26] was used to measure the amount of each gene,

with *Gapdh* as a normalizer. Details of the primer sequences used are provided in S1 Table.

Welch's *t*-test was applied to account for potential heterogeneity of variance between the ISO

and ENR groups, providing more reliable inference when the number of biological replicates

is limited. Click-qPCR [27] was used for the statistical analysis and visualization of gene

expression using the ΔCq values.

Over-representation analysis (ORA)

ORA   of   ISO-ENR   DEGs   was   conducted   using   g:Profiler   [28]

(https://biit.cs.ut.ee/gprofiler,   version e113_eg59_p19_f6a03c19,   database   updated,

23/05/2025). Significant enrichment was tested across multiple categories, with Gene

Ontology (GO) [29] (GO:Molecular Function [MF], GO:Biological Process [BP],

GO:Cellular Component [CC]), Reactome [30], Kyoto Encyclopedia of Genes and Genomes

(KEGG) [31], and TRANSFAC [32] as the transcription factor target databases and

miRTarBase [33] as a miRNA motif database. Notably, driver GO terms were identified by

grouping significant GO terms into relational clusters and selecting representative terms that

explained the enrichment of the surrounding functions, whereas other emphasized terms were

manually selected. The significance threshold was set at 0.05, following the g:SCS method,

which adjusts *p*-values for GO enrichment by accounting for the hierarchical structure of GO

terms. This approach provides a more accurate multiple testing correction than standard

methods, such as the Bonferroni and Benjamini–Hochberg corrections for dependent tests.

This reduces redundancy and highlights key biological themes by retaining multiple leading

terms for functional groups [34].

Gene co-expression and protein–protein interaction (PPI) network analysis

DEGs identified from environmental comparisons using Ensembl gene IDs were

converted to gene symbols and annotated using the g:Convert tool in g:Profiler. Next,

Cytoscape [35] (version 3.10.2) and its plugins GeneMANIA [36] (application version: 3.5.3)

and STRING [37] (application version: 2.2.0) were used to analyze gene co-expression and

PPI networks among the converted DEGs. In the network visualization, the size and color of

each node indicate the calculated betweenness centrality in the network, with larger and more

intensely colored nodes representing a relatively higher centrality. The thickness of each edge

indicates the normalized maximum weight score, which reflects the strength or confidence of

the association between two genes or proteins. To improve clarity, isolated nodes were

manually removed from the network.

Gene set enrichment analysis (GSEA)

GSEA [38] was performed on the environmental comparison data using

clusterProfiler [39-41] (version 4.14.6) and fgsea [42] (version 1.32.4), with the R package

org.Rn.eg.db (version 3.20.0) used for genomic annotation. GO was used to characterize the

biological features of the gene sets using the clusterProfiler:gseGO() function. To enhance the

interpretability of the enrichment analysis, the Molecular Signatures Database (MSigDB)

[43] and its R package msigdbr (version 24.1.0) were used for pathway enrichment analysis

using the clusterProfiler::GSEA() function. The log_2FC values calculated using edgeR were

used as input, and the following parameters were specified for GSEA: minGSSize = 10,

maxGSSize = 500, eps = 0, nPermSimple = 10000, pvalueCutoff = 0.1, pAdjustMethod =

"BH" (Benjamini-Hochberg), and by = "fgsea." In the resulting dot plot, the GeneRatio (i.e.,

the ratio of core enrichment genes to the total number of genes in each gene set) was plotted

on the x-axis, indicating the proportion of genes that most strongly contributed to enrichment.

In GSEA, core-enriched genes are defined as the subset of genes that contribute most

significantly to the enrichment score, thereby representing the central elements driving the

biological changes within each gene set. These genes were then used to calculate the

GeneRatio in the dot plot, indicating their contribution to primary changes in the pathway.

Data processing, statistics, and drawing

R (version 4.4.2) and its tidyverse package (version 2.0.0) were used for data

preprocessing, statistical computing, and data visualization.

Results

Comparison of transcriptomic similarities

Transcript quantification and summarization were successfully performed, resulting

in the identification of 23,978 genes; each was annotated with a unique Ensembl gene ID.

The raw gene count dataset was normalized to scaled TPM values, which were used for the

analyses described in this section (S1 Data). To examine the overall transcriptomic similarity

between samples, multivariate analyses were conducted based on the scaled TPM values (S2

Fig). The hierarchical clustering dendrogram revealed a clear separation between the ENR

and ISO groups and the left and right CA1 region samples from the same individual clustered

as pairs. Furthermore, the multidimensional scaling plot based on the scaled TPM values

illustrated the relative distances among the samples.

DEG analysis and RT-qPCR

DEG analysis showed that no genes met the significance criteria in the left–right

comparison under the ENR conditions, whereas only one DEG, *Pklm1*, was identified under

the ISO conditions (S3 Fig). The complete edgeR results for the left–right comparison are

provided in S2 Data. Conversely, the ISO–ENR comparison revealed significant differences

in the gene expression profiles in both the left and right CA1 regions. In the left CA1 region,

189 DEGs were identified: 25 genes were highly expressed in the ENR group, and 164 genes

were highly expressed in the ISO group (Fig 1A). In the right CA1 region, 94 DEGs were

detected: 13 genes were highly expressed in the ENR group and 81 genes in the ISO group

(Fig 1B). The full edgeR results for the ISO–ENR comparison are available in the S3 Data,

and the corresponding DEG lists are compiled in the S4 Data.

Fig 1. Differentially expressed genes (DEGs) in the housing condition comparison

between the left and right CA1 regions. (A) and (B) show volcano plots illustrating the

distribution of statistically tested genes. FDR: false discovery rate; FC: fold-change.

Thresholds for DEG analysis (FDR = 0.1 and $\log_2$FC = $\pm\log_2 1.5$) are indicated using green

dashed lines. (A) ISO–ENR comparison in the left CA1 region. (B) ISO–ENR comparison in

the right CA1 region. ISO, isolated housing conditions; ENR, enriched housing conditions.

The number of overlapping ISO–ENR DEGs between the left and right CA1 regions

was visualized using Venn diagrams (Fig 2A). A total of 70 DEGs were shared, with seven

upregulated in the ENR group and 63 upregulated in the ISO group. All shared DEGs are

listed in Table 1. To validate the DEG analysis results, RT-qPCR experiments were carried

out using the selected genes from Table 1 according to the following criteria: $\log_{10}$CPM > 3

and FDR < 0.05. Among the genes that met these criteria, *Cdkl5* and *Fos* genes were selected

for RT-qPCR analysis. *Cdkl5* and *Fos* showed significantly lower relative expression in the

ENR group than in the ISO group in both the left and right CA1 regions (Fig 2B).

Fig 2. Summary of bilaterally shared ISO–ENR DEGs and RT-qPCR validation. (A)

Venn diagrams showing the overlap between the ISO–ENR DEG sets in the left and right

CA1 regions. (B) RT-qPCR quantification of *Fos* and *Cdkl5* expression, normalized to *Gapdh*.

*$*p < 0.05$, $**p < 0.01$, $***p < 0.001$. ISO, isolated housing conditions; ENR, enriched

housing conditions; DEGs, differentially expressed genes; RT-qPCR, reverse

transcription-quantitative polymerase chain reaction.

Table 1. Shared DEGs by comparing rearing environments in bilateral CA1.

Gene	Description	logFC (L)	logCPM (L)	FDR (L)	logFC (R)	logCPM (R)	FDR (R)
Abca5	ATP binding cassette subfamily A member 5	-2.600	1.191	0.066	-9.905	1.185	0.014
Adgrf5	adhesion G protein-coupled receptor F5	-0.697	3.738	0.089	-0.919	3.630	0.032
Ago3	argonaute RISC catalytic component 3	-0.750	3.282	0.037	-0.653	3.294	0.026
Ankrd52	ankyrin repeat domain 52	-0.695	4.779	0.003	-0.596	4.785	0.048
Apold1	apolipoprotein L domain containing 1	-1.702	2.150	0.023	-2.317	2.306	0.023
Arfgef3	ARFGEF family member 3	-0.588	5.949	0.009	-0.629	5.886	0.016
Bach2	BTB domain and CNC homolog 2	-0.803	1.885	0.052	-0.772	1.879	0.056
Bmp3	bone morphogenetic protein 3	-0.684	2.255	0.070	-1.332	2.081	0.038
Btg2	BTG anti-proliferation factor 2	-1.224	2.917	0.013	-1.244	2.908	0.026
Cacna1e	calcium voltage-gated channel subunit alpha1 E	-0.891	7.262	0.002	-0.832	7.217	0.015
Caln1	calneuron 1	-0.728	4.474	0.005	-0.778	4.363	0.039
Cbl	Cbl proto-oncogene	-0.714	4.648	0.004	-0.690	4.675	0.018
Cchcr1	coiled-coil alpha-helical rod protein 1	0.829	2.310	0.025	0.821	2.174	0.081
Cdkl5	cyclin-dependent kinase-like 5	-0.840	5.585	0.003	-0.769	5.549	0.014
Cds2	CDP-diacylglycerol synthase 2	-0.662	5.235	0.010	-0.737	5.171	0.014
Clmn	calmin	-0.724	5.496	0.005	-0.639	5.435	0.032
Csrnp3	cysteine and serine rich nuclear protein 3	-0.781	4.124	0.007	-0.602	4.129	0.099
Dgkh	diacylglycerol kinase, eta	-1.414	3.131	0.003	-0.931	3.096	0.086
Egfr	epidermal growth factor receptor	-0.748	2.294	0.096	-0.790	2.227	0.068
Egr2	early growth response 2	-1.455	2.856	0.006	-1.673	2.710	0.036
Elfn2	extracellular leucine-rich repeat and fibronectin type III domain containing 2	-0.917	7.317	0.006	-0.832	7.217	0.032
Erbb4	erb-b2 receptor tyrosine kinase 4	-1.030	3.940	0.013	-0.757	3.845	0.014
Flt1	Fms related receptor tyrosine kinase 1	-0.676	4.561	0.089	-0.882	4.485	0.073

Gene	Description	logFC (L)	logCPM (L)	FDR (L)	logFC (R)	logCPM (R)	FDR (R)
Fos	Fos proto-oncogene, AP-1 transcription factor subunit	-1.835	3.075	0.002	-1.876	3.045	0.009
Fzd3	frizzled class receptor 3	-0.656	4.324	0.014	-0.597	4.313	0.018
Gabrb1	gamma-aminobutyric acid type A receptor subunit beta1	-0.839	4.376	0.010	-0.825	4.264	0.014
Gan	gigaxonin	-1.038	1.889	0.026	-1.137	1.829	0.060
Grin2a	glutamate ionotropic receptor NMDA type subunit 2A	-0.932	7.760	0.002	-0.717	7.752	0.032
Grin2b	glutamate ionotropic receptor NMDA type subunit 2B	-1.411	5.835	0.0004	-1.123	5.771	0.005
Gucy1a2	guanylate cyclase 1 soluble subunit alpha 2	-1.066	2.955	0.007	-1.007	3.005	0.014
Hivep3	HIVEP zinc finger 3	-0.681	5.667	0.006	-0.587	5.594	0.067
Igsf9b	immunoglobulin superfamily, member 9B	-1.368	2.790	0.002	-1.083	2.644	0.051
Ino80d	INO80 complex subunit D	-1.175	2.272	0.004	-1.025	2.323	0.024
Junb	JunB proto-oncogene, AP-1 transcription factor subunit	-0.745	5.479	0.004	-0.788	5.419	0.018
Kdr	kinase insert domain receptor	-0.923	2.770	0.091	-1.324	2.725	0.057
Klf10	KLF transcription factor 10	-0.603	5.020	0.020	-0.652	5.014	0.038
Mecp2	methyl CpG binding protein 2	-0.836	3.695	0.003	-0.764	3.624	0.039
Mrtfb	myocardin related transcription factor B	-0.655	6.576	0.005	-0.617	6.494	0.032
Mtcl2	microtubule crosslinking factor 2	-0.953	4.782	0.003	-0.758	4.723	0.051
Npas4	neuronal PAS domain protein 4	-1.267	2.819	0.070	-1.429	2.877	0.090
Nsd3	nuclear receptor binding SET domain protein 3	-0.755	4.227	0.004	-0.753	4.185	0.014
Pdzrn4	PDZ domain containing RING finger 4	-1.111	1.705	0.015	-0.973	1.732	0.065
Phc3	polyhomeotic homolog 3	-0.732	3.975	0.004	-0.615	3.983	0.043
Ppp1r9a	protein phosphatase 1, regulatory subunit 9A	-0.625	6.169	0.003	-0.604	6.112	0.014
Ptar1	protein prenyltransferase alpha subunit repeat containing 1	-0.931	2.587	0.007	-0.806	2.737	0.027
Ptprb	protein tyrosine phosphatase, receptor type, B	-0.695	4.413	0.084	-0.947	4.258	0.086
Rc3h1	ring finger and CCCH-type domains 1	-0.662	3.984	0.007	-0.593	3.936	0.032

Gene	Description	logFC (L)	logCPM (L)	FDR (L)	logFC (R)	logCPM (R)	FDR (R)
RT1-CE15	RT1 class I, locus CE15	-2.765	1.491	0.024	-4.398	1.570	0.015
RT1-CE16	RT1 class I, locus CE16	0.770	4.842	0.010	0.844	4.844	0.014
RT1-N1	RT1 class Ib, locus N1	1.451	2.212	0.003	1.430	2.082	0.014
RT1-N3	RT1 class Ib, locus N3	1.024	1.610	0.010	1.170	1.582	0.034
Scn8a	sodium voltage-gated channel alpha subunit 8	-0.720	7.846	0.002	-0.679	7.784	0.014
Slc1a2	solute carrier family 1 member 2	-0.836	9.126	0.003	-0.762	9.074	0.043
Slc4a8	solute carrier family 4 member 8	-0.834	4.395	0.004	-0.750	4.283	0.051
Slc7a14	solute carrier family 7, member 14	-0.623	4.700	0.004	-0.602	4.645	0.042
Ston2	stonin 2	-0.796	4.162	0.004	-0.881	4.052	0.038
Tanc2	tetratricopeptide repeat, ankyrin repeat and coiled-coil containing 2	-0.714	6.028	0.002	-0.618	5.989	0.030
Tead1	TEA domain transcription factor 1	-0.647	2.047	0.042	-0.841	1.952	0.066
Tet3	tet methylcytosine dioxygenase 3	-0.863	4.300	0.003	-0.727	4.165	0.032
Tmem245	transmembrane protein 245	-0.609	4.184	0.032	-0.857	4.199	0.026
Tnrc6b	trinucleotide repeat containing adaptor 6B	-1.121	4.268	0.002	-0.918	4.229	0.014
Xkr4	XK related 4	-0.648	4.784	0.013	-0.668	4.755	0.014
Zbtb20	zinc finger and BTB domain containing 20	-1.572	3.396	0.002	-1.178	3.305	0.015
Zfp704	zinc finger protein 704	-1.001	3.440	0.004	-0.794	3.400	0.043
ENSRNOG00000066523	None	1.330	3.232	0.013	1.397	3.164	0.027
ENSRNOG00000066876	None	-0.991	1.982	0.071	-1.279	1.981	0.014
ENSRNOG00000067970	None	-1.034	2.914	0.011	-0.911	2.874	0.065
ENSRNOG00000069024	None	-0.908	3.174	0.026	-1.060	3.201	0.047
ENSRNOG00000069624	None	1.712	1.187	0.039	1.280	0.904	0.049
ENSRNOG00000070648	None	0.842	5.558	0.028	0.964	5.551	0.053

logCPM, log$_{10}$CPM; logFC, log$_2$(fold-change); FDR, false discovery rate. L: left, R: right; DEGs, differentially expressed genes.

To further characterize the ISO–ENR DEGs, correlation and multivariate analyses

were conducted using $\log_2$FC values from the edgeR analysis of the left and right CA1

regions (S4 Fig). The correlation plot of the $\log_2$FC values for environmental comparisons

between the left and right CA1 regions yielded a Pearson correlation coefficient of 0.587.

Additionally, a heatmap based on the top 500 genes ranked by absolute $\log_2$FC values

showed a high degree of directional similarity in gene expression in the bilateral CA1

regions.

Taken together, these results indicate that although the number of

environment-responsive genes differs between hemispheres (left: 189; right: 94), the overall

direction of gene expression changes in response to the environment is highly correlated

between the left and right CA1 regions.

291 ORA of bilateral DEGs in the environmental comparison

To elucidate the biological significance of the effects of the housing environment on

bilateral CA1 regions, ORA was performed using the ISO–ENR DEGs identified in the left

and right hippocampal CA1 regions. All results obtained using g:Profiler are provided in the

S5 Data.

Fig 3A shows the functional associations of the DEGs in the left hippocampal CA1

region as determined with GO enrichment and pathway analyses. A total of 189 DEGs were

analyzed based on Ensembl gene IDs, of which 180 were annotated. This analysis revealed

significant enrichment of GO-MF terms related to gene transcription and DNA binding (e.g.,

sequence-specific DNA binding and DNA-binding transcription factor activity), as well as

membrane-associated protein functions (e.g., transmembrane signaling receptor activity,

voltage-gated ion channel activity, and glutamate-gated calcium ion channel activity). Among

BP terms, those associated with nervous system development (e.g., axonogenesis), synaptic

function, and memory formation (e.g., long-term memory and synaptic vesicle localization)

were identified. For CC, synapse- and axon-related terms (e.g., synaptic membrane, Schaffer

collateral-CA1 synapse, and axon initial segment) were significantly enriched. Pathway

analysis indicated significant enrichment of the MAPK signaling pathway (KEGG:04010)

and the nuclear receptor transcription pathway (REAC:R-NOR-383280).

Fig 3. Manhattan plots showing significantly enriched terms from over-representation

analyses of the left and right ISO–ENR DEGs. The numbered and emphasized points have

the following characteristics: Driver terms, representing the most relevant GO terms in a

tabular format, are highlighted in GO:MF, GO:BP, and GO:CC. Other highlighted points

were manually selected based on their significance in additional databases (KEGG Pathway,

REAC: Reactome Pathway, TF: TRANSFAC, and MIRNA: miRTarBase). The adjusted

p-values (p_{adj}) were calculated using the G:SCS method implemented in g:Profiler. The

threshold for statistical significance was set at 0.05. (A) Manhattan plot and the highlighted

term details for the left ISO–ENR DEGs. (B) Right ISO-ENR Manhattan plot and the

highlighted term details for the right ISO–ENR DEGs. ISO, isolated housing conditions;

ENR, enriched housing conditions; DEGs, differentially expressed genes; GO, Gene

Ontology; MF, Molecular function; BP, Biological process; CC, Cellular component; KEGG,

Kyoto Encyclopedia of Genes and Genomes.

Fig 3B shows the functional associations of the DEGs identified in the right

hippocampal CA1 region using GO enrichment and pathway analyses. A total of 94 DEGs

were analyzed based on Ensembl gene IDs, of which 87 were annotated. This analysis

revealed significant enrichment of transmembrane receptor protein kinase activity and DNA

methylation-related GO-MF terms (e.g., DNA 5-methylcytosine dioxygenase activity). In

GO-BP, the regulation of RNA metabolic processes was highly enriched. For GO-CC, the

terms related to cell junctions and postsynaptic density were significantly represented.

Pathway analysis revealed significant enrichment of NPAS4-mediated transcriptional

regulation.

Fig 4 illustrates the functional associations between shared DEGs identified through

GO enrichment and pathway analyses. From the 70 shared Ensembl gene IDs, 64

corresponding annotated gene symbols were used in this analysis. This analysis revealed

significant enrichment in GO-MF terms related to transcriptional regulator activity and

transmembrane receptor protein tyrosine kinase activity. In terms of GO-BP, the most

significantly enriched term was positive regulation of nucleobase-containing compound

metabolic processes, with the smallest adjusted p-value. The presynaptic membrane and

receptor complex were significantly represented in GO-CC. Pathway analysis revealed

significant enrichment in the calcium signaling pathway (KEGG:04020) and

post-transcriptional silencing by small RNAs (REAC:R-RNO-426496). Additionally, the

transcription factor E2F and microRNA miR-132-3p were found to be enriched.

Fig 4. Manhattan plot showing significantly enriched terms from over-representation

analysis of the bilaterally shared ISO-ENR DEGs. The numbered and emphasized points

have the following characteristics: Driver terms, representing the most relevant GO terms in a

tabular format, are highlighted in GO:MF, GO:BP, and GO:CC. Other highlighted points

were manually selected based on their significance in additional preset databases (KEGG

Pathway, REAC: Reactome Pathway, TF: TRANSFAC, and MIRNA: miRTarBase). The

adjusted p-values (p_{adj}) were calculated using the G:SCS method implemented in g:Profiler.

The threshold for statistical significance was set at 0.05. ISO, isolated housing conditions;

ENR, enriched housing conditions; DEGs, differentially expressed genes; GO, Gene

Ontology; MF, Molecular function; BP, Biological process; CC, Cellular component; KEGG,

Kyoto Encyclopedia of Genes and Genomes.

Collectively, these ORA results suggest that the bilateral CA1 regions respond to environmental stimuli through distinct biological processes: the left hemisphere is enriched for pathways related to synaptic function and memory formation, whereas the right hemisphere is associated with transcriptional and epigenetic regulatory mechanisms.

Gene co-expression and PPI networks of DEGs in the environmental comparison

To evaluate the functional associations among ISO–ENR DEGs, gene co-expression network analyses were performed under three different conditions: left, right, and shared ISO–ENR DEGs. Gene co-expression networks were constructed using the GeneMANIA database in Cytoscape (Figs 5A, 6A, and 7A). All results from the Cytoscape analysis are provided in S6 Data. To highlight the common features of the bilaterally shared DEGs, highly correlated genes were predicted and visualized using a query gene-based weighting algorithm (Fig 7A). Across the three co-expression networks, a cluster of immediate early genes (IEGs), such as *Arc*, *Fos*, and *Junb,* was connected by thick edges. Additionally, key hub DEGs were highlighted in each network: *Shc3*, *Tnr*, and *Taok1* in the left CA1 region; *Ppp1r9a* and *Erbb4* in the right CA1 region; *Ppp1r9a*, *Jun*, and *Fosb* in the shared DEGs.

Fig 5. Gene co-expression and protein–protein interaction networks of the left CA1

region DEGs in the ISO-ENR comparison. A total of 180 DEGs annotated with gene

symbols were included in the analysis. In network visualization, node size and color represent

betweenness centrality, with larger and more intensely colored nodes indicating higher

centrality. Edge thickness represents the normalized maximum weight score, which reflects

the strength of the association between connected genes or proteins. Isolated nodes were

manually excluded to ensure visual clarity. (A) Gene co-expression network constructed

using the GeneMANIA database. (B) Protein–protein interaction network constructed using

the STRING database. ISO, isolated housing conditions; ENR, enriched housing conditions;

DEGs, differentially expressed genes.

Fig 6. Gene co-expression and protein–protein interaction networks of the right CA1

region DEGs in the ISO-ENR comparison. A total of 87 DEGs annotated with gene

symbols were included in the analysis. In network visualization, node size and color represent

betweenness centrality, with larger and more intensely colored nodes indicating higher

centrality. Edge thickness represents the normalized maximum weight score, which reflects

the strength of the association between connected genes or proteins. Isolated nodes were

manually excluded to ensure visual clarity. (A) Gene co-expression network constructed

using the GeneMANIA database. (B) Protein-protein interaction network constructed using

the STRING database. ISO, isolated housing conditions; ENR, enriched housing conditions;

DEGs, differentially expressed genes.

Fig 7. Gene co-expression and protein–protein interaction networks of bilaterally

shared ISO-ENR DEGs. A total of 64 DEGs annotated with gene symbols were included in

the analysis. In network visualization, node size and color represent betweenness centrality,

with larger and more intensely colored nodes indicating higher centrality. Edge thickness

represents the normalized maximum weight score, which reflects the strength of the

association between connected genes or proteins. Isolated nodes were manually excluded to

ensure visual clarity. (A) Gene co-expression network constructed using the GeneMANIA

database. A query gene-based weighting algorithm was applied to highlight highly correlated

genes (marked with red borders) within the network, thereby emphasizing common features

among shared DEGs. (B) Protein–protein interaction network constructed using the STRING

database. ISO, isolated housing conditions; ENR, enriched housing conditions; DEGs,

differentially expressed genes.

To examine interactions at the protein level, PPI network analyses were also

conducted under the same three conditions. PPI networks were generated using the STRING

database in Cytoscape (Figs 5B, 6B, and 7B). In these networks, Fos consistently appeared as

a central hub protein. In addition to the IEG protein cluster, PPI network analysis revealed

strong connections that were not observed in the co-expression networks, including

Fos–Npas4, Grin2a–Grin2b–Slc1a2, and Grin2b–Mecp2–Cdkl5–Tet3.

GSEA in the environmental comparison

416   To biologically interpret the transcriptomic alterations, gene lists ranked by their

corresponding log_2FC values were used to perform GSEA. The directionality of GO

enrichment was evaluated using the log_2FC values as a ranking metric. All GSEA results

obtained using clusterProfiler are provided in S7 Data. The dot plots in Fig 8 show the

GO-GSEA results focusing on the environmental conditions in the left and right CA1 regions.

The analyses identified core enrichment genes, defined as the subset of genes that contributed

most significantly to the enrichment score and potentially act as key drivers of the biological

changes within each gene set (see S7 Data).

Fig 8. Biological profiles of environmental conditions based on gene set enrichment

analysis (GSEA). In the ISO–ENR comparison, GSEA using Gene Ontology terms was

conducted with log_2(fold change) values from the edgeR analysis as input. Adjusted *p*-values

are shown below each dot. The x-axis of the dot plot represents GeneRatio, which is defined as

the proportion of genes within a gene set that belong to the core enrichment subset. (A) Dot

plot for the left ISO–ENR. (B) Dot plot for the right ISO–ENR. ISO, isolated housing

conditions; ENR, enriched housing conditions

In total, 1,293 gene sets were enriched in the left ISO–ENR analysis. Fig 8A

illustrates the GO-GSEA results for the left CA1 region, showing the top 10 gene sets ranked

by GeneRatio. Gene sets related to mitochondria and their functions showed a high degree of

enrichment in the activated direction, indicating enrichment under ENR conditions. In the

suppressed direction, corresponding to enrichment in the ISO condition, gene sets related to

synapses (e.g., asymmetric synapses, postsynaptic density) and DNA-binding transcription

activator activity were observed. Contrastingly, 496 gene sets were enriched in the right

ISO–ENR analysis. Fig 8B shows the GO-GSEA results in the right CA1 region, also

presenting the top 10 gene sets ranked by GeneRatio. Gene sets associated with protein

folding and RNA splicing were strongly enriched under the ENR conditions. Under the ISO

conditions, gene sets related to synapses and transcription activator activity were enriched,

consistent with the findings in the left CA1 region. Separate analyses were conducted for the

GO-BP, GO-CC, and GO-MF categories. The corresponding dot plots are presented in S5 Fig

(GO-BP), S6 Fig (GO-CC), and S7 Fig (GO-MF).

Finally, pathway-based enrichment analyses were performed using MSigDB's C2

Reactome and KEGG databases to complement the GO-based results. These results are

shown in Figs S8 (Reactome) and S9 (KEGG). The numbers of significant pathway gene sets

are as follows: Reactome, 197 on the left and 28 on the right; and KEGG, 93 on the left and

29 on the right (S7 data). These results strengthen the GO enrichment features described

above. In the ENR condition, the left CA1 region showed activation of mitochondrial

respiration, and the right CA1 region showed enrichment of RNA splicing. Under the ISO

conditions, both regions showed stimulations of synaptic development and function.

These GSEA results complement the ORA findings, suggesting that the left CA1

response under the ENR condition involves enhanced mitochondrial metabolic activity,

whereas the right CA1 response under the same condition is characterized by

post-transcriptional regulation through RNA splicing. Notably, synaptic processes were

commonly enriched in the ISO condition across both hemispheres, further reinforcing the

ORA results.

Discussion

This study aimed to elucidate how housing conditions influence gene expression in

the hippocampal CA1 region, revealing not only distinct transcriptomic landscapes but also

common molecular frameworks in the left and right hemispheres. Through comprehensive

analyses of DEGs, including their numbers, enrichment profiles, and co-expression/PPI

networks, we delineated both unique and shared molecular responses of the bilateral CA1

regions to environmental stimuli.

Transcriptomic landscapes of the bilateral CA1 regions in response to housing environments

While a direct left–right comparison revealed high baseline similarity (0–1 DEGs),

the subsequent ORA and GSEA analyses of environmental responses (ISO vs. ENR)

uncovered coherent and biologically distinct functional themes in each hemisphere. These

findings suggest that the observed qualitative asymmetry represents a genuine biological

response rather than a statistical artifact. This interpretation holds even though the overall

log_2FC values of all tested genes were moderately correlated between hemispheres (Pearson's

coefficient: $r = 0.587$, S4 Fig), indicating that despite a shared overall trend, the subset of

most significantly regulated genes diverged into functionally distinct pathways. In the

comparison between the housing conditions, the number of DEGs in the right CA1 region

was 94, whereas that in the left CA1 region was 189 (Fig 1). This suggests that the degree of

gene expression changes in response to the housing environment differs markedly between

the two hemispheres. To further interpret these differences, we performed an ORA on the

DEGs from both the left and right hippocampal CA1 regions (ISO vs. ENR conditions),

which revealed distinct biological processes and molecular mechanisms of each group.

Notably, both regions were enriched in the transcription factors E2F and miR-132-3p,

suggesting shared transcriptional and post-transcriptional control mechanisms in response to

environmental stimuli in the CA1 region [44,45].

Despite these similarities, significant differences were observed between the two

groups. In the left CA1 region, GO terms related to glutamate receptor activity, synapse

function, and long-term memory (e.g., glutamate-gated calcium ion channel activity and

long-term memory), along with the MAPK signaling pathway, were significantly enriched

(Fig 3A). These findings suggest that the activation of gene groups is crucial for

neuroplasticity and synaptic transmission, supporting the notion that environmental stimuli

predominantly activate mechanisms related to nervous system function and memory

formation in the left hemisphere [46]. Conversely, the right CA1 region exhibited enrichment

in pathways associated with RNA metabolism and cellular structural reorganization (e.g.,

regulation of RNA metabolic processes and cell periphery), as illustrated in Fig 3B.

Furthermore, the pathways regulated by the neuron-specific transcription factor NPAS4,

known for controlling activity-dependent gene expression [47], were enriched, highlighting

responses primarily involved in intrinsic cell regulation and homeostasis and suggesting a

focus on gene expression regulation in the right hemisphere response.

Next, we employed GSEA to evaluate functional trends at the gene set level in both

the left and right hippocampal CA1 regions ISO–ENR groups. The results provided deeper

insights into the molecular signatures of the two groups (Fig 8, S8 Fig, and S9 Fig). In the left

CA1 region, metabolic activities, including the mitochondrial respiratory chain, oxidative

phosphorylation, and cellular respiration, were significantly activated by environmental

enrichment. These functional categories primarily focus on energy production and align with

the previously observed activation of synaptic transmission, long-term memory, and MAPK

pathways (Fig 3A), suggesting that increased neural activity is linked to increased metabolic

demand. In contrast, post-transcriptional regulations within the nucleus and cytoplasm, such

as RNA splicing and protein refolding, were selectively activated in the right CA1 region.

This finding aligns with the activation of transcription factors such as NPAS4 and E2F, as

shown in Fig 3B, reflecting a central role for intracellular regulatory mechanisms in response

to stimuli. Notably, we observed similar enrichment patterns of asymmetric synaptic

structures and increased postsynaptic density gene sets under the ISO conditions in both the

left and right CA1 regions. Social isolation during adolescence increases gene and protein

expression of NR2B in the rodent hippocampus [48,49]. This trend may reflect the

significantly higher expression of *Grin2b*, which encodes GluN2B (NR2B), in the bilateral

ISO group.

Collectively, these ORA and GSEA results consistently demonstrate that distinct

molecular mechanisms underpin the effects of environmental treatments on the left and right

hippocampal CA1 regions. The left CA1 region appears to promote structural plasticity, such

as the remodeling of synaptic ultrastructure, through coordinated metabolic activity, whereas

the right CA1 region preferentially engages endogenous gene regulatory pathways that may

suppress experience-driven neural plasticity under enriched environmental conditions. These

findings are in line with functional evidence demonstrating that environmental experience

enhances interhemispheric asymmetry in hippocampal gamma oscillations [9]. The molecular

pathways identified in this study may therefore constitute a biological substrate for such

experience-dependent functional lateralization.

Commonality of environment-induced differential gene expression between the left and right CA1 regions

Co-expression and PPI network analyses of environmentally responsive DEGs in the

left and right CA1 regions revealed a shared molecular framework in both areas (Figs 5A, 6A,

and 7A). Within the co-expression network, IEGs, such as *Fos*, *Junb*, *Egr1*, and *Btg2,* formed

central clusters in both left and right CA1 regions. This suggests that rapid transcriptional

responses to environmental stimuli and early memory formation are common initial

responses in both hemispheres [50,51].

Beyond IEGs, in the PPI network, molecules involved in neural development and

plasticity, specifically Grin2b, Mecp2, Cdkl5, and Tet3, formed functionally connected

clusters in both the left and right CA1 regions (Fig 5B, 6B, and 7 B). This

Grin2b–Mecp2–Cdkl5–Tet3 cluster has been implicated in the maintenance of synaptic

function, epigenetic regulation, and neurodevelopmental disorders [52-54], and likely

constitutes a core basis for environment-dependent gene regulation common to both the left

and right CA1 regions. Indeed, our ORA results (Fig 3) showed a significant enrichment of

GO items related to synaptic structure in the left CA1 region and transcriptional regulation,

including DNA 5-methylcytosine activity in the right CA1 region. The network centered on

this cluster may function as a molecular interface that integrates the responses of the two

hemispheres through epigenetic control-based neural regulation. The GSEA results using the

Reactome Pathway partly support this speculation: the left CA1 region shows enrichment of

the gene set "transcriptional regulation by MECP2" in the ISO group (S8 Fig). However,

further studies are needed to confirm this hypothesis.

Furthermore, the ORA of the shared DEGs between the left and right CA1 regions in

response to environmental exposure (Fig 4) revealed the enrichment of terms related to

transcriptional regulation, NMDA receptor activity, synaptic structure, and cellular signaling

pathways. Notably, categories such as "sequence-specific DNA binding" and "transcription

regulator activity" underscore the activation of shared transcriptional networks involving

IEGs such as *Fos*, *Junb*, and *Egr1*, observed in both co-expression and PPI network analyses.

These results indicate the presence of a common transcriptional response module in response

to environmental stimuli in both hemispheres. Enrichment of terms such as "NMDA

glutamate receptor activity" and "presynaptic membrane" further supports the shared

activation of neural plasticity-related pathways, consistent with the GSEA results showing

synaptic-related processes. Moreover, the detection of pathways such as the "calcium

signaling pathway" and the involvement of miR-132-3p—both known to be linked to

activity-dependent synaptic remodeling and long-term transcriptional control

[45,55]—suggest that core molecular mechanisms are symmetrically recruited in

experience-dependent circuit reorganization across the bilateral CA1 regions.

In summary, while the left and right CA1 regions employ distinct molecular

strategies (e.g., structural vs. transcriptional emphasis), they may share a common

foundational response. This shared basis is evident in acute IEG responses and overlapping

networks integral to neuroplasticity and epigenetic regulation. Therefore, brain laterality

should not be considered a complete dichotomy but rather a graded asymmetry, where diverse

response patterns are superimposed on a shared core mechanism. These findings, together

with earlier network-based analyses, highlight that despite hemispheric asymmetries, a

conserved molecular framework likely orchestrates transcriptional and synaptic responses to

environmental input.

Limitations and perspectives

Careful consideration of sex-specific differences in lateralization is required.

Following a previous study, we used male rats to exclude the potential influence of female

sex hormones on lateralization [9]. However, hippocampal asymmetry has also been observed

in female rats [56,57]. Recent studies have revealed sex-related differences in NMDA

receptor function and synaptic plasticity [58,59]. For example, estrogen modulates NMDA

receptor signaling and long-term potentiation in females, whereas males may rely more on

GluN2B-mediated pathways [60]. These molecular differences may underlie the distinct

patterns of hippocampal asymmetry and memory processing between sexes, highlighting the

importance of including both sexes in future studies.

Although the present study used the 4-week period following weaning as the

window for environmental exposure, the appropriateness of this timeframe remains unclear.

Postnatal brain development, including that of the hippocampus, involves a critical period

that profoundly influences learning and experience [61,62]. Previous studies on mice have

also examined the relationship between the duration of environmental enrichment and

subsequent structural and functional changes, as well as alterations in hippocampal gene

expression [18]. To gain a more comprehensive understanding, future research should

incorporate cross-sectional studies across diverse developmental stages, including old age,

and longitudinal studies spanning multiple life phases.

The statistical power of the current study for detecting DEGs may be insufficient

because each group included only three animals. Experimental validation using RT-qPCR

was performed for some DEGs; however, more detailed and robust statistical analyses would

require larger sample sizes. For example, in the left–right comparison, only one DEG, *Pkml1*,

was identified in the ISO group. The limited sample size raises concerns about reduced

sensitivity; therefore, it is difficult to determine based on this study alone whether the limited

detection of significant genes in the left–right comparison is due to insufficient statistical

power or to underlying biological factors. Additionally, to compensate for the reduced

sensitivity due to the small sample size, a less stringent statistical criterion was used for DEG

detection, raising concerns about an increased false-positive rate. Thus, large-scale studies

with adequate statistical power are essential for robust and reliable identification of

lateralized and environment-induced gene expression profiles.

As suggested by the enrichment analysis results, future investigations on the

molecular-level left–right differences should extend beyond the mRNA layer. NMDA

receptor subunit genes, including *Grin2b*, contribute to neural functional diversity and precise

formation and maintenance of neural circuits through splice variation arising from alternative

exon usage [63]. Additionally, *Mecp2*, which was highlighted by network analysis, is

involved not only in epigenetic regulation in the brain but also in the regulation of alternative

splicing [64]. Furthermore, isoform-specific alterations in Mecp2 protein levels have been

implicated in Rett syndrome [65]. To further advance our findings, the identification of novel

isoforms through long-read sequencing technologies and the accurate quantification of splice

variants using deep sequencing, together with genome-wide epigenetic profiling, is required.

Finally, this study employed bulk RNA-seq analysis of the entire CA1 region. While

this approach provides an overall transcriptomic profile, it inherently averages gene

expression across diverse cell populations and therefore cannot distinguish between gene

expression changes occurring within specific cell types (e.g., pyramidal neurons or

astrocytes) and those arising from alterations in cellular composition. Future studies utilizing

single-cell RNA sequencing (scRNA-seq) or spatial transcriptomics will be essential to

elucidate cell-type-specific and spatially resolved responses to environmental stimuli in the

bilateral hippocampus.

Acknowledgments

627 The authors thank Dr. Takatoshi Ueki and Dr. Hajime Hirase for their kind advice. The authors

628 thank Ms. Yoko Iwauchi for technical assistance. We would like to thank Editage

629 (www.editage.jp) for English language editing.

References

1. Corballis MC. The evolution and genetics of cerebral asymmetry. Philos Trans R Soc

Lond B Biol Sci. 2009;364(1519):867-79. doi: 10.1098/rstb.2008.0232. PubMed PMID:

19064358; PubMed Central PMCID: PMCPMC2666079.

2. Corballis MC. Left brain, right brain: facts and fantasies. PLoS Biol.

2014;12(1):e1001767. Epub 20140121. doi: 10.1371/journal.pbio.1001767. PubMed

PMID: 24465175; PubMed Central PMCID: PMCPMC3897366.

3.   Morillon B, Lehongre K, Frackowiak RS, Ducorps A, Kleinschmidt A, Poeppel D, et al.

Neurophysiological origin of human brain asymmetry for speech and language. Proc

Natl Acad Sci U S A. 2010;107(43):18688-93. Epub 20101018. doi:

10.1073/pnas.1007189107. PubMed PMID: 20956297; PubMed Central PMCID:

PMCPMC2972980.

4.   Nonaka S, Shiratori H, Saijoh Y, Hamada H. Determination of left-right patterning of the

mouse embryo by artificial nodal flow. Nature. 2002;418(6893):96-9. doi:

10.1038/nature00849. PubMed PMID: 12097914.

5.   Kawakami R, Dobi A, Shigemoto R, Ito I. Right isomerism of the brain in inversus

viscerum mutant mice. PLoS One. 2008;3(4):e1945. Epub 20080416. doi:

10.1371/journal.pone.0001945. PubMed PMID: 18414654; PubMed Central PMCID:

PMCPMC2291575.

6.   Goto K, Kurashima R, Gokan H, Inoue N, Ito I, Watanabe S. Left-right asymmetry

defect in the hippocampal circuitry impairs spatial learning and working memory in iv

mice. PLoS One. 2010;5(11):e15468. Epub 20101117. doi:

10.1371/journal.pone.0015468. PubMed PMID: 21103351; PubMed Central PMCID:

PMCPMC2984506.

7.   Kawakami R, Shinohara Y, Kato Y, Sugiyama H, Shigemoto R, Ito I. Asymmetrical

allocation of NMDA receptor epsilon2 subunits in hippocampal circuitry. Science.

2003;300(5621):990-4. doi: 10.1126/science.1082609. PubMed PMID: 12738868.

8.   Shinohara Y, Hirase H, Watanabe M, Itakura M, Takahashi M, Shigemoto R. Left-right

asymmetry of the hippocampal synapses with differential subunit allocation of glutamate

receptors. Proc Natl Acad Sci U S A. 2008;105(49):19498-503. Epub 20081203. doi:

10.1073/pnas.0807461105. PubMed PMID: 19052236; PubMed Central PMCID:

PMCPMC2593619.

9.   Shinohara Y, Hosoya A, Hirase H. Experience enhances gamma oscillations and

interhemispheric asymmetry in the hippocampus. Nat Commun. 2013;4:1652. doi:

10.1038/ncomms2658. PubMed PMID: 23552067; PubMed Central PMCID:

PMCPMC3644069.

10.   Feng T, Alicea C, Pham V, Kirk A, Pieraut S. Experience-Dependent Inhibitory Plasticity

Is Mediated by CCK+ Basket Cells in the Developing Dentate Gyrus. J Neurosci.

2021;41(21):4607-19. Epub 20210427. doi: 10.1523/JNEUROSCI.1207-20.2021.

PubMed PMID: 33906898; PubMed Central PMCID: PMCPMC8260240.

11.   Biggio F, Mostallino MC, Talani G, Locci V, Mostallino R, Calandra G, et al. Social

enrichment reverses the isolation-induced deficits of neuronal plasticity in the

hippocampus of male rats. Neuropharmacology. 2019;151:45-54. Epub 20190329. doi:

10.1016/j.neuropharm.2019.03.030. PubMed PMID: 30935859.

12. Gunturkun O, Ocklenburg S. Ontogenesis of Lateralization. Neuron. 2017;94(2):249-63.

doi: 10.1016/j.neuron.2017.02.045. PubMed PMID: 28426959.

13. Gunturkun O, Strockens F, Ocklenburg S. Brain Lateralization: A Comparative

Perspective. Physiol Rev. 2020;100(3):1019-63. doi: 10.1152/physrev.00006.2019.

PubMed PMID: 32233912.

14. Moskal JR, Kroes RA, Otto NJ, Rahimi O, Claiborne BJ. Distinct patterns of gene

expression in the left and right hippocampal formation of developing rats. Hippocampus.

2006;16(8):629-34. doi: 10.1002/hipo.20198. PubMed PMID: 16847945.

15. Shinohara Y, Yahagi K, Kawano M, Nishiyori H, Kawazu C, Suzuki N, et al. miRNA

profiling of bilateral rat hippocampal CA3 by deep sequencing. Biochem Biophys Res

Commun. 2011;409(2):293-8. Epub 20110514. doi: 10.1016/j.bbrc.2011.05.004.

PubMed PMID: 21575607.

16. Rampon C, Jiang CH, Dong H, Tang YP, Lockhart DJ, Schultz PG, et al. Effects of

environmental enrichment on gene expression in the brain. Proc Natl Acad Sci U S A.

2000;97(23):12880-4. doi: 10.1073/pnas.97.23.12880. PubMed PMID: 11070096;

PubMed Central PMCID: PMCPMC18858.

17. Huttenrauch M, Salinas G, Wirths O. Effects of Long-Term Environmental Enrichment

on Anxiety, Memory, Hippocampal Plasticity and Overall Brain Gene Expression in

C57BL6 Mice. Front Mol Neurosci. 2016;9:62. Epub 20160803. doi:

10.3389/fnmol.2016.00062. PubMed PMID: 27536216; PubMed Central PMCID:

PMCPMC4971077.

18. Singhal G, Morgan J, Jawahar MC, Corrigan F, Jaehne EJ, Toben C, et al. The effects of

short-term and long-term environmental enrichment on locomotion, mood-like behavior,

cognition and hippocampal gene expression. Behav Brain Res. 2019;368:111917. Epub

20190417. doi: 10.1016/j.bbr.2019.111917. PubMed PMID: 31004685.

19. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018;34(17):i884-i90. doi: 10.1093/bioinformatics/bty560. PubMed

PMID: 30423086; PubMed Central PMCID: PMCPMC6129281.

20. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and

bias-aware quantification of transcript expression. Nat Methods. 2017;14(4):417-9. Epub

20170306. doi: 10.1038/nmeth.4197. PubMed PMID: 28263959; PubMed Central

PMCID: PMCPMC5600148.

21. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level

estimates improve gene-level inferences. F1000Res. 2015;4:1521. Epub 20151230. doi:

10.12688/f1000research.7563.2. PubMed PMID: 26925227; PubMed Central PMCID:

PMCPMC4712774.

22. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139-40.

Epub 20091111. doi: 10.1093/bioinformatics/btp616. PubMed PMID: 19910308;

PubMed Central PMCID: PMCPMC2796818.

23. Chen Y, Lun AT, Smyth GK. From reads to genes to pathways: differential expression

analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood

pipeline. F1000Res. 2016;5:1438. Epub 20160620. doi: 10.12688/f1000research.8987.2.

PubMed PMID: 27508061; PubMed Central PMCID: PMCPMC4934518.

24. Chen Y, Chen L, Lun ATL, Baldoni PL, Smyth GK. edgeR v4: powerful differential

analysis of sequencing data with expanded functionality and improved support for small

counts and larger datasets. Nucleic Acids Res. 2025;53(2). doi: 10.1093/nar/gkaf018.

PubMed PMID: 39844453; PubMed Central PMCID: PMCPMC11754124.

25. Bardou P, Mariette J, Escudié F, Djemiel C, Klopp C. jvenn: an interactive Venn diagram

viewer. BMC Bioinformatics. 2014;15(1):293. Epub 20140829. doi:

10.1186/1471-2105-15-293. PubMed PMID: 25176396; PubMed Central PMCID:

PMCPMC4261873.

26. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25(4):402-8. doi:

10.1006/meth.2001.1262. PubMed PMID: 11846609.

27. Kubota A, Tajima A. Click-qPCR: A Simple Tool for Interactive qPCR Data Analysis.

Bio-protocol 15(22):e5513. doi: 10.21769/BioProtoc.5513.

28. Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. g:Profiler-interoperable

web service for functional enrichment analysis and gene identifier mapping (2023

update). Nucleic Acids Res. 2023;51(W1):W207-W12. doi: 10.1093/nar/gkad347.

PubMed PMID: 37144459; PubMed Central PMCID: PMCPMC10320099.

29. Gene Ontology C, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, et al.

The Gene Ontology knowledgebase in 2023. Genetics. 2023;224(1). doi:

10.1093/genetics/iyad031. PubMed PMID: 36866529; PubMed Central PMCID:

PMCPMC10158837.

30. Milacic M, Beavers D, Conley P, Gong C, Gillespie M, Griss J, et al. The Reactome

Pathway Knowledgebase 2024. Nucleic Acids Res. 2024;52(D1):D672-D8. doi:

10.1093/nar/gkad1025. PubMed PMID: 37941124; PubMed Central PMCID:

PMCPMC10767911.

31. Kanehisa M, Furumichi M, Sato Y, Matsuura Y, Ishiguro-Watanabe M. KEGG:

biological systems database as a model of the real world. Nucleic Acids Res.

2025;53(D1):D672-D7. doi: 10.1093/nar/gkae909. PubMed PMID: 39417505; PubMed

Central PMCID: PMCPMC11701520.

32. Matys V, Kel-Margoulis OV, Fricke E, Liebich I, Land S, Barre-Dirrie A, et al.

TRANSFAC and its module TRANSCompel: transcriptional gene regulation in

eukaryotes. Nucleic Acids Res. 2006;34(Database issue):D108-10. doi:

10.1093/nar/gkj143. PubMed PMID: 16381825; PubMed Central PMCID:

PMCPMC1347505.

33. Cui S, Yu S, Huang HY, Lin YC, Huang Y, Zhang B, et al. miRTarBase 2025: updates to

the collection of experimentally validated microRNA-target interactions. Nucleic Acids

Res. 2025;53(D1):D147-D56. doi: 10.1093/nar/gkae1072. PubMed PMID: 39578692;

PubMed Central PMCID: PMCPMC11701613.

34. Reimand J, Kull M, Peterson H, Hansen J, Vilo J. g:Profiler--a web-based toolset for

functional profiling of gene lists from large-scale experiments. Nucleic Acids Res.

2007;35(Web Server issue):W193-200. Epub 20070503. doi: 10.1093/nar/gkm226.

PubMed PMID: 17478515; PubMed Central PMCID: PMCPMC1933153.

35. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a

software environment for integrated models of biomolecular interaction networks.

Genome Res. 2003;13(11):2498-504. doi: 10.1101/gr.1239303. PubMed PMID:

14597658; PubMed Central PMCID: PMCPMC403769.

36. Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, et al. The

GeneMANIA prediction server: biological network integration for gene prioritization

and predicting gene function. Nucleic Acids Res. 2010;38(Web Server issue):W214-20.

doi: 10.1093/nar/gkq537. PubMed PMID: 20576703; PubMed Central PMCID:

PMCPMC2896186.

37. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The

STRING database in 2023: protein-protein association networks and functional

enrichment analyses for any sequenced genome of interest. Nucleic Acids Res.

2023;51(D1):D638-D46. doi: 10.1093/nar/gkac1000. PubMed PMID: 36370105;

PubMed Central PMCID: PMCPMC9825434.

38. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.

Gene set enrichment analysis: a knowledge-based approach for interpreting

genome-wide expression profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50.

Epub 20050930. doi: 10.1073/pnas.0506580102. PubMed PMID: 16199517; PubMed

Central PMCID: PMCPMC1239896.

39. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological

themes among gene clusters. OMICS. 2012;16(5):284-7. Epub 20120328. doi:

10.1089/omi.2011.0118. PubMed PMID: 22455463; PubMed Central PMCID:

PMCPMC3339379.

40. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal

enrichment tool for interpreting omics data. Innovation (Camb). 2021;2(3):100141. Epub

20210701. doi: 10.1016/j.xinn.2021.100141. PubMed PMID: 34557778; PubMed

Central PMCID: PMCPMC8454663.

41. Xu S, Hu E, Cai Y, Xie Z, Luo X, Zhan L, et al. Using clusterProfiler to characterize

multiomics data. Nat Protoc. 2024;19(11):3292-320. Epub 20240717. doi:

10.1038/s41596-024-01020-z. PubMed PMID: 39019974.

42. Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast gene

set enrichment analysis. bioRxiv. 2021:060012. doi: 10.1101/060012.

43. Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP.

Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011;27(12):1739-40.

Epub 20110505. doi: 10.1093/bioinformatics/btr260. PubMed PMID: 21546393;

PubMed Central PMCID: PMCPMC3106198.

44. Swiss VA, Casaccia P. Cell-context specific role of the E2F/Rb pathway in development

and disease. Glia. 2010;58(4):377-90. doi: 10.1002/glia.20933. PubMed PMID:

19795505; PubMed Central PMCID: PMCPMC2882865.

45. Qian Y, Song J, Ouyang Y, Han Q, Chen W, Zhao X, et al. Advances in Roles of

miR-132 in the Nervous System. Front Pharmacol. 2017;8:770. Epub 20171025. doi:

10.3389/fphar.2017.00770. PubMed PMID: 29118714; PubMed Central PMCID:

PMCPMC5660991.

46. Jordan JT. The rodent hippocampus as a bilateral structure: A review of hemispheric

lateralization.   Hippocampus.   2020;30(3):278-92.   Epub   20191227.   doi:

10.1002/hipo.23188. PubMed PMID: 31880377.

47. Fu J, Guo O, Zhen Z, Zhen J. Essential Functions of the Transcription Factor Npas4 in

Neural Circuit Development, Plasticity, and Diseases. Front Neurosci. 2020;14:603373.

Epub 20201201. doi: 10.3389/fnins.2020.603373. PubMed PMID: 33335473; PubMed

Central PMCID: PMCPMC7736240.

48. Zhao X, Sun L, Jia H, Meng Q, Wu S, Li N, et al. Isolation rearing induces social and

emotional function abnormalities and alters glutamate and neurodevelopment-related

gene expression in rats. Prog Neuropsychopharmacol Biol Psychiatry. 2009;33(7):1173-7.

Epub 20090627. doi: 10.1016/j.pnpbp.2009.06.016. PubMed PMID: 19563853.

49. Chang CH, Hsiao YH, Chen YW, Yu YJ, Gean PW. Social isolation-induced increase in

NMDA receptors in the hippocampus exacerbates emotional dysregulation in mice.

Hippocampus. 2015;25(4):474-85. Epub 20141105. doi: 10.1002/hipo.22384. PubMed

PMID: 25348768.

50. Lara Aparicio SY, Laureani Fierro AJ, Aranda Abreu GE, Toledo Cardenas R, Garcia

Hernandez LI, Coria Avila GA, et al. Current Opinion on the Use of c-Fos in

Neuroscience. NeuroSci. 2022;3(4):687-702. Epub 20221219. doi:

10.3390/neurosci3040050. PubMed PMID: 39483772; PubMed Central PMCID:

PMCPMC11523728.

51. Tregub PP, Komleva YK, Kukla MV, Averchuk AS, Vetchinova AS, Rozanova NA, et al.

Brain Plasticity and Cell Competition: Immediate Early Genes Are the Focus. Cells.

2025;14(2). Epub 20250119. doi: 10.3390/cells14020143. PubMed PMID: 39851571;

PubMed Central PMCID: PMCPMC11763428.

52. Lee S, Kim W, Ham BJ, Chen W, Bear MF, Yoon BJ. Activity-dependent NR2B

expression is mediated by MeCP2-dependent epigenetic regulation. Biochem Biophys

Res Commun. 2008;377(3):930-4. Epub 20081024. doi: 10.1016/j.bbrc.2008.10.082.

PubMed PMID: 18952054.

53. Rao VK, Swarnaseetha A, Tham GH, Lin WQ, Han BB, Benoukraf T, et al.

Phosphorylation of Tet3 by cdk5 is critical for robust activation of BRN2 during

neuronal differentiation. Nucleic Acids Res. 2020;48(3):1225-38. doi:

10.1093/nar/gkz1144. PubMed PMID: 31807777; PubMed Central PMCID:

PMCPMC7026633.

54. Zito A, Lee JT. Variable expression of MECP2, CDKL5, and FMR1 in the human brain:

Implications for gene restorative therapies. Proc Natl Acad Sci U S A.

2024;121(9):e2312757121. Epub 20240222. doi: 10.1073/pnas.2312757121. PubMed

PMID: 38386709; PubMed Central PMCID: PMCPMC10907246.

55. Yasuda R, Hayashi Y, Hell JW. CaMKII: a central molecular organizer of synaptic

plasticity, learning and memory. Nat Rev Neurosci. 2022;23(11):666-82. Epub 20220902.

doi: 10.1038/s41583-022-00624-2. PubMed PMID: 36056211.

56. Diamond MC, Murphy GM, Jr., Akiyama K, Johnson RE. Morphologic hippocampal

asymmetry in male and female rats. Exp Neurol. 1982;76(3):553-65. doi:

10.1016/0014-4886(82)90124-8. PubMed PMID: 7084374.

57. Wolff SC, Hruska Z, Nguyen L, Dohanich GP. Asymmetrical distributions of muscarinic

receptor binding in the hippocampus of female rats. Eur J Pharmacol.

2008;588(2-3):248-50. Epub 20080409. doi: 10.1016/j.ejphar.2008.04.002. PubMed

PMID: 18499095.

58. Yagi S, Galea LAM. Sex differences in hippocampal cognition and neurogenesis.

Neuropsychopharmacology. 2019;44(1):200-13. Epub 20180907. doi:

10.1038/s41386-018-0208-4. PubMed PMID: 30214058; PubMed Central PMCID:

PMCPMC6235970.

59. Narattil NR, Maroun M. Differential role of NMDA receptors in hippocampal-dependent

spatial memory and plasticity in juvenile male and female rats. Hippocampus.

2024;34(11):564-74. Epub 20240815. doi: 10.1002/hipo.23631. PubMed PMID:

39143939.

60. Copenhaver AE, LeGates TA. Sex-Specific Mechanisms Underlie Long-Term

Potentiation at Hippocampus-->Medium Spiny Neuron Synapses in the Medial Shell of

the Nucleus Accumbens. J Neurosci. 2024;44(27). Epub 20240703. doi:

10.1523/JNEUROSCI.0100-24.2024. PubMed PMID: 38806250; PubMed Central

PMCID: PMCPMC11223474.

61. Sakai A, Sugiyama S. Experience-dependent transcriptional regulation in juvenile brain

development. Dev Growth Differ. 2018;60(8):473-82. doi: 10.1111/dgd.12571. PubMed

PMID: 30368782.

62. Sakimoto Y, Shintani A, Yoshiura D, Goshima M, Kida H, Mitsushima D. A critical

period for learning and plastic changes at hippocampal CA1 synapses. Sci Rep.

2022;12(1):7199. Epub 20220503. doi: 10.1038/s41598-022-10453-z. PubMed PMID:

35504922; PubMed Central PMCID: PMCPMC9065057.

63. Tumdam R, Hussein Y, Garin-Shkolnik T, Stern S. NMDA Receptors in

Neurodevelopmental Disorders: Pathophysiology and Disease Models. Int J Mol Sci.

2024;25(22). Epub 20241118. doi: 10.3390/ijms252212366. PubMed PMID: 39596430;

PubMed Central PMCID: PMCPMC11594297.

64. Liu Y, Whitfield TW, Bell GW, Guo R, Flamier A, Young RA, et al. Exploring the

complexity of MECP2 function in Rett syndrome. Nat Rev Neurosci. 2025;26(7):379-98.

Epub 20250513. doi: 10.1038/s41583-025-00926-1. PubMed PMID: 40360671.

65. Rodrigues DC, Mufteev M, Ellis J. Regulation, diversity and function of MECP2 exon

and   3'UTR   isoforms.   Hum   Mol   Genet.   2020;29(R1):R89-R99.   doi:

10.1093/hmg/ddaa154.   PubMed   PMID: 32681172;   PubMed   Central   PMCID:

PMCPMC7530521.

Supporting information

S1 Fig. Brief workflow chart of this study.

S2 Fig. Multivariate analyses using the TPM-normalized gene count data.

S3 Fig. Volcano plot visualizations of the left–right comparison.

S4 Fig. Correlation plots focusing on the $\log_2$FC values calculated using edgeR.

S5 Fig. GSEA for the environmental comparisons in the left and right CA1 regions

using the GO-BP database.

S6 Fig. GSEA for the environmental comparisons in the left and right CA1 regions

using the GO-CC database.

S7 Fig. GSEA for the environmental comparisons in the left and right CA1 regions

using the GO-MF database.

S8 Fig. GSEA for the environmental comparisons in the left and right CA1 regions

using the MSigDB's C2 Reactome Pathway database.

S9 Fig. GSEA for the environmental comparisons in the left and right CA1 regions

using the MSigDB's C2 KEGG Pathway database.

S1 Data. Raw gene count data for the edgeR analysis.

S2 Data. TPM-normalized gene count data generated by tximport.

S3 Data. The edgeR calculation results of the left–right comparison.

S4 Data. The edgeR calculation results of the ISO–ENR comparison.

S5 Data. The g:Profiler calculation results of over-representation analysis.

S6 Data. The Cytoscape calculation results of network analysis.

S7 Data. The clusterProfiler calculation results of gene set enrichment analysis.

Fig 1

A

ID	Source	Term ID	Term Name	p_{adj} (query_1)
1	GO:MF	GO:0043565	sequence-specific DNA binding	5.082×10^{-5}
2	GO:MF	GO:0003700	DNA-binding transcription factor activity	6.695×10^{-5}
3	GO:MF	GO:0004888	transmembrane signaling receptor activity	3.594×10^{-4}
4	GO:MF	GO:0022849	glutamate-gated calcium ion channel activity	9.003×10^{-3}
5	GO:MF	GO:0022824	transmitter-gated monoatomic ion channel activity	3.432×10^{-2}
6	GO:MF	GO:0022836	gated channel activity	3.611×10^{-2}
7	GO:MF	GO:0015171	amino acid transmembrane transporter activity	4.677×10^{-2}
8	GO:BP	GO:0065007	biological regulation	4.369×10^{-9}
9	GO:BP	GO:0014706	striated muscle tissue development	1.496×10^{-3}
10	GO:BP	GO:0034330	cell junction organization	5.691×10^{-3}
11	GO:BP	GO:0035418	protein localization to synapse	1.167×10^{-2}
12	GO:BP	GO:0051716	cellular response to stimulus	2.319×10^{-2}
13	GO:BP	GO:0035914	skeletal muscle cell differentiation	2.828×10^{-2}
14	GO:BP	GO:0098609	cell-cell adhesion	4.073×10^{-2}
15	GO:BP	GO:0007616	long-term memory	4.203×10^{-2}
16	GO:CC	GO:0097060	synaptic membrane	5.815×10^{-9}
17	GO:CC	GO:0043235	receptor complex	1.869×10^{-3}
18	GO:CC	GO:0045121	membrane raft	1.725×10^{-2}
19	GO:CC	GO:0098685	Schaffer collateral - CA1 synapse	2.082×10^{-2}
20	GO:CC	GO:0043194	axon initial segment	2.192×10^{-2}
21	GO:CC	GO:0009986	cell surface	3.381×10^{-2}
22	KEGG	KEGG:04010	MAPK signaling pathway	5.685×10^{-5}
23	REAC	REAC:R-RNO-383280	Nuclear Receptor transcription pathway	3.362×10^{-2}
24	TF	TF:M00803_1	Factor: E2F; motif: GGCGSG; match class: 1	2.416×10^{-8}

Left ISO_ENR: Of the 189 Ensembl gene ID-based DEGs,180 annotated genes were used.

B

ID	Source	Term ID	Term Name	p_{adj} (query_1)
1	GO:MF	GO:0019199	transmembrane receptor protein kinase activity	1.831×10^{-4}
2	GO:MF	GO:0070579	DNA 5-methylcytosine dioxygenase activity	2.535×10^{-2}
3	GO:MF	GO:0019838	growth factor binding	2.813×10^{-2}
4	GO:BP	GO:0065007	biological regulation	7.802×10^{-5}
5	GO:BP	GO:0051252	regulation of RNA metabolic process	2.471×10^{-4}
6	GO:CC	GO:0030054	cell junction	1.014×10^{-4}
7	GO:CC	GO:0043235	receptor complex	4.075×10^{-4}
8	GO:CC	GO:0098839	postsynaptic density membrane	3.638×10^{-2}
9	GO:CC	GO:0071944	cell periphery	3.965×10^{-2}
10	REAC	REAC:R-RNO-9768919	NPAS4 regulates expression of target genes	3.533×10^{-2}
11	REAC	REAC:R-RNO-9634815	Transcriptional Regulation by NPAS4	3.533×10^{-2}
12	TF	TF:M00803_1	Factor: E2F; motif: GGCGSG; match class: 1	6.074×10^{-3}
13	MIRNA	MIRNA:rno-mir-132-3p	rno-mir-132-3p	4.999×10^{-2}

Right ISO-ENR: Of the 94 Ensembl gene ID-based DEGs, 87 annotated genes were used.

ID	Source	Term ID	Term Name	p_{adj} (query_1)
1	GO:MF	GO:0043565	sequence-specific DNA binding	9.048×10^{-3}
2	GO:MF	GO:0004714	transmembrane receptor protein tyrosine kinase activity	9.686×10^{-3}
3	GO:MF	GO:0000976	transcription cis-regulatory region binding	1.696×10^{-2}
4	GO:MF	GO:0140110	transcription regulator activity	3.982×10^{-2}
5	GO:MF	GO:0004972	NMDA glutamate receptor activity	4.988×10^{-2}
6	GO:BP	GO:0045935	positive regulation of nucleobase-containing compound metabolic process	7.383×10^{-5}
7	GO:BP	GO:0065007	biological regulation	1.105×10^{-2}
8	GO:BP	GO:0014706	striated muscle tissue development	4.992×10^{-2}
9	GO:CC	GO:0042734	presynaptic membrane	7.808×10^{-5}
10	GO:CC	GO:0043235	receptor complex	3.274×10^{-3}
11	KEGG	KEGG:04020	Calcium signaling pathway	6.858×10^{-4}
12	REAC	REAC:R-RNO-426496	Post-transcriptional silencing by small RNAs	4.979×10^{-2}
13	MIRNA	MIRNA:rno-mir-132-3p	rno-mir-132-3p	4.999×10^{-2}
14	TF	TF:M00803_1	Factor: E2F; motif: GGCGSG; match class: 1	2.737×10^{-3}
15	TF	TF:M00244	Factor: NGFI-C; motif: WTGCGTGGGYGG	3.668×10^{-3}

Of the 70 Ensembl gene IDs as shared DEGs, 64 annotated gene symbols were used.

A

B