No association between metabolic rate and an exaggerated sexually

selected trait.

Kentarou Matsumura[1,4], Tomohito Noda[1,2,3], Manmohan D. Sharma[1], Kensuke Okada[4],

Takahisa Miyatake[4] and David J. Hosken[1*]

[1]Centre for Ecology and Conservation, University of Exeter, Cornwall, Tremough, Penryn

TR10 9EZ, UK.

[2]Department of Biological Sciences, Graduate School of Science, The University of
Tokyo, Tokyo, Japan

[3]Molecular Biosystem Research Institute, National Institute of Advanced Industrial
Science and Technology (AIST), Tsukuba, Japan

[4]Faculty of Environmental, Life, Natural Science and Technology, Okayama

University, Okayama 700-8530, Japan.

*corresponding author – d.j.hosken@exeter.ac.uk

Acknowledgements

We thank many members of FlyLab who helped in various ways with this study,

especially Sarah Daisy Kendall for help with the imaging and Ruth Archer for support.

DJH and MDS were funded by the Leverhulme Trust.

Conflict of interests

The authors have no conflicts of interest to declare and have not used AI in the

preparation of this manuscript.

Ethics

This study has ethical approval from the Ethics Committee of the Centre for Ecology

and Conservation, University of Exeter.

Key words: sexual selection, costs, energetics, weapon, genetics

Lay Summary

The large weapons used in competition between males are thought to be energetically costly. We investigate these potential costs by testing for associations between metabolic rate and weapon size in a beetle. We found no relationship between weapons and metabolism, indicating weapons are not always energetically costly.

Abstract

Exaggerated sexually selected traits are thought to be energetically costly to produce and to maintain. This is especially true of sexually selected weapons, traits that are frequently positively allometric, and therefore likely to be especially energetically costly. However, there have been few studies that directly measure these putative costs. Here we tested for the associations between resting metabolic rate, during development and adulthood, and the size of a weapon used during adult male-male competition in the beetle *Gnatocerus cornutus*. We found that metabolic rate was not statistically associated with weapon size, but there were effects of genotype and mass during development (pupal size) on weapons. Additionally, age, mass and genotype all affected metabolic rate in various ways. Despite no evidence of energetic costs to weapon development or maintenance, direct predation costs associated with them have been detected previously, and hence enlarged weapons are costly in this system.

Introduction

Sexual selection is a major mechanism of evolution and has caused some of the most exaggerate behaviours and morphologies in nature (Andersson 1994). While sexually selected traits are typically thought to be costly, evidence for this is somewhat limited, even though understanding putative costs is important because they set limits on trait exaggeration and determine trade-offs with other fitness components (Kotiaho 2001).

Predation costs have been reasonably well documented for sexually selected characters as they can increase conspicuousness to predators (e.g. Endler 1987; Reznick and Endler

1982), and predators and parasitoids frequently target sexually signalling males (e.g.
Ryan 1985; Hosken et al. 1984; Grey and Cade 1999; reviewed in Zuk and Kolluru 1998),
even preferring the same call parameters favoured by females in some cases (Ryan 1985,
2018). Additionally, investing in sexually selected traits can generate immune function
costs (e.g. Folstad and Karter 1992; Salvador et al. 1996; Hosken 2001; McKean and
Nunney 2008), although it is not always clear whether reductions in elements of immune
function have significant fitness impacts (Kotiaho 2001). Other physiological costs of
exaggerated sexually selected traits have arguably been less studied (Halliday 1987),
although energetic costs of sexual signalling have been documented (e.g. Bailey et al.
1993; Kotiaho et al. 1998; Reinhold et al. 1998), and fights can also be energetically
expensive (Hack 1997; Neat et al. 1998). For example, signalling can increase metabolic
rate by more than 50% (e.g. Bailey et al. 1993; Hoback and Wagner 1997), while male
fighting can increase energy expenditure 6 times above resting metabolic rate (Hack
1997).
78
The weapons used in fighting for territories and mates are amongst the most exaggerated
morphological characters, frequently scaling positively with body size, such that large
individuals have disproportionately large weapons – weapons show positive static
allometry (e.g. Eberhard et al. 2018; Kodric-Brown et al. 2006). This positive allometry is
found in the weapons of stag beetles, earwigs and fiddler crabs for example (Kodric-
Brown et al. 2006), and steep scaling relationships at least partly explains why weapons
are thought to be especially costly to produce and maintain (Somjee 2021). Consistent
with this, there can be considerable metabolic cost to morphological-weapon
maintenance, with metabolic expenditure decreasing by about 25% when weapons are
lost (Somjee et al. 2018; also see O'Brian et al. 2019). Additionally, mass specific
metabolic rates are higher in crabs with intact weapons (Allen & Levinton 2007), and more
generally, sexual traits are thought to be metabolically demanding (reviewed in Kotiaho
2001), and this may be especially true of long-lasting morphological characters (Allen &
Levinton 2007). However, it has also been suggested that costs of weapons will depend
on the tissue-types that form them (Dinh 2022; Somjee 2021). For example, rhinoceros
beetle horns are hollow and therefore metabolically inactive, while fiddler crab claws are
full of muscle and therefore energetically expensive (Somjee 2021). Note this relates to

maintenance costs, and even when these costs are low, developmental costs, the costs

of actually building large weapons, can be very significant (Rosenfeld et al. 2015). One

step toward understanding the costs of sexually selected traits like weapons - noting

these costs have to impact fitness to be evolutionarily significant (Kotiaho 2001) - is to

assess the metabolic expenditure associated with them.

Resting metabolic rate (RMR), which measures energetic expenditure on somatic

maintenance (McNab 2005; Rezende et al. 2004; Snodgrass et al. 2005; Sinclair et al.

2011), could influence the resources available for investment in energetically expensive

traits like weapons, with the "residual" energy not needed for basic maintenance being

free to invest in these characters (Rowe and Houle 1996; Reinhold 1999; Tomkins et al.

2004; Ketola and Kotiaho 2009). Thus, RMR could be negatively associated with weapon

size, since lower maintenance costs could generate greater energy pools left to invest in

larger weapons. Alternatively, a high RMR could enable greater energy acquisition and

conversion into weapons, and thus RMR could be positively associated with weapon size

(Bennett and Ruben 1979; Biro and Stamps 2010; Ketola and Kotiaho 2010). In either

case, metabolic rate and weapon development could be linked, but the metabolic costs

of weapons are highly variable (O'Brian et al. 2019), requiring case-by-case assessment.

Here we tested for associations between resting metabolic rate and weapon size in the

broad-horned flour beetle, *Gnatocerus cornutus*. Male beetles have enormously

enlarged mandibles that are used in male-male fights (Okada and Miyatake 2009; Harano

et al. 2010), and females lack this exaggerated character completely (Okada et al. 2006;

Harano et al. 2010; Matsumura et al. 2024). Like many sexually-selected weapons

(Kodric-Brown et al. 2006), the mandibles are positively allometric (Okada and Miyatake

2010; Shinohara et al. in review). Furthermore, males with larger mandibles have higher

fighting and mating success (Okada and Miyatake 2009; Harano et al. 2010; Yamane et

al. 2010; Katsuki et al. 2012), and there are also age-effects on fighting ability and a

genetic component to aggression in younger males (Nishitani et al. 2024). We have

previously documented predation costs to larger mandibles, with the smaller mandibles

evolving in populations subjected to male predation (Okada et al. 2021), and this

predation reduces male (but not female) longevity (Pennell et al. 2024). Thus, there are

direct costs to mandibles that have life-history impacts (c.f. Kotiaho 2001), but it is not

clear whether metabolic rate (MR) affects mandible size in any way. Mandible

development occurs during pupation with a number of gene knockdowns just prior to

pupation able to induce mandible development in females or mandible suppression in

males (Sugiyama et al. 2023). We therefore assessed mandible-size/metabolic rate

associations in both pupae, to assess putative energetic costs to mandible development,

and adults, to test for putative maintenance costs. Additionally, we employed iso-female

lines (Hoffmann and Parson 1989; David et al. 2005) in tests of trait associations. Iso-

female lines are effectively distinct but replicable genotypes, which allowed us to test for

genetic effects in our traits of interest (Hoffmann and Parson 1989; David et al. 2005;

Ingleby et al. 2013; Archer et al. 2024; Nishitani et al. 2024). While we found no

associations between metabolic rate and weapon size, genotype and pupal size

influenced weapon size, and age, mass and genotype all impacted metabolic rate in

various ways.

Materials and methods

The *G. cornutus* beetle culture originated from adults collected in Miyazaki city, Japan, in

1957 (Okada et al. 2006) and has subsequently been maintained in the laboratory of

National Agriculture and Food Research Organisation, Japan. The beetles used in this

study were derived from that culture. They have been reared on 90% organic white flour

(Doves Farm Foods) enriched with 10% brewer's yeast (Brewers' yeast, Thermo Fisher

Scientific). Both rearing and experimentation were conducted in climate chambers

maintained at 27 ℃, 60% relative humidity and with a photoperiod of 14h:10h, light to

dark cycle. These laboratory conditions mimic the natural conditions where *G. cornutus*

was found (Katsuki et al. 2012).

In this study, we used 18 iso-female lines established from the stock population

(Matsumura et al. 2020; Nishitani et al. 2024). To initiate these lines, 18 males and 18

females from the stock population were selected at random and paired. Subsequently,

sib-sib matings within each family were used to propagate isolines for about 30

generations until the present study was conducted. As noted above, isolines represent

distinct genotypes, and significant variation among them indicates genetic variation in

the base population for the character of focus (Hoffmann and Parsons 1989). Previous

work has shown these isolines harbour significant genetic variation for all the traits

investigated even after 40 generations of propagation (e.g. Matsumura et al. 2020;

Nishitani et al. 2024).

Metabolic rate (MR) and trait measurement follow standard procedures. In brief, the

resting metabolic rate, assessed via CO_2 production (see methods in Arnqvist et al. 2010;

Sinclair et al. 2010; Okada et al. 2011), of know-age male pupae (N = 205) and adults (N

= 178) was measured using flow-through respirometry (Sable Systems, Las Vegas, USA)

(Lighton 2008) along with animal mass and used as our indicator of MR. Briefly,

compressed zero-air (79% N_2 and 21% O_2) was first scrubbed of residual CO_2 and water

(Ascarite II and Magnesium Perchlorate) and regulated at a flow rate of 100ml min-1 ±1%

with a Sierra Instruments (Monterey, CA, USA) model 840L mass-flow meter connected

to a mass-flow control unit (Sable Systems Intelligent Mass Flow Control Unit, MFC-2,

Las Vegas, USA). Air was then fed into a recently calibrated LI-COR LI- 7000 infra-red gas

analyzer (IRGA) capable of resolving differences of 1ppm of CO_2 in the air, plumbed in

differential mode (see Lighton 2008). Air exiting the IRGA reference cell was passed

through a Sable Systems (Las Vegas USA) RM8 eight-chamber multiplexer and then

forwarded to the sample cell for measurements. A PC running Sable Systems Expedata

software (ver.1.8.2) and connected to a Sable Systems UI2 analog–digital interface was

used for data acquisition. Within the multiplexer, one empty chamber acted as a negative

control and beetles were haphazardly assigned to other chambers to minimize positional

bias. Baseline measurements were made before and after each set of recordings to test

and control for any drift between calibrations. VCO_2 was measured for 7 minutes/beetle

(with a sample read every second), and data from the most stable section (Lighton 2008)

of the last two minutes of measurement (to control for wash-out) were used to calculate

the average CO_2 production for each individual (ranging from 60-100 seconds of

measurement/beetle). Identifying the stable section was done manually using Expedata

software (ver.1.8.2).

After measurement of metabolic rate, pupae were placed into 24-well plate separately

with food and maintained until they became adults, and RMR was measured again at

adulthood using the methods described above. To avoid an effect of feeding, adult
beetles (10-15 days old: when males are sexually mature) were isolated without food for
two hours before the measurement of MR. All measurements of RMR were conducted
between 12:00 and 19:00 in an incubator maintained at 27°C (SD±0.4°C) and RMR was
repeatable: the intraclass correlation coefficient (ICC), estimated using the irr R package
(vers. 0.84.1; Gamer et al. 2019) for 49 beetles measured thrice was 92% (ICC(C,3) =
.920, 95% CI = [.872, .952] estimated using an average-score, consistency-based, two-
way random effects model: $F_{48,96}$ = 12.5; P < 0.001). This is consistent with previous
repeatability tests (Matsumura et al. 2025).

Following measurement of adult MR, mandible size was measured using standard
approaches (e.g. Okada & Miyatake 2009; Harano et al. 2010; Okada et al. 2014). Briefly,
male beetles were euthanized by freezing and stored at -20°C until processing for
morphological measurement. Each beetle was placed on a slide and oriented
consistently under a stereomicroscope (Leica M125) to ensure standardized imaging.
Digital images of the whole body were captured using a mounted digital camera (Leica
DFC 295) connected to a PC running ImageJ software (version 1.48). Images were
calibrated to scale using a 1000μm reference standard. Mandible length, defined as the
distance from the base to the tip of the mandible was quantified using ImageJ's
measurement tools. All measurements were taken by a single observer to minimize
variation, and repeatability was confirmed through blind re-measurements of a subset of
individuals (repeatability > 0.95, following Okada & Miyatake 2009).

Statistics
Visual inspection of the data did not indicate significant violations of normality (and see
Kneif & Forstmeier 2021; Hosken et al. 2018), and although the data-set contained one
very small individual, removal of this datum did not change model outcomes (we
therefore present analyses that include this male). We use MANOVA, mixed model
GLMs and least squares linear regression to test for association and all analyses were
carried out in SPSS Version 29.0.2.0 (20) (R-code can be provided on request). Isoline
was entered as a random effect in GLMs, but outcomes are unchanged if it was treated
as a fixed effect. Additionally, we also calculated mass-specific MR (Reinhold 1999) for

some analyses even though (ANCOVA-type) models that include mass and MR are

recommended (Lighton 2008). Note sample sizes differ between pupal and adult

analyses because not all pupae (N = 205) survived to adulthood (N = 178), and for some

individuals, data were missing.

Results.

We initially conducted a MANOVA to test for genetic effects on our traits of interest, with

genotype (isoline) as the predictor, and pupal and adult metabolic rates and mass, and

adult mandible length as dependents. This showed a strong genetic effect on the

multivariate combination of these traits (isoline: Wilks' Lambda $F_{85,647.4}$ = 1.612; P < 0.001).

This was driven by genotype (isoline) effects on adult MR (F = 1.750; P = 0.041), mandible

length (F = 1.857; P = 0.027) and pupal mass (F = 2.007; P = 0.015). Using mass-specific

MR (Reinhold 1999) rather than mass and MR, did not change univariate conclusions –

isoline still affects mandible length (F = 1.857; P = 0.027) and adult mass-specific MR (F

= 1.79; P = 0.035) – although the multivariate effect was no longer significant (Wilks'

Lambda = 0.661; $F_{51,402.7}$ = 1.18; P = 0.196).

We then tested for associations between mass and MR in adults and pupae using GLMs

(that included isoline as a random effect and age as a fixed effect) and least squares

linear regression (MR vs Mass). For pupal MR, the only significant effect in the GLM was

pupal age ($F_{3,140}$ = 59.258; P < 0.001; all other effects F < 0.001; P > 0.99), with MR tending

to increase with pupal age (Figure 1). However, regression of pupal MR on mass (i.e. the

phenotypic association ignoring age and genetic affects) found a significant positive

association between MR and mass (standardised β = 0.234; $F_{1,204}$ = 11.750; P < 0.001). In

the GLM for adult MR, adult mass did not impact adult MR ($F_{1,154}$ = 2.092; P = 0.150), but

age and genotype (isoline) had significant effects (Age: $F_{5,154}$ = 2.698; P = 0.023. Isoline:

$F_{17,154}$ = 2.053; P = 0.011). However, the age affect was driven by the elevated MR of two 15

day-old individuals, and when they were removed from the analysis, pupal age no longer

significantly impacted MR (isoline remained significant). Regression of adult MR on adult

mass (i.e. the phenotypic association ignoring age and genetic affects) revealed a non-

significant but positive association between MR and mass (standardised β = 0.115; $F_{1,176}$

= 2.357; P = 0.126).

To test for effects of MR on mandible length, we used a univariate GLM with isoline as a

random effect and pupal and adult mass and MR as covariates. This revealed strong

positive effects of pupal mass on mandible length ($F_{1,132}$ = 5.893; P = 0.017. Figure Y) and

an effect of genotype ($F_{17,132}$ = 1.855; P = 0.028. Figure Z), with no other predictor

significantly affecting mandible length (all F < 1.63; all P > 0.2). The removal of one very

small individual (Figure 2) did not alter these findings. Similarly, removal of individuals

with small mandibles for their isolines (Figure 3) did not change model outcomes.

Additionally, analyses using pupal and adult mass-specific MR (with isoline as a random

effect) also recovered the genotype effect ($F_{17,135}$ = 1.739; P = 0.043), but again found no

effect of MR on mandible size (all F < 0.35; P > 0.54), and multiple regression of

phenotypic values (i.e. that did not include isoline) was significant overall ($F_{4,150}$ = 9.203;

P < 0.001) because of an effect of pupal mass on mandible size (standardised β = 0.315;

t = 3.124; P = 0.002) with no other significant effects (adult and pupal MR, adult mass: all

t < 1.3; P > 0.19). Finally, given associations between mass and MR (see above), we also

ran a GLM that only included MR (adult pupae) and isoline (fixed effect) as predictors and

still found no MR-mandible associations (all F < 1.67; P > 0.19).

Discussion

Despite expected associations between metabolic rate and weapon size (e.g. Tomkins et

al. 2004; Biro and Stamps 2010), we did not find any statistically significant relationships

between the two either during pupation, when mandibles develop (Sugiyama et al. 2023),

or during adulthood when they are used in fights (Harano et al. 2010). This is despite large

sample sizes and finding a range of other statistically significant associations. Most

notable of these was the significant effect of isoline on all the traits we measure, which

is consistent with genetic underpinnings to these characters (David et al. 2005). We

discuss these findings in turn.

There are good reasons to expect resting (basal) metabolic rate to be associated with

weapons size, either because a low resting metabolic rate uses less energy and hence

more energy is available to be allocated to weapons, or because a higher metabolic rate

allows individuals to garner more resource to be allocated to armaments (e.g. Tomkins et

al. 2004; Biro and Stamps 2010). Additionally, many weapons are positively allometric

(Kodric-Brown et al. 2006), including the mandibles at the centre of our study (Okada and

Miyatake 2010), and metabolic expenditure has been positively associated with weapon

size in other taxa (e.g. Somjee et al. 2018). Furthermore, morphological characters are a

fixed cost once developed and hence may be especially likely to be energetically

expensive (Allen & Levinton 2007). However, despite all this, we did not find any

associations, and MR – weapon associations are generally highly variable and appear to

be influenced by the muscle mass associated with the weapons (O'Brien et al. 2019) and

the tissue type that weapons are built from (Somjee 2021). In *G. cornutus*, the muscle

largely used to operate the mandible sits within the pronotum and this is highly

genetically and phenotypically correlated with mandible size (e.g. Harano et al. 2010). So

despite not measuring muscle mass in our study, mandible size itself should capture

much of the variation in muscle mass and hence reflect this association if it existed.

However, the mandibles themselves are exoskeleton and this should reduce their

metabolic maintenance costs (Dinh 2022) which is consistent with the lack of

association between adult MR and weapon size in *G. cornutus*.

While there might not be significant metabolic costs associated with bearing mandibles

once they have developed (Dinh 2022; Somjee 2021), we thought there could well be

elevated metabolic expenditure associated with developing larger mandibles as has

been found in other taxa (Somjee 2021). For this reason MR was measured during

pupation – when mandibles develop (e.g. Sugiyama et al. 2023). Again however, we found

no associations between (pupal) MR and weapon size, although larger pupal mass, which

probably reflects resource accumulation prior to trait development, was positively

associated with mandible size. If pupal mass reflects resource accumulation, then it

could be argued that there are energetic links to mandible size, but these are independent

of pupal (and adult) metabolic rate. Consistent with this, the balance of macronutrients

in the larval diet has a strong effect on mandible size in adult beetles (House et al. 2016).

Additionally, in populations selected for increased mandible size, there is a reduction in

survival during pupation (Okada et al. 2011), which could mean we have underestimated

metabolic expenditure on mandible growth if those individuals with the largest

mandibles do not survive through pupation to be measured. In any case, previous

findings of predation costs associated with larger mandibles, and knock-on longevity

impacts (Okada et al. 2021; Pennell et al. 2024). Thus there are life-history costs to larger

mandibles, as required to demonstrate "real" costs associated with sexually selected

characters (Kotiaho 2001).

Like all non-significant associations, it is possible that we simply lacked the statistical

power to find any effect. However, our sample sizes were large, MR was repeatable, and

we did find other associations, so we are reasonably confident that the lack of an

association between MR and mandible size is real. This conclusion is also consistent

with findings from other taxa where larger weapons are composed of more energetically

inexpensive exoskeleton (Dinh 2022). However, comparing male and female mass-

specific metabolic rates would provide another means of exploring potential energetic

costs to mandibles (Oufiero & Garland 2007) to ensure these conclusions are robust, and

this is something for future work, especially since survival through the pupal stage could

be impacted by mandible size (Okada et al. 2011).

Despite finding no MR-mandible associations, we nonetheless identified isoline effects

for mandible size, pupal mass and adult metabolic rate, reflecting genetic variation in all

these characters (Hoffman and Parsons 1988; David et al. 2005). It was not surprising

that we found evidence for a genetic basis to mandible size, since this trait has a high

heritability (Okada et al. 2014), responds readily to artificial selection (Okada and

Miyatake 2009) and to male specific predation (Okada et al. 2021). For example,

bidirectional selection on mandible size generates rapid microevolutionary responses for

both larger and smaller size, and also generates as raft of correlated response in body

shape (Okada and Miyatake 2009; Harano et al. 2010). Likewise, there were apparent

genetic effects on pupal mass which is not surprising as size measures are generally

heritable (e.g. Messina 1993; Iyengar and Eisner 1999; Blanckenhorn and Hosken 2003).

Genetic variation for MR has also been found in other taxa (e.g. Nilsson et al. 2009; Einum

et al. 2019), although this at times depends on cyto-nuclear interactions (Arnqvist et al.

2010), but in any case, there appears to be a genetic component to adult MR in *G.*

cornutus. The importance of this insight is not clear as we do not have a thorough

understanding of MR-fitness links at this point. However, genotype had no effect on pupal

MR, but there were strong age effects which likely reflect the effects of development stage on energy expenditure, as has been documented in other studies (e.g. Krams et al. 2017, 2018). Additionally, these age effects may have obscured any genetic effect on pupal MR. We also found no significant genotype effect on adult mass which was somewhat surprising because adult mass has a high heritability in *G. cornutus* when estimated with a paternal half-sib breeding design (Okada et al. 2014). Of course such designs are specifically to enable genetic parameter estimation and will return estimates with greater accuracy than our approach here.

We also documented mass-MR associations in some analyses but not others, with pupal MR having the strongest associations with mass (although MR and mass were positively associated in all analyses). While mass-MR associations are well established across species (Withers 1992; Nespolo et al. 2005), intra-specific associations can be weaker. Additionally, while some insect studies find mass-MR relationships (e.g. Lighton 2008; Okada et al. 2011; Sinclair et al. 2011; Matsumara et al. 2025), others do not (Van Voorhies et al. 2004; Canzano et al. 2006; Oikawa et al. 2006), which may reflect differences in factors like starvation resistance and time in captivity (e.g. Harshman et al. 1999; Van Voorhies et al. 2004). As noted above, age and genetic effects were also implicated in MR variation. Finally, the MR estimates we obtained here are similar to those for other similarly sized beetles (Matsumara et al. 2025).

In summary, we found no evidence for associations between metabolic rate and an exaggerated sexually selected trait used in male-male combat. This is despite the fact that this weapon shows positive allometry and has other costs associated with its expression. This finding is consistent with suggestions that tissue type affects the energetic costs of weapons (Somjee 2021). Additionally, we record genotypic effects on metabolic rate, but whether metabolic rate relates to life-history characters and fitness remains to be elucidated, as does potential associations between MR and fighting ability.

References

Allen NJ, Levinton JS. 2007. Costs of bearinfg a sexually selected ornamental weapon in
a fiddle crab. Funt Ecol 21:154-161.

Andersson, M. 1994. *Sexual selection*. Princeton University Press.

Archer CR, Alper C, Mack L, Weedon M, Sharma MD, Sutter A, Hosken DJ. 2024. Alcohol
reduces choosiness and relaxes mate preference in female *Drosophila simulans*.
Biol J Linn Soc 143:1-11.

Arnqvist G, Dowling DK, Eady P, Gay L, Tregenza T, Tuda M, Hosken DJ. 2010. The genetic
architecture of metabolic rate: environment specific epistasis between
mitochondrial and nuclear genes in an insect. Evolution 64:3354-3363.

Bailey WJ, Withers PC, Endersby M, Gaull K. 1993. The energetic cost of calling in the
bushcricket *Requena verticalis* (Orthoptera: Tettigoniidae: Listrocelidinae). J Exp
Biol 178:21–37.

Bennet AF, Ruben JA. 1979. Endothermy and activity in vertebrates. Science 206:649–
654.

Blanckenhorn WU, Hosken DJ. 2003. Heritable condition in yellow dung flies. Behav
Ecol 14:612-618.

Biro PA, Stamps JA. 2010. Do consistent individual differences in metabolic rate
promote consistent individual differences in behaviour? Trends Ecol Evol 2:653–
659.

Canzano AA, Krockenberger AA, Jones RE, Seymour JE. 2006. Rates of metabolism in
diapausing and reproductively active butterflies, *Euploea core* and *Euploea*
sylverster (Lepidoptera: Nymphalidae). Physiol Entomol 31:184-189.

David JR, Gilbert P, Legout H, Pétavy G, Capy P, Moreteau B. 2005. Isofemale lines in
Drosophila: an empirical approach to quantitative trait analysis in natural
populations. Heredity 94:3-12.

Dinh JP. 2022. Large and exaggerated sexually selected weapons comprise high
proportions of metabolically inexpensive exoskeleton. Biol Lett 18:20210550.

Eberhard WG, Rodriguez RL, Huber BA, Speck B, Miller H, Buzatto BA, Machado G.
2018. Sexual selection and static allometry: the importance of function. Qrt Rev
Biol 93:207-250.

Einum S, Fossen EIF, Parry V, Pelabon C. 2019. Genetic variation in metabolic rate and

correlations with other energy budget components and life history in *Daphnia*

magna. Evol Biol 46:170-178.

Endler JA. 1987. Predation, light intensity and courtship behavior in *Poecilia reticulata*

(Pisces: Poeciliidae). Anim Behav 35:1376 -1385.

Folstad I, Karter AJ. 1992. Parasites, bright males and the immunocompetence

handicap hypothesis. Am Nat 139:603-622.

Gamer M, Lemon J, Fellows I, Singh P. 2019. *irr: Various coefficients of interrater*

reliability and agreement. Retrieved from https://CRAN.R-project.org/package=irr

Gray DA, Cade WH. 1999. Sex, death and genetic variation: natural and sexual selection

on cricket song. Proc R Soc Lond B 266:707–709.

Hack MA.1997. The energetic costs of fighting in the house cricket, *Acheta domesticus*

L. Behav Ecol 8:28-36.

Halliday TR. 1987. Physiological constraints on sexual selection. In: Bradbury JW and

Andersson MB, editors, *Sexual selection: testing the alternatives*. John Wiley and

Sons. Pp 247-264.

Harano T, Okada K, Nakayama S, Miyatake T, Hosken DJ. 2010. Intralocus sexual conflict

unresolved by sex-limited trait expression. Curr Biol 20:2036–2039.

Harshman LG, Hoffmann AA, Clarke AG. 1999. Selection for starvation resistance in

Drosophila melanogaster: physiological correlates, enzyme activities and multiple

stress responses. J Evol Biol 12:370-379.

Hoback WW, Wagner WE. 1997. The energetic costs of calling in the variable field

cricket, *Gryllus lineaticeps*. Physiol Entomol 22:286-290.

Hoffmann AA, Parsons PA. 1988. The analysis of quantitative variation in natural

populations with isofemale strains. Genet Sel Evol 20:87-98.

Hosken DJ. 2001. Sex and death: microevolutionary trade-offs between reproductive

and immune investment in dung flies. Curr Biol 11:R379-R380.

Hosken DJ, Bailey WJ, O'Shea JE, Roberts JD. 1994. Localisation of insect calls by the

bat *Nyctophilus geoffroyi* (Chiroptera: Vespertilionidae): a laboratory study. Aust J

Zool 42:177-184.

Hosken DJ, Buss DL, Hodgson DJ. 2018. Beware the F-test (or how to compare variances).

Anim Behav 136:119-126.

House CM, Jensen K, Rapkin J, Lane S, Okada K, Hosken DJ, Hunt J. 2016. Macronutrient
balance mediates the growth of sexually selected weapons but not genitalia in male
broad-horned beetles. Funct Ecol 30:769-779.

Ingleby FC, Hunt J, Hosken DJ. 2013. Genotype-by-environment interactions for female
mate choice of male cuticular hydrocarbons in *Drosophila simulans*. PLoS One
8:e67623.

Iyengar VK, Eisner T. 1999. Heritability of body mass, a sexually selected trait, in an
arctiid moth (*Utetheisa ornatrix*). Proc Natl Acad Sci USA 96:9169-9171.

Katsuki M, Harano T, Miyatake T, Okada K, Hosken DJ. 2012. Intralocus sexual conflict
and offspring sex ratio. Ecol Lett15:193–197.

Ketola T, Kotiaho JS. 2009. Inbreeding, energy use and condition. J Evol Biol 22:770-781.

Ketola T, Kotiaho JS. 2010. Inbreeding, energy use and sexual signalling. Evol Ecol
24:761–772.

Knief U, Forstmeier W. 2021. Violating the normality assumption may be the lesser of
two evils. Behav Res Methods 53:2576-2590.

Kodric-Brown A, Sibly RM, Brown JH. 2006. The allometry of ornaments and weapons.
Proc Natl Acad Sci USA 103:8733-8738.

Kotiaho JS. 2001. Cost of sexual traits: a mismatch between theoretical considerations
and empirical evidence. Biol Rev 76:365–376.

Kotiaho JS, Alatalo RV, Mappes J, Nielsen MG, Parri S, Rivero A. 1998. Energetic costs of
size and sexual signalling in a wolf spider. Proc R Soc Lond B 265:2203–2209.

Krams I, Trakimas G, Kecko S, Elferts D, Krams R, Luoto S, Rantala MJ, Mänd M, Kuusik
A, Kekäläinen J, Jõers P, Kortet R, Krama T. 2018. Linking organismal growth, coping
styles, stress reactivity, and metabolism via responses against a selective
serotonin reuptake inhibitor in an insect. Sci Reports 8:8599.

Krams IA, Niemelä PT, Trakimas G, Krams R, Burghardt GM, Krama T, Kuusik A, Mänd M,
Rantala MJ, Mänd R, Kekäläinen J, Sirkka I, Luoto S, Kortet R. 2017. Metabolic rate
associates with, but does not generate covariation between, behaviours in western
stutter-trilling crickets, *Gryllus integer*. Proc R Soc B 284:20162481.

Lighton JRB. 2008. *Measuring metabolic rates*. Oxford University Press.

Matsumura K, Abe MS, Sharma MD, Hosken DJ, Yoshii T, Miyatake T. 2020. Genetic
variation and phenotypic plasticity in circadian rhythms in an armed beetle,
Gnatocerus cornutus (Tenebrionidae). Biol J Linn Soc 130:34-40.

Matsumura K, Yamamoto Y, Yoshimura K, Miyatake T. 2024. Effect of temperature on
sexual size dimorphism during the developmental period in the broad-horned flour
beetle. J Thermal Biol 124:103962.

Matsumura K, Hosken DJ, Noda T, Miyatake T, Sharma MD. 2025. Behaviour, behavioural
syndromes and metabolism: the effects of artificial selection for death feigning on
metabolic rate. J Insect Sci 25:16; ieaf007.

McKean KA, Nunney L. 2008. Increased sexual activity reduces male immune function
in *Drosophila melanogaster*. Proc Natl Acad Sci USA 98:7904-7909.

McNab BK. 2005. Food habits and the evolution of energetics in birds of paradise
(Paradisaeidae). J Comp Physiol B 175:117–132.

Messina FJ. 1993. Heritability and "evolvability" of fitness components in
Callosobruchus maculatus. Heredity 71:623-629.

Neat FC, Taylor AC, Huntingford FA. 1998. Proximate costs of fighting in cichlid fish: the
role of injuries and energy metabolish. Anim Behav 55:875-882.

Nishitani T, Matsumura K, Postma E, Sharma MD, Hosken DJ, Miyatake T. 2024.
Experimental quantification of genetic and ontogenetic effects on fighting
behaviour in the broad-horned flour beetle. Behav Ecol Sociobiol 78:34.

Nespolo RF, Lardies MA, Bozinovic F. 2003. Intrapopulation variation in the standard
metabolic rate of insects: repeatability, thermal dependence and sensitivity (Q_{10})
of oxygen consumption in a cricket. J Exp Biol 206:4309–4315.

Nilsson J-A, Åkesson M, Nilsson JF. 2009. Heritability of resting metabolic rate in a wild
population. J Evol Biol 22:1867-1874.

O'Brian DM, Boisseau RP, Duell M, McCullough E, Powell EC, Somjee U, Solie S, Hickey
AJ, Howell GI, Painting CJ, Emlen DJ. 2019. Muscle mass drives costs in sexually
selected arthropod weapons. Proc R Soc B 286: 20191063.

Okada K, Katsuki M, Okada Y, Miyatake T. 2011. Immature performance linked with
exaggeration of a sexually selected trait in an armed beetles. J Evol Biol 24:1737-
1743.

Okada K, Katsuki M, Sharma MD, House CM, Hosken DJ. 2014. Sexual conflict over
mating in *Gnatocerus cornutus*: females prefer lovers not fighters. Proc R Soc B
281:20140388.

Okada K, Katsuki M, Sharma MD, Kiyose K, Seko T, Okada Y, Wilson AJ, Hosken DJ. 2021.
Natural selection increases female fitness by reversing the exaggeration of a
male sexually selected trait. Nat Commun 12:3420.

Okada K, Miyanoshita A, Miyatake T. 2006. Intra-sexual dimorphism in male mandibles
and male aggressive behavior in the broad horned flour beetle *Gnatocerus*
cornutus (Coleoptera: Tenebrionidae). J Insect Behav 19:457-467.

Okada K, Miyatake T. 2009. Genetic correlations between weapons, body shape and
fighting behaviour in the horned beetle *Gnatocerus cornutus*. Anim Behav
77:1057-1065.

Okada K, Miyatake T. 2010. Plasticity of size and allometry in multiple sexually selected
traits in an armed beetle *Gnatocerus cornutus*. Evol Ecol 24:1339-1351.

Okada K, Pitchers WR, Sharma MD, Hunt J, Hosken DJ. 2011. Longevity, calling effort and
metabolic rate in two populations of cricket. Behav Ecol Sociobiol 65:1773-1778.

Oikawa A, Mori N, Kimura MT. 2006. A comparison of oxygen consumption in *Drosophila*
flies from different climates. Entomol Sci 9:347-354.

Oufiero CE, Garland T Jr. 2007. Evaluating performance costs of sexually selected traits.
Funct Ecol 21: 667-689.

Pennell T, Katsuki M, Archer CR, Sharma MD, Okada K, Hosken DJ. 2024. Predation affects
the evolution of sex-specific longevity. Biol Lett 20:20240451.

Reinhold K. 1999. Energetically costly behaviour and the evolution of resting metabolic
rate in insects. Funct Ecol 13:217-224.

Reinhold K, Greenfield MD, Jang Y, Broce A. 1998. Energetic cost of sexual
attractiveness: ultrasonic advertisement in wax moths. Anim Behav 55:905–913.

Rezende EL, Bozinovic F, Garland T Jr. 2004. Climatic adaptation and the evolution of
basal and maximum rates of metabolism in rodents. Evolution 58:1361–1374.

Reznick DA, Endler JA. 1982. The impact of predation on life history evolution in
Trinidadian guppies (*Poecilia reticulata*). Evolution 36:160-177.

Rosenfeld J, Van Leeuwen T, Richards J, Allen D. 2015. Relationship between growth and

standard metabolic rate: measurement artefacts and implications for habitat use

and life-history in salmonids. J Anim Ecol 84:4-20.

Rowe L, Houle D.1996. The lek paradox and the capture of genetic variance by condition

dependent traits. Proc R Soc Lond B 263:1415–1421.

Ryan MJ. 1985. *The Túngara Frog*. University of Chicago Press.

Ryan MJ. 2018. *A taste for the beautiful: the evolution of attraction*. Princeton University

Press.

Salvador A, Veiga JP,Martin J, Lopez P, Abelenda M, Puerta M. 1996. The cost of

producing a sexual signal: testosterone increases the susceptibility of male lizards

to ectoparasitic infestation. Behav Ecol 7: 145-150.

Sinclair BJ, Bretman A, Tregenza T, Tomkins JL, Hosken DJ. 2011. Metabolic rate does not

increase with starvation in *Gryllus bimaculatus* when changing fuel use is taken into

account. Physiol Entomol 36:84-89.

Snodgrass JJ, Leonard WR, Tarskaia LA, Alekseev VP, Krivoshapkin VG. 2005. Basal

metabolic rate in the Yakut (Sakha) of Siberia. Am J Hum Biol 17:155–172.

Somjee U. 2021. Positive allometry of sexually selected traits: do metabolic

maintenances costs play a important role? BioEssays 43:2000183.

Somjee U, Woods HA, Duell M, Miller CW. 2018. The hidden cost of sexually selected

traits: the metabolic expense of maintaining a sexually selected weapon. Proc R

Soc B 285:20181685.

Sugiyama M, Ozawa T, Ohta K, Okada K, Niimi T, Yamaguchi K, Shigenobu S, Okada Y.

2023. Transcriptomic and functional screening of weapon formation genes implies

significance of cell adhesion molecules and female-biased genes in broad-horned

flour beetles. PLoS Genetics 19:e1011069.

Tomkins JL, Radwan J, Kotiaho JS, Tregenza T. 2004. Genic capture and resolving the lek

paradox. Trends Ecol Evol 19:323–328.

Van Voorhies WA, Khazaelia AA, Curtsinger JW. 2004. Lack of correlation between body

mass and metabolic rate in *Drosophila melanogaster*. J Insect Physiol 50:445-453.

Withers PC. 1992. *Comparative animal physiology*. Saunders.

Yamane T, Okada K., Nakayama, S, Miyatake T. 2010. Dispersal and ejaculatory

strategies associated with exaggeration of weapon in an armed beetle. Proc R Soc B

277:1705-1710.

Zuk M, Kolluru GR. 1998. Exploitation of sexual signals by predators and parasitoids. Qrt

Rev Biol 73:415–438.

Zuk M, Rotenberry JT, Tinghitela RM. 2006. Silent night: adaptive disappearance of a

sexual signal in a parasitized population of field crickets. Biol Lett 2:521-524.

Figure 1. The effect of pupal age on pupal metabolic rate (MR). MR tends to increase as

development proceeds. Shown are medians, 1st and 3rd quartile, minimum and maximum

and outliers. Interpretation does not change if outliers (2-day old individuals) are

removed from the analysis.

582 **Figure 2.** The statistically significant variation in mandible size (um) across the 18

583 genotypes (isoline) used in the study - which have been given arbitrary numbers. Shown

584 are medians, 1st and 3rd quartile, minimum and maximum and outliers. Not the effect of

585 isoline on mandible size is not changed with the removal of outliers (individuals with

586 small mandible size for their genotype).

591 **Figure 3.** The positive association between pupal mass (mg) and the length of mandibles

592 (μm) in adult males. This relationship is statistically significant and this is not altered by

593 the removal of the single very small male.