Characterisation of putative retrotrapezoid nucleus (RTN) chemoreceptor neurons in the adult human brainstem

Yazhou Liu[1], Rita Machaalani[2], Irit Markus[1], Claire E. Shepherd*[1,3], Natasha N. Kumar*[1]

[1] Faculty of Medicine and Health, The University of New South Wales, NSW, 2052, Australia

[2] Sydney Medical School, Faculty of Medicine and Health, University of Sydney, NSW, 2006, Australia

[3] Neuroscience Research Australia, Barker Street, NSW, Australia

Corresponding author: Natasha N Kumar (natasha.kumar@unsw.edu.au)

Authors ORCIDS:

Yazhou Liu: 0000-0002-2096-4064

Natasha Kumar: 0000-0003-4712-8792

Rita Machaalani: 0000-0002-7675-1121

Irit Markus: None

Claire Shepherd: 0000-0002-0399-3218

*Co-senior authors

Abstract

The retrotrapezoid nucleus (RTN) of rodents is located ventral to the facial motor nucleus (7N) and

consists of acid-sensitive neurons that activate breathing and mediate the central component of the

ventilatory response to hypercapnia. In rodents, RTN neurons can be histologically identified by the

presence of paired-like homeobox 2B positive nuclei (Phox2b+) and the absence of cytoplasmic choline

acetyltransferase (ChAT-) and tyrosine hydroxylase (TH-). Up to 50% of rodent RTN neurons

synthesise galanin, and 88% express pituitary adenylate cyclase activating polypeptide (PACAP). The

human RTN (hRTN) has not been mapped to date. This study aimed to map the location and

cytoarchitecture of the adult hRTN and compare the findings to the homologies of rodents, macaques

and human infants. Formalin-fixed, paraffin-embedded tissue blocks from two adult cases, spanning

the medulla-pons, were serially sectioned (10μm thick) and every four in thirty sections was assayed

for immunohistochemistry for ChAT, or double-labelled Phox2b/TH, Phox2b/galanin and

Phox2b/PACAP, followed by analysis using QuPath software. hRTN neurons, identified as

Phox2b+/TH-/ChAT-, were located ventral to 7N and lateral to the superior olive, overlapped with the

C1 or A5 catecholaminergic population and extended rostrocaudally from Obex +13 to +17 mm. In the

parafacial area, 90% of Phox2b immunoreactive (-ir) neurons are hRTN neurons, totaling around 5000

bilaterally, and were surrounded by numerous TH-ir fibers. Galanin- and PACAP-ir was identified in

43% and 39% of Phox2b-ir parafacial neurons, respectively. This is the first study to characterise and

quantitatively map the adult human RTN using a series of neurochemical markers.

Keywords

Human RTN, Phox2b, chemoreception, immunohistochemistry, galanin, PACAP

Abbreviations

4V, 4th ventricle; 6N, abducens nucleus; 6n, abducens nerve; 7N, facial motor nucleus; 7n, facial nerve;

8n, vestibulocochlear nerve; A5, A5 catecholaminergic neurons; Biotin-SP-conjugated, biotin-

succinimidyl ester conjugated; C1, C1 adrenergic neurons; CCHS, congenital hypoventilation

syndrome; ChAT, choline acetyltransferase; CO_2, carbon dioxide; D, dorsal; DAB, 3,3'-

diaminobenzidine; DPX, dibutylphthalate polystyrene xylene; FFPE, formalin-fixed, paraffin-

embedded; H_2O_2, hydrogen peroxide; H&E, hematoxylin & eosin; hRTN, human retrotrapezoid nucleus;

HIER, heat-induced epitope retrieval; icp, inferior cerebellar peduncle; IC, inferior colliculus; IF,

immunofluorescence; IgG, immunoglobulin G; IHC, immunohistochemistry; IO, inferior olive; IOP,

inferior olive, principal nucleus; -ir, immunoreactivity; IRt, intermediate reticular nucleus; IS, inferior

salivatory nucleus; ISH, in situ hybridisation; mcp, medial cerebellar peduncle; ml, medial lemniscus;

MSO, medial superior olive; NDS, normal donkey serum; NK1R, neurokinin-1 receptor; NMB,

neuromedin B; NTS, nucleus of the solitary tract; PACAP, pituitary adenylate cyclase activating

polypeptide; PBS, phosphate-buffered saline; PCRt, parvocellular reticular nucleus; PCRtA,

parvocellular reticular nucleus, alpha; Phox2b, paired-like homeobox 2B; PPGAL, preprogalanin; Pn,

pontine nuclei; py, pyramidal tract; R, rostral; Rn, Red nucleus; RNA, ribonucleic acid; ROI, regions

of interest; RRID, Research Resource Identifier; RTN, retrotrapezoid nucleus; SC, superior colliculus;

SLC17A6, solute carrier family 17 member A6 (vesicular glutamate transporter 2 gene); SO, superior

olive; Sp5O, spinal trigeminal nucleus, oral part; sp5, spinal trigeminal tract, nerve; TBS, Tris-buffered

saline; TH, tyrosine hydroxylase; Thal, thalamus; V, ventral; vGlut2, vesicular glutamate transporter 2;

VRC, ventral respiratory column; WB, Western blotting.

Introduction

In mammals, breathing is regulated by central respiratory nuclei and a neural network in the brainstem

(Ikeda et al., 2017). Central respiratory chemoreceptors are clusters of cells in the lower brainstem

respiratory network which regulate breathing stimulated by carbon dioxide (CO_2) or its proxy, protons

(H^+). Acid sensing chemoreceptor neurons in the retrotrapezoid nucleus (RTN) (Guyenet and Bayliss,

2015, Guyenet and Bayliss, 2022) project to the ventral respiratory column (VRC) which drives the

central respiratory chemoreflex response (Guyenet et al., 2019). Impaired central respiratory

chemoreception results in an impaired hypercapnic ventilatory response, as occurs in congenital

hypoventilation syndrome (CCHS) and central sleep apnoea (Kasi et al., 2018, Meylemans et al., 2021,

Degl'Innocenti et al., 2018, Amiel et al., 2009).

The RTN comprises about 2000 neurons in the rat, and 700 - 800 neurons in mice, bilaterally

(Takakura et al., 2008, Guyenet and Bayliss, 2015, Shi et al., 2017, Lazarenko et al., 2009). Phox2b is

the most widely recognised biomarker for RTN chemoreceptors, and changes in its expression are

regarded as an indicator of RTN chemoreceptor dysfunction associated with various central respiratory

disorders in both rodents and humans (Amiel et al., 2009, Goridis and Brunet, 2010, Meylemans et al.,

2021, Lavezzi et al., 2012). In addition to Phox2b, alterations in neuropeptides, such as galanin and

PACAP, are important for regulation of central chemoreception in the RTN. In rodents, galanin is

expressed in approximately 50% of RTN neurons (Stornetta et al., 2009) and the PreBotzinger complex

receives glutamatergic innervation from galaninergic RTN neurons (Bochorishvili et al., 2012, Dereli

et al., 2024). Long term hypercapnia induces an increase in preprogalanin (PPGAL) expression in the

RTN (Dereli et al., 2019) and microinjection of galanin into the VRC, specifically in the PreBotzinger

and Botzinger complexes, impairs the chemoreflex response to hypercapnia (Abbott et al., 2009). In

neonatal mice, PACAP is expressed in 100% of RTN chemoreceptor neurons and the deletion of

PACAP in RTN neurons leads to elevated incidence of apnoea's and reduced respiratory responses to

CO_2 stimulation in the postnatal period (Shi et al., 2021). Although PACAP expression in the human

brainstem has been studied (Huang et al., 2017), the RTN was not investigated due to the lack of

knowledge regarding its precise location.

To date, four studies have investigated the location of the RTN in primates; 1 in monkeys and

3 in humans (Table 1). Levy and colleagues aimed to identify Phox2b mRNA and protein in formalin-

fixed, paraffin-embedded (FFPE) adult human brainstem, but it failed to be detected and as such, the

data was on a population of PPGAL/ Solute Carrier Family 17 Member A6 (SLC17A6) mRNA

containing neurons in the parafacial region (Levy et al., 2021). The other two studies were conducted

on FFPE tissue from foetuses or infants (Rudzinski and Kapur, 2010, Lavezzi et al., 2012). Rudzinski

and Kapur (2010), described the putative human RTN (hRTN) as a discontinuous cluster of small- to

medium-sized Phox2b positive/ Neurokinin 1 receptor positive/Tyrosine hydroxylase negative

(Phox2b+/NK1R+/TH-) neurons situated ventral to the facial motor nucleus (7N) and lateral to the

superior olivary nucleus (SO) spanning from the caudal pons to the rostral medulla (Rudzinski and

Kapur, 2010). Lavezzi et al (2012) described the RTN as a cluster of Phox2b immunoreactive (-ir)

neurons situated between the 7N and SO in the caudal pons (Lavezzi et al., 2012). While both infant

studies sought to characterise the hRTN, they were restricted by limited tissue sampling and inconsistent

Phox2b-ir patterns (Table 1).

Since the RTN has not been systematically mapped using Phox2b in the human brainstem, our

overall study goal was to fill this research gap. This immunohistochemical study used post-mortem

FFPE tissue from 2 adult humans to: 1) characterize Phox2b-ir in the RTN region, 2) establish the

neurochemical signature of hRTN neurons and quantify these neurons, 3) establish the neuroanatomical

location and dimensions of the hRTN relative to 7N, C1 or A5 catecholaminergic neurons, 4) determine

the proportion of parafacial Phox2b-ir neurons expressing the neuropeptides galanin and PACAP, and

5) describe the morphological features (size and density) of parafacial Phox2b-ir neurons. Our findings

provide valuable insights for investigations into Phox2b-associated disorders, particularly those

involving respiratory and autonomic dysfunction including CCHS and sleep apnoea.

 Table 1: Summary of data collected on primate RTN in chronological order.

Developmental age, number of cases, Sex ratio	Tissue Frozen or FFPE, and plane of cut (coronal vs sagittal)	Sections/case analysed; Obex/bregma level	Antibody/probe details & dilution	RTN Phox2b subcellular labelling pattern & anatomical location	Number of RTN neurons (bilateral)	Reference
Human fetuses and infants (19-44w, n=8)	5 μm coronal (n=5) and sagittal (n=3) FFPE sections	1-2 sections/case (level not specified but in the presence of the 7N).	Rabbit anti-Phox2b (gift of J.-F. Brunet, 1:1000). NK1R (Abcam, 1:200); TH (Chemicon, 1:1000)	IHC: cell nucleus. RTN was ventral to 7N and lateral to the SO in the pontomedullary junction.	Not provided	(Rudzinski and Kapur, 2010)
Human fetuses (35-40w, n=9) and infant (2-8 months, n=11)	4 μm coronal FFPE sections	3-6 sections/case: 1-2 sections from each brainstem level of upper pons, upper medulla and from Obex -2/3 mm to +2/3 mm.	Rabbit anti-Phox2b (gift of J.-F. Brunet, 1:750)	IHC: cell nucleus and cytoplasm. RTN located between the 7N and SO in caudal pons.	980 ± 23 Phox2b-ir neurons However, equation applied not provided	(Lavezzi et al., 2012)
Macaca fascicularis (5- & 6-years old males and 16-year-old female, n=2 IHC; n=1 ISH)	40 μm coronal frozen and paraformaldehyde fixed free-floating sections	60 sections/case (1/8 serial sections); Obex -1.40mm to +13mm.	IHC: Rabbit anti-Phox2b antibody (1:2,000; gift from J-F. Brunet) ISH: PPGAL and Phox2b mRNAs	IHC: subcellular location of Phox2b immunoreactivity not described. Parafacial region located ventral and lateral to 7N; ISH: cell cytoplasm. PPGAL+/Phox2b+ RTN neurons located caudal, ventral and lateral to 7N spanning from Obex +4.56 to +10.16 mm.	2400 PPGAL+/ Phox2b+ neurons. PPGAL-/Phox2b neurons were not included. Calculated as an estimation by automated method yet cautions on possibility of double counting.	(Levy et al., 2019)
Human adults (n=2)	10 μm coronal FFPE sections	60 sections/case (1/10 serial sections), Obex +14 mm to +17 mm.	mRNA probes (targeting 237–492 sequence of the galanin message-associated peptide mRNA (i.e. PPGAL) and 469–1173 sequence of the vGlut2 mRNA (SLC17A6) and NMB.	IHC: N/A since Phox2b did not work. ISH: N/A since Phox2b did not work. Based on PPGAL, SLC17A6 and NMB, RTN located in parafacial region.	Not provided	(Levy et al., 2021)
Human adults (n=2)	10 μm coronal FFPE sections	90 sections/case (5/30 serial sections), Obex +12 mm to +18 mm.	Mouse anti-Phox2b (Bio-Strategy Laboratory, cat#sc-376997, 1:1000)	IHC: cell nucleus RTN was identified ventral to 7N, lateral to the SO and medial to facial nerve and Sp5O.	Approximately 5000 Phox2b+TH-ChAT- RTN neurons.	Study herein

Methods

Human Tissue Selection

Post-mortem, formalin-fixed brainstem specimens from two adult human cases (76-year-old female, and 87-year-old male; Supplementary Table 1), were sourced from the Sydney Brain Bank through regional brain donor programs in Sydney, Australia. Specimens were acquired from asymptomatic individuals exhibiting solely age-related brain changes and devoid of significant neuropathology. Exclusion criteria included documented respiratory drive disorders, post-mortem delays exceeding 72 hours and formalin fixation exceeding ten years. The demographic and autopsy characteristics are presented in Supplementary Table 1.

Brainstem tissue blocks spanning from the rostral medulla oblongata (Obex +12 mm) to the caudal pons (Obex +18 mm) were selected (Fig. 1), as this region contains the 7N and is expected to include the hRTN. This Obex range was guided by a human brainstem atlas (Paxinos et al., 2020).

Tissue sectioning

Ten-micron serial sections were cut from 2-3 mm thick transverse oriented FFPE brainstem blocks, using a rotary microtome. We collected approximately 700 sections per case, extending rostrocaudally from Obex +18 to +12 mm (Fig. 1). One in every thirty serial sections (300 μm apart) was stained with Hematoxylin & Eosin (H&E), to obtain representative sections containing the 7N. After mapping the full extent of the 7N in H&E sections, four of every thirty serial sections were selected for immunohistochemistry using antibodies against ChAT, Phox2b+TH, Phox2b+PACAP and Phox2b+galanin, respectively (Supplementary Table 2).

Immunohistochemistry (IHC) Protocol for Human FFPE Brain Tissue

The IHC protocol was adapted from our previous methods (Huang et al., 2017), with necessary optimisations for Phox2b detailed in Supplementary Material 1. Antigen retrieval was performed using

heat-induced epitope retrieval (HIER) with citrate buffer (pH 6, #S2369, Agilent DAKO) for 15 minutes.

After a 2-hour block in 10% normal donkey serum (NDS) in tris buffered saline (TBS) with 0.3% Tween

20, sections were incubated overnight with goat anti-ChAT followed by a 2-hour incubation with

biotinylated secondary antibody, in 2% NDS /TBS with 0.1% Tween 20 (Table 2). Signal amplification

used the Vectastain Elite ABC HRP Kit (#PK-6100, Vector Laboratories), and visualisation was

achieved using 3,3'-diaminobenzidine (DAB Substrate Kit, #SK-4100, Vector Laboratories).

For dual labelling (Phox2b+TH, Phox2b+PACAP, or Phox2b+galanin), Phox2b was detected

first and visualised using DAB with nickel enhancement (black) (#SK-4100, Vector Laboratories). In

the sequential round of immunohistochemistry, the tissue was incubated with a primary antibody against

TH, PACAP, or galanin, followed by the corresponding secondary antibody and DAB (brown)

visualisation. After dehydration and clearing, sections were coverslipped using dibutylphthalate

polystyrene xylene (DPX). Antibody details and concentrations are listed in Table 2.

  Table 2. Primary and secondary antibodies used for immunohistochemistry (IHC)

Antibody	Source (Catalogue Number), RRID	Immunogen	Validation	Target & expected subcellular labelling	Dilution
Primary (1°) Antibodies					
Mouse monoclonal anti-Phox2b IgG antibody	Bio-Strategy Laboratory (sc-376997), RRID: AB_2813765	Human origin N-term amino acids 11-70	IHC: cross-reactivity with mouse brain tissue ISH: co-localisation with Phox2b mRNA labelled using RNAscope (Dereli et al., 2021)	Phox2b transcription factor (e.g. nucleus of RTN and C1 neurons)	1:1000
Chicken anti-Tyrosine Hydroxylase (TH)	Aves Labs (TYH-0020), RRID: AB_10013440	Human tyrosine hydroxylase (E.C.1.14.16.2) protein	IF: human gastric tissue (Eicher et al., 2022).	Catecholaminergic neurons (e.g. cytoplasm of C1 neurons)	1:6000
Goat anti-Choline Acetyltransferase (ChAT)	Millipore Sigma (AB144P), RRID: RRID AB_2079751	Human placental enzyme	IHC: rat brain tissue (1:200) (Bhagwandin et al., 2006); IHC: ape brain tissue (1:3000) (Williams et al., 2022)	Cholinergic neurons (e.g. cytoplasm of 7N)	1:1000
Rabbit anti-Galanin	Peninsula Laboratories International (T-4325), RRID: not provided	Synthetic peptide H-Gly-Trp-Thr-Leu-Asn-Ser-Ala-Gly-Tyr-Leu-Leu-GlyPro-His-Ala-Val-Gly-Asn-His-Arg-Ser-Phe-Ser-Asp-Lys-Asn-Gly-LeuThr-Ser-OH coupled to carrier protein.	IHC: human brain tissue (1:300)(Falkenstetter et al., 2020)	Galaninergic neurons (e.g. cytoplasm of NTS and RTN neurons)	1:300
Rabbit anti-PACAP	Abcam (ab216627), RRID: Not provided	Human ADCYAP1 aa 150 to C-terminus	IF: Pig nerve fibers in the aortic arch wall (1:4000) (Rytel et al., 2022); IHC: mouse brain tissue (1:700) (Hu et al., 2020).	PACAP neuropeptide (e.g. cytoplasm of NTS and RTN neurons)	1:300
Secondary (2°) Antibodies					
Biotin-SP-conjugated Donkey anti-Mouse IgG	Jackson ImmunoResearch (715-035-150), RRID: AB_2340770	Whole molecule mouse IgG	WB: Human HeLa cells (1:10000) (Srinivasan et al., 2024)	Mouse anti-Phox2b 1° Ab	1:500
Biotin-SP-conjugated Donkey anti-Chicken IgG	Jackson ImmunoResearch (703-065-155), RRID: AB_2313595	Whole molecule chicken IgY	IHC: mouse eyes (dilution not provided) (Little et al., 2020)	Chicken anti-TH 1° Ab	1:500
Biotin-SP-conjugated Donkey anti-Goat IgG	Jackson ImmunoResearch (705-065-147), RRID: AB_2340397	Whole molecule goat IgY	IHC: Mouse brain and spine tissue (1:500) (Agostinelli et al., 2021)	Goat anti-ChAT 1° Ab	1:500
Biotin-SP-conjugated Donkey anti-Rabbit IgG	Jackson ImmunoResearch (711-065-152), RRID: AB_2340593	Whole molecule rabbit IgY	WB: Human HeLa cells (1:10000) (Srinivasan et al., 2024)	Rabbit anti-PACAP 1° Ab	1:500

Microscopy and imaging

Slides were digitised using a slide scanner (Aperio AT2, Leica BioSystems) at 20× magnification. The scanned files were viewed using QuPath Software (Version 0.2.3) (Bankhead et al., 2017). Coronal hemi-sections approximating from Obex +12 to +18 mm were assessed.

The parafacial area was defined as follows: rostrocaudally, spans from the rostral ventrolateral medulla to the caudal ventrolateral pons, located adjacent to the facial motor nucleus (Obex +13 to +17 mm),dorsoventrally and mediolaterally, it is ventral to the spinal trigeminal nucleus, oral part (Sp5O), dorsal to ventral medullary surface, lateral to the intermediate reticular nucleus (IRt) and inferior olive (IO), and medial to the facial nerve (7n) (Fig. 1). Schematic diagrams were drawn using Adobe Illustrator 2024 (Adobe Inc., San Jose, CA, USA).

Fig. 1 Tissue sectioning and parafacial area localisation (a) Sagittal schematic of the adult human brainstem and thalamus indicating the angle of tissue sectioning and location spanning from Obex (indicated by blue arrow) +12 to +18 mm (between red lines). (b & c) Representative H&E stained caudal (Obex +14 mm) and rostral (Obex +16 mm) hemi-sections overlaid with brainstem schematics. (b) Caudally, the parafacial (red outline) is encapsulated by a region medial to sp5 and Sp5O, ventral to 7N, lateral to the SO and overlapping with C1 or A5 neurons (blue outline). (c) Rostrally, the parafacial area is medial to 7n, ventral to 7N, lateral to MSO and overlaps with C1 or A5 neurons (blue outline). Adapted from (Paxinos et al., 2020)

Quantitative analyses

The cell diameter, and area of the cell soma and nucleus of RTN neurons, were measured using QuPath

tools (See details in Supplementary Material 2). This data was collated using Microsoft Excel v14.0

(2010, Microsoft, USA), and exported into GraphPad Prism (Version 9, CA, USA).

Neurons with the following neurochemical signatures were counted:

1- Phox2b immunoreactivity (-ir) parafacial neurons, located ventrolateral to the 7N, as:

a. Phox2b-ir RTN neurons: Phox2b+ nucleus, TH- and ChAT- cytoplasm,

b. PACAP+/Phox2b+ parafacial neurons and PACAP-/Phox2b+ parafacial neurons,

c. Galanin+/Phox2b- parafacial neurons and galanin-/Phox2b+ parafacial neurons,

2- 7N neurons as ChAT+ cytoplasm

3- C1 neurons: Phox2b+ nucleus, TH+ cytoplasm (Kang et al., 2007, Agostinelli et al., 2023),

4- A5 neurons: Phox2b- nucleus, TH+ cytoplasm (Kang et al., 2007, Agostinelli et al., 2023).

Phox2b-negative RTN neurons were not investigated in this study. The RTN neurons referred in this

study were Phox2b-ir RTN neuron. Number of RTN neurons was calculated by applying the formula

derived from (Abercrombie, 1946):

Phox2b-ir RTN Cell Count (N) $= RTN\ (Phox2b - ir)\ hemisection\ cell\ count\ (n) \times \frac{T}{T+D} \times 2 \times 30$

N, Total Phox2b-ir RTN cell count; n, RTN cell count/hemi-section; T, section thickness; D, average

diameter of the cell.

Morphological data from the 2 adult cases was averaged and presented as mean ± standard deviation.

No statistical analysis was performed due to the study being an anatomical description of the hRTN,

without comparison between a control and experimental group.

Results

Immunoreactive labelling profile of antibodies used in this study

The subcellular localisation of labelling for each primary antibody was as expected in positive control

regions, including the 7N, NTS and C1 (Fig. 2) (Huang et al., 2017, Kordower et al., 1992, Levy et al.,

2019, Agostinelli et al., 2023). ChAT, TH, PACAP and galanin-ir were observed in the cytoplasm and

fibers, while Phox2b-ir was confined to the nucleus of neurons (Fig. 2).

Fig. 2 Exemplary H&E and primary antibody labelling. (a) H&E staining was performed to identify

7N neurons, based on their anatomical location. (b) ChAT-ir cytoplasm and fibers in motor neurons

within the 7N. (c) TH-ir cytoplasm and fibers in catecholaminergic C1 neurons. (d) Phox2b-ir nuclei in

NTS neurons. (e) PACAP-ir cytoplasm and fibers of hematoxylin counterstained NTS neurons. (f)

Galanin-ir cytoplasm and fibers in the NTS. Arrowheads point to examples of positive

immunoreactivity; arrowheads in (a) and (b) point to the same 7N cells

Phox2b-ir was subcategorised as either strong or weak (Supplementary Fig. 1), with cells in each

category counted separately before totals were combined (see Supplementary Table 3).

Neurochemical signature of putative hRTN neurons

Identification of RTN neurons according to their biochemical signature of Phox2b+ nucleus, TH-

/ChAT- cytoplasm, was clearly discernable from neighboring 7N (ChAT+) and overlapping C1 or A5

(TH+) neurons (Fig. 3a, e, f and Supplementary Fig. 2). TH+ fibers coursed through the RTN region

(Fig. 3a). In a high magnification image (Fig. 3b), the TH-positive axons were clearly delineated, and

TH+ boutons made close appositions with RTN neurons. The subcellular labelling of these

neurochemical markers in the RTN and nearby nuclei (7N, C1 or A5) is documented in Table 3. A few

Phox2b+/TH- neurons were also detected in Sp5O (Fig. 3c), IRt and parvocellular reticular nucleus

(PCRt) (Fig. 3d).

Fig. 3 RTN neurochemical signature and spatial relationship to 7N, C1 or A5. (a) Low-magnification image showing the anatomical locations of the RTN (red outline), C1 or A5 region (blue outline), Sp5O (black outline), and 7N (green outline) within the brainstem. (b) Dual immunohistochemistry for Phox2b and TH identified RTN neurons as Phox2b+ with nuclear localisation (asterisks), and TH- cytoplasm, distinguishing RTN neurons from C1 or A5 neurons. Numerous TH-ir fibers (arrowheads) were present in the RTN region and closely apposed Phox2b-ir neurons (asterisks). (c) Sp5O neurons exhibited Phox2b-ir (arrowhead). (d) A few sparsely distributed Phox2b+/TH- neurons were detected in the IRt, dorsomedial to the 7N (arrowhead). (e) A5 neuron expressing TH-ir in the cytoplasm. (f) Low power image illustrating the location of the RTN (red outline) ventral to the 7N (green outline) in the brainstem (Obex +15 mm). (g) Phox2b and ChAT double

immunohistochemical detection demonstrate RTN neurons as Phox2b+ and ChAT- (arrowheads). In

contrast, the 7N neurons (asterisk) were identified by their ChAT+ soma and relatively large diameter

Phox2b+ parafacial neurons are immunoreactive for galanin and PACAP

Although TH immunohistochemistry was not performed in combination with Phox2b/PACAP or

Phox2b/galanin labelling, our quantitative data show that 90% of Phox2b-ir neurons located ventral to

7N are devoid of TH-ir (Fig. 3b). This strongly suggests that these neurons are part of the RTN

population. Therefore, in this context, we confidently identified RTN neurons based on their Phox2b

expression and anatomical location within the parafacial region. Based on the double labelling for

Phox2b and galanin as well as the double labelling for Phox2b and PACAP, a total of 5295 Phox2b+

parafacial neurons were counted (Table 4).

Galanin-ir was detected in the soma and dendrites of Phox2b-ir parafacial neurons (Fig. 4a-c)

and represented 43% of total Phox2b-ir population (2262 galanin+/Phox2b+ neurons out of 5295

Phox2b+ parafacial neurons) (n=2, with 14 sections counted per case) (Supplementary Table 3).

Similarly, PACAP-ir was confined to cell soma and dendrites of 39% Phox2b-ir parafacial neurons

(2090 PACAP+/Phox2b+ parafacial neurons out of 5295 Phox2b-ir parafacial neurons, Fig. 4d-f,

Supplementary Table 3).

Fig. 4 Galanin and PACAP immunoreactivities are detectable in large subsets of Phox2b-ir parafacial neurons. (a and d) Low magnification image of the RTN region immediately ventral to the 7N. (b) Many neurons in the parafacial region are both galaninergic (brown cytoplasmic labelling) and Phox2b-ir (black nucleus labelling). (c) Magnification of inset in b) demonstrating a galanin+/Phox2b+ neuron (arrow) and a galanin-/Phox2b+ neuron (arrowhead). (e) A subgroup of neurons in the parafacial region display PACAP-ir (brown cytoplasmic labelling) and Phox2b-ir (black nucleus labelling). (f) Magnification of inset in (e) demonstrating PACAP+/Phox2b+ neuron (arrow) and PACAP-/Phox2b+ neurons (arrowheads)

Anatomical characteristics of the hRTN based on the distribution of Phox2b+/TH- neurons

Putative hRTN neurons were located ventral and ventrolateral to the 7N, lateral to the SO, dorsomedial

to the pontomedullary junction, medial to the Sp5O, spinal trigeminal tract (sp5) and the facial nerve,

and overlapping with C1 or A5 catecholaminergic neurons (Fig. 3).

RTN neurons were found from Obex +13 to +17 mm (Fig. 5 and Fig. 6) of the medulla

oblongata. At caudal levels, Obex +13 to +13.8 mm, RTN neurons were sparsely distributed (Fig. 5 and

Fig. 6a). By contrast, within the range of +13.8 to +14.8 mm, RTN neurons formed a distinct and

prominent cluster (Fig. 5), indicated by a concentrated presence of Phox2b+/TH- neurons. More

rostrally, (+14.8 to +16.8 mm), the RTN neurons were again more sparsely distributed, devoid of a

distinct cluster (Fig. 5 and Fig. 6c). Anatomically, rostral RTN neurons were distributed medial to the

facial nerve, ventrolateral to the 7N, dorsal to the ventral surface of medulla and lateral to the SO (Fig.

5).

Fig. 5 Schematic coronal hemi-sections illustrating the distribution of parafacial Phox2b+/TH-neurons, representative of the RTN (from Obex +13 to +17 mm). Red dots represent Phox2b+/TH- neurons. RTN neurons are located immediately ventral to the 7N, lateral to the SO, and medial to the facial nerve (rostrally) or spinal trigeminal nucleus (caudally). The most densely packed

266 cluster of RTN neurons is situated 14 mm rostral to Obex. Drawings were made with reference to

267 (Paxinos et al., 2020)

Obex +13.1 mm   Obex +15 mm   Obex +16.8 mm

Fig. 6 RTN neurons extend along the entire rostrocaudal length of the 7N. (a1 and a2) Caudal pole

of the 7N (Obex +13.1 mm). (a3) C1 neuron (TH-ir, asterisk) next to an RTN neuron (arrowhead). (b1

and b2) 7N neurons are distinguished from the RTN region by the green outline. (b3) RTN neurons

(Phox2b+/TH-, arrowheads) and TH-ir fibers (brown puncta) are notably abundant at this level of the

RTN. (c1 and c2) Rostral pole of the 7N (Obex +16.8 mm). (c3) Few RTN neurons (Phox2b+/TH-,

arrowhead) were present. An A5 neuron (Phox2b-/TH+, asterisk) is visible in the rostral RTN region

Morphology of hRTN neurons

The soma of RTN neurons was typically round, or fusiform, with one or two dendrites extending in

various directions (Fig. 4). RTN neurons were significantly smaller in size compared to 7N, C1 or A5

neurons, with an average soma area of 309.9 ± 61.8 μm² (n=241 cells), and an average nucleus area of

69.0 ± 23.3 μm² (Table 3).

281 Table 3. Morphological characteristics of RTN, 7N, C1 and A5 neurons, based on their
282 respective subcellular neurochemical signature

Cell Type	Morphology	Nucleus: Area (μm²)	Cell Area (μm²)	Cell Diameter (μm)	Subcellular neurochemical profile		
					Cytoplasm	Nucleus	Axon
7N neurons (n=68, 1 section/case)	Large multipolar neurons with short dendrites	274.6 ± 108	1194.0 ± 371.8	41.8 ± 10.7	ChAT+, TH-, PACAP+/-, galanin-	Phox2b+/-	ChAT+
RTN neurons (n=241, 3 sections/case)	Medium round or fusiform neurons with short dendrites	69.0 ± 23.3	309.9 ± 61.8	19.9 ± 4.0	ChAT-, TH-, PACAP+/-, Galanin+/-	Phox2b+	Not detected
C1 neurons (n=38, 4 sections/case)	Medium multipolar neuron with short dendrites and long projecting axons	98.0 ± 51.2	528.2 ± 165.2	35.1±8.9	ChAT-, TH+, PACAP+/-, galanin-	Phox2b+	TH+
A5 neurons (n=12, 4 sections/case)	Medium multipolar neuron with short dendrites and long projecting axons	101.3 ± 44.9	468.7 ± 169.4	33.9 ±11.7	ChAT-, TH+, PACAP+/-, galanin-	Phox2b-	TH+

Mean ± standard deviation.

284 Quantification of hRTN neurons

Based on Phox2b+/TH- immunoreactivity, the number of putative RTN neurons ranged from 5 to 65

per hemi-section, with a peak observed between Obex +14.0 mm and +14.3 mm (Fig. 7a). Considering

both weak and strong Phox2b-ir, the total number of RTN neurons was 5300 and 4903 (n=2 cases,

Table 4), spanning 4 mm in rostrocaudal length (Fig. 7a) and ~1.5-2 mm in dorsoventral length (Fig. 5)

The rostrocaudal distribution of galanin+/Phox2b+ and PACAP+/Phox2b+ parafacial neurons

followed the same trend, showing a peak in counts at Obex 14 mm and spanning 4 mm in rostrocaudal

length (Fig. 7b and 7c). From these double labelled sections, a total of 5295 Phox2b-ir parafacial

neurons were counted (Table 4). The galanin population was 43% of the total Phox2b-ir population

while PACAP was 39% (Table 4).

a.

b.

c.

Fig. 7 Cell count of the RTN, 7N, C1 and A5 neurons. (a) Rostral-caudal distribution and quantification RTN (Phox2b+/TH-), C1 (Phox2b+/TH+) and A5 (Phox2b-/TH+) neurons within the region of interest specified in Fig. 1b and Fig. 1c. Additionally, total 7N (ChAT+) neurons were counted. The rostral-caudal distribution of (b) Galanin+/Phox2b+ neurons and (c) PACAP+/Phox2b+ neurons, in the RTN region. The curves represent the average number of cells/hemi-section (n=2). The vertical bars indicate interindividual variability in RTN, 7N, C1, and A5 neuron counts observed between the two human cases.

Table 4. Total Phox2b-ir parafacial neurons and the proportion that are galanin-ir or PACAP-ir*

	Phox2b+ RTN neurons *	Phox2b+ parafacial neurons	Phox2b+/ PACAP+ parafacial neurons	Phox2b+/gala nin+ parafacial neurons	% of Phox2b+ parafacial PACAP+ neurons	% of Phox2b+ parafacial galanin+ neurons
Case 1	5300	5960	2420	2985	41	50
Case 2	4903	4630	1760	1540	38	33
Average	5102	5295	2090	2262	39	43

*Refer to Supplement Table 3 for the breakdown of the counts according to labelling intensity.

Discussion

This study provides a comprehensive characterisation of the adult human RTN using histological and neuroanatomical means. Leveraging previous reports from infant, macaque and rodent studies, we identified a large cluster of Phox2b+/TH- neurons ventral to the 7N, aligning with the expected location of the RTN. We present the first in-depth analysis of Phox2b+ RTN neurons, detailing their rostrocaudal distribution, cell counts, morphology and expression of respiratory-related neuropeptides - galanin and PACAP. We also demonstrate puncta from TH+ fibers in close apposition to RTN neurons. These findings establish a crucial baseline for assessing age-related or disease-related changes in the RTN, such as neuronal loss, or altered neuropeptide expression, in respiratory and neurological disorders.

Adult human RTN neurons are Phox2b positive and subsets express galanin and PACAP immunoreactivity

In this study, the adult hRTN was identified using histological criteria established for identification of RTN neurons in rodents; namely, parafacial neurons that express the transcription factor Phox2b and are devoid of TH and ChAT immunoreactivity (Guyenet and Bayliss, 2022). We identified a highly circumscribed cluster of Phox2b neurons in adult hRTN, in agreement with findings in infants. Approximately 43% of these neurons co-expressed galanin, closely aligning with findings from adult rats (Stornetta et al., 2009). Furthermore, the number of galanin-ir neurons we identified (2,262) is comparable to the ~2,400 reported in the macaque RTN (Levy et al., 2019). Our study is the first to demonstrate PACAP expression in hRTN, with 40% of Phox2b-ir parafacial neurons co-expressing PACAP. This proportion is substantially lower than what has been reported in adult mice, where

PACAP is expressed in 100% of RTN chemoreceptors in neonates and 88% in adulthood (Shi et al., 2017, Shi et al., 2021).

The location of hRTN neurons in the adult is highly consistent with the distribution reported in human infants and other species

In this study, the spatial distribution of adult hRTN neurons closely aligns with findings from rodents, non-human primates, and human infants (Levy et al., 2019, Levy et al., 2021, Lavezzi et al., 2012, Rudzinski and Kapur, 2010, Shi et al., 2017). Dorsoventrally, the hRTN neurons are situated dorsal to the pontomedullary junction, lateral to the superior olive, and ventrolateral to the 7N. Rostrocaudally, hRTN neurons extended from Obex +13 to +17 mm. A previous study in adults reported PPGAL+/VGlut2+ putative hRTN neurons extending from Obex +14 to +17 mm (Levy et al., 2021). This shorter rostrocaudal extent observed is potentially due to discrepancies between versions of human brainstem atlases. Notably, the Paxinos 2012 atlas used by Levy et al, describes the facial motor nucleus as 3 mm long, while the updated 2020 version used in our study reports it as 4 mm in rostrocaudal length. Nevertheless, both studies demonstrate close proximity between the RTN and 7N throughout their rostrocaudal extent, underscoring the reliability of the findings.

Another key difference between our study and Levy's (Levy et al., 2021) is the biomarkers used to identify RTN neurons. While our study successfully detected Phox2b-ir, Levy et al. relied on vGlut2 and PPGAL co-expression, however, they did not define the non-galaninergic RTN neurons (Levy et al., 2021) leading to an underestimation of RTN neurons. Thus, our dataset represents the most complete count of RTN neurons to date including both galanin+ and galanin- RTN neurons.

The adult human contains more RTN neurons than adult rodents and macaque

We identified approximately 5000 Phox2b+/TH- neurons within the putative hRTN, bilaterally, a count significantly higher than the approximately 980 Phox2b-ir neurons reported in human infants (Lavezzi et al., 2012). The discrepancy may stem from differences in immunohistochemistry protocol, or the exclusion of "weakly" Phox2b-ir neurons from the RTN population in earlier studies (Lavezzi et al., 2012, Levy et al., 2019). However, even when counting only strongly Phox2b-ir neurons, we found an

average of 2530 per case bilaterally, which is still significantly greater than the 980 reported by Lavezzi et al. (Lavezzi et al., 2012). Since RTN neuron numbers remain stable from late prenatal stages to adulthood in rodents, developmental stage is unlikely the cause. Direct comparisons across studies would require consistent tissue preservation, antigen retrieval and immunohistochemical protocols. Species comparisons report bilateral RTN counts of ~2,000 in rats (Stornetta et al., 2009, Takakura et al., 2008), ~700 in mice (Lazarenko et al., 2009, Shi et al., 2017), and ~2,400 galaninergic neurons in macaques (Levy et al., 2019). Our higher count in humans likely reflects both true interspecies differences and methodological factors such as more extensive serial sectioning and variations in RTN rostrocaudal length across atlases (Paxinos, 2012, Paxinos et al., 2020). Some earlier studies did not assess serial sections and based total cell counts on only 1-2 sections per case (Lavezzi et al., 2012, Rudzinski and Kapur, 2010). Notably, our female case had 29% more RTN neurons than the male, suggesting possible sex or inter-individual differences that warrant further investigation.

C1-RTN circuit and its role in breathing control: evidence of neuronal interactions

In this study, we show that TH-ir presumptive C1 neurons are intermingled with RTN neurons, and our data suggest close appositions between TH-ir terminals and RTN neurons. Rodent studies have demonstrated that the RTN is dense with PNMT-ir terminals (Rosin et al., 2006) and vigorously activated by adrenaline via α1 adrenoceptors (Oliveira et al., 2016). Furthermore, optogenetic stimulation of C1 neurons in rats produces respiratory effects similar to RTN photo-stimulation (Souza et al., 2020), suggesting a functional C1-RTN circuit in the central control of breathing. These findings suggest that hRTN neurons receive projections from C1 catecholaminergic neurons, however, further studies are needed to confirm the origin of the TH-ir axons and boutons apposing hRTN neurons.

Factors affecting RTN neuronal counts and future research directions

The RTN neuronal cell count may be influenced by several factors. First, the neurochemical signature used to define RTN neurons could inadvertently include Phox2b+/TH- PCRt, IRt or Sp5O neurons located in the region dorsal and medial to 7N (Fig. 3d), However these were excluded from hRTN because their distribution was anatomically discontinuous from the RTN cluster. Secondly, although

this study identified hRTN neurons based on the neurochemical signature established in rodent

functional studies (Lazarenko et al., 2009), further research is needed to confirm that these

Phox2b+/TH- neurons are indeed chemosensory (Shi et al., 2017). Third, we have reported RTN neuron

counts in older adults, and it is uncertain whether Phox2b-ir neurons undergo age-related degeneration.

If such degeneration occurs, our quantitative results may underestimate the number of Phox2b-ir RTN

neurons in young adults. Finally, although this study was limited by a small sample size, a key strength

lies in our thorough analysis of the entire length of the RTN (14 sections per case, spanning 4mm

rostrocaudally), rather than focussing on a single plane as is common in studies using post-mortem

human brain tissue. A further strength is the consistent rostral-caudal distribution and the quantified

peaks of RTN neurons observed across the two cases. Future studies with a larger sample size are

needed to evaluate changes in this neuronal population with age, sex and disease, strengthening the

robustness and generalisability of our findings. Our study lays the groundwork for these future

investigations and for future studies investigating changes in disorders affecting central respiratory

chemoreception.

Conclusion

This study mapped the full extent of the adult hRTN using established neurochemical and anatomical

markers, locating it ventral to 7N, lateral to the superior olive, and overlapped with C1 or A5

catecholaminergic neurons. Extending rostrocaudally from Obex +13 to +17 mm, the hRTN contains

around 5000 neurons bilaterally - comprising 90% of Phox2b-ir neurons in the parafacial area -

surrounded by dense TH-ir fibers. Approximately 40% of these neurons express galanin and PACAP.

The data herein provides baseline values for future studies in clinical contexts such as CCHS and sleep

apnoea.

Acknowledgements

This work was supported by grants from Australian Research Council (grant no. DP180101890).

Tissues were received from the Sydney Brain Bank which is supported by Neuroscience Research

Australia and a special gift in memory of Jim Raftos from the Shaw family. We thank Prof Patrice G

Guyenet and Prof Ruth Stornetta for providing feedback on an advanced version of the manuscript.

References

ABBOTT, S. B., BURKE, P. G. & PILOWSKY, P. M. 2009. Galanin microinjection into the
PreBotzinger or the Botzinger Complex terminates central inspiratory activity and reduces responses to
hypoxia and hypercapnia in rat. *Respir Physiol Neurobiol,* 167**,** 299-
306.http://dx.doi.org/10.1016/j.resp.2009.06.003
ABERCROMBIE, M. 1946. Estimation of nuclear population from microtome sections. *The
Anatomical Record,* 94**,** 239-247.http://dx.doi.org/https://doi.org/10.1002/ar.1090940210
AGOSTINELLI, L. J., SEAMAN, S. C., SAPER, C. B., et al. 2023. Human Brainstem and Cerebellum
Atlas: Chemoarchitecture and Cytoarchitecture Paired to MRI. *J Neurosci,* 43**,** 221-
239.http://dx.doi.org/10.1523/JNEUROSCI.0587-22.2022
AMIEL, J., DUBREUIL, V., RAMANANTSOA, N., et al. 2009. PHOX2B in respiratory control:
lessons from congenital central hypoventilation syndrome and its mouse models. *Respir Physiol
Neurobiol,* 168**,** 125-32.http://dx.doi.org/10.1016/j.resp.2009.03.005
BANKHEAD, P., LOUGHREY, M. B., FERNANDEZ, J. A., et al. 2017. QuPath: Open source
software for digital pathology image analysis. *Sci Rep,* 7**,** 16878.http://dx.doi.org/10.1038/s41598-017-
17204-5
BOCHORISHVILI, G., STORNETTA, R. L., COATES, M. B., et al. 2012. Pre-Bötzinger complex
receives glutamatergic innervation from galaninergic and other retrotrapezoid nucleus neurons. *J Comp
Neurol,* 520**,** 1047-61.http://dx.doi.org/10.1002/cne.22769
DEGL'INNOCENTI, D., BECATTI, M., PERUZZI, M., et al. 2018. Systemic oxidative stress in
congenital central hypoventilation syndrome. *Eur Respir J,*
52.http://dx.doi.org/10.1183/13993003.01497-2018
DERELI, A. S., OH, A. Y. S., MCMULLAN, S., et al. 2024. Galaninergic and hypercapnia-activated
neuronal projections to the ventral respiratory column. *Brain Struct Funct,* 229**,** 1121-
1142.http://dx.doi.org/10.1007/s00429-024-02782-8
DERELI, A. S., YASEEN, Z., CARRIVE, P., et al. 2019. Adaptation of Respiratory-Related Brain
Regions to Long-Term Hypercapnia: Focus on Neuropeptides in the RTN. *Front Neurosci,* 13**,**
1343.http://dx.doi.org/10.3389/fnins.2019.01343
GORIDIS, C. & BRUNET, J. F. 2010. Central chemoreception: lessons from mouse and human
genetics. *Respir Physiol Neurobiol,* 173**,** 312-21.http://dx.doi.org/10.1016/j.resp.2010.03.014
GUYENET, P. G. & BAYLISS, D. A. 2015. Neural Control of Breathing and CO2 Homeostasis.
Neuron, 87**,** 946-61.http://dx.doi.org/10.1016/j.neuron.2015.08.001
GUYENET, P. G. & BAYLISS, D. A. 2022. Central respiratory chemoreception. *Handb Clin Neurol,*
188**,** 37-72.http://dx.doi.org/10.1016/b978-0-323-91534-2.00007-2
GUYENET, P. G., STORNETTA, R. L., SOUZA, G., et al. 2019. The Retrotrapezoid Nucleus: Central
Chemoreceptor and Regulator of Breathing Automaticity. *Trends Neurosci,* 42**,** 807-
824.http://dx.doi.org/10.1016/j.tins.2019.09.002
HUANG, J., WATERS, K. A. & MACHAALANI, R. 2017. Pituitary adenylate cyclase activating
polypeptide (PACAP) and its receptor 1 (PAC1) in the human infant brain and changes in the Sudden
Infant Death Syndrome (SIDS). *Neurobiol Dis,* 103**,** 70-77.http://dx.doi.org/10.1016/j.nbd.2017.04.002
IKEDA, K., KAWAKAMI, K., ONIMARU, H., et al. 2017. The respiratory control mechanisms in the
brainstem and spinal cord: integrative views of the neuroanatomy and neurophysiology. *J Physiol Sci,*
67**,** 45-62.http://dx.doi.org/10.1007/s12576-016-0475-y
KANG, B. J., CHANG, D. A., MACKAY, D. D., et al. 2007. Central nervous system distribution of
the transcription factor Phox2b in the adult rat. *J Comp Neurol,* 503**,** 627-
41.http://dx.doi.org/10.1002/cne.21409
KASI, A. S., KUN, S. S., KEENS, T. G., et al. 2018. Adult With PHOX2B Mutation and Late-Onset
Congenital Central Hypoventilation Syndrome. *J Clin Sleep Med,* 14**,** 2079-
2081.http://dx.doi.org/10.5664/jcsm.7542

KORDOWER, J. H., LE, H. K. & MUFSON, E. J. 1992. Galanin immunoreactivity in the primate
central nervous system. *J Comp Neurol,* 319**,** 479-500.http://dx.doi.org/10.1002/cne.903190403
LAVEZZI, A. M., WEESE-MAYER, D. E., YU, M. Y., et al. 2012. Developmental alterations of the
respiratory human retrotrapezoid nucleus in sudden unexplained fetal and infant death. *Auton Neurosci,*
170**,** 12-9.http://dx.doi.org/10.1016/j.autneu.2012.06.005
LAZARENKO, R. M., MILNER, T. A., DEPUY, S. D., et al. 2009. Acid sensitivity and ultrastructure
of the retrotrapezoid nucleus in Phox2b-EGFP transgenic mice. *J Comp Neurol,* 517**,** 69-
86.http://dx.doi.org/10.1002/cne.22136
LEVY, J., DROZ-BARTHOLET, F., ACHOUR, M., et al. 2021. Parafacial neurons in the human
brainstem express specific markers for neurons of the retrotrapezoid nucleus. *J Comp Neurol,* 529**,**
3313-3320.http://dx.doi.org/10.1002/cne.25191
LEVY, J., FACCHINETTI, P., JAN, C., et al. 2019. Tridimensional mapping of Phox2b expressing
neurons in the brainstem of adult Macaca fascicularis and identification of the retrotrapezoid nucleus.
J Comp Neurol, 527**,** 2875-2884.http://dx.doi.org/10.1002/cne.24713
MEYLEMANS, A., DEPUYDT, P., DE BAERE, E., et al. 2021. Adult-onset congenital central
hypoventilation syndrome due to PHOX2B mutation. *Acta Neurol Belg,* 121**,** 23-
35.http://dx.doi.org/10.1007/s13760-020-01363-w
OLIVEIRA, L. M., MOREIRA, T. S., KUO, F. S., et al. 2016. alpha1- and alpha2-adrenergic receptors
in the retrotrapezoid nucleus differentially regulate breathing in anesthetized adult rats. *J Neurophysiol,*
116**,** 1036-48.http://dx.doi.org/10.1152/jn.00023.2016
PAXINOS, G., FURLONG, T. & WATSON, C. 2020. *Human brainstem: cytoarchitecture,*
chemoarchitecture, myeloarchitecture, Academic Press.
PAXINOS, G., XU-FENG, H., SENGUL, G., & WATSON, C 2012. Organization of brainstem nuclei.
In The human nervous system *Elsevier***,** 260-327
ROSIN, D. L., CHANG, D. A. & GUYENET, P. G. 2006. Afferent and efferent connections of the rat
retrotrapezoid nucleus. *J Comp Neurol,* 499**,** 64-89.http://dx.doi.org/10.1002/cne.21105
RUDZINSKI, E. & KAPUR, R. P. 2010. PHOX2B immunolocalization of the candidate human
retrotrapezoid nucleus. *Pediatr Dev Pathol,* 13**,** 291-9.http://dx.doi.org/10.2350/09-07-0682-OA.1
SHI, Y., STORNETTA, D. S., REKLOW, R. J., et al. 2021. A brainstem peptide system activated at
birth protects postnatal breathing. *Nature,* 589**,** 426-430.http://dx.doi.org/10.1038/s41586-020-2991-4
SHI, Y., STORNETTA, R. L., STORNETTA, D. S., et al. 2017. Neuromedin B Expression Defines the
Mouse Retrotrapezoid Nucleus. *J Neurosci,* 37**,** 11744-
11757.http://dx.doi.org/10.1523/JNEUROSCI.2055-17.2017
SOUZA, G., STORNETTA, R. L., STORNETTA, D. S., et al. 2020. Differential Contribution of the
Retrotrapezoid Nucleus and C1 Neurons to Active Expiration and Arousal in Rats. *J Neurosci,* 40**,**
8683-8697.http://dx.doi.org/10.1523/jneurosci.1006-20.2020
STORNETTA, R. L., SPIROVSKI, D., MOREIRA, T. S., et al. 2009. Galanin is a selective marker of
the retrotrapezoid nucleus in rats. *J Comp Neurol,* 512**,** 373-83.http://dx.doi.org/10.1002/cne.21897
TAKAKURA, A. C., MOREIRA, T. S., STORNETTA, R. L., et al. 2008. Selective lesion of
retrotrapezoid Phox2b-expressing neurons raises the apnoeic threshold in rats. *J Physiol,* 586**,** 2975-
91.http://dx.doi.org/10.1113/jphysiol.2008.153163

Statements & Declarations

Funding

This work was supported by Australian Research Council (Grant number DP180101890, NNK),

Research Training Program Domestic Fee Offset (RSAI8000, YL) and Research Training Program

Domestic Stipend (RSAP1000, YL).

Competing Interests

The authors do not have relevant financial or non-financial interests to disclose.

Author Contributions

This study was designed by YL, CES, RM and NNK. IM trained YL in tissue preparation and immunohistochemistry on FFPE tissues. All authors contributed to the methodology validation and experiment troubleshooting. The tissue preparation, experiments, data collection, data analysis and first manuscript draft were produced by YL. NNK, YL, RM and CES revised and modified manuscript drafts.

Data Availability

The datasets generated during the current study are available from the corresponding author on reasonable request.

Ethics approval

Donor tissue was previously collected by the Sydney Brain Bank under UNSW ethics project no. HC#200026. This research project (Project no. HC#200901) was approved by the Human Research Ethics Advisory Panel of the University of New South Wales.

Consent to participate

Informed consent was obtained from individual participants included in the study prior to tissue donation.

Consent to publish

The authors affirm that human research participants provided informed consent for use of their data for publication.

a. Phox2b & TH

Sp5O

7N

RTN

C1 or A5

D

M

800μm

b

*

*

*

20μm

c

20μm

d

20μm

e

20μm

f. Phox2b & ChAT

7N

RTN

g

500μm

g

*

100μm

a. Phox2b & galanin
7N
400µm

d. Phox2b & PACAP
7N
400µm
D
M

b. Phox2b & galanin
100µm

e. Phox2b & PACAP
100µm

c. Phox2b & galanin
50µm

f. Phox2b & PACAP
50µm

Obex +13mm

Obex +13.5mm

Obex +14mm

Obex +14.5mm

Obex +15mm

Obex +15.5mm

Obex +16mm

Obex +16.5mm

Obex +17mm

Obex +13.1 mm

Obex +15 mm

Obex +16.8 mm

a.

b.

c.

Developmental age, number of cases, Sex ratio	Tissue Frozen or FFPE, and plane of cut (coronal vs sagittal)	Sections/case analysed; Obex/bregma level	Antibody/probe details & dilution	RTN Phox2b subcellular labelling pattern & anatomical location	Number of RTN neurons (bilateral)	Reference
Human fetuses and infants (19-44w, n=8)	5 µm coronal (n=5) and sagittal (n=3) FFPE sections	1-2 sections/case (level not specified but in the presence of the 7N).	Rabbit anti-Phox2b (gift of J.-F. Brunet, 1:1000). NK1R (Abcam, 1:200); TH (Chemicon, 1:1000)	IHC: cell nucleus. RTN was ventral to 7N and lateral to the SO in the pontomedullary junction.	Not provided	(Rudzinski and Kapur, 2010)
Human fetuses (35-40w, n=9) and infant (2-8 months, n=11)	4 µm coronal FFPE sections	3-6 sections/case: 1-2 sections from each brainstem level of upper pons, upper medulla and from Obex -2/3 mm to +2/3 mm.	Rabbit anti-Phox2b (gift of J.-F. Brunet, 1:750).	IHC: cell nucleus and cytoplasm. RTN located between the 7N and SO in caudal pons.	980 ± 23 Phox2b-ir neurons. However, equation applied not provided	(Lavezzi et al., 2012)
Macaca fascicularis (5- & 6-years old males and 16-year-old female, n=2 IHC; n=1 ISH)	40 µm coronal frozen and paraformaldehyde fixed free-floating sections	60 sections/case (1/8 serial sections); Obex -1.40mm to +13mm.	IHC: Rabbit anti-Phox2b antibody (1:2,000; gift from J.-F. Brunet) ISH: PPGAL and Phox2b mRNAs	IHC: subcellular location of Phox2b immunoreactivity not described. Parafacial region located ventral and lateral to 7N; ISH: cell cytoplasm. PPGAL+/Phox2b+ RTN neurons located caudal, ventral and lateral to 7N spanning from Obex +4.56 to +10.16 mm.	2400 PPGAL+/ Phox2b+ neurons. PPGAL-/Phox2b neurons were not included. Calculated as an estimation by automated method yet cautions on possibility of double counting.	(Levy et al., 2019)
Human adults (n=2)	10 µm coronal FFPE sections	60 sections/case (1/10 serial sections), Obex +14 mm to +17 mm.	mRNA probes (targeting 237–492 sequence of the galanin message-associated peptide mRNA (i.e. PPGAL) and 469–1173 sequence of the vGlut2 mRNA (SLC17A6) and NMB.	IHC: N/A since Phox2b did not work. ISH: N/A since Phox2b did not work. Based on PPGAL, SLC17A6 and NMB, RTN located in parafacial region.	Not provided	(Levy et al., 2021)
Human adults (n=2)	10 µm coronal FFPE sections	90 sections/case (5/30 serial sections), Obex +12 mm to +18 mm.	Mouse anti-Phox2b (Bio-Strategy Laboratory, cat#sc-376997, 1:1000)	IHC: cell nucleus RTN was identified ventral to 7N, lateral to the SO and medial to facial nerve and Sp5O.	Approximately 5000 Phox2b+TH-ChAT- RTN neurons.	Study herein

Antibody	Source (Catalogue Number), RRID	Immunogen	Validation	Target & expected subcellular labelling	Dilution
Primary (1°) Antibodies					
Mouse monoclonal anti-Phox2b IgG antibody	Bio-Strategy Laboratory (sc-376997), RRID: AB_2813765	Human origin N-term amino acids 11-70	IHC: cross-reactivity with mouse brain tissue ISH: co-localisation with Phox2b mRNA labelled using RNAscope ([Dereli et al., 2021](#))	Phox2b transcription factor (e.g. nucleus of RTN and C1 neurons)	1:1000
Chicken anti-Tyrosine Hydroxylase (TH)	Aves Labs (TYH-0020), RRID: AB_10013440	Human tyrosine hydroxylase (E.C.1.14.16.2) protein	IF: human gastric tissue ([Eicher et al., 2022](#)).	Catecholaminergic neurons (e.g. cytoplasm of C1 neurons)	1:6000
Goat anti-Choline Acetyltransferase (ChAT)	Millipore Sigma (AB144P), RRID: RRID AB_2079751	Human placental enzyme	IHC: rat brain tissue (1:200) ([Bhagwandin et al., 2006](#)); IHC: ape brain tissue (1:3000) ([Williams et al., 2022](#))	Cholinergic neurons (e.g. cytoplasm of 7N)	1:1000
Rabbit anti-Galanin	Peninsula Laboratories International (T-4325), RRID: not provided	Synthetic peptide H-Gly-Trp-Thr-Leu-Asn-Ser-Ala-Gly-Tyr-Leu-Leu-GlyPro-His-Ala-Val-Gly-Asn-His-Arg-Ser-Phe-Ser-Asp-Lys-Asn-Gly-LeuThr-Ser-OH coupled to carrier protein.	IHC: human brain tissue (1:300)([Falkenstetter et al., 2020](#))	Galaninergic neurons (e.g. cytoplasm of NTS and RTN neurons)	1:300
Rabbit anti-PACAP	Abcam (ab216627), RRID: Not provided	Human ADCYAP1 aa 150 to C-terminus	IF: Pig nerve fibers in the aortic arch wall (1:4000) ([Rytel et al., 2022](#)); IHC: mouse brain tissue (1:700) ([Hu et al., 2020](#)).	PACAP neuropeptide (e.g. cytoplasm of NTS and RTN neurons)	1:300
Secondary (2°) Antibodies					
Biotin-SP-conjugated Donkey anti-Mouse IgG	Jackson ImmunoResearch (715-035-150), RRID: AB_2340770	Whole molecule mouse IgG	WB: Human HeLa cells (1:10000) ([Srinivasan et al., 2024](#))	Mouse anti-Phox2b 1° Ab	1:500
Biotin-SP-conjugated Donkey anti-Chicken IgG	Jackson ImmunoResearch (703-065-155), RRID: AB_2313595	Whole molecule chicken IgY	IHC: mouse eyes (dilution not provided) ([Little et al., 2020](#))	Chicken anti-TH 1° Ab	1:500
Biotin-SP-conjugated Donkey anti-Goat IgG	Jackson ImmunoResearch (705-065-147), RRID: AB_2340397	Whole molecule goat IgY	IHC: Mouse brain and spine tissue (1:500) ([Agostinelli et al., 2021](#))	Goat anti-ChAT 1° Ab	1:500
Biotin-SP-conjugated Donkey anti-Rabbit IgG	Jackson ImmunoResearch (711-065-152), RRID: AB_2340593	Whole molecule rabbit IgY	WB: Human HeLa cells (1:10000) ([Srinivasan et al., 2024](#))	Rabbit anti-PACAP 1° Ab	1:500

Cell Type	Morphology	Nucleus: Area (μm^2)	Cell Area (μm^2)	Cell Diameter (μm)	Subcellular neurochemical profile		
					Cytoplasm	Nucleus	Axon
7N neurons (n=68, 1 section/case)	Large multipolar neurons with short dendrites	274.6 ± 108	1194.0 ± 371.8	41.8 ± 10.7	ChAT+, TH-, PACAP+/-, galanin-	Phox2b+/-	ChAT+
RTN neurons (n=241, 3 sections/case)	Medium round or fusiform neurons with short dendrites	69.0 ± 23.3	309.9 ± 61.8	19.9 ±4.0	ChAT-, TH-, PACAP+/-, Galanin+/-	Phox2b+	Not detected
C1 neurons (n=38, 4 sections/case)	Medium multipolar neuron with short dendrites and long projecting axons	98.0 ± 51.2	528.2 ± 165.2	35.1±8.9	ChAT-, TH+, PACAP+/-, galanin-	Phox2b+	TH+
A5 neurons (n=12, 4 sections/case)	Medium multipolar neuron with short dendrites and long projecting axons	101.3 ± 44.9	468.7 ± 169.4	33.9 ±11.7	ChAT-, TH+, PACAP+/-, galanin-	Phox2b-	TH+

Mean ± standard deviation.

	Phox2b+ RTN neurons *	Phox2b+ parafacial neurons	Phox2b+/ PACAP+ parafacial neurons	Phox2b+/gala nin+ parafacial neurons	% of Phox2b+ parafacial PACAP+ neurons	% of Phox2b+ parafacial galanin+ neurons
Case 1	5300	5960	2420	2985	41	50
Case 2	4903	4630	1760	1540	38	33
Average	5102	5295	2090	2262	39	43

a. Phox2b

50μm

b.

a. Obex + 13.2mm
7N
RTN
a2
a1
D
M
400µm

a1
20µm

a2
20µm

b. Obex + 14.5mm
7N
b3
b2
b1
RTN
400µm

b1
20µm

b2
20µm

b3
20µm

c. Obex + 16.2mm
7N
c1
RTN
400µm

c1
20µm