1

Optimizing Single-Cell Long-Read Sequencing for Enhanced Isoform Detection

in Pancreatic Islets

Running title: Single-cell Long-Read Sequencing in Pancreatic Islets

Maria S. Hansen, Christopher J. Hill, Lori Sussel, Kristen L. Wells*

Barbara Davis Center, University of Colorado Anschutz Medical Campus, Aurora CO 80045

*Corresponding Author: Kristen.wells-wrasman@cuanschutz.edu

Keywords: single cell long read RNA-sequencing, transcriptomics, islet biology, RNA

isoforms, RNA splicing

Word count: 4,218

Figures: 3

Tables: 4

Abstract

Alternative splicing is an essential mechanism for generating protein diversity by producing distinct isoforms from a single gene. Dysregulation of splicing that affects pancreatic function and immune tolerance has been linked to both type 1 and type 2 diabetes. Next-generation sequencing technologies, with their short read lengths, are limited in their ability to accurately detect splice variants. Long-read sequencing technologies offer the potential to overcome these limitations by providing full-length transcript information; however, their application in single-cell RNA sequencing has been hindered by technical challenges, including insufficient read lengths and higher error rates. Furthermore, cell types that produce high levels of a single transcript, such as islet endocrine cells, can obscure identification of lower abundance transcripts. In this study, we optimized a protocol for single-cell long-read sequencing in pancreatic islets to improve read length and transcript detection. Our findings demonstrate that 5' library preparation protocols outperform 3' protocols, resulting in better transcript identification. Furthermore, we show that targeted depletion of insulin transcripts enhances the detection of informative reads, highlighting the utility of transcript depletion strategies. This optimized protocol enables isoform-specific gene expression analysis and reveals differential transcript usage across the various cell types in pancreatic islets. By leveraging this approach, we gain deeper insights into the transcriptomic complexity and cellular heterogeneity within pancreatic islets.

Article Highlights

• This study addresses the limitations of current single-cell long-read RNA-sequencing

(sclrRNA-seq) technologies in detecting full-length transcripts and isoform diversity,

particularly in pancreatic islets.

• We demonstrate that optimizing single-cell library preparation protocols

reproducibly enhances read length and transcript identification in pancreatic islets.

• 5' capture methods, combined with targeted insulin depletion and extended reverse

transcription, significantly improved read length and isoform detection compared to

standard protocols, while maximizing the number of informative reads.

• These improvements yield longer reads in single-cell experiments, substantially

enhancing transcript identification and enabling more accurate analysis of isoform

diversity.

Introduction

Alternative splicing (AS) plays a critical role in generating protein diversity from the

~22,000 known protein-coding genes, leading to the production of over 140,000 distinct

transcripts [1]. This process allows for the generation of proteins with different amino acid

sequences, impacting their functions and localization within the cell, and allowing them to

respond readily to changes in the environment [2, 3]. Splicing dysregulation is a key factor in

many diseases, including diabetes, either due to inherent mutations in splice sites or RNA-

binding proteins (RBPs), or in response to changes in environmental conditions, such as

inflammatory stress or hyperglycemia [4, 5]. In the context of type 1 diabetes (T1D), diversity

in isoform expression has significant implications for pancreatic function and immune

tolerance. For example, differential isoform expression of autoantigens IA-2 and G6pc2

between the pancreas and thymus has been proposed to contribute to the generation of

autoreactive T cells in T1D [6, 7]. Furthermore, dysregulated splicing events have been

observed in islets from individuals with type 2 diabetes (T2D), underscoring the importance

of splicing regulation in maintaining proper cellular function and immune homeostasis [5].

As one specific example, SNAP-25, a component of the SNARE complex responsible for

vesicle fusion and exocytosis, exists as two isoforms (SNAP-25a and SNAP-25b). In SNAP-

25b-deficient mice, $[Ca^{2+}]$ elevations are prematurely activated and delayed in termination,

and insulin secretion is increased [8].

Despite the critical need to detect splice variants in the context of diabetes, next

generation sequencing (NGS) technologies remain insufficient for this task. Identifying

isoform-specific gene expression requires sequencing reads that span multiple exons of the

mRNA transcript. In the human genome, transcript lengths are estimated to average

between 1,800 and 4,900 bp, with the mode of the distribution around 2,000 bp [9]. NGS

technologies have read lengths of 150 base pairs, making it difficult to identify isoforms. In

contrast, long-read sequencing technologies, such as PacBio and Oxford Nanopore

Technologies, offer the generation of full-length reads that can capture the full RNA

molecule, thereby providing a clearer picture of isoform diversity. Published single-cell long-

read RNA sequencing (sclrRNA-seq) libraries often report shorter read lengths than

expected, which may limit transcript coverage and isoform detection. For instance, a recent

study reported a median read length of 900 bp for sclrRNA-seq of two cancer cell lines [10].

Advances in sequencing technologies, especially single-cell approaches, have

revealed the complex heterogeneity within the pancreas, uncovering distinct functional and

transcriptomic subpopulations across different cell types. In pancreatic islets, single-cell

genomics and patch-seq have identified transcriptionally and functionally distinct beta-cell

subpopulations directly linking gene expression to key physiological processes such as

vesicle exocytosis [11]. This heterogeneity underscores the importance of characterizing

splicing events and their resulting isoforms at the single-cell level. However, sclrRNA-seq

technologies come with inherent limitations. Nanopore flow cells produce fewer reads than

Illumina, with around 20,000 reads per cell for a 5,000-cell experiment, well below the

typical 30,000–50,000 reads per cell common in NGS. Moreover, Nanopore's higher error

rate (1%) increases the likelihood of incorrect barcode and UMI assignments. To overcome

these challenges, we have optimized a protocol for pancreatic islets that improves read

length, advancing the utility of long-read sequencing for single-cell transcriptomics.

Research Design and Methods

Dissociation of pancreatic islets

10-week-old female C57BL/6 mice were obtained from Jackson laboratories.

Pancreatic islets were isolated from mice under ketamine/xylazine/acepromazine

anesthesia by collagenase delivery into the pancreas via injection into the bile duct. The

collagenase-inflated pancreas was surgically removed and digested. After isolation, islets

were dissociated using Accutase in a 37°C bead bath for 25-30 minutes. Single-cell

suspension was filtered through a 40 mm filter and quenched in RPMI media + 10% FBS.

Cells were washed again with RPMI +10% FBS and with PBS + 0.1% BSA. Single-cell

suspensions were loaded into a Genomics Chromium targeting 4000 cells per sample.

Dissociation of spleens

Spleens were isolated from mice under ketamine/xylazine/acepromazine

anesthesia. Spleens were dissociated through a 70 μm strainer in cIMDM using a 3 mL

syringe plunger. Cells were washed, centrifuged, and treated with 1 mL Ammonium-

Chloride-Potassium (ACK) lysis buffer for 30 seconds, followed by dilution in cIMDM and a

second spin. After one additional cIMDM wash, cells were resuspended in PBS + 0.1% BSA.

Single-cell suspensions were loaded into a Genomics Chromium targeting 4000 cells per

sample.

scRNA-seq library preparation and insulin depletion

Single-cell libraries were prepared using either the Chromium Next GEM Single Cell

3' Kit v3.1 or the Chromium Next GEM Single Cell 5' Kit v2 following the protocol up to and

including step 2.4, stopping just before fragmentation. For the optimized libraries, the

following modifications were applied to the 5' library preparation: 1 ul 10 mM dNTP solution

(Thermo Scientific FERR0191) was added to the reaction in step 1.1. The extension time was

increased from 45 minutes to 2 hours in step 1.5. 1 ul 10 mM dNTP solution was added to

the reaction in step 2.2 and the extension time was increased from 1 minute to 3 minutes.

Insulin depletion was performed on cDNA from step 2.4 of the 10X Genomics Chromium

library preparation using the DepleteX™ RNA Depletion Panel (Insulin) kit from Jumpcode

Genomics. We followed the PacBio MAS-IsoSeq protocol (December 2022, Version 1.0) with

the following modifications: During RNP Complex Formation (Step A), we used 0.9 ul Cas9

instead of 2.3 ul, and 1.6 ul Insulin Guide RNA instead of 4.0 ul Single Cell Boost Guide RNA.

During Bead Cleanup (Step D), we used 50 ul (1X) AMPure XP Beads instead of 75 ul 1.5X

SMRTbell Cleanup Beads.

Following insulin depletion, long-read libraries were prepared from the cDNA using

Sequencing Kit V14 (Nanopore SQK-LSK114) and the PCR Expansion (Nanopore EXP-

PCA001). For 3' libraries, the *Ligation sequencing V14 — single-cell transcriptomics with 3'*

cDNA prepared using 10X Genomics on PromethION (SQK-LSK114) protocol was used. For

5' libraries, the *Ligation sequencing V14 - Single-cell transcriptomics with 5' cDNA prepared*

using 10X Genomics on PromethION (SQK-LSK114) protocol was used. Short Fragment

Buffer (SFB) was used for library preparation instead of Long Fragment buffer (LFB). Library

Beads (LIB) were used for the flow cell priming mix stead of Library Solution (LIS). Libraries

were sequenced on R10.4.1 flow cells on either a PromethION 2 Solo (P2S) or PromethION

2 Integrated (P2i).

Bulk RNA-seq library preparation

Islets were isolated from two male C57BL/6J mice as described above. RNA was

purified using the Qiagen RNeasy micro kit (Qiagen Cat # 74004). Bulk RNA-seq libraries

were prepared using KAPA mRNA Hyper Prep kit (Roche 8098123702), following the KAPA

mRNA HyperPrep Kit protocol (KR1352 – v7.21). Insulin depletion was applied between step

8 and 9 in the KAPA library prep protocol.

Single-cell RNA-seq preprocessing

Single-cell libraries were processed using the EPI2ME single-cell workflow v1.1.0

from Oxford Nanopore Technologies for the identification of cell and UMI barcodes

(https://github.com/epi2me-labs/wf-single-cell) using the mm10-2020-A mouse reference

provided by 10x Genomics. The kit name was either 3prime or 5prime depending on the 10x

library preparation and we provided the kit version v3 and v1 for the 3' and 5' samples we

prepared respectively. We provided the kit version that was indicated in the publications

associated with each published dataset. All parameters can be found on github

(https://github.com/CUAnschutzBDC/sc-islet-longread-

analysis/blob/61d52b6ef94ff974a394a9335dd65d28b9e6360e/01_wf_single_cell/trial_par

ameters.tsv) The pipeline was altered to bypass the StringTie step to use a consistent GTF

file from the mm10 genome. These jobs were executed using Singularity and Nextflow. Only

the reads that had a sequence quality higher than 7 were input into this pipeline.

We manually created a gene count matrix from the BAM outputs that were produced

by the EPI2ME pipeline, and then utilized the emptyDrops function from the DropletUtils

package [12] to identify cell barcodes. Using these barcodes, we subset the gene and

transcript count matrices from the single-cell workflow and calculated the read lengths of

cells that were identified using the barcodes. Read lengths were calculated from the BAM

files by looking at the length of the query sequences for all primary reads and were subset

further by matching barcodes (CB), whether the read was tagged with a gene (GN), and

whether the read was tagged with a transcript (TR). Finally, we examined the BAM files and

calculated the number of reads that were tagged with a gene and a transcript, before creating

the final output plots.

Detailed analysis of the 5' modified islet dataset

The filtered data for the 5' modified islet dataset was input into Seurat (v4.1.3) [13-

17], using R (v4.2.3). Quality control metrics were calculated using the perCellQCMetrics

function from the Scuttle package (v1.8.4) [18]. We used the DoubletFinder package (v2.0.3)

[19] to remove cells that were identified as doublets, before running PCA. We tested several

resolutions of UMAP before generating final clustering.

Cell types were initially identified using a reference dataset [20] and further refined

based on the expression of known islet cell type markers. Dimensionality reduction and

clustering was first performed using gene-level expression data, followed by a second round

of clustering based on transcript-level expression. To evaluate the consistency between

gene- and transcript-based cell type annotations, we calculated concordance between the

two methods and visualized the results as confusion matrices using the Matrix (v1.5-4) [21]

and pheatmap (v1.0.12) packages. Clusters identified by transcript-level expression were

used for downstream analysis. To assess differential splicing, we conducted pseudobulk

differential transcript usage (DTU) analysis using DTUrtle (v1.0.2) [22] alongside pseudobulk

differential gene expression (DGE) analysis. Pseudobulk count matrices were generated by

aggregating counts per replicate and cell type using Seurat. Differential expression was

tested at the gene level using DESeq2 (v1.38.3), and differential transcript usage was tested

at the transcript level using DTUrtle with sparseDRIMSeq (v0.1.2) for model fitting. Batch

effects across replicates were corrected using ComBat-seq from the sva package (v3.50.0).

Most figures were generated using the dplyr (v1.1.0), ggplot2 (v3.4.1), readr (v2.1.4),

tidyverse (v2.0.0), MetBrewer (v0.2.0), and scAnalysisR (v0.0.0.9000) [23] packages. Further

details of this pipeline can be found at our code repository:

https://github.com/CUAnschutzBDC/maria_long_read_081224/

Docker images can be found here:

RNA-seq preprocessing (STAR, Fastqc, cutadapt, featurecounts):

https://hub.docker.com/r/kwellswrasman/rnaseq_general

scRNA-seq (DTurtle, scAnalysisR, Seurat):

https://hub.docker.com/r/kwellswrasman/sclrseq_methods_r_docker

Bulk RNA-seq analysis

Raw reads were trimmed using cutadapt (v4.8) and python (v3.10.14) [24] and aligned

to the mouse GRCm38 genome using STAR (v2.7.11b) [25]. Gene expression was quantified

with featureCounts (v2.0.6) [26] and differential expression analysis was conducted using

DESeq2 (v1.44.0) [27] with R version 4.4.1. All plots were made using ggplot2 (v3.5.1) in R.

NGS coverage

The 10k Human DTC Melanoma, Chromium GEM-X Single Cell 5' and 10k Human DTC

Melanoma, Chromium GEM-X Single Cell 3' aligned bam files were downloaded from 10x

genomics (https://www.10xgenomics.com/datasets). Coverage plots were generated using

NGS plot (v2.63) [28].

Transcript Coverage

To visualize the transcript coverage for each of our samples, we utilized the coverage

function from the GenomicFeatures (v1.60.0) package. The reads output from our single-cell

post-processing pipeline were aligned to a transcriptome reference generated from the

GRCm38 Ensembl GTF, and subset based on matching barcodes and unique UMIs from our

processed Seurat objects. For each sample, a coverage matrix of counts per transcript

position was created for every transcript in the GTF file, then each normalized to be a length

of 100. The counts were then normalized by read depth, and then again by applying a softmax

transformation to generate matrix values that summed to 1 across each transcript, before a

column mean provided a value per each 1-100 position per sample. These coverages were

subset based on the number of exons per transcript, supplied by the GTF file, as well as the

length of the transcripts, with a prerequisite that the transcript have >1 exon.

Gviz plots

For visualizing our transcriptome-aligned reads on the genome, we utilized Gviz

(v1.52.0) and the bam files output from our single-cell post-processing pipeline. Briefly,

reads for each sample were aligned to a transcriptome reference generated using the

GRCm38 Ensembl GTF file, and subsequently subset based on matching barcodes and

unique UMIs from our processed Seurat objects. The final output bam files were generated

by subsetting the genome-aligned bam files to only those reads that uniquely mapped to one

transcript in our transcriptome alignment. These bams were then merged by replicate. For

each of these merged bam files, an alignments track was created utilizing Gviz and displayed

alongside the GRCm38 transcripts.

Overrepresentation analysis (ORA)

Differential transcript usage (DTU) genes were subjected to KEGG pathway

overrepresentation analysis (ORA) using gprofiler2 (v0.2.1) [29] with a custom background of

all expressed genes. KEGG pathways were extracted from the enrichment results and ranked

by p-value. The top 25 pathways were visualized as bubble plots.

3' versus 5' bias plot

Isoform-defining splice junctions were identified across datasets. Transcripts were

classified based on whether their unique junctions fell in the 3' region (last 25% of the

transcript), 5' region (first 25% of the transcript), central (middle 50% of the transcript) or

both 5' and 3', and binned accordingly

Data availability

All data will be made available on GEO upon publication.

Code Availability

All source data supporting the findings of this study are provided. Custom analysis pipelines

are available on GitHub at:

https://github.com/CUAnschutzBDC/sc-islet-longread-analysis.

Results

Evaluation of read lengths and isoform detection in published single-cell long read datasets

We aimed to identify isoform differences between islet cell types and subtypes using

single-cell RNA-sequencing. To evaluate the ability of single-cell sequencing technologies

to generate full-length reads, we reanalyzed previously published sclrRNA-seq datasets

generated using 10x Genomics and Oxford Nanopore Technologies, focusing on their ability

to capture full-length transcripts and detect isoform-specific transcript expression. Our

analysis included seven sclrRNA-seq libraries from five different studies [10, 30-33]. The

reanalysis revealed an average read length of 794 bp, and an average mode of 582 bp,

compared to the expected mode distribution of ~2,000 bp in the human genome [9] (Figure

1A). This discrepancy between the average read length and the expected transcript length

underscores the ongoing challenge of capturing full-length transcripts. This shortfall in read

length is important because it limits the transcript detection ability. Where gene detection

ranges from 60-75% of total reads, transcript detection ranges from 30-60% of total reads

(Figure 1B). These findings highlight the limitations of current sclrRNA-seq technologies in

achieving comprehensive transcript-level resolution.

<u>Efficient and specific depletion of insulin from islet sequencing libraries generates</u>

<u>enhanced read diversity</u>

Analyzing transcript expression requires a higher overall read depth than gene

expression analysis, as each gene is associated with multiple transcripts. Initial analysis of

our sclrRNA-seq libraries of mouse pancreatic islets led to the discovery that the two mouse

insulin genes, *Ins1* and *Ins2* made up 25% of the total reads, impeding our ability to achieve

optimal read depth (Figure 1C). To overcome this issue, we incorporated an insulin depletion

step into the protocol and validated the specificity and efficiency of the depletion in a bulk

short-read RNA-sequencing library of mouse pancreatic islets. The depletion was

remarkably efficient and highly specific: insulin transcripts were uniquely depleted, while all

other genes remained unaffected (Figure 1D and Supplementary Table 1). The same insulin

depletion was then applied to a single-cell pancreatic islet library followed by long-read

Nanopore sequencing. Importantly, the insulin depletion was as efficient as in the bulk

sample (Figure 1C). This strategy was applied to all subsequent pancreatic islet libraries

generated for this study. Notably, the classification of beta cells does not rely on the

presence of insulin transcripts. We demonstrate this by computationally removing insulin

reads from a scRNA-seq library and repeating cell clustering (Supplementary Figure 1A-D).

<u>Protocol modifications enhance read length and transcript identification in islet single-cell</u>

<u>long read libraries</u>

289   Most high-throughput sclrRNA-seq methods rely on 10x Genomics single-cell

capture and library preparation, which was originally optimized to generate and amplify

shorter sequences, raising the question of whether it can effectively amplify full-length

transcripts. 10x Genomics offers two types of transcriptomic profiling for single-cell RNA-

seq: one that captures the 3' end of transcripts and another that captures the 5' end. Studies

have shown that 3' RNA libraries frequently contain internal priming artifacts [34] that would

prevent the amplification of full length reads. To test for internal priming in 3' vs 5' libraries,

we downloaded libraries generated using each technology in human melanoma samples

from datasets created by 10x Genomics and analyzed the genomic coverage [28]. 3' libraries

exhibited a notably higher degree of internal priming compared to 5' libraries, as evidenced

by an increased number of reads mapping to the central regions of transcripts in genomic

coverage plots (Figure 1E). Because reads generated through internal priming cannot span

the full length of a transcript, this phenomenon likely contributes to the shorter read lengths

observed in these libraries. To test this, sclrRNA-seq libraries were prepared from mouse

pancreatic islets in parallel using both 3' and 5' capture technologies (n = 2 independent

biological replicates per capture method). The libraries prepared with 5' technology resulted

in a subtle increase in longer reads, compared to the 3' libraries (one-sided Wilcoxon test, p

$< 2 \times 10^{-16}$) (Figure 1F and Supplementary Figure 2B). Furthermore, the 5' library provided

substantially improved transcript identification, increasing from 46.4 ± 0.1% to 59.7 ± 1.3%

(Figure 1B).

To further improve the read length, several additional optimization steps were

introduced into the islet 5' library preparation protocol (Chromium Next GEM Single Cell 5'

Reagent Kits v2), including increasing the extension time from 45 minutes to 2 hours during

GEM-RT Incubation and from 1 to 3 minutes during cDNA amplification, based on the

approach outlined by Lebrigand et al. [30] and increasing the amount of dNTPs. These

modifications resulted in longer reads than those from the 5' library without modifications

(one-sided Wilcoxon test, $p < 2 \times 10^{-16}$) (Figure 1F and Supplementary Figure 2B) and enabled

far better transcript identification than any of the published datasets (62.0 ± 1.2%; Figure

1B). Additionally, the 5' modified libraries showed improved transcript coverage (Figure 1G

and Supplementary Figure 2C), with the most pronounced improvement observed in longer

transcripts (Figure 1H and Supplementary Figure 2D). Overall, this emphasizes the

preference for 5' capture over 3' capture and highlights the necessity for library prep

optimizations to enhance the amplification of full-length reads.

Isolating high-quality RNA from pancreatic islets is notoriously difficult, primarily due

to the presence of digestive enzymes, including RNases that are secreted by the exocrine

pancreas. To explore whether a different cell type might yield still longer reads, we applied

3', 5', and 5' optimized library preparations, as described above, to lymphocytes isolated

from dissociated mouse spleens (n = 2 independent biological replicates per capture

method). The 5' lymphocyte samples demonstrated better transcript identification

compared to the 3' pancreatic islet samples (Supplementary Figure 2A). However, the 10x

Genomics Chromium library prep modifications for the 5' samples did not yield the same

improvements in the lymphocyte samples as observed with the pancreatic islet samples

(Supplementary Figure 2A, E-F). This suggests that the benefits of these optimizations might

be tissue-specific and highlights the need for further refinements tailored to different tissue

types.

<u>Isoform variants identified between alpha and beta cells and within beta cell</u>

<u>subpopulations</u>

With the improved library preparation, the optimized 5' sclrRNA-seq datasets from

mouse pancreatic islets were used to explore whether splicing changes could be detected

from different cell types and cell states. Importantly, the 5' modified sclrRNA-seq dataset,

merged across replicates, allowed clear identification of all expected cell populations

(Figure 2A-B and Supplementary Table 2). Importantly, despite depletion ~95% of insulin

transcripts, enough transcripts remain to identify insulin-expressing cells (Figure 2B).

Furthermore, the analysis revealed that cell type identification remains robust whether using

gene-level or transcript-level expression data for dimensionality reduction and clustering,

with over 90% concordance between the two approaches (Figure 2D-G). This stability in

broad cell type classification aligns with the understanding that major cell types are defined

by distinct gene expression patterns. However, when examining substructure within these

cell types, substantial differences emerged between gene-level and transcript-level

analyses, with consistency ranging from 43% to 90% across subclusters (Figure 2I,

Supplementary Figure 3). These findings suggest that while gene-level expression is

sufficient for identifying major cell types, transcript-level analysis provides crucial insights

into subtle variations within cell populations. Such variations may reflect different cell

states, functions, or responses that are not captured by gene-level analysis alone.

The primary strength of sclrRNA-seq lies in its ability to capture cell-specific isoform

expression. To assess differential splicing, we performed pseudobulk differential transcript

usage (DTU) analysis on merged 5' modified replicates, alongside pseudobulk differential

gene expression (DGE) analysis. A pseudobulk approach was chosen because it treats the

biological sample, rather than individual cells, as the unit of replication, reducing pseudo-

replication and yielding more robust and reproducible results [35, 36]. DTU analysis [22]

identifies proportional differences in the transcript composition of a gene, comparing how

much each transcript contributes to the total gene expression across conditions. Using this

analysis, 218 DTU events were identified between alpha and beta cells (Figure 3A). We also

detected 2,095 DGE genes, of which 56 overlapped between the DTU and DGE analyses

(Supplementary Table 3). As a representative example of beta cell subpopulation analysis,

we focused on the comparison between the 1_beta and 4_beta subpopulations, where we

identified 316 DGE genes and 9 DTU genes, with 2 genes overlapping between the two

analyses (Supplementary Table 4). Overrepresentation analysis (ORA) of DTU genes revealed

enrichment in pathways such as synaptic vesicle cycle, calcium reabsorption, and oxidative

phosphorylation in both alpha versus beta cells and 1_beta versus 4_beta comparisons

(Figure 3B and Supplementary Figure 4A). Specifically, when comparing alpha and beta

cells, we identified isoform-specific differences in *Gnas*, a key regulator of cAMP signaling

and insulin secretion (Figure 3C and Supplementary Figure 4B). Additional DTU genes of

interest in this comparison include *Pcsk2*, which encodes a prohormone convertase

required for insulin processing, as well as *Prdx2*, and *Calm2*, which contribute to antioxidant

defense and calcium signaling, respectively. Analysis of two beta cell subpopulations

(1_beta and 4_beta) revealed distinct isoform usage of *Ndufs2***,** a central component of

oxidative phosphorylation and ATP production (Figure 3D and Supplementary Figure 4C).

This comparison also highlighted isoform differences in *Stxbp1***,** a gene important for insulin

granule exocytosis. To assess whether 3' or 5' library preparation introduces bias in detecting

transcript isoforms, such as preferential capture of variation near the transcript ends, we

examined transcript identification based on splice junctions (Supplementary Figure 4D).

This analysis showed that there was limited 3' or 5' bias, with differences largely driven by a

few highly expressed transcripts. Representative gene plots illustrate that 5' libraries

generally provide reads spanning more exon junctions (Supplementary Figure 5), whereas 3'

libraries better detect certain long transcripts with substantial 3' end variation

(Supplementary Figure 6) and may disproportionately enhance detection of some

transcripts due to internal priming (Supplementary Figure 7).

<u>Discussion</u>

This study demonstrates how an improved sclrRNA-seq library preparation protocol

from isolated islets produces longer reads and increases the proportion of reads that can be

confidently assigned to specific transcripts, improving the utility of long-read sequencing

data for identifying splice variants and cellular heterogeneity in pancreatic endocrine cell

populations. Specifically, this study demonstrates that islet sclrRNA-seq libraries prepared

with 5' protocols outperform those prepared with 3' protocols. While neither library type

showed a strong overall 3' or 5' bias, 3' libraries are prone to internal priming, whereas 5'

reads tend to cover more exon junctions, providing more complete isoform identification.

Enhancements to the 5' library preparation further improves read length and transcript

tagging efficiency in pancreatic islets. Furthermore, depleting insulin transcripts from the

pancreatic islet libraries proved to be a highly effective strategy for maximizing informative

reads, demonstrating the broader potential of targeted transcript depletion in single-cell

RNA-sequencing experiments.

While the modified 5' protocol significantly improved read length in islet samples,

lymphocyte samples showed significant improvement only with the unmodified 5' protocol,

with no additional benefit from the modifications. This indicates that individual cell types will

require unique modifications and further optimizations. Despite the significant

improvements in read length achieved with the modified protocol, it did not meet

expectations for full-length transcript coverage. Achieving this goal will require further

modifications to the 10x chemistry, including adjustments to the master mix and reverse

transcriptase.

Although full-length coverage was not achieved for all transcripts, we successfully

analyzed transcript expression and identified differential transcript usage across cell types

and cell subpopulations. These advancements are critical for uncovering the full complexity

of transcriptomes and hold immense potential for broad application across tissues,

enabling deeper insights into cellular heterogeneity, isoform regulation, and functional

diversity. While scRNA-seq allows the identification of distinct cell populations and

subpopulations, the precise number and composition of clusters can be influenced by

analytical choices such as clustering resolution, preprocessing, and dimensionality

reduction, supporting previous analyses showing that subpopulations may not be

consistently defined across studies [37]. In addition, transcript capture and read depth can

influence subpopulation detection: while adequate coverage suffices for gene-level

conclusions, even greater depth is needed to resolve transcript-level differences, as each

gene is often associated with multiple transcripts. Despite these limitations, single-cell

transcriptomic analyses still provide valuable insights into isoform-level regulation within

specific cell types. Investigating these variations at the single-cell level allows us to uncover

the intricate heterogeneity within tissues, offering a deeper understanding of the functional

and transcriptional diversity that would otherwise go unnoticed. Understanding splicing

dysregulation in pancreatic islets is particularly important, as it may reveal how alternative

splicing shapes beta cell function, immune tolerance, and beta cell susceptibility in

diabetes.

Acknowledgments

We thank Laura White, PhD, and Jay Hesselberth, PhD, for their guidance and support with

Nanopore long-read sequencing technologies. We thank Mia Smith, PhD, for guidance on

spleen sample preparation. We also thank Scott Beard, BDC Cytometer Core Manager, for

islet and spleen isolations.

Funding. This work was supported by grants from the National Institutes of Health

(P30DK116073 [Lori Sussel] , R01 DK082590 [Lori Sussel], and U01 DK127505 [Lori Sussel]).

Duality of interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. M.S.H. was responsible for data acquisition and prepared the

original manuscript. K.L.W. and L.S. reviewed and edited the manuscript. C.J.H. and K.L.W.

developed the computational pipelines. M.S.H., C.J.H., and K.L.W. contributed to data

analysis and the graphical presentation of results. All authors contributed to the study's

methodology and conceptualization. K.L.W. is the guarantor of this work and had full access

to all data in the study and takes responsibility for the integrity of the data and the accuracy

of the data analysis.

References

1. Gonzalez-Porta, M., et al., *Transcriptome analysis of human tissues and cell lines*
reveals one dominant transcript per gene. Genome Biol, 2013. **14**(7): p. R70.
2. Black, D.L., *Mechanisms of alternative pre-messenger RNA splicing*. Annu Rev
Biochem, 2003. **72**: p. 291-336.
3. Piazzi, M., et al., *Alternative Splicing, RNA Editing, and the Current Limits of Next*
Generation Sequencing. Genes (Basel), 2023. **14**(7).
4. Juan-Mateu, J., O. Villate, and D.L. Eizirik, *MECHANISMS IN ENDOCRINOLOGY:*
Alternative splicing: the new frontier in diabetes research. Eur J Endocrinol, 2016.
174(5): p. R225-38.
5. Jeffery, N., et al., *Cellular stressors may alter islet hormone cell proportions by*
moderation of alternative splicing patterns. Hum Mol Genet, 2019. **28**(16): p. 2763-
2774.
6. Diez, J., et al., *Differential splicing of the IA-2 mRNA in pancreas and lymphoid*
organs as a permissive genetic mechanism for autoimmunity against the IA-2 type 1
diabetes autoantigen. Diabetes, 2001. **50**(4): p. 895-900.
7. Dogra, R.S., et al., *Alternative splicing of G6PC2, the gene coding for the islet-*
specific glucose-6-phosphatase catalytic subunit-related protein (IGRP), results in
differential expression in human thymus and spleen compared with pancreas.
Diabetologia, 2006. **49**(5): p. 953-7.
8. Daraio, T., et al., *SNAP-25b-deficiency increases insulin secretion and changes*
spatiotemporal profile of Ca(2+)oscillations in beta cell networks. Sci Rep, 2017.
7(1): p. 7744.
9. Lopes, I., et al., *Gene Size Matters: An Analysis of Gene Length in the Human*
Genome. Front Genet, 2021. **12**: p. 559998.
10. Shiau, C.K., et al., *High throughput single cell long-read sequencing analyses of*
same-cell genotypes and phenotypes in human tumors. Nat Commun, 2023. **14**(1):
p. 4124.
11. Camunas-Soler, J., et al., *Patch-Seq Links Single-Cell Transcriptomes to Human*
Islet Dysfunction in Diabetes. Cell Metab, 2020. **31**(5): p. 1017-1031 e4.
12. Lun, A.T.L., et al., *EmptyDrops: distinguishing cells from empty droplets in droplet-*
based single-cell RNA sequencing data. Genome Biol, 2019. **20**(1): p. 63.
13. Hao, Y., et al., *Dictionary learning for integrative, multimodal and scalable single-cell*
analysis. Nat Biotechnol, 2024. **42**(2): p. 293-304.
14. Hao, Y., et al., *Integrated analysis of multimodal single-cell data*. Cell, 2021.
184(13): p. 3573-3587 e29.
15. Stuart, T., et al., *Comprehensive Integration of Single-Cell Data*. Cell, 2019. **177**(7):
p. 1888-1902 e21.
16. Butler, A., et al., *Integrating single-cell transcriptomic data across different*
conditions, technologies, and species. Nat Biotechnol, 2018. **36**(5): p. 411-420.
17. Satija, R., et al., *Spatial reconstruction of single-cell gene expression data*. Nat
Biotechnol, 2015. **33**(5): p. 495-502.

18. McCarthy, D.J., et al., *Scater: pre-processing, quality control, normalization and visualization of single-cell RNA-seq data in R.* Bioinformatics, 2017. **33**(8): p. 1179-1186.

19. McGinnis, C.S., L.M. Murrow, and Z.J. Gartner, *DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors.* Cell Syst, 2019. **8**(4): p. 329-337 e4.

20. Baron, M., et al., *A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas Reveals Inter- and Intra-cell Population Structure.* Cell Syst, 2016. **3**(4): p. 346-360 e4.

21. Jagan, D.B.a.M.M.a.M. *Matrix: Sparse and Dense Matrix Classes and Methods.* 2024; Available from: https://CRAN.R-project.org/package=Matrix.

22. Tekath, T. and M. Dugas, *Differential transcript usage analysis of bulk and single-cell RNA-seq data with DTUrtle.* Bioinformatics, 2021. **37**(21): p. 3781-3787.

23. Wells, K.L., et al., *Combined transient ablation and single-cell RNA-sequencing reveals the development of medullary thymic epithelial cells.* Elife, 2020. **9**.

24. Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.* EMBnet. journal, 2011. **17**(1): p. 10-12.

25. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-21.

26. Liao, Y., G.K. Smyth, and W. Shi, *featureCounts: an efficient general purpose program for assigning sequence reads to genomic features.* Bioinformatics, 2014. **30**(7): p. 923-30.

27. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

28. Shen, L., et al., *ngs.plot: Quick mining and visualization of next-generation sequencing data by integrating genomic databases.* BMC Genomics, 2014. **15**: p. 284.

29. Raudvere, U., et al., *g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update).* Nucleic Acids Res, 2019. **47**(W1): p. W191-W198.

30. Lebrigand, K., et al., *High throughput error corrected Nanopore single cell transcriptome sequencing.* Nat Commun, 2020. **11**(1): p. 4025.

31. Tian, L., et al., *Comprehensive characterization of single-cell full-length isoforms in human and mouse with long-read sequencing.* Genome Biol, 2021. **22**(1): p. 310.

32. Wang, Q., et al., *Single cell transcriptome sequencing on the Nanopore platform with ScNapBar.* RNA, 2021. **27**(7): p. 763-70.

33. You, Y., et al., *Identification of cell barcodes from long-read single-cell RNA-seq with BLAZE.* Genome Biol, 2023. **24**(1): p. 66.

34. Svoboda, M., H.R. Frost, and G. Bosco, *Internal oligo(dT) priming introduces systematic bias in bulk and single-cell RNA sequencing count data.* NAR Genom Bioinform, 2022. **4**(2): p. lqac035.

35. Zimmerman, K.D., M.A. Espeland, and C.D. Langefeld, *A practical solution to pseudoreplication bias in single-cell studies.* Nat Commun, 2021. **12**(1): p. 738.

36.   Squair, J.W., et al., *Confronting false discoveries in single-cell differential*
expression. Nat Commun, 2021. **12**(1): p. 5692.
37.   Mawla, A.M. and M.O. Huising, *Navigating the Depths and Avoiding the Shallows of*
Pancreatic Islet Cell Transcriptomes. Diabetes, 2019. **68**(7): p. 1380-1393.

535
536
537

<u>Figure legends</u>

<u>Figure 1</u>

Read length and transcript identification comparison between single-cell long-read RNA-

sequencing (sclrRNA-seq) libraries. (A) Read length distribution of published sclrRNA-seq

libraries prepared with 10x Genomics and Nanopore technology. Biological replicates are

included for datasets from Lebrigand et al., 2020 and Wang et al., 2021; other datasets are

shown as single samples. (B) Proportion of reads across datasets where the gene is

identified, the transcript is identified, or neither is identified. Shown are published

reanalyzed datasets and six mouse pancreatic islet samples: two replicates prepared with

3' 10x Genomics technology, two replicates with 5' 10x Genomics technology, and two

replicates with 5' modified 10x Genomics technology (incorporating library preparation

optimizations). (C) Proportion of reads aligned to *Ins1* or *Ins2* in a single-cell RNA-seq

analysis of mouse pancreatic islets pre- and post-insulin depletion. (D) Volcano plot

depicting differential gene expression between non-depleted and insulin-depleted bulk

RNA-seq libraries from mouse pancreatic islets. A total of 23,431 genes is shown in the plot.

(E) NGS coverage plot indicating read start sites across the genomic region. Libraries shown

are 3' and 5' single-cell 10x Genomics preparations derived from human DTC melanoma

cells. (F) Read length distribution comparison across mouse pancreatic islet sclrRNA-seq

libraries, including two replicates prepared with 3' 10x Genomics technology, two replicates

with 5' 10x Genomics technology, and two replicates with 5' modified 10x Genomics

technology. (G) Coverage plot showing Softmax-normalized read coverage across relative

transcript positions for all multi-exon transcripts in mouse pancreatic islet scRNA-seq

libraries prepared with 3', 5', or 5' modified 10x Genomics technology. (H) Coverage plot

showing Softmax-normalized read coverage across relative transcript positions for all

transcripts with ≥9 exons in mouse pancreatic islet scRNA-seq libraries prepared with 3', 5',

or 5' modified 10x Genomics technology.

<u>Figure 2</u>

Comparison of single-cell clustering based on gene expression versus transcript-level

expression. (A) UMAP projection of single cells from insulin-depleted mouse pancreatic islet

sclrRNA-seq libraries prepared with 5' modified 10x Genomics technology (two biological

replicates merged), based on gene-level expression. Cells are colored by gene expression

profiles reflecting major pancreatic cell types. (B) Heatmap showing expression of cell type-

specific markers across single cells insulin-depleted 5' modified libraries (two biological

replicates merged), based on gene-level clustering. (C) UMAP projection of single cells

based on gene-level expression, colored by gene expression profiles reflecting cell

subpopulations. (D) UMAP projection of single cells based on transcript-level (isoform)

expression. Colored by transcript expression profiles reflecting major pancreatic cell types.

(E) UMAP projection of single cells based on transcript-level expression, colored by

transcript expression profiles reflecting cell subpopulations. (F) Confusion matrix showing

concordance in cell type identification between gene-based and transcript-based

clustering. (G) Bar plot quantifying cell type concordance between clustering methods. (H)

Confusion matrix showing low concordance in beta cell subpopulation identification

between clustering methods. (I) Bar plot quantifying beta cell subpopulation concordance

between clustering methods. C1–C5 indicate comparison 1 through comparison 5.

<u>Figure 3</u>

Differential transcript usage between cell types and cell subpopulations. (A) Heatmap

showing differential transcript usage (DTU) between alpha and beta cells. Each row

represents a transcript (significant at FDR < 0.05), and each column represents a cell type.

Colors indicate the fraction of a gene's expression contributed by each transcript within each

cell type. DTU genes were identified from pseudobulk analysis of two biological replicates of

mouse pancreatic islet sclrRNA-seq libraries prepared with 5' modified 10x Genomics

technology. (B) KEGG pathway overrepresentation analysis (ORA) for DTU genes between

alpha and beta cells. The x-axis shows recall, i.e., the proportion of functionally annotated

genes in the query that are associated with each pathway. Bubble size represents the

number of DTU genes in the pathway, and bubble color indicates statistical significance (-

log10 p-value). The top 25 enriched KEGG pathways are shown. (C) Differential gene

expression (DGE), differential transcript expression (DTE), and differential transcript usage

(DTU) analysis of *Gnas* between alpha and beta cells. DTU shows the relative contribution of

each transcript to the gene's overall expression. (C) DGE, DTE, DTU analysis of *Ndufs2*

between beta cell subpopulations 1_beta and 4_beta.

Cell type identification in mouse pancreatic islets with and without insulin transcripts. (A) Heatmap showing expression of cell type–specific markers across clusters defined by gene expression in a dataset including insulin compared to the same dataset from which insulin transcripts have been computationally removed. (B) Confusion matrix comparing cell type assignments between datasets with and without insulin transcripts. (C) UMAP projection of single cells from dataset with insulin, colored by cell type. (D) UMAP projection of single cells from dataset without insulin, colored by cell type.

(A) Proportion of reads across datasets where the gene is identified, the transcript is identified, or neither is identified. Shown are six mouse pancreatic islet samples and six mouse spleen samples generated in this study, alongside published reanalyzed datasets. Each tissue from this study includes two biological replicates prepared with 3′ 10x Genomics technology, two with 5′ 10x Genomics technology, and two with 5′ modified 10x Genomics technology. (B) Read length distribution of mouse pancreatic islet scRNA-seq libraries, separated by replicate. Includes two replicates prepared with 3′ 10x Genomics technology, two replicates with 5′ 10x Genomics technology, and two replicates with 5′ 10x Genomics technology with library preparation optimizations. (C–F) Coverage plots showing Softmax-normalized read coverage across relative transcript positions in mouse pancreatic islet (C, D) and spleen (E, F) scRNA-seq libraries prepared with 3′, 5′, or 5′ modified 10x Genomics technology. Samples separated by replicate. Panels C and E include all multi-exon transcripts, while panels D and F include only transcripts with ≥9 exons.

(A) Confusion matrix showing concordance in alpha cell subpopulation identification between gene-based and transcript-based clustering. (B) Bar plot showing concordance of alpha cell subpopulations across clustering methods. Concordance is quantified for two comparisons: C1 (comparison 1) and C2 (comparison 2). (C) Confusion matrix showing low concordance in delta cell subpopulation identification between clustering methods. (D) Bar plot quantifying delta cell subpopulation concordance between clustering methods quantified for comparisons C1-C6.

(A) KEGG pathway enrichment for DTU genes between beta cell subpopulations 1_beta and 4_beta. The x-axis shows the proportion of functionally annotated genes in the query that are associated with each pathway. Bubble size represents the number of DTU genes in the pathway, and bubble color indicates statistical significance (-log10 p-value). The top 25 enriched KEGG pathways are shown. (B) Differential transcript usage (DTU) analysis of Gnas between alpha and beta cells, separated by biological replicate. (C) DTU analysis of Ndufs2 between beta cell subpopulations 1_beta and 4_beta, separated by biological replicate. (D) Bar plot showing 3′ versus 5′ bias across libraries prepared with different technologies. Transcripts are classified by whether the junctions that uniquely identify each isoform fall in the 3′ region or 5′ region of the transcript, or both.

G6pc2 gene and transcript expression across mouse pancreatic islet libraries. (A) Gene-level expression of G6pc2 in 3′ modified, 5′ standard, and 5′ modified libraries (two biological replicates merged for each library). (B) Transcript-level expression of ENSMUST00000112317, isoform of G6pc2, in the same libraries. (C) Transcript-level expression of ENSMUST00000005364, isoform of G6pc2, in the same libraries. (D) Gviz plot showing a gene model track of select transcripts of G6pc2, and an alignments track including all reads uniquely aligned to any transcript of G6pc2.

Gng12 gene and transcript expression across mouse pancreatic islet libraries. (A) Gene-level expression of Gng12 in 3′ modified, 5′ standard, and 5′ modified libraries (two biological replicates merged for each library). (B) Transcript-level expression of ENSMUST00000043148, isoform of Gng12, in the same libraries. (C) Gviz plot showing a gene model track of select transcripts of Gng12, and an alignments track including all reads uniquely aligned to any transcript of Gng12. Arrow indicates a long exon at the 3′ end of the transcript.

Nktr gene and transcript expression across mouse pancreatic islet libraries. (A) Gene-level expression of Nktr in 3′ modified, 5′ standard, and 5′ modified libraries (two biological replicates merged for each library). Transcript-level expression of ENSMUST00000182904, isoform of Nktr, in the same libraries. (C) Gviz plot showing a gene model track of select transcripts of Nktr, and an alignments track including all reads uniquely aligned to any transcript of Nktr. Arrow indicates retained intron captured by internal priming.

Read length and transcript identification comparison between single-cell long-read RNA-sequencing (sclrRNA-seq) libraries. (A) Read length distribution of published sclrRNA-seq libraries prepared with 10x Genomics and Nanopore technology. Biological replicates are merged for datasets from Lebrigand et al., 2020 and Wang et al., 2021; other datasets are shown as single samples. (B) Proportion of reads across datasets where the gene is identified, the transcript is identified, or neither is identified. Shown are published reanalyzed datasets and six mouse pancreatic islet samples: two replicates prepared with 3′ 10x Genomics technology, two replicates with 5′ 10x Genomics technology, and two replicates with 5′ 10x Genomics technology incorporating library preparation optimizations. (C) Proportion of reads aligned to Ins1 or Ins2 in a single-cell RNA-seq analysis of mouse pancreatic islets pre- and post-insulin depletion. (D) Volcano plot depicting differential gene expression between non-depleted and insulin-depleted bulk RNA-seq libraries from mouse pancreatic islets. A total of 23,431 genes is shown in the plot. (E) NGS coverage plot indicating read start sites across the genomic region. Libraries shown are 3′ and 5′ single-cell 10x Genomics preparations from human DTC melanoma cells. (F) Read length distribution comparison across mouse pancreatic islet scRNA-seq libraries, including two replicates prepared with 3′ 10x Genomics technology, two replicates with 5′ 10x Genomics technology, and two replicates with 5′ 10x Genomics technology with library preparation optimizations, alongside published datasets from Lebrigand et al., 2020 and Wang et al., 2021. Biological replicates are merged for the 10x libraries as well as for datasets from Lebrigand et al., 2020 and Wang et al., 2021. (G) Coverage plot showing Softmax-normalized read coverage across relative transcript positions for all multi-exon transcripts in mouse pancreatic islet scRNA-seq libraries prepared with 3′, 5′, or 5′ modified 10x Genomics technology. (H) Coverage plot showing Softmax-normalized read coverage across relative transcript positions for all transcripts with ≥9 exons in mouse pancreatic islet scRNA-seq libraries prepared with 3′, 5′, or 5′ modified 10x Genomics technology.

Comparison of single-cell clustering based on gene expression versus transcript-level expression. (A) UMAP projection of single cells from insulin-depleted mouse pancreatic islet sclrRNA-seq libraries prepared with 5′ modified 10x Genomics technology (two biological replicates merged). Cells are colored by gene expression profiles reflecting major pancreatic cell types. (B) Heatmap showing expression of cell type–specific markers across insulin-depleted 5′ modified libraries (two biological replicates merged), based on gene-level clustering. Cell type abbreviations: EC, endothelial cell; MΦ, macrophage; qSC, quiescent stellate cell. (C) UMAP projection of single cells based on gene-level expression, colored by gene expression profiles reflecting cell subpopulations. (D) UMAP projection of single cells based on transcript-level (isoform) expression. Colored by transcript expression profiles reflecting major pancreatic cell types. (E) UMAP projection of single cells based on transcript-level expression, grouped by transcript expression profiles reflecting cell subpopulations. (F) Confusion matrix showing concordance in cell type identification between gene-based and transcript-based clustering. (G) Bar plot quantifying cell type concordance between clustering methods. (H) Confusion matrix showing low concordance in beta cell subpopulation identification between clustering methods. (I) Bar plot quantifying beta cell subpopulation concordance between clustering methods. C1–C5 indicate comparison 1 through comparison 5.

Differential transcript usage between cell types and cell subpopulations. (A) Heatmap showing differential transcript usage (DTU) between alpha and beta cells. Each row represents a transcript (significant at FDR < 0.05), and each column represents a cell type. Colors indicate the fraction of a gene's expression contributed by each transcript within each cell type. DTU genes were identified from pseudobulk analysis of two biological replicates of mouse pancreatic islet sclrRNA-seq libraries prepared with 5′ modified 10x Genomics technology. (B) KEGG pathway overrepresentation analysis (ORA) for DTU genes between alpha and beta cells. The x-axis shows recall, i.e., the proportion of functionally annotated genes in the query that are associated with each pathway. Bubble size represents the number of DTU genes in the pathway, and bubble color indicates statistical significance (-log10 p-value). The top 25 enriched KEGG pathways are shown. (C) Differential gene expression (DGE), differential transcript expression (DTE), and differential transcript usage (DTU) analysis of Gnas between alpha and beta cells. DTU shows the relative contribution of each transcript to the gene's overall expression. (C) DGE, DTE, and DTU analysis of Ndufs2 between beta cell subpopulations 1_beta and 4_beta.