Rapid hypothalamic-pituitary recovery after chronic glucocorticoid therapy
enables strategies that prevent adrenal suppression in mice

Lindsey S. Gaston[1], Brenna C. Jorgensen[1], Hannah R. Friedman[1], Marc S. Sherman[2], Joseph A.
Majzoub[1]

[1]Division of Endocrinology, Department of Pediatrics, Boston Children's Hospital, Harvard
Medical School, Boston, MA, USA

[2]Division of Gastroenterology, Department of Medicine, Massachusetts General Hospital, Harvard
Medical School, Boston, MA, USA

*Joseph Majzoub, MD

300 Longwood Avenue, Boston, MA 02115; 617-919-2930

Email: Joseph.Majzoub@childrens.harvard.edu

*Lindsey Gaston, MD

300 Longwood, Avenue, Boston, MA 02115; 617-355-7476

Email: Lindsey.Gaston@childrens.harvard.edu

Author Contributions: LG and JM conceived the project and designed experiments. LG, BJ, and
HF conducted experiments, and LG analyzed the data. LG, MS, and BJ acquired microscopic
images. LG wrote the manuscript, which all authors have reviewed, edited, and approved.

Competing Interest Statement: The authors have no competing interests to disclose.

Classification: Biological sciences: medical sciences

Keywords: Glucocorticoids, adrenal suppression, prevention

31   **This PDF file includes:**

32   Main Text

33   Figures 1 to 4

Abstract

Glucocorticoid-induced adrenal insufficiency (GIAI) can persist for months after cessation of chronic corticosteroid therapy, predisposing patients to life-threatening adrenal crises. The mechanisms underlying this prolonged suppression are poorly understood and have been primarily attributed to delayed recovery of hypothalamic-pituitary function (Graber *et al.,* 1965). We examined the timing of hypothalamic, pituitary, and adrenal (HPA) recovery after 8 weeks of dexamethasone (DEX) treatment in mice. During DEX exposure, HPA axis activity was fully suppressed. Within one week of withdrawal, *Crh* mRNA and plasma ACTH surprisingly rebounded above control levels, yet corticosterone (CORT) remained low for an additional seven weeks. DEX-treated adrenals were atrophic and contained large clusters of lipid-associated macrophages. Even after accounting for these macrophages, CORT secretion was disproportionately low relative to the size of the adrenocortical cell mass for at least two weeks after DEX withdrawal despite supraphysiologic ACTH stimulation. Thus, the adrenal is the primary site of post-withdrawal HPA axis dysfunction, involving both cortical atrophy and impaired steroidogenesis. We then assessed whether trophic stimulation of the adrenal using either a pharmacologic or genetic approach prevented GIAI. Co-treatment with DEX and daily cosyntropin, an ACTH analog, failed to prevent GIAI. However, mice with non-suppressible endogenous ACTH due to deletion of *Nr3c1* (the glucocorticoid receptor gene) in hypothalamic neurons maintained normal adrenal function and histology despite long-term DEX. Localizing the mechanisms of post-withdrawal GIAI to the adrenal may permit the development of strategies that prevent GIAI by delivering sustained trophic support to the adrenal during the period of glucocorticoid exposure.

Significance Statement

Glucocorticoids are among the most commonly prescribed anti-inflammatory agents, yet chronic exposure can cause sustained suppression of the hypothalamic-pituitary-adrenal (HPA) axis, predisposing patients to life-threatening adrenal crises. This glucocorticoid-induced adrenal insufficiency (GIAI) has been attributed primarily to delayed hypothalamic-pituitary recovery after treatment withdrawal, limiting the development of preventive strategies. Using a murine model of GIAI, we found that hypothalamic-pituitary function recovers rapidly after cessation of chronic glucocorticoids, whereas adrenal suppression persists for eight weeks. Continuous adrenal stimulation during glucocorticoid exposure, achieved through a targeted genetic approach, preserved adrenal function. These findings identify the adrenal as the principal site of post-withdrawal HPA axis dysfunction and suggest that therapies sustaining adrenal trophic signaling could prevent GIAI in patients requiring long-term glucocorticoid therapy.

Main Text

Introduction

Systemic glucocorticoids are prescribed to up to 3% of the population in developed countries for treatment of autoimmune and inflammatory diseases, malignancies, and to prevent allograft rejection[1,2]. Many additional patients are chronically treated with topical and inhaled formulations that can exert systemic effects even at moderate doses[3-5]. Like endogenous cortisol in humans and corticosterone in rodents, exogenous glucocorticoids engage the glucocorticoid receptor (GR) to suppress hypothalamic-pituitary-adrenal (HPA) axis activity through negative feedback. This feedback reduces the synthesis and secretion of corticotropin-releasing hormone (CRH) from the paraventricular nucleus of the hypothalamus and adrenocorticotropic hormone (ACTH) from the anterior pituitary[6]. In the absence of ACTH, a key trophic factor for the adrenal cortex,

cortisol-producing cells of the zona fasciculata (zF) undergo both suppression of steroidogenesis
and apoptotic cell loss[7,8].

Hypothalamic-pituitary-adrenal (HPA) axis suppression is typically transient in patients
treated with short-term glucocorticoids. However, treatment exceeding four weeks can cause
protracted adrenal insufficiency that can persist for up to a year after steroid withdrawal[9-11]. While
awaiting HPA axis recovery, individuals are at risk for adrenal crisis with otherwise routine
illnesses or trauma, which has been associated with longer hospitalizations, higher rates of ICU
admissions, and even death in patients who are not promptly administered stress-dose
steroids[12,13].

Despite the ubiquity of glucocorticoid use and the morbidity of glucocorticoid-induced adrenal
insufficiency (GIAI), there are currently no available treatments to prevent or hasten recovery
from adrenal suppression. This is due, in part, to the prevailing view that prolonged GIAI is
mediated by delayed, post-withdrawal recovery of the hypothalamus and pituitary, which are
relatively inaccessible therapeutic targets. Post-withdrawal central suppression was first
described by Graber et al., who reported that ACTH levels were inappropriately low to normal for
at least two months in patients withdrawn from chronic glucocorticoids despite release from
negative feedback[14]. Since direct ACTH assays had not yet been developed, systemic ACTH
levels in this study were interpolated from the corticosterone responses of hypophysectomized
rats injected with extracted patient plasma[14,15]. Another study cited inadequate ACTH responses
to CRH stimulation testing in patients withdrawn from long-term prednisone as evidence of post-
withdrawal central suppression. However, testing was conducted 24 hours or two biological half-
lives from the last prednisone dose, thus limiting differentiation between active, glucocorticoid-
mediated negative feedback and genuine, post-withdrawal suppression[16,17]. Post-withdrawal
central suppression has also been described after surgical cure of Cushing's syndrome, a state of
endogenous glucocorticoid excess, though clinical studies have disagreed as to whether this is
mediated by persistent hypothalamic or pituitary dysfunction[18-20].

Interestingly, patients with congenital adrenal hyperplasia, who also require chronic,
suppressive glucocorticoid therapy, exhibit elevated levels of both ACTH and ACTH-dependent
steroid precursors at their morning hydrocortisone trough[21,22]. This suggests that some patients
have rapid normalization of hypothalamic-pituitary function after release from glucocorticoid
feedback inhibition. The mammalian HPA axis is highly conserved evolutionarily, and all rodent
studies have reported rapid restoration of hypothalamic-pituitary function after glucocorticoid
treatment. Basal and stimulated ACTH levels (following lipopolysaccharide, arecoline, and CRH
administration) were recovered within days of discontinuing 5 or 7 days of methylprednisolone or
dexamethasone, respectively, in rats[23,24]. In mice treated with dexamethasone for 30 days,
plasma ACTH and corticotrope *Pomc* mRNA levels were elevated above control levels by 1-week
post-withdrawal[25]. In all studies, corticosterone (CORT) levels and/or steroidogenic pathway gene
expression remained low 5-7 days after discontinuing steroids despite normalization of adrenal
weights[23-25]. Importantly, no animal studies to date have examined HPA axis recovery after longer
courses of glucocorticoids known to cause protracted adrenal suppression in humans.

Given these discrepancies in the existing clinical and preclinical literature, we developed a
murine model of GIAI to define the locus and mechanisms of persistent HPA axis dysfunction
following withdrawal of long-term glucocorticoids. After we discovered that adrenal dysfunction
persists for months after steroid cessation despite rapid hypothalamic-pituitary recovery, we next
tested whether sustained trophic stimulation of the adrenal during the period of glucocorticoid
exposure could preserve steroidogenic function. Together, these studies identify the adrenal as

the principal site of post-withdrawal GIAI and highlight it as an accessible target for future
preventative interventions.

Results

Mice treated with long-term dexamethasone (DEX) have protracted adrenal insufficiency despite
early hypothalamic-pituitary recovery

We first established when hormonal function at each node of the HPA axis recovered following
long-term DEX treatment. Adult, male, WT mice were treated with either continuous DEX or vehicle
(DEX-naïve, or DN) in the drinking water for 8 weeks. Starting at the time of DEX withdrawal (8+0)
and at 1-2 week intervals through 8+8 (where A weeks of treatment plus B weeks of withdrawal =
A+B), separate cohorts of mice underwent serial cosyntropin stimulation and insulin-induced
hypoglycemia (IIH) testing followed by necroscopy (Model 1, Fig. S1).

As expected, long-term DEX treatment suppressed *Crh* mRNA within the paraventricular
nucleus of the hypothalamus (PVH; 28±5 in 8+0 vs. 156±32 arbitrary units [AU] in DEX-naïve (DN)
control animals, p<0.05). However, *Crh* expression recovered to control levels 1 week after DEX
was discontinued (8+1; 159±28 AU vs. DN, p>0.9999, Fig. 1*A*, *B*). Similarly, IIH-stimulated plasma
ACTH concentrations were significantly suppressed during the period of glucocorticoid exposure
(4.2±3.6 pg/mL at 8+0) but rebounded significantly above control levels by 1-week post-withdrawal
(1707±85 pg/mL at 8+1 vs. 875±44 pg/mL). IIH-stimulated ACTH levels remained elevated through
4 weeks post-withdrawal and reached a peak two weeks after completion of DEX treatment (Fig.
1*C*). Despite this, both IIH- and cosyntropin-stimulated CORT levels remained significantly lower
than DN controls until 8 weeks after DEX cessation (Fig. 1*D*, *E*). Thus, mice withdrawn from long-
term DEX have delayed recovery of CORT secretion despite rapid restoration of hypothalamic-
pituitary function.

Long-term DEX induces adrenocortical atrophy with accumulation of large macrophage clusters

Having found the adrenal to be the limiting site of post-withdrawal GIAI, we then assessed
adrenal mass and histology at select timepoints after chronic DEX. At 8+0, adrenal mass was
reduced by 60% compared to controls (1.0±0.041 mg vs. 2.5±0.063 mg in DN, p<0.0001, Fig. 2*A*).
Although adrenals from DEX-exposed adrenals weighed significantly less than those from DN
controls through 6 weeks post-withdrawal, divergent patterns of recovery were noted for adrenal
mass vs. function. In particular, adrenal mass nearly doubled between 0 to 1 weeks post-withdrawal
to 75% of controls (1.9±0.11 mg at 8+1) and remained relatively stable over the remaining recovery
period. In contrast, CORT secretion was 75% lower than control values at 8+1 and more gradually
recovered over the subsequent 7 weeks (Fig. S2*A*, *B*; raw data derived from Figs. 1*D*, *E* and 2*A*).

Hematoxylin and eosin staining revealed that DEX caused striking adrenocortical atrophy
followed by gradual re-accumulation of cortical tissue in the 8 weeks post-withdrawal (Fig. 2*B*).
Interestingly, DEX-exposed adrenals contained large, vacuolated structures with reduced nuclear
density near the corticomedullary junction that persisted through 8 weeks post-withdrawal (Fig. 2*B*,
magnification in 2*E*). These clusters were intensely autofluorescent, which was markedly
attenuated with the addition of TrueBlack, a lipofuscin autofluorescence quencher[26] (Fig. S2*A*, *B*).
ISH with a 5'-Cy5-Oligod(T)30 demonstrated similar, nuclear mRNA poly(A) content in these
structures compared to surrounding cortical cells (Fig. S2*C*)[27,28]. The large, vacuolated,
extranuclear space in these clusters was not strongly labeled with 568 NHS Ester (SE), a total
protein stain (Fig S2*D*)[29,30]. Together, this suggests that the multinucleated clusters observed in
DEX-exposed adrenals contain live, lipofuscin-rich, biologically active cells rather than acellular,

proteinaceous debris. These cells were negative for the steroidogenic enzyme *Cyp11b1* mRNA, a
marker of zona fasciculata (zF) cells.

Immunostaining with the myeloid-specific marker, CD68, and macrophage-specific marker,
F4/80, identified these cells as clusters of adrenal macrophages (AMs). Rare, mononuclear CD68-
and F4/80-positive cells were visualized in the adrenal cortex (labeled with fluorescent 488-
conjugated streptavidin[31]) of DN mice (Fig. 2*C, D*). Increased CD68- and F4/80-positive
macrophages were present in the adrenal cortex after 2 weeks of DEX treatment (2+0). After 8
weeks of DEX (8+0), large AM clusters accumulated at the corticomedullary junction (Fig. 2*C, D*,
magnified in 2*F, G*) and appeared to fragment and migrate into the adrenal medulla by 8 weeks
post-withdrawal (8+8). AMs contained lipid detected by both oil red O and BODIPY staining but
were negative for the lipid droplet associated proteins perilipin-1 and 2 (PLIN 1 and 2; Fig. S4*A-D*).
AMs were also positive for CD11c and *Trem2* by IF and ISH, respectively, both of which are
expressed by lipid-associated macrophages in the context of metabolic associated steatohepatitis
and atherosclerosis (Fig. S4*E, F*)[32-35].

CORT secretion is disproportionately low relative to steroidogenic cell mass for at least 2 weeks
post-DEX

As AMs likely contribute to adrenal mass but not biosynthetic function, we then asked
specifically how the steroidogenic cell population contracted and recovered after chronic DEX
treatment. To do this, we performed morphometric analysis on thick, immunostained adrenal
sections, from which we selected the largest, equatorial one from each gland. Steroidogenic
adrenocortical cells were labeled with 488-streptavidin[31] (green), and the medulla was identified by
immunofluorescence for tyrosine hydroxylase (red). Sections were not treated with TrueBlack so
that AMs could be identified by endogenous autofluorescence (cyan; Fig. 3*A*).

As observed with adrenal weights, pan-cortical area, which includes both streptavidin-positive
and autofluorescent regions, decreased with DEX treatment (764 ± 284 mm^2 in 8+0 vs. 1596 ± 319.4
mm^2 in DN) but nearly normalized by 1-week post-withdrawal (1100 ± 72 mm^2 in 8+1, Fig. 3*A, B*).
Medullary (TH-positive) area was comparable at all time points, indicating that relatively equivalent,
equatorial adrenal sections were used for morphometric analyses ($p=0.15$, Fig. 3*C*).
Autofluorescent AM area increased with DEX treatment and remained stably elevated through 8
weeks post-withdrawal (Fig. 3*A, D*). Steroidogenic cell type area, computed by subtracting AMs
(cyan autofluorescence) from the pan-cortical (green) area, markedly decreased with DEX
treatment (272 ± 47 mm^2 in 8+0 vs. 1286 ± 153 mm^2 in DN, $p<0.001$) but more than tripled in the first
week post-withdrawal (887 ± 70mm^2; Fig. 3*A, E*).

To examine how adrenal function relates to adrenocortical cell mass, we calculated the ratio of
cosyntropin-stimulated CORT secretion to the steroidogenic cell type area, normalized to DN
controls (Fig. 3*F*). This ratio showed a nonsignificant decrease at the end of DEX treatment
(0.60 ± 0.14) but reached its lowest values at 1- and 2-weeks post-DEX (0.58 ± 0.05 and 0.44 ± 0.05,
$p<0.05$ vs. DN). By 8+6, the ratio had returned to baseline (0.99 ± 0.01) and was elevated relative
to DN by 8+8 (2.0 ± 0.18, $p<0.001$ vs DN). These findings indicate that, for at least 2 weeks after
DEX withdrawal, CORT secretion is disproportionately low compared to the size of the non-
macrophage adrenal cortex, even under supraphysiologic ACTH stimulation.

Immunofluorescence for CD31 revealed preserved vascularity in both DEX-treated (8+0) and
withdrawn (8+1) adrenals (Fig. S5*A*). The zona glomerulosa, labeled by Dab2, thickened during
chronic DEX exposure (45 ± 2.6 µm at 8+0 vs. 36 ± 0.96 µm in DN) but returned to baseline by 8+1
(36 ± 2.2 µm; Fig. S5B and C). Ki-67 staining demonstrated a robust increase in adrenocortical
proliferation within one week of DEX withdrawal ($17\pm1.5\%$ of 8+1 cells vs. $2.0 \pm 1.0\%$ of DN cells;
Fig. S5B and D), consistent with the known mitogenic effects of ACTH[36]. Together, these findings
indicate that disproportionately low CORT secretion relative to steroidogenic cell type area cannot

be explained by impaired adrenal vascularity, zonation, or trophic signaling but instead reflects a
selective defect in CORT biosynthesis.

Maintenance of endogenous ACTH secretion but not intermittent cosyntropin administration during
the period of DEX exposures prevents GIAI

Having localized post-withdrawal GIAI to the adrenal, we next tested whether maintaining
trophic stimulation during glucocorticoid exposure could prevent adrenal suppression. Wild-type
mice were treated with DEX as above and received daily, intraperitoneal injections of cosyntropin
(ACTH [1-24]; DEX+COS) for either 4 days (short-term) or 8 weeks (long-term). Long-term cohorts
underwent either a) serial cosyntropin stimulation testing at 1–2-week intervals from 0-8 weeks
after withdrawal, or b) necroscopy at 8+1 (Model 2, Fig. S1). Short-term DEX+COS treatment
significantly but variably reduced DEX-induced adrenocortical apoptosis (5.7±1.7% vs. 11±0.8% in
4D DEX glands; Fig. S6*A*, *B*). In contrast, after 8 weeks of treatment (8+0), cosyntropin-stimulated
CORT levels remained equally suppressed in DEX+COS and DEX-only mice (2.0±1.0 µg/dL vs.
2.0±0.61 µg/dL, p=0.99). Following withdrawal, CORT recovery was significantly delayed in
DEX+COS mice (doubling time=5.2 weeks) compared to DEX-only mice (2.5 weeks; p=0.009, Fig.
S6*C*). Morphometric analysis at 8+1 revealed comparable steroidogenic cell type area between
DEX+COS and DEX-only adrenals (Fig. S6*D*, *G*) but reduced AM area in the DEX+COS group
(Fig. S6*F*). AM morphology was also heterogeneous, ranging from scattered smaller cells
throughout the zona fasciculata to typical aggregates at the corticomedullary junction (Fig. S6*D*).
Thus, intermittent, pharmacologic ACTH stimulation during glucocorticoid exposure modestly
blocks adrenocortical apoptosis and macrophage accumulation but is insufficient to sustain
steroidogenic capacity or accelerate functional recovery.

We then tested whether sustained, endogenous ACTH signaling during the period of
glucocorticoid exposure prevents GIAI (Model 3, Fig. S1). We created Sim1^{Cre}:GR$^{fl/fl}$ mice
(HGRKO), which have reduced hypothalamic glucocorticoid receptor expression and exhibit
constitutively elevated *Crh* mRNA content and plasma ACTH and corticosterone concentrations[37].
Cre-negative littermates (HGRWT) were used as controls. Mice were treated for either 4 days or 8
weeks with DEX. HPA axis activity was assessed pre-treatment and on chronic DEX (8+0) in the
unstressed state, following physical restraint, and after cosyntropin stimulation (Fig. 4). On long-
term DEX treatment (8+0), GR immunoreactivity was present in 94% of PVH neurons of HGRWT
mice vs. 23% in HGRKO mice (Fig. S7*A*, *B*). *Crh* expression in the PVH was suppressed in these
same DEX-treated HGRWT (49±12 AU) but not HGRKO animals (244±2.2 AU; Fig. S7*C, D*). GR
was thus successfully deleted from the PVH, which conferred resistance DEX-induced feedback
inhibition of *Crh* mRNA expression.

Apoptosis was uniformly rare in short-term DEX-treated HGRKO glands (1.7%±0.9 vs. X% DN,
p=0.74; Fig. 4A, *B*). HGRKO mice had higher basal ACTH concentrations than HGRWT controls
(470±76 vs. 104±55 pg/mL, *p*<0.05) that were not suppressed on long-term DEX (385±64 vs. 0±0
pg/mL; *p*<0.01; Fig. 4*C*). Pre-treatment basal and stimulated CORT levels were as expected higher
in HGRKO than HGRWT mice (Fig. 4*D*). On long-term DEX, both basal, restraint- and cosyntropin-
stimulated CORT were markedly suppressed in HGRWT vs. remaining high in HGRKO animals
(basal: 1.5±0.98 vs. 14±1.6 µg/dL; restraint: 1.9±1.6 vs. 33±4.5 µg/dL; cosyntropin-stimulated:
1.1±0.62 vs. 33±3.9 µg/dL; *p*<0.01–0.001). Adrenal mass and morphology were normal in HGRKO
animals, and these glands notably lacked the large clusters of AMs observed in HGRWT controls
(Fig. 4E-I). Sustained, CRH-driven, endogenous ACTH signaling during long-term DEX treatment
thus prevents GIAI and cortical atrophy.

Discussion

Using a murine model of GIAI, we found that hypothalamic-pituitary function rapidly
recovered after cessation of chronic glucocorticoids, whereas adrenal suppression persisted for

up to 8 weeks. DEX-exposed adrenals exhibited marked cortical atrophy and infiltration of large,
lipid-laden macrophages. Even after accounting for these macrophages, CORT production was
disproportionately low relative to the size of the adrenal cortex, suggesting that post-withdrawal
GIAI involves both adrenocortical cell loss and steroidogenic or secretory dysfunction. The
provision of continuous, CRH-driven, endogenous ACTH signaling to the adrenal during long-
term DEX treatment preserved adrenal size, morphology, and function. These results identify the
adrenal as the principal site of ongoing suppression following withdrawal from chronic
glucocorticoids and point to sustained trophic stimulation of the gland as a promising therapeutic
strategy to prevent GIAI.

Consistent with our findings, prior rodent studies have shown that hypothalamic-pituitary
function recovers within one week of discontinuing short- to moderate-term glucocorticoid
treatment (5-30 days)[23-25]. We extended treatment to a duration known to cause protracted GIAI
in humans, making the rapid central recovery observed in our model particularly striking. Notably,
the 1965 study that established the paradigm of post-withdrawal central suppression included
only six patients with exogenous glucocorticoid exposure. Because ACTH could not be directly
measured, circulating levels were inferred from corticosterone responses in hypophysectomized
rats injected with patient plasma compared to three standard ACTH doses or saline[14,15]. Another
frequently cited study assessed HPA axis function just 24 hours (1-2 biological half lives) after the
final glucocorticoid dose[16]. ACTH suppression in this context could therefore reflect direct,
glucocorticoid-mediated feedback inhibition on the hypothalamus and pituitary rather than
delayed central axis recovery. Interestingly, patients with congenital adrenal hyperplasia, who
require chronic, suppressive glucocorticoid therapy, exhibit elevated levels of ACTH and ACTH-
dependent precursors at their morning hydrocortisone troughs, as would be expected with rapid
normalization of hypothalamic-pituitary function[21,22]. Together, the methodological limitations of
earlier clinical studies and our finding of rapid hypothalamic-pituitary recovery in mice underscore
the need to reexamine the paradigm of post-withdrawal central suppression using modern, well-
timed ACTH assays and adequately powered cohorts of glucocorticoid-treated patients.

The brief or absent post-withdrawal central suppression found in our study encouraged us to
investigate mechanisms of GIAI at the adrenal level. Since ACTH is a well-established trophic
factor for the zF[7,8], the time required for cellular regeneration from DEX-induced cortical atrophy
is an intuitive explanation for persistently low CORT secretion after glucocorticoid withdrawal
despite supraphysiologic ACTH stimulation. The adult zona fasciculata is considered a "modestly
proliferative" tissue with a turnover rate of 120 days in mice[38,39]. While this is longer than the 8
weeks required for functional recovery in our study, more rapid renewal would be expected in
states of increased ACTH stimulation, as occurs after DEX withdrawal[40]. Interestingly, we found
that adrenal mass recovered more rapidly than steroidogenic function. This is consistent with
prior studies that demonstrated that CORT levels and steroidogenic pathway gene expression
were low one week after discontinuing short- to moderate-term glucocorticoids despite
normalization of adrenal weights[23-25]. Using quantitative morphometry, we further demonstrated
that this dissociation between adrenal size and function persisted even after accounting for the
DEX-induced expansion of large, lipid-laden adrenal macrophages. Together, these findings
suggest that post-withdrawal GIAI reflects not only adrenocortical cell loss but also a persistent
defect in CORT biosynthesis or secretion, the mechanisms of which should be a focus of future
studies.

While Wylie and Kerr noted accumulation of apoptotic bodies "within phagosomes of
'histiocytes'" after short-term glucocorticoid treatment[7], macrophage accumulation has not been
reported after long-term glucocorticoid therapy. Large AMs in our model share multiple features
with adrenal macrophages described in acute and chronic stress states. They express classical
macrophage markers, such as CD68, F4/80 and Cd11c, and are intensely autofluorescent due to
lipid accumulation[34,41-43]. Male murine adrenal macrophages are also long-lived[42], consistent with
our continued detection of AMs at least 8 weeks after DEX withdrawal. The AMs in the present
study most strikingly differ from typical adrenal macrophages in both their large size and apparent
multinucleation. Morphologically similar adrenal macrophages have been described in aged, male

Znrf3 knockout mice, which undergo early adrenocortical hyperplasia followed by senescence[44]. It
is therefore plausible that adrenal macrophages undergo expansion, fusion, and multinucleation
when the demand for phagocytic clearance is high, as has been reported in adipose, ovary and
bone tissues[45-47]. Interestingly, Xu et al. reported that adrenal macrophages in atherogenic
conditions restrict glucocorticoid production via Trem2-dependent paracrine suppression of
steroidogenic acute regulatory protein (StAR), the rate limiting enzyme in CORT biosynthesis, in
neighboring zona fasciculata cells[34]. Since AMs in our model strongly express *Trem2,* further
work is needed to determine if they contribute to GIAI via a similar paracrine signaling
mechanism.

Most importantly, our finding of rapid hypothalamic-pituitary recovery after glucocorticoid
cessation has enabled us to begin development of new strategies to prevent GIAI targeted to the
adrenal. We demonstrated that sustained, CRH-driven ACTH signaling but not intermittent
cosyntropin treatment during chronic glucocorticoid treatment prevents GIAI. The differential
efficacy of our genetic and pharmacologic strategies may reflect the short half-life of cosyntropin
or a requirement for physiologic ACTH dynamics to maintain adrenal steroidogenic capacity[48-50].
We could not assess ultradian ACTH rhythms in HGRKO mice because of blood volume
constraints and the stress associated with serial sampling, but corticosterone secretion in these
animals follows a typical circadian pattern[37]. Moreover, continuous CRH infusion restores
circadian ACTH and CORT rhythms in CRH knockout mice[51]. Consistent with our findings in
mice, limited clinical data suggests that CRH-dependent ACTH signaling but not intermittent,
pharmacologic treatment with ACTH may protect against GIAI. Patients with ectopic CRH-
producing tumors, though rare, recover HPA axis function more rapidly after tumor resection than
those with ACTH-secreting adenomas[52]. In contrast, combination therapy with intramuscular
ACTH and glucocorticoids produced more severe HPA axis suppression in 18 patients with
rheumatologic disease compared to those treated with glucocorticoids alone. Together, these
findings identify sustained, potentially CRH-driven, tropic stimulation of the adrenal as a
promising strategy to prevent GIAI. Future translational efforts should define the characteristics of
ACTH signaling that are required to preserve adrenal function during chronic glucocorticoid
therapy.

As a potential limitation, this study only included male mice to minimize biological variability
across a broad, exploratory survey of HPA axis recovery. While there are no reported sex
differences in the risk, severity, or duration of GIAI in humans[10,53-55], HPA axis responsiveness,
rates of adrenocortical turnover, and adrenal macrophage populations are all sexually dimorphic
in rodents[44,56-62]. Future studies will specifically investigate sex differences in the mechanisms of
post-withdrawal GIAI and time to HPA axis recovery in this preclinical model. Additionally, sample
sizes were relatively modest to permit interrogation of HPA axis function at multiple timepoints
after chronic glucocorticoid treatment. Since we interrogated HPA axis function in the stimulated
state, as is done to clinically diagnose adrenal insufficiency, key analyses were all still adequately
powered. Nonetheless, subsequent mechanistic studies will include larger cohorts of animals now
that biologically relevant timepoints have been identified.

In conclusion, our findings demonstrate that post-withdrawal GIAI arises primarily from
adrenal dysfunction rather than delayed hypothalamic-pituitary recovery. This prolonged adrenal
suppression, driven by both adrenocortical cell loss and impaired steroidogenesis, is prevented
by maintaining sustained trophic stimulation of the adrenal during glucocorticoid exposure. These
results establish the adrenal as the principal locus of post-withdrawal GIAI and an accessible
target for therapeutic intervention. Pharmacologic strategies that reproduce physiologic adrenal
stimulation could prevent GIAI, thereby reducing the morbidity associated with one of the most
widely prescribed drug classes while preserving their therapeutic efficacy.

Materials and Methods

Animals

All animal protocols were in accordance with the National Institutes of Health guidelines and
were approved by the Animal Care and Use Committee of Boston Children's Hospital (Protocol
00001983). Mice were housed on a 12-hour/12-hour light-dark cycle (lights on at 7 am) with ad
libitum access to rodent chow and water unless otherwise indicated. Mice used for studies were
all aged either 8 weeks at the time of DEX initiation (long-term studies) or 8-16 weeks (DEX-naïve
controls or short-term studies).

Wild-type (WT) mice: C57BL/6J mice (The Jackson Laboratory, Bar Harbor, ME) were
maintained by breeding within our population.

Sim1^Cre:GR^fl/fl mice (HGRKO): Sim1[Cre] mice (generous gift of Brad Lowell, Beth Israel Lahey
Medical Center, Boston, MA) were bred to floxed GR exon 3 mice (B6.Cg-*Nr3c1^tm1.1Jda*/J, Catalog
#021021, The Jackson Laboratory, Bar Harbor, ME), both maintained on an inbred C57BL/6J
background[37,63]. Littermates that were Cre negative but had GR exon 3 floxed alleles were used
as controls. Genotyping was performed as per Friedman *et al*[64] using the primers and conditions
detailed in Table SI.

Pharmacologic treatments: Mice were treated with either water-soluble dexamethasone
(Sigma-Aldrich 2915) at a concentration of 3.6 mcg/mL (~20 mcg/day assuming an average water
intake of 5.8 mL/day[65]) or no-drug control via drinking water for either 4 days (short-term) or 8
weeks (long-term studies). When specified, mice were co-treated with 10 mcg/kg cosyntropin,
(synthetic ACTH 1-24, Sandoz, NDC 0781-3440-95), dissolved at 10 mcg/mL in 0.9% saline.
Cosyntropin was administered once daily by intraperitoneal injection at 16:30 to align with the
murine, circadian ACTH peak, for either 4 days or 8 weeks, as above.

Hormonal assays: All testing was performed, as specified, on separate days so that no
mouse was bled more than once on a given day. Plasma for measurement of corticosterone and
ACTH was obtained by rapid retroorbital phlebotomy without anesthesia into EDTA-coated
capillary tubes with a total time from first handling the animal to completion of bleeding not
exceeding 30 seconds. Basal/unstimulated phlebotomy was performed at 16:30 in a ventilated
hood in the same room in which the animals were housed. Cosyntropin stimulation[51], insulin-
induced hypoglycemia testing (1 U/kg dose)[66], and physical restraint (20 minutes)[66] were all
performed as previously published. Blood was collected on ice, and plasma was separated by
centrifugation at 1500 RPM at 4°C and stored at -80°C until assay. Plasma concentrations of
corticosterone (ALPCO, 55-CORMS-E01) and ACTH (21-ACTHU-E01) were measured by ELISA
per manufacturer instructions. For the ACTH assay, plasma was diluted 1:10 with ACTH 0
calibrator (ALPCO; 11-C-270-10-G25). The lower limit of detection is 0.061 mcg/dL and 0.22
pg/mL for the CORT and ACTH assays, respectively.

Necroscopy and tissue preparation: Animals were euthanized by rapid decapitation
without anesthesia followed by necroscopy for isolation of hypothalami and adrenals. Coronal
slabs flanking the hypothalamus were prepared with a surgical blade and then fixed in 10%
formalin overnight before transfer to 70% ethanol while awaiting paraffin embedding. Adrenals
were trimmed of surrounding fat tissue, rinsed in phosphate-buffered saline (PBS), and weighed
before overnight fixation in 10% neutral buffered formalin.

Single-molecule RNA in situ hybridization

Three 5 µm-thick coronal brain sections were mounted onto each slide, and every other slide
was stained with hematoxylin and visualized with brightfield microscopy (Nikon E800). The
paraventricular nucleus was identified histologically based on its characteristic "butterfly-shaped"
appearance in coronal sections, which was used to anchor selection of 2 unstained slides
immediately rostral and caudal to this structure for analysis. Single-molecule RNA in situ
hybridization was performed using a catalog Mus musculus *Crh* probe (ACD #316091) and 2.5
HD RED detection kit (ACD #316091) following manufacturer's protocol. Slides were counter
stained with 50% Gill's Hematoxylin II and mounted with EcoMount. Positive control *Ppib (*ACD

#313911) and negative control *dapB (*ACD #310043) were used in every experiment to ensure
sample quality. Brightfield images at 40x magnification were generated with an Olympus VS120
Virtual Slide Scanner and then analyzed with semi-quantitative densitometry using the Weka
Segmentation plugin for Fiji ImageJ per ACDBio instructions[67]. Analysis was restricted to
freehand sections encompassing the characteristic "butterfly" morphology of PVH cell bodies, and
CRH expression, reported as arbitrary units (AU) was quantified by dividing the area of the *Crh*
densitometric signal (μm^2) by the total area of this region (mm^2).

Total mRNA and protein staining

For quantifying total mRNA *in situ*, FFPE sections were deparaffinized and antigen-retrieved
in buffer TE (pH=9). We then adapted the protocol in Farny *et al.* and Hurt *et al.* to label polyA-
capped RNAs on FFPE sections using Cy5-conjugated polyDT (26-4430-02, GeneLink)[27,28]. Total
protein *in situ* was measured using the N-Hydroxysuccinimide (NHS) ester method; NHS esters
react with primary amines and well-approximate total protein mass[29,30]. Briefly, FFPE sections
were deparaffinized and antigen-retrieved in buffer TE (pH=9). Slides were stained with NHS-
Alexa 568 dye conjugates (Thermo Scientific A20003) in PBS for 30 minutes and then washed 5X
in PBS.

Paraffin section immunofluorescence

After fixation, adrenals were oriented with their broadest surface parallel to the cutting plane
in 4% low-melting-temperature SeaPlaque agarose (Lonza, #50101). Agarose blocks were then
dehydrated in ethanol, xylene, and embedded in paraffin blocks. Paraffin sections were cut at 5
μm thickness, and immunofluorescence was performed as described in Leng *et al.*[68] with the
antigen retrieval conditions and antibody concentrations detailed in Table S2. When indicated,
Streptavidin, Alexa Fluor 488 Conjugate (Invitrogen, #S112233) was added to the secondary
antibody solution at a final concentration of 1:500. In all experiments, sections were incubated in
TrueBlack Lipofuscin Autofluorescence Quencher (Biotium, #23007) in 70% EtOH at RT for 3
minutes and rinsed three times in PBS prior to mounting with Prolong Gold Antifade Mountant
with DAPI (Invitrogen, #P36931). Images were acquired with either a Nikon upright Eclipse 90i
microscope with a 20x/0.75 Plan-Apochromat objective or an Olympus VS120 Virtual Slide
Microscope using a 20x objective and adjusted for brightness and contrast in ImageJ.

Floating section immunofluorescence.

After fixation, adrenals were oriented with their broadest surface parallel to the cutting
plane in 4% low-melting-temperature SeaPlaque agarose (Lonza, #50101) and sectioned at a
thickness of 100 μm with a vibratome. Individual slices were separated by free floating in PBS.
Sections were wet mounted in PBS on Superfrost Plus slides (Fisher Scientific, 12-550-15) and
imaged with a 4x objective on a Nikon E800 brightfield microscope. The area of each section was
measured in ImageJ, and the largest section was selected for subsequent immunostaining.
Floating section immunofluorescence was then performed as per Leng *et al*[68] with the addition of
Streptavidin, Alexa Fluor 488 Conjugate (1:500, Invitrogen, #S112233) to the secondary antibody
solution. Images were acquired with a Nikon Ti2 microscope with a Yokogawa CSUW1 confocal
unit, a Nikon Plan Apo Lambda 20x objective lens and a Zyla 4.2 PLUS sCMOS camera
(ANDOR). All images were taken with identical laser strength, gain and exposure settings for the
405nm (used to capture autofluorescence), 488nm, and 647nm channels. Each image was
acquired using the Large Image function in the NIS elements software. This function allowed us
to take images in a 4x4 grid with 15% overlap, allowing the images to be stitched together.
Brightness and contrast were adjusted in ImageJ.

Image analysis

Fluorescent images were analyzed using QuPath (version 0.4.3), an open-source software
for digital pathology.

Positive Cell Detection: QuPath's cell detection feature was employed to label cells as
either Ki-67 or TUNEL positive based on co-localization of the 647 channel signal with the DAPI
nuclear signal. Images were opened in fluorescence mode, and regions of interest (ROI) were
manually defined to include only the adrenal cortex based on morphologic characteristics. ROI
were further restricted to the zF for Ki-67 analysis, as defined by DAB2-negative cortical cells.
The "cell detection" feature was then run specifying DAPI as the detection channel and the 647
nm channel nuclear (Ki-67 analysis) or cellular (TUNEL analysis) mean as the score
compartment. Nuclear size and intensity threshold parameters for both were adjusted and
inspected visually to ensure accurate positive cell detection. Any cell exhibiting signal overlap
between the DAPI and 647 nm channels above the specified thresholds was classified as either
TUNEL or Ki-67 positive and reported as a percentage among total cells.

Morphometry: Morphometric analysis was performed on thick, floating section
immunofluorescent specimens to quantify the area of either the adrenal cortex (Streptavidin
staining, 488 nm channel), medulla (tyrosine hydroxylase immunostaining, 647 nm channel), or
adrenal macrophages (autofluorescence, 408 nm channel) based on signal intensity across
multiple channels using the "create thresholder" function in QuPath. Images were opened in
fluorescence mode, and brightness/contrast of individual channels were adjusted for optimal
signal visualization. Regions of interest were manually defined to include the entire adrenal slice
but not surrounding adipose tissue. For each channel, QuPath's intensity thresholding was
applied to ensure accurate delineation of stained or autofluorescent areas, as confirmed by visual
inspection. As DEX treatment has been shown to attenuate streptavidin staining in the murine zF,
we applied a lower threshold in 8+0 adrenals to ensure detection of all residual adrenocortical
tissue[31]. The area of tissue with a signal above this threshold for each channel was classified as
positive and reported in square microns.

Statistics

Two-tailed Student's t tests were performed to compare significance among two groups.
Comparisons done beyond 2 groups were performed by ANOVA, and Bonferroni correction was
applied for post-hoc, pairwise multiple comparison testing. Nonlinear regression using an
exponential growth equation was used to model stimulated CORT recovery (dependent variable)
over time (independent variable), and doubling time was computed as $\ln(2)/K$ where K is the rate
constant. When two exponential growth models were compared, the extra-sum-of-squares test
was employed to determine whether the exponential equations provided significantly different fits
to the data. All statistical analyses were performed in GraphPad Prism v.10.3.1. Windows,
GraphPad Software, Boston, Massachusetts USA, www.graphpad.com. *P* values below 0.05
were considered statistically significant.

Study approval

All animal experiments were approved by the Boston Children's Hospital Institutional Animal
Care and Use Committee (Protocol Number 00001983).

Data availability

All raw data associated with findings presented in this manuscript are included in the
Supporting Data Values file and are available from the corresponding author upon reasonable
request.

Acknowledgments

This study was supported by NIH 5T32DK007699-39 (LG), NIH F32DK131795-02 (LG), a Pediatric Endocrine Society Research Fellowship Award (LG), a Pediatric Endocrine Society Clinical Scholar Award (LG), a Boston Children's Hospital Office of Faculty Development/Basic & Clinical Translational Research Executive Committees Faculty Career Development Fellowship (LG), and the Thomas Morgan Rotch Harvard Medical School endowment (JAM). We thank Drs. David Breault, Kleiton Borges, and Mesut Berber for their input on this project and assistance processing adrenal tissue, identifying zonal markers, and performing thick, floating section immunofluorescence. We also thank Suzanne White from the BIDMC Histology core for preparing and sectioning all FFPE specimens, including working closely with the author to isolate the PVH in mouse brains.

References

1. Fardet L, Petersen I, Nazareth I. Prevalence of long-term oral glucocorticoid prescriptions in the UK over the past 20 years. *Rheumatology (Oxford).* 2011;50(11):1982-1990.

2. Overman RA, Yeh JY, Deal CL. Prevalence of oral glucocorticoid usage in the United States: a general population perspective. *Arthritis Care Res (Hoboken).* 2013;65(2):294-298.

3. Daley-Yates P, Brealey N, Thomas S, et al. Therapeutic index of inhaled corticosteroids in asthma: A dose-response comparison on airway hyperresponsiveness and adrenal axis suppression. *Br J Clin Pharmacol.* 2021;87(2):483-493.

4. Woods CP, Argese N, Chapman M, et al. Adrenal suppression in patients taking inhaled glucocorticoids is highly prevalent and management can be guided by morning cortisol. *European journal of endocrinology.* 2015;173(5):633-642.

5. Lam LH, Sugarman JL. Adrenal Suppression With Chronic Topical Corticosteroid Use in Psoriasis Patients. *J Drugs Dermatol.* 2016;15(8):945-948.

6. Keller-Wood ME, Dallman MF. Corticosteroid inhibition of ACTH secretion. *Endocrine reviews.* 1984;5(1):1-24.

7. Wyllie AH, Kerr JF, Currie AR. Cellular events in the adrenal cortex following ACTH deprivation. *The Journal of pathology.* 1972;106(1):Pix.

8. Wyllie AH, Kerr JF, Macaskill IA, Currie AR. Adrenocortical cell deletion: the role of ACTH. *The Journal of pathology.* 1973;111(2):85-94.

9. Younes AK, Younes NK. Recovery of steroid induced adrenal insufficiency. *Translational pediatrics.* 2017;6(4):269-273.

10. Livanou T, Ferriman D, James VH. Recovery of hypothalamo-pituitary-adrenal function after corticosteroid therapy. *Lancet (London, England).* 1967;2(7521):856-859.

11. Beuschlein F, Else T, Bancos I, et al. European Society of Endocrinology and Endocrine Society Joint Clinical Guideline: Diagnosis and therapy of glucocorticoid-induced adrenal insufficiency. *European journal of endocrinology.* 2024;190(5):G25-g51.

12. Ebrahimi F, Widmer A, Wagner U, et al. Association of adrenal insufficiency with patient-oriented health-care outcomes in adult medical inpatients. *European journal of endocrinology.* 2019;181(6):701-709.

13. Mebrahtu TF, Morgan AW, Keeley A, Baxter PD, Stewart PM, Pujades-Rodriguez M. Dose dependency of iatrogenic glucocorticoid excess and adrenal insufficiency and mortality: a cohort study in England. *The Journal of clinical endocrinology and metabolism.* 2019;104(9):3757-3767.

14. Graber AL, Ney RL, Nicholson WE, Island DP, Liddle GW. NATURAL HISTORY OF PITUITARY-ADRENAL RECOVERY FOLLOWING LONG-TERM SUPPRESSION WITH CORTICOSTEROIDS. *The Journal of clinical endocrinology and metabolism.* 1965;25:11-16.

15. Ney RL, Shimizu N, Nicholson WE, Island DP, Liddle GW. CORRELATION OF PLASMA ACTH CONCENTRATION WITH ADRENOCORTICAL RESPONSE IN NORMAL HUMAN

SUBJECTS, SURGICAL PATIENTS, AND PATIENTS WITH CUSHING'S DISEASE. *The Journal of clinical investigation.* 1963;42(11):1669-1677.

16.   Schlaghecke R, Kornely E, Santen RT, Ridderskamp P. The effect of long-term glucocorticoid therapy on pituitary-adrenal responses to exogenous corticotropin-releasing hormone. *The New England journal of medicine.* 1992;326(4):226-230.

17.   Rao PS. Corticosteroids. In: Wexler P, ed. *Encyclopedia of Toxicology (Second Edition).* New York: Elsevier; 2005:669-670.

18.   Avgerinos PC, Chrousos GP, Nieman LK, Oldfield EH, Loriaux DL, Cutler GB, Jr. The corticotropin-releasing hormone test in the postoperative evaluation of patients with cushing's syndrome. *The Journal of clinical endocrinology and metabolism.* 1987;65(5):906-913.

19.   Avgerinos PC, Nieman LK, Oldfield EH, et al. The effect of pulsatile human corticotropin-releasing hormone administration on the adrenal insufficiency that follows cure of Cushing's disease. *The Journal of clinical endocrinology and metabolism.* 1989;68(5):912-916.

20.   Gomez MT, Magiakou MA, Mastorakos G, Chrousos GP. The pituitary corticotroph is not the rate limiting step in the postoperative recovery of the hypothalamic-pituitary-adrenal axis in patients with Cushing syndrome. *The Journal of clinical endocrinology and metabolism.* 1993;77(1):173-177.

21.   Dauber A, Feldman HA, Majzoub JA. Nocturnal Dexamethasone versus Hydrocortisone for the Treatment of Children with Congenital Adrenal Hyperplasia. *Int J Pediatr Endocrinol.* 2010;2010.

22.   Mallappa A, Sinaii N, Kumar P, et al. A phase 2 study of Chronocort, a modified-release formulation of hydrocortisone, in the treatment of adults with classic congenital adrenal hyperplasia. *The Journal of clinical endocrinology and metabolism.* 2015;100(3):1137-1145.

23.   Spiga F, Zhao Z, Lightman SL. Prolonged treatment with the synthetic glucocorticoid methylprednisolone affects adrenal steroidogenic function and response to inflammatory stress in the rat. *Brain Behav Immun.* 2020;87:703-714.

24.   Calogero AE, Kamilaris TC, Johnson EO, Tartaglia ME, Chrousos G. Recovery of the rat hypothalamic-pituitary-adrenal axis after discontinuation of prolonged treatment with the synthetic glucocorticoid agonist dexamethasone. *Endocrinology.* 1990;127(4):1574-1579.

25.   Duncan PJ, McClafferty H, Nolan O, et al. Corticotroph isolation from Pomc-eGFP mice reveals sustained transcriptional dysregulation characterising a mouse model of glucocorticoid-induced suppression of the hypothalamus-pituitary-adrenal axis. *Journal of neuroendocrinology.* 2022;34(7):e13165.

26.   Sakr N, Glazova O, Shevkova L, et al. Characterizing and Quenching Autofluorescence in Fixed Mouse Adrenal Cortex Tissue. *Int J Mol Sci.* 2023;24(4).

27.   Farny NG, Hurt JA, Silver PA. Definition of global and transcript-specific mRNA export pathways in metazoans. *Genes Dev.* 2008;22(1):66-78.

28.   Hurt JA, Obar RA, Zhai B, Farny NG, Gygi SP, Silver PA. A conserved CCCH-type zinc finger protein regulates mRNA nuclear adenylation and export. *J Cell Biol.* 2009;185(2):265-277.

29.   Liu X, Oh S, Kirschner MW. The uniformity and stability of cellular mass density in mammalian cell culture. *Front Cell Dev Biol.* 2022;10:1017499.

30.   Kafri R, Levy J, Ginzberg MB, Oh S, Lahav G, Kirschner MW. Dynamics extracted from fixed cells reveal feedback linking cell growth to cell cycle. *Nature.* 2013;494(7438):480-483.

31.   Paul A, Laufer E. Endogenous biotin as a marker of adrenocortical cells with steroidogenic potential. *Molecular and cellular endocrinology.* 2011;336(1-2):133-140.

32.   Wentworth JM, Naselli G, Brown WA, et al. Pro-inflammatory CD11c+CD206+ adipose tissue macrophages are associated with insulin resistance in human obesity. *Diabetes.* 2010;59(7):1648-1656.

33. Wu H, Gower RM, Wang H, et al. Functional role of CD11c+ monocytes in atherogenesis
associated with hypercholesterolemia. *Circulation.* 2009;119(20):2708-2717.
34. Xu Y, Patterson MT, Dolfi B, et al. Adrenal gland macrophages regulate glucocorticoid
production through Trem2 and TGF-β. *JCI Insight.* 2024;9(14).
35. Ganguly S, Rosenthal SB, Ishizuka K, et al. Lipid-associated macrophages' promotion of
fibrosis resolution during MASH regression requires TREM2. *Proceedings of the National*
Academy of Sciences of the United States of America. 2024;121(35):e2405746121.
36. Dallman MF. Control of adrenocortical growth in vivo. *Endocrine research.* 1984;10(3-
4):213-242.
37. Laryea G, Schütz G, Muglia LJ. Disrupting hypothalamic glucocorticoid receptors causes
HPA axis hyperactivity and excess adiposity. *Molecular endocrinology (Baltimore, Md).*
2013;27(10):1655-1665.
38. Freedman BD, Kempna PB, Carlone DL, et al. Adrenocortical zonation results from lineage
conversion of differentiated zona glomerulosa cells. *Developmental cell.* 2013;26(6):666-
673.
39. Kataoka Y, Ikehara Y, Hattori T. Cell proliferation and renewal of mouse adrenal cortex. *J*
Anat. 1996;188 (Pt 2)(Pt 2):375-381.
40. Yates R, Katugampola H, Cavlan D, et al. Adrenocortical development, maintenance, and
disease. *Curr Top Dev Biol.* 2013;106:239-312.
41. Hume DA, Halpin D, Charlton H, Gordon S. The mononuclear phagocyte system of the
mouse defined by immunohistochemical localization of antigen F4/80: macrophages of
endocrine organs. *Proceedings of the National Academy of Sciences of the United States*
of America. 1984;81(13):4174-4177.
42. O'Brien CJ, Sidarta-Oliveira D, Stawiarski A, et al. Distinct adrenal gland macrophages
regulate corticosteroid production. *bioRxiv.* 2023:2023.2009.2005.556330.
43. González-Hernández JA, Bornstein SR, Ehrhart-Bornstein M, Geschwend JE, Adler G,
Scherbaum WA. Macrophages within the human adrenal gland. *Cell and Tissue Research.*
1994;278(2):201-205.
44. Warde KM, Smith LJ, Liu L, et al. Senescence-induced immune remodeling facilitates
metastatic adrenal cancer in a sex-dimorphic manner. *Nat Aging.* 2023;3(7):846-865.
45. Lazarov T, Juarez-Carreño S, Cox N, Geissmann F. Physiology and diseases of tissue-
resident macrophages. *Nature.* 2023;618(7966):698-707.
46. Foley KG, Pritchard MT, Duncan FE. Macrophage-derived multinucleated giant cells:
hallmarks of the aging ovary. *Reproduction.* 2021;161(2):V5-v9.
47. Braune J, Lindhorst A, Fröba J, et al. Multinucleated Giant Cells in Adipose Tissue Are
Specialized in Adipocyte Degradation. *Diabetes.* 2021;70(2):538-548.
48. Furman BL. Cosyntropin☆. In: *Reference Module in Biomedical Sciences.* Elsevier; 2017.
49. Furman BL. Cosyntropin. In: Enna SJ, Bylund DB, eds. *xPharm: The Comprehensive*
Pharmacology Reference. New York: Elsevier; 2007:1-3.
50. Spiga F, Zavala E, Walker JJ, Zhao Z, Terry JR, Lightman SL. Dynamic responses of the
adrenal steroidogenic regulatory network. *Proceedings of the National Academy of*
Sciences of the United States of America. 2017;114(31):E6466-e6474.
51. Muglia LJ, Jacobson L, Weninger SC, et al. Impaired diurnal adrenal rhythmicity restored
by constant infusion of corticotropin-releasing hormone in corticotropin-releasing hormone-
deficient mice. *The Journal of clinical investigation.* 1997;99(12):2923-2929.
52. Lois KB, Santhakumar A, Vaikkakara S, et al. Phaeochromocytoma and ACTH-dependent
cushing's syndrome: tumour crf secretion can mimic pituitary cushing's disease. *Clinical*
endocrinology. 2016;84(2):177-184.
53. Broersen LH, Pereira AM, Jørgensen JO, Dekkers OM. Adrenal Insufficiency in
Corticosteroids Use: Systematic Review and Meta-Analysis. *The Journal of clinical*
endocrinology and metabolism. 2015;100(6):2171-2180.
54. Prete A, Bancos I. Glucocorticoid induced adrenal insufficiency. *BMJ.* 2021;374:n1380.

55.   Prete A, Paragliola RM, Bottiglieri F, et al. Factors predicting the duration of adrenal
686   insufficiency in patients successfully treated for Cushing disease and nonmalignant primary
687   adrenal Cushing syndrome. *Endocrine.* 2017;55(3):969-980.
688   56.   Gao X, Yamazaki Y, Tezuka Y, et al. Gender differences in human adrenal cortex and its
689   disorders. *Molecular and cellular endocrinology.* 2021;526:111177.
690   57.   Duncko R, Kiss A, Skultétyová I, Rusnák M, Jezová D. Corticotropin-releasing hormone
691   mRNA levels in response to chronic mild stress rise in male but not in female rats while
692   tyrosine hydroxylase mRNA levels decrease in both sexes. *Psychoneuroendocrinology.*
693   2001;26(1):77-89.
694   58.   Viau V, Bingham B, Davis J, Lee P, Wong M. Gender and puberty interact on the stress-
695   induced activation of parvocellular neurosecretory neurons and corticotropin-releasing
696   hormone messenger ribonucleic acid expression in the rat. *Endocrinology.*
697   2005;146(1):137-146.
698   59.   Goel N, Bale TL. Sex differences in the serotonergic influence on the hypothalamic-
699   pituitary-adrenal stress axis. *Endocrinology.* 2010;151(4):1784-1794.
700   60.   Grabek A, Dolfi B, Klein B, Jian-Motamedi F, Chaboissier MC, Schedl A. The Adult Adrenal
701   Cortex Undergoes Rapid Tissue Renewal in a Sex-Specific Manner. *Cell Stem Cell.*
702   2019;25(2):290-296.e292.
703   61.   Hammer GD, Basham KJ. Stem cell function and plasticity in the normal physiology of the
704   adrenal cortex. *Molecular and cellular endocrinology.* 2021;519:111043.
705   62.   Dolfi B, Gallerand A, Firulyova MM, et al. Unravelling the sex-specific diversity and
706   functions of adrenal gland macrophages. *Cell reports.* 2022;39(11):110949.
707   63.   Tronche F, Kellendonk C, Kretz O, et al. Disruption of the glucocorticoid receptor gene in
708   the nervous system results in reduced anxiety. *Nature genetics.* 1999;23(1):99-103.
709   64.   Friedman HR, Gaston LS, Chan LF, Majzoub JA. Absent, but not glucocorticoid-modulated,
710   corticotropin-releasing hormone (Crh) regulates anxiety-like behaviors in mice. *bioRxiv.*
711   2024:2024.2009.2023.614550.
712   65.   Bachmanov AA, Reed DR, Beauchamp GK, Tordoff MG. Food intake, water intake, and
713   drinking spout side preference of 28 mouse strains. *Behav Genet.* 2002;32(6):435-443.
714   66.   Jacobson L, Muglia LJ, Weninger SC, Pacák K, Majzoub JA. CRH deficiency impairs but
715   does not block pituitary-adrenal responses to diverse stressors. *Neuroendocrinology.*
716   2000;71(2):79-87.
717   67.   Advanced Cell Diagnostics. (2019). TS-46-003 Technical Note: How to correctly interpret
718   your RNAscope images. Rev B. Retrieved from https://acdbio.com/2019-mar-26-how-
719   correctly-interpret-your-rnascope-images. In.
720   68.   Leng S, Pignatti E, Khetani RS, et al. β-Catenin and FGFR2 regulate postnatal rosette-
721   based adrenocortical morphogenesis. *Nat Commun.* 2020;11(1):1680.

    Figures and Tables

Figure 1. **Corticosterone (CORT) secretion is suppressed for up to 8 weeks after discontinuing long-term DEX despite rapid hypothalamic-pituitary recovery.** A) *In situ* hybridization (RNAScope) for basal *Crh* (red) in the paraventricular nucleus (PVH) of the hypothalamus in DEX-naïve (DN) animals and those treated with 8 weeks DEX both immediately prior to (8+0) and 1-week post-withdrawal (8+1; top row, scale bar=200 microns). Nuclei are labeled by hematoxylin (purple). Magnification of region within the black box is shown directly below (bottom row, scale bar=50 microns). B) Quantification of *Crh* mRNA expression from RNAScope images of the PVH from DN, 8+0, and 8+1 animals (n=3). $p<0.001$ by 1-way ANOVA. Recovery of insulin-stimulated plasma C) ACTH and D) CORT and E) cosyntropin-stimulated CORT levels 0-8 weeks after withdrawal of an 8-week DEX course. Data shown as individual mice with mean (connecting line) ± SEM (error bars). Grey bar (mean± SEM) indicates the reference range for adult, male, wildtype, untreated (DEX-naïve or DN) controls (n=4-8). * denotes $p<0.05$, ** $p<0.01$, *** $p<0.001$, and ****$p<0.0001$ vs. DN.

Figure 2. Long-term DEX induces adrenocortical atrophy with accumulation of giant macrophage clusters that persist for at least 8 weeks post-withdrawal. A) DEX-treated adrenals weigh significantly less than DN controls through 6 weeks post-withdrawal. P<0.0001 by one-way ANOVA; * indicates p<0.05, ** p<0.01, *** p<0.001, and ****p<0.0001 vs. DN. B) Hematoxylin and eosin staining reveals large, poorly defined, vacuolated structures with decreased nuclear density at the corticomedullary junction in DEX-exposed adrenals that persist through 8-weeks post-withdrawal. Representative structures (outlined with red, dashed circles) correspond with clusters of cells that are positive for the classical macrophage markers C) CD68

and D) F4/80. C) Immunofluorescence for CD68 (magenta) with DAPI (blue) and 488-streptavidin
(green) counterstains, the latter of which labels all biotin-rich cells of the adrenal cortex.
Representative images shown for DN adrenals, those on DEX for 2-8 weeks (e.g. 2+0 to 8+0), as
well as one (8+1) and eight (8+8) weeks after an 8-week DEX course. D) Immunofluorescence for
the mouse macrophage marker F4/80 (green) and medullary marker tyrosine hydroxylase (TH;
magenta) in DN and DEX-exposed adrenals (2+0 and 8+0). Magnification of representative
macrophages in 8+0 glands from E) H&E and immunostaining for F) CD68 and G) F4/80. All
staining performed with TrueBlack lipofuscin autofluorescence quencher (C, D, F, and G).

A.

B. Pan-cortical area (mm²)

C. Medullary area (mm²)

D. AM area (mm²)

Figure 3. Stimulated CORT secretion is disproportionately low relative to adrenocortical cell mass for at least 2 weeks after withdrawal of long-term DEX. A) IF without TrueBlack for tyrosine hydroxylase (TH, red) DAPI (blue) and 488-streptavidin (green; columns 1 and 2) counterstains. Adrenal macrophages (AMs) were detected by autofluorescence (cyan; columns 3 and 4). Mice were either untreated (DN) or treated for 8 weeks DEX followed by withdrawal for 0-8 weeks (8+0, 8+1, 8+2, 8+6, and 8+8). Shown as 100 µM-thick, whole adrenal slices (columns 1 and 3, scale bar=500 microns) with magnification of the region in the white box (columns 2 and 4, scale bar=100 microns). Columns 1 and 2 imaged at wavelengths 405 nm, 488 nm, and 647 nm; columns 3 and 4 imaged at wavelengths 405 nm, 440 nm, and 647 nm. Morphometric analysis of B) pan-cortical area, which includes both streptavidin-positive and autofluorescent regions (p=0.18 by one-way ANOVA), C) Autofluorescent AM area (p=0.18), D) steroidogenic cell type area, calculated as pan-cortical area minus AMs (p<0.01), and E) medullary (TH- positive) area (p=0.17 by one-way ANOVA). F) Ratio of steroidogenic cell type area to cosyntropin-stimulated CORT secretion, normalized to DN controls. * p<0.05, ** p<0.01, *** p<0.001 vs. DN (B-F)

**Figure 4. Selective deletion of the hypothalamic glucocorticoid receptor (GR) confers
protection against DEX-induced adrenal suppression.** A) Short-term (4 days DEX) induces

apoptosis in DEX-treated wild-type (4D DEX) but not Sim1^{Cre}:GR$^{fl/fl}$ (HGRKO) mice, as shown by
TUNEL staining (magenta) with B) quantification. $p<0.0001$ by one-way ANOVA. **** denotes
$p<0.0001$ for select pairwise comparisons. HGRKO mice and Cre-negative, functionally wild-type
littermate controls (HGRWT) mice were then treated with DEX for 8 weeks. Hormonal function
was assessed pre-treatment on long-term DEX. C) Basal (unstressed circadian peak) ACTH
levels in HGRWT vs. HGRKO animals prior to DEX (left of dashed line) and on 8 weeks of DEX
(right of dashed line). $p=0.19$ for DEX status and $p<0.0001$ for genotype by two-way ANOVA. H)
CORT secretion in the basal state, following 20 minutes of physical restraint (Restraint), and
following cosyntropin stimulation (COS-stim.) in HGRWT vs HGRKO animals, either prior to DEX
treatment (left of dashed line) vs. on 8 weeks of DEX (right of dashed line). E) Adrenal weights
and morphometric analyses for F) pan-cortical area, G) AM area, and H) steroidogenic cell type
area in HGRWT and HGRKO adrenals on long-term DEX (8+0), all computed as per Fig. 3 and
normalized to DN adrenals from Fig. 3. I) Representative thick, floating section IF without
TrueBlack for tyrosine hydroxylase (TH, red), with DAPI (blue), and 488-streptavidin (green);
methods and visualization as per Fig. 3. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$ by t-test
(HGRWT vs. HGRKO (C-H).