Rapid hypothalamic-pituitary recovery after chronic glucocorticoid therapy enables

strategies that prevent adrenal suppression

Lindsey S. Gaston[1], Brenna C. Jorgensen[1], Hannah R. Friedman[1], Marc S. Sherman[2], Joseph

A. Majzoub[1]

Affiliations:

[1]Division of Endocrinology, Department of Pediatrics, Boston Children's Hospital, Harvard

Medical School, Boston, MA, USA

[2]Division of Gastroenterology, Department of Medicine, Massachusetts General hospital,

Harvard Medical School, Boston, MA, USA

Conflict-of-interest Statement: The authors have declared that no conflict of interest exists

ABSTRACT

Glucocorticoid-induced adrenal insufficiency (GIAI) persists for months to years after long-term

treatment, which past studies suggest is mediated by initial, post-withdrawal hypothalamic-

pituitary suppression. We examined the timing of hypothalamic, pituitary, and adrenal (HPA)

recovery after 8 weeks of dexamethasone (DEX) treatment in mice. At withdrawal, HPA function

was fully suppressed, but *Crh* mRNA and plasma ACTH surprisingly rebounded above control

levels within 1 week. Despite this, corticosterone (CORT) remained suppressed for an additional

7 weeks. DEX-exposed adrenals were atrophic and contained previously undescribed, giant,

multinucleated histiocytes. CORT secretion was disproportionately low relative to the non-

histocyte adrenocortical area in the first 1-2 weeks post-DEX, suggesting that post-withdrawal

GIAI is mediated by adrenocortical atrophy with superimposed, reduced ACTH responsiveness.

We then tested whether trophic stimulation of the adrenal during glucocorticoid exposure

prevented GIAI. Daily co-treatment with DEX and cosyntropin (an ACTH analog) failed to

prevent GIAI. However, mice with non-suppressible endogenous ACTH due to knockout of the

hypothalamic glucocorticoid receptor maintained normal adrenal function and histology despite

long-term DEX. Localizing the mechanisms of post-withdrawal GIAI to the adrenal may permit

the development of preventative strategies that mimic native HPA signaling during glucocorticoid

exposure, thus reducing the morbidity of a widely prescribed drug class.

INTRODUCTION

Up to 3% of the United States population is prescribed systemic glucocorticoids, and additional patients are chronically treated with topical and inhaled formulations that can exert systemic toxicity even at moderate doses[1-6]. Exogenous glucocorticoids, like native cortisol in humans and corticosterone in rodents, exert glucocorticoid-receptor (GR)-mediated feedback inhibition on the central nodes of the HPA axis, leading to decreased synthesis and secretion of hypothalamic corticotropin-releasing hormone (CRH) and pituitary adrenocorticopic hormone (ACTH)[7]. ACTH not only acutely drives steroidogenesis but is also a key trophic factor for the adrenal gland[8,9]. In fact, Wyllie and Kerr's earliest morphologic descriptions of apoptosis were in ACTH-deprived adrenals, via either hypophysectomy or short-term prednisolone treatment, which was rescued by concurrent, intraperitoneal ACTH injections[8,10].

Hypothalamic-pituitary-adrenal (HPA) axis suppression is typically transient in patients treated with short-term glucocorticoids; however, treatment exceeding four weeks can cause protracted adrenal insufficiency that can persist for weeks up to years after steroid withdrawal[11-13]. While the axis awaits recovery, individuals are at risk for adrenal crisis with otherwise routine illnesses or trauma, which has been associated with longer hospitalizations, higher rates of ICU admissions, and even death in patients who are not promptly administered stress-dose steroids[14,15]. Despite the ubiquity of glucocorticoid use and the healthcare burden of glucocorticoid-induced iatrogenic adrenal insufficiency (GIAI), the mechanisms of persistent HPA axis dysfunction after discontinuation of steroid treatment remain unknown[14]. Elucidation of these processes may lead to novel therapies that either prevent or hasten recovery from GIAI.

Humans are thought to recover from GIAI by reversal of sequential phases of post-withdrawal central and adrenocortical dysfunction. This is based on the work of Graber and colleagues, who found that patients exposed for a year or more to glucocorticoids had inappropriately low to normal ACTH levels for the first 1-2 months after steroid withdrawal despite release from negative feedback. Basal cortisol required another 5-9 months to normalize even after ACTH rebounded

above control levels[16]. Post-withdrawal central suppression has also been described in patients

treated for Cushing disease, a state of chronic, endogenous ACTH excess driven by an ACTH-

secreting pituitary adenoma[17-20]

Given the evolutionary conservation of the mammalian HPA axis, animal models may address

the limitations of clinical studies in defining the anatomic locus and mechanisms of post-

withdrawal GIAI[21]. Interestingly, all rodent studies-to date have demonstrated rapid restoration of

hypothalamic-pituitary function after steroid treatment. Basal and stimulated ACTH levels

(following lipopolysaccharide, arecholine, and CRH administration) were recovered within days of

discontinuing 5 or 7 days of methylprednisolone or dexamethasone, respectively, in rats[22,23]. In

mice treated with dexamethasone for 30 days, plasma ACTH and corticotroph *Pomc* mRNA levels

were elevated above control animals by 1-week post-withdrawal[24]. In all studies, corticosterone

(CORT) levels and/or steroidogenic pathway gene expression remained low 5-7 days after

discontinuing steroids depiste normalization of adrenal weights[22-24]. No preclinical studies to date

have examined HPA axis recovery after longer courses of glucocorticoids known to cause

protracted adrenal suppression in humans.

To identify the rate-limiting steps in HPA axis recovery and the mechanisms underlying

protracted GIAI following long-term glucocorticoid treatment, we evaluated the temporal dynamics

of hypothalamic, pituitary, and adrenal recovery in mice withdrawn from an 8-week

dexamethasone (DEX) regimen. After finding that CORT secretion remained supressed for 8

weeks after DEX withdrawal despite rapid restoration of hypothalamic-pituitary function, we

attempted to prevent GIAI by providing sustained trophic stimulation to the adrenal during

glucocorticoid exposure using either a pharmacologic or genetic approach. Overall, we found that

the adrenal is the major anatomic locus of post-withdrawal GIAI and is an accessible therapeutic

target for strategies to prevent long-term, post-withdrawal adrenal suppression.

RESULTS

Mice treated with long-term DEX have protracted adrenal insufficiency despite early hypothalamic-pituitary recovery

We first established when hormonal function recovered at each node of the HPA axis following long-term DEX treatment. Adult, male, WT mice were treated with continuous DEX in drinking water for 8 weeks. Starting at the time of DEX withdrawal (8+0) and at 1-2 week intervals through 8+8 (where 8 weeks of treatment plus N weeks of withdrawal = 8+N), ACTH and corticosterone (CORT) were measured in the basal, unstimulated state at 16:30 followed by necroscopy for evaluation of hypothalamic *Crh* mRNA expression by *in situ hybridization* (Model 1, Fig. 1).

As expected, long-term DEX treatment suppressed *Crh* mRNA within the paraventricular nucleus of the hypothalamus (PVH; 28±5 in 8+0 vs. 156±32 arbitrary units [AU] in DEX-naïve (DN) animals, $p<0.05$). Expression recovered to control levels 1 week after DEX was discontinued (8+1; 159±28 AU, $p>0.9999$ vs. DN, Fig. 2A, B). Plasma ACTH levels in the unstimulated, basal state were similarly suppressed on DEX (8+0; 6.1±6.1 pg/mL) but rebounded above controls (DN; 306±125) by 1-week post-withdrawal (1450±207, $p<0.001$, Fig. 2C). This elevation in basal ACTH secretion persisted until 8 weeks post-withdrawal. Basal CORT levels were undetectable on long-term DEX and gradually recovered in the 6 weeks following treatment completion, although differences between DEX-withdrawn and DEX-naïve animals did not reach statistical significance owing to expected variability in unstimulated hormone measurements (Fig. 2D).

In separate, terminal cohorts of mice, we then performed insulin-induced hypoglycemia and cosyntropin-stimulation testing followed by necroscopy at 1-2 week intervals after withdrawal of 8 weeks DEX, as above (Model 1, Fig. 1). Each of these provocative tests has been used to clinically diagnose AI by evaluating HPA axis responsiveness to a physiologically-relevant stressor and maximal CORT secretion to a standardized, pharmacologic stimulus, respectively. As expected, IIH-stimulated ACTH (4.2±3.6 pg/mL) and CORT (1.0±0.24 mcg/dL) as well as cosyntropin-stimulated CORT (2.0±0.61 mcg/dL) levels were suppressed at the completion of

DEX treatment (8+0) relative to DN controls (875±44 pg/mL, 27±1.1 mcg/dL, and 15.7±1.2 mcg/dL for IIH-stimulated ACTH, IIH-stimulated CORT, and cosyntropin-stimulated CORT, respectively; $p<0.001$ for all, Fig 1E-G). However, by one week post-withdrawal (8+1), IIH-stimulated ACTH levels (1707±85 pg/mL) were surprisingly elevated above DN control values (875±44 pg/mL, $p<0.001$). IIH-stimulated ACTH levels remained elevated through 4 weeks post-withdrawal and reached a peak two weeks after completion of DEX treatment (Fig. 1E).

Despite this supraphysiologic ACTH stimulation, both IIH-stimulated (Fig. 1F) and cosyntropin-stimulated CORT levels (Fig. 1G) remained significantly lower than DN-controls until 8 weeks post-DEX withdrawal. Recovery of both IIH-stimulated and cosyntropin-stimulated CORT secretion was modeled with exponential growth equations yielding doubling times of 2.4 ($R^2=0.89$; replicates test for lack of fit $p=0.0001$) and 2.5 weeks ($R^2=0.86$; replicates test for lack of fit $p=0.1056$), respectively (Fig. S1A-B). Mice withdrawn from long-term DEX thus have delayed recovery of CORT secretion despite rapid restoration of hypothalamic-pituitary function.

Long-term DEX induces adrenocortical atrophy with accumulation of giant histiocyte clusters

As the above studies implicate the adrenal rather than the hypothalamus or pituitary as the major anatomic locus of post-withdrawal HPA axis dysfunction, we then assessed adrenal mass and morphology at select time points after long-term DEX treatment. Long-term DEX treatment reduced adrenal weights to 40% of controls (1.0±0.041 mg in 8+0 vs. 2.5±0.063 mg in DN, $p<0.0001$, Fig. 3A). Although adrenals from DEX-exposed adrenals weighed significantly less than those from DN controls through 6 weeks post-withdrawal, distinct patterns of recovery were noted for adrenal mass vs. function. In particular, adrenal mass nearly doubled between 0 to 1 weeks post-withdrawal to 75% of controls (1.9±0.11 mg at 8+1) and remained relatively stable over the remaining recovery period. In contrast, CORT secretion remained 75% lower than control values at 8+1 and more gradually recovered over the subsequent 7 weeks (Fig. S1A and B; raw data derived from Fig. 2E and F and 3A).

Hematoxylin and eosin staining revealed that long-term DEX caused massive, adrenocortical

atrophy followed by gradual reaccumulat ion of cortical tissue in the 8 weeks following

glucocorticoid withdrawal. Interestingly, DEX-exposed adrenals contained large, vacuolated

structures with reduced nuclear density near the corticomedullary junction that persisted through

8 weeks post-withdrawal (Fig. 3B, magnification in 3C). These clusters were intensely

autofluorescent, which was markedly attenuated with the addition of TrueBlack, a lipofuscin

autofluorescence quencher[25] (Fig. 3D). ISH with a 5'-Cy5-Oligod(T)30 demonstrated similar,

nuclear mRNA poly(A) content in these structures compared to surrounding cortical cells[26,27]. The

large, vacuolated, extranuclear space in these clusters was not strongly labeled with 568 NHS

Ester (SE), a total protein stain[28,29]. Together, this suggests that the multinucleated clusters

observed in DEX-exposed adrenals contain live, lipofuscin-containing, biologically active cells

rather than acellular, proteinaceous debris.

We then asked whether these cells represented an atypical steroidogenic population vs.

histiocytes, the latter of which have been identified using morphologic characteristics in rat

adrenals following short-term glucocorticoid treatment[10,30]. Immunofluorescence for the myeloid-

specific surface marker CD68 was performed on adrenals exposed to 2-8 weeks DEX (2+0, 4+0,

and 8+0), as well as those withdrawn from 8-weeks DEX for 1 and 8 weeks (8+1 and 8+8). Glands

were counterstained with fluorescent (488)-conjugated streptavidin, which labels all

steroidogenic cells of the adrenal cortex[31], and TrueBlack to quench lipofuscin autofluorescence

(Fig. 4). Rare, mononuclear, CD68-positive cells were visualized in dexamethasone naïve glands.

More numerous CD68 positive cells were present throughout the zona fasciculata (zF) after 2

weeks of DEX treatment (2+0), which was followed by larger, apparently mutinuclear CD68+

accumulations at the corticomedullary junction with ongoing treatment (at 4+0 and 8+0; Fig. 4A

and B). Large, corticomedullary, CD68-positive accumulations remained present at 1-week post-

withdrawal (8+1), although these clusters appreared to fragment and migrate centripedally by 8

weeks off treatment (8+8; Fig. 4A). Subsequent immunofluorescence with the membrane-bound

mouse macrophage marker F4/80 and adrenal medulla marker tyrosine hydroxylase (TH)

demonstrated expansion of smaller, single-nucleated macrophage-like cells after 2 weeks of DEX.

After 8 weeks of treatment, these cells appeared to aggregate into large, multinuclear, F4/80-

positive clusters at the corticomedullary junction, which we herein term "Giant Histiocytes of

Steroid Treatment" (GHOSTs; Fig. 4C and D).

CORT secretion is disproportionately low relative to steroidogenic cell mass in the first 2 weeks

after DEX withdrawal

Reasoning that GHOSTs might contribute to adrenal mass but not biosynthetic function, we

then asked specifically how the steroidogenic cell population contracted and recovered with

prolonged DEX treatment and withdrawal. To do this, we performed morphometric analysis on

thick, immunostained adrenal sections, which allowed for comparison of the largest, equatorial

adrenal section from each gland. Since no antibodies reliably distinguished adrenocortical cell

types in thick, floating section IF pilot studies (data not shown), we first performed standard IF

on FFPE tissue using Dab2 and Ki-67, markers of the zG and proliferative cells, respectively[32]

(Fig S2). There was no significant change in zona glomerulosa (zG) thickness after long-term

DEX treatment (Fig. S2B). The zF had a slight increase in nuclear density at the completion of

DEX treatment (8+0 vs. DN, Fig. S2C) but was otherwise unchanged in the period after DEX-

withdrawal. A more than 5-fold increase in the proliferative cellular marker, Ki-67, appeared in

the zF starting at 1-week post-withdrawal and continued throughout the recovery period (Fig.

S2D). Taken together, these data demonstrate that adrenals withdrawn from long-term DEX

recover primarily by proliferation of the zF rather than cellular hypertrophy or alterations in zG

compartment size, making the pan-cortical area a reasonable surrogate for zF cell counts.

Moreover, GHOSTs did not contain the mRNA encoding the CORT biosynthetic enzyme

Cyp11b1, suggesting that GHOSTs are not capable of steroidogenisis (Fig. S2E).

Following the above FFPE studies, we then performed morphometric analyses of whole,

equatorial adrenal slices from DN, 8+0, 8+1, 8+2, 8+6, and 8+8 animals to quantify

steroidogenic cell type loss and recovery after long-term DEX treatment (Fig. 5). 488-

streptavidin[31] (green) and IF for TH (red), both as above, were used to label steroidogenic vs.

medullary cells, respectively. Since sections were not treated with TrueBlack, GHOSTs were

labeled by endogenous autofluorescence (cyan; Fig. 5A). Pan-cortical area, which includes both

streptavidin-positive and autofluorescent regions, decreased significantly with DEX treatment

(480 ± 7 mm^2 in 8+0 vs. 1290 ± 156 mm^2 in DN, $p<0.01$) but normalized by 1-week post-

withdrawal (1096 ± 65 mm^2 in 8+1, Fig. 5A and B). Autofluorescent GHOST area increased

significantly with DEX treatment and remained stably elevated through 8 weeks post-withdrawal,

with fragmentation and centripetal migration at later timepoints (Fig. 5A and C). Steroidogenic

cell type area was computed by subtracting GHOSTs (cyan autofluorescence) from the pan-

cortical (green) area, which markedly decreased with DEX treatment (272 ± 47 mm^2 in 8+0 vs.

1286 ± 153 mm^2 in DN, $p<0.001$) but more than tripled in the first week post-withdrawal

(887 ± 70mm^2; Fig. 5D). Medullary (TH-positive) area was comparable at all timepoints, indicating

that relatively equivalent, equatorial adrenal sections were used for morphometric analyses

($p=0.15$, Fig. 5E).

We next assessed the relationship between maximal CORT secretion to cosyntropin

stimulation and the size of the steroidogenic cell population in the recovering adrenal. This was

first measured as the ratio of stimulated CORT secretion to the steroidogenic area normalized to

this same ratio for DN controls (Fig. 5F). This ratio insignificantly decreased at the completion of

DEX treatment (from 1.0 ± 0.12 in DN to 0.62 ± 0.11 in 8+0, $p=0.08$) but was surprisingly lowest at

1- and 2- weeks post-DEX (0.46 ± 0.036 and 0.42 ± 0.010; $p<0.05$; Fig. 5F). Conversely, this ratio

of stimulated CORT secretion to steroidogenic area was significantly higher at 8 weeks post-

DEX (1.8 ± 0.11, $p<0.01$) vs. DN.

To examine this relationship another way, we conducted a linear regression analysis of

steroidogenic cell type area (independent variable) vs. cosyntropin-stimulated CORT

(dependent variable) at these same timepoints. This yielded a regression model of $Y =$

$0.016X - 3.9$ (R^2=0.43, p=0.0012; Fig. 5G). As visualized by an actual vs. predicted residual

plot, measured cosyntropin stimulated CORT levels were >50% lower at 1-2 weeks post-

withdrawal than predicted by the steroidogenic cell area at these same timepoints (5.0 vs. 10

mcg/dL at 8+1 and 5.0 vs. 11 mcg/dL at 8+2, Fig. 5H). Measured stimulated CORT secretion

roughly matched and then exceeded the predicted values 6 and 8-weeks post-withdrawal,

respectively (11 vs. 12 mcg/dL at 8+6 and 25 vs. 13 mcg/dL at 8+8). Overall, these analyses

demonstrate that CORT secretion is disproportionately low relative to the size of the recovering

non-histiocyte, adrenocortical cell compartment in the first two weeks after long-term DEX

treatment. As both basal and stimulated ACTH were elevated at these same time points (Fig. 2C

and E), this is consistent with adrenocortical atrophy with superimposed, secretory hypo-

responsiveness to ACTH.

Maintenance of endogenous ACTH secretion but not intermittent cosyntropin administration

during the period of DEX exposures prevents GIAI

After identifying the adrenal as the primary anatomical site of post-withdrawal HPA axis

dysfunction, we investigated whether GIAI could be prevented by providing trophic stimulation to

the gland during DEX exposure, a period when hypothalamic-pituitary function is otherwise

suppressed (Fig. 2A, B, C, and E, 8+0 timepoint). We first did this pharmacologically by

administering daily, intraperitoneal injections of cosyntropin, a synthetic ACTH analog (1-24) at

16:30, which coincides with the physiologic, circadian ACTH peak in rodents[33]. Mice were

concurrently treated with DEX via drinking water, as above (Model 2, Fig. 1). Following 8 weeks

of combined DEX+COS treatment, both agents were withdrawn, and adrenal recovery was

assessed with serial cosyntropin-stimulation testing at 1–2-week intervals until 8 weeks after

treatment cessation. At the time of treatment withdrawal (8+0), stimulated CORT secretion was

as low in DEX+COS-treated (2.0±1.0 mcg/dL) vs. DEX-treated animals (2.0±0.61 mcg/dL, 0.99).

Post-withdrawal, stimulated CORT secretion recovered more slowly in mice treated with

DEX+COS vs. DEX alone, with doubling times of 5.2 weeks ($Y = 3.8 * e^{0.13X}$; r^2=0.4) vs. 2.5

weeks ($2.6 * e^{0.28X}$, r_2=0.9), respectively, in exponential growth models (p=0.0009 for comparison

of fits by extra sum-of-squares F test; Fig. 6A).

Next, we tested whether maintenance of endogenous ACTH secretion, unlike intermittent,

pharmacologic administration, prevented DEX-induced adrenal suppression (Model 3, Fig. 1).

To do this, we bred Sim1[Cre]:GR[fl/fl] mice (HGRKO), which have selective, hypothalamic deletion

of the glucocorticoid receptor (GR), leading to high basal Crh, ACTH, and CORT levels, the

latter of which are not diminished by an overnight dexamethasone-suppression test[34]. By

immunofluorescent analysis, GR was absent in 77% of PVH cells in HGRKO mice vs. 6% in

their Cre-negative (HGRWT) or functionally WT littermates (p<0.001, Fig. 6B and D). As

expected, PVH *Crh* mRNA expression was significantly higher in DEX-treated (8+0) HGRKO

(244±2.2 arbitrary units [AU]) vs. HGRWT animals (49±12 AU, p<0.01, Fig. 6C and E). Basal,

pre-treatment ACTH levels were higher in HGRKO (470±76 pg/mL) vs. HGRWT mice (104±55

pg/mL, p<0.05). At the completion of 8 weeks DEX treatment, ACTH levels were undetectable in

HGRWT animals (0±0 pg/mL) but preserved in HGRKO mice (385±64 pg/mL, p<0.01, Fig. 6F).

In the presence of this ongoing ACTH stimulation, adrenal weights were significantly higher

at the completion of DEX treatment in HGRKO (2.3±0.2 mg) vs. HGRWT animals (1.1±0.06 mg,

p≤ 0.001, Fig. 6G). Prior to DEX treatment, basal and stimulated CORT levels, via both a

psychological stressor (20 minutes physical restraint) and cosyntropin-stimulation testing, were

higher in HGRKO compared to HGRWT animals (Fig. 6H). Importantly, basal and stimulated

CORT levels on long-term DEX were markedly suppressed in HGRWT mice but robust in

HGRKO animals (basal: 1.5±0.98 in HGRWT vs. 14±1.6 mcg/dL in HGRKO, p<0.01; restraint

1.9±1.6 vs. 33±4.5 mcg/dL, p<0.01; cosyntropin-stimulated 1.1±0.62 vs. 33±3.9 mcg/dL,

p<0.001).

Cosyntropin-stimulated CORT levels were then directly compared for all 8+0 animals in

models 1, 2, and 3 (Fig. 1) to assess the efficacy of these pharmacologic or genetic strategies

for preventing GIA (Fig. 6I). At the time of DEX±COS withdrawal, CORT secretion was

suppressed in all treatment groups (WT DEX 8+0: 2.0±0.6; WT DEX+COS 8+0: 2.0±1.0;

HGRWT DEX 8+0 1.1±0.62 mcg/dL, p<0.01 for all vs. DN) except HGRKO mice (33±3.9

mcg/dL), which was significantly higher than DN controls (p <0.01).Thus, long-term DEX does

not suppress the HPA axis of HGRKO mice but does so in WT mice, including those co-treated

with intermittent cosyntropin.

Short-term DEX-induced adrenocortical apoptosis is attenuated in cosyntropin-treated mice and

absent in HGRKO mice, but only HGRKO mice have preserved adrenal mass and morphology

on long-term DEX

We next assessed whether our pharmacologic or genetic strategies protected mice against

DEX-induced adrenocortical apoptosis and atrophy. TUNEL staining was performed on adrenals

from WT animals treated for 4 days with either vehicle (DN), DEX, or DEX+COS as well as from

HGRKO animals treated for 4 days with DEX (Fig. 7A, B). This short-term timepoint permitted

investigation of active, DEX-induced adrenocortical apoptosis prior to major zF tissue loss. As

expected, 4 days of DEX induced significant adrenocortical apoptosis (TUNEL positive

cells=11±0.7% vs. 0.63±0.19% for DN, p<0.001). There were significantly fewer TUNEL-positive

cells in glands treated with DEX+COS (5.7±1.7%) vs. DEX alone (11±0.7%, p<0.05), albeit with

a bimodal distribution in which apoptosis was reduced by >65% in n=4 animals but comparable

to DEX-treated glands in the remaining n=3. TUNEL cells were uniformly fewer in HGRKO

animals (1.7±0.90%) such that there was no significant difference between these glands vs. DN

controls (p=0.97).

Adrenal morphology after long-term DEX treatment in either the pharmacologic (Model 2,

Fig. 1) or genetic (Model 3, Fig. 1) paradigms was then examined using thick, floating section

immunofluorescence for TH with 488-streptavidin, (as in Fig. 5), in WT DEX+COS 8+1, HGRWT

8+0, and HGRKO 8+0 glands (Fig. 7C). Pan-cortical, GHOST, and steroidogenic cell type areas

were all measured by morphometric analysis and normalized to DN controls from Model 1 (DN

data from Fig. 5B-D). Pan-cortical area decreased in WT DEX+COS 8+1 (0.68±0.15) but not

HGRKO 8+0 adrenals (1.3±0.09, p=0.02; Fig. 7D). Autofluorescent or GHOST area increased

>20x in 8+1 (22±6.4) yet was negligible HGRKO glands (2.8±2.3; Fig. 7E). Steroidogenic cell

type area was 62% (0.62+0.17) vs. 99% (0.99±0.02) of DN controls in WT DEX+COS 8+1 vs.

HGRKO 8+0 glands, respectively (Fig. 7F). Of note, 8+1 glands were heterogenous in their

cortical size and distribution of GHOSTs, such that one animal had a normal steroidogenic cell

type area with small GHOSTs interspersed throughout the zF (Fig. S3). HGRKO glands, on the

other hand, were uniformly normal in size and lacked GHOSTs. Our genetic but not

pharmacologic strategy of delivering trophic stimulation to the adrenal thus consistently prevents

short-term, DEX-induced adrenocortical apoptosis and preserves glandular size and

morphology on long-term treatment.

DISCUSSION

In this study, we investigated the timing of hypothalamic, pituitary, and adrenal recovery in

mice after 8 weeks of treatment with DEX to probe the mechanisms of protracted GIAI after

discontinuation of exogenous glucocorticoid therapy. Post-withdrawal central suppression of

hypothalamic-pituitary function is generally thought to be the limiting factor in the recovery of

adrenal function, and that once this occurs, adrenal function rapidly normalizes in response to

hypothalamic-pituitary stimulation. We found that hypothalamic-pituitary dysfunction recovers

rapidly after cessation of long-term, exogenous glucocorticoid treatment, whereas adrenal

function remains subnormal for a much longer period of time despite hypothalamic-pituitary

stimulation. We also found that long-term DEX treatment induces marked adrenal atrophy with

accumulation of giant, CD68- and F4/80-positive histiocyte clusters (GHOSTs), which persist

beyond recovery of adrenal function. However, even after accounting for the likely inability of

GHOSTs to produce glucocorticoids, CORT secretion is disproportionately low relative to the

remaining adrenocortical cell mass for at least 2 weeks post-DEX withdrawal despite

supraphysiologic ACTH stimulation during this time.

After we established the adrenal as the primary site of post-withdrawal HPA axis

dysfunction, we then asked whether trophic stimulation of the gland during the period of DEX

exposure can prevent GIAI, using both genetic and pharmacologic approaches. Mice with

feedback-resistant, sustained ACTH secretion due to deletion of the glucocorticoid receptor

gene (Gr, *Nr3c1*) from hypothalamic neurons (HGRKO) maintained normal adrenal function and

morphology throughout the entire period of long-term DEX treatment, while WT animals co-

treated with DEX and cosyntropin (DEX+COS) for the same amount of time had even slower

CORT recovery than those treated with DEX alone. In our short-term studies, DEX-induced

adrenocortical apoptosis was attenuated in both groups of mice, although this rescue was more

robust in DEX-treated HGRKO than in DEX+COS animals.

Our finding of early hypothalamic-pituitary recovery aligns with previous rodent studies

showing normal plasma ACTH and/or corticotroph *Pomc* mRNA levels within 1 week of

discontinuing 5-30 days of methylprednisolone or dexamethasone in mice and rats[22-24]. Ours is

the first preclinical study to examine HPA axis recovery after long term glucocorticoid therapy,

which is typically defined as greater than four weeks of supraphysiologic exposure and is most

strongly associated with GIAI development in humans[13]. Our model successfully reproduced

prolonged, severe CORT deficiency but lacked the initial period of hypothalamic-pituitary

dysfunction reported in prior clinical studies[16]. Of note, we did not evaluate hypothalamic-

pituitary status in the first week after steroid withdrawal, as any suppression at this time could

have been due to DEX-mediated negative feedback rather than to GIAI, given the long half-life

of the drug. Nonetheless, the lack of clinically significant, prolonged hypothalamic-pituitary

dysfunction in our model warrants investigation of the differences between the rodent and

human HPA axes as well as re-evaluation of the methodologies used to test for post-withdrawal

central suppression.

The 1965 Graber and Liddle study included only 6 patients with hypercortisolism from

exogenous glucocorticoid treatment who were studied after completing an undefined steroid

taper. Most importantly, ACTH levels were unable to be directly measured and were instead

derived from the CORT responses of hypophysectomized rats injected with extracted patient

plasma versus three standard doses of ACTH or saline[16,35]. During long-term prednisone

treatment in humans, ACTH responses to CRH stimulation and insulin-induced hypoglycemia

are suppressed. However, all testing in Graber et al. was performed 24 hours (~1-2 biological

half-lives) after prednisone administration, thus limiting the ability of the authors to distinguish

between active, glucocorticoid feedback inhibition vs. post-withdrawal central suppression[13,36].

In favor of early hypothalamic-pituitary recovery after steroid-induced adrenal suppression in

humans, patients with congenital adrenal hyperplasia, who also require chronic, suppressive

glucocorticoid therapy, have elevated levels of both ACTH and ACTH-dependent steroid

precursors just 12 hours (~3-4 have-lives) following a hydrocortisone dose[37,38]. The

discrepancies and methodologic limitations of the existing clinical literature, combined with our

finding of early, post-withdrawal hypothalamic-pituitary recovery in mice, highlight a need to

revisit this question with modern ACTH assays and larger numbers of glucocorticoid-treated and

-withdrawn patients .

Re-evaluating the paradigm of post-withdrawal central suppression has important

implications for defining the mechanisms underlying GIAI and developing effective preventative

strategies. First, if both basal and stress-induced ACTH secretion are rapidly restored after

discontinuing long-term glucocorticoids in humans, the latency in adrenal recovery that follows

reflects a period of ACTH hypo-responsiveness. This would be most likely presumed secondary

to adrenocortical atrophy given the well-established role of ACTH in preventing zF apoptosis

and maintaining proper adrenal zonation[9,10]. While we indeed found that DEX-treatment led to

adrenocortical apoptosis followed by major zF atrophy, adrenal weights recovered quickly

compared to CORT secretion after DEX was withdrawn. This aligns with prior preclinical studies

in which rodents had normal adrenal weights but low plasma CORT and/or adrenal

steroidogenic pathway gene expression within a week of discontinuing short-term (5-7 days)

glucocorticoid treatment[22-24]. With our finding of large, CD68+ and F4/80+ histiocyte clusters in

DEX-exposed glands, adrenal weights may overestimate functional cortical regeneration after

long-term glucocorticoid treatment. However, even after excluding this myeloid compartment by

morphometric analysis, recovery of CORT secretion lagged behind that of adrenocortical mass

after DEX withdrawal. As a potential limitation, this finding is based on 2-dimensional (2D) areas

as a surrogate for 3-dimensional (3D) cell counts. We established that zF cellular density was

similar between DEX-withdrawn and untreated control mice at all examined post-withdrawal

time points, making adrenocortical area a reasonable proxy for cellularity. Moreover, 2D area

overestimates 3D volume early in recovery when the glandular radius is reduced[39]. Our

conclusion that CORT secretion is disproportionately low relative to the adrenocortical cell mass

in the first 2 weeks after DEX withdrawal is thus conservative and warrants future investigation

into the mechanisms of ACTH hypo-responsiveness in early, post-withdrawal GIAI.

We have found giant, multinuclear histiocyte clusters after long-term glucocorticoid

treatment. Using electron microscopy, Wyllie and Kerr observed apoptotic cell bodies within

histiocyte phagosomes in rat adrenals after two days of prednisolone treatment[10]. Large, CD68-

and F4/80-positive histiocytes resembling GHOSTs were reported in aging male Znrf3 knockout

mice, which exhibit cortical hyperplasia followed by senescence[40]. In our studies, numerous

mononuclear macrophages were observed throughout the zF by 1-week of DEX, which

coincided with massive adrenocortical apoptosis by TUNEL staining. With ongoing DEX

treatment, these cells appeared to aggregate into giant clusters at the corticomedullary junction.

GHOSTs resembled multinucleated giant cells by both H&E and CD68 staining, but IF with the

membrane-bound macrophage marker F4/80 revealed a morphologic pattern consistent with

fusion of multiple, smaller multinucleated histiocytes. Interestingly, multinucleation and fusion of

phagocytic cells has been described in adipose tissue, ovary, and bone when demand for

cellular clearance is high[41-43]. We hypothesize that the massive adrenocortical apoptosis that

accompanies corticosteroid treatment leads to recruitment and/or expansion of an adrenal

macrophage population, which subsequently undergo fusion and multinucleation, perhaps for

enhanced phagocytic capacity.

Notably, GHOSTs remained abundant in the early weeks after DEX treatment when the

hypothalamus and pituitary had recovered but adrenal secretory capacity remained

disproportionally low for cortical size, raising the possibility that these histiocytes may disrupt

steroidogenesis. Adrenal macrophages have been reported to restrict CORT production via

paracrine signaling after undergoing activation in response to acute and chronic stress[44].

Specifically, intra-adrenal cortisol upregulates triggering receptor on myeloid cells 2 (Trem2) in

adrenal macrophages in a GR-dependent manner[44]. This promotes TGFβ secretion and

ultimately downregulation of steroidogenic acute regulatory protein (stAR) in adrenocortical

cells, which is the rate limiting enzyme in CORT biosynthesis[44-46]. In our studies, rising ACTH

and thus intra-adrenal CORT after DEX withdrawal may activate an expanded adrenal

macrophage population, providing one potential mechanism of secretory hypo-responsiveness

to ACTH after long-term glucocorticoid treatment. Future studies will further characterize the

origin, phagocytic, and regulatory functions of this macrophage population and investigate their

contribution to prolonged GIAI.

Most importantly, refocusing the mechanisms of post-withdrawal GIAI to the adrenal, rather

than the hypothalamus or pituitary, suggests that we may be able to prevent prolonged

suppression by providing effective, trophic stimulation to the gland during the period of

glucocorticoid exposure. In this study, we were able to preserve adrenal function and

morphology with a genetic approach that provided constant, CRH-driven, endogenous ACTH

stimulation during long-term glucocorticoid therapy but not with daily, intraperitoneal cosyntropin

injections. There are several key differences between these two strategies that should be

considered in subsequent attempts to develop an effective, preventative pharmacologic

treatment. First, we selected cosyntropin (ACTH 1-24) rather than purified ACTH (1-39) for initial

long-term studies, as the truncated peptide is thought to be as biologically potent but less

antigenic than the full-length hormone[47,48]. While cosyntropin is widely used diagnostically to

acutely stimulate steroidogenesis, its trophic effects are largely unstudied[49].

ACTH is also likely constantly elevated in the HGRKO model vs. intermittent in our

pharmacologic model. Cosyntropin, like native ACTH, has a biologic half-life of only 10-30

minutes[47,48]. We had hoped to overcome this by by achieving an initial systemic ACTH activity

>200x greater than that of physiologic ACTH at the circadian peak or in response to stress, akin

to what is used for high-dose stimulation testing in humans[50,51]. Cosyntropin treatment statistically

significantly reduced short-term, DEX-induced adrenocortical apoptosis, suggesting that we

successfully delivered biologically active cosyntropin to the gland. Interestingly, we observed a

heterogenous effect of cosyntropin on both short-term DEX-induced adrenocortical apoptosis and

long-term zF size and morphology, whereas CORT secretion was uniformly suppressed after

long-term DEX+COS treatment. Future studies will investigate whether different properties of

ACTH exposure are necessary to maintain the steroidogenic machinery vs. promote zF growth

and/or survival. The failure of intermittent cosyntropin to prevent GIAI may be due to an

inadequate duration of ACTH exposure, as adrenal *Mc2r* mRNA is increased after 6-48h of ACTH

treatment but not within an hour of ACTH injection or 4 hours of physicial restraint[52-56].

Alternatively, ultradian pulsatility may be necessary for the full biologic activity of ACTH, as it is

for GnRH and PTH[57,58]. Consistent with our interventions using CRH vs. ACTH to prevent GIAI in

mice, in patients with Cushing syndrome due to either ectopic CRH or ACTH secretion, two with

ectopic CRH section had rapid recovery of HPA axis function after tumor removal without

perceptible HPA axis suppression, whereas a third patient with ectopic ACTH secretion had

prolonged adrenal suppression after tumor resection[59]. Future studies will explore the

mechanisms by which activation of the endogenous HPA axis but not intermittent cosyntropin

administration preventions GIAI as a prelude to the development of future, preventative

pharmacologic strategies.

As a potential limitiation, our sample sizes were relatively small to enable a broad, agnostic

survey of HPA axis recovery across multiple timepoints. Most analyses were still adequately

powered given relative homogeneity among experimental animals, particularly when hormonal

function was interrogated with provocative tasting rather than basal measurements, as is done in

clinical endocrine practice. Nonetheless, subsequent mechanistic and interventional studies will

include larger cohorts of animals now that key, biologically relevant timepoints have been

identified. Similarly, the present study included only male mice to minimize biological variability

across this broad, initially exploratory survey of HPA axis recovery. While there are no reported

sex differences in the risk, severity, or duration of GIAI in humans[12,19,60,61], female rodents have

have more responsive HPA axes to stress as well as more rapid adrenocortical turnover, the latter

of which uniquely involes a capsular stem cell pool[62-67]. Warde et al. further reported more robust

adrenal CD68+ myeloid cell accumulation following cellular senescence in male vs. female Znrf3

knockout mice, which raises the possibility that the presence and functional consequences of

adrenal macrophages could be sexually dimorphic in our model[40,68]. Future studies will thus

specificially focus on sex-differences in the mechanisms of post-withdrawal GIAI and time to axis

recovery.

In conclusion, we have developed a physiological, preclinical model of GIAI in which long-term

glucocorticoids induce prolonged, adrenocortical dysfuntion in excess of glandular atrophy

despite rapid restoration of hypothalamic-pituitary function. Trophic stimulation of the gland during

the period of glucocorticoid exposure by nonsuppressible activation of the HPA axis, but not by

intermittent cosyntropin administration, prevents DEX-induced adrenal suppression. Future

studies will examine the mechanisms underlying adrenal hyporesponsiveness to ACTH in the

initial weeks after steroid withdrawal as well as those by which these two modes of adrenal

stimulation differentially protect against GIAI. This work may collectively guide the development

of novel therapies that prevent GIAI, thus decreasing the morbidity of one of our most widely

prescribed drug classes without sacrificing therapeutic efficacy.

MATERIALS AND METHODS

Sex as a biological variable

Our study exclusively examined male mice for reasons discussed (see Discussion above) and

to minimize biologic variability and thus adequately power a broad survey of HPA axis recovery

across multiple timepoints. Is it unknown whether the findings are relevant for female mice.

Animals

All animals protocols were in accordance with the National Institutes of Health guidelines and

were approved by the Animal Care and Use Committee of Boston Children's Hospital (Protocol

00001983). Mice were housed on a 12-hour/12-hour light-dark cycle (lights on at 7 am) with ad

libitum access to rodent chow and water unless otherwise indicated. Mice used for studies were

all aged either 8 weeks at the time of DEX initiation (long-term studies) or 8-16 weeks (DEX-naïve

controls or short-term studies).

Wild-type (WT) mice: C57BL/6J mice (The Jackson Laboratory, Bar Harbor, ME) were

maintained by breeding within our population.

Sim1^Cre:GR^fl/fl mice (HGRKO): Sim1[Cre] mice (generous gift of Brad Lowell, Beth Israel Lahey

Medical Center, Boston, MA) were bred to floxed GR exon 3 mice (B6.Cg-*Nr3c1^tm1.1Jda*/J, Catalog

#021021, The Jackson Laboratory, Bar Harbor, ME), both maintained on an inbred C57BL/6J

backround[34,69]. Littermates that were Cre negative but had GR exon 3 floxed alleles were used

as controls. Animals were genotyped using tail DNA extracted with 0.05 M NaOH, heated at 95°C

for 30 min and then neutralized with Tris-HCl (pH 7.4). All genotyping protocols used a 2x Taq

RED Master Mix, 1.5 mM MgCl$_2$ (Apex, 5200300-1250) with 10 μM diluted primers (IDT). Primers

and PCR protocols for each genotype are detailed in Table SI.

Pharmacologic treatments: Mice were treated with either water-soluble dexamethasone

(Sigma-Aldrich 2915) at a concentration of 3.6 mcg/mL (~20 mcg/day assuming an average water

intake of 5.8 mL/day[70]) or no-drug control via drinking water for either 4 days (short-term) or 8

weeks (long-term studies). When specified, mice were co-treated with 10 mcg/kg cosyntropin,

(synthetic ACTH 1-24, Sandoz, NDC 0781-3440-95), dissolved at 10 mcg/mL in 0.9% saline.

Cosyntropin was administered once daily by intraperitoneal injection at 16:30 p.m. to align with

the murine, circadian ACTH peak, for either 4 days or 8 weeks, as above.

Hormonal assays: All testing was performed, as specified, on separate days so that no

mouse was bled more than once on a given day. Plasma for measurement of corticosterone and

ACTH was obtained by rapid retroorbital phlebotomy without anesthesia into EDTA-coated

capillary tubes with a total time from first handling the animal to completion of bleeding not

exceeding 30 seconds. Blood was collected on ice, and plasma was separated by centrifugation

at 1500 RPM at 4°C and stored at -80°C until assay. Plasma concentrations of corticosterone

(ALPCO, 55-CORMS-E01) and ACTH (21-ACTHU-E01) were measured by ELISA per

manufaturer instructions. For the ACTH assay, plasma was diluted 1:10 with ACTH 0 calibrator

(ALPCO; 11-C-270-10-G25). The lower limit of detection is 0.061 mcg/dL and 0.22 pg/mL for the

CORT and ACTH assays, respectively.

• *Basal measurements:* Animals underwent rapid retroorbital phlebotomy, as above, at

16:30 in a ventilated hood in the same room as which the animals were housed.

Plasma ACTH and corticosterone were measured by ELISA as above.

• *Cosyntropin-stimulation testing:* Cosyntropin stimulation testing was performed 2

hours after lights on in ad libitum fed mice. Mice were bled by retroorbital phlebotomy

60 min after intraperiteoneal inection of 10 mcg/kg cosyntropin (10 mcg/mL in 0.9%

saline). Plasma CORT was measured by ELISA, as above[33].

• *Insulin-induced hypoglyemia:* Mice were deprived of food at lights off. The next

morning, within 2 hours of lights on, mice were intraperitoneally injected with

recombinant regular human insulin (Eli Lilly) at a dose of 1 u/kg, which was estimated

from previously published data to produce sustained hypoglycemia and consistent

pituitary-adrenal activation at 30-60 min after injection[71,72]. Mice were bled via

retroorbital phlebotomy for plasma hormone analysis (both ACTH and CORT) 40

minutes after injection[73]. Tail vein blood glucose was measured with a point-of-care

glucometer (FreeStyle Lite, Abbott) 60 minutes after insulin injection to document a

blood glucose nadir of <60 mg/dL, which occurred in all experimental animals.

• *Physical restraint:* Animals were confined in ventilated 50-mL conical tubes for 20

minutes followed by rapid retroorbital phlebotomy for plasma corticosterone

measurement.

Necroscopy and tissue preparation: Animals were euthanized by rapid decapitation without

anesthesia followed by necroscopy for isolation of hypothalami and adrenals. Coronal slabs

flanking the hypothalamus were prepared with a surgical blade and then fixed in 10% formalin

overnight before transfer to 70% ethanol while awaiting paraffin embedding. Adrenals were

trimmed of surrounding fat tissue, rinsed in phosphate-buffered saline (PBS), and weighed

before overnight fixation in 10% neutral buffered formalin.

Single-molecule RNA in situ hybridization

Three 5mM-thick coronal brain sections were mounted onto each slide, and every other

slide was stained with hematoxylin and visualized with brightfield microscopy (Nikon E800). The

paraventricular nucleus was identified histologically based on its characteristic "butterfly-

shaped" appearance in coronal sections, which was used to anchor selection of 2 unstained

slides immediately rostral and caudal to this structure for analysis. Single-molecule RNA in situ

hybridization was performed using a catalog Mus musculus *Crh* probe (ACD #316091) and 2.5

HD RED detection kit (ACD #316091) following manufacturer's protocol. Slides were counter

stained with 50% Gill's Hematoxylin II and mounted with EcoMount. Positive control *Ppib (*ACD

#313911) and negative control *dapB (*ACD #310043) were used in every experiment to ensure

sample quality. Brightfield images at 40x magnification were generated with an Olympus VS120

Virtual Slide Scanner and then analyzed with semi-quantitative, densitometry using the Weka

Segmentation plugin for Fiji ImageJ per ACDBio instructions[74]. Analysis was restricted to free-

handed sections encompassing the characteristic "butterfly" morphology of PVH cell bodies, and

CRH expression, reported as arbitrary units (AU) was quantified by diving the area of the *Crh*

densitometric signal (μm^2) by the total area of this region (mm^2)

Total mRNA and protein staining

For quantifying total mRNA *in situ*, FFPE sections were deparaffinized and antigen-retrieved

in buffer TE (pH=9). We then adapted the protocol in Farny *et al.* and Hurt *et al.* to label polyA-

capped RNAs on FFPE sections using Cy5-conjugated polyDT (26-4430-02, GeneLink)[26,27].

Total protein *in situ* was measured using the N-Hydroxysuccinimide (NHS) ester method; NHS

esters react with primary amines and well-approximate total protein mass[28,29]. Briefly, FFPE

sections were deparaffinized and antigen-retrieved in buffer TE (pH=9). Slides were stained with

NHS-Alexa 568 dye conjugates (Thermo Scientific A20003) in PBS for 30 minutes and then

washed 5X in PBS.

Paraffin section immunofluorescence

After fixation, adrenals were oriented with their broadest surface parallel to the cutting plane

in 4% low-melting-temperature SeaPlaque agarose (Lonza, #50101). Agarose blocks were then

dehydrated in ethanol, xylene, and embedded in paraffin blocks. Paraffin sections were cut at 5

µm thickness. Sections were rinsed in xylene, an ethanol gradient, and then PBS. Antigen retrieval

was performed in either 10 mM Sodium Citrate pH 6.0 or Tris-EDTA pH 9.0 (Table SII). Sections

were blocked in 5% Normal Donkey Serum (NDS), 0.1% Tween-20 in PBS for 2h at RT. Slides

were then incubated in primary anitbodies diluted as below (Table SII) in 5% NDS at 4°C

overnight. Slides were washed three times for 5 min in 0.1% Tween-20 in PBS. Sections were

incubated for 2h at RT in secondary antibodies diluted 1:200 in PBS. For cortical counterstaining

in the CD68 experiment, Streptavidin, Alexa Fluor 488 Conjugate (Invitrogen, #S112233) was

added to the secondary antibody solution at a final concentration of 1:500. After three 5-min

washes with 0.1% Tween-20 in PBS, sections were incubated in TrueBlack Lipofuscin

Autofluorescence Quencher (Biotium, #23007) in 70% EtOH at RT for 3 minutes. Slides were

rinsed thee times in PBS and then mounted with Prolong Gold Antifade Mountant with DAPI

(Invitrogen, #P36931). For CD68 IF, Images were acquired with a Nikon Ti2 microscope with a

Yokogawa CSUW1 confocal unit, a Nikon Plan Apo Lambda 20x objective lens and a Zyla 4.2

PLUS sCMOS camera (ANDOR). All images were taken with identical laser strength, gain and

exposure settings for the 488nM and 647nM channels. For all other experiments, fluorescent

images were obtained with an Olympus VS120 Virtual Slide Microscope using a 20x objective.

Brightness and contrast were adjusted for all images in ImageJ.

Floating section immunofluorescence.

After fixation, adrenals were oriented with their broadest surface parallel to the cutting plane

in 4% low-melting-temperature SeaPlaque agarose (Lonza, #50101) and sectioned at a

thickness of 100 μm with a vibratome. Individual slices were separated by free floating in PBS.

Sections were wet-mounted in PBS on Superfrost Plus slides (Fisher Scientific, 12-550-15) and

imaged with a 4x objective on a Nikon E800 brightfield microscope. The area of each section

was measured in ImageJ, and the largest section was selected for subsequent immunostaining.

Slices were washed three times with 0.1% Tween-20 in PBS and blocked in 5% normal donkey

serum, 1% bovine serum albumin, and 0.1% Tween-20 in PBS for 1h at RT with gentle rocking.

Slices were incubated with Sheep anti-tyrosine hydroxylase (1:500; Sigma Aldrich, #AB1542)

diluted in blocking solution at 4°C overnight. After three 20 min washes with 0.1% Tween-20 in

PBS, slices were incubated with Alexa Fluor 647-conjugated donkey anti-sheep IgG (1:100,

Invitrogen, #A-21448) and Streptavidin, Alexa Fluor 488 Conjugate (1:500, Invitrogen,

#S112233) diluted in 0.1% Tween-20 in PBS for 2h at RT. For nuclear staining, DAPI (4'6-

diamino-2-phenylindole) was added to secondary antibody mix for the last 5 min at a final

concentration of 1:1000. Slices were washed with 0.1% Tween-20 in PBS for 20 min three times

and then directly mounted on Superfrost Plus Slides with Prolong Gold Antifade Mountant

(thermo Fisher Scientific, #P36930). Images were acquired with a Nikon Ti2 microscope with a

Yokogawa CSUW1 confocal unit, a Nikon Plan Apo Lambda 20x objective lens and a Zyla 4.2

PLUS sCMOS camera (ANDOR). All images were taken with identical laser strength, gain and

exposure settings for the 405nm (used to capture autofluorescence), 488nm, and 647nm

channels. Each image was acquired using the Large Image function in the NIS elements

software. This function allowed us to take images in a 4x4 grid with 15% overlap, allowing the

images to be stitched together. Brightness and contrast were adjusted for all images in ImageJ.

Image analysis

Fluorescent images were analyzed using QuPath (version 0.4.3), an open-source software

for digital pathology.

Positive Cell Detection

QuPath's cell detection feature was employed to label cells as either Ki-67 or TUNEL positive

based on co-localization of the 647 channel signal with the DAPI nuclear signal. Images were

opened in fluorescence mode, and regions of interest (ROI) were mannually defined to include

only the adrenal cortex based on morphologic characteristics. ROI were further restricted to the

zF for Ki-67 analysis, as defined by DAB2-negative cortical cells. The "cell detection" feature was

then run specifying DAPI as the detection channel and the 647 nm channel nuclear (Ki-67

analysis) or cellular (TUNEL analysis) mean as the score compartment. Nuclear size and intensity

threshold parameters for both were adjusted and inspected visually to ensure accurate positive

cell detection. Any cell exhibiting signal overlap between the DAPI and 647 nm channels above

the specified thresholds was classified as either TUNEL or Ki-67 positive and reported as a

percentage among total cells.

Morphometry

Morphometric analysis was performed on thick, floating section immunofluorescent

specimens to quantify the area of either the adrenal cortex (Strepatividin staining, 488 nm

channel), medulla (tyrosine hydroxylase immunostaining, 647 nm channel), or GHOSTs

(autofluorescence, 408 nm channel) based on signal intensity across multiple channels using the

"create thresholder" function in QuPath. Images were opened in fluorescence mode, and

brightness/contrast of individual channels were adjusted for optimal signal visualization. Regions

of interest were mannually defined to include the entire adrenal slice but not surrounding adipose

tissue. For each channel, QuPath's intensity thresholding was applied to ensure accurate

delineation of stained or autofluorescent areas, as confirmed by visual inspection. As DEX

treatment has been shown to attenuate streptavidin staining in the murine zF, we applied a lower

threshold in 8+0 adrenals to ensure detection of all residual adrenocortical tissue[31]. The area of

tissue with a signal above this threshold for each channel was classified as positive and reported

in square microns.

Statistics

Two-tailed Student's t test was performed to compare significance among two groups.

Comparisons done beyond 2 groups were performed by ANOVA, and Bonferroni correction was

applied for post-hoc, pairwise multiple compraison testing. Nonlinear regression using an

exponential growth equation was used to model stimulated CORT recovery (dependent variable)

over time (independent variable), and doubling time was computed as ln(2)/K where K is the rate

constant. When two exponential growth models were compared, the extra-sum-of squares test

was employed to determine whether the exponential equations provided significantly different fits

to the data. The relationship between steroidogenic area (independent variable) vs. stimulated

CORT (dependent variable) was modeled by simple linear regression. Goodness of fit for all

regression analyses was reported as R^2 values. *P* values below 0.05 were considered statistically

significant. All statistical analyses were performed in Graphpad Prism v.10.3.1. Windows,

GraphPad Software, Boston, Massachusetts USA, www.graphpad.com

Study approval

All animal experiments were approved by the Boston Children's Hospital Institutional Animal

Care and Use Committee (Protocol Number 00001983).

Data availability

All raw data associated with findings presented in this manuscript are included in the

Supporting Data Values file and are available from the corresponding author upon reasonable

request.

AUTHOR CONTRIBUTIONS

LG and JM conceived the project and designed experiments. LG, BJ, and HF conducted

experiments, and LG analyzed the data. LG, MS, and BJ acquired microscopic images. LG wrote

the manuscript, which was edited by JM and MS. All authors approved the final version of the

submitted manuscript.

ACKNOWLEDGEMENTS

This study was supported by NIH 5T32DK007699-39 (LG), NIH F32DK131795-02 (LG), a

Pediatric Endocrine Society Research Fellowship Award (LG), a Pediatric Endocrine Society

Clinical Scholar Award (LG), a Boston Children's Hospital Office of Faculty Development/Basic &

Clinical Translational Research Executive Committees Faculty Career Development Fellowship

(LG), and the Thomas Morgan Rotch Harvard Medical School endowment (JAM). We thank Drs.

David Breault, Kleiton Borges, and Mesut Berber for their input on this project and assistance

processing adrenal tissue, identifying zonal markers, and performing thick, floating section

immunofluorescence. We also thank Suzanne White from the BIDMC Histology core for preparing

and sectioning all FFPE specimens, including working closely with the author to isolate the PVH

in mouse brains.

 FIGURES AND LEGENDS

Figure 1. Experimental design for dexamethasone (DEX) ± cosyntropin (COS) treatment and withdrawal in wild-type (WT, Models 1 and 2) and HGRKO ($Sim1^{Cre}$:$GR^{fl/fl}$) (Model 3) mice. In model 1, the natural history of HPA axis recovery after long-term DEX was studied by treating WT mice with 8 weeks of DEX (red bar) followed by serial assessments of axis function and histology at 1-2 week intervals after DEX was withdrawn (blue bar). In cohort 1, basal, unstimulated ACTH and CORT were serially measured at the above timepoints (yellow dots). Subsequent cohorts of animals underwent provocative HPA axis assessment with cosyntropin stimulation and insulin-induced hypoglycemia testing on successive days followed by necroscopy at these same timepoints (yellow dots). Pharmacologic (model 2) and genetic strategies (model 3) were then studied for their ability to prevent GIAI. In model 2, mice were co-treated for 8 weeks with DEX and cosyntropin, an ACTH analog, followed by cosyntropin-stimulation testing to assess adrenal function after both agents were withdrawn (green dots). Necroscopy for histologic studies was performed in a separate cohort of mice 1-week post-DEX+COS (8+1; split yellow and green dot). In model 3, HPA axis activity was assessed upon completion of an 8-week DEX course (8+0) in mice with selective hypothalamic deletion of the glucocorticoid receptor (HGRKO), which allows for ongoing, endogenous ACTH secretion on GC therapy. ACTH and CORT were measured in the basal, unstimulated state followed by provocative assessments (restraint stress and cosyntropin-stimulation testing on successive days) and necroscopy[75].

Figure 2. Stimulated CORT secretion requires 8 weeks to normalize after long-term DEX treatment despite early hypothalamic-pituitary recovery. A) *In situ* hybridization for basal *Crh* (red) in the paraventricular nucleus (PVH) of the hypothalamus in DEX-naïve (DN) animals and those treated with 8 weeks DEX both immediately prior to (8+0) and 1-week post-withdrawal (8+1; top row, scale bar=200 microns). Magnification of region within the black box is shown directly below (bottom row, scale bar=50 microns). B) Quantification of *Crh* mRNA expression from RNAScope images of the PVH from DN, 8+0, and 8+1 animals (n=2). p<0.05 by 1-way ANOVA. C) Recovery of plasma levels of basal (unstimulated), late afternoon ACTH and D) corticosterone (CORT) as well as E) insulin-induced hypoglycemia-stimulated ACTH, F) insulin-induced hypoglycemia-stimulated CORT, and G) cosyntropin-stimulated CORT 0-8 weeks after withdrawal of an 8-week DEX course. Data shown as individual mice with mean (connecting line) ± SEM (error bars). Grey bar (mean± SEM) indicates the reference range for adult, male, wildtype, untreated (DEX-naïve or DN) controls (n=4-8). * denotes p<0.05, ** p<0.01, *** p<0.001, and ****p<0.0001 vs. DN.

Figure 3. 8 weeks DEX induces adrenocortical atrophy with accumulation of poorly defined, vacuolated, cellular structures that persist through 8 weeks post-withdrawal. A) DEX-treated adrenals weigh significantly less than DN controls through 6 weeks post-withdrawal. P<0.0001 by one-way ANOVA; * indicates p<0.05, ** p<0.01, *** p<0.001, and ****p<0.0001 vs. DN. B) Hematoxylin and eosin staining reveals large, poorly defined, vacuolated structures with decreased nuclear density at the corticomedullary junction in DEX-exposed adrenals that persist through 8-weeks post-withdrawal. Representative structures are outlined with red, dashed circles and C) magnified (black box). D) These accumulations are present in DEX-exposed (8+1, bottom) but not DN adrenals (top) and are intensely autofluorescent (green). Autofluorescence is eliminated with the addition of TrueBlack, a lipofuscin autofluorescence quencher. Like surrounding adrenocortical cells, nuclei within these clusters contain mature mRNA (*in situ* hybridization for total polyA with 5'-Cy5-Oligo d(T)30; magenta) and protein (568 NHS Ester staining; red; both with TrueBlack). All scale bars=100 microns (B-D).

Figure 4. Giant histiocyte clusters accumulate with prolonged glucocorticoid treatment and persist through at least 8-weeks post-withdrawal. A) Immunofluorescence for the myeloid immune marker CD68 (magenta) with DAPI (blue) and 488-streptavidin (green) counterstains, the latter of which labels all biotin-rich steroidogenic cells of the adrenal cortex. Representative images shown for DN adrenals, those on DEX for 2-8 weeks (e.g. 2+0 to 8+0), as well as one (8+1) and eight (8+8) weeks after an 8-week DEX course. B) Magnification of representative CD68- positive cluster from an 8+0 gland. C) Immunofluorescence for the mouse macrophage marker F4/80 (green) and medullary marker tyrosine hydroxylase (TH; magenta) in DN and DEX-exposed adrenals (2+0 and 8+0). D) Magnification of F4/80-positive cluster from an 8+0 gland. All staining performed with TrueBlack lipofuscin autofluorescence quencher (A-D). All scale bars = 100 microns.

Figure 5. Stimulated CORT secretion is disproportionately low relative to adrenocortical cell mass in the first 1-2 weeks after withdrawal of long-term DEX. A) IF without TrueBlack for tyrosine hydroxylase (TH, red) DAPI (blue) and 488-streptavidin (green; columns 1 and 2) counterstains. Giant Histiocytes of Steroid Treatment (GHOSTs) were detected by autofluorescence (cyan; columns 3 and 4). Mice were either untreated (DN) or treated for 8 weeks DEX followed by withdrawal for 0-8 weeks (8+0, 8+1, 8+2, 8+6, and 8+8). Shown as 100 uM-thick, whole adrenal slices (columns 1 and 3, scale bar=500 microns) with magnification of the region in the white box (columns 2 and 4, scale bar=100 microns). Columns 1 and 2 imaged at wavelengths 405 nm, 488 nm, and 647 nm; columns 3 and 4 imaged at wavelengths 405 nm, 440 nm, and 647 nm. Morphometric analysis of B) pan-cortical area, which includes both streptavidin-positive and autofluorescent regions (p=0.0017cby one-way ANOVA), C) Autofluorescent GHOST area (p=0.0053 by one-way ANOVA), D) steroidogenic cell type area, calculated as pan-cortical area minus GHOSTs (p=0.0011 by one-way ANOVA), and E) medullary (TH- positive) area (p=0.15 by one-way ANOVA). F) Ratio of steroidogenic cell type area to cosyntropin-stimulated CORT secretion, normalized to DN controls. * p<0.05, ** p<0.01,

810   *** p<0.001 vs. DN (B-F). G). Linear regression analysis of steroidogenic cell type area
811   (independent variable) vs. cosyntropin-stimulated CORT secretion (dependent variable;
812   R^2=0.43, p=0.0012). H) residual plot from G) of actual vs. predicted cosyntropin-stimulated
813   CORT.

Figure 6. Endogenous ACTH secretion but not intermittent cosyntropin administration preserves adrenal mass and secretory function on long-term DEX treatment. A) Recovery of cosyntropin-stimulated CORT secretion in WT mice withdrawn from 8 weeks DEX+COS (red) vs. DEX-monotherapy (black). Data shown as individual mice with error bars centered on mean value ± SEM. Gray bar (mean ± SEM) indicates the reference range for adult, male, wildtype, DEX-naïve controls. Stimulated CORT recovery was modeled with exponential functions yielding doubling times of 5.2 weeks after DEX+COS ($Y = 3.8 * e^{0.13X}$ R^2=0.4, red curve) vs. 2.5 weeks after DEX ($Y = 2.6 * e^{0.28X}$; R^2=0.9, black curve). p= 0.0009 by extra sum-of-squares F test. B) IF for the glucocorticoid receptor (GR, red; DAPI, blue) in HGRWT (left panel) and HGRKO (right panel). C) *in situ* hybridization for *Crh* mRNA in the PVH of long-term DEX-treated Cre-negative control HGRWT (left panel) and Sim1^Cre^:GR^fl/fl^ HGRKO (right panel) animals. D) Quantification of the percentage of GR-positive nuclei and E) *Crh* mRNA expression in the PVH of HGRWT vs. HGRKO animals. F) Basal (unstressed circadian peak) ACTH levels in HGRWT vs. HGRKO animals prior to DEX (left of dashed line) and on 8 weeks of DEX (right of dashed line). p=0.19 for DEX status and p<0.0001 for genotype by two-way ANOVA. G) Adrenal weights on 8 weeks of DEX in HGRWT vs. HGRKO animals. H) CORT secretion in the

basal state, following 20 minutes of physical restraint (Restraint), and following cosyntropin stimulation (COS-stim.) in HGRWT vs HGRKO animals, either prior to DEX treatment (left of dashed line) vs. on 8 weeks of DEX (right of dashed line). * $p<0.05$, ** $p<0.01$, *** $p<0.001$ by t-test (HGRWT vs. HGRKO (D-H). I) Comparison of cosyntropin-stimulated CORT levels after 8 weeks DEX treatment from experiments in Models 1, 2, and 3 vs. DN controls: DEX in WT (8+0), DEX+COS in WT (DEX+COS 8+0), DEX in HGRWT (8+0), and DEX in HGRKO mice (8+0). $p<0.0001$ by one-way ANOVA; ** $p<0.01$ and *** $p<0.001$ vs. DN.

Figure 7. Intermittent cosyntropin attenuates short-term DEX-induced adrenocortical apoptosis, but only endogenous ACTH secretion maintains adrenal mass and morphology on long-term DEX treatment. A) Terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) in adrenals from WT mice treated for 4 days with vehicle (DN), DEX alone (4D DEX), or DEX with daily, intraperitoneal cosyntropin (4D DEX+COS), as well as DEX-treated HGRKO animals (4D HGRKO). B) quantitation of A) p=0.0004 by one-way ANOVA. *p<0.05, ** p<0.01, *** p<0.001 for post-hoc multiple comparison testing. C) IF without TrueBlack for tyrosine hydroxylase (TH, red), with DAPI (blue), and 488-streptavidin (green) counterstains in columns 1 and 2. Giant Histiocytes of Steroid Treatment (GHOSTs) were detected by autofluorescence (cyan) in columns 3 and 4. Representative glands shown from WT animals 1-week withdrawn from 8 weeks combined DEX+COS (DEX+COS 8+1), see Fig. 1, Model 2; HGRWT on 8 weeks of DEX (8+0), and HGRKO on 8 weeks DEX (8+0), see Fig. 1, Model 3. 100 micron-thick, whole adrenal slices (columns 1 and 3, scale bar=500 microns) with magnification of the region in the white box (columns 2 and 4, scale bar=100 microns). Columns 1 and 2 are merged images at wavelengths 405 nm, 488 nm, and 647 nm; columns 3 and 4 are merged images at wavelengths 405 nm, 440 nm, and 647 nm. Morphometric analysis of D) pan-cortical area, which includes both streptavidin-positive and autofluorescent regions (p=0.0017

by one-way ANOVA), E) autofluorescent GHOST area (p=0.0053 by one-way ANOVA), F) steroidogenic cell type area, calculated as pan-cortical area minus GHOSTs (p=0.0011 by one-way ANOVA). Data in 7D are normalized to DN controls in Figure 5B; Data 7E are normalized to DN controls in Figure 5C; Data in 7F are normalized to DN controls in Figure 5D. * p<0.05 by t-test.

References

1. Fardet L, Petersen I, Nazareth I. Prevalence of long-term oral glucocorticoid prescriptions in the UK over the past 20 years. *Rheumatology (Oxford).* 2011;50(11):1982-1990.

2. Overman RA, Yeh JY, Deal CL. Prevalence of oral glucocorticoid usage in the United States: a general population perspective. *Arthritis Care Res (Hoboken).* 2013;65(2):294-298.

3. Daley-Yates P, Brealey N, Thomas S, et al. Therapeutic index of inhaled corticosteroids in asthma: A dose-response comparison on airway hyperresponsiveness and adrenal axis suppression. *Br J Clin Pharmacol.* 2021;87(2):483-493.

4. Woods CP, Argese N, Chapman M, et al. Adrenal suppression in patients taking inhaled glucocorticoids is highly prevalent and management can be guided by morning cortisol. *European journal of endocrinology.* 2015;173(5):633-642.

5. Lam LH, Sugarman JL. Adrenal Suppression With Chronic Topical Corticosteroid Use in Psoriasis Patients. *J Drugs Dermatol.* 2016;15(8):945-948.

6. Kachroo P, Stewart ID, Kelly RS, et al. Metabolomic profiling reveals extensive adrenal suppression due to inhaled corticosteroid therapy in asthma. *Nature medicine.* 2022;28(4):814-822.

7. Keller-Wood ME, Dallman MF. Corticosteroid inhibition of ACTH secretion. *Endocrine reviews.* 1984;5(1):1-24.

8. Wyllie AH, Kerr JF, Currie AR. Cellular events in the adrenal cortex following ACTH deprivation. *The Journal of pathology.* 1972;106(1):Pix.

9. Pignatti E, Leng S, Carlone DL, Breault DT. Regulation of zonation and homeostasis in the adrenal cortex. *Molecular and cellular endocrinology.* 2017;441:146-155.

10. Wyllie AH, Kerr JF, Macaskill IA, Currie AR. Adrenocortical cell deletion: the role of ACTH. *The Journal of pathology.* 1973;111(2):85-94.

945   11.   Younes AK, Younes NK. Recovery of steroid induced adrenal insufficiency. *Translational*
946   *pediatrics.* 2017;6(4):269-273.

947   12.   Livanou T, Ferriman D, James VH. Recovery of hypothalamo-pituitary-adrenal function
948   after corticosteroid therapy. *Lancet (London, England).* 1967;2(7521):856-859.

949   13.   Beuschlein F, Else T, Bancos I, et al. European Society of Endocrinology and Endocrine
950   Society Joint Clinical Guideline: Diagnosis and therapy of glucocorticoid-induced adrenal
951   insufficiency. *European journal of endocrinology.* 2024;190(5):G25-g51.

952   14.   Ebrahimi F, Widmer A, Wagner U, et al. Association of adrenal insufficiency with patient-
953   oriented health-care outcomes in adult medical inpatients. *European journal of*
954   *endocrinology.* 2019;181(6):701-709.

955   15.   Mebrahtu TF, Morgan AW, Keeley A, Baxter PD, Stewart PM, Pujades-Rodriguez M. Dose
956   dependency of iatrogenic glucocorticoid excess and adrenal insufficiency and mortality: a
957   cohort study in England. *The Journal of clinical endocrinology and metabolism.*
958   2019;104(9):3757-3767.

959   16.   Graber AL, Ney RL, Nicholson WE, Island DP, Liddle GW. NATURAL HISTORY OF
960   PITUITARY-ADRENAL RECOVERY FOLLOWING LONG-TERM SUPPRESSION WITH
961   CORTICOSTEROIDS. *The Journal of clinical endocrinology and metabolism.* 1965;25:11-
962   16.

963   17.   Avgerinos PC, Chrousos GP, Nieman LK, Oldfield EH, Loriaux DL, Cutler GB, Jr. The
964   corticotropin-releasing hormone test in the postoperative evaluation of patients with
965   cushing's syndrome. *The Journal of clinical endocrinology and metabolism.*
966   1987;65(5):906-913.

967   18.   Avgerinos PC, Nieman LK, Oldfield EH, et al. The effect of pulsatile human corticotropin-
968   releasing hormone administration on the adrenal insufficiency that follows cure of
969   Cushing's disease. *The Journal of clinical endocrinology and metabolism.* 1989;68(5):912-
970   916.

971   19.   Prete A, Paragliola RM, Bottiglieri F, et al. Factors predicting the duration of adrenal
972   insufficiency in patients successfully treated for Cushing disease and nonmalignant
973   primary adrenal Cushing syndrome. *Endocrine.* 2017;55(3):969-980.

974   20.   Gomez MT, Magiakou MA, Mastorakos G, Chrousos GP. The pituitary corticotroph is not
975   the rate limiting step in the postoperative recovery of the hypothalamic-pituitary-adrenal
976   axis in patients with Cushing syndrome. *The Journal of clinical endocrinology and*
977   *metabolism.* 1993;77(1):173-177.

978   21.   Denver RJ. Structural and functional evolution of vertebrate neuroendocrine stress
979   systems. *Annals of the New York Academy of Sciences.* 2009;1163:1-16.

980   22.   Spiga F, Zhao Z, Lightman SL. Prolonged treatment with the synthetic glucocorticoid
981   methylprednisolone affects adrenal steroidogenic function and response to inflammatory
982   stress in the rat. *Brain Behav Immun.* 2020;87:703-714.

983   23.   Calogero AE, Kamilaris TC, Johnson EO, Tartaglia ME, Chrousos G. Recovery of the rat
984   hypothalamic-pituitary-adrenal axis after discontinuation of prolonged treatment with the
985   synthetic glucocorticoid agonist dexamethasone. *Endocrinology.* 1990;127(4):1574-1579.

986   24.   Duncan PJ, McClafferty H, Nolan O, et al. Corticotroph isolation from Pomc-eGFP mice
987   reveals sustained transcriptional dysregulation characterising a mouse model of
988   glucocorticoid-induced suppression of the hypothalamus-pituitary-adrenal axis. *Journal of*
989   *neuroendocrinology.* 2022;34(7):e13165.

990   25.   Sakr N, Glazova O, Shevkova L, et al. Characterizing and Quenching Autofluorescence in
991   Fixed Mouse Adrenal Cortex Tissue. *Int J Mol Sci.* 2023;24(4).

992   26.   Farny NG, Hurt JA, Silver PA. Definition of global and transcript-specific mRNA export
993   pathways in metazoans. *Genes Dev.* 2008;22(1):66-78.

994   27.   Hurt JA, Obar RA, Zhai B, Farny NG, Gygi SP, Silver PA. A conserved CCCH-type zinc
995   finger protein regulates mRNA nuclear adenylation and export. *J Cell Biol.*
996   2009;185(2):265-277.

28. Liu X, Oh S, Kirschner MW. The uniformity and stability of cellular mass density in mammalian cell culture. *Front Cell Dev Biol.* 2022;10:1017499.

29. Kafri R, Levy J, Ginzberg MB, Oh S, Lahav G, Kirschner MW. Dynamics extracted from fixed cells reveal feedback linking cell growth to cell cycle. *Nature.* 2013;494(7438):480-483.

30. Almeida H, Ferreira J, Neves D. Macrophages of the adrenal cortex: a morphological study of the effects of aging and dexamethasone administration. *Annals of the New York Academy of Sciences.* 2004;1019:135-140.

31. Paul A, Laufer E. Endogenous biotin as a marker of adrenocortical cells with steroidogenic potential. *Molecular and cellular endocrinology.* 2011;336(1-2):133-140.

32. Romero DG, Yanes LL, de Rodriguez AF, et al. Disabled-2 is expressed in adrenal zona glomerulosa and is involved in aldosterone secretion. *Endocrinology.* 2007;148(6):2644-2652.

33. Muglia LJ, Jacobson L, Weninger SC, et al. Impaired diurnal adrenal rhythmicity restored by constant infusion of corticotropin-releasing hormone in corticotropin-releasing hormone-deficient mice. *The Journal of clinical investigation.* 1997;99(12):2923-2929.

34. Laryea G, Schütz G, Muglia LJ. Disrupting hypothalamic glucocorticoid receptors causes HPA axis hyperactivity and excess adiposity. *Molecular endocrinology (Baltimore, Md).* 2013;27(10):1655-1665.

35. Ney RL, Shimizu N, Nicholson WE, Island DP, Liddle GW. CORRELATION OF PLASMA ACTH CONCENTRATION WITH ADRENOCORTICAL RESPONSE IN NORMAL HUMAN SUBJECTS, SURGICAL PATIENTS, AND PATIENTS WITH CUSHING'S DISEASE. *The Journal of clinical investigation.* 1963;42(11):1669-1677.

36. Schlaghecke R, Kornely E, Santen RT, Ridderskamp P. The effect of long-term glucocorticoid therapy on pituitary-adrenal responses to exogenous corticotropin-releasing hormone. *The New England journal of medicine.* 1992;326(4):226-230.

37.   Dauber A, Feldman HA, Majzoub JA. Nocturnal Dexamethasone versus Hydrocortisone

1024   for the Treatment of Children with Congenital Adrenal Hyperplasia. *Int J Pediatr*

1025   *Endocrinol.* 2010;2010.

1026   38.   Mallappa A, Sinaii N, Kumar P, et al. A phase 2 study of Chronocort, a modified-release

1027   formulation of hydrocortisone, in the treatment of adults with classic congenital adrenal

1028   hyperplasia. *The Journal of clinical endocrinology and metabolism.* 2015;100(3):1137-

1029   1145.

1030   39.   Stewart J, Clegg D, Watson S. *Calculus.* Cengage Learning, Incorporated; 2021.

1031   40.   Warde KM, Smith LJ, Liu L, et al. Senescence-induced immune remodeling facilitates

1032   metastatic adrenal cancer in a sex-dimorphic manner. *Nat Aging.* 2023;3(7):846-865.

1033   41.   Lazarov T, Juarez-Carreño S, Cox N, Geissmann F. Physiology and diseases of tissue-

1034   resident macrophages. *Nature.* 2023;618(7966):698-707.

1035   42.   Foley KG, Pritchard MT, Duncan FE. Macrophage-derived multinucleated giant cells:

1036   hallmarks of the aging ovary. *Reproduction.* 2021;161(2):V5-v9.

1037   43.   Braune J, Lindhorst A, Fröba J, et al. Multinucleated Giant Cells in Adipose Tissue Are

1038   Specialized in Adipocyte Degradation. *Diabetes.* 2021;70(2):538-548.

1039   44.   Xu Y, Patterson MT, Dolfi B, et al. Adrenal gland macrophages regulate glucocorticoid

1040   production through Trem2 and TGF-β. *JCI Insight.* 2024;9(14).

1041   45.   Miller WL. StAR search--what we know about how the steroidogenic acute regulatory

1042   protein mediates mitochondrial cholesterol import. *Molecular endocrinology (Baltimore,*

1043   *Md).* 2007;21(3):589-601.

1044   46.   Miller WL, Bose HS. Early steps in steroidogenesis: intracellular cholesterol trafficking. *J*

1045   *Lipid Res.* 2011;52(12):2111-2135.

47.   Furman BL. Cosyntropin☆. In: *Reference Module in Biomedical Sciences.* Elsevier; 2017.

48. Furman BL. Cosyntropin. In: Enna SJ, Bylund DB, eds. *xPharm: The Comprehensive*

Pharmacology Reference. New York: Elsevier; 2007:1-3.

49. Hamilton DD, Cotton BA. Cosyntropin as a diagnostic agent in the screening of patients

for adrenocortical insufficiency. *Clin Pharmacol.* 2010;2:77-82.

50. Dekkers OM, Timmermans JM, Smit JW, Romijn JA, Pereira AM. Comparison of the

cortisol responses to testing with two doses of ACTH in patients with suspected adrenal

insufficiency. *European journal of endocrinology.* 2011;164(1):83-87.

51. Dickstein G, Spigel D, Arad E, Shechner C. One microgram is the lowest ACTH dose to

cause a maximal cortisol response. There is no diurnal variation of cortisol response to

submaximal ACTH stimulation. *European journal of endocrinology.* 1997;137(2):172-175.

52. Mountjoy KG, Bird IM, Rainey WE, Cone RD. ACTH induces up-regulation of ACTH

receptor mRNA in mouse and human adrenocortical cell lines. *Molecular and cellular*

endocrinology. 1994;99(1):R17-20.

53. Lebrethon MC, Jaillard C, Naville D, Bégeot M, Saez JM. Regulation of corticotropin and

steroidogenic enzyme mRNAs in human fetal adrenal cells by corticotropin, angiotensin-

II and transforming growth factor beta 1. *Molecular and cellular endocrinology.*

1994;106(1-2):137-143.

54. Lebrethon MC, Naville D, Begeot M, Saez JM. Regulation of corticotropin receptor number

and messenger RNA in cultured human adrenocortical cells by corticotropin and

angiotensin II. *The Journal of clinical investigation.* 1994;93(4):1828-1833.

55. Rehman KS, Sirianni R, Parker CR, Jr., Rainey WE, Carr BR. The regulation of

adrenocorticotrophic hormone receptor by corticotropin-releasing hormone in human fetal

adrenal definitive/transitional zone cells. *Reprod Sci.* 2007;14(6):578-587.

56. Liu Y, Smith LI, Huang V, et al. Transcriptional regulation of episodic glucocorticoid

secretion. *Molecular and cellular endocrinology.* 2013;371(1-2):62-70.

57. Stamatiades GA, Kaiser UB. Gonadotropin regulation by pulsatile GnRH: Signaling and
gene expression. *Molecular and cellular endocrinology.* 2018;463:131-141.

58. Chiavistelli S, Giustina A, Mazziotti G. Parathyroid hormone pulsatility: physiological and
clinical aspects. *Bone Res.* 2015;3:14049.

59. Lois KB, Santhakumar A, Vaikkakara S, et al. Phaeochromocytoma and ACTH-dependent
cushing's syndrome: tumour crf secretion can mimic pituitary cushing's disease. *Clinical*
endocrinology. 2016;84(2):177-184.

60. Broersen LH, Pereira AM, Jørgensen JO, Dekkers OM. Adrenal Insufficiency in
Corticosteroids Use: Systematic Review and Meta-Analysis. *The Journal of clinical*
endocrinology and metabolism. 2015;100(6):2171-2180.

61. Prete A, Bancos I. Glucocorticoid induced adrenal insufficiency. *BMJ.* 2021;374:n1380.

62. Gao X, Yamazaki Y, Tezuka Y, et al. Gender differences in human adrenal cortex and its
disorders. *Molecular and cellular endocrinology.* 2021;526:111177.

63. Duncko R, Kiss A, Skultétyová I, Rusnák M, Jezová D. Corticotropin-releasing hormone
mRNA levels in response to chronic mild stress rise in male but not in female rats while
tyrosine hydroxylase mRNA levels decrease in both sexes. *Psychoneuroendocrinology.*
2001;26(1):77-89.

64. Viau V, Bingham B, Davis J, Lee P, Wong M. Gender and puberty interact on the stress-
induced activation of parvocellular neurosecretory neurons and corticotropin-releasing
hormone messenger ribonucleic acid expression in the rat. *Endocrinology.*
2005;146(1):137-146.

65. Goel N, Bale TL. Sex differences in the serotonergic influence on the hypothalamic-
pituitary-adrenal stress axis. *Endocrinology.* 2010;151(4):1784-1794.

66. Grabek A, Dolfi B, Klein B, Jian-Motamedi F, Chaboissier MC, Schedl A. The Adult Adrenal
Cortex Undergoes Rapid Tissue Renewal in a Sex-Specific Manner. *Cell Stem Cell.*
2019;25(2):290-296.e292.

67.   Hammer GD, Basham KJ. Stem cell function and plasticity in the normal physiology of the

adrenal cortex. *Molecular and cellular endocrinology.* 2021;519:111043.

68.   Dolfi B, Gallerand A, Firulyova MM, et al. Unravelling the sex-specific diversity and

functions of adrenal gland macrophages. *Cell reports.* 2022;39(11):110949.

69.   Tronche F, Kellendonk C, Kretz O, et al. Disruption of the glucocorticoid receptor gene in

the nervous system results in reduced anxiety. *Nature genetics.* 1999;23(1):99-103.

70.   Bachmanov AA, Reed DR, Beauchamp GK, Tordoff MG. Food intake, water intake, and

drinking spout side preference of 28 mouse strains. *Behav Genet.* 2002;32(6):435-443.

71.   Schreyer SA, Chua SC, Jr., LeBoeuf RC. Obesity and diabetes in TNF-alpha receptor-

deficient mice. *The Journal of clinical investigation.* 1998;102(2):402-411.

72.   Daneva T, Spinedi E, Hadid R, Gaillard RC. Impaired hypothalamo-pituitary-adrenal axis

function in Swiss nude athymic mice. *Neuroendocrinology.* 1995;62(1):79-86.

73.   Jacobson L, Muglia LJ, Weninger SC, Pacák K, Majzoub JA. CRH deficiency impairs but

does not block pituitary-adrenal responses to diverse stressors. *Neuroendocrinology.*

2000;71(2):79-87.

74.   Advanced Cell Diagnostics. (2019). TS-46-003 Technical Note: How to correctly interpret

your RNAscope images. Rev B. Retrieved from https://acdbio.com/2019-mar-26-how-

correctly-interpret-your-rnascope-images. In.

75.   Created in BioRender. Gaston, L. (2025) https://BioRender.com/v66r332

1117