Support for the social buffering hypothesis, especially under

unpredictable precipitation regimes

Roberto Salguero-Gómez[1,*], Gabriel S Santos[1,2,3], Lauren Brent[4], Samuel J L

Gascoigne[1,5], Michael B Bonsall[1], Josh A Firth[1,6], Gregory Albery[7], Rahul Mondal[1,8],

Maja Kajin[1], & Katrina J Davis[1]

1 Department of Biology, University of Oxford, South Parks Road, Oxford, OX1 3RB, United Kingdom

8
9 2 Vale Institute of Technology - Sustainable Development, Rua Boaventura da Silva 955, Belém, Pará 66055-090, Brazil

10
11 3 Department of Biology, Espirito Santo Federal University, Avenida Fernando Ferrari, 514 Vitória, Espírito Santo 29075-910, Brazil

12 4 Centre for Research in Animal Behaviour, University of Exeter, Exeter, EX4 4QG, United Kingdom

13
5 School of Biological Sciences, University of Aberdeen,Tillydrone Ave, Aberdeen, AB24 2TZ, United Kingdom

6 School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, UK

7 School of Natural Sciences, Trinity College Dublin, Dublin, Republic of Ireland

8 Department of Biostatistics and Epidemiology, International Institute for Population Sciences, India

* Correspondence: rob.salguero@biology.ox.ac.uk

Total word count: 7,681 (Introduction through Discussion)

Total tables: 1

Total figures: 5

Total references: 78

SOMs: 3 tables, 5 figures

Acknowledgements This work was supported by a NERC Pushing the Frontiers

grant (NE/X013766/1) to RSG. Some ideas for this paper originated in discussions with

colleagues during the Royal Society meeting "Understanding age and society using

natural populations".

Conflicts of interest None

Author Contributions RSG conceived the ideas and designed the methodological

approach with input from KD and LB. RSG collected the pertinent data or accessed it

via open-access repositories. RSG curates COMADRE. RSG performed the analyses

with input from GS. RSG wrote the first draft with input from KD. All coauthors

contributed to subsequent versions of the manuscript.

Statement of inclusivity The authorship is composed of researchers of all academic

stages, genders, and several nationalities, including multiple languages.

Abstract

Sociality is hypothesised to buffer populations against environmental stochasticity by

stabilising key vital rates, yet this has rarely been tested across diverse taxa. We

compiled demographic time series from 87 populations for 66 animal species spanning

12 taxonomic classes to test whether sociality promotes demographic buffering and

whether this relationship depends on environmental context. Using stochastic

elasticities and phylogenetically comparative analyses, we find broad support for the

social buffering hypothesis: more social species exhibit lower sensitivity of population

growth to vital rate variance (lower stochastic elasticity to vital rate variances, $|T_\sigma|$),

particularly in juvenile survival and reproduction. Contrary to expectations, juvenile vital

rates, but not adult survival, showed the strongest buffering patterns in social taxa,

suggesting life-stage-specific social mechanisms. Furthermore, we show that social

species invest more in maximising the average performance of adult survival and

reproduction (higher stochastic elasticity to vital rate means, T_μ), consistent with

adaptive strategies favouring canalisation of fitness-related traits. This buffering benefit

is especially relevant in highly unpredictable environments: in climates with erratic

precipitation, social species have greater demographic advantages, exhibiting even

lower T_σ values. Together, these findings provide support for the social buffering

hypothesis, especially in highly stochastic environments. Social traits confer some

demographic stability under moderate climatic conditions, but may become more

relevant under extreme stochasticity—an insight particularly important for understanding

population resilience in a rapidly changing world.

Keywords: comparative demography; cooperative breeding; demographic buffering;

environmental stochasticity; phylogenetic analysis; sociality continuum.

Introduction

Sociality is crucial for the persistence of natural populations. The way in which individuals interact within social groups influences not only individual fitness (Silk, 2007b; Snyder-Mackler et al., 2020), but also shapes population-level processes such as survival (Creel & Christianson, 2008; Salguero-Gómez, 2024), reproduction (Lukas & Clutton-Brock, 2012; Salguero-Gómez, 2024), dispersal (Greenwood, 1980), and resistance to environmental pressures (Oro, 2020; Rubenstein & Lovette, 2007; Salguero-Gómez, 2024). Social organisations, from ephemeral aggregations to highly coordinated closed social groups, have evolved across diverse taxa as adaptive responses to predation (Krause & Ruxton, 2002), resource limitation (Koenig & Dickinson, 2016), or reproductive constraints (Emlen, 1982). Interactions that occur between members of a social organisation often mediate access to resources and mates (Clutton-Brock, 2009), affect stress physiology (Young & Bennett, 2013), and enable individuals to cope with environmental uncertainty (Bourke, 2011). As such, sociality is a central feature of the viability of natural populations.

Social organisation is a multifaceted trait. However, this characteristic has been historically often boiled down to a binary trait that is mostly examined in a few taxonomic groups. Indeed, while much of the previous literature distinguishes species as either social or non-social (Lukas & Clutton-Brock, 2012), this dichotomy is now often considered as insufficient for capturing the complexity and gradation of social behaviours observed across the animal kingdom. For instance, some spiders exhibit facultative sociality, forming colonies under high resource availability but remaining solitary otherwise (Avilés et al., 2012). Moreover, most comparative research in this

area has focused on birds (Jetz et al., 2012) and mammals (Lukas & Clutton-Brock,

2012), where social systems have been extensively studied. This narrow focus has left

a gap regarding the generality of how social organisations influence demography across

animals, including in less well-studied taxa such as reptiles, amphibians, invertebrates

(but see (Crump, 2015; Doody et al., 2013; Wong & Balshine, 2011)), and marine

organisms. A growing consensus now acknowledges the need for integrative

frameworks that account for the dimensionality of sociality (Albery et al., 2024; Doody et

al., 2013; Firth et al., 2024; Salguero-Gómez, 2024), including the kinds, intensity, and

frequency of said interactions.

There are clear implications regarding how sociality should allow populations to

persist via the social buffering hypothesis (Klug & Bonsall, 2014; Rubenstein & Lovette,

2007; Shine, 1978). This hypothesis posits that social interactions, particularly those

involving cooperative care, resource sharing, or communal defence, can mitigate the

negative demographic impacts of environmental fluctuations. By pooling resources,

enhancing survival, or smoothing reproductive success over time, sociality may reduce

demographic variance and stabilise population growth. However, the strength and

consistency of such buffering effects are expected to vary depending on the type of

social organisation, with cooperative breeders hypothesised to benefit more strongly

from such mechanisms than solitary, loosely gregarious, or communal species

(Salguero-Gómez, 2024).

Population ecologists have extensively examined the drivers of demographic

buffering (J.-M. Gaillard & Yoccoz, 2003; Morris et al., 2008; Pfister, 1998; Tuljapurkar,

1982), a life-history strategy whereby natural selection reduces temporal variability in

vital rates that most affect population fitness. This strategy is predicted to evolve under

environmental stochasticity, as a means of stabilising population growth in the face of

unpredictable ecological pressures (Bruijning et al., 2020; Hilde et al., 2020; Le Coeur et

al., 2022; Tuljapurkar, 1982). Traditionally, demographic buffering has been assessed,

among other ways (Lewontin & Cohen, 1969), by examining the coefficient of variation

in vital rates or their temporal variance (Hilde et al., 2020; Pfister, 1998). However,

recent theoretical and empirical work has emphasised the utility of stochastic elasticity

analysis in explicitly quantifying how changes in the mean and variance of vital rates

affect long-term population growth rate (λ_s) (Ezard & Coulson, 2010; S. Gascoigne et

al., 2024; S. J. L. Gascoigne et al., 2023; Giaimo & Traulsen, 2023; Haridas &

Tuljapurkar, 2005).

Stochastic elasticities directly quantify the buffering capacity of a population due

to its vital rates. In this framework, the stochastic elasticity to the mean of vital rates (T_μ)

reflects the relative importance of selection for improving average demographic

performance, whereas the stochastic elasticity to vital rate variance (T_σ) captures the

sensitivity of λ_s to interannual variability in those same vital rates. Importantly, T_μ and

T_σ probe alternative buffering mechanisms by which populations can persist in variable

environments (S. J. L. Gascoigne et al., 2024). Across species, the impacts of mean

vital rates on population growth rate in variable environments covary with life history

strategies. For example, mean adult survival is especially important in species with

longer generation times (Oli & Dobson, 2003; Saether & Bakke, 2000). Furthermore, T_σ

also covaries with life history strategies. Specifically, the quantification of T_σ across taxa

has demonstrated the increased sensitivity of species with shorter lifespans to variation

in demographic rates relative to those with longer lifespans (Morris et al., 2008).

Together, these metrics provide a powerful and evolutionarily relevant view of how

organisms manage trade-offs between maximising fitness and buffering environmental

uncertainty. Despite their theoretical relevance, T_μ and T_σ have rarely been applied in

macroecological or behavioural studies—particularly not to test the role of sociality in

modulating selection on demographic stability. Few studies have explicitly linked

behavioural traits such as social organisation to either of these components of

population dynamics, despite predictions that social behaviour may act as a buffering

mechanism (Rubenstein & Lovette, 2007; Salguero-Gómez, 2024). Bridging this gap is

essential to understand whether social systems merely enhance survival and

reproduction, or also reduce vulnerability to environmental stochasticity—a distinction

with major implications for forecasting population resilience under global change.

Here, we collate data on 87 populations and 66 animal species spanning 12

taxonomic classes, thus encompassing a wide diversity of life histories and social

systems, to examine the demographic consequences of sociality through the lens of

demographic buffering. We test four key hypotheses. (H1) Social species should exhibit

stronger demographic buffering, operationalised as lower temporal variance vital rates,

relative to solitary taxa, in line with the social buffering hypothesis (Rubenstein &

Lovette, 2007). This buffering is predicted to be especially pronounced in cooperative

breeders, where group-living and alloparental care redistribute energetic costs and

reduce reproductive uncertainty. (H2) This buffering effect should be primarily mediated

via adult stages, particularly adult survival, which tends to be more canalised in long-

lived species (J.-M. Gaillard & Yoccoz, 2003; Morris et al., 2008; Pfister, 1998), while

juvenile rates should contribute minimally to buffering patterns. (H3) More social species

should exhibit higher stochastic elasticity to the mean of adult survival and reproduction

than to juvenile survival and maturation, indicating stronger selection for maximising the

average performance of key adult vital rates in social species compared to solitary

ones. Finally, (H4) the hypothesised relationship between sociality and demographic

buffering should be modulated by environmental stochasticity, such that buffering

effects in social species diminish under highly variable conditions, *e.g.,* in arid regions

with high rainfall variability. Indeed, recent evidence has emerged suggesting that

demographic buffering can be ineffective in extremely stochastic environments

(Rodriguez-Caro et al., 2021; Santos et al., 2024). To test these four hypotheses, we

develop a comparative phylogenetic framework that incorporates time series of

demographic information, species-specific climatic variability, and a recently introduced

sociality continuum, which classifies species from more solitary to fully social (Salguero-

Gómez, 2024).

Methods

To test the hypotheses that sociality modulates demographic buffering in animals via

adult survival and its interaction with environmental stochasticity, we performed a suite

of phylogenetically-informed comparative analyses. Our analyses build on and extend

two recent methodological frameworks: one developing novel demographic metrics to

quantify the degree of demographic buffering in natural populations using matrix

population models (MPMs, hereafter) (Santos et al., 2023), and another one introducing

a continuum of social organisation for animal species, from more solitary to more social

(Salguero-Gómez, 2024). By incorporating longitudinal demographic data specifically

designed to capture temporal variation in survival and reproduction, and linking them to

the climatic patterns associated with each studied location, we also test the hypothesis

that the expected relationships between sociality and demography break down in

extreme environments.

<u>Demographic data</u>

To quantify the strength of demographic buffering across species, we first compiled

demographic time series data for animal populations from the COMADRE Animal Matrix

Database (v.4.32.3.1) (Salguero-Gomez et al., 2016), a global open-access repository

of age- and stage-structured matrix population models (MPMs) for animals. We

complemented this database with demographic data on *Suricata suricatta* (meerkat)

from (Conquet et al., 2023), which also includes an annual long-term series of MPMs.

To ensure the reliability, comparability, and biological realism of these MPMs, we

applied a stringent series of selection criteria. Indeed, in its version 4.23.3.1,

COMADRE contains 3,488 MPMs from 429 populations across 415 peer-reviewed

studies. These studies, though peer-reviewed, were not necessarily built with

comparative demography in mind by each single set of authors (Salguero-Gomez et al.,

2021). As such, we imposed the following set of selection criteria using R (v. 4.3.2) and

the R packages Rcompadre and Rage (Jones et al., 2022), and popbio (Stubben &

Milligan, 2007). We retained only MPMs that satisfied the following conditions:

- Wild populations only: We excluded MPMs parameterised with data from

laboratory settings, captive populations (*e.g.*, zoos), or experimentally

manipulated populations by filtering the metadata of the COMADRE R object by

the variables 'Captivity' and 'MatrixTreatment'. This step ensured that emergent

demographic traits would reflect natural ecological and social dynamics.

- Proper matrix decomposition: We required that each MPM be decomposed into

its survival-development (***U***) and reproduction (***F***) submatrices such that the

overall MPM ***A*** = ***U*** + ***F*** (Caswell 2001). This structure allowed us to calculate

standard life history traits and to perform matrix algebra reliably across species.

- Biological plausibility of vital rates: Using the 'cdb_flag' function in the R package

Rcompadre, we excluded matrices with missing values or with biologically

implausible values (*i.e.*., survival probabilities $0 < \sigma \leq 1$). MPMs containing ***F***

submatrices solely composed of zero-values were also excluded.

- Animals only: We retained only MPM associated with the kingdom *Animalia* by

filtering the 'Kingdom' metadata, and thus excluding any entries from *Bacteria* or

Fungi.

- Time series completeness: We grouped MPMs by species, study (based on the

'Authors', 'Source' and 'YearPublication' metadata), and population

('MatrixPopulation'), and retained only those populations for which at least three

annual MPMs were available. This step allowed us to estimate the impact of

interannual variability in vital rates on population growth while minimising bias

due to small sample sizes. We note that the primary results we show are largely

insensitive to the available data duration of each study (Table S1).

-    Population selection for multi-population species: For species represented by

multiple studies (3 species in this study), we retained the studies containing the

longest and best spatially replicated data.

-    Humans: In COMADRE, we have access to temporally replicated MPMs for

humans (*Homo sapiens sapiens*) for 42 countries (Nicol-Harper et al., 2018).

However, to strike a compromise between having humans represented in this

analysis, but not overwhelm the analyses with highly unbalanced population

replications for any one species, and also due to the difficulties in assigning

buffering abilities against environmental stochasticity to humans (Mondal et al.,

2024), we retained only one human population: Spain.

-    Standardising projection intervals: Because MPMs in COMADRE span different

projection intervals (*e.g.*, 6 months, 5 years; Figure S1), we back-transformed all

MPMs to a 1-year time step using the methods by elevating each matrix element

a_{ij} to the power of the frequency of study, following Salguero-Gómez and

Gamelon (Salguero-Gomez & Gamelon, 2021) and Salguero-Gómez (Salguero-

Gómez, 2024).

This set of selection criteria resulted in a subset of COMADRE for the next

analytical steps, comprising 87 populations from 66 animal species, and totalling 955

MPMs (Figure S2).

<u>Estimates of demographic buffering</u>

To quantify demographic buffering across the resulting subset of animal species, we

focused on interannual variability in stage-specific survival (σ), maturation (γ), and

reproduction (φ). However, as the dimension of the MPMs varied considerably (Figure

S3), we collapsed all MPMs with dimension >2 (*i.e.*, representing more than two stages

in the life cycle of the species) down to 2×2 MPMs, where the first stage represents

juveniles (*J*) and the second stage adults (*A*). To do so, we implemented the collapsing

criterion developed by Salguero-Gómez and Plotkin (2010), which allows for the

collapsing of any MPM while retaining its eigenstructure. Following suggestions from

Salguero-Gómez and Plotkin (2010), we kept the first stage unaltered, and collapsed

into the adult stage from the second life cycle stage onwards. As such, the overall

structure of the resulting MPMs contain four vital rates, shown in Eq. 1: juvenile survival

(σ_J), juvenile maturation (γ), adult survival (σ_A), and reproduction (φ).

$$\boldsymbol{A} = \begin{pmatrix} \sigma_J(1-\gamma) & \varphi \\ \sigma_J\gamma & \sigma_A \end{pmatrix}$$ (Eq. 1)

Next, we quantified the stochastic elasticities of population growth rate (λ_s) to

both changes in the mean and in the variance of vital rates across each population's

time series of MPMs. Briefly, stochastic elasticities describe the proportional change in

long-term stochastic growth rate in response to small, proportional changes in a given

vital rate, thus making them ideal tools to examine how natural populations respond to

environmental stochasticity (Haridas & Tuljapurkar, 2005; Tuljapurkar et al., 2003).

Following Haridas and Tuljapurkar (2005), we distinguished between two forms of

elasticity: (i) the elasticity of the stochastic population growth rate (λ_s) to mean of vital

rates, E_{vr}^{μ}, which isolates the impact of changing the mean value of a given vital rate *vr*

in the MPM (Eq. 1) while holding its variability constant; and (ii) the elasticity of λ_s to the

changes in the variance of a given vital rate, E_{vr}^{σ}, which isolates the impact of changing

variability while holding the mean of said vital rate (and all others) constant. These two

types of stochastic elasticities are critical for evaluating demographic buffering, which is

defined as selection to reduce the sensitivity of λ_s to variability in vital rates (Hilde et al.,

2020; Pfister, 1998; Santos et al., 2023).

We calculated these elasticities using population-specific MPMs constructed from

each of the four annual vital rate estimates shown in Eq. 1: juvenile survival (σ_J),

juvenile maturation (γ), adult survival (σ_A), and reproduction (φ). For each population,

we implemented the approach detailed in Haridas and Tuljapurkar (Haridas &

Tuljapurkar, 2005), which expresses λ_s as a function of the full time series of MPMs and

uses perturbation analysis to compute the stochastic elasticities. Specifically, we

calculated E_{vr}^{μ} as the derivative of log(λ_s) with respect to the mean of the vital rate vr,

and E_{vr}^{σ} as the derivative with respect to its standard deviation, both obtained via

numerical approximations using finite perturbations that are relative to the value of the

vital rates under examination (S. J. L. Gascoigne et al., 2023; Haridas & Tuljapurkar,

2005). By separating effects of means and variances, this approach allows us to

estimate the total elasticity of λ_s to changes in the mean of vital rates (T_{μ}) and the total

elasticity of λ_s to changes in the variance of vital rates (T_{σ}) by summing across all vital

rate elasticities in each population:

282                             $$T_{\mu} = \sum E_{vr}^{\mu} \qquad \text{(Eq. 2)}$$

283                             $$T_{\sigma} = \sum E_{vr}^{\sigma} \qquad \text{(Eq. 3)}$$

To compute these elasticities empirically, we used the 'stoch.elas' function in the

popbio R package (Stubben & Milligan, 2007) as a baseline, which we then modified to

generate separate elasticities for the mean and standard deviation of each vital rate

using Monte Carlo simulation across MPM resamples. Specifically, for each population,

we calculated the stochastic growth rate λ_s from 10,000 iterations of matrix multiplication

through the random time series of MPMs (one per year). Then, we introduced a small

perturbation (1%) to either the mean or the variance of each vital rate independently

while keeping all others fixed. We then recalculated λ_s and took the finite-difference

approximation of each elasticity (Haridas & Tuljapurkar, 2005). The separate

components of the sum of mean elasticities (T_μ) thus measures the aggregate selective

importance of vital rate averages and helps us test (H3) whether more social species

show higher stochastic elasticity to adult survival and reproduction than to juvenile

rates, suggesting stronger selection on adult vital rate averages in social than solitary

species. In contrast, the sum of variance elasticities (T_σ) measures the aggregate

selective importance of vital rate variances , and as such is directly relevant to our test

(H1) of whether more social species exhibit reduced sensitivity to vital rate variance. As

the distributions of T_μ and T_σ were right-skewed, we log$_{10}$-transformed them before

using them as response variables in our phylogenetic comparative analyses (below).

We also applied a log$_{10}$-transformation to the four vital rate stochastic elasticities to the

mean ($E_{\sigma J}{}^\mu$, $E_\gamma{}^\mu$, $E_{\sigma A}{}^\mu$ and $E_\varphi{}^\mu$) and to the variance ($E_{\sigma J}{}^\sigma$, $E_\gamma{}^\sigma$, $E_{\sigma A}{}^\sigma$ and $E_\varphi{}^\sigma$).

It is worth noting that, ecologically and evolutionarily, T_μ and T_σ (and their

respective underlying vital rate components: $E_{vr}{}^\mu$ and $E_{vr}{}^\sigma$) capture complementary

forces. The stochastic elasticity to changes in mean vital rates reflects selection for life

history traits that increase average performance (*e.g.*, survival, reproduction), while the

stochastic elasticity to changes in the variance of vital rates reflects selection against

variability (*i.e.*, demographic buffering). Under the demographic buffering hypothesis,

we expect a higher degree of sociality to correlate with lower T_σ, indicating reduced

exposure of λ_s to environmental noise. The framework from Haridas and Tuljapurkar

(Haridas & Tuljapurkar, 2005) also contains a unique property associated with elasticity

analyses, shown in Eq. 4. Specifically, with equal proportional changes in mean and

variance in vital rates, the stochastic elasticity E_{vr} is the summation of E_{vr}^μ and E_{vr}^σ.

Since, with all levels of variability in vital rates the sum of stochastic elasticity E_{vr} sums

to one, the summations of elasticity values T_μ and T_σ also equal to one. This property

holds for all possible life histories. In turn, just like their deterministic counterparts

(Takada et al., 2018), T_μ and T_σ are valuable tools for comparative analysis as their

values are unbiased by the matrix dimensionality (*i.e.*, the number of stages in the MPM

prior to downscaling to a 2×2 MPM, as done here), life history complexity (*e.g.*,

iteroparous *vs.* semelparous), or asymptotic properties (*e.g.*, declining, stable or

increasing populations as per λ_s) of the MPMs.

$$T_\mu + T_\sigma = 1 \tag{Eq. 4}$$

Importantly, T_μ and T_σ inhabit different numeric domains. T_μ is invariably positive

as the summed impacts of minor increases in the mean of all vital rates yield an

increase to the stochastic population growth rate λ_s (Haridas & Tuljapurkar, 2005). On

the other hand, T_σ is invariably negative as the summed impacts of minor increases in

the variance of all vital rates lead to a decrease in λ_s (Haridas & Tuljapurkar, 2005). In turn, to

allow for the $\log_{10}$-transformation of T_σ, $E_{\sigma J}^\sigma$, E_γ^σ, $E_{\sigma A}^\sigma$ and E_ϕ^σ, we used absolute

values for comparative analysis as in Santos et al. (2023). Consequently, the absolute-

value transformation yields values that negatively relate to the degree of demographic

buffering, with higher values of T_σ and $|E_{vr}^\sigma|$ identifying less buffered populations.

<u>Sociality classification</u>

To classify the degree of sociality across the 66 animal species included in this study,

we employed a five-level sociality continuum previously introduced and justified by

Salguero-Gómez (2024). This continuum captures gradations in social organisation

across taxa by integrating spatial cohesion, temporal stability, and the frequency of

social interactions. The five ordinal levels are: (1) Solitary: individuals live alone except

during brief breeding encounters (*e.g.*, tigers, some wasps); (2) Gregarious: individuals

form temporary or fluid groups but engage in limited social coordination (*e.g.*,

wildebeest, schooling fish); (3) Communal: individuals cohabit or share nesting sites but

do not engage in cooperative care (*e.g.*, purple martins, some reef fish); (4) Colonial:

individuals consistently share nesting or living spaces, often in dense aggregations

(*e.g.*, seabirds, coral polyps); and (5) Social: individuals form stable, cooperative groups

with persistent social bonds and behavioural coordination, including cooperative

breeding or hierarchical organisations (*e.g.*, meerkats, female elephants, baboons).

These categories were designed to be taxonomically agnostic, allowing consistent

application across a wide range of animal taxa.

We assigned sociality scores based on species-level information gathered from a

combination of curated databases (*e.g.*, Animal Diversity Web (Dewey et al., 2010),

FishBase (Froese & Pauly, 2002), IUCN Red List (Gearty & Chamberlain, 2025)), peer-

reviewed literature, and expert consultation. When available, we relied on published

ethological reviews and species-specific studies documenting group size, social

behaviour, and breeding systems. Each classification was independently validated by at

least one taxonomic expert. In cases of ambiguity, we adopted a conservative approach

by assigning the species to the lower of two adjacent categories unless strong evidence

supported otherwise. See Salguero-Gómez (2024) and Table S2 for further details on

our scoring criteria.

<u>Adult body mass</u>

As life history traits, vital rates, and demographic buffering metrics often covary

allometrically with body size (Calder 1984; Charnov 1993; Blueweiss et al. 1978), we

explicitly discounted the effect of body mass in our phylogenetic comparative models.

For each species in our dataset, we obtained adult body mass estimates from curated

trait databases specific to major vertebrate and invertebrate taxa. Mammalian and avian

body mass data were extracted primarily from the AnAge (de Magalhães & Costa,

2009) and AVONET databases (Tobias et al., 2022), respectively, while data for

reptiles, amphibians, and fish were sourced from the Amniote Database (Myhrvold et

al., 2015) and FishBase (Froese & Pauly, 2002), the latter using the rfishbase R

package (Boettiger et al., 2012). For invertebrates and less-studied groups, we relied on

MOSAIC (Bernard et al., 2023), data from Healy et al. (2019), and primary literature

searches via Web of Science using species names and the keywords "adult body mass"

or "adult weight." All mass values were converted to grams and $\log_{10}$-transformed prior

to analysis to meet assumptions of linearity and homoscedasticity. When multiple mass

estimates were available for a species, we took the mean of adult female body mass

when reported, as this tends to better reflect demographic contributions in iteroparous

animals (J.-M. Gaillard et al., 2005; Isaac et al., 2007).

<u>Phylogenetic tree</u>

To account for non-independence due to shared evolutionary history, we constructed a

phylogenetic tree spanning all 66 species in our dataset using the Open Tree of Life as

implemented in the rotl R package (Michonneau et al., 2016). First, we matched species

names to OTL taxonomy using the 'tnrs_match_names' function, resolving any

synonyms or misspellings manually. Using the matched taxon IDs, we retrieved a

synthetic, ultrametric phylogeny. This resulting tree incorporates curated backbone

information from published phylogenies across major animal clades. Branch lengths

were scaled using divergence time estimates where available in the OTL backbone. For

compatibility with downstream comparative models, we ensured the tree was fully

bifurcating and resolved any polytomies using the 'multi2di' function from the ape R

package (Paradis et al., 2004).

For species represented by multiple populations (n = 12 species, Table S2), we

modified the tree above to create population-level phylogenetic tips by duplicating the

species-level branch and assigning each population a unique identifier (e.g.,

Species_1_Pop_1, *Species_1_Pop_2*). These within-species duplicates were assigned

to nearly zero-length terminal branches, thereby preserving the species-level topology

and divergence times while accommodating population-level replication in the data. This

approach allowed us to test hypotheses using the full population-level dataset (n = 87

populations) while still accounting for phylogenetic structure at the species level. The

order of introduction of the populations within the tip of each species does not affect

assessments in macroecological studies using COMADRE (Merrien et al., 2021). The

resulting population-expanded tree was used for all comparative analyses, including

phylogenetic ANOVAs, PGLS models, and estimation of phylogenetic signal (Pagel's λ),

using the R packages caper (Orme et al., 2013), nlme (Heisterkamp et al., 2017), and

phytools (Revell, 2012).

Climatic data and environmental stochasticity

To assess whether (H4) the demographic buffering effects of sociality vary across

environments with differing climatic regimes, we obtained long-term climate data for

each population using the NASA POWER (Prediction of Worldwide Energy Resources)

database (*NASA Prediction Worldwide Energy Resources (POWER) Climate Data:*

Data Access Viewer, 2020). This resource offers high resolution climatic products at

approx. 4 km^2 resolution from January 1958. For each of the 87 populations in our

dataset, for which we have GPS coordinates in COMADRE, we extracted monthly mean

temperature (T2M) and total precipitation (PRECTOTCORR) over the duration of the

corresponding demographic time series and the 30 years preceding the start of the

study, using the nasapower R package (Sparks, 2018). We then computed interannual

summary statistics for each location and variable, including mean, maximum, minimum, and variance of precipitation and temperature, which formed the basis for estimating environmental predictability. Here, in regards to aquatic species, it is important to note that, even though species such as fish and corals are buffered on the short-term from fast fluctuations in precipitation, this abiotic factor remains a strong proxy for key hydrological processes—such as stream flow, turbidity, nutrient input, and water temperature, which in turn directly influence vital rates in freshwater and coastal environments (Buisson et al., 2008; Comte & Grenouillet, 2013; Fabricius, 2005).

We calculated three standardised metrics of climatic predictability following Colwell's framework (1974), adapted for time series climate data: constancy, contingency, and predictability. Constancy quantifies the extent to which a climatic variable remains stable over time (1 - variance/mean), while contingency reflects the extent to which fluctuations are structured and recurrent. Predictability is then defined as the sum of constancy and contingency. These metrics were calculated separately for our records of monthly temperature and precipitation. We merged these values with each population's demographic data, allowing us to explore how environmental predictability modulates the relationship between sociality and demographic buffering.

To construct a parsimonious and interpretable climatic PCA, we first reduced collinearity among the aforementioned environmental variables. Starting with a comprehensive set of temperature and precipitation-derived metrics (e.g. mean, variance, predictability), we computed all pairwise Spearman correlation coefficients to identify highly correlated variables. We visualised these correlations using custom pairwise scatterplots (Figure S4) and used a threshold of $|\rho| > 0.85$ to flag and remove

strongly collinear variables using the function 'findCorrelation' in the caret R package

(Irizarry, 2019). This threshold balances the need to retain meaningful ecological

variation while minimizing redundancy and inflation of variance along principal

component axes (Dormann et al., 2013). This step dropped off the following variables

from our next steps: contingency and predictability of temperature, as well as constancy

and contingency of precipitation. Next, for the climatic variables that we retained (mean,

variance, and constancy of temperature, as well as mean, variance, and predictability of

precipitation) we conducted a PCA using the 'prcomp' function in base R, centering and

scaling all variables. We retained only the first two principal component (PC) axes, as

only they raised associated eigenvalues > 1 (Figure S5) in agreement with Kaiser's

criterion (Legendre & Legendre, 2012).

This multivariate approach revealed how our 87 examined populations across 66

animal species are organised along two axes of climatic variability. The climatic

principal component analysis reveals two axes (Figure 2) whose associated eigenvalue

> 1 (Figure S5), and thus we retained these axes to test (H4) that the relationship

between degree of sociality and demographic buffering would break apart in more

stochastic environments. The first principal component (PC1), which explains 51% of

the variance, positions populations along a continuum of climates with more variable

precipitation (right) *vs.* more predictable precipitation (left). The second principal

component (PC2), explaining 18% of variance, places populations along a continuum of

higher mean annual precipitation (bottom) *vs.* climates with higher constancy in

temperature (top).

Comparative phylogenetic models

To test our four main hypotheses, we implemented a suite of phylogenetically-informed

comparative models that account for the shared evolutionary history of the species in

our dataset. As a first step, to discount the effect of adult body mass on the stochastic

elasticities, a standard approach in comparative demography (J. M. Gaillard et al., 1989;

Healy et al., 2019), we constructed a comparative data object using the

'comparative.data' function from the caper R package (Orme et al., 2013) to obtain the

residuals of the phylogenetic generalised least squares (PGLS) models between each

of the stochastic elasticities (Total: T_μ , T_σ ; and vital rate specific: $|E_{vr}^\mu|$, $|E_{vr}^\sigma|$) and body

mass. We then used these residuals for the next analytical steps.

To evaluate H1, which posits that more social species exhibit stronger

demographic buffering, we examined variation in the sum of stochastic elasticities to

variance (T_σ) and in the coefficients of variation of individual vital rates. We ran

phylogenetic ANOVAs with sociality (ordinal, 1–5) as the predictor and each buffering

metric as a response. These phylogenetic ANOVAs were implemented by fitting linear

models followed by post hoc Tukey HSD tests, and their outputs were interpreted within

a phylogenetic context based on trait alignment to the tree. To ensure results were

phylogenetically robust, we repeated these tests in a PGLS framework, comparing

results across both model types.

To test H2, which predicts that demographic buffering is primarily driven by adult

vital rates, we used both phylogenetic ANOVAs and PGLS models with stochastic

elasticities of the stochastic population growth rate to changes in the temporal variance

of specific vital rates, separately. Specifically, we modelled the interannual CV of

juvenile survival ($|E^{\sigma}_{\sigma J}|$), maturation ($|E^{\sigma}_{\gamma}|$), adult survival ($|E^{\sigma}_{\sigma A}|$), and reproduction

($|E^{\sigma}_{\varphi}|$) separately, using the degree of sociality as a predictor. Following significant

ANOVA results, we carried out post-hoc Tukey test comparisons to determine which

sociality levels differed most in their buffering patterns. By combining these models with

the analysis of stochastic elasticities to variance (E^{σ}) by life cycle stage, we further

tested whether adult rates were the primary targets of buffering in more social taxa.

For H3, which posits that social species exhibit greater stochastic elasticity to the

mean of adult vital rates, we analysed both total elasticity to the mean (T_{μ}) and vital-

rate-specific ($|E^{\mu}_{vr}|$) values as response variables. We used PGLS models with the

degree of sociality as predictor, and again followed up with phylogenetic ANOVAs and

Tukey tests where model residuals supported group-wise comparisons. This allowed us

to assess whether more social species rely more heavily on maximising the mean of

critical demographic parameters—particularly adult survival and reproduction—rather

than buffering their variance compared to less social species.

To test H4, that social buffering is weaker in highly stochastic environments, we

extended the above model. Specifically, we ran two separate models evaluating the

interaction between the degree of sociality and each of the two principal component

axes of our climatic PCA above—our proxy for environmental stochasticity. These were

implemented in PGLS models where response variables included either T_{σ} or $|E^{\sigma}_{vr}|$. We

centred and scaled both predictors and examined interaction terms to assess whether

the buffering benefit of sociality diminished as environments became more variable. We

also conducted model comparisons with and without interaction terms using AIC to

evaluate support for climate-modulated effects. A significant interaction term between

the degree of sociality and a climatic driver would indicate that any potential relationship

between demographic buffering and sociality is indeed modulated by the climate. A

positive effect of this interaction would imply that climatic extremes render the

demographic buffering of more social species less effective, as demographic buffering

is highest when its symbol is smallest.

Across all analyses, we tested assumptions of model normality, checked for

heteroscedasticity, and confirmed the presence of phylogenetic signal in the residuals.

When strong effects were detected, we visualised results using standardised model

coefficients and plotted predicted means from Tukey post-hoc tests to highlight

contrasts across the sociality continuum. This multi-model approach ensured that our

conclusions were robust to model structure and phylogenetic correction, and allowed us

to rigorously test how sociality interacts with demography and environment across

animal taxa.

Results

Our results support the hypothesis (H1) that more social species exhibit enhanced

demographic buffering compared to more solitary species. Indeed, more social species

show a lower stochastic elasticity to vital rate variance (T_σ) (Figure 3A). We found

significant differences in T_σ across the sociality continuum, with social species exhibiting

the lowest T_σ values, followed by communal, and then colonial, gregarious, and solitary

(P = 0.0049). Solitary species consistently showed the highest sensitivity to changes in

the variance of all vital rates.

We found partial support for the prediction (H2) that adult stages drive

demographic buffering patterns in more social species. Across the full dataset,

stochastic elasticities to the variance of reproduction ($|E^{\sigma}_{\varphi}|$) were consistently lower in

more social species (Figure 4D, P = 0.0012). However, we found no significant

difference across our social continuum regarding the stochastic elasticity to the variance

of adult survival ($|E^{\sigma}_{\sigma A}|$; Figure 4C, P = 0.6666). Unexpectedly, we found a lower value

of the stochastic elasticity to variance in juvenile survival ($|E^{\sigma}_{\sigma J}|$) in more social species

compared to less social ones (Figure 4A, P = 0.0128), implying that more social species

buffer more against stochastic environments in early stages of their development. The

stochastic elasticity to variance in maturation ($|E^{\sigma}_{\gamma}|$) showed no significant differences

across sociality groups.

We found support for our hypothesis (H3) that more social species invest more in

maximising the mean of adult vital rates than more solitary species. Indeed, total

stochastic elasticity to the mean values ($|T_{\mu}|$) differ significantly across sociality levels (P

= 0.0062; Figure 3B), with social species having the lowest values. The vital-rate-

specific analyses also reveal that social species exhibit significantly higher stochastic

elasticity values to changes in the mean of adult survival ($|E^{\mu}_{\sigma A}|$; Figure 5C, P = 0.0017)

and reproduction ($|E^{\mu}_{\varphi}|$; Figure 5D, P = 0.0125). However, the stochastic elasticity to

mean juvenile survival is also lowest in more social species ($|E^{\mu}_{\sigma J}|$; Figure 5A, P =

0.0046). There is no effect for mean maturation on the stochastic population growth rate

(λ_s) ($|E^{\mu}_{\gamma}|$; Figure 5B, P = 0.5481). The differences in vital-rate specific stochastic

elasticities are clearest between solitary and social species in post-hoc comparisons.

These patterns imply that more social species achieve population stability not only by

buffering variance, as discussed above, but also by enhancing the average

performance of adult and juvenile vital rates.

We found evidence that (H4) the benefits of social buffering are climatically-

dependent. However, we did not find evidence to support that these potential benefits

break down under highly unpredictable climatic regimes, but in fact increase non-

linearly. We found a negative, statistically significant coefficient for the interaction

between sociality and precipitation predictability (PC1 in Figure 2) in the PGLS model

predicting T_σ (P = 0.027; Table 1A). This means that, as environments become less

climatically predictable (left to right in PC1; *e.g.*, more erratic rainfall), highly social

species diminish their stochastic elasticity to variance even further compared to highly

social species in more predictable environments, indicating enhanced demographic

buffering in highly stochastic environments. The interaction between sociality and PC2

(Temperature constancy) did not significantly predict T_σ (Table 1). The fact that the

positioning of solitary, gregarious, communal, colonial, and social species along this

climatic PCA is not significantly associated with either axis (Table S3) indicates that this

key finding is not driven by environmental filtering on each group in different regions of

the climatic space.

Discussion

Understanding how sociality influences demographic performance under environmental

stochasticity is essential for predicting population responses to ongoing global change.

Although social behaviour is widely recognised as a key axis of animal life history

(Clutton-Brock, 2009; Lukas & Clutton-Brock, 2012; Silk, 2007a), its role in modulating

the demographic consequences of environmental variability remains poorly quantified

across taxa. This gap is especially important given the increasing recognition that social

traits can act as buffering mechanisms against ecological uncertainty (Albery et al.,

2022; Doody et al., 2013; Rubenstein & Lovette, 2007). Our study provides the first

macroecological and phylogenetically informed test of the social buffering hypothesis

across the animal kingdom, using time-series demographic data from 87 populations of

66 animal species spanning 12 taxonomic classes. We find broad support for the idea

that more social species exhibit stronger demographic buffering (H1), especially in adult

stages (H2), and partially support the life history theory expectation that social species

maximise mean demographic performance in adult vital rates (H3). Crucially, we show

that these buffering benefits are conditional and become more important in highly

stochastic environments (H4), thus adding a critical environmental contingency to the

social buffering hypothesis.

Our results support the social buffering hypothesis (Rubenstein & Lovette, 2007),

showing that population growth in more social species is less sensitive to interannual

fluctuations in vital rates, as measured by lower total stochastic elasticity to variance

(T_σ; (Haridas & Tuljapurkar, 2005). This finding suggests that features of systems that

are organised socially—such as alloparental care, food and information sharing,

cooperative hunting, and defence—confer demographic stability by reducing the

influence of environmental noise on fitness-related traits. These results echo and extend

previous work in specific taxonomic groups (*e.g.*, birds: (Jetz et al., 2012); mammals:

(Silk, 2007b); reptiles: (Doody et al., 2013)) and are consistent with theoretical

predictions that selection should favour strategies that reduce the fitness costs of

environmental variance (Hilde et al., 2020; Morris et al., 2008; Pfister, 1998). By

demonstrating this pattern across 12 taxonomic classes, our findings indicate that the

demographic benefits of sociality are widespread and likely not restricted to well-studied

animal groups.

The ability of social species to buffer against environmental stochasticity is not

uniformly distributed across their vital rates or life cycle stages. Rather, we provide

evidence that their vital rates can exert different levels of demographic buffering. The

most prominent buffering mechanism we show here in the context of sociality is related

to the forces of natural selection that shape adult survival and reproduction through

environmental canalisation. Our finding is consistent with theoretical predictions

(Tuljapurkar, 1982) and empirical evidence (J.-M. Gaillard & Yoccoz, 2003; Morris et al.,

2008; Santos et al., 2023) that traits contributing most to population growth rate are

often the most canalised (*i.e.*, buffered). In this sense, it is not surprising that, in our

study, juvenile survival and maturation show little evidence of buffering (*i.e.*, greater

values of $|E_{vr}^{\sigma}|$) in more social species, as juvenile vital rates are typically under weaker

selection—and thus more variable—in long-lived species (McDonald et al., 2016;

Stearns, 1976, 1999). Social systems likely reinforce this canalisation by promoting

adult stability through mechanisms such as cooperative breeding, protection from

predation, and shared resource acquisition (Bourke, 2011; Koenig & Dickinson, 2016).

In addition, we also provide evidence of a second buffering mechanism that acts on

juvenile survival—where canalisation is less expected. Although juvenile survival is

generally under weak natural selection, we show that increasing sociality significantly

reduces the extent to which variation in this vital rate affects population growth. As

sociality increases, juvenile survival becomes more buffered—a pattern not detected in

vital rates that are already highly buffered, such as adult survival. This secondary

buffering mechanism reinforces the idea that demographic buffering is life cycle stage

dependent, but also challenges the assumption that demographic buffering occurs

exclusively at vital rates where stability most enhances population growth (Hilde et al.,

2020; Morris et al., 2008; Pfister, 1998).

In partial support of H3, we found that social species do not exhibit higher total

stochastic elasticity to mean vital rates (T_μ; (Haridas & Tuljapurkar, 2007) overall, but do

show higher elasticity to the mean of adult survival and reproduction specifically

compared to less social species. This finding suggests that, while sociality may not

increase the overall importance of mean performance across all vital rates, it does

amplify the role of key adult traits in determining population growth. This pattern is

consistent with life-history theory predicting that species should maximise the

performance of traits most important to fitness (Stearns, 1999). The elevated $|E^\mu_{\sigma A}|$ and

$|E^\mu_\square|$ values in more social species likely reflect adaptive strategies that enhance adult

demographic performance in stable or cooperative contexts. However, this pattern did

not hold across juvenile traits, again pointing to a stage-specific effect of sociality on

demographic   investment.   Indeed,   social   buffering   might   be   expected   to

disproportionately benefit adults over juveniles if social relationships establish over the

life course (Firth et al. 2024). Specifically, if adults typically hold stronger and more consistent social relationships compared to juveniles (Woodman et al. 2024), this may enhance their ability to leverage group support under environmental stress. For instance, in cooperatively breeding mammalian species (such as meerkats, or baboons), dominant adult breeders often hold established social relationships to other group members and benefit from priority access to shared resources, reduced predation risk through increased vigilance, and assistance in offspring care; these mechanisms would all directly stabilise adult survival and fecundity more than they would affect juvenile traits (Clutton-Brock et al. 2016; Silk et al. 2016). Similarly, adult individuals in stable avian breeding groups often benefit from load-lightening (via helper contributions to territory defense, and joint nest provisioning), thus potentially reducing adult mortality and reproductive variability relative to juvenile birds, who are often less integrated into established social roles in cooperatively breeding systems (Rubenstein & Abbot 2017). These differences in social roles across the life-course suggest that demographic buffering by sociality might be particularly pronounced for adult vital rates.

Importantly, our results reveal that the demographic buffering effects of sociality are context-dependent and may be more important in highly stochastic environments (H4). This finding adds a novel macroecological insight to the social buffering hypothesis. Contrary to some work suggesting that demographic buffering mechanisms can break down under extreme environmental conditions (Rodriguez-Caro et al., 2021; Santos et al., 2024), our results indicate that sociality may become even more effective in buffering demographic processes against environmental stochasticity when conditions are highly unpredictable. This pattern echoes long-standing predictions about

the evolution of cooperative systems in harsh environments, such as the development

of eusociality in termites, (semi)social cockroaches, and naked mole rats, where group

living is thought to have evolved in response to arid, variable conditions (Alexander,

1974; Faulkes & Bennett, 2013; Field & Toyoizumi, 2020; Lubin & Bilde, 2007). Rather

than social organisation being strained or cooperative behaviours faltering under

extreme variability (Hayes, 2017), our findings suggest that sociality can enhance

demographic resilience precisely where environmental unpredictability is greatest. Thus,

we highlight the importance of considering environmental context in assessments of

social evolution and suggest that sociality may play a crucial role in mediating species'

demographic responses to ongoing climate change.

Together, our findings suggest that sociality promotes demographic stability by

buffering the variance of adult vital rates, and that these benefits are environmentally

dependent. In doing so, we bridge theoretical work on demographic buffering (Hilde et

al., 2020; Pfister, 1998; Tuljapurkar, 1982) with recent calls to integrate social behaviour

into eco-evolutionary frameworks (Albery et al., 2022; Firth et al., 2024; Salguero-

Gómez, 2024). Our use of time-series demographic data across a broad range of taxa,

combined with a structured classification of sociality and high-resolution climate

products, offers a general framework to assess how social and environmental factors

interact to shape population dynamics. In an era of increasing environmental

unpredictability (Bathiany et al., 2018), understanding the limits of social buffering is

vital for predicting which species are most vulnerable to demographic destabilisation—

and why.

References

Albery, G. F., Bansal, S., & Silk, M. J. (2024). Comparative approaches in social

network ecology. *Ecology Letters*, *27*(1), e14345. https://doi.org/10.1111/ele.14345

Albery, G. F., Clutton-Brock, T. H., Morris, A., Morris, S., Pemberton, J. M., Nussey, D.

H., & Firth, J. A. (2022). Ageing red deer alter their spatial behaviour and become

less social. *Nature Ecology & Evolution*, *6*(8), 1231–1238.

https://www.nature.com/articles/s41559-022-01817-9

Alexander, R. D. (1974). The evolution of social behavior. *Annual Review of Ecology*

and Systematics, *5*(1), 325–383.

https://doi.org/10.1146/annurev.es.05.110174.001545

Avilés, L., Harwood, G., & Koenig, W. (2012). A quantitative index of sociality and its

application to group-living spiders and other social organisms. *Ethology: Formerly*

Zeitschrift Für Tierpsychologie, *118*(12), 1219–1229.

https://doi.org/10.1111/eth.12028

Bathiany, S., Dakos, V., Scheffer, M., & Lenton, T. M. (2018). Climate models predict

increasing temperature variability in poor countries. *Science Advances*, *4*(5),

eaar5809. https://doi.org/10.1126/sciadv.aar5809

Bernard, C., Santos, G. S., Deere, J. A., Rodriguez-Caro, R., Capdevila, P., Kusch, E.,

Gascoigne, S. J. L., Jackson, J., & Salguero-Gómez, R. (2023). MOSAIC - A

unified trait database to complement structured population models. *Scientific Data*,

10(1), 335. https://doi.org/10.1038/s41597-023-02070-w

Boettiger, C., Lang, D. T., & Wainwright, P. C. (2012). rfishbase: exploring, manipulating

and visualizing FishBase data from R. *Journal of Fish Biology*, *81*(6), 2030–2039.

https://doi.org/10.1111/j.1095-8649.2012.03464.x

Bourke, A. F. G. (2011). Social group maintenance. In *Principles of Social Evolution* (pp. 129–161). Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199231157.003.0005

Bruijning, M., Metcalf, C. J. E., Jongejans, E., & Ayroles, J. F. (2020). The evolution of variance control. *Trends in Ecology & Evolution*, *35*(1), 22–33. https://doi.org/10.1016/j.tree.2019.08.005

Buisson, L., Thuiller, W., Lek, S., Lim, P., & Grenouillet, G. (2008). Climate change hastens the turnover of stream fish assemblages. *Global Change Biology*, *14*(10), 2232–2248. https://doi.org/10.1111/j.1365-2486.2008.01657.x

Clutton-Brock, T. (2009). Cooperation between non-kin in animal societies. *Nature*, *462*(7269), 51–57. https://doi.org/10.1038/nature08366

Colwell, R. K. (1974). Predictability, constancy, and contingency of periodic phenomena. *Ecology*, *55*(5), 1148–1153. https://doi.org/10.2307/1940366

Comte, L., & Grenouillet, G. (2013). Do stream fish track climate change? Assessing distribution shifts in recent decades. *Ecography*, *36*(11), 1236–1246. https://doi.org/10.1111/j.1600-0587.2013.00282.x

Conquet, E., Ozgul, A., Blumstein, D. T., Armitage, K. B., Oli, M. K., Martin, J. G. A., Clutton-Brock, T. H., & Paniw, M. (2023). Demographic consequences of changes in environmental periodicity. *Ecology*, *104*(3), e3894. https://doi.org/10.1002/ecy.3894

Creel, S., & Christianson, D. (2008). Relationships between direct predation and risk effects. *Trends in Ecology & Evolution*, *23*(4), 194–201.

https://doi.org/10.1016/j.tree.2007.12.004

Crump, M. L. (2015). Anuran reproductive modes: Evolving perspectives. *Journal of*

Herpetology, *49*(1), 1–16. https://doi.org/10.1670/14-097

de Magalhães, J. P., & Costa, J. (2009). A database of vertebrate longevity records and

their relation to other life-history traits. *Journal of Evolutionary Biology*, *22*(8), 1770–

1774. https://doi.org/10.1111/j.1420-9101.2009.01783.x

Dewey, T., Shefferly, N., & Havens, A. (2010). Animal Diversity Web. *University of*.

https://animaldiversity.ummz.umich.edu/site/accounts/information/Felis_silvestris.ht

ml

Doody, J. S., Burghardt, G. M., & Dinets, V. (2013). Breaking the social–non‐social

dichotomy: A role for reptiles in vertebrate social behavior research? *Ethology:*

Formerly Zeitschrift Für Tierpsychologie, *119*(2), 95–103.

https://doi.org/10.1111/eth.12047

Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R.

G., Gruber, B., Lafourcade, B., Leitão, P. J., Münkemüller, T., McClean, C.,

Osborne, P. E., Reineking, B., Schröder, B., Skidmore, A. K., Zurell, D., &

Lautenbach, S. (2013). Collinearity: a review of methods to deal with it and a

simulation study evaluating their performance. *Ecography*, *36*(1), 27–46.

https://doi.org/10.1111/j.1600-0587.2012.07348.x

Emlen, S. T. (1982). The evolution of helping. I. an ecological constraints model. *The*

American Naturalist, *119*(1), 29–39. https://doi.org/10.1086/283888

Ezard, T. H. G., & Coulson, T. (2010). How sensitive are elasticities of long-run

stochastic growth to how environmental variability is modelled? *Ecological*

Modelling, *221*(2), 191–200. https://doi.org/10.1016/j.ecolmodel.2009.09.017

Fabricius, K. E. (2005). Effects of terrestrial runoff on the ecology of corals and coral

reefs: review and synthesis. *Marine Pollution Bulletin*, *50*(2), 125–146.

https://doi.org/10.1016/j.marpolbul.2004.11.028

Faulkes, C. G., & Bennett, N. C. (2013). Plasticity and constraints on social evolution in

African mole-rats: ultimate and proximate factors. *Philosophical Transactions of the*

Royal Society of London. Series B, Biological Sciences, *368*(1618), 20120347.

https://doi.org/10.1098/rstb.2012.0347

Field, J., & Toyoizumi, H. (2020). The evolution of eusociality: no risk-return tradeoff but

the ecology matters. *Ecology Letters*, *23*(3), 518–526.

https://doi.org/10.1111/ele.13452

Firth, J. A., Albery, G. F., Bouwhuis, S., Brent, L. J. N., & Salguero-Gómez, R. (2024).

Understanding age and society using natural populations. *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences,

379(1916), 20220469. https://doi.org/10.1098/rstb.2022.0469

Froese, R., & Pauly, D. (2002). FishBase: A Global Information System on Fishes.

World Wide Web Electronic Publication: Www.fishbase.org., 000–000.

https://oceanrep.geomar.de/id/eprint/2964

Gaillard, J. M., Pontier, D., Allaine, D., Lebreton, J. D., Trouvilliez, J., & Clobert, J.

(1989). An analysis of demographic tactics in birds and mammals. *Oikos* , *56*(1),

59–76. https://doi.org/10.2307/3566088

Gaillard, J.-M., & Yoccoz, N. G. (2003). Temporal variation in survival of mammals: A

case of environmental canalization? *Ecology*, *84*(12), 3294–3306.

https://doi.org/10.1890/02-0409

Gaillard, J.-M., Yoccoz, N. G., Lebreton, J.-D., Bonenfant, C., Devillard, S., Loison, A.,

Pontier, D., & Allaine, D. (2005). Generation time: a reliable metric to measure life-

history variation among mammalian populations. *The American Naturalist*, *166*(1),

119–123; discussion 124–128. https://doi.org/10.1086/430330

Gascoigne, S. J. L., Kajin, M., & Salguero-Gómez, R. (2024). Criteria for buffering in

ecological modeling. *Trends in Ecology & Evolution*, *39*(2), 116–118.

https://doi.org/10.1016/j.tree.2023.11.006

Gascoigne, S. J. L., Kajin, M., Tuljapurkar, S., Santos, G. S., Compagnoni, A., Steiner,

U. K., Vinton, A. C., Jaggi, H., Sepil, I., & Salguero-Gómez, R. (2023). Structured

demographic buffering: A framework to explore the environment drivers and

demographic mechanisms underlying demographic buffering. In *bioRxiv* (p.

2023.07.20.549848). https://doi.org/10.1101/2023.07.20.549848

Gascoigne, S., Kajin, M., Sepil, I., & Salguero-Gomez, R. (2024). Testing for efficacy in

four measures of demographic buffering. In *EcoEvoRxiv*.

https://doi.org/10.32942/x23911

Gearty, W., & Chamberlain, S. (2025). *rredlist: "IUCN" Red List Client*.

Giaimo, S., & Traulsen, A. (2023). Generation time in stage-structured populations

under fluctuating environments. *The American Naturalist*, *201*(3), 404–417.

https://doi.org/10.1086/722608

Greenwood, P. J. (1980). Mating systems, philopatry and dispersal in birds and

mammals. *Animal Behaviour*, *28*(4), 1140–1162. https://doi.org/10.1016/s0003-

3472(80)80103-5

Haridas, C. V., & Tuljapurkar, S. (2005). Elasticities in variable environments: properties

and implications. *The American Naturalist*, *166*(4), 481–495.

https://doi.org/10.1086/444444

Haridas, C. V., & Tuljapurkar, S. (2007). Time, transients and elasticity. *Ecology Letters*,

10, 1143–1153.

Hayes, L. D. (2017). Cooperative Breeding in Vertebrates: Studies of Ecology,

Evolution, and Behavior. Edited by Walter D. Koenig and Janis L. Dickinson;

illustrated by Stef den Ridder. Cambridge and New York: Cambridge University

Press. $140.00. x + 379 p. + 12 pl.; ill.; index. ISBN: 978-1-107-04343-5. 2016. *The*

Quarterly Review of Biology, *92*(1), 99–100. https://doi.org/10.1086/690874

Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-Gomez, R., & Buckley, Y. M. (2019).

Animal life history is shaped by the pace of life and the distribution of age-specific

mortality and reproduction. *Nature Ecology & Evolution*, *3*(8), 1217–1224.

https://doi.org/10.1038/s41559-019-0938-7

Heisterkamp, S., Willigen, E., Diderichsen, P.-M., & Maringwa, J. (2017). Update of the

nlme package to allow a fixed standard deviation of the residual error. *The R*

Journal, *9*(1), 239. https://doi.org/10.32614/rj-2017-010

Hilde, C. H., Gamelon, M., Saether, B. E., Gaillard, J. M., Yoccoz, N. G., & Pelabon, C.

(2020). The Demographic Buffering Hypothesis: Evidence and Challenges. *Trends*

in Ecology & Evolution, *35*(6), 523–538. https://doi.org/10.1016/j.tree.2020.02.004

Irizarry, R. A. (2019). The caret package. In *Introduction to Data Science* (pp. 523–528).

Chapman and Hall/CRC. https://doi.org/10.1201/9780429341830-30

Isaac, N. J. B., Turvey, S. T., Collen, B., Waterman, C., & Baillie, J. E. M. (2007).

Mammals on the EDGE: conservation priorities based on threat and phylogeny. *PloS One*, *2*(3), e296. https://doi.org/10.1371/journal.pone.0000296

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity of birds in space and time. *Nature*, *491*(7424), 444–448. https://doi.org/10.1038/nature11631

Jones, O. R., Barks, P., Stott, I., James, T. D., Levin, S., Petry, W. K., Capdevila, P., Che-Castaldo, J., Jackson, J., Romer, G., Schuette, C., Thomas, C. C., & Salguero-Gomez, R. (2022). Rcompadre and Rage-Two R packages to facilitate the use of the COMPADRE and COMADRE databases and calculation of life-history traits from matrix population models. *Methods in Ecology and Evolution / British Ecological Society*. https://doi.org/10.1111/2041-210x.13792

Keesey, M. T. (2020). *PhyloPic*. PhyloPic. http://phylopic.org

Klug, H., & Bonsall, M. B. (2014). What are the benefits of parental care? The importance of parental effects on developmental rate. *Ecology and Evolution*, *4*(12), 2330–2351. https://doi.org/10.1002/ece3.1083

Koenig, W. D., & Dickinson, J. L. (2016). *Cooperative Breeding in Vertebrates: Studies of Ecology, Evolution, and Behavior*. Cambridge University Press.

Krause, J., & Ruxton, G. D. (2002). Evolutionary considerations. In *Living in Groups* (pp. 104–122). Oxford University PressOxford. https://doi.org/10.1093/oso/9780198508175.003.0007

Le Coeur, C., Yoccoz, N. G., Salguero-Gómez, R., & Vindenes, Y. (2022). Life history adaptations to fluctuating environments: Combined effects of demographic buffering and lability. *Ecology Letters*, *25*(10), 2107–2119.

https://doi.org/10.1111/ele.14071

Legendre, P., & Legendre, L. (2012). *Numerical Ecology* (3rd ed.). Elsevier.

Lewontin, R. C., & Cohen, D. (1969). On population growth in randomly varying
environments. *Proceedings of the National Academy of Sciences of the United*
States of America, *62*(4), 1056 – &. https://doi.org/10.1073/pnas.62.4.1056

Lubin, Y., & Bilde, T. (2007). The evolution of sociality in spiders. In *Advances in the*
Study of Behavior (pp. 83–145). Elsevier. https://doi.org/10.1016/s0065-
3454(07)37003-4

Lukas, D., & Clutton-Brock, T. (2012). Cooperative breeding and monogamy in
mammalian societies. *Proceedings. Biological Sciences*, *279*(1736), 2151–2156.
https://doi.org/10.1098/rspb.2011.2468

McDonald, J. L., Bailey, T., Delahay, R. J., McDonald, R. A., Smith, G. C., & Hodgson,
D. J. (2016). Demographic buffering and compensatory recruitment promotes the
persistence of disease in a wildlife population. *Ecology Letters*, *19*(4), 443–449.
https://doi.org/10.1111/ele.12578

Merrien, T., Davis, K., Di Marco, M., Capdevila, P., & Salguero-Gomez, R. (2021).
Human pressures filter out the less resilient demographic strategies in natural
populations of plants and animals worldwide. *bioRxiv*, 2021.09.29.462372.

Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl: an R package to interact with
the Open Tree of Life data. *Methods in Ecology and Evolution / British Ecological*
Society, *7*(12), 1476–1481. https://doi.org/10.1111/2041-210x.12593

Mondal, R., Aburto, J. M., Sear, R., Tuljapurkar, S., Mishra, U. S., & Salguero-Gómez,
R. (2024). Human populations with low survival at advanced ages and postponed

fertility reduce long-term growth in high inflation environments. In *bioRxiv*.

https://doi.org/10.1101/2024.07.31.606043

Morris, W. F., Pfister, C. A., Tuljapurkar, S., Haridas, C. V., Boggs, C. L., Boyce, M. S.,

Bruna, E. M., Church, D. R., Coulson, T., Doak, D. F., Forsyth, S., Gaillard, J. M.,

Horvitz, C. C., Kalisz, S., Kendall, B. E., Knight, T. M., Lee, C. T., & Menges, E. S.

(2008). Longevity can buffer plant and animal populations against changing climatic

variability. *Ecology*, *89*(1), 19–25.

Myhrvold, N. P., Baldridge, E., Chan, B., Sivam, D., Freeman, D. L., & Ernest, S. K. M.

(2015). An amniote life-history database to perform comparative analyses with

birds,   mammals,   and   reptiles.   *Ecology*,   *96*(11),   3109–3000.

https://doi.org/10.1890/15-0846r.1

NASA Prediction Worldwide Energy Resources (POWER) Climate Data: Data Access

Viewer. (2020).

Nicol-Harper, A., Dooley, C., Packman, D., Mueller, M., Bijak, J., Hodgson, D., Townley,

S., & Ezard, T. (2018). Inferring transient dynamics of human populations from

matrix   non□normality.   *Population   Ecology*,   *60*(1-2),   185–196.

https://doi.org/10.1007/s10144-018-0620-y

Oli, M. K., & Dobson, F. S. (2003). The relative importance of life-history variables to

population growth rate in mammals: Cole's prediction revisited. *The American*

Naturalist, *161*(3), 422–440. https://doi.org/10.1086/367591

Orme, D., Freckleton, R., Thomson, G., Petzoldt, T., Fritz, S., Isaac, N., & Pearse, W.

(2013). *The caper package: comparative analysis of phylogenetics and evolution in*

R. http://cran.r-project.org/web/packages/caper/

Oro, D. (2020). *Perturbation, Behavioural Feedbacks, and Population Dynamics in*

Social Animals: When to Leave and Where to Go. Oxford University Press.

https://play.google.com/store/books/details?id=f4_UDwAAQBAJ

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and

Evolution in R language. *Bioinformatics* , *20*(2), 289–290.

https://doi.org/10.1093/bioinformatics/btg412

Pfister, C. A. (1998). Patterns of variance in stage-structured populations: evolutionary

predictions and ecological implications. *Proceedings of the National Academy of*

Sciences of the United States of America, *95*(1), 213–218.

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and

other things). *Methods in Ecology and Evolution*, *3*(2), 217–223.

https://doi.org/10.1111/j.2041-210x.2011.00169.x

Rodriguez-Caro, R. C., Capdevila, P., Gracia, E., Barbosa, J. M., Gimenez, A., &

Salguero-Gomez, R. (2021). The limits of demographic buffering in coping with

environmental variation. *Oikos* , *130*(8), 1346–1358.

https://doi.org/10.1111/oik.08343

Rubenstein, D. R., & Lovette, I. J. (2007). Temporal environmental variability drives the

evolution of cooperative breeding in birds. *Current Biology: CB*, *17*(16), 1414–1419.

https://doi.org/10.1016/j.cub.2007.07.032

Saether, B. E., & Bakke, O. (2000). Avian life history variation and contribution of

demographic traits to the population growth rate. *Ecology*, *81*(3), 642–653.

Salguero-Gómez, R. (2024). More social species live longer, have longer generation

times and longer reproductive windows. *Philosophical Transactions of the Royal*

Society of London. Series B, Biological Sciences, *379*(1916), 20220459.

https://doi.org/10.1098/rstb.2022.0459

Salguero-Gomez, R., & Gamelon, M. (2021). *Demographic Methods across the Tree of*

Life. Oxford University Press.

https://play.google.com/store/books/details?id=wtg9EAAAQBAJ

Salguero-Gomez, R., Jackson, J., & Gascoigne, S. J. L. (2021). Four key challenges in

the open-data revolution. *The Journal of Animal Ecology*, *90*(9), 2000–2004.

https://doi.org/10.1111/1365-2656.13567

Salguero-Gomez, R., Jones, O. R., Archer, C. R., Bein, C., de Buhr, H., Farack, C.,

Gottschalk, F., Hartmann, A., Henning, A., Hoppe, G., Romer, G., Ruoff, T.,

Sommer, V., Wille, J., Voigt, J., Zeh, S., Vieregg, D., Buckley, Y. M., Che-Castaldo,

J., … Vaupel, J. W. (2016). COMADRE: a global data base of animal demography.

The Journal of Animal Ecology, *85*(2), 371–384. https://doi.org/10.1111/1365-

2656.12482

Salguero-Gómez, R., & Plotkin, J. B. (2010). Matrix dimensions bias demographic

inferences: implications for comparative plant demography. *The American*

Naturalist, *176*(6), 710–722. https://doi.org/10.1086/657044

Santos, G. S., Gascoigne, S. J. L., Dias, A. T. C., Kajin, M., & Salguero-Gómez, R.

(2023). A unified framework to identify demographic buffering in natural

populations. In *bioRxiv* (p. 2023.07.03.547528).

https://doi.org/10.1101/2023.07.03.547528

Santos, G. S., Yang, X., L Gascoigne, S. J., Compagnoni, A., Dias, A. T. C.,

Tuljapurkar, S., Kajin, M., & Salguero-Gómez, R. (2024). Population responses to

environmental stochasticity are primarily driven by survival-reproduction trade-offs

and mediated by aridity. In *bioRxiv*. https://doi.org/10.1101/2024.07.24.604949

Shine, R. (1978). Propagule size and parental care: The "safe harbor" hypothesis.

Journal of Theoretical Biology, *75*(4), 417–424. https://doi.org/10.1016/0022-

5193(78)90353-3

Silk, J. B. (2007a). Social components of fitness in primate groups. *Science*, *317*(5843),

1347–1351. https://doi.org/10.1126/science.1140734

Silk, J. B. (2007b). The adaptive value of sociality in mammalian groups. *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences,

362(1480), 539–559. https://doi.org/10.1098/rstb.2006.1994

Snyder-Mackler, N., Burger, J. R., Gaydosh, L., Belsky, D. W., Noppert, G. A., Campos,

F. A., Bartolomucci, A., Yang, Y. C., Aiello, A. E., O'Rand, A., Harris, K. M., Shively,

C. A., Alberts, S. C., & Tung, J. (2020). Social determinants of health and survival

in humans and other animals. *Science (New York, N.Y.)*, *368*(6493), eaax9553.

https://doi.org/10.1126/science.aax9553

Sparks, A. (2018). Nasapower: A NASA POWER global meteorology, surface solar

energy and climatology data client for R. *Journal of Open Source Software*, *3*(30),

1035. https://doi.org/10.21105/joss.01035

Stearns, S. C. (1976). Life-history tactics: a review of the ideas. *The Quarterly Review*

of Biology, *51*(1), 3–47. https://doi.org/10.1086/409052

Stearns, S. C. (1999). *The Evolution of Life Histories*. Oxford University Press.

Stubben, C., & Milligan, B. (2007). Estimating and analyzing demographic models using

the popbio package in R. *Journal of Statistical Software*, *22*(11), 1–23.

Takada, T., Kawai, Y., & Salguero-Gómez, R. (2018). A cautionary note on elasticity

analyses in a ternary plot using randomly generated population matrices.

Population Ecology, *60*(1-2), 37–47. https://doi.org/10.1007/s10144-018-0619-4

Tobias, J. A., Sheard, C., Pigot, A. L., Devenish, A. J. M., Yang, J., Sayol, F., Neate-

Clegg, M. H. C., Alioravainen, N., Weeks, T. L., Barber, R. A., Walkden, P. A.,

MacGregor, H. E. A., Jones, S. E. I., Vincent, C., Phillips, A. G., Marples, N. M.,

Montaño-Centellas, F. A., Leandro-Silva, V., Claramunt, S., … Schleuning, M.

(2022). AVONET: morphological, ecological and geographical data for all birds.

Ecology Letters, *25*(3), 581–597. https://doi.org/10.1111/ele.13898

Tuljapurkar, S. (1982). Population-dynamics in variable environments. 2. Correlated

environments, sensitivity analsyis and dynamics. *Theoretical Population Biology*,

21(1), 114–140. https://doi.org/10.1016/0040-5809(82)90009-0

Tuljapurkar, S., Horvitz, C. C., & Pascarella, J. B. (2003). The many growth rates and

elasticities of populations in random environments. *The American Naturalist*,

162(4), 489–502. https://doi.org/10.1086/378648

Wong, M., & Balshine, S. (2011). The evolution of cooperative breeding in the African

cichlid fish, Neolamprologus pulcher. *Biological Reviews of the Cambridge*

Philosophical Society, *86*(2), 511–530. https://doi.org/10.1111/j.1469-

185X.2010.00158.x

Young, A. J., & Bennett, N. C. (2013). Intra-sexual selection in cooperative mammals

and birds: why are females not bigger and better armed? *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences,

368(1631), 20130075. https://doi.org/10.1098/rstb.2013.0075

Table 1. Populations located in climates with low predictability for precipitation regimes show a disproportionate

increase in their demographic buffering abilities. Estimates and associated P-values (P < 0.05 in bold) for a battery of

phylogenetic generalised least square (pgls) models evaluating the effects of the degree of sociality and climate on the

elasticity of the stochastic population growth rate (λ_s) to (**A**) the total change in mean and variance of all vital rates in each

population, (**B**) changes in the mean of each vital rate, and (**C**) change in the variance of each vital rate, separately. PC1

and PC2 are defined in Figure 2, and correspond to a continuum of predictability in precipitation and of constancy of

temperature, respectively. A significant interaction between sociality and either principal component axis indicates that the

relationship between demographic buffering and sociality changes along climatic regimes.

Stochastic elasticity to		Predictor	Estimate	P	Predictor	Estimate	P
A. Total	Mean - T_μ	Sociality	0.221	0.541	Sociality	0.052	0.878
		PC1	0.894	0.165	PC2	-0.268	0.770
		Soc × PC1	-0.231	0.286	Soc × PC2	0.22	0.472
	Variance - T_σ	Soc	1.778	**0.016**	Soc	0.634	0.497
		PC1	4.233	**0.026**	PC2	-1.362	0.585

		Soc × PC1	-1.431	**0.027**	Soc × PC2	1.290	0.131
B. Changes in mean	Juvenile survival - $|E_{\sigma_J}^{\mu}|$	Soc	0.423	0.446	Soc	0.052	0.911
		PC1	0.768	0.424	PC2	-0.816	0.516
		Soc × PC1	-0.411	0.216	Soc × PC2	0.631	0.141
	Maturation - $|E_{\gamma}^{\mu}|$	Soc	-0.503	0.538	Soc	-0.724	0.337
		PC1	1.027	0.468	PC2	-1.278	0.524
		Soc × PC1	-0.345	0.475	Soc × PC2	0.551	0.408
	Adult survival - $|E_{\sigma_A}^{\mu}|$	Soc	0.183	0.751	Soc	0.028	0.960
		PC1	1.150	0.257	PC2	0.032	0.983
		Soc × PC1	-0.235	0.491	Soc × PC2	0.069	0.889
	Reproduction - $|E_{\varphi}^{\mu}|$	Soc	0.382	0.656	Soc	-0.216	0.778
		PC1	2.262	0.141	PC2	0.216	0.916
		Soc × PC1	-0.677	0.193	Soc × PC2	0.482	0.481
C.	Juvenile	Soc	0.518	0.631	Soc	0.112	0.911

Changes in variance	survival - $|E_{\sigma J}^{\sigma}|$	PC1	1.791	0.342	PC2	-1.321	0.625
		Soc × PC1	-0.607	0.346	Soc × PC2	0.595	0.507
	Maturation - $|E_{\gamma}^{\sigma}|$	Soc	-0.156	0.888	Soc	-0.413	0.691
		PC1	1.790	0.356	PC2	-1.711	0.540
		Soc × PC1	-0.497	0.451	Soc × PC2	0.569	0.538
	Adult survival - $|E_{\sigma A}^{\sigma}|$	Soc	0.283	0.384	Soc	0.150	0.613
		PC1	0.525	0.352	PC2	0.151	0.849
		Soc × PC1	-0.152	0.425	Soc × PC2	0.056	0.832
	Reproduction - $|E_{\varphi}^{\sigma}|$	Soc	1.223	0.334	Soc	0.430	0.708
		PC1	3.401	0.129	PC2	-0.574	0.852
		Soc × PC1	-0.976	0.197	Soc × PC2	0.879	0.393

Figure legends

Figure 1. Representation of the 87 natural populations of the 66 animal species examined in this study to test the social buffering hypothesis. A. Geographic location of the populations. B. Phylogenetic relationships of the species. The phylogeny was modified to accommodate intra-specific spatial replication, indicated by the name of the species followed by the number of the population. In total, we examined patterns of demographic buffering across 12 animal taxonomic classes. Each species was classified into one of the five levels of sociality shown in the insert in panel A, following Salguero-Gómez (Salguero-Gómez, 2024).

Figure 2. The examined 87 animal populations are organised along two axes of climatic variation: (PC1) predictability in precipitation and (PC2) constancy of temperature. Principal component analyses of the first two principal components, displaying the loadings of climatic products derived from the NASA POWER database. Climatic variables are: (1) Mean annual total precipitation ($\bar{P}$), (2) Variance annual total precipitation (ΔT), (3) Predictability of annual precipitation (P_{pred}), (4) Mean annual temperature ($T\square$), (5) Variance annual temperature (ΔT); and (6) Constancy of temperature (T_{cons}). The first principal component (PC1), which explains 50% of the variance, positions populations along a continuum of climates with more variable precipitation (right) *vs.* more predictable precipitation (left). The second principal component (PC2), explaining 18% of variance, places populations along a continuum of higher mean annual precipitation (bottom) *vs.* climates with higher constancy in temperature (top). Constancy quantifies the extent to which a climatic variable remains

1022   stable over time. Predictability is the sum of constancy and contingency, where

1023   contingency reflects the extent to which fluctuations are structured and recurrent. The

1024   colours of each dot represent the classification of sociality (See Figure 1). Silhouettes

1025   represent a subset of the animal species shown on the PCA, obtained from phylopic

1026   (Keesey, 2020).

1027

Figure 3. Increases in sociality are associated with canalisation in the selective

pressures of demographic performance and increases in demographic buffering.

Sum of stochastic elasticities of population growth rate (λ_s) to changes in the (**A**)

variance (T_σ) and (**B**) mean (T_μ) of all vital rates in a given population, classified by

degree of sociality. Response variable is the absolute value of the residuals of T_σ and T_μ

against adult body mass of each animal species. T_σ is $\log_{10}$-transformed. P values (top-

right) correspond to a phylogenetic ANOVA, and post-hoc Tukey test letters positioned

on top of each group indicate whether groups are significantly different, after

phylogenetic corrections.

1037

Figure 4. As species become more social, their population performance becomes

less sensitive to changes in juvenile survival and reproduction. Stochastic

elasticity of population growth rate (λ_s) to changes in the variance of four vital rates (vr)

($|E_{vr}^{\sigma}|$): ($|E_{vr}^{\sigma}|$): juvenile survival ($|E_{\sigma J}^{\sigma}|$), maturation ($|E_{\gamma}^{\sigma}|$), adult survival ($|E_{\sigma A}^{\sigma}|$), and

reproduction ($|E_{\varphi}^{\sigma}|$), grouped by degree of sociality. Response variables are the

residuals of E_{vr}^{σ} against adult body mass of each animal species, and $\log_{10}$-

transformed. P values correspond to a phylogenetic ANOVA, and post-hoc Tukey test

letters positioned on top of each group indicate whether groups are significantly

different, after phylogenetic corrections.

1047

Figure 5. In more social species, vital rates are more tightly regulated. Stochastic

elasticity of population growth rate (λ_s) to changes in the mean of four vital rates (*vr*)

($|E_{vr}^{\mu}|$): juvenile survival ($|E_{\sigma_J}^{\mu}|$), maturation ($|E_{\gamma}^{\mu}|$), adult survival ($|E_{\sigma_A}^{\mu}|$), and

reproduction ($|E_{\varphi}^{\mu}|$), grouped by degree of sociality. Response variables are the

absolute values of the residuals of E_{vr}^{σ} against adult body mass of each animal species.

P values correspond to a phylogenetic ANOVA, and post-hoc Tukey test letters

positioned on top of each group indicate whether groups are significantly different, after

phylogenetic corrections.

1056

Figure 1

1060 **Figure 2**

1061

1062

1063

1064

1065 **Figure 3**

1066

Figure 4

Figure 5

 Supplementary Online Materials

1077 Support for the social buffering hypothesis, especially under

1078 unpredictable precipitation regimes

1079

1080 **Table of contents**

1081

Table S1. Sensitivity of results to temporal replication in studies

Our overall results are mostly insensitive to the duration of the study. Here, study duration is defined as the number of matrix population models (MPMs) available in each examined population. Battery of pgls models examining the relationships between the different stochastic elasticities of stochastic population growth rate (λ_s) and sociality, with study duration as a covariate. Note that the latter is only borderline significant in three occasions: T_σ, $|E_{\sigma A}^{\mu}|$ and $|E_{\sigma A}^{\sigma}|$.

Stochastic elasticity to		Predictor	Estimate	P
A. Total		Sociality	0.079	0.768
	Mean - T_μ	Duration	-0.200	0.040
		Sociality	0.807	0.399
	Variance - T_σ	Duration	-0.451	0.062
B. Changes in mean	Juvenile survival - $|E_{\sigma J}^{\mu}|$	Sociality	0.117	0.818
		Duration	-0.086	0.487
		Sociality	-0.738	0.318
	Maturation - $|E_\gamma^{\mu}|$	Duration	-0.023	0.894
	Adult survival - $|E_{\sigma A}^{\mu}|$	Sociality	0.052	0.911
		Duration	-0.241	0.046
	Reproduction - $|E_\varphi^{\mu}|$	Sociality	-0.062	0.938
		Duration	-0.144	0.458
C. Changes in	Juvenile survival -	Sociality	0.108	0.906

variance	$\lvert E_{\sigma J}^{\sigma}\rvert$	Duration	-0.334	0.141
		Sociality	-0.474	0.630
	Maturation - $\lvert E_{\gamma}^{\sigma}\rvert$	Duration	-0.206	0.389
	Adult survival - $\lvert E_{\sigma A}^{\sigma}\rvert$	Sociality	0.185	0.470
		Duration	-0.134	0.042
	Reproduction - $\lvert E_{\varphi}^{\sigma}\rvert$	Sociality	0.592	0.593
		Duration	-0.417	0.131

1100

Table S2. Metadata and source of matrix population models

Sources of the demographic data obtained from COMADRE for the 66 examined species, together with other pertinent details. *Population* details the number of populations available per species in this study. *Dimension* details the number of stages the original matrix population models (MPMs) had before being collapsed to a set of 2×2 MPMs (see Methods). *Sociality* details the assigned level of our sociality continuum, with the explanation for each of the five levels provided in the Methods too.

Species	name	Phylum	Class	Order	Family	Population	Authors	Journal	DOI_ISBN	Year	Dimension	Sociality
Acyrthosiphon pisum	Pea aphid	Arthropoda	Insecta	Aphidomorpha	Aphididae	1	Hamda; Jevtic; Laskowski	Ecotoxicology	10.1007/s10646-012-0904-5	2012	6	Gregarious

Agaricia agaricites	Tan lettuce-leaf coral	Cnidaria	Anthozoa	Scleractinia	Agariciidae	1	Hughes; Tanner	Ecology	10.1890/0012-9658(2000)081[2250:RFLHAL]2.0.CO;2	2000	3	Colonial
Alces alces	Moose	Chordata	Mammalia	Artiodactyla	Cervidae	1	Ballard; Whitman; Reed	Wildlife Monogr	https://www.jstor.org/stable/3830713	1991	3	Solitary
Amblo	Rock	Chorda	Actinop	Percifo	Centrar	1	People	Master	NA	2010	3	Gregari

plites rupestris	bass	ta	terygii	rmes	chidae		s	Thesis				ous
Ampelisca abdita	Amphipod	Arthropoda	Malacostraca	Amphipoda	Ampeliscidae	1	Kuhn; Munns; Serbst; Edwards; Cantwell; Gleason; Pelletier; Berry	Environ Toxicol Chem	10.1002/etc.5620210425	2002	8	Gregarious
Anser caerule	Snow goose	Chordata	Aves	Anseriformes	Anatidae	1	Cooch; Rockw	Ecol Monog	10.1890/0012	2001	5	Colonial

scens							ell; Brault	r	-9615(2001)071[0377:RAODRT]2.0.CO;2			
Anthropoides paradiseus	Blue crane	Chordata	Aves	Gruiformes	Gruidae	1	Altweg; Anderson	Funct Ecol	10.1111/j.1365-2435.2009.01563.x	2009	5	Gregarious
Astroblepus ubidiai	Andean catfish	Chordata	Actinopterygii	Siluriformes	Astroblepidae	1	Vélez-Espino	Ecol Freshw Fish	10.1111/j.1600-	2005	7	Gregarious

									0633.2005.00084.x			
Boloria eunomia	Bog fritillary	Arthropoda	Insecta	Lepidoptera	Nymphalidae	1	Radchuk; Turlure; Schtickzelle	J Anim Ecol	10.1111/j.1365-2656.2012.029.x	2012	5	Solitary
Brachyteles hypoxanthus	Northern muriqui	Chordata	Mammalia	Primates	Atelidae	2	Morris; Altmann; Brockman; Cords; Fediga	Am Nat	10.1086/657443	2011	10	Social

						n; Pusey; Stoinski; Bronikowski; Alberts; Strier					
Caenorhabditis elegans	NA	Nematoda	Secernentea	Rhabditida	Rhabditidae	1	Li; Ju; Liao; Liao	Ecotoxicology	10.1007/s10646-014-1267-x)	2014	6	Gregarious
Callospermop-	Golden-	Chordata	Mammalia	Rodentia	Sciuridae	1	Hostetler;	PLOS ONE	10.1371/jourN	2012	6	Solitary

hilus lateralis	mantled ground squirrel						Kneip; Van Vuren; Oli		Al.pone.0034379			
Capitella capitata	Polychaete	Annelida	Polychaeta	NA	Capitellidae	1	Hansen; Forbes; Forbes	Funct Ecol	10.1046/j.1365-2435.1999.00299.x	1999	2	Solitary
Cebus capucinus	White-faced capuchin monkey	Chordata	Mammalia	Primates	Cebidae	2	Morris; Altman; Brockman; Cords;	Am Nat	10.1086/657443	2011	8	Social

							Fedigan; Pusey; Stoinski; Bronikowski; Alberts; Strier						
Centrocercus minimus	Gunnison sage-grouse	Chordata	Aves	Charadriiformes	Stercorariidae	1	Davis; Hooten; Phillips; Doherty	Ecol Evol	10.1002/ece3.1290	2014	4		Solitary

							Morris; Altmann; Brockman; Cords; Fedigan; Pusey; Stoinski; Bronikowski; Alberts; Strier					
Cercopithecus mitis	Blue monkey	Chordata	Mammalia	Primates	Cercopithecidae	1		Am Nat	10.1086/657443	2011	9	Social
Chloro	Vervet	Chorda	Mamm	Primat	Cercop	1	Isbell;	Int J	10.100	2009	2	Social

cebus aethiops s		ta	alia	es	ithecid ae		Young; Jaffe; Carlson; Chanc ellor	Primat ol	7/s107 64- 009- 9332-7			
Chroso mus oreas	Mounta in redbell y dace	Chorda ta	Actinop terygii	Cyprini formes	Cyprini dae	1	People s	Master Thesis	NA	2010	2	Comm unal
Ciconia ciconia	White stork	Chorda ta	Aves	Ciconiif ormes	Ciconii dae	1	Schau b; Pradel; Lebret on	Biol Conser v	10.101 6/j.bioc on.200 3.11.00 2	2004	2	Gregari ous
Clinoco	Woolly	Chorda	Actinop	Scorpa	Cottida	2	Davis;	Mar	10.335	2002	3	Solitary

ttus analis	sculpin	ta	terygii	eniform es	e		Levin	Ecol Prog Ser	4/meps 234229			
Clinost omus fundulo ides	Rosysi de dace	Chorda ta	Actinop terygii	Cyprini formes	Cyprini dae	1	People s	Master Thesis	NA	2010	2	Comm unal
Colias alexan dra	Queen Alexan dra's sulphur	Arthrop oda	Insecta	Lepido ptera	Pierida e	1	Hayes	Oecolo gia	10.100 7/BF00 349187	1981	7	Solitary
Cottus bairdii	Mottled sculpin	Chorda ta	Actinop terygii	Scorpa eniform es	Cottida e	1	People s	Master Thesis	NA	2010	3	Solitary
Erythro cebus	Patas monke	Chorda ta	Mamm alia	Primat es	Cercop ithecid	1	Isbell; Young;	Int J Primat	10.100 7/s107	2009	2	Social

patas	y				ae		Jaffe; Carlson; Chancellor	ol	64-009-9332-7			
Etheostoma flabellare	Fantail darter	Chordata	Actinopterygii	Perciformes	Percidae	1	Peoples	Master Thesis	NA	2010	3	Solitary
Eulamprus tympanum	Water skink	Chordata	Reptilia	Squamata	Scincidae	1	Blomberg; Shine	Austral Ecol	10.1046/j.1442-9993.2001.01120.x	2001	5	Solitary
Falco	Lesser	Chordata	Aves	Falconi	Falconi	1	Hiraldo	J Appl	10.230	1996	2	Comm

nauma nni	kestrel	ta		formes	dae		; Negro; Donaz ar; Gaona	Ecol	7/2404 688			unal
Falco peregri nus	Peregri ne falcon	Chorda ta	Aves	Falconi formes	Falconi dae	1	Altweg g; Jenkin s; Abadi	Ibis	10.111 1/ibi.12 125	2013	5	Solitary
Forpus passeri nus	Green- rumpe d parrotl ets	Chorda ta	Aves	Psittaci formes	Psittaci dae	1	Sander cock; Beissin ger	J Appl Stat	10.108 0/0266 476012 010881 8	2002	2	Social
Gopher	Desert	Chorda ta	Reptilia	Testudi	Testudi	1	Perez-	Oikos	10.111	2011	4	Comm

us agassizii	tortoise	ta		nes	nidae		Heydrich; Oli; Brown		1/j.1600-0706.2011.19735.x			unal
Gorilla beringei beringei	Mountain gorilla	Chordata	Mammalia	Primates	Hominidae	1	Morris; Altmann; Brockman; Cords; Fedigan; Pusey; Stoinski;	Am Nat	10.1086/657443	2011	11	Social

							Bronikowski; Alberts; Strier					
Helioseris cuculata	Sunray lettuce coral	Cnidaria	Anthozoa	Scleractinia	Agariciidae	1	Hughes; Tanner	Ecology	10.1890/0012-9658(2000)081[2250:RFLHAL]2.0.CO;2	2000	3	Colonial
Homo sapiens	Human	Chordata	Mammalia	Primates	Hominidae	1	Nicol-Harper; Dooley	Popul Ecol	10.1007/s10144-	2018	18	Social

sapien s							; Packm an; Mueller ; Bijak; Hodgs on; Townle y; Ezard		018- 0620-y			
Lagopu s leucura	White- tailed ptarmig an	Chorda ta	Aves	Gallifor mes	Phasia nidae	1	Wilson; Martin	BMC Ecol	10.118 6/1472 -6785- 12-9	2012	2	Solitary
Lagopu s muta	Japane se rock	Chorda ta	Aves	Gallifor mes	Phasia nidae	1	Wilson; Martin	BMC Ecol	10.118 6/1472	2012	2	Solitary

	ptarmigan								-6785-12-9			
Lepus americanus	Snowshoe hare	Chordata	Mammalia	Lagomorpha	Leporidae	1	Meslow; Keith	J Wildlife Manage	10.2307/3799557	1968	4	Solitary
Macaca mulatta	Rhesus macaque	Chordata	Mammalia	Primates	Cercopithecidae	1	Kessler; Pacheco; Rawlings; Ruiz-Lambrides; Delgad	Am J Primatol	10.1002/ajp.22323	2014	5	Social

							o; Sabat					
Membranipora membranacea	Sea mat	Bryozoa	Gymnolaemata	Cheilostomida	Membraniporidae	2	Harvell; Caswell; Simpson	Oecologia	10.1007/BF00323539	1990	5	Colonial
Microtus oeconomus	Root vole	Chordata	Mammalia	Rodentia	Muridae	1	Johannesen; Aars; Andreassen; Ims	Popul Ecol	10.1007/s10144-003-0139-7	2003	3	Solitary
Nocomis	Bluehead	Chordata	Actinopterygii	Cypriniformes	Cyprinidae	1	Peoples	Master Thesis	NA	2010	3	Communal

leptoce phalus	chub											
Notam acropu s eugenii	Tamm ar wallaby	Chorda ta	Mamm alia	Diproto dontia	Macrop odidae	3	Chamb ers; Bencini	Wildlife Res	10.107 1/WR1 0080	2010	2	Gregari ous
Orbicel la annular is	Caribb ean star coral	Cnidari a	Anthoz oa	Sclerac tinia	Faviida e	2	Hughe s; Tanner	Ecolog y	10.189 0/0012 - 9658(2 000)08 1[2250: RFLHA L]2.0.C O;2	2000	3	Coloni al

Orcinus orca	Killer whale	Chordata	Mammalia	Cetacea	Delphinidae	2	Vélez-Espino; Ford; Araújo; Ellis; Parken; Balcomb	Can Tech Report Fish & Aq Sci	978-1-100-23563-9	2014	7	Social
Ovis aries	Soay sheep	Chordata	Mammalia	Artiodactyla	Bovidae	1	Clutton-Brock; Price; Albon; Jewell	J Anim Ecol	10.2307/5330	1992	6	Gregarious
Pagurus	Long-clawed	Arthropoda	Malacostraca	Decapoda	Paguridae	1	Damiani	Ecology	10.1890/04-	2005	3	Solitary

longicarpus	hermit crab								0956			
Pan troglodytes schweinfurthii	Eastern chimpanzee	Chordata	Mammalia	Primates	Hominidae	1	Morris; Altmann; Brockman; Cords; Fedigan; Pusey; Stoinski; Bronikowski; Alberts	Am Nat	10.1086/657443	2011	17	Social

							; Strier				
Papio cynoce phalus	Olive baboon	Chorda ta	Mamm alia	Primat es	Cercop ithecid ae	1	Morris; Altman n; Brock man; Cords; Fediga n; Pusey; Stoinsk i; Bronik	Am Nat	10.108 6/6574 43	2011	8	Social

							owski; Alberts; Strier					
Paramuricea clavata	Violescent sea-whip; Red gorgonian	Cnidaria	Anthozoa	Alcyonacea	Plexauridae	2	Linares; Doak	Mar Ecol Prog Ser	10.3354/meps08437	2010	7	Colonial
Pimephales promelas	Fathead minnow	Chordata	Actinopterygii	Cypriniformes	Cyprinidae	1	Schwindt	NA	NA	2013	4	Communal
Plexau	Gorgon	Cnidaria	Anthozoa	Alcyon	Plexau	1	Lasker	Oecolo	10.100	1991	3	Colonial

ra homom alla	ian coral	a	oa	acea	ridae			gia	7/BF00 318316			al
Porites astreoides	Caribbean coral reef	Cnidaria	Anthozoa	Scleractinia	Poritidae	1	Edmunds	Mar Ecol Prog Ser	10.3354/meps 08595	2010	3	Colonial
Propithecus verreauxi	Verreaux's sifaka	Chordata	Mammalia	Primates	Indriidae	1	Morris; Altmann; Brockman; Cords; Fedigan; Pusey;	Am Nat	10.1086/6574 43	2011	8	Social

							Stoinski; Bronikowski; Alberts; Strier				
Pseudodiploria strigosa	Symmetrical brain coral	Cnidaria	Anthozoa	Scleractinia	Faviidae	1	Edmunds	Mar Ecol Prog Ser	10.3354/meps08595	2010	3	Colonial
Pygoscelis adeliae	Adelie penguin	Chordata	Aves	Sphenisciformes	Spheniscidae	1	Hinke; Trivelpiece; Trivelpiece	Ecosphere	10.1002/ecs2.1666	2017	2	Colonial

Rana temporaria	European common frog	Chordata	Amphibia	Anura	Ranidae	1	Campbell; Garner; Tessa; Scheele; Griffiths; Wilfert; Harrison	PeerJ	NA	2018	11	Gregarious
Sceloporus grammicus	Mesquite lizard	Chordata	Reptilia	Squamata	Phrynosomatidae	2	Méndez–de la Cruz; Zúñiga	Can J Zool	10.1139/Z08-124	2008	3	Solitary

							-Vega; Cuellar					
Scolytus ventralis	Fir engraver beetle	Arthropoda	Insecta	Coleoptera	Curculionidae	1	Berryman	Can Entomol	10.4039/Ent1051465-11	1973	6	Solitary
Strix occidentalis	Northern spotted owl	Chordata	Aves	Strigiformes	Strigidae	1	LaHaye; Zimmerman; Gutiérrez	Auk	10.1642/0004-8038(2004)121[1056:TVITVR]2.0.CO;2	2004	3	Solitary

Suricata suricatta	Meerkat	Meerkat	Mammalia	Carnivora	Herpestidae	1	Conquet,Ozgul, Blumstein, Armitage, Oli, Martin, Clutton-Brock, Paniw	Ecosphere	10.1002/ecy.3894	2023	4	Social
Turdus torquatus	Ring ouzel	Chordata	Aves	Passeriformes	Turdidae	1	Sim; Rebecca; Ludwig; Grant;	J Anim Ecol	10.1111/j.1365-2656.2010.01	2011	2	Solitary

									750.x			
							Reid		750.x			
Umbonium costatum	NA	Mollusca	Gastropoda	Vetigastropoda	Trochidae	1	Noda; Nakao	J Anim Ecol	10.2307/5722	1996	6	Communal
Ursus maritimus	Polar bear	Chordata	Mammalia	Carnivora	Ursidae	1	Hunter; Caswell; Runge; Regehr; Amstrup; Stirling	Ecology	10.1890/09-1641	2010	6	Solitary
Vireo atricapi	Black-capped	Chordata	Aves	Passeriformes	Vireonidae	11	Walker;	Biol Conser	10.1016/j.bioc	2016	2	Solitary

							Marzluf f; Cimprich	v	on.201 6.09.01 6.			
Xenos aurus agreno n	Knob- scaled lizard	Chorda ta	Reptilia	Squam ata	Xenos auridae	1	Zamor a- Abrego ; Chang; Zúñiga -Vega; Nieto- Montes de Oca; Johnso	Herpet ologica	10.165 5/09- 005.1	2010	4	Solitary

							n					
Xenosaurus grandis	Crevice-dwelling lizard	Chordata		Reptilia	Squamata	Xenosauridae	1	Zúñiga-Vega; Valverde; Rojas-Gonzalez; Lemos-Espinal	Copeia	10.1643/0045-8511(2007)7[324:AOTPDO]2.0.CO;2	2007	4	Solitary
Xenosaurus platyceps	Flathead knob-scaled lizard	Chordata		Reptilia	Squamata	Xenosauridae	2	Rojas-Gonzalez; Jones;	Amphibia-Reptilia	10.1163/156853087841249	2008	4	Solitary

						Zúñiga -Vega; Lemos- Espinal	92			

1109

1110

1111 **Table S3. Degree of sociality across the climatic space**

1112 **The degree of sociality is not predicted by the climatic principal component axis.**

1113 Results of a pgls model predicting the degree of sociality as a function of the positioning

1114 of each population along PC1 (Precipitation predictability; Figure 2) and PC2

1115 (temperature constancy).

	Estimate	S. E.	t	P
Intercept	2.616	1.1876	2.2032	0.0436
PC1	0.1150	0.2581	0.4453	0.6624
PC2	0.2682	0.3579	0.7493	0.4652
PC1 × PC2	0.0165	0.1845	0.0894	0.9299

1116

1117

Figure S1. Sampling frequency of the matrix population models

Most of the matrix population models (MPMs) in our selected set of matrix

population models from the COMADRE Animal Matrix Database were sampled

once a year. Species in our study with an original sampling frequency that is not

annual: *Caenorhabditis elegans* (0.000114), *Capitella capitata* (0.0192), *Xenosaurus*

agrenon (0.08), *Membranopira mebranaceae* (0.12), *Acyrthosiphon pisum* (0.02),

Clinocottus analis (0.25), *Ampelisca abdita* (0.03), *Eulamprus tympanum* (0.08),

Sceloporus grammicus (0.5), *Montastrae annularis* (5), and Homo sapiens sapiens (5).

1126

1127

1128 **Figure S2. Matrix population model availability per study**

1129 **Number of matrix population models (MPMs) per studied population in our**

1130 **selected subset of 66 species from the COMADRE Animal Matrix.** Total MPM =

1131 955. Total number of separate populations = 87.

1132

1133 **Figure S3. Matrix population model dimensionality**

1134 **The dimensionality of our examined matrix population models (MPMs) varied**

1135 **considerably between 2 and 18 stages.** As such, we collapsed all models to a two-

1136 stage MPM to allow for stage comparability in vital rates.

1137

 Figure S4. Climate driver collinearities

Some of the examined climatic drivers are strongly collinear. Pairwise correlations

(top-right) and scatterplots (bottom-left) of the statistics of monthly precipitation and

temperature derived from the NASA POWER database to quantify the degree of

environmental stochasticity to which our 87 natural populations have been exposed.

Temperature statistics include: : mean; ΔT: variance: T_{cons}: constancy; T_{cont}:

contingency; and T_{pred}: predictability. Ditto for precipitation (P). Constancy quantifies the

extent to which a climatic variable remains stable over time (1 - variance/mean);

contingency quantifies the extent to which fluctuations are structured and recurrent;

predictability = constancy + contingency, as per Colwell (Colwell, 1974). Font size of

spearman correlation coefficient is proportional to its values.

1149

1150   **Figure S5. PCA screeplot**

1151   **Screeplot of the six principal components of the climatic PCA.** The blue dashed

1152   line separates the principal components axis with associated eigenvalue > 1, which we

1153   retained for the next steps in our analyses, as per the Kaiser criterion.

1154

1155

1156

Figure S6. Climatic predictability of demographic buffering

The predictability of annual precipitation patterns predicts the ability of animal

populations to buffer against extreme climatic events. Scaled effect sizes (negative:

red; positive: blue) and P values for the correlations between the climatic PCAs (Figure

2 - PC1: precipitation predictability, PC2: temperature constancy, and their interaction)

and: (**A**) the sum of total stochastic elasticities to the mean ($|T_\mu|$) and the variance ($|T_\sigma|$),

(**B**) the stochastic elasticities to changes in vital rate means ($|E^\mu|$), and (**C**) to changes in

vital rate variances ($|E^\sigma|$). Vital rates are: juvenile survival (σ_J), maturation (γ), adult

survival (σ_A), and reproduction (φ) (Eq. 1).

1166